March 2021 Volume 28 Number 0

Editors:

Kaye Wilson & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

2	Introducing PHARMAC	
5	General Rules	Section A
6	Alimentary Tract & Metabolism	Section B
41	Blood & Blood Forming Organs	
51	Cardiovascular System	
65	Dermatologicals	
75	Genito Urinary System	
82	Hormone Preparations – Systemic	
93	Infections – Agents For Systemic Use	
114	Musculoskeletal System	
122	Nervous System	
148	Oncology Agents & Immunosuppressants	
223	Respiratory System & Allergies	
232	Sensory Organs	
237	Various	
239	Extemporaneous Compounds (ECPs)	Section C
242	Special Foods	Section D
263	National Immunisation Schedule	Section I
273	Index	

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	✓	Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓	Entocort CIR
⇒SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
1 Mild to moderate ileal, ileocaecal or proximal Crohn's dise	ease; and			

0.4 Dishetes as

2 Any of the following:

2.1 Diabetes; or

2.2 Cushingoid habitus; or

2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	26.55	21.1 g OP	✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLO	RIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg	49.50	100	✓ Asamax
Tab long-acting 500 mg	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g	118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml		7	✓ Pentasa
Suppos 500 mg	22.80	20	✓ Asacol
Suppos 1 g	54.60	30	✓ Pentasa
OLSALAZINE			
Tab 500 mg	93.37	100	Dipentum
Cap 250 mg		100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mgSULFASALAZINE	92.91	100	✓	Nalcrom
* Tab 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g6.35	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg2.66	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy 30 q OP ✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

* Tab 200 mcg - Up to 120 tab available on a PSO41.50 ✓ Cytotec

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement......10.40 14 ✓ Apo-Clarithromycin

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FAI	MOTIDINE - Only on a prescription			
*	Tab 20 mg4.	.91	100	✓ Famotidine
				Hovid S29
*	Tab 40 mg8.	48	100	✓ Famotidine
	•			Hovid S29
*	Inj 10 mg per ml, 4 ml – Subsidy by endorsement57.	.02	10	✓ Mylan S29
	Subsidy by endorsement – Subsidised for patients receiving treatn	nent as part	of palliative	care.

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the
 prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record
 of prior dispensing of ranitidine.

*	Oral liq 150 mg per 10 ml5.14	300 ml	✓ Peptisoothe
*	Inj 25 mg per ml, 2 ml	5	✓ Zantac
/D	anticoath a Oval lin 150 may now 10 miles be delicted 1 Contambay 2001)		

(Peptisoothe Oral liq 150 mg per 10 ml to be delisted 1 September 2021) (Zantac Inj 25 mg per ml, 2 ml to be delisted 1 June 2021)

Proton Pump Inhibitors

100	✓ Lanzol Relief
100	✓ Lanzol Relief
90	Omeprazole actavis10
90	Omeprazole actavis20
90	Omeprazole actavis 40
5 q	✓ Midwest
Ü	
5	✓ Dr Reddy's
	Omeprazole
	✓ Ocicure S29
100	✓ Panzop Relief
100	✓ Panzop Relief
	90 90 90 5 g 5

	Subsidy (Manufacturer's Price \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mgSUCRALFATE	14.51	50	√ (Gastrodenol 629
Tab 1 g	35.50 (48.28)	120	(Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Reta	ail pharmacy			
Tab 550 mg	625.00	56	✓ <u>></u>	<u> (ifaxan</u>
➤SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatol epatologist. Approvals valid for 6 months where the patiolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Piepatologist. Approvals valid without further renewal unleasenefiting from treatment.	ent has hepatic encephalop Practitioner on the recomme	athy d	espite an a	dequate trial of maximun penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Ref Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 0 ml 0	✓ F	Proglicem \$29 Proglicem \$29 Proglycem \$29
■ SA1320 Special Authority for Subsidy nitial application from any relevant practitioner. Approvypoglycaemia caused by hyperinsulinism.	als valid for 12 months whe	re use	d for the tre	atment of confirmed
Renewal from any relevant practitioner. Approvals valid of ppropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE				
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓ (Glucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL Inj human 100 u per ml	25.26 1	0 ml C		Actrapid Humulin R
Inj human 100 u per ml, 3 ml	42.66	5	√	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE				
Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ 1	NovoMix 30 FlexPen

	Subsidy		Fully Brand or
	(Manufacturer's Prices)	ce) Subs Per	idised Generic Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH
,			✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
,			✓ Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30
			✓ PenMix 40
			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml	,		
3 ml		5	Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,	,		
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE			
▲ Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin David Astina Damentians			
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
▲ Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
▲ Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE			
▲ Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
▲ Inj 100 u per ml, 3 ml	46.07	5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ Apidra SoloStar
NSULIN LISPRO			
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
▲ Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
* Tab 50 mg	3.50	90	✓ Glucobay
-	10.47		✓ Accarb
* Tab 100 mg	6.40	90	✓ Glucobay
	20.23		✓ Accarb
Blood Glucose Lowering Agents			
blood didcose Loweling Agents			
EMPAGLIFLOZIN - Special Authority see SA2019 on the next p		nacy	
* Tab 10 mg		30	✓ Jardiance
* Tab 25 mg	58.56	30	Jardiance

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

⇒SA2019 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Any of the following:
 - 2.1 Patient is Maaori or any Pacific ethnicity*: or
 - 2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.5 Patient has diabetic kidney disease (see note b)*; and
- 3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months; and
- 4 Treatment will not be used in combination with a funded GLP-1 agonist.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2020 on the next page - Retail pharmacy

*	Tab 5 mg with 1,000 mg metformin hydrochloride		60	✓ Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	✓ Jardiamet

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2020 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Any of the following:
 - 2.1 Patient is Maaori or any Pacific ethnicity*: or
 - 2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.5 Patient has diabetic kidney disease (see note b)*; and
- 3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months; and
- 4 Treatment will not be used in combination with a funded GLP-1 agonist.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

GLIBENCLAMIDE		
* Tab 5 mg	100	✓ Daonil
GLICLAZIDE		
* Tab 80 mg	500	✓ Glizide
GLIPIZIDE		
* Tab 5 mg	100	✓ Minidiab
METFORMIN HYDROCHLORIDE		
* Tab immediate-release 500 mg	1,000	✓ Apotex
* Tab immediate-release 850 mg	500	✓ Apotex
PIOGLITAZONE		
* Tab 15 mg	90	✓ Vexazone
* Tab 30 mg	90	✓ Vexazone
* Tab 45 mg	90	✓ <u>Vexazone</u>
VILDAGLIPTIN		
Tab 50 mg35.00	60	✓ Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE		
Tab 50 mg with 1,000 mg metformin hydrochloride35.00	60	✓ Galvumet
Tab 50 mg with 850 mg metformin hydrochloride35.00	60	Galvumet

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips ______20.00 1 OP

CareSens Dual

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

20.00

✓ CareSens N Premier

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRC

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

IIVC	oblini bin nebelo inaximam of 200 dev per prescripti	OH		
*	29 g × 12.7 mm	10.50	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	00 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.00	100	✓ B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year p	eriod.		
Min basal rate 0.025 U/h		1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- Fither
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic	
(Wallalacaters i nee)	Per	J	Manufacturer	
Ψ	rei	•	Manuacturei	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment: and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

continued...

pump therapy; and

- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

INSULIN PUMP CARTRIDGE - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of cartridge sets will be funded per year.

Cartridge 300 U, t:lock × 10......50.00 1 OP ✓ Tandem Cartridge

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

- 6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×
 10 with 10 needles......130.00 1 OP Paradigm Sure-T
 MMT-864
- 6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock.......130.00 1 OP ✓ Sure-T MMT-863
- 6 mm steel needle; 29 G; manual insertion; 80 cm tubing ×
 10 with 10 needles.......130.00 1 OP Paradigm Sure-T

- 8 mm steel needle; 29 G; manual insertion; 80 cm tubing ×
 10 with 10 needles......130.00 1 OP ✓ Paradigm Sure-T
 MMT-876

(Paradigm Sure-T MMT-884 10 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-886 10 mm steel needle; 29 G; manual insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

Clear and Sure-T MMT-864 6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-866 6 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-874 8 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Sure-T MMT-876 8 mm steel needle; 29 G; manual insertion; 80 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	. 130.00	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	. 130.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- a) Maximum of 12 infusion sate will be funded nor year

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A
			IVIIVI I -OOOA

Subsidy		Fully	Brand or	
(Manufacturer's Price)	nufacturer's Price) Subsidised		Generic	
\$	Per	✓	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon cannula; angle insertion; 120 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-382
13 mm teflon cannula; angle insertion; 45 cm line \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-368
13 mm teflon cannula; angle insertion; 60 cm line \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-381
13 mm teflon cannula; angle insertion; 80 cm line \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-383
17 mm teflon cannula; angle insertion; 110 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-377
17 mm teflon cannula; angle insertion; 60 cm line × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Silhouette MMT-378
17 mm teflon cannula; angle insertion; 60 cm line × 10 with			
10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-373
17 mm teflon cannula; angle insertion; 80 cm line × 10 with			
10 needles	130.00	1 OP	 Paradigm Silhouette

(Paradigm Silhouette MMT-382 13 mm teflon cannula; angle insertion; 120 cm line x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Silhouette MMT-368 13 mm teflon cannula; angle insertion; 45 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-381 13 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-383 13 mm teflon cannula; angle insertion; 80 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-377 17 mm teflon cannula; angle insertion; 110 cm line × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Silhouette MMT-378 17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles to be delisted 1 April 2021) (Paradigm Silhouette MMT-384 17 mm teflon cannula; angle insertion; 80 cm line × 10 with 10 needles to be delisted 1 April 2021)

6 mm teflon cannula; straight insertion; insertion device;

9 mm teflon cannula; straight insertion; insertion device;

6 mm teflon cannula: straight insertion: insertion device: 60 cm

110 cm line × 10 with 10 needles140.00

line × 10 with 10 needles......140.00

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	/	Manufacturer	

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.		
6 mm teflon cannula; straight insertion; insertion device; 45 cm		
blue tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-941
6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-921
6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-943
6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-923
6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-945
6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-965
6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing × 10 with 10 needles	1 OP	✓ Paradigm Mio MMT-925
9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing × 10 with 10 needles130.00	1 OP	✓ Paradigm Mio MMT-975

1 OP

1 OP

1 OP

✓ AutoSoft 90

✓ AutoSoft 90

✓ AutoSoft 90

Subs (Manufactur	. ,	Fully lised	Brand or Generic	
\$	Per	1	Manufacturer	

9 mm teflon cannula; straight insertion; insertion device; 60 cm

(Paradigm Mio MMT-941 6 mm teflon cannula; straight insertion; insertion device; 45 cm blue tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-921 6 mm teflon cannula; straight insertion; insertion device; 45 cm pink tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-943 6 mm teflon cannula; straight insertion; insertion device; 60 cm blue tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-923 6 mm teflon cannula; straight insertion; insertion device; 60 cm pink tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-945 6 mm teflon cannula; straight insertion; insertion device; 80 cm blue tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-965 6 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-925 6 mm teflon cannula; straight insertion; insertion device; 80 cm pink tubing \times 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Mio MMT-975 9 mm teflon cannula; straight insertion; insertion device; 80 cm clear tubing \times 10 with 10 needles to be delisted 1 April 2021)

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-398
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with			45 " 6 116.
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-399
6 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with			
10 needles; luer lock	130.00	1 OP	Quick-Set MMT-393
6 mm teflon cannula; straight insertion; 80 cm tubing \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-387
9 mm teflon cannula; straight insertion; 106 cm tubing \times 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-396
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-397
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with			
10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-392
9 mm teflon cannula; straight insertion; 80 cm tubing × 10 with			
10 needles	130.00	1 OP	✓ Paradigm Quick-Set MMT-386

(Paradigm Quick-Set MMT-398 6 mm teflon cannula; straight insertion; 110 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-399 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-387 6 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-396 9 mm teflon cannula; straight insertion; 106 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-397 9 mm teflon cannula; straight insertion; 60 cm tubing × 10 with 10 needles to be delisted 1 April 2021)

(Paradigm Quick-Set MMT-386 9 mm teflon cannula; straight insertion; 80 cm tubing x 10 with 10 needles to be delisted 1 April 2021)

INSULIN PUMP RESERVOIR - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 packs of reservoir sets will be funded per year.

10 × luer lock conversion cartridges 1.8 ml for Paradigm pumps50.00	1 OP	✓ ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 1050.00	1 OP	✓ MiniMed
		1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 1050.00	1 OP	✓ MiniMed 3.0 Reservoir MMT-332A

Fully

منطنممط

Brand or

	(Manufacturer's Price)	Per Su	Desidised Generic ✓ Manufacturer	
Digestives Including Enzymes				
PANCREATIC ENZYME				
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ <u>Creon 10000</u>	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))	*	100	✓ Panzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph	1			
Eur U)	34.93	20 g OP	Creon Micro	
URSODEOXYCHOLIC ACID – Special Authority see SA1739 be Cap 250 mg	•	cy 100	✓ <u>Ursosan</u>	
<u> </u>				

Subsidy

(Manufacturaria Drica)

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

ner.

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soln	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		-	
* Dry	6.02	500 g OP	
·	(17.32)	•	Normacol Plus
	2.41	200 g OP	
	(8.72)	•	Normacol Plus

Faecal Softeners

DOCOGATE GODIOW - Only of a prescription		
* Tab 50 mg	100	✓ Coloxyl
* Tab 120 mg	100	✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES		
* Tab 50 mg with sennosides 8 mg3.10	200	✓ Laxsol
POLOXAMER – Only on a prescription		
Not funded for use in the ear.		
* Oral drops 10%	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below – Retail	oharmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
,	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$) S Per	ubsidised	Generic Manufacturer
	Ψ	1 61	<u> </u>	Manuacturei
Osmotic Laxatives				
GLYCEROL				
* Suppos 3.6 g - Only on a prescription	9.25	20	✓ <u>P</u>	<u>SM</u>
LACTULOSE - Only on a prescription				
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ <u>L</u>	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI	CARBONATE AND	SODIUN	A CHLORII	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 n	ng,			
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg6.70	30	✓ N	lolaxole
SODIUM ACID PHOSPHATE - Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	√ F	leet Phosphate
·				Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE	- Only on a prescri	iption		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	'		
5 ml		50	✓ N	licolette
Stimulant Laxatives				
BISACODYL – Only on a prescription				
* Tab 5 mg	5.99	200	√ L	ax-Tab
* Suppos 10 mg		10	_	ax-Suppositories
SENNA – Only on a prescription			_	
* Tab, standardised	2.17	100		
	(8.21)		S	enokot
	0.43	20	_	
	(2.06)		S	enokot

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

75.00 180 g OP **Cystadane**

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency: or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1988 below - Retail pharmacy

Inj 1 mg per ml, 5 ml vial......2,234.00

1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
IDURSULFASE – Special Authority see SA1623 below – Retail pharmacy Inj 2 mg per ml, 3 ml vial		1	√ E	laprase

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE − Special Authority see SA1695 below − Retail pharmacy
Inj 100 U per ml, 5 ml vial.......1,335.16

✓ Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

Soln 100 mg per mlCBS 100 ml ✓ Amzoate S29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 below - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

1 ✓ Elelvso

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Email: gaucherpanel@pharmac.govt.nz Wellington

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific
 deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomeoaly: or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

	Subsidy (Manufacturer's P \$	Price) Subs	Fully Brand or sidised Generic Manufacturer	
Mouth and Throat				
Agents Used in Mouth Ulceration				
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with				
Endorsement	9.00 (20.31)	500 ml	Difflam	
Additional subsidy by endorsement for a patient who hap rescription is endorsed accordingly.	as oral mucositis a	as a result of tre	eatment for cancer, and the	
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			_	
Paste	17.20 4.55	56 g OP 15 g OP	✓ Stomahesive	
	(7.90)	10 9 01	Orabase	
	1.52	5 g OP		
2. 1	(3.60)	00 00	Orabase	
Powder	8.48 (10.95)	28 g OP	Stomahesive	
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	(10.93)		Stomanesive	
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP		
The framework got 0.7 /0 with obtained and office of 1/2	(6.00)	10 9 01	Bonjela	
TRIAMCINOLONE ACETONIDE				
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase	
Oropharyngeal Anti-infectives				
AMPHOTERICIN B				
Lozenges 10 mg	5.86	20	✓ Fungilin	
MICONAZOLE				
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>	
NYSTATIN			4	
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>	
Other Oral Agents				
For folinic mouthwash, pilocarpine oral liquid or saliva substitute	formula refer Sta	ndard Formula	ne, page 239	
THYMOL GLYCERIN				
* Compound, BPC	9.15	500 ml	✓ PSM	
Vitamins				
Vitamin B				
HYDROXOCOBALAMIN				
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a F	PSO 1.89	3	✓ Neo-B12	
		-	✓ Vita-B12	
	3.15	5	✓ Hydroxocobalamin	
			Mercury Pharma	

Subsidy

Fully

Brand or

(M	Subsidy anufacturer's Price \$	e) Per	Fully Subsidised	
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription				
♣ Tab 25 mg — No patient co-payment payable ♣ Tab 50 mg		90 500		Vitamin B6 25 Apo-Pyridoxine
HIAMINE HYDROCHLORIDE – Only on a prescription Tab 50 mg		100		Max Health
★ Tab, strong, BPC	7.15	500	✓	Bplex
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription * Tab 100 mg	9.90	500	√	<u>Cvite</u>
Vitamin D				
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml	87.98	100 100 20 ml C	•	One-Alpha One-Alpha One-Alpha
CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg		100 100		Calcitriol-AFT Calcitriol-AFT
COLECALCIFEROL Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription. Coral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml (<u>Vit.D3</u> Puria
Multivitamin Preparations				
# Cap* Cap		30	/	Clinicians Renal Vit

⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

200 g OP ✓ Paediatric Seravit

⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	Brand or Generic Manufacturer
VITAMINS				
* Tab (BPC cap strength) * Cap (fat soluble vitamins A, D, E, K) – Special Authority see	11.45	1,000) ✓ <u>M</u>	<u>lvite</u>
SA1720 below – Retail pharmacy	23.40	60	✓ V	itabdeck

⇒SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

	V	in	er	a	S
--	---	----	----	---	---

Calcium
CALCILIM CARRONATE

CA	LCIUM CARBONATE		
*	Tab eff 1.75 g (1 g elemental)	20	✓ Calcium Sandoz S29
*	Tab 1.25 g (500 mg elemental)6.69	250	✓ Calci-Tab 500
	7.52		✓ Arrow-Calcium
	Calci-Tab 500 to be Sole Supply on 1 May 2021		
*	Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement54.60	76	✓ Cacit S29
	Subsidy by endorsement – Only when prescribed for paediatric patier	its (< 5 years) wl	here calcium carbonate oral liquid is

considered unsuitable. (Calcium Sandoz S29 Tab eff 1.75 g (1 g elemental) to be delisted 1 April 2021) (Arrow-Calcium Tab 1.25 g (500 mg elemental) to be delisted 1 May 2021)

CALCIUM GLUCONATE

*	Inj 10%, 10 ml ampoule	32.00	10	Max Health -
				Hameln \$29
		64.00	20	✓ May Health 920

Fluoride

SODIUM FLUORIDE

*	Tab 1.1 mg (0.5 mg elementa)5.75	100	✓ PSM

lodine

POTASSIUM IODATE

Tab 253 mcg (150 mc	g elemental iodine)4.58	90	✓ NeuroTabs
---------------------	--------------------	-------	----	-------------

Iron

FERRIC CARBOXYMALTOSE – Special Authority see S/	A1840 below – Retail pharmacy	1
Ini 50 mg per ml. 10 ml	150.00	✓ Feriniect

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
 - 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

* Tab 200 mg (65 mg elemental)3.09	100	✓ Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg4.68	60	✓ <u>Ferro-F-Tabs</u>
FERROUS SULFATE * Oral liq 30 mg (6 mg elemental) per 1 ml	500 ml	✓ Ferodan
FERROUS SULPHATE * Tab long-acting 325 mg (105 mg elemental)2.06	30	✓ Ferrograd
IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule	5	✓ Ferrosig

Magnesium

For magnesium hydroxide mixture refer Standard Formulae, page 239

MAGNESIUM HYDROXIDE

(1	Subsidy Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
MAGNESIUM SULPHATE				
* Inj 2 mmol per ml, 5 ml ampoule	25.53 28.00	10	✓	Martindale DBL DBL S29 S29
(DBL Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 July 2021) (DBL S29 S29 Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 Ju	ıly 2021)			
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	/	<u>Zincaps</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — **(chronic renal failure)** from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу			
Wastage claimable			_		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	<u>Binocrit</u>	
Inj 2,000 iu in 1 ml, syringe	100.00	6	✓	<u>Binocrit</u>	
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit	
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit	
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit	
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	1	Binocrit	
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	✓	Binocrit	
Inj 10,000 iu in 1 ml, syringe	197.50	6	✓	Binocrit	
Inj 40,000 iu in 1 ml, syringe		1	✓]	Binocrit	

Megaloblastic

FOL	IC	AC	ID

*	Tab 0.8 mg	21.84	1,000	1	Apo-Folic Acid
*	Tab 5 mg	12.12	500	1	Apo-Folic Acid
	Oral lig 50 mcg per ml	26.00	25 ml OP	1	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	· 1	✓ Alprolix
Inj 500 iu vial		1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
lnį 2,000 iu vial		1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
ELTROMBOPAG – Special Authority see SA174	•		·

Tab 50 mg3,100.00 **➤ SA1743** Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

continued...

✓ Revolade

✓ Revolade

28 28

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
 \$	Per	✓	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, cubiact to critaria

Subject to criteria.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

NONACOG GAMMA. [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

nj 500 iu vial	435.00	1	✓ RIXUBIS
nj 1,000 iu vial	870.00	1	✓ RIXUBIS
nj 2,000 iu vial		1	✓ RIXUBIS
nj 3,000 iu vial	·	1	✓ RIXUBIS
) -/ · · · · · · · · · · · · · · · · ·	,		

	Cubaidu		South	Drand or
	Subsidy (Manufacturer's Price) \$	Subsidi Per	=ully ised ✓	d Generic
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [•			-
For patients with haemophilia. Preferred Brand of short half-l	ife recombinant facto	r VIII. Acces	ss t	o funded treatment is
managed by the Haemophilia Treaters Group in conjunction v	with the National Hae			
Inj 250 iu vial		1	1	Advate
Inj 500 iu vial		1		Advate
Inj 1,000 iu vial		1		Advate
Inj 1,500 iu vial	•	1		Advate
Inj 2,000 iu vial	•	1	_	Advate
Inj 3,000 iu vial	•	1	•	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE I				Section VIIII Access to C. C. C.
For patients with haemophilia. Rare Clinical Circumstances E				
treatment is managed by the Haemophilia Treaters Group in a subject to criteria.	Conjunction with the P	vauonai Hät	eil)(oprillia ivianagement Group,
subject to criteria. Ini 250 iu vial	237 50	1	/	Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial	,	1		Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				-
For patients with haemophilia A receiving prophylaxis treatme		treatment	is n	nanaged by the Haemonhilia
Treaters Group in conjunction with the National Haemophilia				5,
Inj 250 iu vial		1	•	Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1	✓	Adynovate
lnj 2,000 iu vial	2,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml	28.50	5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	•	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO		5	1	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Konakion MM
			F	
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	•	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
-				Multichem
DIPYRIDAMOLE				_
* Tab long-acting 150 mg	10.90	60	/	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next page		•		
* Tab 90 mg		56	/	Brilinta
1 ab 00 mg		50	•	Simila .

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

S	Subsidy	Fully	Brand or
(Manufa	cturer's Price)	Subsidised	Generic
	\$ Per	r √	Manufacturer

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome: and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

10

Fully

Brand or

Clexane

	(Manufacturer's Price)	Subsidised		Generic		
	\$	Per	1	Manufacturer		
Heparin and Antagonist Preparations						
ENOXAPARIN SODIUM - Special Authority see SA1646 below						
Inj 20 mg in 0.2 ml syringe	27.93	10	√ C	lexane		

Subsidy

Inj 60 mg in 0.6 ml syringe	56.18	10	Clexane
Inj 80 mg in 0.8 ml syringe	74.90	10	Clexane
Inj 100 mg in 1 ml syringe		10	✓ Clexane
Inj 120 mg in 0.8 ml syringe	116.55	10	 Clexane Forte
Inj 150 mg in 1 ml syringe	133.20	10	Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Low molecular weight heparin treatment is required during a patients pregnancy; or

- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

LIEDADINI CODILINA

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPAKIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml	32.66	5	✓ DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	✓ Hospira
	42.40		✓ Heparin DBL S29
HEPARINISED SALINE			
Ini 10 ju per ml. 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg	76.36	60	✓	Pradaxa
Cap 150 mg		60	1	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	✓	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	1	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
WARFARIN SODIUM Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	1	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg		100	1	Marevan
* Tab 5 mg		50	✓	Coumadin
•	11.48	100	1	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail p	harmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe		10	1	Nivestim
OA4050 On a stat Authority for Outside.				

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC $< 0.5 \times 10^9$ /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price \$	e) Si Per	Fully ubsidised	Brand or Generic Manufacturer
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE] # Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO # Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE		5 1	_	Biomed Biomed
* Inj 75 mg per ml, 10 ml	55.00	50	✓ J	AstraZeneca Juno §29 Potassium Chloride Aguettant §29
SODIUM BICARBONATE Inj 8.4%, 50 ml a) Up to 5 inj available on a PSO b) Not in combination	19.95	1	√ E	Biomed
Inj 8.4%, 100 ml	20.50	1	√ E	Biomed
SODIUM CHLORIDE Not funded for use as a nasal drop. Not funded for nebulise for nebuliser use.	er use except when u	used in co	njunction	with an antibiotic intended
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.23 1.26	500 ml 1.000 ml		Baxter Baxter
Only if prescribed on a prescription for renal dialysis, more for emergency use. (500 ml and 1,000 ml packs)	aternity or post-natal	care in th	ne home (of the patient, or on a PSO
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5 239	✓ E	Biomed
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO		20	✓ <u>F</u>	resenius Kabi
Inj 0.9%, 10 ml ampoule - Up to 5 inj available on a PSO		50	_	resenius Kabi
Inj 0.9%, 20 ml ampoule TOTAL PARENTERAL NUTRITION (TPN)	5.00	20	✓ <u>F</u>	resenius Kabi
Infusion	CBS	1 OP	√ 1	'PN
WATER 1) On a prescription or Practitioner's Supply Order only was Schedule requiring a solvent or diluent; or 2) On a bulk supply order; or 3) When used in the extemporaneous compounding of eyel When used for the dilution of sodium chloride soln 7%	ye drops; or		•	isted in the Pharmaceutical
lai 5 ml anna cula . Ha ta 5 ini availahla an 3 200	7.00	50		unto vDlo o vuo o
Inj 5 ml ampoule – Up to 5 inj available on a PSOInj 10 ml ampoule – Up to 5 inj available on a PSO		50 50		nterPharma Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO		20	•	riizer Fresenius Kabi
ing 20 mil amposite — Op to 3 mil available on a 1 30		20		Aultichem
(InterPharma Inj 5 ml ampoule to be delisted 1 June 2021) (InterPharma Inj 20 ml ampoule to be delisted 1 June 2021)	7.50	30	✓ I	nterPharma

	Subsidy (Manufacturer's F \$	Price) Subsi	Fully Brand or idised Generic Manufacturer
Oral Administration			
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	9.77	50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)		1,000 ml OP	✓ <u>Pedialyte -</u> Bubblegum
PHOSPHORUS			
Tab eff 500 mg (16 mmol)	82.50	100	✓ Phosphate Phebra
POTASSIUM CHLORIDE			
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (11.85)	60	Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90 [′]	200	✓ Span-K
SODIUM BICARBONATE			
Cap 840 mg	8.52	100	✓ Sodibic✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE			

454 g OP

✓ Resonium-A

			CARDIO	VASC	ULAR SYSTEM
		Subsidy (Manufacturer's Price	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
A	Alpha-Adrenoceptor Blockers				
Δ	Alpha Adrenoceptor Blockers				
DC	DXAZOSIN				
*		8.95	500	✓ A	po-Doxazosin
*	°		500		po-Doxazosin
PH	HENOXYBENZAMINE HYDROCHLORIDE				
*	Cap 10 mg	65.00	30	✓ B	NM S29
		216.67	100	✓ D	ibenzyline \$29
PE	RAZOSIN				,
*	Tab 1 mg	5.53	100	✓ A	po-Prazosin
*	Tab 2 mg		100		po-Prazosin
*	Tab 5 mg	11.70	100	✓ A	po-Prazosin
TE	RAZOSIN - Subsidy by endorsement				
	Subsidy by endorsement – Subsidised for patients who were endorsed accordingly. Pharmacists may annotate the presc dispensing of terazosin.				
	Tab 2 mg	7.50	500	✓ A	po-Terazosin
		14.20	28		eva S29
	Tab 5 mg	10.90	500	✓ A	po-Terazosin
	•	24.80	28	✓ To	eva S29
A	Agents Affecting the Renin-Angiotensin Systen	ı			
Δ	ACE Inhibitors				
C 4	APTOPRIL				
	Oral liq 5 mg per ml		95 ml OP 100 ml OP		apoten aptopril-Mylan §29)
CII	Oral liquid restricted to children under 12 years of age. LAZAPRIL				

CILAZAPRIL * Tab 0.5 mg 2.09 * Tab 2.5 mg 4.80 Tab 5 mg 8.35	90 90 90	✓ <u>Zapril</u> ✓ <u>Zapril</u> ✓ Zapril
ENALAPRIL MALEATE		
* Tab 5 mg	100	✓ Acetec
* Tab 10 mg2.02	100	✓ Acetec
* Tab 20 mg	100	✓ Acetec
LISINOPRIL		
* Tab 5 mg	90	✓ Ethics Lisinopril
* Tab 10 mg	90	✓ Ethics Lisinopril
* Tab 20 mg	90	 Ethics Lisinopril
PERINDOPRIL		
Tab 2 mg	30	✓ Apo-Perindopril
4.95		✓ Coversyl
Tab 4 mg4.80	30	✓ Apo-Perindopril
6.30		✓ Coversyl

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
QUINAPRIL * Tab 5 mg * Tab 10 mg * Tab 20 mg	3.16	90 90 90	✓ Arrow-Quinapril 5 ✓ Arrow-Quinapril 10 ✓ Arrow-Quinapril 20
ACE Inhibitors with Diuretics			
CILAZAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy Subsidy by endorsement – Subsidised for patients who 2020 and the prescription is endorsed accordingly. Phaexists a record of prior dispensing of cilazapril with hydrocentric substance.	were taking cilazapril with armacists may annotate the ochlorothiazide.	pres	scription as endorsed where there
* Tab 5 mg with hydrochlorothiazide 12.5 mg	10.18	100	✓ Apo-Cilazapril/ Hydrochlorothiazide
 (Apo-Cilazapril/ Hydrochlorothiazide Tab 5 mg with hydroch QUINAPRIL WITH HYDROCHLOROTHIAZIDE Tab 10 mg with hydrochlorothiazide 12.5 mg * Tab 20 mg with hydrochlorothiazide 12.5 mg 	3.57 3.83	28 30 30	✓ Accuretic ✓ Accuretic 10 ✓ Accuretic 20
Angiotensin II Antagonists			
CANDESARTAN CILEXETIL * Tab 4 mg * Tab 8 mg * Tab 16 mg * Tab 32 mg LOSARTAN POTASSIUM * Tab 12.5 mg	2.28 3.67 6.39	90 90 90 90	✓ Candestar ✓ Candestar ✓ Candestar ✓ Candestar ✓ Candestar ✓ Losartan Actavis
* Tab 25 mg * Tab 50 mg * Tab 100 mg	2.25	84 84 84	 ✓ <u>Losartan Actavis</u> ✓ <u>Losartan Actavis</u> ✓ <u>Losartan Actavis</u>
Angiotensin II Antagonists with Diuretics			
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZID Tab 50 mg with hydrochlorothiazide 12.5 mg	_	30	✓ <u>Arrow-Losartan &</u> <u>Hydrochlorothiazide</u>
Angiotensin II Antagonists with Neprilysin I	nhibitors		
SACUBITRIL WITH VALSARTAN - Special Authority see S Note: Due to the angiotensin II receptor blocking activit			

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/10:

Subsidy	Ful	ly Brand or	_
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per •	Manufacturer	

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 122

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lighticalitie hydrochionide refer to NERVOUS SYSTEM, Anaesthetics, Local,	page 122	
AMIODARONE HYDROCHLORIDE	00	Awataa
▲ Tab 100 mg	30	✓ <u>Aratac</u>
▲ Tab 200 mg	30	✓ <u>Aratac</u>
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSO16.37	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO12.07	10	✓ Martindale
	10	· <u>martinaalo</u>
DIGOXIN	040	(Lawrente BO
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ <u>Lanoxin PG</u>
* Tab 250 mcg – Up to 30 tab available on a PSO	240	Lanoxin
* Oral liq 50 mcg per ml	60 ml	✓ Lanoxin
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	✓ Rythmodan
FLECAINIDE ACETATE		•
▲ Tab 50 mg	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide
		Controlled
		Release Teva
▲ Cap long-acting 200 mg	90	✓ Flecainide
Cap long acting 200 mg	30	Controlled
		Release Teva
Ini 10 ma nor ml 15 ml amnorda	-	
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor

	Subsidy		Fully	Brand or
	Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100	1	ANI S29
			1	Mexiletine
				Hydrochloride
				USP S29
▲ Cap 250 mg	202.00	100	/	Mexiletine
, ,				Hydrochloride
				USP S29
PROPAFENONE HYDROCHLORIDE				
▲ Tab 150 mg	40.90	50	1	Rytmonorm
·				,
Antihypotensives				
· · · · · · · · · · · · · · · · · · ·				
MIDODRINE - Special Authority see SA1474 below - Retail pharr	nacy			
Tab 2.5 mg	53.00	100	✓	Gutron
Tab 5 mg	79.00	100	✓	Gutron

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL	4.26	500	✓ Mylan Atenolol
* Tab 100 mg		500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml		300 ml OP	✓ Atenolol AFT ✓ Atenolol AFT S29 S29
Restricted to children under 12 years of age.			
BISOPROLOL FUMARATE			
* Tab 2.5 mg	1.84	90	 Bisoprolol Mylan
-	3.53		✓ Bosvate
Bisoprolol Mylan to be Sole Supply on 1 April 2021			
* Tab 5 mg	2.55	90	 Bisoprolol Mylan
-	5.15		✓ Bosvate
Bisoprolol Mylan to be Sole Supply on 1 April 2021			
* Tab 10 mg	3.62	90	 Bisoprolol Mylan
-	9.40		✓ Bosvate
Bisoprolol Mylan to be Sole Supply on 1 April 2021			

(Bosvate Tab 2.5 mg to be delisted 1 April 2021)

(Bosvate Tab 5 mg to be delisted 1 April 2021)

(Bosvate Tab 10 mg to be delisted 1 April 2021)

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	✓ Manufacturer
CARVEDILOL			
* Tab 6.25 mg	2.24	60	✓ Carvedilol Sandoz
* Tab 12.5 mg		60	✓ Carvedilol Sandoz
* Tab 25 mg	2.95	60	✓ Carvedilol Sandoz
CELIPROLOL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who we endorsed accordingly. Pharmacists may annotate the pres dispensing of celiprolol.			
* Tab 200 mg	21.40	180	✓ Celol
(Celol Tab 200 mg to be delisted 1 April 2021)			
LABETALOL			
* Tab 100 mg	14 50	100	✓ Trandate
* Tab 200 mg		100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	- <u>ITANAACE</u>
The first porting porting 20 his ampoulo	(88.60)	Ü	Trandate
* inj 5 mg per ml, 20 ml vial		1	a dato
, , , , , , , , , , , , , , , , , , , ,	(48.20)		Alvogen S29
METOPROLOL SUCCINATE	(10.20)		7 og 6.1.
	1 45	30	✓ Betaloc CR
* Tab long-acting 23.75 mg * Tab long-acting 47.5 mg		30	✓ Betaloc CR
* Tab long-acting 95 mg		30	✓ Betaloc CR
* Tab long-acting 190 mg		30	✓ Betaloc CR
METOPROLOL TARTRATE		00	- Bottaleo ori
	E 66	100	✓ Apo-Metoprolol
* Tab 50 mg Tab 100 mg		60	✓ Apo-Metoprolol ✓ Apo-Metoprolol
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial		5	✓ Metoprolol IV Mylan
, , ,	20.30	J	• <u>metoproloriv mylan</u>
NADOLOL * Tob 40 mg	16.60	100	✓ Apo-Nadolol
* Tab 40 mg Tab 80 mg		100	✓ Apo-Nadolol
•	20.43	100	Apo-Nadoloi
PINDOLOL	40.00	400	4 A DI 111
* Tab 5 mg		100	Apo-Pindolol
* Tab 10 mg		100	✓ Apo-Pindolol
* Tab 15 mg	33.31	100	✓ Apo-Pindolol
PROPRANOLOL			
* Tab 10 mg		100	✓ Apo-Propranolol
* Tab 40 mg		100	✓ Apo-Propranolol
* Cap long-acting 160 mg		100	✓ Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 below			
Retail pharmacy	CBS	500 m	
			Propranolol S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

continued...

2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTAL OL

*	Tab 80 mg	.32.58	500	✓ Mylan
*	Tab 160 mg	.10.98	100	Mylan

TIMOLOL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking timolol prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol.

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE		
Tab 2.5 mg1.08	90	✓ Vasorex
1.72	100	✓ Apo-Amlodipine
16.20	28	✓ Bristol \$29
Vasorex to be Sole Supply on 1 June 2021		
Tab 5 mg	90	✓ Vasorex
1.56	28	✓ Sandoz S29
		✓ Teva S29
3.33	250	✓ Apo-Amlodipine
Vasorex to be Sole Supply on 1 June 2021		
Tab 10 mg1.19	90	✓ Vasorex
1.66	28	✓ Sandoz S29
4.40	250	✓ Apo-Amlodipine
Vasorex to be Sole Supply on 1 June 2021		
(Apo-Amlodipine Tab 2.5 mg to be delisted 1 June 2021)		
(Apo-Amlodipine Tab 5 mg to be delisted 1 June 2021)		
(Apo-Amlodipine Tab 10 mg to be delisted 1 June 2021)		
FELODIPINE		
	30	✓ Plendil ER
	90	✓ Felo 5 ER
★ Tab long-acting 5 mg 3.93 ★ Tab long-acting 10 mg 4.32	90	✓ Felo 3 ER
78 Tab long-acting to my4.32	90	FEID IU EN

CARDIOVASCULAR SYSTEM				
	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer	
NIFEDIPINE				
* Tab long-acting 10 mg	10.63	60	✓ Adalat 10 ✓ Adefin \$29	
	18.80	56	✓ Adeim \$29	
	10.00	50	• Tellsipille win 10 329	
* Tab long-acting 20 mg		100	✓ Nyefax Retard	
* Tab long-acting 30 mg		30	✓ Adalat Oros	
* Tob long acting 60 mg	34.10	100 30	✓ Mylan S29 ✓ Adalat Oros	
* Tab long-acting 60 mg		30	✓ Adefin XL	
	52.81	100	✓ Mylan S29	
(Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021) (Adefin 329 Tab long-acting 10 mg to be delisted 1 August 2021) (Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021) (Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) (Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021))		·	
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
* Tab 30 mg		100	✓ Dilzem	
* Tab 60 mg * Cap long-acting 120 mg		100 500	✓ Dilzem✓ Apo-Diltiazem CD	
* Cap long-acting 120 mg		500	✓ Apo-Diltiazem CD	
* Cap long-acting 240 mg		500	✓ Apo-Diltiazem CD	
(Dilzem Tab 30 mg to be delisted 1 June 2021)				
(Dilzem Tab 60 mg to be delisted 1 January 2022)				
PERHEXILINE MALEATE				
* Tab 100 mg	62.90	100	✓ Pexsig	
VERAPAMIL HYDROCHLORIDE	7.01	100		
* Tab 40 mg * Tab 80 mg		100 100	✓ Isoptin✓ Isoptin	
* Tab long-acting 120 mg		100	✓ Isoptin Retard \$29	
			✓ Isoptin SR	
* Tab long-acting 240 mg	15.12	30	✓ Isoptin SR	
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a		_		
PSO	25.00	5	✓ Isoptin	
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day - Only on a prescription		4	✓ <u>Mylan</u>	
* Patch 5 mg, 200 mcg per day – Only on a prescription		4	✓ <u>Mylan</u>	
* Patch 7.5 mg, 300 mcg per day – Only on a prescription	10.93	4	✓ <u>Mylan</u>	
CLONIDINE HYDROCHLORIDE * Tab 25 mcg	0.75	112	✓ Clonidine BNM	
* Tab 25 mcg* * Tab 150 mcg		100	✓ Condine blviii ✓ Catapres	
* Inj 150 mcg per ml, 1 ml ampoule		10	✓ <u>Medsurge</u>	
•				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
METHYLDOPA * Tab 250 mg	15.10 52.85	100 500		Methyldopa Mylan Methyldopa Mylan S29 S29
Diuretics				
Loop Diuretics				
# Inj 500 mcg per ml, 4 ml vial	16.36 7.95 7.24 25.00 11.20 60.65	30 100 5 1,000 50 80 ml C 6 5		Burinex S29 S29 Burinex Burinex Apo-Furosemide Urex Forte Lasix Lasix Furosemide-Baxter
Potassium Sparing Diuretics				
AMILORIDE HYDROCHLORIDE Oral liq 1 mg per ml EPLERENONE – Special Authority see SA1728 below – Retail Tab 50 mg	pharmacy	25 ml C 30		Biomed Inspra
Tab 25 mg		30		Inspra Inspra
■ SA1728 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals val the following criteria: Both: 1 Patient has heart failure with ejection fraction less than 4: 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolar 2.2 Patient has experienced a clinically significant adv	0%; and ctone; or			
METOLAZONE	000			
Tab 5 mg	CBS	1 50	_	Metolazone S29 Zaroxolyn S29
SPIRONOLACTONE * Tab 25 mg	4.38	100	1	Spiractin

ıb 5 mg	.CBS	1	✓ Metolazone S29
		50	✓ Zaroxolyn S29
NOLACTONE			
ıb 25 mg	4.38	100	✓ Spiractin
		100	✓ Spiractin
		25 ml OP	✓ Biomed
)	NOLACTONE b 25 mgb 100 mg	b 25 mg	50 NOLACTONE b 25 mg

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE * Tab 5 mg with furosemide 40 mg	8.63	28	✓ Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE * Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ Moduretic

	Subsidy (Manufacturer's Pric	۵۱	Fully Subsidised	Brand or Generic
	\$	Per	√	Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
* Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than eme	ergency.			
* Tab 5 mg	34.55	500	✓	Arrow- Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	26.00	25 ml C	P 🗸	Biomed
CHLORTALIDONE [CHLORTHALIDONE]				
Tab 25 mg		30		Igroton \$29
INDADAMIDE	6.50	50	•	<u>Hygroton</u>
INDAPAMIDE ★ Tab 2.5 mg	10.45	90	1	<u>Dapa-Tabs</u>
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE			_	
* Tab long acting 400 mg		90 30		Bezalip Bezalip Retard
* Tab long-acting 400 mg	12.09	30	<u> </u>	bezalip netaru
Other Lipid-Modifying Agents				
ACIPIMOX	04.50			.
* Cap 250 mg	21.56	30		Olbetam Olbetam S29 S29
NICOTINIC ACID			•	Olbetaili 529 529
Tab 50 mg	4.12	100	1	Apo-Nicotinic Acid
Tab 500 mg		100		Apo-Nicotinic Acid
(Apo-Nicotinic Acid Tab 50 mg to be delisted 1 May 2021) (Apo-Nicotinic Acid Tab 500 mg to be delisted 1 May 2021)				
Resins				
COLESTIPOL HYDROCHLORIDE				
Grans for oral liq 5 g	32.89	30	✓	Colestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg		500		Lorstat
* Tab 20 mg		500		<u>Lorstat</u>
* Tab 40 mg* Tab 80 mg		500 500		<u>Lorstat</u> Lorstat
Tab ov my		500	•	<u>= 5tut</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PRAVASTATIN * Tab 10 mg * Tab 20 mg		28 28 100	√ P	ravastatin Mylan ravastatin Mylan po-Pravastatin
Pravastatin Mylan to be Sole Supply on 1 April 2021 * Tab 40 mg Pravastatin Mylan to be Sole Supply on 1 April 2021 (Pravastatin Mylan Tab 10 mg to be delisted 1 April 2021) (Apo-Pravastatin Tab 20 mg to be delisted 1 April 2021) (Apo-Pravastatin Tab 40 mg to be delisted 1 April 2021)		28 100	√ P	ravastatin Mylan po-Pravastatin
SIMVASTATIN * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 80 mg	2.03 3.58	90 90 90 90	✓ <u>S</u>	imvastatin Mylan imvastatin Mylan imvastatin Mylan imvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see SA1045 below – Retail	pharmacy	
* Tab 10 mg	1.95 30	Ezetimibe Sandoz

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

		p	
Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

Subsidy			Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump
				Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISC	DSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Tab long-acting 60 mg		90	✓ Duride

Sympathomimetics

ADRENALINE

Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

Vasodilators

HYDRAI AZINE HYDROCHI ORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail		
	pharmacyCBS	1	Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM S29
		100	✓ Onelink S29
*	Inj 20 mg ampoule25.90	5	✓ Apresoline

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
MINOXIDIL A Tab 10 mg	70.00	100	1	Loniten
NICORANDIL ▲ Tab 10 mg ▲ Tab 20 mg		60 60		<u>Ikorel</u> Ikorel
PAPAVERINE HYDROCHLORIDE * Inj 12 mg per ml, 10 ml ampoule	217.90	5	•	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg	42.26	50	✓	Trental 400
Endothelin Receptor Antagonists				

AMBRISENTAN - Special Authority see SA1702 below - Retail pharmacy		
Brand switch fee payable (Pharmacode 2605309) - see page 237 for details		
Tab 5 mg1,550.00	30	 Ambrisentan Mylan
Tab 10 mg1,550.00	30	✓ <u>Ambrisentan Mylan</u>

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1991 below - Retail pharmacy Tab 62.5 mg141.00 Tab 125 mg141.00

✓ Bosentan Dr. Reddy's

✓ Bosentan Dr Reddv's

60

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1. 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Fither:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy: and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail pharma	су		
Tab 25 mg	0.64	4	✓ Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg		12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*: and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dvn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail	il pharmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri
⇒SA1696 Special Authority for Subsidy			
Special Authority approved by the Pulmonary Arterial Hypertensio	n Panel		
Notes: Application details may be obtained from PHARMAC's we	bsite schedule.pl	harmac.gov	t.nz/SAForms or:
The Coordinator, PAH Panel			
PHARMAC, PO Box 10-254, WELLINGTON			
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.	govt.nz		
ILOPROST - Special Authority see SA1705 below - Retail pharm	nacy		
Nebuliser soln 10 mcg per ml, 2 ml	,	30	✓ Ventavis
⇒SA1705 Special Authority for Subsidy			
Special Authority approved by the Bulmanany Arterial Hypertensia	n Danal		

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC. PO Box 10-254. WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - F	Retail pharmacy	· ·	
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	✓ Oratane
Cap 20 mg	20.49	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription......13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 93

HYDROGEN PEROXIDE

DERMATOLOGICALS

	Subsidy)	Fully	Brand or
	(Manufacturer's F \$	rice) Subs Per	idised •	Generic Manufacturer
MUPIROCIN				
Oint 2%		15 g OP	_	
	(10.50)		В	actroban
a) Only on a prescription b) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%	1.59	5 g OP	√ F	oban
a) Maximum of 5 g per prescription		- 3 -	_	
b) Only on a prescription				
c) Not in combination	1.50	r - OD		aha
Oint 2%	1.59	5 g OP	V <u>r</u>	<u>oban</u>
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%	10.80	50 g OP	√ F	lamazine
a) Up to 250 g available on a PSO				
b) Not in combination				
Antifungals Topical				
AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14 93	5 ml OP	✓ M	lycoNail
CICLOPIROX OLAMINE		0 1111 01	• •	<u>iyoonuu</u>
a) Only on a prescription				
b) Not in combination				
Nail-soln 8%	5.72	7 ml OP	✓ A	po-Ciclopirox
CLOTRIMAZOLE	2 ==	00 - 00		
* Crm 1%	0.77	on a UD	<i>-</i>	
		20 g OP	•	lomazol
a) Only on a prescription		20 y Oi		ilomazol
a) Only on a prescriptionb) Not in combination		20 g Ol		iomazoi
a) Only on a prescriptionb) Not in combination		•		canesten
a) Only on a prescription b) Not in combination Soln 1%	4.36	•		
a) Only on a prescription b) Not in combination Soln 1%	4.36	•		
a) Only on a prescription b) Not in combination Soln 1%	4.36 (7.55)	20 ml OP		
a) Only on a prescription b) Not in combination Soln 1%	4.36 (7.55)	•	C	
a) Only on a prescription b) Not in combination Soln 1%	4.36 (7.55)	20 ml OP	C	anesten
a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination ECONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination b) Not in combination	4.36 (7.55) 1.00 (7.48)	20 ml OP 20 g OP	C	anesten
a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination ECONAZOLE NITRATE Crm 1% a) Only on a prescription	4.36 (7.55) 1.00 (7.48)	20 ml OP	C P	anesten Yevaryl
a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination ECONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination b) Not in combination	4.36 (7.55) 1.00 (7.48)	20 ml OP 20 g OP	C P	anesten

	L	DERIVIA I OLOGICALS	
	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic ✓ Manufacturer
MICONAZOLE NITRATE			
* Crm 2%	0.81	15 g OP	✓ <u>Multichem</u>
a) Only on a prescriptionb) Not in combination			
* Lotn 2%	4.36	30 ml OP	
	(10.03)		Daktarin
a) Only on a prescription			
b) Not in combination			
* Tinct 2%	4.36 (12.10)	30 ml OP	Daktarin
a) Only on a prescriptionb) Not in combination			
Antipruritic Preparations			
CALAMINE			
a) Only on a prescription b) Not in combination Crm, aqueous, BP	1.26	100 g	✓ <u>healthE Calamine</u> <u>Aqueous Cream</u>
			BP

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%......3.29

20 g OP

✓ Itch-Soothe

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Crystals	6.92	25 g	✓ MidWest
	29.60	100 g	✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 83

Corticosteroids - Plain

BE	TAMETHASONE DIPROPIONATE		
	Crm 0.05%2.9	6 15 g OP	✓ Diprosone
	36.0	0 50 g OP	✓ Diprosone
	Oint 0.05%2.9	6 15 g OP	✓ Diprosone
	36.0	0 50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base4.3	30 g OP	Diprosone OV
BE	TAMETHASONE VALERATE		
*	Crm 0.1%3.4	5 50 g OP	✓ Beta Cream
	Oint 0.1%3.4		✓ Beta Ointment
*	Lotn 0.1%	0 50 ml OP	✓ Betnovate
CL	OBETASOL PROPIONATE		
*	Crm 0.05%2.1	8 30 g OP	✓ Dermol
*	Oint 0.05%2.1	2 30 g OP	✓ Dermol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's P		sidised	Generic
	\$	Per		Manufacturer
LOBETASONE BUTYRATE				
Crm 0.05%		30 g OP		
	(10.00)		E	Eumovate
IFLUCORTOLONE VALERATE				
Fatty oint 0.1%	8.97	50 g OP		
•	(15.86)	ŭ	N	Verisone
Nerisone Fatty oint 0.1% to be delisted 1 August 2021)	, ,			
YDROCORTISONE				
Crm 1% – Only on a prescription	3.70	100 g OP	✓	lydrocortisone
		9	_	(PSM)
	17.15	500 g	✓ F	lydrocortisone
		000 g		(PSM)
Powder – Only in combination	49 95	25 g	1	ABM
Up to 5% in a dermatological base (not proprietary Topic				
galenicals	oai contioostorioa	r iairi, wiar	JI WILLIO	at other definationogies
<u> </u>				
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only		0501		ND 1 -4 110
a prescription	10.57	250 ml	• [OP Lotn HC
YDROCORTISONE BUTYRATE			_	
Lipocream 0.1%		100 g OP		ocoid Lipocream
Oint 0.1%		100 g OP	_	ocoid.
Milky emul 0.1%	13.70	100 ml OP	✓ <u>L</u>	ocoid Crelo
ETHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	√ <u> </u>	<u>Advantan</u>
Oint 0.1%	4.46	15 g OP	√ <u>I</u>	<u>Advantan</u>
IOMETASONE FUROATE				
Crm 0.1%	1.51	15 g OP	✓ E	locon Alcohol Free
	2.50	50 g OP	✓ E	locon Alcohol Free
Oint 0.1%	1.51	15 g OP	✓ E	locon
	2.90	50 g OP	✓	locon
Lotn 0.1%	6.30	30 ml OP	✓ E	locon
RIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	✓ I	Aristocort
Oint 0.02%		100 g OP		Aristocort
Corticosteroids - Combination				
ETAMETUA CONE VALEDATE MUTU CUO CUINO CON CO	a properintian			
ETAMETHASONE VALERATE WITH CLIOQUINOL — Only on Crm 0.1% with clioquinol 3%		15 g OP		
OIIII 0. 1 /0 WILLI GIIOQUILIOI 3 /0	(4.90)	15 y OF	_	Betnovate-C
Betnovate-C Crm 0.1% with clioquinol 3% to be delisted 1 June				Join 0 Valo-0
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU		4E = OD		
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
A Martinum of 45 may	(10.45)		ŀ	ucicort
a) Maximum of 15 g per prescription				
b) Only on a prescription				
	ption			

	Subsidy (Manufacturer's P \$	Price) Subsi Per	Fully Brand or idised Generic Manufacturer
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	3.35 3.35 CIN AND NYSTAT	15 g OP 15 g OP	✓ Pimafucort ✓ Pimafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 r and gramicidin 250 mcg per g - Only on a prescription	•	15 g OP	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 10%</u>
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm	1.92	500 g	✓ Basic AquaCream ✓ Boucher
CETOMACROGOL	5.75		✓ Topiderm
* Crm BP	2.48	500 g	✓ <u>healthE</u>
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ Boucher
	3.10	1,000 ml OP	✓ Kenkay Sorbolene✓ ADE✓ Boucher
EMULSIFYING OINTMENT * Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION * Crm	2.19	500 g	✓ <u>O/W Fatty Emulsion</u> <u>Cream</u>
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50% UREA	5.35	500 ml OP	✓ <u>healthE</u>
* Crm 10%	1.37	100 g OP	✓ healthE Urea Cream

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

	Subsidy (Manufacturer's P \$		Fully dised	Brand or Generic Manufacturer
WOOL FAT WITH MINERAL OIL - Only on a prescription				
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
•	(11.95)			P Lotion
	1.40	250 ml OP		
	(4.53)			P Lotion
	5.60	1,000 ml		
	(20.53)		Α	Ipha-Keri Lotion
	(23.91)		В	K Lotion
	1.40	250 ml OP		
	(7.73)		В	SK Lotion
OIL D				

Other Dermatological Bases

PARAFFIN

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIN			

		Lotion	
IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy			
Tab 3 mg - Up to 100 tab available on a PSO 17.20	4	✓ Stromectol	

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

200 ml OP

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

healthE

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical mi

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or



Subsidy	Fully	Brand or
(Manufacturer's	s Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%	0 -	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN Shamooo 0.5%	200 ml OP	✓ Parasidasa
Sharipoo 0.5%11.36	200 m OP	 Parasidose

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see SA2024 below – Retail pharmacy		
Cap 10 mg17.86	60	✓ Novatretin
Cap 25 mg41.36	60	✓ <u>Novatretin</u>

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment;
- 2 Patient is not of child bearing potential.

RETAMETHASONE DIDRODIONATE WITH CALCIDOTRIOL

BETAMETHASONE DIFNOFICINATE WITH CALCIFOTHIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	52.24	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP	Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ <u>Midwest</u>

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

		•		ATOLOGICALO
	Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully sidised	Brand or Generic Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP	HUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%		75 g OP 30 g OP		Egopsoryl TA Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP		Coco-Scalp Coco-Scalp
PIMECROLIMUS – Special Authority see SA1970 below – Retail a) Maximum of 15 g per prescription b) Note: a maximum of 15 g per prescription and no more th Cream 1%	an one prescripti	15 g OP	√ <u>E</u>	<u>Elidel</u>
Initial application only from a dermatologist, paediatrician, ophtho of a dermatologist, paediatrician or ophthalmologist. Approvals vameeting the following criteria: Both:				
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications to documented epidermal atrophy, documented allergy to top pressure. 				
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.		a prescriptio 500 ml		Pinetarsol Pinetarsol
SALICYLIC ACID Powder – Only in combination	18.88	250 g		Midwest PSM
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	proprietary Topica	al Corticoster	oid – Pl	ain or collodion flexible
SULPHUR				
Precipitated – Only in combination	6.35	100 g	✓ I	Midwest

Scalp Preparations

BETAMETHASONE VALERATE		
* Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	100 ml OP	✓ <u>Locoid</u>
KETOCONAZOLE Shampoo 2%	100 ml OP	✓ <u>Sebizole</u>

1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain

2) With or without other dermatological galenicals.

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

DERMATOLOGICALS

Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

✓ Marine Blue Lotion SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 72

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	_
(Manufacturer's Price) \$	Per	Subsidised ✓	Generic Manufacturer	
· · · · · · · · · · · · · · · · · · ·				_

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic

Manufacturer

Contraceptives - Non-hormonal

Condoms

CO	NDOMS			
*	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
*	53 mm	0.95	10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
K	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
		11.42	144	✓ Moments
	 a) Up to 60 dev available on a PSO 			
	b) Maximum of 60 dev per prescription			
ĸ	53 mm, chocolate, brown	0.95	10	✓ <u>Moments</u>
		11.64	144	✓ <u>Moments</u>
	 a) Up to 60 dev available on a PSO 			
	 b) Maximum of 60 dev per prescription 			
K	53 mm, strawberry, red		10	✓ Moments
		11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
K	56 mm		10	✓ <u>Moments</u>
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			4.6.114.1.1
K	56 mm, 0.05 mm thickness		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	44.04		40.114.114
F	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO	0.07	40	/ U
K	56 mm, 0.08 mm thickness		10	✓ Moments
) II + 00 I	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
Ŀ	b) Maximum of 60 dev per prescription	0.07	10	√ Mamarta
•	56 mm, 0.08 mm thickness, red	0.97 11.64	10 144	✓ <u>Moments</u>✓ Moments
	a) Un to 60 day available as a DCO	11.04	144	* INIOINICITES
	a) Up to 60 dev available on a PSO b) Maximum of 60 dev per prescription			
k-	56 mm, chocolate	1 20	12	✓ Gold Knight
N.	JU IIIII, GIUGUIAIC	15.57	12 144	✓ Gold Knight
	a) Up to 60 dev available on a PSO	10.07	144	- Gold Killgill
	b) Maximum of 60 dev per prescription			
K	56 mm, strawberry	1 20	12	✓ Gold Knight
	50 mm, snawpeny	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO	10.01	177	- doid Killylit
	b) Maximum of 60 dev per prescription			
ĸ	60 mm	1 49	12	✓ Gold Knight XL
,-	VV 11111111111111111111111111111111111	14.87	144	✓ Shield XL
		17.02		✓ Gold Knight XL

a) Maximumosidisedev per prescription b) substantification a PSO

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
*	60 mm (bulk pack)	14.87	144	1	Gold Knight XL
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

	b) Only on a PSO			
*	IUD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓ Choice
	-			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to)		
	84 tab available on a PSO	19.80	84	Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	6.62	84	
		(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	✓	Microgynon 20 ED
	6.45	112	✓	Femme-Tab ED
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - U	p			
to 84 tab available on a PSO	9.45	84	✓	Microgynon 50 ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	hority see SA0500 on	the	orevious p	age
b) Up to 63 tab available on a PSO				
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	1.77	84	✓	Levlen ED
	6.45	112	✓	Femme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to)			
84 tab available on a PSO		84	✓	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - L	J p			
to 84 tab available on a PSO	6.62	84	✓	Necon
	8.29		•	Norimin

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO	16.50 22.00	84 112	✓ <u>Microlut</u> ✓ <u>Microlut</u>
*	Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO	.106.92	1	✓ Jadelle

Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer	
SO7.98 6.25	1 84	_		
4.95	1 Part Lof S	·		
	(Manufacturer's Price) \$ SO7.986.25	(Manufacturer's Price) Substitute Subs	(Manufacturer's Price) Subsidised Per \$ 1 ✓ □ ✓ □ 6.25 84 ✓ № 4.95 1 ✓ Р	(Manufacturer's Price) \$ Subsidised Generic Manufacturer SO7.98 1 ✓ Depo-Provera 6.25 84 ✓ Noriday 28

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

*	Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs - Up			
	to 168 tab available on a PSO	4.98	168	Ginet
	Ginet to be Sole Supply on 1 April 2021			

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE		Acroci
* Vaginal crm 1% with applicators	35 g OP	✓ <u>Clomazol</u>
* Vaginal crm 2% with applicators3.00 MICONAZOLE NITRATE	20 g OP	✓ <u>Clomazol</u>
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
PSO	160.00	5	DBL Ergometrine
OESTRIOL			
* Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
* Pessaries 500 mcg	6.86	15	✓ Ovestin
OXYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	 Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	✓ Oxytocin BNM

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Subs	Fully sidised	Brand or Generic Manufacturer	
OXYTOCIN WITH ERGOMETRINE MALEATE — Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	✓ <u>s</u> y	/ntometrine	

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

40 test OP ✓ David One Step Cassette

> **Pregnancy Test** ✓ Smith BioMed Rapid **Pregnancy Test**

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 110

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy ✓ Ricit * Tab 5 mg4.81 100 Ricit to be Sole Supply on 1 April 2021

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Fither:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy ✓ Tamsulosin-Rex

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN * Tab 5 mg	500 473 ml	✓ Apo-Oxybutynin ✓ Apo-Oxybutynin
POTASSIUM CITRATE Oral lig 3 mmol per ml - Special Authority see SA1083 on the		
next page – Retail pharmacy31.80	200 ml OP	✓ Biomed

GENITO-URINARY SYSTEM

Albustix

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	•	Manufacturer

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

* Grans eff 4 g sachets	28	✓ Ural
SOLIFENACIN SUCCINATE		
Tab 5 mg3.00	30	✓ Solifenacin Mylan
Tab 10 mg5.50	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORT	HO-	LOI I	IDIN	F

UΠ	THO-TOLIDINE			
*	Compound diagnostic sticks	.7.50	50 test OP	
	, ,	(8.25)		Hemastix
TE	FRABROMOPHENOL			
*	Blue diagnostic strips	.7.02	100 test OP	

Obstetric Preparations

Antiprogesterones

MIFFPRISTONE

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

(13.92)

✓ Mifegyne	. 1	Tab 200 mg60.00
✓ Mifegyne	3	180.00

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	1	Manufacturer

Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA1618 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	210.30	28	✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g., painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial - Special Authority see SA1687 below -✓ Zoledronic acid Mylan

⇒SA1687 Special Authority for Subsidy

Initial application — (bone metastases) only from an oncologist, haematologist or palliative care specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) only from an oncologist or medical practitioner on the recommendation of a oncologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
 - 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
 - 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

	IAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATI Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5 5	
	(36.96)		Celestone
DE	VAMETUACONE		Chronodose
*	XAMETHASONE Tab 0.5 mg - Up to 60 tab available on a PSO	30	✓ Dexmethsone
	Tab 4 mg — Up to 30 tab available on a PSO	30	✓ Dexmethsone
	Oral liq 1 mg per ml45.00	25 ml OP	✓ Biomed
DE	XAMETHASONE PHOSPHATE		
	Dexamethasone phosphate injection will not be funded for oral use.		
*	Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
*	Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
	JDROCORTISONE ACETATE		
*	Tab 100 mcg14.32	100	✓ Florinef
	DROCORTISONE		
	Tab 5 mg	100	✓ <u>Douglas</u>
	Tab 20 mg	100	 ✓ <u>Douglas</u> ✓ Solu-Cortef
不	Inj 100 mg vial	1	▼ Solu-Cortei
	a) Up to 5 inj available on a PSO b) Only on a PSO		
ME	THYLPREDNISOLONE		
	Tab 4 mg	100	✓ Medrol
	Tab 100 mg	20	✓ Medrol
	THYLPREDNISOLONE (AS SODIUM SUCCINATE)	_0	<u></u>
IVIL	Inj 40 mg vial18.90	1	✓ Solu-Medrol-Act-
	11, 10 mg 11d	•	O-Vial
	Inj 125 mg vial	1	✓ <u>Solu-Medrol-Act-</u> <u>O-Vial</u>
	Inj 500 mg vial	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
	Inj 1 g vial27.83	1	✓ Solu-Medrol
ME	THYLPREDNISOLONE ACETATE		- Join monto
IVIL	Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol
DD		J	- Depo-Medioi
	EDNISOLONE Oral lig 5 mg per ml – Up to 30 ml available on a PSO6.00	30 ml OP	✓ Redipred
~	Restricted to children under 12 years of age.	JU IIII UF	• <u>Ileuipieu</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

^{*}Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
	\$	Per		Manufacturer
PREDNISONE				
★ Tab 1 mg	10.68	500	✓	Apo-Prednisone
★ Tab 2.5 mg		500	✓	Apo-Prednisone
Fab 5 mg - Up to 30 tab available on a PSO	11.09	500	✓	Apo-Prednisone
★ Tab 20 mg – Up to 30 tab available on a PSO	29.03	500	✓	Apo-Prednisone
ETRACOSACTRIN				
	75.00	1	/	UK Synacthen S29
, , , ,				AU Synacthen
				Synacthen
Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Depot
, 31			1	Synacthene
				Retard S29
RIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓	Kenacort-A 10
	26.62		/	Adcortyl S29
Kenacort-A 10 to be Sole Supply on 1 April 2021				
Inj 40 mg per ml, 1 ml ampoule	11.30	1	/	Triaver \$29
,,	51.10	5	/	Kenacort-A 40
	70.62	-	1	Kenalog \$29
Kenacort-A 40 to be Sole Supply on 1 April 2021	7 3.02			Tronding

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CYPROTERONE ACETATE		
Tab 50 mg13.17	50	✓ Siterone
Tab 100 mg26.75	50	✓ Siterone
TESTOSTERONE		
Patch 5 mg per day90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE		
Inj 100 mg per ml, 10 ml vial85.00	1	✓ Depo-Testosterone
TESTOSTERONE ESTERS		
Inj 250 mg per ml, 1 ml12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE		
Cap 40 mg21.00	60	Andriol Testocaps
Inj 250 mg per ml, 4 ml vial86.00	1	✓ Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy (Manufacturer's Price) \$) Su Per	Fully bsidised	Brand or Generic Manufacturer
Oestrogens				
DESTRADIOL - See prescribing guideline on the previous page				
F Tab 1 mg		28 OP		
ŭ	(11.10)			Estrofem
₭ Tab 2 mg	4.12 [′]	28 OP		
Ç	(11.10)			Estrofem
Patch 100 mcg per 24 hours	7.91 [′]	4	1	Climara
a) No more than 1 patch per week				
b) Only on a prescription				
Fatch 50 mcg per 24 hours	7.04	4	1	Climara
a) No more than 1 patch per week				
b) Only on a prescription				
Patch 25 mcg per day	6.12	8	/	Estradot
01	7.85		/	Estradiol TDP
				Mylan S29
a) No more than 2 patch per week				,
b) Only on a prescription				
Patch 50 mcg per day	7 04	8	1	Estradot 50 mcg
Tator of may per day	9.22	Ŭ		Estradiol TDP
	V.LL			Mylan S29
a) No more than 2 patch per week				Wylanozo
b) Only on a prescription				
Patch 75 mcg per day	7 01	8	1	Estradot
1 atom 70 mag per day	10.60	U		Estradiol TDP
	10.00		•	Mylan S29
a) No more than 2 notes per week				Wylair 323
a) No more than 2 patch per weekb) Only on a prescription				
Patch 100 mcg per day	7.01	8	1	Estradot
3 1 ,	7.91	0	•	EStrauot
a) No more than 2 patch per weekb) Only on a prescription				
ESTRADIOL VALERATE – See prescribing guideline on the pr			_	_
F Tab 1 mg		84	_	<u>Progynova</u>
F Tab 2 mg	12.36	84	•	<u>Progynova</u>
ESTROGENS - See prescribing guideline on the previous page				
Conjugated, equine tab 300 mcg	3.01	28		
	(17.50)			Premarin
Conjugated, equine tab 625 mcg		28		
	(17.50)			Premarin
Progestogens				
IEDROXYPROGESTERONE ACETATE - See prescribing guid	laling on the previou	e nage		
← Tab 2.5 mg		30	/	Provera
€ Tab 5 mg		100		Provera
₭ Tab 10 mg		30		Provera
· 100 10 Hig	0.34	50	•	1 10VC10

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$	e) S Per	Fully Brand or Subsidised Generic ✓ Manufacturer					
Progestogen and Oestrogen Combined Preparations								
OESTRADIOL WITH NORETHISTERONE - See prescribing gr	uideline on page 84							
* Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP						
	(18.10)		Kliovance					
* Tab 2 mg with 1 mg norethisterone acetate		28 OP						
	(18.10)		Kliogest					
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	5 40	00 OD						
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Trianguage					
	(18.10)		Trisequens					
Other Oestrogen Preparations								
ETHINYLOESTRADIOL								
* Tab 10 mcg	17.60	100	✓ <u>NZ Medical and</u> Scientific					
OESTRIOL								
* Tab 2 mg	7.00	30	✓ Ovestin					
			<u></u>					
Other Progestogen Preparations								
LEVONORGESTREL								
* Intra-uterine device 52 mg	269.50	1	✓ Mirena					
* Intra-uterine device 13.5 mg	215.60	1	Jaydess					
MEDROXYPROGESTERONE ACETATE								
Tab 100 mg	116.15	100	✓ Provera HD					
NORETHISTERONE								
* Tab 5 mg - Up to 30 tab available on a PSO	18.29	100	✓ Primolut N					
PROGESTERONE								
Cap 100 mg - Special Authority see SA1609 below - Retail	il							
cap rooms openial realiting decontroop below riotal			4					

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

pharmacy......16.50

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

30

✓ Utrogestan

_		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
ī	hyroid and Antithyroid Agents				
CA	RBIMAZOLE				
*	Tab 5 mg	10.80	100		Neo-Mercazole Neo-Mercazole S29 S29
ΙF	VOTHYROXINE				
*	Tab 25 mcg	5.55	90	1	Synthroid
*	Tab 50 mcg		28		Mercury Pharma
	•	5.79	90	✓	Synthroid
		64.28	1,000) /	Eltroxin
*	Tab 100 mcg	1.78	28	✓	Mercury Pharma
		6.01	90	✓	Synthroid
		66.78	1,000) /	Eltroxin
PR	OPYLTHIOURACIL – Special Authority see SA1199 below – Propylthiouracil is not recommended for patients under the a treatments are contraindicated.		the	patient is p	regnant and other
	Tab 50 mg	35.00	100	✓	PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA1629 below	/ – Retail pha	rmacy	
*	Inj 5 mg cartridge	34.88	1	 Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	104.63	1	✓ Omnitrope

⇒SA1629 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

Subsidy	•	Fully	Brand or
(Manufacturer's Price)		osidised	Generic
\$	Per		Manutacturer

continued...

children who are 5 years or older, GH testing with sex steroid priming is required; and

- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) x 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manutacturer	

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and

Subsidy (Manufactured Price)		Fully ubsidised	Brand or
(Manufacturer's Price) \$	Per	ubsidised •	Generic Manufacturer

continued...

G

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
1 3, 7 3	66.48		✓ Zoladex
Teva to be Sole Supply on 1 May 2021			
Implant 10.8 mg, syringe	122.37	1	✓ Teva
, , , ,	177 50		✓ Zoladex

Teva to be Sole Supply on 1 May 2021

(Zoladex Implant 3.6 mg, syringe to be delisted 1 May 2021)

(Zoladex Implant 10.8 mg, syringe to be delisted 1 May 2021)

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Denot 3-month

Vasopressin Agonists

DESMOPRESSIN Wafer 120 mcg	47.00	30	✓ Minirin Melt
DESMOPRESSIN ACETATE			
Tab 100 mcg	25.00	30	✓ Minirin
Tab 200 mcg	54.45	30	✓ Minirin
▲ Nasal drops 100 mcg per ml		2.5 ml OP	✓ Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ <u>Desmopressin-</u> <u>PH&T</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓ Minirin

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
✓ Dostinex	2	waived by Special Authority see SA1370 on the next page3.75
✓ Dostinex	8	15.20

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

\sim 1 $^{\prime}$	~ A A	IFEN	ı	\neg ı \pm	\square	

Tab 50 mg	29.84	10	Mylan
			Clomiphen S29
DANAZOL			
Cap 100 mg	19.13	28	✓ Mylan S29
Cap 200 mg	97.83	100	✓ Azol
(Mylan S29 Cap 100 mg to be delisted 1 April 2021)			
(Azol Cap 200 mg to be delisted 1 April 2021)			
METYRAPONE			
Cap 250 mg	558.00	50	✓ <u>Metopirone</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 ✓ Eskazole S29 **⇒SA1318** Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription Tab 100 mg7.97 6 ✓ Vermox 15 ml (7.53)Vermox **PRAZIQUANTEL** Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 65
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 232

Cephalosporins and Cephamycins			
CEFACLOR MONOHYDRATE			
Cap 250 mg		100	✓ Ranbaxy-Cefactor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefactor
CEFALEXIN			
Cap 250 mg	3.33	20	✓ Cephalexin ABM
Cap 500 mg		20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	8.75	100 ml	✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable	11.75	100 ml	✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with a Di accordingly.	HB approved p	rotocol and t	he prescription is endorsed
Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Inj 1 g vial	3.49	5	✓ <u>AFT</u>
CEFTRIAXONE - Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
 Subsidised only if prescribed for a dialysis or cystic fibrosis p pelvic inflammatory disease, or the treatment of suspected mendorsed accordingly. 			
Inj 500 mg vial	0.80	1	✓ Ceftriaxone-AFT
Inj 1 g vial		5	✓ Ceftriaxone-AFT
, 0		Ü	- Oldinadilo Al I
CEFUROXIME AXETIL — Subsidy by endorsement	ntion io andoro	ad according	dv
Only if prescribed for prophylaxis of endocarditis and the prescri	•	ed according 50	ııy. ✓ Zinnat
Tab 250 mg		50	Lilliat

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

Macrolides

AZITHROMYCIN - Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority

Tab 250 mg	.8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable1	14.38	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*: or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

✓ Apo-Clarithromycin 14 ✓ Klacid Grans for oral lig 250 mg per 5 ml - Wastage claimable......192.00 50 ml

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

1 Atypical mycobacterial infection; or

ERYTHROMYCIN (AS LACTORIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO			•
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
 a) Up to 200 ml available on a PSO 			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg	29.90	100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-
			<u>Roxithromycin</u>
Tab 300 mg	16 33	50	✓ Arrow-
Tab 500 Hig	10.33	50	Roxithromycin
			1 to Aittii Oili yolii

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) Subs	sidised	
	\$	Per	1	Manufacturer
Penicillins				
Penicillins				
AMOXICILLIN			_	
Cap 250 mg	22.50	500	/	<u>Alphamox</u>
 a) Up to 30 cap available on a PSO 				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500	/	<u>Alphamox</u>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	1	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15.97	10	/	Ibiamox
Inj 500 mg vial		10	1	Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	1	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO	0.89	10	1	Curam Duo 500/125
available on a r Go	5.00	20		Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25			-	, tuginonini
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO		100 1111		, tuginonini
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml – Up to 200 ml available on a PSO		I00 ml OP	1	Curam
(Augmentin Tab 500 mg with clavulanic acid 125 mg to be delisted		100 1111 01	•	Outain
	a rodry 2021)			
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj	044.00	4.0	,	D: :::: 1.4
available on a PSO	344.93	10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10	1	Sandoz
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO	16.83	250	/	Staphlex
Cap 500 mg - Up to 30 cap available on a PSO	56.61	500	1	Staphlex
Grans for oral liq 25 mg per ml	2.29	100 ml		AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	1	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial	17.56	10	1	Flucloxin
Inj 500 mg vial	18.87	10	1	Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	1	Flucil

·	Subsidy		Fully	
	(Manufacturer's Price) \$	Subside Per	isea •	Generic Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	2.59	50	1	Cilicaine VK
Cap 500 mg	4.26	50	1	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP			_	
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	/	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable	2.00	100 ml	./	AFT
Grans for oral liq 250 mg per 5 ml		100 1111	٠	<u>AFI</u>
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	1	Cilicaine
, 19 1 1, 31 1, 11 1, 11 11 11				
Tetracyclines				
DOXYCYCLINE				
* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	1	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
	(12.05)			Mino-tabs
* Cap 100 mg		100		
OMOSS On a lat Authority for Manufacturers Date	(52.04)			Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 65

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	
	<u> </u>	Per		Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.61	24	✓	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	•	Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and the			accordingly	y.
Inj 150 mg	65.00	1	✓	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	25.00	5	1	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection a	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	182.00	10	✓	Teligent \$29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection a	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓	Pfizer
	87.50	50	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	t infection :	and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retail No patient co-payment payable	pharmacy			
Tab 400 mg	42.00	5	1	Avelox
⇒SA1740 Special Authority for Subsidy				

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications:
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufactured Price)		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacturer
PAROMOMYCIN - Special Authority see SA1689 below - Retail	pharmacy			
Cap 250 mg		16	✓ Hu	matin \$29
⇒SA1689 Special Authority for Subsidy				
Initial application only from an infectious disease specialist, clini	ical microbiologist or	gastro	enterologist.	Approvals valid for 1
month for applications meeting the following criteria: Either:				
Patient has confirmed cryptosporidium infection; or				
2 For the eradication of Entamoeba histolyica carriage.				
Renewal only from an infectious disease specialist, clinical micro	biologist or gastroent	erolog	gist. Approva	ls valid for 1 month for
applications meeting the following criteria:				
Either:				
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE – Special Authority see SA1328 below – Reta	ail pharmaou			
Tab 25 mg		30	√ Da	raprim \$29
⇒SA1328 Special Authority for Subsidy		00	• 50	партт
Initial application from any relevant practitioner. Approvals valid	d without further rene	wal ur	nless notified	for applications meetin
the following criteria:				
Any of the following:				
1 For the treatment of toxoplasmosis in patients with HIV for2 For pregnant patients for the term of the pregnancy; or	r a period of 3 months	s; or		
3 For infants with congenital toxoplasmosis until 12 months	of age.			
SODIUM FUSIDATE [FUSIDIC ACID]	g			
Tab 250 mg	34.50	12	✓ Fu	cidin
SULFADIAZINE SODIUM - Special Authority see SA1331 below				
Tab 500 mg	543.20	56	✓ Wo	ockhardt S29
⇒ SA1331 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	d without further rene	wal ur	nless notified	for applications meetin
the following criteria: Any of the following:				
For the treatment of toxoplasmosis in patients with HIV for	a period of 3 months	s: or		
2 For pregnant patients for the term of the pregnancy; or		-,		
3 For infants with congenital toxoplasmosis until 12 months	of age.			
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement		5		bramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient an Solution for inhalation 60 mg per ml, 5 ml - Subsidy by	a the prescription is e	endors	sea according	lly.
endorsement	395.00 5	6 dos	e 🗸 To	bramycin BNM
	2,200.00		✓ T0	
a) Wastage claimable				
b) Only if prescribed for a cystic fibrosis patient and the	prescription is endors	sed ac	cordingly.	
c) Tobramycin BNM to be Sole Supply on 1 May 2021 (TOBI Solution for inhalation 60 mg per ml, 5 ml to be delisted 1 I	May 2021)			
TRIMETHOPRIM	,,			
* Tab 300 mg - Up to 30 tab available on a PSO	16.50	50	✓ TN	<u>IP</u>
•				

	Subsidy (Manufacturer's Price \$) Si Per	Fully ubsidised	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
★ Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - l to 30 tab available on a PSO		500	✓ Ti	risul
Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 available on a PSO		100 ml	✓ D	eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is Inj 500 mg vial	endorsed according			ment of Clostridium

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 66
- b) For topical antifungals refer to GENITO URINARY, page 79

FLUCONAZOLE

Cap 50 mg2.75	5 28	✓ Mylan
Cap 150 mg		✓ Mylan
Cap 200 mg12.89		✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy109.34	4 35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 on the			
next page – Retail pharmacy	141.80	150 ml OP	✓ Sporanox

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOI E

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
· ·			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
•	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - R	etail pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml		105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

* Tab 250 mg	8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the next page	e – Retail phar	macy	
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,437.00	70 ml	✓ Vfend

	Subsidy	Fully	Brand or
(Manufa	,	sidised	Generic
	\$ Per	· ·	Manutacturer

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail p	harmacy		
Tab 7.5 mg	117.00	56	✓ Primacin S29
Tab 15 mg	400.00	100	✓ Sanofi
			Primaquine \$29

(Primacin S29 Tab 7.5 mg to be delisted 1 June 2021)

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SULPHATE			
* Tab 300 mg	61.91	500	✓ Q 300
(O 300 Tah 300 mg to be delisted 1, July 2021)			

Q 000 Tab 000 Tig to be delisted Today 2021

INFECTIONS - AGENTS FOR SYSTEMIC USE					
	Subsidy (Manufacturer's Pric \$	e) Sub Per	Fully osidised	Brand or Generic Manufacturer	
Antitrichomonal Agents					
METRONIDAZOLE Tab 200 mg — Up to 30 tab available on a PSO Tab 400 mg — Up to 15 tab available on a PSO Oral liq benzoate 200 mg per 5 ml Suppos 500 mg	5.23 25.00	250 21 100 ml 10	✓ <u>N</u> ✓ F	letrogyl letrogyl lagyl-S lagyl	
ORNIDAZOLE Tab 500 mg		10		rrow-Ornidazole	
Antituberculotics and Antileprotics					
Note: There is no co-payment charge for all pharmaceuticals I immigration status. CLOFAZIMINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend dermatologist.	lation of, an infectious	s disease pl	nysician,	clinical microbiologist or	
* Cap 50 mg	442.00	100	✓ L	amprene S29	
CYCLOSERINE – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician. Cap 250 mg		s disease pl		clinical microbiologist or	
DAPSONE – Retail pharmacy-Specialist				,	
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend dermatologist Tab 25 mg	268.50	100	✓ D	apsone	
Tab 100 mg ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specia		100	• 0	apsone	
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend respiratory physician		s disease pl	nysician,	clinical microbiologist or	
Tab 100 mg	85.73	100	√ E	MB Fatol S29	
Tab 400 mg	49.34	56	✓ N	lyambutol S29	
ISONIAZID – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician		nedicine ph	ysician, p	paediatrician, clinical	
* Tab 100 mg		100	✓ <u>P</u>	<u>SM</u>	
${\sf ISONIAZID\ WITH\ RIFAMPICIN\ -Retail\ pharmacy-Specialist}$					
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommend microbiologist, dermatologist or public health physician 		nedicine ph	ysician, p	paediatrician, clinical	
* Tab 100 mg with rifampicin 150 mg * Tab 150 mg with rifampicin 300 mg	85.54	100 100	_	ifinah ifinah	

	Subsidy (Manufacturar's Price)		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Special	list			
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recomme	ndation of, an infectious d	iseas	e specialist,	clinical microbiologist or
respiratory physician Grans for oral liq 4 g sachet	200.00	30	./ 0	aser S29
	280.00	30	V P	aser 529
PROTIONAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recomme	ndation of an infectious d	iseas	e specialist	clinical microbiologist or
respiratory physician	aa.iioii oi, aii iiiooiioao a		o opoolaliot,	ommout timer obtaining for or
Tab 250 mg	305.00	100	✓ P	eteha S29
PYRAZINAMIDE - Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recomme	ndation of, an infectious d	iseas	e physician,	clinical microbiologist or
respiratory physician * Tab 500 mg	50.00	100	1	AFT-Pyrazinamide
RIFABUTIN – Retail pharmacy-Specialist		100	• •	ii 1-r yrazillallilide
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recomme	ndation of, an infectious d	iseas	e physician,	respiratory physician or
gastroenterologist	, , , , , , , , , , , , , , , , , , , ,		, , , , , ,	, , , , , , , , , , , , , , , , , , ,
* Cap 150 mg	299.75	30	✓ N	lycobutin
RIFAMPICIN – Subsidy by endorsement				
a) No patient co-payment payable				
b) For confirmed recurrent Staphylococcus aureus infe- antimicrobial based on susceptibilities and the presc				1 /
Retail pharmacy - Specialist. Specialist must be an				
paediatrician, or public health physician.	, ,			0 /
* Cap 150 mg		100	_	Rifadin
* Cap 300 mg		100	_	<u>Rifadin</u>
* Oral liq 100 mg per 5 ml	12.00	60 m	₩ <u>F</u>	<u>Rifadin</u>
Antivirals				
For eye preparations refer to Eye Preparations, Anti-Infective	e Preparations, page 232			
Hepatitis B Treatment				
ENTECAVIR				
* Tab 0.5 mg	52.00	30	√ E	intecavir Sandoz
LAMIVUDINE – Special Authority see SA1685 below – Reta				
Tab 100 mg		28	√ Z	etlam
Oral liq 5 mg per ml	270.00 24	0 ml (OP 🗸 Z	effix
⇒SA1685 Special Authority for Subsidy				
Initial application only from a relevant specialist or medical		mend	ation of a re	levant specialist.
Approvals valid for 1 year where used for the treatment or preserved from any relevant practitioner. Approvals valid for		o troc	tmont or nr	yention of honotitic B
TENOFOVIR DISOPROXIL	2 years where used for th	e nec	uneni oi pie	evention of nepatitis b.
Tenofovir disoproxil prescribed under endorsement for t	he treatment of HIV is incl	uded	in the count	of up to 4 subsidised
antiretrovirals for the purposes of Special Authority SA1				I. Sp to . Cabolalood
* Tab 245 mg (300.6 mg as a succinate)		30	√ <u>T</u>	enofovir Disoproxil
				<u>Teva</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Herpesvirus Treatments				
ACICLOVIR				
* Tab dispersible 200 mg	1.60	25	✓ L	ovir
* Tab dispersible 400 mg	5.38	56	✓ L	ovir
* Tab dispersible 800 mg		35	✓ L	ovir
VALACICLOVIR				
Tab 500 mg	5.75	30	✓ V	aclovir
Tab 1,000 mg		30	✓ V	aclovir
VALGANCICLOVIR - Special Authority see SA1993 below - Re	etail pharmacy		_	
Tab 450 mg	' '	60	_	<u>alganciclovir</u> Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis: and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis: and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive: and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient is immunocompromised; and

Subsidy Fully (Manufacturer's Price) Subsidised		Brand or Generic	
 Per	✓	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient is immunocompromised; and
 - 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 ✓ Harvoni

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see SA1994 on the next page

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 108 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

⇒SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or

Subsidy (Manufacturer's Price)	S	Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:

	ubsidy cturer's Price) Su	Fully	Brand or Generic
<u> </u>	\$ Per	✓	Manufacturer

continued...

- 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previo	us page – Retail pharr	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	ous page – Retail phai	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	ous page - Retail pha	rmacy	
Tab 200 mg	60.00	60	✓ <u>Nevirapine</u>
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

Nucleosides Reverse Transcriptase Inhibitors

Tab 300 mg Oral liq 20 mg per ml	180.00	60 240 ml OP	y ✓ <u>Ziagen</u> ✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority ser Note: abacavir with lamivudine (combination tablets) counts as anti-retroviral Special Authority.	two anti-retro		•
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPROX Retail pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil coun anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil	•	•	
245 mg (300 mg as a maleate)	106.88	30	✓ Mylan
EMTRICITABINE – Special Authority see SA1651 on the previous p	age – Retail	pharmacy 30	✓ <u>Emtriva</u>
LAMIVUDINE - Special Authority see SA1651 on the previous page	– Retail pha	ırmacy	
Tab 150 mg	84.50	60	✓ <u>Lamivudine</u> Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA1651 on the previou Cap 100 mg	s page – Ret	ail pharmacy 100	✓ Retrovir

200 ml OP

✓ Retrovir

	Subsidy		Fully	Brand or
	(Manufacturer's Price			Generic
	\$	Per	✓	Manufacturer
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets) the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg) counts as two anti-		medication	
Tab 500 mg with lamivadine 150 mg			, Vib	парпапп
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA1651 on pa	age 108 - Retail pha	armacy		
Cap 150 mg	141.68	60	✓ Tev	a
Cap 200 mg	188.91	60	✓ Tev	ra
DARUNAVIR - Special Authority see SA1651 on page 108 - Ref	tail pharmacy			
Tab 400 mg		60	✓ Dar	unavir Mylan
ů	335.00		✓ Pre	
Darunavir Mylan to be Sole Supply on 1 April 2021				
Tab 600 mg	196.65	60	Dar	unavir Mylan
	476.00		Pre	zista
Darunavir Mylan to be Sole Supply on 1 April 2021				
(Prezista Tab 400 mg to be delisted 1 April 2021)				
(Prezista Tab 600 mg to be delisted 1 April 2021)				
LOPINAVIR WITH RITONAVIR - Special Authority see SA1651	on page 108 – Reta	ail pharma	су	
Tab 100 mg with ritonavir 25 mg		60	Kal	etra
Tab 200 mg with ritonavir 50 mg	463.00	120	Kal	etra
Oral liq 80 mg with ritonavir 20 mg per ml	735.00 3	00 ml OP	✓ Kal	etra
RITONAVIR - Special Authority see SA1651 on page 108 - Reta	ail pharmacy			
Tab 100 mg	43.31	30	✓ No.	<u>vir</u>
Strand Transfer Inhibitors				
Strana Transier inimbitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 108 - Tab 50 mg		30	✓ Tiv	icay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 or	•	nharmacy		•
Tab 400 mg		60	∕ ✓ Ise	ntress
Tab 600 mg	,	60		ntress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - . PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
 \$	Per	1	

continued...

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA1995 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.
- Inj 180 mcg prefilled syringe......500.00

) 4 Yegasys

⇒SA1995 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has chronic hepatitis C, genotype 1; and
 - 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
 - 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
 - 4 Patient is to be treated in combination with boceprevir; and
 - 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

1 Patient has chronic hepatitis C, genotype 1; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

continued...

- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C. genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive: or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*: and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- - 3.1 Patient has a cutaneous T cell lymphoma*: or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

continued...

- 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Notes:

- · Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Wastage claimable	86.40	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	Arrow-Norfloxacin
Only if proceribed for a nationt with an uncomplicated u	rinany tract infactio	n that ic unr	acananciva to a first line agent or

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
				THE TELEVISION
Autichalinestereses				
Anticholinesterases				
NEGOTIONINE METHOUGH EATE				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	19.60	10	✓	Juno S29
, , , ,	29.40		/	Max Health
	98.00	50		AstraZeneca
	50.00	50	•	AStrazeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.79	100	✓	Mestinon
Non Staraidal Anti Inflammatory Drugo				
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
	4.00		,	5111
* Tab EC 25 mg		50		Diclofenac Sandoz
* Tab 50 mg dispersible	1.50	20	•	Voltaren D
* Tab EC 50 mg	1.23	50	✓	Diclofenac Sandoz
* Tab long-acting 75 mg		500	/	Apo-Diclo SR
* Tab long-acting 100 mg		500		Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5		Voltaren
* Suppos 12.5 mg	2.04	10		Voltaren
* Suppos 25 mg	2.44	10	•	Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10	/	Voltaren
* Suppos 100 mg		10	/	Voltaren
				· ontai on
IBUPROFEN				
* Tab 200 mg	21.40	1,000	•	Relieve
* Tab long-acting 800 mg	5.99	30	/	Ibuprofen SR BNM
* Oral liq 20 mg per ml		200 m		Ethics
		_00		
KETOPROFEN			_	
* Cap long-acting 200 mg	12.07	28	•	Oruvail SR
MEFENAMIC ACID				
	1.05	EΟ		
* Cap 250 mg		50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
NADDOVEN	. ,			
NAPROXEN	00.00	F00		N - fl 050
* Tab 250 mg		500		Noflam 250
* Tab 500 mg	22.19	250		Noflam 500
* Tab long-acting 750 mg	6.16	28	✓	Naprosyn SR 750
* Tab long-acting 1 g	8.21	28		Naprosyn SR 1000
5 5 5				
SULINDAC				
* Tab 100 mg	9.57	56	✓	Mylan S29
* Tab 200 mg	15.10	50	1	Aclin
3	16.91	56		Sulindac Mylan S29
	10.01	50	•	Junitual wylan 22
TENOXICAM				
* Tab 20 mg	9.15	100	1	Tilcotil
* Inj 20 mg vial		1		AFT
- · · · · · · · · · · · · · · · · · · ·		•	,	•

	MUSCULOSKELETAL SYSTEM					
	Subsidy (Manufacturer's Pr \$	ice) Subs	Fully Brand or sidised Generic ✓ Manufacturer			
NSAIDs Other						
CELECOXIB Cap 100 mg Cap 200 mg		60 30	✓ Celecoxib Pfizer ✓ Celebrex ✓ Celecoxib Pfizer			
Topical Products for Joint and Muscular Pain						
CAPSAICIN Crm 0.025% – Special Authority see SA1289 below – Retail pharmacy		45 g OP 60 g OP	✓ Zostrix ✓ Rugby Capsaicin Topical Cream S29			
Zostrix to be Sole Supply on 1 April 2021 Rugby Capsaicin Topical Cream 229 Crm 0.025% to be deliste SA1289 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid steeparthritis that is not responsive to paracetamol and oral non-subsequents.	d without further r					
Antirheumatoid Agents						
HYDROXYCHLOROQUINE — Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, system suppression, relevant dermatological conditions (cutaneous f mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmo Pharmacists may annotate the prescription as endorsed whe hydroxychloroquine. Note: Indication marked with a * is an	orms of lupus and nary)*, and the pare there exists a sunapproved indicate	d lichen planu rescription is e record of prior ation.	s, cutaneous vasculitides and endorsed accordingly. dispensing of			
* Tab 200 mg LEFLUNOMIDE	7.98	100	✓ <u>Plaquenil</u>			
Tab 10 mg	6.00	30	✓ Arava			
Tab 20 mg		30	✓ Arava			
PENICILLAMINE	67.00	100	✓ D Damamina			
Tab 125 mg Tab 250 mg		100 100	✓ D-Penamine✓ D-Penamine			
Drugs Affecting Bone Metabolism						
Alendronate for Osteoporosis						
ALENDRONATE SODIUM * Tab 70 mg	2.44	4	✓ Fosamax			
ALENDRONATE SODIUM WITH COLECALCIFEROL * Tab 70 mg with colecalciferol 5,600 iu	1.51	4	✓ Fosamax Plus			
Other Treatments						

DENOSUMAB - Special Authority see SA1777 on the next page - Retail pharmacy

Inj 60 mg prefilled syringe......326.00

✓ Prolia

1

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

1

Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	17.05	1	Pamisol

RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779 on the next page - Retail pharmacy ✓ Evista

\$29 Unapproved medicine supplied under Section 29

✓ fully subsidised

	Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
	\$	Per	✓	Manufacturer

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg3.10	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml490.00	1	✓ Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

Sub	osidy	Fully	Brand or
(Manufactu	urer's Price) Subsic	lised	Generic
(\$ Per	✓	Manufacturer

continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

	Subsidy Fully (Manufacturer's Price) Subsidised		r
(Manufact		sed Generic	
	\$ Per	Manufac	turer

continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA19	63 below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	 Benzbromaron AL
			100 829

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	9.58	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA1996 below - Retai	l pharmacy		
Tab 80 mg	39.50	28	✓ Adenuric
Tab 120 mg	39.50	28	✓ Adenuric

⇒SA1996 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout: and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

	Subsidy Fully		Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

2 Patient has a documented history of allopurinol intolerance.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

Muscle Relaxants

BACLOFE	

*	Tab 10 mg4.20	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients where oral an	tispastic ag	ents have been ineffective or have
	caused intolerable side effects and the prescription is endorsed accordingly	у.	

Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement............372.98 5 Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

DANTDOLENE

DANTROLENE Con 05 mm	07.50	100	/ Dambulum
Cap 25 mg	97.50	100	Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg	18.54	100	✓ Norflex

121

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised Per Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE A Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			•
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule1	21.84	5	✓ Movapo

BROMOCRIPTINE MESYLATE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of

	prior dispensing of bromocriptine mesylate.			
*	Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ΕN	TACAPONE			
\blacktriangle	Tab 200 mg	22.00	100	✓ Entapone
LE'	VODOPA WITH BENSERAZIDE			
*	Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
*	Cap 50 mg with benserazide 12.5 mg	13.75	100	✓ Madopar 62.5
*	Cap 100 mg with benserazide 25 mg	15.80	100	✓ Madopar 125
*	Cap long-acting 100 mg with benserazide 25 mg	22.85	100	✓ Madopar HBS
*	Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
LE'	VODOPA WITH CARBIDOPA			
*	Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
*	Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
*	Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
PR	AMIPEXOLE HYDROCHLORIDE			
\blacktriangle	Tab 0.25 mg	6.12	100	✓ Ramipex
\blacktriangle	Tab 1 mg		100	✓ Ramipex
RO	PINIROLE HYDROCHLORIDE			
A	Tab 0.25 mg	2.85	84	✓ Ropin
	,	3.39	100	✓ Mylan S29
\blacktriangle	Tab 1 mg	3.95	84	✓ Ropin
	•	4.70	100	✓ Mylan S29
\blacktriangle	Tab 2 mg	5.48	84	✓ Ropin
\blacktriangle	Tab 5 mg	12.50	84	✓ Ropin
SE	LEGILINE HYDROCHLORIDE			
*	Tab 5 mg	22.00	100	✓ Apo-Selegiline S29 S29
ТО	LCAPONE			
A	Tab 100 mg	152.38	100	✓ Tasmar

Anticholinergics

ENZATROPINE MESYLATE			
Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml		5	✓ Phebra

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

				VOOS STSTEIN
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ K	(emadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	•	56	./ c	Rilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:			_	
1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.				initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow.	onths for applications	s mee	ting the follo	owing criteria:
TETRABENAZINE Tab 25 mg	91.10	112	✓ <u>N</u>	<u>Notetis</u>
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement	dministration and the	30 ml pres 10	cription is e	(ylocaine 2% Jelly ndorsed accordingly. nstillagel Lido

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

a) Up to 5 each available on a PSO

accordingly.

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 m	ı 🗸	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	✓	Lidocaine-Claris
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓	Lidocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓	Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓	Lidocaine-Claris
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	103.32	10	1	Pfizer
a) Up to 5 each available on a PSO		. •		
b) On heiding a heid and a firm and the standard and a second and a				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see	SA0906 above – Retail pharn	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - S	Special Authority see SA0906	above – Reta	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 114

Non-opioid Analgesics

ASPIRIN * Tab dispersible 300 mg - Up to 30 tab available on a PS	SO4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia accordingly.	or diabetic periphera	I neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream S29
Zostrix HP to be Sole Supply on 1 April 2021			
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price \$) Sul	Fully osidised	Brand or Generic Manufacturer
ARACETAMOL				
Tab 500 mg - blister pack	0.50	20	✓ F	/ledco Paracare Pharmacy Health
	1.12			Ethics Paracetamol Classic
	2.48	100	-	Paracare Pharmacy Health
	11.75	96		Panadol Mini Caps
	24.82	1,000		Paracetamol Pharmacare
			✓ F	harmacare
 a) Maximum of 300 tab per prescription; can be waived b) Up to 30 tab available on a PSO c) 	by endorsement			
 Subsidy by endorsement for higher quantities i regular daily dosing for one month or greater, a annotate the prescription as endorsed where d Maximum of 100 tab per dispensing for non-en (for non-endorsed patients), then dispense in r 	and the prescription is ispensing history sup dorsed patients. If q	s annotate ports a lo juantities į	ed accor ng-term orescribe	dingly. Pharmacists ma condition. ed for more than 100 tab
Tab 500 mg - bottle pack – Maximum of 300 tab per	epeat disperisings no	n exceedi	ily 100 t	ab per dispensing.
prescription; can be waived by endorsement	24.82	1,000	√ F	Paracetamol Pharmacare

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

*	Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓ Paracare
	a) Up to 200 ml available on a PSO			
	b) Not in combination			
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double Strength
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3 29	10	✓ Gacet
*			10	✓ Gacet
-	Suppos 250 mg			
*	Suppos 500 mg	12.40	50	✓ <u>Gacet</u>
^	minid Ampleonian			
U	pioid Analgesics			
CO	DEINE PHOSPHATE - Safety medicine; prescriber may determin	e dispensina f	requency	
-	Tab 15 mg		100	✓ PSM
	Tab 30 mg	7 45		✓ PSM
	Tau ou my	1.40	100	
	Tab 60 mg	14.25	100	✓ <u>PSM</u>

60

✓ DHC Continus

Tab long-acting 60 mg......8.60

DIHYDROCODEINE TARTRATE

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	e) Su Per	bsidised	Generic Manufacturer
ETALT AADV	Ψ	1 61		Wandacturer
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable	,			
c) Safety medicine; prescriber may determine dispensing		4.0		
Inj 50 mcg per ml, 2 ml ampoule		10	-	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	-	Boucher and Muir
Patch 12.5 mcg per hour		5		Fentanyl Sandoz
Patch 25 mcg per hour		5		Fentanyl Sandoz
Patch 50 mcg per hour		5		Fentanyl Sandoz
Patch 75 mcg per hour		5		Fentanyl Sandoz
Patch 100 mcg per hour	11.40	5	✓	Fentanyl Sandoz
a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing d) Extemporaneously compounded methadone will only b (methadone powder, not methadone tablets). e) For methadone hydrochloride oral liquid refer Standard Tab 5 mg Oral liq 2 mg per ml Oral liq 5 mg per ml Oral liq 10 mg per ml Inj 10 mg per ml, 1 ml	e reimbursed at the ra Formulae, page 2391.405.795.796.79	10 200 ml 200 ml 200 ml 200 ml 10	V V	t form available Methatabs Biodone Biodone Forte Biodone Extra Forte AFT
b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing d) Extemporaneously compounded methadone will only b (methadone powder, not methadone tablets). e) For methadone hydrochloride oral liquid refer Standard Tab 5 mg Oral liq 2 mg per ml Oral liq 5 mg per ml Oral liq 10 mg per ml	e reimbursed at the ra Formulae, page 2391.405.795.796.79	10 200 ml 200 ml 200 ml	V V	Methatabs Biodone Biodone Forte Biodone Extra Forte

Oral liq 10 mg per ml27.74

✓ Ordine S29

✓ RA-Morph

200 ml

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)		Subsidised		
	\$	Per	1	Manufacturer	
MORPHINE SULPHATE					
a) Only on a controlled drug form					
b) No patient co-payment payable					
c) Safety medicine; prescriber may determine dispensing for		40	,	0	
Tab immediate-release 10 mg		10		Sevredol	
Tab immediate-release 20 mg		10		Sevredol	
Tab long-acting 30 mg		10		Arrow-Morphine LA	
Tab long-acting 60 mg		10		Arrow-Morphine LA	
Cap long-acting 10 mg		10		m-Eslon	
Cap long-acting 30 mg		10		m-Eslon	
Cap long-acting 60 mg		10		m-Eslon	
Cap long-acting 100 mg		10		m-Eslon	
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	'SO6.99	5	/	DBL Morphine	
				Sulphate	
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO5.61	5	✓	DBL Morphine	
				Sulphate	
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO7.08	5	1	DBL Morphine	
,				Sulphate	
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 7.28	5	/	DBL Morphine	
ing oo mg por mi, i mi ampoulo op to o mg available on a	. 00	Ü	_	Sulphate	
(Arrow-Morphine LA Tab long-acting 30 mg to be delisted 1 Jun	0.2021)			Ouiphate	
(Arrow-Morphine LA Tab long-acting 60 mg to be delisted 1 Apr	11 2021)				
OXYCODONE HYDROCHLORIDE					
a) Only on a controlled drug form					
b) No patient co-payment payable					
c) Safety medicine; prescriber may determine dispensing for	requency				
Tab controlled-release 5 mg	2.15	20	✓	Oxycodone Sandoz	
•	3.01	28	✓	Oxycodone Sandoz	
				S29 S29	
Tab controlled-release 10 mg	2 15	20	/	Oxycodone Sandoz	
Tab controlled-release 20 mg		20		Oxycodone Sandoz	
Tab controlled-release 40 mg		20		Oxycodone Sandoz	
Tab controlled-release 80 mg		20		Oxycodone Sandoz	
Cap immediate-release 5 mg		20	•	OxyNorm OxyNorm	
		20			
Cap immediate-release 10 mg				OxyNorm OxyNorm	
Cap immediate-release 20 mg		20		OxyNorm OxyNorm	
Oral liq 5 mg per 5 ml		250 m		OxyNorm	
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm OxyNorm	
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm OxyNorm	
Inj 50 mg per ml, 1 ml ampoule		5		<u>OxyNorm</u>	
PARACETAMOL WITH CODEINE - Safety medicine; prescribe		ensin	g frequenc	у	
* Tab paracetamol 500 mg with codeine phosphate 8 mg	26.51	1,000	/	Paracetamol +	

Codeine (Relieve)

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
PETHIDINE HYDROCHLORIDE	•			
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f	requency			
Tab 50 mg		10	✓	PSM
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO29.88	5	✓	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO30.72	5	✓	DBL Pethidine Hydrochloride
TRAMADOL HYDROCHLORIDE				•
Tab sustained-release 100 mg	1.52	20	✓.	Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20	✓:	Tramal SR 200
Cap 50 mg	2.80	100	✓	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
MITRIPTYLINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 10 mg	2.49	100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100		Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; presc	riber may determine d	isper	ising freque	ency
Tab 10 mg	13.99	100	✓.	Anafranil S29
				Apo-Clomipramine
Tab 25 mg	9.46	100	✓ ,	Apo-Clomipramine
Anafranil S29 Tab 10 mg to be delisted 1 May 2021)				
OOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by e	ndorsement			
a) Safety medicine; prescriber may determine dispensing f	requency			
b) Subsidy by endorsement – Subsidised for patients who	were taking dosulepin	[doth	iepin] hydro	ochloride prior to 1 June
2019 and the prescription is endorsed accordingly. Pha	rmacists may annotate	the	prescription	as endorsed where the
exists a record of prior dispensing of dosulepin [dothiepi				
Tab 75 mg		30		Dosulepin Mylan
Cap 25 mg	7.83	50	•	Dosulepin
				Mylan S29
MIPRAMINE HYDROCHLORIDE - Safety medicine; prescribe	r may determine dispe	nsing	frequency	
Tab 10 mg	5.48	50		Tofranil
	10.96	100	_	Tofranil
Tab 25 mg	8.80	50	•	Tofranil
MAPROTILINE HYDROCHLORIDE - Subsidy by endorsement				
a) Safety medicine; prescriber may determine dispensing f				
 Subsidy by endorsement – Subsidised for patients who 2020 and the prescription is endorsed accordingly. Pha 				
exists a record of prior dispensing of maprotiline hydrocl				
Tab 75 mg	14.01	20		Ludiomil
	21.01	30	✓	Ludiomil
Ludiomil Tab 75 mg to be delisted 1 August 2021)				
(Ludiomil Tab 75 mg to be delisted 1 August 2021)				

			<u> </u>
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presc	· · · · · · · · · · · · · · · · · · ·		
Tab 10 mg	•		Norpress
Tab 25 mg			Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective		
TRANYLCYPROMINE SULPHATE			
Tab 10 mg	12.85	28	Parnate S29 S29
·	22.94	50	Parnate
	45.88	100	Parnate S29 S29
	96.00	•	Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg	6.40	60	Aurorix
* Tab 300 mg	9.80	60	Aurorix
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	1.52	84	PSM Citalopram
ESCITALOPRAM			
* Tab 10 mg	1.40	28	Escitalopram- Apotex
* Tab 20 mg	2.49	28	Escitalopram- Apotex
FLUOXETINE HYDROCHLORIDE			
* Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	1.98	30	Fluox
When prescribed for a patient who cannot swallow accordingly; or	whole tablets or caps	sules and the p	rescription is endorsed
2) When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with	•	•	•
Cap 20 mg	2.91	84	Fluox
PAROXETINE			
* Tab 20 mg	1.20	30	Paxtine
•	3.61	90	Loxamine
SERTRALINE			
Tab 50 mg	0.92		Setrona Setrona All
	3.05		Setrona AU Arrow-Sertraline
Tab 100 mg		_	Setrona
145 100 mg			Setrona AU
	5.25	90	Arrow-Sertraline

MIRTAZAPINE		Subsidy (Manufacturer's Price)	Per	Fully Brand or Subsidised Generic r Manufacturer
Tab 45 mg	Other Antidepressants			
# Cap 37.5 mg	Tab 30 mg			
Antiepilepsy Drugs Agents for Control of Status Epilepticus CLONAZEPAM — Safety medicine; prescriber may determine dispensing frequency Inj 1 mg per ml, 1 ml. — 21.00	* Cap 37.5 mg * Cap 75 mg	8.11	84	✓ Enlafax XR
CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency				
Inj 1 mg per ml, 1 ml	Agents for Control of Status Epilepticus			
Inj 5 mg per ml, 2 ml ampoule — Subsidy by endorsement	· · · · · · · · · · · · · · · · · · ·		5	✓ Rivotril
Rectal tubes 5 mg - Up to 5 tube available on a PSO	Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO		5	✓ Hospira
★ Inj 5 ml	Rectal tubes 5 mg - Up to 5 tube available on a PSO		5	✓ Stesolid
★ Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO	* Inj 5 ml	1,500.00	5	✓ AFT S29
CARBAMAZEPINE * Tab 200 mg	* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a			·
CARBAMAZEPINE		133.92	5	✓ Hospira
★ Tab 200 mg 14.53 100 ✓ Tegretol ★ Tab long-acting 200 mg 16.98 100 ✓ Tegretol CR ★ Tab 400 mg 34.58 100 ✓ Tegretol ★ Tab long-acting 400 mg 39.17 100 ✓ Tegretol CR ★ Oral liq 20 mg per ml 26.37 250 ml ✓ Tegretol CLOBAZAM - Safety medicine; prescriber may determine dispensing frequency Tab 10 mg 9.12 50 ✓ Frisium CLONAZEPAM - Safety medicine; prescriber may determine dispensing frequency Oral drops 2.5 mg per ml 7.38 10 ml OP ✓ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg 2.65 100 ✓ Apo-Gabapentin ★ Cap 300 mg 4.07 100 ✓ Apo-Gabapentin				
Tab 10 mg	* Tab 200 mg	16.98 34.58 39.17 26.37 2	100 100 100	✓ Tegretol CR ✓ Tegretol ✓ Tegretol CR
Oral drops 2.5 mg per ml 7.38 10 ml OP ✔ Rivotril ETHOSUXIMIDE Cap 250 mg 140.88 100 ✔ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✔ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg 2.65 100 ✔ Apo-Gabapentin ★ Cap 300 mg 4.07 100 ✔ Apo-Gabapentin	Tab 10 mg	9.12	50	✓ Frisium
Cap 250 mg 140.88 100 ✓ Zarontin Oral liq 250 mg per 5 ml 56.35 200 ml ✓ Zarontin GABAPENTIN Note: Not subsidised in combination with subsidised pregabalin * Cap 100 mg 2.65 100 ✓ Apo-Gabapentin * Cap 300 mg 4.07 100 ✓ Apo-Gabapentin	Oral drops 2.5 mg per ml		ml C	OP
Note: Not subsidised in combination with subsidised pregabalin ★ Cap 100 mg	Cap 250 mg			
★ Cap 300 mg		alin		
	* Cap 300 mg	4.07	100	✓ Apo-Gabapentin

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
LACOSAMIDE - Special Authority see SA1125 below - Retail p	harmacy				
▲ Tab 50 mg	25.04	14	✓ V	impat	
▲ Tab 100 mg	50.06	14	✓ V	impat	
•	200.24	56	✓ V	impat	
▲ Tab 150 mg	75.10	14	✓ V	impat	
ŭ	300.40	56	✓ V	impat	
▲ Tab 200 mg	400.55	56		impat	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

L/ \	MOTHGINE		
lack	Tab dispersible 2 mg55.00	30	✓ Lamictal
lack	Tab dispersible 5 mg50.00	30	✓ Lamictal
*	Tab dispersible 25 mg2.76	56	✓ Logem
*	Tab dispersible 50 mg	56	✓ Logem
*	Tab dispersible 100 mg4.40	56	Logem
ΙF	VETIRACETAM		
	Tab 250 mg4.99	60	✓ Everet
	Tab 500 mg	60	✓ Everet
	Tab 750 mg	60	✓ Everet
	Tab 1,000 mg	60	✓ Everet
	Oral lig 100 mg per ml	300 ml OP	✓ Levetiracetam-AFT
DL	ENOBARBITONE		
ЕП	For phenobarbitone oral liquid refer Standard Formulae, page 239		
*	Tab 15 mg40.00	500	✓ PSM
*	Tab 30 mg	500	✓ PSM
	· ·	300	▼ <u>FSWI</u>
	ENYTOIN SODIUM		
*	Tab 50 mg75.00	200	✓ Dilantin Infatab
	Cap 30 mg74.00	200	✓ Dilantin
	Cap 100 mg37.00	200	Dilantin
*	Oral liq 30 mg per 5 ml22.03	500 ml	Dilantin
PR	EGABALIN		
	Note: Not subsidised in combination with subsidised gabapentin		
*	Cap 25 mg2.25	56	✓ Pregabalin Pfizer
*	Cap 75 mg	56	✓ Pregabalin Pfizer
*	Cap 150 mg4.01	56	✓ Lyrica
			✓ Pregabalin Pfizer
*	Cap 300 mg7.38	56	✓ Pregabalin Pfizer

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	I Generic
	\$	Per		Manufacturer
PRIMIDONE				
* Tab 250 mg	17.25	100	✓	Apo-Primidone
	62.00	200	•	Mysoline S29 S29
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC		100	✓	Epilim
* Oral liq 200 mg per 5 ml		300 m	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	•	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail p	harmacy			
Cap 250 mg	509.29	60	✓	Diacomit S29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

▲ Tab 25 mg	11.07	60	✓ Arrow-Topiramate
v			✓ Topiramate Actavis
	26.04		✓ Topamax
Tab 50 mg	18.81	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	44.26		✓ Topamax
Tab 100 mg	31.99	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	75.25		✓ Topamax
Tab 200 mg	55.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
1	29.85		✓ Topamax
Sprinkle cap 15 mg	20.84	60	✓ Topamax
Sprinkle cap 25 mg	26.04	60	✓ Topamax
GABATRIN - Special Authority see SA1997 below - Retail pharmacy	,		•
Tab 500 mg		100	✓ Sabril
04400			

⇒SA1997 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Either:

TOPIRAMATE

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

continued...

\$29 Unapproved medicine supplied under Section 29

NERVOUS SYSTEM

Subsidy Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

Acute Migraine Treatment

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 114

Acute wiigianie Treatment			
RIZATRIPTAN			
Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	24.44	100	✓ Apo-Sumatriptan
Tab 100 mg	46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj	per		
prescription	34.00	2 OP	✓ <u>Imigran</u>
Burnhalanda af Manada a			

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 54 PIZOTIFEN

* Tab 500 mcg.......23.21 100 ✓ Sandomigran

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 on the next page – Retail pharmacy

Cap 2 × 80 mg and 1 × 125 mg......84.00 3 OP **Emend Tri-Pack**

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
φ.	Dox /	Manufacturer	

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

	TAHISTINE DIHYDROCHLORIDE	
*	Tab 16 mg	

* Tab 16 mg	3.88	84	✓ <u>Vergo 16</u>
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg	0.55	10	✓ Nausicalm
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml	14.95	5	✓ Nausicalm
	21.53	10	✓ Hameln
Hameln to be Sole Supply on 1 May 2021			
(Nausicalm Inj 50 mg per ml, 1 ml to be delisted 1 May 202	21)		
DOMPERIDONE			
* Tab 10 mg	2.25	100	✓ Pharmacy Health
HYOSCINE HYDROBROMIDE			
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1998 below -	- Retail		
pharmacy	14.11	2	✓ Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

* Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
* Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO9.50	10	✓ <u>Pfizer</u>
ONDANSETRON		
* Tab 4 mg2.68	50	✓ Onrex
* Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ Ondansetron
		ODT-DRLA
* Tab 8 mg4.57	50	✓ Onrex
* Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ Ondansetron
		ODT-DRLA
PROCHLORPERAZINE		
* Tab 3 mg buccal5.97	50	
(30.00)		Buccastem
* Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacture

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dis	spensing frequence	CV	
Tab 100 mg	5.15	30	✓ Sulprix
•	17.16	100	✓ Amisulpride
			Mylan S29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg		60	✓ Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine d	isnensina freauer	nev	
Tab 5 mg		30	✓ Aripiprazole Sandoz
	28.58	49	✓ Aripiprazole 1A
	20.00		Pharma S29
Tab 10 mg	17 50	30	✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	✓ Aripiprazole Sandoz
(Aripiprazole 1A Pharma S29) Tab 5 mg to be delisted 1 June 20		30	Alipipiazole Jaliuoz
. , ,	,		
CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; pr			
Tab 10 mg – Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	✓ Largactil
CLOZAPINE – Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequ	ency		
Tab 25 mg	5.69	50	✓ Clozaril
	6.69		✓ Clopine
	11.36	100	✓ Clozaril
	13.37		✓ Clopine
Tab 50 mg	8.67	50	✓ Clopine
	17.33	100	✓ Clopine
Tab 100 mg	14.73	50	Clozaril
	17.33		✓ Clopine
	29.45	100	Clozaril
	34.65		✓ Clopine
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	Clopine
Suspension 50 mg per ml	17.33	100 ml	✓ Clopine
	67.62		✓ Versacloz
HALOPERIDOL - Safety medicine; prescriber may determine di	spensing frequer	icv	
Tab 500 mcg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50	✓ Serenace
	29.72	100	✓ Serenace
Oral liq 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a P		10	✓ Serenace

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
LEVOMEPROMAZINE - Safety medicine; prescriber may dete	rmine dispensing frequency	uency		
Tab 25 mg (33.8 mg as a maleate)	16.10	100	1	Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100	•	Nozinan
Tab 100 mg (135 mg as a maleate)	41.75	100	✓	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	•	<u>Nozinan</u>
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may determ	nine di	spensing	frequency
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
LITHIUM CARBONATE – Safety medicine; prescriber may dete		uonov		
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100		Douglas
, ,		100	•	Douglas
OLANZAPINE – Safety medicine; prescriber may determine dis			_	
Tab 2.5 mg		28	_	Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28	_	Zypine
Tab orodispersible 10 mg	2.38	28	•	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg	10.49	84	✓	Neulactil
	12.49	100	✓	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
	44.45	100	✓	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	_	Quetapel
v				
RISPERIDONE – Safety medicine; prescriber may determine di		60	./	Risperidone (Teva)
Tab 0.5 mg Tab 1 mg		60		Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 3 mg		60	_	Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral lig 1 mg per ml		30 ml		Risperon
		00 1111	•	порстоп
ZIPRASIDONE – Safety medicine; prescriber may determine di			,	
Cap 20 mg		60	_	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60		Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	escriber may determin	e disp	•	' '
Tab 10 mg	31.45	100	•	Clopixol
Depot Injections				
FILIDENTINO DECANOATE Cofety mandial and a second beautiful and a se		da a f		
FLUPENTHIXOL DECANOATE – Safety medicine; prescriber n	•	-		Fluencel
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	•	Fluanxol

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	av determine dispensi	na freau	encv	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	•	5		laldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		laldol Concentrate laldol Decanoas \$29
OLANZAPINE – Special Authority see SA1428 below – Retail p Safety medicine; prescriber may determine dispensing frequ	ency			Decanoas 220
Inj 210 mg vial	252.00	1	√ <u>Z</u>	'yprexa Relprevv
Inj 300 mg vial	414.00	1	√ <u>Z</u>	Zyprexa Relprevy
Inj 405 mg vial	504.00	1	√ <u>Z</u>	yprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing fre	quency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency			
Inj 25 mg vial	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

NERVOUS SYSTEM

	Subsidy		Fully	Brand or
((Manufacturer's Price)	Subsi	dised	Generic
	\$	Per	✓	Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 **✓ Clopixol**

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine dispe	ensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ <u>Paxam</u>
DIAZEPAM - Safety medicine; prescriber may determine dispensir	ng frequency		
Tab 2 mg		500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispen	sing frequency		
Tab 1 mg	9.72	250	✓ <u>Ativan</u>
Tab 2.5 mg	12.50	100	✓ <u>Ativan</u>
OXAZEPAM - Safety medicine; prescriber may determine dispensi	ing frequency		
Tab 10 mg	6.17	100	✓ Ox-Pam
Tab 15 mg	8.53	100	✓ Ox-Pam

Multiple Sclerosis Treatments

⇒SA2026 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

past 24 months; and

- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) at any time in the last six months (i.e. the patient has walked 100 metres or more with

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2026 on the previous page - Retail pharmacy

a) Wastage claimable

or without aids in the last six months).

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

 Cap 120 mg
 520.00
 14
 ✓ Tecfidera

 Cap 240 mg
 2,000.00
 56
 ✓ Tecfidera

FINGOLIMOD - Special Authority see SA2026 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Cap 0.5 mg......2,200.00 28 Gilenya

GLATIRAMER ACETATE - Special Authority see SA2026 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

lnj 40 mg prefilled syringe.................2,275.00 12 **✓ Copaxone**

INTERFERON BETA-1-ALPHA - Special Authority see SA2026 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer \$ INTERFERON BETA-1-BETA - Special Authority see SA2026 on page 138 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. NATALIZUMAB - Special Authority see SA2026 on page 138 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. OCRELIZUMAB - Special Authority see SA2026 on page 138 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Inj 30 mg per ml, 10 ml vial......9,346.00 ✓ Ocrevus TERIFLUNOMIDE - Special Authority see SA2026 on page 138 - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 28 ✓ Aubagio Aubagio to be Sole Supply on 1 June 2021 Sedatives and Hypnotics MELATONIN - Special Authority see SA1666 below - Retail pharmacy Tab modified-release 2 mg - No more than 5 tab per day28.22 30 ✓ Circadin ⇒SA1666 Special Authority for Subsidy Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and 4 Patient is aged 18 years or under*. Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient is aged 18 years or under*; and

- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine disp	ensing frequency		
Inj 1 mg per ml, 5 ml ampoule	2.98	10	✓ Mylan Midazolam
	4.30		✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj availab	le		
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	e endorsed for stat	tus epilepticu	ıs use only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available	e on		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	e endorsed for stat	tus epilepticu	ıs use only.

			NERVOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
PHENOBARBITONE SODIUM - Special Authority see SA1386	below – Retail pharm	асу	
Inj 200 mg per ml, 1 ml ampoule	78.20	10	✓ Max Health S29
➤ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria:	d without further rene	wal u	nless notified for applications meet
Both: 1 For the treatment of terminal agitation that is unresponsiv 2 The applicant is part of a multidisciplinary team working ir	• •	ł	
TEMAZEPAM – Safety medicine; prescriber may determine disp Tab 10 mg	. ,	25	✓ <u>Normison</u>
TRIAZOLAM – Safety medicine; prescriber may determine dispertable 125 mcg	5.10	100	
Tab 250 mcg	(9.85) 4.10 (11.20)	100	Нурат
ZOPICLONE – Safety medicine; prescriber may determine dispe	, ,		, μ
Tab 7.5 mg		500	✓ Zopiclone Actavis
Stimulants/ADHD Treatments			
ATOMOXETINE			
Cap 10 mg	18.41	28	✓ Generic Partners
Cap 18 mg	27.06	28	✓ Generic Partners
Cap 25 mg	29.22	28	✓ Generic Partners
Cap 40 mg		28	✓ Generic Partners
Cap 60 mg	46.51	28	 Generic Partners
Cap 80 mg	56.45	28	Generic Partners
Cap 100 mg	58.48	28	Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see SA1149 b	oelow – Retail pharma	асу	

a) Only an a sandralled drop form

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg20.00 100 ✓ PSM

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a) Only on a controlled drug form

 b) Safety medicine; prescriber may determine dispen 	sing frequency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
•			✓ Rubifen
Tab extended-release 18 mg	7.75	30	 Methylphenidate ER
· ·			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg		30	✓ Rubifen SR
Ç	50.00	100	✓ Ritalin SR
Tab extended-release 27 mg	11.45	30	 Methylphenidate ER
· ·			- Teva
Tab extended-release 36 mg	15.50	30	✓ Methylphenidate ER
3			- Teva
Tab extended-release 54 mg	22.25	30	✓ Methylphenidate ER
		• • •	- Teva

(Ritalin SR Tab sustained-release 20 mg to be delisted 1 June 2021)

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg		30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg		30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below -	Retail pharmacy		
Patch 4.6 mg per 24 hour	48.75	30	✓ Generic Partners
Patch 9.5 mg per 24 hour	48.75	30	✓ Generic Partners

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

NERVOUS SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

28

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ <u>Buprenorphine</u> Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg.......11.00 30 ✓ Zyban

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Su Per	Fully bsidised	Brand or Generic Manufacturer	
DISULFIRAM Tab 200 mg	250.00	100		ıntabuse	
NALTREXONE HYDROCHLORIDE – Special Authority see SA1 Tab 50 mg		harmacy 30		laltraccord	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

e provisions in P	art I of Secti	on A.
18.14	28	Habitrol
3.94	7	Habitrol
19.95	28	Habitrol
4.52	7	Habitrol
22.86	28	Habitrol
5.18	7	Habitrol
19.18	216	Habitrol
3.20	36	Habitrol
21.02	216	Habitrol
3.24	36	Habitrol
38.21	384	Habitrol
8.64	96	Habitrol
38.21	384	Habitrol
8.64	96	Habitrol
44.17	384	Habitrol
10.01	96	Habitrol
44.17	384	Habitrol
10.01	96	Habitrol
	e provisions in P18.143.9419.954.5222.865.1819.183.2021.023.2438.218.6444.1710.01	

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	25.64	53 OP	✓ Varenicline Pfizer
Tab 1 mg	27.10	56	✓ Varenicline Pfizer

NERVOUS SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)		bsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA1667 below

Inj 25 mg vial	 271.35 1	✓ Ribomustin
Inj 100 mg vial	 1,085.38 1	✓ Ribomustin
Inj 1 mg for ECP	 11.40 1 mg	✓ Baxter

⇒SA1667 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.
 Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

BUSULFAN – PCT – Retail pharmacy-Specialist Tab 2 mg	90.25	100	✓ Myleran
<u> </u>	09.23	100	• Wylcian
CARBOPLATIN – PCT only – Specialist Inj 10 mg per ml, 45 ml vial	20 50	1	✓ DBL Carboplatin
mj to mg per mi, 45 mi viai	45.20	1	✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP		1 mg	✓ Baxter
· · · ·		i ilig	Dunio
CARMUSTINE – PCT only – Specialist	1 207 00	4	✓ DICNII
Inj 100 mg vial	1,367.00	1	✓ BiCNU
In: 100 mm for FOD	1 007 00	100 OD	✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	12.29	1	 DBL Cisplatin
	15.00		 Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1	 DBL Cisplatin
	21.00		 Cisplatin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
(DBL Cisplatin Inj 1 mg per ml, 50 ml vial to be delisted 1 April	2021)		
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
, , , , , , , , , , , , , , , , , , ,	158.00	100	✓ Procytox S29
Wastage claimable	100.00	100	1 Tooytox
Inj 1 g vial – PCT – Retail pharmacy-Specialist	35.65	1	✓ Endoxan
, 3 , ., ., ., ., .,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter
LOMUSTINE – PCT – Retail pharmacy-Specialist		9	
Cap 10 mg	132 50	20	✓ CeeNU
Cap 40 mg		20	✓ CeeNU
· · · · · ·		20	- 300110
MELPHALAN Tob 2 mg	40.70	0.5	. Allrayan
Tab 2 mg — PCT — Retail pharmacy-Specialist		25 1	✓ Alkeran✓ Alkeran
Inj 50 mg - PCT only - Specialist	07.80	ı	
	100.00		✓ Alkeran S29 S29
	420.00		✓ Tillomed S29

	Subsidy Manufacturer's Price \$) Per	Fully Subsidised	
OXALIPLATIN – PCT only – Specialist Inj 100 mg vial	25.01	1	•	Oxaliplatin Actavis
Inj 5 mg per ml, 20 ml vial	110.00 46.32 0.48	1 1 mg	/	Oxaliplatin Ebewe Oxaliplatin Accord Baxter
THIOTEPA – PCT only – Specialist Inj 15 mg vial		1	•	Bedford \$29 THIO-TEPA \$29
Inj 100 mg vial	CBS	1		Tepadina S29 Tepadina S29
Antimetabolites				
AZACITIDINE – PCT only – Specialist – Special Authority see SA Inj 100 mg vial		1		Azacitidine Dr Reddy's Vidaza

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or

1 ma

✓ Baxter

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and

- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	2			
	Subsidy (Manufacturer's Pri	(a)	Fully ubsidised	Brand or Generic
	\$	Per	ubsidised ✓	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	1	DBL Leucovorin
, , , , , , , , , , , , ,				Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia	alist7.28	1	✓	Calcium Folinate
				<u>Sandoz</u>
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate
				Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	/	Calcium Folinate
			_	Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	/	Calcium Folinate
			_	Ebewe
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	•	Calcium Folinate
				Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	•	Calcium Folinate
				Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	•	Calcium Folinate
Init was few FOD DOT only. Consciolist	0.00	4		Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	•	Baxter
CAPECITABINE – Retail pharmacy-Specialist				
Tab 150 mg		60	_	Capercit
Tab 500 mg	49.00	120	•	Capercit
CLADRIBINE – PCT only – Specialist	740.00			
Inj 1 mg per ml, 10 ml		10 01	_	Leustatin
Inj 10 mg for ECP	749.96	10 mg OF	•	Baxter
CYTARABINE	"	_		D."
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	alist 400.00	5	•	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail	41.06		./	Pfizer
pharmacy-SpecialistInj 1 mg for ECP - PCT only - Specialist		1 10 mg		Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Specialst		100 mg O	_	Baxter
FLUDARABINE PHOSPHATE		roo mg o		Duxtor
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	1	Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		50 mg OF		Baxter
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	12.00	1	1	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1		Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist		100 mg	1	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist		Ū		
Inj 1 g, 26.3 ml vial	62.50	1	/	DBL Gemcitabine
Inj 1 g		1	1	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg		Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist		-		
Inj 20 mg per ml, 5 ml vial	71.44	1	/	Irinotecan
. •				Accord S29
			/	Irinotecan Actavis
				100
	400.00		./	
	100.00		•	Irinotecan-Rex

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pric	e) Per	Fully Subsidised	Brand or Generic Manufacturer
MERCAPTOPURINE Tab 50 mg - PCT - Retail pharmacy-Specialist		25	√ <u>F</u>	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist Special Authority see SA1725 below		100 ml (OP 🗸 A	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

ME	THOTREXATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Hospira
			✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe14.77	1	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ DBL Methotrexate Onco-Vial
			Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
* *	Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist25.00 Inj 100 mg per ml, 50 ml vial – PCT – Retail	1	✓ Methotrexate Ebewe
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
	ospira Inj 2.5 mg per ml, 2 ml to be delisted 1 May 2021) BL Methotrexate Onco-Vial Inj 25 mg per ml, 2 ml vial to be delisted 1 May 202:	1)	
PE	METREXED - PCT only - Specialist - Special Authority see SA1679 below		
	Inj 100 mg vial60.89	1	✓ Juno Pemetrexed
	Inj 500 mg vial217.77	1	 Juno Pemetrexed
_	Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	ç	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
- 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	✓ Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg	CBS	100	✓ Agrylin S29 S29
			✓ Teva S29
	1,175.87		✓ Agrylin
(Agrylin S29 S29 Cap 0.5 mg to be delisted 1 April 2021)			
(Teva S29 Cap 0.5 mg to be delisted 1 April 2021)			
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Ini 10 mg for ECP	481.70	10 ma OP	✓ Baxter

	Subsidy (Manufacturer's Pr	ice) Sub	Fully sidised	Brand or Generic
	\$	Per	•	Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	161.01	1	✓ [DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ E	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see S	SA1889 below			
Inj 3.5 mg vial		1	√ E	Bortezomib Dr-Reddy's
Inj 1 mg for ECP	31.20	1 mg	✓ E	Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

Note: Indications marked with * are unapproved indications.			
DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	 DBL Dacarbazine
	580.60	10	 Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist			
Inj 2 mg per ml, 10 ml	149.50	1	✓ Pfizer
Inj 20 mg for ECP		20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist		Ü	
Inj 10 mg per ml, 2 ml vial	12.40	1	✓ DBL Docetaxel
Inj 20 mg		1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1	✓ Docetaxel
, , , , , , , , , , , , , , , , , , , ,			Accord \$29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
(DBL Docetaxel Inj 10 mg per ml, 2 ml vial to be delisted 1 June		3	
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist	,		
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebewe
.,	17.00	-	✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Doxorubicin Ebewe
	65.00		✓ Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	 Epirubicin Ebewe
2			

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Fully Subsidised	Generic
	\$	Per		Manufacturer
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	✓	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Special	ist7.90	1	1	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	✓	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	_	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phat Brand switch fee payable (Pharmacode 2603187) - see page Cap 500 mgIDARUBICIN HYDROCHLORIDE Inj 5 mg vial – PCT only – Specialist	237 for details 23.82	100		<u>Devatis</u> Zavedos
Inj 10 mg vial - PCT only - Specialist		1	1	Zavedos
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓	Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authorit Wastage claimable	ty see SA1897 below			
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg		21	1	Revlimid
	6,207.00	28	✓	Revlimid
Cap 15 mg	5,429.39	21	✓	Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg	7,627.00	21	✓	Revlimid

⇒SA1897 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
 - Fither
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 The patient has ECOG performance score of 0-1; and
- 5 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

haematologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

MESTO C		
Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 5 mg vial851.37	1	✓ Teva
Inj 20 mg vial	1	✓ Omegapharm \$29
, ,		✓ Teva
Inj 1 mg for ECP	1 mg	✓ Baxter
(Teva Inj 5 mg vial to be delisted 1 June 2021)	· ·	
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1883 belo	w	
Tab 100 mg3,701.00	56	✓ Lynparza
Tab 150 mg3,701.00	56	✓ Lynparza
Cap 50 mg - Wastage claimable	448	✓ Lynparza
(Lynparza Cap 50 mg to be delisted 1 July 2021)		

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based

Subsid	dy Fi	ılly Brand or	
(Manufacture	r's Price) Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

regimen: and

- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist

47.30	5	Paclitaxel Ebewe
24.00	1	✓ Paclitaxel Ebewe
91.67		✓ Paclitaxel Actavis
26.69	1	✓ Paclitaxel Ebewe
137.50		✓ Anzatax
		✓ Paclitaxel Actavis
44.00	1	✓ Paclitaxel Ebewe
275.00		✓ Anzatax
		✓ Paclitaxel Actavis
0.20	1 mg	✓ Baxter
SA1979 below	_	
3,455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN]	- PCT	only - Specialist
-------------------------------	-------	-------------------

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharma	cy-Specialist		
Cap 50 mg		50	✓ Natulan S29

	Subsidy (Manufacturer's Price)	Per	Fully Brand or Subsidised Generic or Manufacturer
TEMOZOLOMIDE - Special Authority see SA1741 below - Reta	ail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ Temaccord
	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ Temaccord
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg		5	✓ Temaccord
•	688.00		✓ Amneal S29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Subsidy (Manufacturer		Fully idised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE -	Retail pharmacy-Specialist – Special Authority see SA1124 be	elow	
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRFTINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 bel	OW	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	✓ Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓ DBL Vinblastine S29
		✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓ Navelbine
210.00		✓ Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	✓ Baxter

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

continued...

Alecensa

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

for 6 months for applications meeting the following criteria:

Roth:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

✓ Sprycel	60	Tab 20 mg3,774.06
✓ Sprycel	60	Tab 50 mg6,214.20
✓ Sprycel	60	Tab 70 mg7,692.58

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA2000 below

✓ Tarceva	30	100 mg764.00	Tab 100 mg
✓ Tarceva	30	150 mg 1.146.00	Tab 150 mg

⇒SA2000 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued defitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
 - 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2001 below

Tab 250 mg1,700.00 30 ✓ Iressa

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fitha
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESII ATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460

	below	2,400.00	60	✓ Glivec
*	Cap 100 mg	58.23	60	✓ Imatinib-Rex
	3	98.00		✓ Imatinib-AFT
	Imatinib-Rex to be Sole Supply on 1 June 2021			
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex
		197.50		✓ Imatinib-AFT

Imatinib-Rex to be Sole Supply on 1 June 2021

(Imatinib-AFT Cap 100 mg to be delisted 1 June 2021)

(Imatinib-AFT Cap 400 mg to be delisted 1 June 2021)

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

continued...

PHARMAC Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Phone: (04) 460 4990

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.

The CML/GIST Co-ordinator

- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

⇒SA1191 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 1.2 The patient has not previously received trastuzumab treatment for HER 2 positive metastatic breast cancer; and
 - 1.3 Lapatinib not to be given in combination with trastuzumab; and
 - 1.4 Lapatinib to be discontinued at disease progression; or
- 2 All of the following:
 - 2.1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
 - 2.2 The patient started trastuzumab for metastatic breast cancer but discontinued trastuzumab within 3 months of starting treatment due to intolerance; and
 - 2.3 The cancer did not progress whilst on trastuzumab; and
 - 2.4 Lapatinib not to be given in combination with trastuzumab; and
 - 2.5 Lapatinib to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 on the next page - Retail pharmacy

 Wastage claimable
 4,680.00
 120
 ✓ Tasigna

 Cap 200 mg
 6,532.00
 120
 ✓ Tasigna

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable			
Cap 75 mg	4,000.00	21	Ibrance
Cap 100 mg	4,000.00	21	✓ Ibrance
Cap 125 mg	4,000.00	21	✓ Ibrance
, ,			

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓ ✓	
PAZOPANIB - Special Authority see SA1190 below - Retail pha	rmacy			
Tab 200 mg	1,334.70	30	✓	Votrient
Tab 400 mg	2,669.40	30	✓	Votrient

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5,000.00	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or 2.2 Both:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy;
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2002 below - Retail pharmacy

Cap 12.5 mg2,315.38	28	✓ Sutent
Cap 25 mg	28	✓ Sutent
Cap 50 mg	28	✓ Sutent

⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 87

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 4.2.2 Patient has ECOG performance score of 0-2; and
- 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg	1.36	10	✓ Calutide-50 S29
· ·	4.07	30	✓ Binarex
	4.21	28	✓ Binarex
Binarex to be Sole Supply on 1 April 2021			
FLUTAMIDE			
Tab 250 mg	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authorit	ty see SA1895 bel	ow	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL	ACFTATE
MEGEOTIOL	AULIAIL

Tab 160 mg	63.53	30	✓ Apo-Megestrol
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial	30.64	5	✓ Octreotide
			MaxRx S29
	56.87		✓ DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule	72.50	5	✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓ DBL Octreotide
	222.00		Octreotide
			(Sun) \$29

Fully

Subsidised

Brand or

Generic

Subsidy

(Manufacturer's Price)

	\$	Per	✓ Manufacturer
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Spi	ecial Authority see SA20	004 below -	Retail pharmacy
Inj LAR 10 mg prefilled syringe	1,772.50	1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe	2,358.75	1	Sandostatin LAR
Inj LAR 30 mg prefilled syringe		1	Sandostatin LAR

⇒SA2004 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery: or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:

	Subsidy (Manufacturer's		Fı Subsidis	ully	Brand or Generic	
	\$	P	er	✓	Manufacturer	
continued						
5.1	Carcinoid syndrome (diagnosed by tissue pathology and/or urina	ry 5HIAA	analysis);	and		

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

TAMOXIFEN CITRATE

*	Tab 10 mg15.00	60	✓ <u>Tamoxifen Sandoz</u>
*	Tab 20 mg6.65	60	✓ <u>Tamoxifen Sandoz</u>

Aromatase Inhibitors

ANASTROZOLE			
* Tab 1 mg4	.55	30	✓ Anatrole
5	.04		✓ Rolin
Anatrole to be Sole Supply on 1 April 2021			
(Rolin Tab 1 mg to be delisted 1 April 2021)			
EXEMESTANE			
* Tab 25 mg14	.50	30	✓ Pfizer Exemestane
LETROZOLE			
* Tab 2.5 mg4	.68	30	✓ Letrole

Immunosuppressants

AZATI HODDINE

Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg7.35	60	Azamun
* Tab 50 mg7.60	100	✓ Azamun
* Inj 50 mg vial199.00	1	✓ Imuran
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	✓ Cellcept
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

etail pharmacy		
690.00	4	✓ Enbrel
690.00	4	✓ Enbrel
1,050.00	4	✓ Enbrel
1,050.00	4	✓ Enbrel
	etail pharmacy 690.00 690.00 1,050.00 1,050.00	

⇒SA1974 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

Fither:

- 1 Both: 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD): or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD: or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
 - 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for psoriatic arthritis; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price)			Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

2.6 Either:

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

2.7 Either:

- 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
- 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - S	pecialist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only - Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG \$29 Ini 40 mg per ml vial to be delisted 1	April 2022)		

 "			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	` ¢ ′	Por ./	Manufacturar

Monoclonal Antibodies

ADALIMUMAB - Special Authority see SA1975 below - Retail pharmacy

Inj 20 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Ini 40 mg per 0.8 ml prefilled syringe	1.599.96	2	✓ Humira

⇒SA1975 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and

- 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
- 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment: or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Fither
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less: or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plague psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	Full		
 (Manufacturer's Price) \$	Subsidise Per 🗸	d Generic Manufacturer	

continued...

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 below

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA1982 below

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA1982 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses: or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria: Fither:

- - 1 A withdrawal period has been tried and the patient has relapsed; or
 - 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis; or

Subsidy (Manufacturer's Price)	Fu Subsidis	,	and or eneric
 \$	Per	✓ M	anufacturer

continued...

- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis; or
- 2.12 Neurosarcoidosis: or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Eithor
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:

Subsidy	Fully	/ Brand or
(Manufacturer's Price)	Subsidised	d Generic
 \$	Per 🗸	Manufacturer

continued...

- 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
- 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fullminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 The patient has had a good clinical response following 3 initial doses; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Au	uthority see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	•	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

Inj 100 mg per 10 ml vial	1,075.50 2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30 1	✓ Mabthera
Inj 1 mg for ECP	5.64 1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept: or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - S	special Authority see SA2028 b	elow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2028 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Fither

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months: or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

- All of the following:
 - 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective: and
 - 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
 - 3 Genetic causes of nephrotic syndrome have been excluded; and
 - 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Fu Subsidis	,	and or eneric
 \$	Per	✓ M	anufacturer

continued...

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has had a rituximab treatment-free interval of 12 months or more; and

Sul	bsidy Fu	ully Brand of	or
	,	,	
(Manufact	turer's Price) Subsidis	ed Generio	2
(· M	_4
	\$ Per	Manufa	icturer

continued...

- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Fither:
 - 2.1 Both:

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
- 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
- 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Roth:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*: and
- 2 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
- 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

SECUKINUMAB – Special Authority see SA1754 below – Retail pharmacy
Inj 150 mg per ml, 1 ml prefilled syringe........1,599.00 2 ✓ Cosentyx

⇒SA1754 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1977 below

✓ Actemra	1	4 ml vial	lnj 20 mg per ml, 4 ml vial
✓ Actemra	1	10 ml vial550.00	Inj 20 mg per ml, 10 ml vial
✓ Actemra	1	20 ml vial1,100.00	
✓ Baxter	1 ma	2.85	Ini 1 mg for ECP

SA1977 | Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 All of the following:

- 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
- 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
- 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or

2 All of the following:

- 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
- 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
- 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

(Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses. **Initial application — (previous use)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules: and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy: and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

6 Either:

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Roth:

1.1 Fither:

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule: and

1.2 Fither:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

		•	
Inj 150 mg vial	1,350.00	1	Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce) Si	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — **(early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
- 3.2.2 The cancer did not progress whilst on lapatinib; or
- 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial	2,320.00	1	✓ Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	23.20	1 mg	✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*: or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

IIVC	DLUMAB - PCT	only - Specialist - Special Authority	y see SA2006 on the next pa	age	
- 1	nj 10 mg per ml	, 4 ml vial	1,051.98	1	Opdivo
		, 10 ml vial		1	✓ Opdivo
	, , ,	,) 	·	1 mg	✓ Baxter
				-	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Either:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
must have reduction in short axis to < 10 mm.

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Either:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease

 Subsidy	Fully		Brand or
Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

progression; and

- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS – Special Authority see SA2008 below – R Wastage claimable	etail pharmacy		
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor
OACOOO On a stat Austh and to day Out at the			

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA2005 on the next page - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease: and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 a	bove – Retail pharmacy	
Initiation kit - 5 vials freeze dried venom with diluent	5.00 1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent305	5.00 1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		
diluent285	5.00 1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		
9 ml, 3 diluent 1.8 ml305	5.00 1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305	5.00 1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA1367	above – Retail pharma	су
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml305	5.00 1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305	5.00 1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		
dried venom, with diluent	5.00 1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		
dried venom, with diluent	5.00 1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml305	5.00 1 OP	✓ Albey
).00 TOP	Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent305	5.00 1 OP	✓ Venomil \$29
unca venom, with anaent		· Venonin ozo

	Subsidy		Fully Brand or
	(Manufacturer's Price	e) Subs	idised Generic
	\$	Per	✓ Manufacturer
Autibistanius			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral liq 1 mg per ml		200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE		200	
	0.07	500 ml	✓ Histafen
* Oral liq 2 mg per 5 ml	9.37	300 1111	▼ nistaleli
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg		40	
	(8.40)		Polaramine
	1.01	20	
	(5.99)		Polaramine
* Oral liq 2 mg per 5 ml		100 ml	5
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
	(8.23)		Telfast
* Tab 120 mg	4.74	10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg	1.69	100	✓ Lorafix
* Oral liq 1 mg per ml		120 ml	✓ Lorfast
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1 68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	✓ Hospira
and the state of t			
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose		00 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		00 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	00 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
, 			Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's	Price) Subs	idised Generic
	\$	Per	✓ Manufacturer
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	sts		
EFORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose dev	rice20.64	60 dose	
· · · · · · · · · · · · · · · · · · ·	(35.80)		Foradil
EFORMOTEROL FUMARATE DIHYDRATE	(55.55)		
Powder for inhalation 4.5 mcg per dose, breath activated	-) 10.00	CO dana OD	
(equivalent to eformoterol fumarate 6 mcg metered dos	,	60 dose OP	Orio Trutubalas
	(16.90)		Oxis Turbuhaler
INDACATEROL			
Powder for inhalation 150 mcg		30 dose OP	Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	✓ Serevent Accuhaler
Inhalad Andra at and the with Land Anton Bata	A .l		
Inhaled Corticosteroids with Long-Acting Beta-	-Aarenocept	or Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide v	with		
6 mcg eformoterol fumarate metered dose)	41.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma	rate		
per dose (equivalent to 400 mcg budesonide with 12 mc			
eformoterol fumarate metered dose) - No more than 2	J		
dose per day	82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	18.23	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6	mcg33.74	120 dose OP	✓ Symbicort
ř	-		Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40	120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6		120 dose OP	✓ Symbicort
ř	-		Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	44.08	60 dose OP	✓ Symbicort
, ,			Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	<i>44</i> 08	30 dose OP	✓ Breo Ellipta
1 5 macri for initialiation 100 mag with vitalitorol 25 mag		50 dosc OI	- Dieo Empla

	Subsidy		Fully	Brand or
	(Manufacturer's \$	Price) Subsi Per	dised ✓	Generic Manufacturer
THE CASCALE WITH CALMETERS!	Ψ	rei		Ivianulacturei
LUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.70	120 dose OP	√ So	retide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP		retide
Powder for inhalation 100 mcg with salmeterol 50 mcg – No		120 0000 01	• 00	Tottao
more than 2 dose per day		60 dose OP	✓ Se	retide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No		00 0000 01	- 00	retide Addunater
more than 2 dose per day		60 dose OP	✓ Se	retide Accuhaler
, ,		00 0000 0.		
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml		150 ml	✓ Ve	<u>ntolin</u>
Infusion 1 mg per ml, 5 ml		10	✓ Ve	ntolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	53.00	5	✓ Ve	ntolin
Inhaled Beta-Adrenoceptor Agonists				
ALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000				
dose available on a PSO	3.80	200 dose OP	✓ Ro	spigen
dose available off a f 30		200 dose Oi	✓ Sa	
	(6.00)			ntolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb			•••	110111
available on a PSO		20	✓ As	thalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO		20	✓ As	thalin
ERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22 20	120 dose OP	✓ Br	icanyl Turbuhaler
200 mag motorou accop, croadir activatou		120 0000 01	- 5	iodily: runbundioi
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	✓ At	rovent
a) Up to 400 dose available on a PSO				
b) No patient co-payment payable				
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne	eb			
available on a PSO		20	✓ <u>Un</u>	<u>ivent</u>
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg	ner			
dose CFC-free		200 dose OP	✓ Du	olin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per		200 0000 01	. 50	
vial, 2.5 ml ampoule – Up to 20 neb available on a PSC	5 20	20	✓ Du	ıolin
Trai, 2.0 mi amposio Op to 20 nob available on a 1 oc	,	20	- 50	· · · · · · ·

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Seebri Breezhaler

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

30 dose Spiriva Soln for inhalation 2.5 mcg per dose50.37 60 dose OP ✓ Spiriva Respimat

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Incruse Ellipta

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg......81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP ✓ Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg......2,554.00 60 OP ✓ Ofev 60 OP ✓ Ofev

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

Leukotriene Receptor Antagonists

МО	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg3.95	28	✓ Montelukast Mylan

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

(Tilade Aerosol inhaler, 2 mg per dose CFC-free to be delisted 1 September 2021)

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

Methylxanthines

AMINOPHYLLINE

* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj availabl		5	✓ DBL Aminophylline
THEOPHYLLINE			,
* Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
* Oral liq 80 mg per 15 ml	16.60	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA – Special Authority see SA1978 below – Reta	ail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
IVACAFTOR - PCT only - Specialist - Special Authority see SA	2017 below			
Tab 150 mg	29,386.00	56	✓	Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓	Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓	Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

DUDECONIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

Metered aqueous nasal spray, 50 mcg per dose	200 dose OP 200 dose OP	
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%5.23 Univent to be Sole Supply on 1 April 2021	15 ml OP	✓ Univent

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

mall.......2.20 1 ✓ e-chamber Mask

	Subsidy	_	Fully	Brand or
	(Manufacturer's Price) \$	Su Per	bsidised •	Generic Manufacturer
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	lini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)		1	•	-chamber Turbo
510 ml (single patient)	5.12	1	✓ e	-chamber La Grande
800 ml	6.50	1	✓ V	olumatic
Respiratory Stimulants				
CAFFEINE CITRATE				
Oral liq 20 mg per ml (10 mg base per ml)	15.10 2	5 ml OP	✓ <u>B</u>	iomed

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	ica) Su	bsidised	Generic
	\$	Per	Joiulocu	Manufacturer
	Ψ	r ei		Manuacturei
Ear Preparations				
ACETIC ACID WITH A C. PROPANIEDICI PIACETATE AND RE	NIZETLIONILINA			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE				
For Vosol ear drops with hydrocortisone powder refer Standa	ard Formulae, <mark>pag</mark>	je 239		
Ear drops 2% with 1, 2-Propanediol diacetate 3% and				
benzethonium chloride 0.02%	6.97	35 ml OP	✓ V	/osol
		00 1111 01	•	0301
FLUMETASONE PIVALATE				
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	√ L	.ocacorten-Viaform
				ED's
			./ 1	.ocorten-Vioform
			V L	.ocorten-violoriii
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate				
	F 40	7.5		
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	• K	(enacomb
Ear/Eye Preparations				
, ,				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
	4.50	0 1 OD		
gramicidin 50 mcg per ml		8 ml OP	_	
	(9.27)		S	Sofradex
FRAMYCETIN SULPHATE				
Ear/Eve drops 0.5%	4 10	8 ml OP		
Ear/Eye drops 0.5%		8 IIII OP		
	(8.65)		5	Soframycin
Eye Preparations				
Eye preparations are only funded for use in the eye, unless explicit	oitly stated athorn	ico		
Lye preparations are only funded for use in the eye, unless expin	citiy stated official	156.		
Anti-Infective Preparations				
Anti-infective r reparations				
ACICLOVIR				
	44.00	4.5 0.0		//P00
* Eye oint 3%	14.92	4.5 g OP	• 1	/iruPOS
CHLORAMPHENICOL				
Eye oint 1%	1 55	5 g OP	√ Γ	Devatis
		10 ml OP	_	Chlorafast
Eye drops 0.5%			• •	illoralast
Funded for use in the ear*. Indications marked with * are	e unapproved ind	ications.		
CIPROFLOXACIN				
Eye drops 0.3% - Subsidy by endorsement	12 15	5 ml OP	10	Ciprofloxacin Teva
				•
When prescribed for the treatment of bacterial keratitis o				
for the second line treatment of chronic suppurative otitis		; and the pre	escription	is endorsed accordingly.
Note: Indication marked with a * is an unapproved indic	ation.			
GENTAMICIN SULPHATE				
	11 40	E ml OD		Conontio
Eye drops 0.3%	11.40	5 ml OP	• (Genoptic
PROPAMIDINE ISETHIONATE				
* Eye drops 0.1%	2 97	10 ml OP		
= -j = 3/0 po 0.1 /0		10 1111 01		Brolene
	(14.55)		Е	NOIGHE
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	√ F	ucithalmic
, ,		ŭ		

	Subsidy (Manufacturer's F	Price)	Fully Subsidised	
	\$	Pe		
OBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g	OP 🗸	Tobrex
Eye drops 0.3%	11.48	5 ml (OP 🗸	Tobrex
Corticosteroids and Other Anti-Inflammatory Pro	eparations			
EXAMETHASONE				
Eye oint 0.1%		3.5 g	•.	Maxidex
F Eye drops 0.1%	4.50	5 ml (OP 🗸	Maxidex
Ocular implant 700 mcg - Special Authority see SA1680 belo)W			
- Retail pharmacy	4 444 50	4	./	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family: and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b		
	sulphate 6,000 u per g5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM		
	Eye drops 0.1%	5 ml OP	✓ Voltaren Ophtha

233

	Subsidy (Manufacturer's Price)	Subs	Fully sidised	Brand or Generic Manufacturer
FLUOROMETHOLONE	3	Per		Manufacturer
* Eye drops 0.1%	3.09 5 5.20	5 ml OP	√ F	ML Tucon
KETOROLAC TROMETAMOL – Special Authority see SA1981 Eye drops 0.5%		acy 5 ml OP	✓ A	Acular

⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71	4 ml OP	
• , •	(10.34)		Livostin
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eye drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
Eye drops 1%	5.93	10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Auth	ority see SA1715 below	- Retail pharn	nacy
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	✓ Minims
			Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

Eye drops 2%	1.79	5 ml OP	Rexacrom
Lye 010p3 2 /6	1.1 J	3 1111 01	• ITCAACIO

Glaucoma Preparations - Beta Blockers

BE	TAXOLOL		
*	Eye drops 0.25%	5 ml OP	✓ Betoptic S
	Eye drops 0.5%	5 ml OP	✓ Betoptic
TIN	MOLOL		
*	Eye drops 0.25%1.81	5 ml OP	✓ Arrow-Timolol
	Eye drops 0.5%	5 ml OP	✓ Arrow-Timolol
*	Eye drops 0.5%, gel forming	2.5 ml OP	✓ Timoptol XE

	Subsidy (Manufacturer's Pric \$	ce) Subsi Per	Fully Brand or idised Generic ✓ Manufacturer
Glaucoma Preparations - Carbonic Anhydrase Ir	hibitors		
ACETAZOLAMIDE	17.00	100	. Diamou
* Tab 250 mg	17.03	100	✓ Diamox
BRINZOLAMIDE k Eye drops 1%	9 77	5 ml OP	✓ Azopt
ORZOLAMIDE HYDROCHLORIDE		0 1111 01	7,2001
Eye drops 2%	9.77	5 ml OP	
	(17.44)		Trusopt
OORZOLAMIDE WITH TIMOLOL			
F Eye drops 2% with timolol 0.5%	2.87	5 ml OP	✓ Dortimopt
Glaucoma Preparations - Prostaglandin Analogu	ies		
BIMATOPROST			
€ Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost
			Multichem
ATANOPROST			. _
€ Eye drops 0.005%	1.57	2.5 ml OP	✓ <u>Teva</u>
RAVOPROST	7.00	5l OD	/ T
k Eye drops 0.004%	10.50	5 ml OP	✓ Travopt ✓ Mylan S29
	19.50	2.5 ml OP	✓ Mylan 323
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE			
F Eye drops 0.2%	12.25	5 ml OP	✓ Arrow-Brimonidine
RIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
Fye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
ILOCARPINE HYDROCHLORIDE			
₭ Eye drops 1%		15 ml OP	✓ Isopto Carpine
Eye drops 2%		15 ml OP	✓ Isopto Carpine
Eye drops 4%		15 ml OP	✓ Isopto Carpine
Eye drops 2% single dose — Special Authority see SA0895			
below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine
⇒SA0895 Special Authority for Subsidy			•
nitial application from any relevant practitioner. Approvals valid	for 2 years for ap	plications me	eting the following criteria:
ither:		-	- -
1 Patient has to use an unpreserved solution due to an allero	v to the preserva	tive: or	

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE

15 ml OP ✓ Atropt

	Subsidy (Manufacturer's F \$	Price) Subs	Fully Brand or idised Generic Manufacturer
* Eye drops 1% * Eye drops 1%, single dose (preservative free) – Only on a prescription		15 ml OP 20 dose	✓ Cyclogyl ✓ Minims Cyclopentolate
TROPICAMIDE * Eye drops 0.5% * Eye drops 1%		15 ml OP 15 ml OP	✓ Mydriacyl ✓ Mydriacyl
Preparations for Tear Deficiency			
For acetylcysteine eye drops refer Standard Formulae, page 23 HYPROMELLOSE	9		
* Eye drops 0.5%	2.00 (3.92)	15 ml OP	Methopt
HYPROMELLOSE WITH DEXTRAN * Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ Poly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA1388 above – Retail pha	rmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authori	ty see SA1388 ab	ove – Retai	l pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Author	ority see SA1388	above – Re	tail pharmacy
Eye drops 1 mg per ml	22.00	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pha month is not relevant and therefore only the prescribed do			
month is not relevant and therefore only the prescribed di	bage to the hear	cot Or may	be damied.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE		
Eye drops 0.1%2.20	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT		
* Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE		
Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS

				VARIOUS
	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
Various				
PHARMACY SERVICES May only be claimed once per patient. * Brand switch fee	4.50	1 fee	✓ B\$	SF Ambrisentan Mylan SF Hydroxycarbamide Devatis
a) The Pharmacode for BSF Hydroxycarbamide Devatis b) The Pharmacode for BSF Ambrisentan Mylan is 260 (BSF Ambrisentan Mylan Brand switch fee to be delisted 1 July 2 (BSF Hydroxycarbamide Devatis Brand switch fee to be delisted	5309 - see also <mark>pag</mark> 2021)		55	
Agents Used in the Treatment of Poisonings				
Antidotes				
ACETYLCYSTEINE Inj 200 mg per ml, 10 ml ampoule	58.76	10	✓ Ma	BL Acetylcysteine artindale Pharma S29
NALOXONE HYDROCHLORIDE a) Up to 5 inj available on a PSO b) Only on a PSO	00.00	-	4 D	DI Malaura
* Inj 400 mcg per ml, 1 ml ampoule	22.60	5	_	<u>BL Naloxone</u> Hydrochloride
Removal and Elimination				
CHARCOAL * Oral liq 50 g per 250 ml a) Up to 250 ml available on a PSO b) Only on a PSO	43.50 2	50 ml OP	✓ Ca	arbosorb-X

С

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or



Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below -	 Retail pharmacy 		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> <u>Desferrioxamine</u> <u>Mesylate for Inj</u> BP
SODIUM CALCIUM EDETATE			_
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium
			Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol	60 mg 40 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Preservative Water	qs to 100 ml	Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
CODEINE LINCTUS (15 mg per 5 ml)		Water	10 40 1111
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol Preservative	40 ml	Pilocarpine 4% eye drops Preservative	qs
Water	qs to 100 ml	Water	qs to 500 ml
FOLINIC MOUTHWASH Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 500 ml	Methylcellulose	5 g
(Preservative should be used if quantity supplied is	for more	Preservative	qs
than 5 days. Maximum 500 ml per prescription.)		Water (Preservative should be used if quantity supplied is	to 500 ml
MAGNESIUM HYDROXIDE 8% MIXTURE		than 5 days. Maximum 500 ml per prescription.)	oi illoie
Magnesium hydroxide paste 29%	275 g	, , , , ,	
Methyl hydroxybenzoate	1.5 g	SODIUM CHLORIDE ORAL LIQUID	
Water	to 1,000 m	Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatra	
Methadone powder	qs		,
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	10:- -
Water	to 100 ml	Vancomycin 500 mg injection Glycerol BP	10 vials 40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridiu	
Propylene glycol	to 100 ml	following metronidazole failure)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	id mixture)	VOSOL EAR DROPS	
OMEPRAZOLE SUSPENSION		WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder	qs	Hydrocortisone powder	1%
Sodium bicarbonate powder BP	8.4 g	Vosol Ear Drops	to 35 ml
Water	to 100 ml	·	

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Extemporaneously Compounded Preparations and Galenicals

CODEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing	frequency	
Powder - Only in combination	63.09	25 g	
	(90.09)	-	Douglas
Only in extemporaneously compounded codeine linctus.			
COLLODION FLEXIBLE			

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

100 ml ✓ PSM

COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

Soln30.00 100 ml Midwest

GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus.

473 ml ✓ Ora-Sweet SF

GLYCERIN WITH SUCROSE - Only in combination

Only in combination with Ora-Plus.

473 ml ✓ Ora-Sweet

GI YCFROI

500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations.

METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency

d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

1 GWGCI		ı y	* 711
METHYL HYDROXYBENZOATE Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE		·	
Powder	36.95	100 g	✓ MidWest
Suspension – Only in combination		473 ml	✓ Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA		combination	
Suspension	,	473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only			<u> </u>
		473 ml	✓ Ora-Blend
Suspension	30.95	4/3 1111	▼ <u>Ora-Diellu</u>
PHENOBARBITONE SODIUM			
Powder – Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years			
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenzo	oate 10% solutio	n.	
Lig		500 ml	✓ Midwest
SODIUM BICARBONATE			
Powder BP - Only in combination	10.05	500 g	✓ Midwest
Only in extemporaneously compounded omeprazole and			· INITIANCS!
Only in extemporaneously compounded offeprazole and	iansoprazole su	isperision.	

1 a

✓ AFT

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	✓ <u>M</u>	idwest	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 243

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacture

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

(Glucerna Select RTH Liquid to be delisted 1 September 2021)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
. , ,	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Resource Diabetic
	(2.10)		Sustagen Diabetic

(Glucerna Select Liquid (vanilla) to be delisted 1 September 2021) (Resource Diabetic Liquid (vanilla) to be delisted 1 May 2021)

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 245



Subsidy (Manufacturer's Price) Sub

Subsidised

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

 Liquid
 54.00
 400 g OP
 ✓ Kindergen

 Powder
 54.00
 400 g OP
 ✓ Kindergen

(Kindergen Liquid to be delisted 1 August 2021)

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or

	Subsidy	F	ully	Brand or
(Man	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	•	Manufacturer

continued...

- 2.3 faltering growth in an infant/child; or
- 2.4 increased nutritional requirements; or
- 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see \$ Liquid		he previous pag 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA Liquid		previous page 500 ml OP	− Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Apharmacy [HP3]	Authority see	SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	379 on the r	revious page -	Hospital pharmacy [HP3]
Liquid (strawberry)		200 ml OP	✓ Fortini
Liquid (vanilla)		200 ml OP	✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA137	9 on the pre	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)		200 ml OP	✓ Pediasure
Liquid (strawberry)		200 ml OP	✓ Pediasure
Liquid (vanilla)		200 ml OP	✓ Pediasure
, ,	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authorharmacy [HP3]	ority see SA	1379 on the pre	evious page - Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)		200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the	e previous r	age - Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 247

	Subsidy (Manufacturer's Pri \$	ce) Subs	Fully idised	Brand or Generic Manufacturer
RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid		evious page – 500 ml OP		al pharmacy [HP3] lepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		us page – Hos 220 ml OP	✓ N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110	1 on the previous	page – Hospi	tal pha	rmacy [HP3]
Liquid (apricot) 125 ml	(3.31) 11.52	237 ml OP 4 OP	✓ F	lovaSource Renal
Liquid (caramel) 125 ml	11.52	4 OP	√ F	Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LiquidLiquid	,					
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see	e SA1377 above	- Hospital phar	macy [HP3]			
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above –	Hospital pharma	acy [HP3]			
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN			
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]						
Liquid	12.04	1.000 ml OP	✓ Peptisorb			

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 249



Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
 \$	Per	•	Manufacturer

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 249 - Hospital pharmacy [HP3]

Liquid7.00 1,000 ml OP ✓ Nutrison Energy

✓ fully subsidised 251

	Subsidy		Fully Brand or
	(Manufacturer's		sidised Generic
	\$	Per	✓ Manufacturer
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	page 249 – Ho	spital pharmacy	/ [HP3]
Liquid	1.24	250 ml OP	✓ Isosource Standard
	5.29	1,000 ml OP	Nutrison Standard
			RTH
			✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority	see SA1859 o	on page 249 – H	Hospital pharmacy [HP3]
Liquid	5.29	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority se	e SA1859 on p	oage 249 – Hos	pital pharmacy [HP3]
Liquid	5.29	1,000 ml OP	✓ Jevity ŘŤH
·			✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s	see SA1859 on	nage 249 – Ho	spital pharmacy [HP3]
Liquid		250 ml OP	✓ Ensure Plus HN
<u> </u>	7.00	1,000 ml OP	✓ Ensure Plus RTH
	7.00	1,000 1111 01	✓ Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
ODAL FEED (DOWNER) Chariel Authority and CA18EO on nor	a 040 - Llaanit	الاليومسوطواه	
ORAL FEED (POWDER) - Special Authority see SA1859 on pag		, , , ,	•
Note: Higher subsidy for Sustagen Hospital Formula will only	be reimbursed	i for patients wit	in both a valid Special Authorit
number and an appropriately endorsed prescription.			
Powder (chocolate) – Higher subsidy of up to \$26.00 per 850 with Endorsement		050 ~ OD	✓ Ensure
with Endorsement		850 g OP	Ensure
	9.54	840 g OP	0
	(26.00)		Sustagen Hospital
A delikional and aids by and are one of it and labor for a stion			Formula Active
Additional subsidy by endorsement is available for patien prescription must be endorsed accordingly.	is with fat maia	ibsorption, iai ir	itolerance or chyle leak. The
Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g	0.54	057 - 00	/ Football
with Endorsement		857 g OP	✓ Fortisip
	26.00	850 g OP	✓ Ensure
	9.54	840 g OP	Occation and House's 1
	(26.00)		Sustagen Hospital
A 1 1991			Formula Active
Additional subsidy by endorsement is available for patient	ts with fat mala	apsorption, fat in	itolerance or chyle leak. The

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

(Fortisip Powder (vanilla) to be delisted 1 August 2021)

(Mar	Subsidy nufacturer's Price)	Ful Subsidise	,	Brand or Generic
(Iviai			_	
	\$	Per •	/	Manutacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 249 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with	` ,		·
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see SA1859 on page 249 – Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with

Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

✓ fully subsidised 253

	Subsidy	Fully	Brand or
(Manut	facturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer	
FOOD THICKENER - Special Authority see SA1106 on the p			,		
Powder	6.53 3	00 g OP	✓ N	lutilis	
	7.25 3	80 g OP		eed Thickener Karicare Aptamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX - Special Authority see SA1729 above - Hospita Powder2.81	l pharmacy [HP3] 1,000 g OP	
(5.15)		Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 above - Hospital	pharmacy [HP3]	
Powder	1,000 g OP	
(7.32)		NZB Low Gluten Bread Mix
3.51		
(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR – Special Authority see SA1729 above – Hospital pha	rmacy [HP3] 2,000 g OP	
(18.10)	_, 9 0.	Horleys Flour

✓ fully subsidised 255

	Subsidy		Fully	Brand or
	(Manufacturer's Prices)	ce) Sub Per	sidised	Generic Manufacturer
	*			
GLUTEN FREE PASTA – Special Authority see SA1729 on the			macy [HF	23]
Buckwheat Spirals		250 g OP	_	
	(3.11)		0	rgran
Corn and Vegetable Shells		250 g OP		
	(2.92)		0	rgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		0	rgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		0	rgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)	-	0	rgran
Rice and Corn Penne	2.00	250 g OP		•
	(2.92)	•	0	rgran
Rice and Maize Pasta Spirals	2.00 [°]	250 g OP		
·	(2.92)	Ū	0	rgran
Rice and Millet Spirals	2.00 [′]	250 g OP		
•	(3.11)	J	0	rgran
Rice and corn spaghetti noodles	` '	375 g OP		3
1	(2.92)		0	rgran
Vegetable and Rice Spirals	` '	250 g OP		3 ··
- g - · · · · · · · · · · · · · · · · · ·	(2.92)		0	rgran
Italian long style spaghetti	` '	220 g OP	·	· J· · · ··
	(3.11)		0	rgran
	(3.11)		·	. 3

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING I	MIX – Special Authority see SA1108 on the previous	s page – Hospital	pnarmacy [HP3]
Powder	8.22	500 g OP	Loprofin Mix
LOW PROTEIN PASTA _	- Special Authority see SA1108 on the previous page	– Hospital pharm	nacy [HP3]

LOW PROTEIN PASTA - Special Authority see SA11	08 on the previous page - I	Hospital pharm	acy [HP3]
Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni		250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirale	11.01	500 a OP	✓ Lonrofin

✓ fully subsidised 257

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see SA194 Powder	' '	nacy [HP3] 400 g OP	✓ Alfamino Junior
Powder (unflavoured)		400 g OP	✓ Elecare
		-	✓ Elecare LCP
			Neocate Gold
			 Neocate Junior Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ Elecare
			Neocate Junior Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short aut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	/	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

1 Fither:

1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

✓ fully subsidised 259

SPECIAL FOODS

-			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut: or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	 Special Authority see SA1953 below 	/ – Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption: or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

Powder	450 g OP	✓ Aptamil Gold+ Pepti Junior
30.42	900 g OP	✓ Aptamil AllerPro SYNEO 1
		 Aptamil AllerPro SYNEO 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

✓ fully subsidised 261



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > 0 requal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
 Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE				
	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AI	ND HAEMOPHILUS	INFLUENZ	AE TY	PE B VACCINE -
[Xpharm] Funded for patients meeting any of the following criteria:				
Up to four doses for children up to and under the age of	10 for primary immu	nisation: o	r	
2) An additional four doses (as appropriate) are funded for 10 who are patients post haematopoietic stem cell trans post solid organ transplant, renal dialysis and other seve 3) Up to five doses for children up to and under the age of Note: A course of up-to four vaccines is funded for catch up process.	(re-)immunisation fo plantation, or chemo erely immunosuppres 10 receiving solid or	r children u therapy; pu ssive regim gan transp	ip to ar e or po ens; or antatio	est splenectomy; pre- or n.
to complete full primary immunisation. Please refer to the Improgrammes.				
Inj 30IU diphtheriatoxoid with 40IU tetanustoxoid, 25mcg pertussistoxoid, 25mcg				
pertussisfilamentoushaemagglutinin, 8 mcgpertactin, 80 D-AgUpoliovirus, 10mcghepatitisBsurfaceantigen in 0.5ml syringe	0.00	10	√ <u>In</u>	fanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm] One dose for patients meeting any of the following:				
 For primary vaccination in children; or An additional dose (as appropriate) is funded for (re-)im transplantation, or chemotherapy; functional asplenic; p or post cochlear implants, renal dialysis and other sever For use in testing for primary immunodeficiency disease paediatrician. 	re or post splenector ely immunosuppress	ny; pre- or sive regime	post so ns; or	olid organ transplant, pre-
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg prefilled syringe plus vial 0.5 ml		1	✓ H	iberix
HEPATITIS A VACCINE - [Xpharm]				
Funded for patients meeting any of the following criteria:				
 Two vaccinations for use in transplant patients; or Two vaccinations for use in children with chronic liver di One dose of vaccine for close contacts of known hepatit 	,			
Inj 1440 ELISA units in 1 ml syringe	0.00	1	✓ H	avrix
Inj 720 ELISA units in 0.5 ml syringe		1		avrix Junior

	NATIONA	L HVHVI	IUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Pric \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xphar Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the folk	0.00	1	√ E	Engerix-B
 for household or sexual contacts of kr for children born to mothers who are l for children up to and under the age of serology and require additional vaccin for HIV positive patients; or for patients following non-consensual for patients following immunosuppres for solid organ transplant patients; or for post-haematopoietic stem cell tran following needle stick injury. 	nown acute hepatitis B patients on the patitis B surface antigen (HBs) of 18 years inclusive who are contaction or require a primary cours sexual intercourse; or sion; or	Ag) pos nsidered	itive; or I not to have	e achieved a positive
Inj 20 mcg per 1 ml prefilled syringe	owing criteria: nown acute hepatitis B patients on patients on the patitis B surface antigen (HBs of 18 years inclusive who are contaction or require a primary cours sexual intercourse; or sion; or	Ag) pos nsidered	titis B carrier itive; or I not to have	e achieved a positive
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, Any of the following: 1) Maximum of two doses for children aged 2) Maximum of three doses for patients mee 1) People aged 15 to 26 years inclusive 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell 3) Maximum of four doses for people aged 9	14 years and under; or ting any of the following criteria: e; or) patients: or to 26 years inclusive post chem		·	
Inj 270 mcg in 0.5 ml syringe	0.00	10	√ <u>c</u>	Gardasil 9

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Sul	Fully osidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10	✓ A	fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- c) children aged four years or less (but over three years) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine)......90.00

(2021 Formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

A) INFLUENZA VACCINE - people 65 years and over

is available each year for patients aged 65 years and over

- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

	NATIONAL	IMMUNIS	SATIC	ON SCHEDULE
(Man	Subsidy ufacturer's Price) \$	Subsid Per	Fully dised	Brand or Generic Manufacturer
MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VA Either:	CCINE – [Xpha	arm]		
A) Any of the following:				
 Up to three doses and a booster every five years for pa functional or anatomic asplenia, HIV, complement defic transplant; or One dose for close contacts of meningococcal cases; or 	iency (acquired r			
 A maximum of two doses for bone marrow transplant pa A maximum of two doses for patients following immuno 		r		
B) Both:	suppression, o	'		
Person is aged between 13 and 25 years, inclusive; and Either:	t			
i) One dose for individuals who are entering within the boarding school hostels, tertiary education halls of ii) One dose for individuals who are currently living in residence, military barracks, or prisons, from 1 De	f residence, mili n boarding scho	tary barracl	ks, or p tertiary	orisons; or education halls of
Note: children under seven years of age require two doses 8 weel	ks apart, a boos	ter dose the	ree yea	ars after the primary
series and then five yearly.				
*Immunosuppression due to steroid or other immunosuppressive the	nerapy must be	for a period	d of gre	ater than 28 days.
Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier				
per 0.5 ml vial	0.00	1	✓ <u>M</u> e	enactra
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both:				
 The child is under 9 months of age; and Any of the following: 				
 Up to three doses for patients pre- and post splenector HIV, complement deficiency (acquired or inherited), or present the complement of the co	ore or post solid or atients; or	organ tran		
Note: children under nine months of age require two doses 8 booster schedules with meningococcal ACWY vaccine.			: Immu	nisation Handbook for
*Immunosuppression due to steroid or other immunosuppres	sive therapy mu	st be for a	period	of greater than 28 days
Inj 10 mcg in 0.5 ml syringe	0.00	1	✓ Ne	eisvac-C
PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]				
1) A primary course of three doses for previously unvaccinated	individuals up to	the age of	59 mo	nths inclusive
Note: please refer to the Immunisation Handbook for the appropria	ate schedule for	catch up p	rogram	ımes
Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal				
polysaccharide serotypes 4, 18C and 19F in 0.5 ml prefilled syringe	0.00	10	✓ <u>S</u> y	<u>/nflorix</u>

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously
 received two doses of the primary course of PCV10: or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISAT	ION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE	– [Xpharm]		
Either:			
 Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with funcomplement deficiency (acquired or inherited), coch All of the following: 	ctional asplenia, pre- or p	oost-solid organ	transplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)immur b) Treatment is for a maximum of two doses; and c) Any of the following: 	· ·		
 i) on immunosuppressive therapy or radiati immune response; or 	ion therapy, vaccinate wl	nen there is expe	ected to be a sufficient
ii) with primary immune deficiencies; oriii) with HIV infection; or			
iv) with renal failure, or nephrotic syndrome;	or		
v) who are immune-suppressed following or		uding haematop	oietic stem cell transplant);
or vi) with cochlear implants or intracranial shu	inte: or		
vii) with cerebrospinal fluid leaks; or	into, or		
viii) receiving corticosteroid therapy for more			
prednisone of 2 mg/kg per day or greater	r, or children who weigh	more than 10 kg	on a total daily dosage of
20 mg or greater; or ix) with chronic pulmonary disease (includin	g asthma treated with hig	ah-dose corticos	teroid therapy): or
x) pre term infants, born before 28 weeks g		g., acco cocc	10.0.u 1o. up///, 0.
xi) with cardiac disease, with cyanosis or fai	lure; or		
xii) with diabetes; or			
xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	functional asplenia		
xiv) who are pre or poor opionocionity, or with	Turiotional aspionia.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each			
23 pneumococcal serotype)	0.00	1 🗸	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]			
Up to three doses for patients meeting either of the follow			
 For partially vaccinated or previously unvaccinated i For revaccination following immunosuppression. 	ndividuals; or		
Note: Please refer to the Immunisation Handbook for app	propriate schedule for ca	tch-up programn	100
Inj 80D antigen units in 0.5 ml syringe			POL
ROTAVIRUS ORAL VACCINE - [Xpharm]		-	
Maximum of two doses for patients meeting the following:			
 first dose to be administered in infants aged under 1 no vaccination being administered to children aged 2 			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

		Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
VARICELLA V Either:	ACCINE [CHICKENPOX VACCINE] - [Xpharm]				
1) Maxi	imum of one dose for primary vaccination for eithe	er:			
,	Any infant born on or after 1 April 2016; or For previously unvaccinated children turning 11 varicella infection (chickenpox), or	years old on or after 1	July 201	7, who h	nave not previously had a
	imum of two doses for any of the following:				
a)	Any of the following for non-immune patients:				
	ii) with chronic liver disease who may in future iii) with deteriorating renal function before tran iii) prior to solid organ transplant; or	splantation; or	nsplanta	ition; or	
	iv) prior to any elective immunosuppression*,				
L١	v) for post exposure prophylaxis who are imm			_:::	-liet
,	For patients at least 2 years after bone marrow to For patients at least 6 months after completion of				·
	For HIV positive non immune to varicella with mi				
	For patients with inborn errors of metabolism at r varicella, or				
,	For household contacts of paediatric patients wh immune compromise where the household conta For household contacts of adult patients who have	ct has no clinical histo	ry of var	ricella, or	
9/	immunocompromised, or undergoing a procedur has no clinical history of varicella.				
28 days	suppression due to steroid or other immunosuppre	.,	for a tre	eatment p	period of greater than
Inj 1350 P	FU prefilled syringe	0.00	1	✓ \	/arivax
			10	✓ \	<u>/arivax</u>
	OSTER VIRUS (OKA STRAIN) LIVE ATTENUAT r patients meeting either of the following criteria:	ED VACCINE [SHING	LES VA	CCINE]	– [Xpharm]
1) One	dose for all people aged 65 years; or				
	dose for all people aged between 66 and 80 year	s inclusive from 1 Apri	l 2018 a	nd 31 De	ecember 2021.
Ini 19.400	PFU prefilled syringe plus vial	0.00	1	√ 2	ostavax .
,,	- F		10		ostavax
Diagnosti	c Agents				
	PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU po	er 0.1 ml, 1 ml vial	0.00	1	✓]	<u>ubersol</u>

- Symbols -	Afluria Quad Junior	Amoxicillin	
UK Synacthen84	(2021 Formulation) 266	Amoxicillin with clavulanic acid	9
3TC109	AFT-Pyrazinamide104	Amphotericin B	36
- A -	Agents Affecting the	Amsacrine	.150
A-Scabies72	Renin-Angiotensin System 51	AmsaLyo	
Abacavir sulphate109	Agents for Parkinsonism and Related	Amsidine	
Abacavir sulphate with	Disorders 122	Amzoate	3
lamivudine109	Agents Used in the Treatment of	Anaesthetics	
Abiraterone acetate167	Poisonings237	Anafranil	.12
Acarbose11	Agrylin153	Anagrelide hydrochloride	
Accarb11	Agrylin S29 153	Analgesics	
Accuretic52	Albendazole93	Anastrozole	
Accuretic 1052	Albey223	Anatrole	
Accuretic 2052	Albustix81	Andriol Testocaps	
Acetazolamide	Aldurazyme33	Androderm	
Acetec	Alecensa160	ANI	
Acetic acid with 1, 2- propanediol	Alectinib160	Anoro Ellipta	
diacetate and	Alendronate sodium115	Antabuse	
benzethonium232	Alendronate sodium with	Antacids and Antiflatulents	
Acetic acid with hydroxyquinoline and	colecalciferol	Anthelmintics	
ricinoleic acid79	Alfacalcidol	Antiacne Preparations	
Acetylcysteine237	Alfamino Junior258	Antiallergy Preparations	
Aci-Jel79	Alginic acid6	Antianaemics	
Aciclovir	Alglucosidase alfa31	Antiandrogen Oral	7
Infection105	Alkeran149	Contraceptives	70
Sensory232	Alkeran S29	Antiarrhythmics	
Acidex6	Allersoothe224	Antibacterials	
Acipimox 59	Allmercap	Antibacterials Topical	
Acitretin	Allopurinol120	Anticholinergic Agents	
Aclasta 118	•	Anticholinesterases	
Aclin	Alpha-Adrenoceptor Blockers51 Alpha-Keri Lotion70	Antidepressants	
Actemra 212	Alphamox96	Antidepressants	
Actinomycin D	Alphamox 12596	Antiepilepsy Drugs	
Actrapid	Alphamox 25096	Antifibrinolytics, Haemostatics and	. 10
Actrapid Penfill	Alprolix	Local Sclerosants	1
Acular234	Alu-Tab6	Antifibrotics	
Acupan	Aluminium hydroxide	Antifungals	
Adalat 10	Alvogen55	Antifungals Topical	. 10
Adalat Oros	Amantadine hydrochloride122	Antihistamines	22
Adalimumab	Ambrisentan 62	Antihypotensives	
Adapalene	Ambrisentan Mylan	Antimalarials	
Adcortyl84	Amiloride hydrochloride	Antimigraine Preparations	
Adefin57	Amiloride hydrochloride with	Antinausea and Vertigo Agents	
Adefin XL	furosemide58	Antiparasitics	
Adenuric	Amiloride hydrochloride with	Antipruritic Preparations	
ADR Cartridge 1.8	hydrochlorothiazide 58	Antipsychotics	
Adrenaline61	Aminophylline229	Antiretrovirals	
Advantan	Amiodarone hydrochloride53	Antirheumatoid Agents	
Advate	-	Antispasmodics and Other Agents	. 1 13
Advate 45 Advnovate 45	Amisulpride	Antispasmodics and Other Agents Altering Gut Motility	
Adyriovate	Amitriptyline	Antithrombotic Agents	
Allimor	Amlodipine56	Antithymocyte globulin	4
Afluria Quad	Amneal	(equine)	17
(2021 Formulation)	Amorolfine	Antitrichomonal Agents	
(2021 1 01111ulation)207	AITIOTOIIIITE	Anuthonomonal Agents	. 10

Antituberculotics and		Arrow-Amitriptyline	128	Azopt235
Antileprotics	103	Arrow-Bendrofluazide		AZT109–110
Antiulcerants		Arrow-Brimonidine	235	-B-
Antivirals	104	Arrow-Calcium	38	B-D Micro-Fine16
Anxiolytics		Arrow-Diazepam		B-D Ultra Fine16
Anzatax		Arrow-Doxorubicin		B-D Ultra Fine II16
Apidra		Arrow-Losartan &		Bacillus Calmette-Guerin (BCG)
Apidra SoloStar		Hydrochlorothiazide	52	vaccine
Apo-Amlodipine		Arrow-Morphine LA		Bacillus Calmette-Guerin
Apo-Azithromycin		Arrow-Norfloxacin		vaccine
Apo-Bromocriptine		Arrow-Ornidazole		Baclofen
Apo-Ciclopirox		Arrow-Quinapril 10		Bactroban
Apo-Cilazapril/	00	Arrow-Quinapril 20		Barrier Creams and Emollients69
Hydrochlorothiazide	52	Arrow-Quinapril 5		Basic AquaCream69
Apo-Clarithromycin		Arrow-Roxithromycin		BCG Vaccine
•	0			
Alimentary		Arrow-Sertraline		Beclazone 100
Infection		Arrow-Timolol		Beclazone 250
Apo-Clomipramine		Arrow-Topiramate		Beclazone 50
Apo-Diclo SR		Arrow-Tramadol		Beclomethasone dipropionate224
Apo-Diltiazem CD		Arsenic trioxide		Bee venom allergy treatment223
Apo-Doxazosin		Asacol		Bendamustine hydrochloride
Apo-Folic Acid		Asamax		Bendrofluazide59
Apo-Furosemide		Ascorbic acid		Bendroflumethiazide
Apo-Gabapentin		Aspen Adrenaline	61	[Bendrofluazide]59
Apo-Megestrol		Aspirin		Benzathine benzylpenicillin96
Apo-Metoprolol		Blood		Benzatropine mesylate122
Apo-Mirtazapine		Nervous		Benzbromaron AL 100120
Apo-Nadolol		Asthalin	<mark>226</mark>	Benzbromarone120
Apo-Nicotinic Acid		Atazanavir sulphate	110	Benztrop122
Apo-Oxybutynin		Atenolol	54	Benzydamine hydrochloride36
Apo-Perindopril		Atenolol AFT	54	Benzylpenicillin sodium [Penicillin
Apo-Pindolol	<u>55</u>	Atenolol AFT S29	54	G]96
Apo-Pravastatin	<mark>60</mark>	ATGAM	177	Beta Cream67
Apo-Prazosin		Ativan	138	Beta Ointment67
Apo-Prednisone	84	Atomoxetine	141	Beta Scalp73
Apo-Primidone		Atorvastatin	59	Beta-Adrenoceptor Agonists226
Apo-Propranolol		Atropine sulphate		Beta-Adrenoceptor Blockers54
Apo-Pyridoxine		Cardiovascular	53	Betadine70
Apo-Selegiline S29		Sensory	235	Betadine Skin Prep70
Apo-Sumatriptan		Atropt		Betaferon140
Apo-Temozolomide		Atrovent		Betahistine dihydrochloride134
Apo-Terazosin		AU Synacthen		Betaine32
Apo-Timol		Aubagio		Betaloc CR55
Apomorphine hydrochloride		Augmentin		Betamethasone dipropionate67
Aprepitant		Aurorix		Betamethasone dipropionate with
Apresoline		AutoSoft 30		calcipotriol
Aptamil AllerPro SYNEO 1		AutoSoft 90		Betamethasone sodium phosphate
Aptamil AllerPro SYNEO 2		Avelox		with betamethasone acetate
Aptamil Gold+ Pepti Junior		Avonex		Betamethasone valerate
Aqueous cream		Avonex Pen		Betamethasone valerate with
Aratac		Azacitidine		clioquinol68
Arava		Azacitidine Dr Reddy's		Betamethasone valerate with sodium
Aripiprazole 1 A Pharma	105	Azathioprino		fusidate [fusidic acid]
Aripiprazole 1A Pharma		Azathioprine		Betavolol
Aripiprazole Sandoz		Azithromycin		Betnovate
Aristocort	სბ	Azol	92	Betnovate-C68

Betoptic	234	Respiratory	224 230	CareSens N POP	1/
Betoptic S		Budesonide with eformoterol		CareSens N Premier	
Bezafibrate		Bumetanide		CareSens PRO	
Bezalip		Buprenorphine Naloxone BNM		Carmellose sodium with gelatin and	
Bezalip Retard		Buprenorphine with naloxone.		pectinpectin	
Bicalutamide		Bupropion hydrochloride		Carmustine	
Bicillin LA		Burinex		Carvedilol	
BiCNU		Burinex S29		Carvedilol Sandoz	
Bicnu Heritage		Buscopan		Catapres	
Bile and Liver Therapy		Buspirone hydrochloride		CeeNU	
Biltricide		Busulfan	149	Cefaclor monohydrate	
Bimatoprost		- C -		Cefalexin	9:
Bimatoprost Multichem		Cabergoline	91	Cefalexin Sandoz	9:
Binarex		Cacit	38	Cefazolin	
Binocrit		Caffeine citrate		Ceftriaxone	
Biodone		Calamine		Ceftriaxone-AFT	
Biodone Extra Forte		Calci-Tab 500		Cefuroxime axetil	9:
Biodone Forte		Calcipotriol		Celebrex	
Bisacodyl		Calcitonin		Celecoxib	
Bisoprolol fumarate		Calcitriol		Celecoxib Pfizer	11/
Bisoprolol Mylan		Calcitriol-AFT		Celestone Chronodose	 R:
BK Lotion		Calcium carbonate		Celiprolol	
Bleomycin sulphate		Calcium Channel Blockers		Cellcept	
Blood Colony-stimulating		Calcium Disodium Versenate.		Celol	
Factors	48	Calcium folinate		Centrally-Acting Agents	
Blood glucose diagnostic test		Calcium Folinate Ebewe		Cephalexin ABM	o
meter	15	Calcium Folinate Sandoz		Cetirizine hydrochloride	
Blood glucose diagnostic test		Calcium gluconate		Cetomacrogol	
strip	15	Calcium Homeostasis		Cetomacrogol with glycerol	
Blood glucose test strips (visua		Calcium polystyrene sulphona		Cetuximab	
impaired)	•	Calcium Resonium		Charcoal	
Blood Ketone Diagnostic Test		Calcium Sandoz		Chemotherapeutic Agents	
Strip	1/	Calogen		Chickenpox vaccine	27
Bonjela		Calutide-50		Chlorafast	
Boostrix		Candesartan cilexetil		Chlorambucil	
Bortezomib		Candestar		Chloramphenicol	
Bortezomib Dr-Reddy's		Canesten		Chlorothiazide	
Bosentan		Capecitabine		Chlorpheniramine maleate	
Bosentan Dr Reddy's		Capercit		Chlorpromazine hydrochloride	
Bosvate		Capoten		Chlortalidone [Chlorthalidone]	
Bplex	37	Capsaicin		Chlorthalidone	
Breo Ellipta		Musculoskeletal	115	Chlorvescent	
Brevinor 1/28		Nervous		Choice Load 375	
Bricanyl Turbuhaler		Captopril		Choice TT380 Short	
Brilinta		Captopril-Mylan		Choice TT380 Standard	
Brimonidine tartrate		Carafate		Choline salicylate with cetalkonium	/
Brimonidine tartrate with timolo		Carbaccord		chloride	31
maleate		Carbamazepine		Ciclopirox olamine	
Brinzolamide		Carbimazole		Ciclosporin	201
Bristol		Carbomer		Cilazapril	
Brolene		Carboplatin		Cilazapril with	5
Bromocriptine mesylate		Carboplatin Ebewe		hydrochlorothiazide	E
BSF Ambrisentan Mylan		Carbosorb-X		Cilicaine	O
Buccastem		Cardinol LA		Cilicaine VK	შ
Budesonide	104	CareSens Dual		Cinacalcet	
Alimentary	e	CareSens N			
Allineillary	<mark>0</mark>	Caredens in	10	Cipflox	y

Ciprofloxacin	Colistin sulphomethate98	Danazol9
Infection97	Colistin-Link98	Dantrium12
Sensory232	Collodion flexible240	Dantrium S2912
Ciprofloxacin Teva232	Colloidal bismuth subcitrate10	Dantrolene 12
Circadin140	Colofac8	Daonil1
Cisplatin149	Coloxyl30	Dapa-Tabs5
Cisplatin Ebewe149	Combigan235	Dapsone 103
Citalopram hydrobromide129	Compound electrolytes50	Daraprim99
Cladribine151	Compound electrolytes with glucose	Darunavir11
Clarithromycin	[Dextrose]50	Darunavir Mylan11
Alimentary9	Compound hydroxybenzoate240	Dasatinib16
Infection94	Concerta143	Daunorubicin15
Clexane47	Condoms76	David One Step Cassette Pregnancy
Clexane Forte47	Condyline74	Test 80
Climara85	Contraceptives - Hormonal77	DBL Acetylcysteine23
Climara85	Contraceptives - Non-hormonal75	DBL Adrenaline6
Clindamycin98	Copaxone139	DBL Aminophylline22
Clinicians Renal Vit37	Corticosteroids and Related Agents	DBL Bleomycin Sulfate15
Clobazam130	for Systemic Use83	DBL Carboplatin14
Clobetasol propionate67, 73	Corticosteroids Topical67	DBL Cisplatin14
Clobetasone butyrate68	Cosentyx211	DBL Dacarbazine15
Clofazimine103	Cosmegen 154	DBL Desferrioxamine Mesylate for Inj
Clomazol	Coumadin48	BP23
Dermatological66	Coversyl51	DBL Docetaxel15
Genito-Urinary79	Creon 1000029	DBL Ergometrine79
Clomifene citrate92	Creon 2500029	DBL Gemcitabine15
Clomipramine hydrochloride128	Creon Micro29	DBL Gentamicin9
Clonazepam 130, 138	Crotamiton67	DBL Heparin Sodium4
Clonidine57	Crystaderm65	DBL Leucovorin Calcium 15
Clonidine BNM57	Curam96	DBL Methotrexate Onco-Vial15
Clonidine hydrochloride57	Curam Duo 500/12596	DBL Morphine Sulphate12
Clopidogrel45	Cvite37	DBL Naloxone Hydrochloride 23
Clopidogrel Multichem45	Cyclizine hydrochloride134	DBL Octreotide16
Clopine	Cyclizine lactate134	DBL Pethidine Hydrochloride 12
Clopixol136, 138	Cyclogyl236	DBL Vinblastine16
Clotrimazole	Cyclopentolate hydrochloride236	DBL Vincristine Sulfate16
Dermatological66	Cyclophosphamide149	Decozol3
Genito-Urinary79	Cyclorin103	Deferasirox23
Clozapine135	Cycloserine103	Deferiprone23
Clozaril135	Cyproterone acetate84	Denosumab11
Co-trimoxazole100	Cyproterone acetate with	Deolate10
Coal tar72	ethinyloestradiol79	Deoxycoformycin15
Coal tar with allantoin, menthol,	Cystadane32	Depo-Medrol8
phenol and sulphur73	Cytarabine151	Depo-Provera79
Coal tar with salicylic acid and	Cytotec8	Depo-Testosterone8
sulphur73	Cytoxan149	Deprim100
Coco-Scalp	- D -	Dermol67, 75
Codeine phosphate	D-Penamine115	Desferrioxamine mesilate23
Extemporaneous240	Dabigatran48	Desmopressin9
Nervous125	Dacarbazine154	Desmopressin acetate9
Colchicine	Dacarbazine APP154	Desmopressin-PH&T9
Colecalciferol37	Dactinomycin [Actinomycin D]154	Desuric12
Colestid59	Daivobet72	Detection of Substances in
Colestipol hydrochloride59	Daivonex72	Urine8
Colgout120	Daktarin67	Dexamethasone
Colifoam7	Dalacin C98	Hormone83

Sensory233	Diuretics	58	Elaprase	3,
Dexamethasone phosphate83	Docetaxel		Elecare	
Dexamethasone Phosphate	Docetaxel Accord		Elecare LCP	
Panpharma83	Docetaxel Sandoz		Electral	
Dexamethasone with framycetin and	Docusate sodium		Elelyso	
gramicidin232	Docusate sodium with		Elemental 028 Extra	2/1
Dexamethasone with neomycin	sennosides	30	Elidel	
sulphate and polymyxin B	Dolutegravir		Elocon	
sulphate233	Domperidone		Elocon Alcohol Free	
Dexamfetamine sulfate141	Donepezil hydrochloride		Eltrombopag	
Dexmethsone	Donepezil-Rex		Eltroxin	
Dextrochlorpheniramine	Dornase alfa		EMB Fatol	
maleate224	Dortimopt		Emend Tri-Pack	
Dextrose	Dorzolamide hydrochloride		Emicizumab	
DHC Continus	Dorzolamide with timolol		EMLA	
Diabetes	Dostinex		Empagliflozin	
Diabetes Management	Dosulepin [Dothiepin]	91	, 0	
		100	Empagliflozin with metformin	47
Diacomit	hydrochloride		hydrochloride	
Diagnostic Agents272	Dosulepin Mylan		Emtricitabine	10
Diamide Relief	Dothiepin		Emtricitabine with tenofovir	10
Diamox	Doxazosin		disoproxil	
Diasip245	Doxine		Emtriva	
Diason RTH	Doxorubicin Ebewe		Emulsifying ointment	
Diazepam	Doxorubicin hydrochloride		Emulsifying Ointment ADE	
Diazoxide	Doxycycline		Enalapril maleate	
Dibenzyline	DP Lotion		Enbrel	
Diclofenac Sandoz114	DP Lotn HC		Endocrine Therapy	
Diclofenac sodium	DP-Allopurinol		Endoxan	143
Musculoskeletal114	Dr Reddy's Omeprazole	9	Engerix-B	
Sensory233	Drugs Affecting Bone	445	Enlafax XR	
Differin	Metabolism		Enoxaparin sodium	
Difflam	Dual blood glucose and blood keto		Enstilar	
Diflucan	diagnostic test meter		Ensure	
Diflucortolone valerate	Duocal Super Soluble Powder		Ensure Plus	
Digestives Including Enzymes29	Duolin		Ensure Plus HN	
Digoxin	Duolin HFA		Ensure Plus RTH	
Dihydrocodeine tartrate	DuoResp Spiromax		Entacapone	
Dilantin	Duride	61	Entapone	
Dilantin Infatab131	- E -	004	Entecavir	
Diltiazem hydrochloride57	e-chamber La Grande		Entecavir Sandoz	
Dilzem57	e-chamber Mask		Entocort CIR	
Dimethicone 69–70	e-chamber Turbo		Entresto 24/26	5
Dimethyl fumarate	E-Mycin		Entresto 49/51	
Dipentum7	Ear Preparations		Entresto 97/103	
Diphtheria, tetanus and pertussis	Ear/Eye Preparations		Epilim	
vaccine	Easiphen Liquid	257	Epilim Crushable	13
Diphtheria, tetanus, pertussis and	Econazole nitrate		Epilim IV	
polio vaccine	Efavirenz	109	Epilim S/F Liquid	
Diphtheria, tetanus, pertussis, polio,	Efavirenz with emtricitabine and	400	Epilim Syrup	13
hepatitis B and haemophilus	tenofovir disoproxil		Epirubicin Ebewe	
influenzae type B vaccine 264	Eformoterol fumarate		Epirubicin hydrochloride	15
Diprosone67	Eformoterol fumarate dihydrate	225	Eplerenone	5
Diprosone OV67	Eftrenonacog alfa [Recombinant		Epoetin alfa	4
Dipyridamole45	factor IX]		Epoprostenol	6
Disopyramide phosphate53	Efudix		Eptacog alfa [Recombinant factor	
Disulfiram146	Egopsoryl TA	73	VIIa]	44

ERA	95	Feed Thickener Karicare		Fluorometholone	23
Erbitux		Aptamil	255	Fluorouracil	
Ergometrine maleate		FEIBA NF		Fluorouracil Ebewe	
Erlotinib		Felo 10 ER	56	Fluorouracil sodium	
Erythrocin IV		Felo 5 ER		Fluox	
Erythromycin (as lactobionate)		Felodipine		Fluoxetine hydrochloride	
Erythromycin ethyl succinate		Femme-Tab ED		Flupenthixol decanoate	
Erythromycin stearate		Fentanyl		Flutamide	
Esbriet		Fentanyl Sandoz		Flutamin	
		Ferinject		Fluticasone	
Escitalopram Apotox		-			22
Escitalopram-Apotex		Ferodan		Fluticasone furoate with	00
Eskazole		Ferric carboxymaltose		vilanterol	
Essential Prednisolone		Ferriprox		Fluticasone propionate	
Estradiol TDP Mylan		Ferro-F-Tabs		Fluticasone with salmeterol	
Estradot		Ferro-tab		FML	
Estradot 50 mcg		Ferrograd		Foban	
Estrofem	85	Ferrosig	39	Folic acid	
Etanercept	170	Ferrous fumarate	39	Food Thickeners	25
Ethambutol hydrochloride	103	Ferrous fumarate with folic acid.	39	Foods And Supplements For I	nborn
Ethics Aspirin	124	Ferrous sulfate	39	Errors Of Metabolism	25
Ethics Aspirin EC	45	Ferrous sulphate	39	Foradil	22
Ethics Lisinopril		Fexofenadine hydrochloride		Forteo	11
Ethics Paracetamol Classic		Fibro-vein		Fortini	24
Ethinyloestradiol		Filgrastim		Fortini Multi Fibre	
Ethinyloestradiol with		Finasteride		Fortisip	
desogestrel	77	Fingolimod		Fortisip Multi Fibre	
Ethinyloestradiol with		Firazyr		Fosamax	
levonorgestrel	78	Flagyl	103	Fosamax Plus	
Ethinyloestradiol with		Flagyl-S		Framycetin sulphate	
norethisterone	78	Flamazine		Frisium	
Ethosuximide		Flecainide acetate		Frumil	
				Frusemide	
Etopophos		Flecainide BNM	33		
Etoposide		Flecainide Controlled Release		Fucicort	
Etoposide phosphate		Teva		Fucidin	
Etravirine		Fleet Phosphate Enema		Fucithalmic	
Eumovate		Flixonase Hayfever & Allergy		Fulvestrant	
Everet		Flixotide		Fungilin	
Everolimus		Flixotide Accuhaler		Furosemide [Frusemide]	
Evista		Florinef	83	Furosemide-Baxter	5
Exemestane		Fluad Quad		fusidic acid	
Exjade		(2021 Formulation)	267	Dermatological	66, 6
Extemporaneously Compounded	t	Fluanxol		Infection	
Preparations and		Flucil	96	Sensory	23
Galenicals	240	Flucloxacillin	96	- G -	
Eye Preparations		Flucloxin	96	Gabapentin	13
Eylea	188	Flucon	234	Gacet	12
Ezetimibe		Fluconazole	100	Galsulfase	
Ezetimibe Sandoz	60	Fludara Oral	151	Galvumet	
Ezetimibe with simvastatin		Fludarabine Ebewe		Galvus	
- F -		Fludarabine phosphate		Gardasil 9	
Factor eight inhibitor bypassing		Fludrocortisone acetate	83	Gastrodenol	
fraction	44	Fluids and Electrolytes		Gaviscon Double Strength	
Famotidine		Flumetasone pivalate		Gaviscon Infant	
Famotidine Hovid		Fluocortolone caproate with	202	Gazyva	
Faslodex		fluocortolone pivalate and		Gefitinib	
		oinchocoine	0	Gemcitabine Ebewe	
Febuxostat	120	cinchocaine	Ö	Genicitabilie Ebewe	15

Gemcitabine hydrochloride	151	healthE Urea Cream	69	Hydroxyurea	
Genoptic	232	Healtheries Simple Baking Mix	255	[hydroxycarbamide]	155
Gentamicin sulphate		Hemastix		Hygroton	
Infection	98	Hemlibra	43	Hylo-Fresh	
Sensory	232	Heparin sodium	47	Hymenoptera	
Gilenya		Heparinised saline		Hyoscine butylbromide	
Ginet		Heparon Junior		Hyoscine hydrobromide	
Glatiramer acetate		Hepatitis A vaccine		Hypam	
Glecaprevir with pibrentasvir		Hepatitis B recombinant		Hyperuricaemia and Antigout	
Glibenclamide		vaccine	265	Hypromellose	
Gliclazide		Herceptin		Hypromellose with dextran	
Glipizide	13	Hiberix		-1-	
Glivec		Hiprex		Ibiamox	96
Glizide	13	Histaclear		Ibrance	164
Glucagen Hypokit		Histafen		Ibuprofen	
Glucagon hydrochloride		Holoxan		Ibuprofen SR BNM	
Glucerna Select		Horleys Bread Mix		Icatibant	
Glucerna Select RTH		Horleys Flour		Idarubicin hydrochloride	
Glucobay		Hormone Replacement Therapy -	200	Idursulfase	
Glucose [Dextrose]		Systemic	84	Ifosfamide	
Gluten Free Foods		HPV		Igroton	
Glycerin with sodium saccharin		Humalog		lkorel	
Glycerin with sucrose		Humalog Mix 25		llevro	
Glycerol		Humalog Mix 50		lloprost	
Alimentary	31	Human papillomavirus (6, 11, 16, 1		Imatinib mesilate	
Extemporaneous		31, 33, 45, 52 and 58) vaccine	Ο,	Imatinib-AFT	
Glyceryl trinitrate	240	[HPV]	265	Imatinib-Rex	
Alimentary	8	Humatin		Imigran	
Cardiovascular		Humira		Imipramine hydrochloride	
Glycopyrronium		HumiraPen		Imiquimod	
Glycopyrronium bromide		Humulin 30/70		Immune Modulators	
Glycopyrronium with		Humulin NPH		Immunosuppressants	
indacaterol	227	Humulin R		Imuran	
Gold Knight		Hyaluronic acid		Incruse Ellipta	
Gold Knight XL		Hydralazine		Indacaterol	
Goserelin		Hydralazine hydrochloride		Indapamide	
Gutron		Hydrocortisone		Infanrix IPV	
Gynaecological Anti-infectives		Dermatological	68	Infanrix-hexa	
- H -		Hormone		Infant Formulae	
Habitrol	146	Hydrocortisone (PSM)		Infatrini	
Haemophilus influenzae type B		Hydrocortisone acetate		Infliximab	
vaccine	264	Hydrocortisone acetate with		Influenza vaccine	
Haldol		pramoxine hydrochloride	7	Inhaled Corticosteroids	
Haldol Concentrate		Hydrocortisone and paraffin liquid		Inhaled Long-acting	224
Haldol Decanoas		and lanolin	68	Beta-adrenoceptor Agonists	225
Haloperidol		Hydrocortisone butyrate		Inspra	
Haloperidol decanoate		Hydrocortisone with cinchocaine		Instillagel Lido	122
Harvoni		Hydrocortisone with miconazole		Insulin aspart	
Havrix Havrix Junior		Hydrocortisone with natamycin and		Insulin aspart with insulin aspart	10
		neomycin		protamine	
healthE Calamine Aqueous Crea		Hydrogen peroxide		Insulin glargine	
healthE Dimethicone 10%		Hydroxocobalamin Hydroxocobalamin Mercury	30	Insulin glulisine	
			26	Insulin isophane	11
healthE Dimethicone 4% Lotion .		Pharma		Insulin isophane with insulin	44
healthE Dimethicone 5%healthE Glycerol BP		hydroxycarbamide Hydroxychloroquine		neutral	
HEARINE CHYCERUL DY	∠4∪	r ryuruxycriioroquirie	1 10	Insulin lispro	11

Insulin lispro with insulin lispro	Ivacaftor	230	Lantus SoloStar11
protamine11	Ivermectin	70	Lanvis 153
Insulin neutral10	- J -		Lanzol Relief9
Insulin pen needles16	Jadelle	78	Lapatinib ditosylate163
Insulin pump16	Jakavi	165	Largactil135
Insulin pump cartridge21	Jardiamet	12	Laronidase33
Insulin pump infusion set (steel	Jardiance	11	Lasix58
cannula)22	Jaydess		Latanoprost235
Insulin pump infusion set (steel	Jevity HiCal RTH		Lax-Suppositories31
cannula, straight insertion) 23	Jevity RTH		Lax-Tab31
Insulin pump infusion set (teflon	Juno Pemetrexed		Laxatives30
cannula)24	- K -		Laxsol
Insulin pump infusion set (teflon	Kadcyla	217	Ledipasvir with sofosbuvir106
cannula, angle insertion with	Kaletra		Leflunomide115
insertion device)25	Kalydeco		Lenalidomide
Insulin pump infusion set (teflon	Kemadrin		Letrole
cannula, angle insertion)	Kenacomb		Letrozole170
Insulin pump infusion set (teflon	Kenacort-A 10		Leukeran FC149
cannula, straight insertion with	Kenacort-A 40		Leukotriene Receptor
insertion device)26	Kenalog		Antagonists229
Insulin pump infusion set (teflon	Kenalog in Orabase		Leuprorelin91
cannula, straight insertion) 28	Kenkay Sorbolene	69	Leustatin151
Insulin pump reservoir28	Ketocal 3:1	262	Levetiracetam131
Insulin syringes, disposable with	KetoCal 4:1	262	Levetiracetam-AFT131
attached needle16	Ketoconazole		Levlen ED78
Intal Forte CFC Free229	Dermatological	<mark>73</mark>	Levocabastine234
Intelence109	Infection	101	Levodopa with benserazide122
Interferon beta-1-alpha139	Ketogenic Diet	262	Levodopa with carbidopa122
Interferon beta-1-beta140	Ketoprofen		Levomepromazine136
Intra-uterine device77	Ketorolac trometamol	234	Levomepromazine
Invega Sustenna137	KetoSens	14	hydrochloride 136
IPOL271	Keytruda		Levonorgestrel
Ipratropium bromide226, 230	Kindergen		Genito-Urinary78-79
Iressa	Kivexa		Hormone86
Irinotecan Accord151	Klacid		Levothyroxine87
Irinotecan Actavis 100151	Kliogest		Lidocaine [Lignocaine]123–124
Irinotecan hydrochloride	Kliovance		Lidocaine [Lignocaine]
Irinotecan-Rex	Kogenate FS		hydrochloride124
Iron polymaltose39	Konakion MM		Lidocaine [Lignocaine] with
lsentress 110	Konsyl-D		chlorhexidine 124
Isentress HD	Kuvan		
Ismo 20	- L -		Lidocaine [Lignocaine] with
	_		prilocaine
Ismo 40 Retard 61	Labetalol		Lidocaine-Claris124
Isoniazid	Lacosamide		Lignocaine
Isoniazid with rifampicin103	Lactulose		Lioresal Intrathecal
Isoptin57	Laevolac		Lipid-Modifying Agents59
Isoptin Retard57	Lamictal		Liquigen244
Isoptin SR57	Lamivudine	,	Lisinopril51
Isopto Carpine235	Lamivudine Alphapharm		Lithium carbonate136
Isosorbide mononitrate61	Lamotrigine		Livostin234
Isosource Standard252	Lamprene		LMX4124
Isotretinoin65	Lanoxin	53	Locacorten-Viaform ED's232
Ispaghula (psyllium) husk30	Lanoxin PG	53	Local preparations for Anal and
Itch-Soothe67	Lanoxin S29	53	Rectal Disorders8
Itraconazole100	Lansoprazole	9	Locasol258
Itrazole100	Lantus		Locoid68, 73

Locoid Crelo	68	MCT oil (Nutricia)244	Methylprednisolone aceponate	6
Locoid Lipocream	68	Measles, mumps and rubella	Methylprednisolone acetate	8
Locorten-Vioform	232	vaccine 268	Methylxanthines	22
Lodoxamide	234	Mebendazole93	Metoclopramide Actavis 10	
Logem	131	Mebeverine hydrochloride8	Metoclopramide hydrochloride	
Lomide		Medco125	Metolazone	
Lomustine		Medrol83	Metopirone	
Loniten	62	Medroxyprogesterone acetate	Metoprolol IV Mylan	
Loperamide hydrochloride	6	Genito-Urinary79	Metoprolol succinate	
Lopinavir with ritonavir		Hormone85–86	Metoprolol tartrate	
Loprofin		Mefenamic acid114	Metrogyl	
Loprofin Mix		Megestrol acetate168	Metronidazole	10
Lorafix		Melatonin140	Metyrapone	
Loratadine		Melphalan 149	Mexiletine hydrochloride	
Lorazepam		Menactra269	Mexiletine Hydrochloride USP	
Lorfast		Meningococcal (groups A, C, Y and	Miacalcic	
Lorstat		W-135) conjugate vaccine269	Micolette	
Losartan Actavis		Meningococcal C conjugate	Miconazole	
		vaccine269	Miconazole nitrate	
Losartan potassium	52			e.
Losartan potassium with	50	Menthol	Dermatological	
hydrochlorothiazide		Mepolizumab	Genito-Urinary	
Lovir		Mercaptopurine	Micreme	
Loxamine		Mercilon 2877	Micreme H	
Lucrin Depot 1-month		Mesalazine7	Microgynon 20 ED	
Lucrin Depot 3-month		Mesna	Microgynon 30	
Ludiomil		Mestinon114	Microgynon 50 ED	
Lyderm		Metabolic Disorder Agents31	Microlut	
Lynparza		Metformin hydrochloride13	Midazolam	
Lyrica	131	Methadone hydrochloride	Midazolam-Baxter	
- M -	407	Extemporaneous240	Midodrine	
m-Eslon		Nervous126	Mifegyne	
Mabthera		Methatabs126	Mifepristone	
Macrobid	113	Methenamine (hexamine)	Minerals	
Macrogol 3350 with potassium		hippurate113	Mini-Wright AFS Low Range	
chloride, sodium bicarbonate		Methopt236	Mini-Wright Standard	
sodium chloride	31	Methotrexate152	Minidiab	13
Macrogol 400 and propylene		Methotrexate DBL Onco-Vial152	MiniMed 1.8 Reservoir	
glycol		Methotrexate Ebewe152	MMT-326A	2
Madopar 125		Methotrexate Sandoz152	MiniMed 3.0 Reservoir	
Madopar 250	122	Methyl hydroxybenzoate240	MMT-332A	
Madopar 62.5		Methylcellulose240	MiniMed 640G	
Madopar HBS	122	Methylcellulose with glycerin and	MiniMed Mio MMT-921A	2
Madopar Rapid		, , , , , , , , , , , , , , , , , , , ,		2
Magnesium hydroxide		sodium saccharin240	MiniMed Mio MMT-923A	
Maanaaium aulahata	122	, , , , , , , , , , , , , , , , , , , ,	MiniMed Mio MMT-923A MiniMed Mio MMT-925A	2
Magnesium sulphate	122 39	sodium saccharin240	MiniMed Mio MMT-923A MiniMed Mio MMT-925A MiniMed Mio MMT-941A	2
Mantoux	122 39 40	sodium saccharin	MiniMed Mio MMT-923A MiniMed Mio MMT-925A	2
·	122 39 40 272	sodium saccharin	MiniMed Mio MMT-923A MiniMed Mio MMT-925A MiniMed Mio MMT-941A MiniMed Mio MMT-943A MiniMed Mio MMT-945A	24 24 24
Mantoux	122 39 40 272 128	sodium saccharin	MiniMed Mio MMT-923A MiniMed Mio MMT-925A MiniMed Mio MMT-941A MiniMed Mio MMT-943A	24 24 24
Mantoux Maprotiline hydrochloride	122 39 40 272 128	sodium saccharin	MiniMed Mio MMT-923A MiniMed Mio MMT-925A MiniMed Mio MMT-941A MiniMed Mio MMT-943A MiniMed Mio MMT-945A	24 24 24
Mantoux Maprotiline hydrochloride Marevan	122 39 40 272 128 48	sodium saccharin	MiniMed Mio MMT-923AMiniMed Mio MMT-925AMiniMed Mio MMT-941AMiniMed Mio MMT-943AMiniMed Mio MMT-945AMiniMed Mio MMT-965A	24 24 24 24
Mantoux	122 39 40 272 128 48 74 237	sodium saccharin	MiniMed Mio MMT-923AMiniMed Mio MMT-925AMiniMed Mio MMT-941AMiniMed Mio MMT-943AMiniMed Mio MMT-945AMiniMed Mio MMT-965AMiniMed Mio MMT-975A	24 24 24 24
Mantoux	122 39 40 272 128 48 74 237 77	sodium saccharin	MiniMed Mio MMT-923A	2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴
Mantoux	122 39 40 272 128 48 74 237 77	sodium saccharin	MiniMed Mio MMT-923A	242424242424
Mantoux	122 39 40 272 128 48 74 237 77 230 29	sodium saccharin	MiniMed Mio MMT-923A	24 24 24 24 24 24
Mantoux		sodium saccharin 240 Methylcellulose with glycerin and sucrose 240 Methyldopa 58 Methyldopa Mylan 58 Methyldopa Mylan S29 58 Methylnaltrexone bromide 30 Methylphenidate ER - Teva 142 Methylphenidate hydrochloride 142 Methylphenidate hydrochloride 143 Methylprednisolone 83	MiniMed Mio MMT-923A	2 ² 2 ³ 2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴ 2 ⁴
Mantoux		sodium saccharin	MiniMed Mio MMT-923A	24 24 24 24 24 24 24 24 24 24 24 24

MiniMed Silhouette MMT-377A.	24	Mupirocin	66	Nevirapine	10
MiniMed Silhouette MMT-378A.	24	Muscle Relaxants		Nevirapine Alphapharm	10
MiniMed Silhouette MMT-381A.	24	Mvite	38	Nicorandil	
MiniMed Silhouette MMT-382A.	24	Myambutol	103	Nicotine	
MiniMed Silhouette MMT-383A.		Mycobutin		Nicotinic acid	
MiniMed Silhouette MMT-384A.		MycoNail		Nifedipine	
MiniMed Sure-T MMT-864A		Mycophenolate mofetil		Nifuran	
MiniMed Sure-T MMT-866A		Mydriacyl		Nilotinib	
MiniMed Sure-T MMT-874A		Mylan Atenolol		Nilstat	
MiniMed Sure-T MMT-876A		Mylan Clomiphen		Alimentary	3
MiniMed Sure-T MMT-884A		Mylan Midazolam		Genito-Urinary	
MiniMed Sure-T MMT-886A		Myleran		Infection	10
Minims Cyclopentolate		Myometrial and Vaginal Hormo		Nintedanib	
Minims Pilocarpine		Preparations		Nipent	
Minims Prednisolone		Myozyme		Nitrates	
Minirin		Mysoline S29		Nitroderm TTS	
Minirin Melt		- N -		Nitrofurantoin	
Mino-tabs		Nadolol	55	Nitrolingual Pump Spray	
Minocycline hydrochloride		Naglazyme		Nivestim	4
Minomycin		Nalcrom		Nivolumab	
Minor Skin Infections		Naloxone hydrochloride		Nizoral	
Minoxidil		Naltraccord		Nodia	
Mirena		Naltrexone hydrochloride		Noflam 250	
Mirtazapine		Naphazoline hydrochloride		Noflam 500	
•					1 1
Misoprostol		Naphcon Forte	114	Non-Steroidal Anti-Inflammatory	44
Mitomycin C		Naprosyn SR 1000		Drugs Nonacog gamma, [Recombinant	11
Mitozantrone		Naprosyn SR 750			4
Mitozantrone Ebewe		Naproxen		Factor IX]	4
Mixtard 30		Narcaricin mite		Norethisterone	7
MMR II Moclobemide		Nasal Preparations	230	Genito-Urinary Hormone	
		Natalizumab			
Modafinil		Natulan		Norflex	
Modavigil		Nausafix		Norfloxacin	
Moduretic		Nausicalm		Noriday 28	
Molaxole		Navelbine		Norimin	
Moments		Necon		Normacol Plus	
Mometasone furoate		Nedocromil		Normison	
Monogen		Nefopam hydrochloride		Norpress	
Montelukast		Neisvac-C		Nortriptyline hydrochloride	
Montelukast Mylan		Neo-B12		Norvir	
Moroctocog alfa [Recombinant fa		Neo-Mercazole		NovaSource Renal	
VIII]		Neo-Mercazole S29		Novatretin	
Morphine hydrochloride		Neocate Gold		NovoMix 30 FlexPen	
Morphine sulphate		Neocate Junior Unflavoured		NovoRapid	
Motetis		Neocate Junior Vanilla		NovoRapid FlexPen	
Mouth and Throat		Neocate SYNEO		NovoRapid Penfill	
Movapo		Neoral		NovoSeven RT	
Moxifloxacin		Neostigmine metilsulfate		Noxafil	
MSUD Maxamum	256	Nepafenac	234	Nozinan	
Mucilaginous laxatives with		Nepro HP (strawberry)		Nozinan (Swiss)	
stimulants		Nepro HP (vanilla)		Nucala	
Mucolytics		Nepro HP RTH		Nuelin	
Mucosoothe		Nerisone		Nuelin-SR	
Multiple Sclerosis Treatments		Neulactil		Nutilis	
Multivitamin renal		Neulastim		Nutrient Modules	
Multivitamins	37	NeuroTabs	38	Nutrini Energy Multi Fibre	24

Nutrini Energy RTH247	Omnitrope87	Palbociclib	16
Nutrini Low Energy Multi Fibre249	Onbrez Breezhaler225	Paliperidone	13
Nutrini Peptisorb260	Oncaspar LYO157	Pamidronate disodium	
Nutrini Peptisorb Energy260	OncoTICE177		110
Nutrini RTH247	Ondansetron134	Panadol Mini Caps	12
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA134		
Fibre252	One-Alpha37		
Nutrison Concentrated254	Onrex134		
Nutrison Energy251	Opdivo217	Panzytrat	2
Nutrison Energy Multi Fibre252	Ora-Blend240		
Nutrison Multi Fibre252	Ora-Blend SF240	Para-amino salicylic acid	
Nutrison Standard RTH252	Ora-Plus240		
Nyefax Retard57	Ora-Sweet240	Paracare Double Strength	12
Nystatin	Ora-Sweet SF240		
Alimentary36	Orabase36		
Genito-Urinary79	Oral and Enteral Feeds245		. 12
Infection101	Oratane65		
NZB Low Gluten Bread Mix255	Ordine126		
-0-	Orgran256		
O/W Fatty Emulsion Cream69	Ornidazole103		
Obinutuzumab197	Orphenadrine citrate121		
Obstetric Preparations81	Ortho-tolidine81		
Ocicure9	Oruvail SR114		
Ocrelizumab140	Osmolite RTH252		
Ocrevus	Other Endocrine Agents91		
Octocog alfa [Recombinant factor	Other Oestrogen Preparations		
VIII] (Advate)45	Other Progestogen	Paradigm Quick-Set MMT-386	
Octocog alfa [Recombinant factor	Preparations86		
VIII] (Kogenate FS)45	Other Skin Preparations74		
Octreotide	Ovestin	Paradigm Quick-Set MMT-397	
Octreotide (Sun)	Genito-Urinary79		
Octreotide GH	Hormone86		
Octreotide LAR (somatostatin	Ox-Pam138	Paradigm Silhouette MMT-368	
analogue)169	Oxaliplatin150		
Octreotide MaxRx	Oxaliplatin Accord150		
Oestradiol	Oxaliplatin Actavis 100		
Oestradiol valerate85	Oxaliplatin Ebewe		
Oestradiol with norethisterone86	Oxazepam138		
Oestriol	Oxis Turbuhaler 225	•	
Genito-Urinary79	Oxpentifylline		
Hormone86	Oxybutynin80		
Oestrogens85	Oxycodone hydrochloride127		
Ofev227	Oxycodone Sandoz127	Paradigm Sure-T MMT-876	
Oil in water emulsion	Oxycodone Sandoz S29127		
Olanzapine	OxyNorm127		
Olaparib	Oxytocin		
Olbetam59	Oxytocin BNM		
Olbetam S29	Oxytocin with ergometrine	Paraldehyde	
Olopatadine	maleate80		
Olopatadine Teva236	Ozurdex233		
Olsalazine	- P -	Parnate	
Omalizumab	Pacifen121		
	Paclitaxel		
Omeprazole actavis 10	Paclitaxel Actavis		a
Omeprazole actavis 109 Omeprazole actavis 209	Paclitaxel Ebewe		
•	Paediatric Seravit		
Omeprazole actavis 409	ratulatile ottavil3/	гала!!!	13

Paxtine	129	Pindolol	55	Pregabalin Pfizer	13
Pazopanib	165	Pine tar with trolamine laurilsulfate		Pregnancy Tests - hCG Urine	
Peak flow meter		and fluorescein	73	Premarin	
Pedialyte - Bubblegum		Pinetarsol	73	Prevenar 13	27
Pediasure		Pioglitazone	13	Prezista	11
Pediasure RTH		Pirfenidone		Priadel	13
Pegaspargase		Pizotifen		Primacin	
Pegasys		PKU Anamix Infant	257	Primaquine	10
Pegfilgrastim		PKU Anamix Junior	257	Primidone	
Pegylated interferon alfa-2a		PKU Anamix Junior Chocolate	257	Primolut N	8
Pembrolizumab		PKU Anamix Junior LQ	257	Priorix	26
Pemetrexed		PKU Anamix Junior Vanilla	257	Probenecid	12
Penicillamine	115	PKU Lophlex LQ 10	257	Probenecid-AFT	12
Penicillin G	96	PKU Lophlex LQ 20		Procaine penicillin	9
PenMix 30	11	PKU Lophlex Powder		Procarbazine hydrochloride	
PenMix 40	11	PKU Lophlex Sensation 20		Prochlorperazine	13
PenMix 50		Plaquenil		Proctofoam	
Pentasa	7	Plendil ER		Proctosedyl	
Pentostatin [Deoxycoformycin]	157	Pneumococcal (PCV10) conjugate		Procyclidine hydrochloride	
Pentoxifylline [Oxpentifylline]		vaccine	269	Procytox	14
Peptamen Junior		Pneumococcal (PCV13) conjugate		Progesterone	
Peptisoothe		vaccine	270	Proglicem	
Peptisorb		Pneumococcal (PPV23)		Proglycem	1
Perhexiline maleate		polysaccharide vaccine	271	Progynova	8
Pericyazine		Pneumovax 23		Prolia	
Perindopril		Podophyllotoxin		Promethazine hydrochloride	
Perjeta	199	Polaramine	224	Propafenone hydrochloride	
Permethrin	72	Poliomyelitis vaccine		Propamidine isethionate	
Perrigo		Poloxamer		Propranolol	
Pertuzumab		Poly-Gel	236	Propylene glycol	
Peteha	104	Poly-Tears		Propylthiouracil	
Pethidine hydrochloride		Poly-Visc		Protaphane	
Pevaryl		Polycal		Protaphane Penfill	
Pexsig		Ponstan		Protifar	
Pfizer Exemestane		Posaconazole		Protionamide	
Pharmacy Services		Postinor-1		Provera	
Pheburane		Potassium chloride49		Provera HD	
Phenasen		Potassium Chloride Aguettant		PSM Citalopram	
Phenobarbitone		Potassium citrate		Psoriasis and Eczema	
Phenobarbitone sodium		Potassium iodate		Preparations	7
Extemporaneous	240	Povidone iodine		PTU	
Nervous		Pradaxa		Pulmicort Turbuhaler	
Phenothrin	72	Pramipexole hydrochloride		Pulmozyme	22
Phenoxybenzamine		Pravastatin		Puri-nethol	
hydrochloride	51	Pravastatin Mylan	60	Puria	3
Phenoxymethylpenicillin (Penicilli		Praziquantel		Pyrazinamide	
V)		Prazosin		Pyridostigmine bromide	
Phenytoin sodium1		Pred Forte		Pyridoxine hydrochloride	
Phillips Milk of Magnesia		Prednisolone		Pyrimethamine	
Phlexy 10		Prednisolone acetate		Pytazen SR	
Phosphate Phebra		Prednisolone sodium		- Q -	
Phosphorus	50	Prednisolone sodium	-	Q 300	10
Phytomenadione	45	phosphate	234	Quetapel	
Pilocarpine hydrochloride	235	Prednisolone-AFT		Quetiapine	
Pimafucort	69	Prednisone		Quick-Set MMT-392	
Pimecrolimus		Pregabalin		Quick-Set MMT-393	
	-	5	-		

Quinapril	52	Rituximab (Riximyo)	201	Serevent	225
Quinapril with		Rivaroxaban		Serevent Accuhaler	225
hydrochlorothiazide	52	Rivastigmine	144	Sertraline	129
Quinine sulphate	102	Rivotril		Setrona	129
Qvar	224	Riximyo	201	Setrona AU	129
- R -		RIXUBIS	44	Sevredol	127
RA-Morph	126	Rizamelt	133	Sex Hormones Non	
Raloxifene hydrochloride	116	Rizatriptan	133	Contraceptive	84
Raltegravir potassium	110	Rolin	170	Shield XL	
Ramipex		Ropin	122	shingles vaccine	272
Ranbaxy-Cefaclor		Ropinirole hydrochloride	122	SII-Onco-BCG	177
Ranitidine		Rotarix		Sildenafil	63
Rapamune	220	Rotavirus oral vaccine	271	Silhouette MMT-373	25
Reandron 1000		Roxane	6	Siltuximab	212
Recombinant factor IX	42, 44	Roxane-Propranolol	55	Simvastatin	
Recombinant factor VIIa	44	Roxithromycin	95	Simvastatin Mylan	60
Recombinant factor VIII	44–45	Rubifen		Sinemet	
Rectogesic	8	Rubifen SR	142	Sinemet CR	
Redipred		Rugby Capsaicin Topical Cream		Sirolimus	220
Relieve		Musculoskeletal	115	Siterone	
Relistor	30	Nervous	124	Slow-Lopresor	55
Remicade		Rulide D	95	Smith BioMed Rapid Pregnancy	
Renilon 7.5		Rurioctocog alfa pegol [Recombin	ant	Test	80
Resonium-A		factor VIII]		Sodibic	
Resource Beneprotein		Ruxolitinib		Sodium acid phosphate	31
Resource Diabetic		Rythmodan		Sodium alginate	6
Respigen		Rytmonorm		Sodium benzoate	
Respiratory Devices		-S-		Sodium bicarbonate	
Respiratory Stimulants		Sabril	132	Blood	49–50
Retinol palmitate		Sacubitril with valsartan		Extemporaneous	
ReTrieve		SalAir		Sodium calcium edetate	
Retrovir		Salazopyrin		Sodium chloride	
Revlimid		Salazopyrin EN		Blood	49
Revolade		Salbutamol		Respiratory	
Rexacrom		Salbutamol with ipratropium		Sodium citrate with sodium lauryl	
Ribomustin		bromide	226	sulphoacetate	31
Ricit		Salicylic acid		Sodium citro-tartrate	
Rifabutin		Salmeterol		Sodium cromoglicate	
Rifadin		Sandomigran		Alimentary	8
Rifampicin		Sandostatin LAR		Respiratory	
Rifaximin		Sanofi Primaquine		Sensory	234
Rifinah		Sapropterin dihydrochloride		Sodium fluoride	
Rilutek		Scalp Preparations		Sodium Fusidate [fusidic acid]	
Riluzole		Scopoderm TTS		Dermatological	66
Riodine		Sebizole		Infection	
Risedronate Sandoz		Secukinumab		Sensory	
Risedronate sodium		Sedatives and Hypnotics		Sodium hyaluronate [Hyaluronic	202
Risperdal Consta		Seebri Breezhaler		acid]	236
Risperidone	136_137	Selegiline hydrochloride		Sodium phenylbutyrate	ري ري
		Senna		Sodium polystyrene sulphonate	5r
Risperidone (Teva)		Senokot		Sodium tetradecyl sulphate	ال
Ritalin		Sensipar		Sodium valproate	120
Ritalin LA		SensoCard	15	Sofradex	
Ritalin SR		Serenace		Soframycin	
Ritonavir		Seretide		Solifenacin Mylan	
Rituximab (Mabthera)		Seretide Accuhaler		Solifenacin succinate	
mituxiiiiau (iviautiieia)	199	Seretiae Accuriater	∠∠0	Somenacin Succinate	ŏ

Solu-Cortef	83	Systane Unit Dose	236	Timolol	
Solu-Medrol		· -Т-		Cardiovascular	56
Solu-Medrol-Act-O-Vial	83	Tacrolimus	222	Sensory	
Somatropin (Omnitrope)	87	Tacrolimus Sandoz	222	Timoptol XE	
Sotalol		Taliglucerase alfa		Tiotropium bromide	
Spacer device		Tambocor		Tiotropium bromide with	
Span-K		Tamoxifen citrate		olodaterol	227
Spiolto Respimat		Tamoxifen Sandoz		Tivicay	
Spiractin		Tamsulosin hydrochloride		TMP	
Spiriva		Tamsulosin-Rex		TOBI	
Spiriva Respimat		Tandem Cartridge		Tobramycin	
		Tandem t:slim X2 with Basal-I		Infection	oc
Spironolactone				Sensory	
Sporanox		Tap water			
Sprycel		Tarceva		Tobramycin BNM	98
Staphlex		Tasigna		Tobramycin Mylan	
Stemetil		Tasmar		Tobrex	
SteroClear		Tecfidera		Tocilizumab	
Stesolid		Tegretol		Tofranil	
Stimulants/ADHD Treatments .		Tegretol CR		Tolcapone	
Stiripentol		Telfast		Topamax	132
Stocrin		Teligent		Topical Products for Joint and	
Stomahesive		Temaccord		Muscular Pain	
Strides Shasun	101	Temazepam	141	Topiderm	
Stromectol	70	Temozolomide	158	Topiramate	132
Sucralfate	10	Tenofovir disoproxil	104	Topiramate Actavis	132
Sulfadiazine Silver	66	Tenofovir Disoproxil Teva	104	Total parenteral nutrition (TPN)	49
Sulfadiazine sodium	99	Tenoxicam	114	TPN	49
Sulfasalazine	8	Tensipine MR10	57	Tramadol hydrochloride	128
Sulindac	114	Tepadina	150	Tramal SR 100	
Sulindac Mylan	114	Terazosin	51	Tramal SR 150	128
Sulphur		Terbinafine		Tramal SR 200	128
Sulprix		Terbutaline sulphate		Trandate	55
Sumatriptan		Teriflunomide		Tranexamic acid	
Sunitinib		Teriparatide		Tranylcypromine sulphate	
Sunscreens		Testosterone		Trastuzumab	
Sunscreens, proprietary		Testosterone cipionate		Trastuzumab emtansine	
Sure-T MMT-863		Testosterone esters		Travatan	
Sure-T MMT-873		Testosterone undecanoate		Travoprost	
Sustagen Diabetic		Tetrabenazine		Travopt	
Sustagen Hospital Formula	240	Tetrabromophenol		Treatments for Dementia	
Active	252	Tetracosactrin		Treatments for Substance	144
		Tetracycline		Dependence	1/15
Sustanon Ampoules		•			
Sutent		Thalidomide		Trental 400	02
Sylvant		Thalomid		Tretinoin	C.F
Symbicort Turbuhaler 100/6		Theophylline		Dermatological	65
Symbicort Turbuhaler 200/6		Thiamine hydrochloride		Oncology	
Symbicort Turbuhaler 400/12		THIO-TEPA		Trexate	152
Symmetrel		Thioguanine		Triamcinolone acetonide	
Sympathomimetics		Thiotepa		Alimentary	36
Synacthen		Thymol glycerin		Dermatological	
Synacthen Depot	84	Thyroid and Antithyroid Agents		Hormone	84
Synacthene Retard		Ticagrelor		Triamcinolone acetonide with	
Synflorix		Tilade		gramicidin, neomycin and nyst	
Synthroid		Tilcotil	114	Dermatological	
Syntometrine		Tillomed	149	Sensory	232
Syrup (pharmaceutical grade).	241				

Triover	0.4	Vanalovta	150	XMET Maxamum	256
Triaver Triazolam		Venclexta		Xolair	
Trimethoprim		VenetoclaxVenlafaxine		XP Maxamum	
	99	Venomil		Xylocaine	
Trimethoprim with		VENOX			
sulphamethoxazole	100	Ventavis		Xylocaine 2% JellyXyntha	
[Co-trimoxazole]		Ventolin		- Z -	44
Trisequens		VentoiiiiVepesid		Zantac	
Trophic Hormones		Verapamil hydrochloride		Zapril	
Tropicamide		Vergo 16		Zarontin	
Trusopt		Vermox		Zaroxolyn	
TruSteel		Versacloz		Zavedos	
Tuberculin PPD [Mantoux] tes		VersaciozVesanoid		Zeffix	
Tubersol		Vexazone		Zetlam	
Two Cal HN		Vfend		Ziagen	
Two Cal HN RTH		Viaderm KC		Zidovudine [AZT]	
		Vidaza			108
Tykerb				Zidovudine [AZT] with	440
Tysabri	140	Vigabatrin		lamivudine	
	007	Vildagliptin	13	Zimybe	
Ultibro Breezhaler		Vildagliptin with metformin	10	Zinc and castor oil	
Ultraproct		hydrochloride		Zinc sulphate	
Umeclidinium		Vimpat		Zincaps	
Umeclidinium with vilanterol		Vinblastine sulphate		Zinnat	
Univent	,	Vincristine sulphate		Ziprasidone	
Ural		Vinorelbine		Zista	
Urea		Vinorelbine Ebewe		Zithromax	
Urex Forte		Viramune Suspension		Zoladex	91
Urinary Agents		ViruPOS		Zoledronic acid	00
Urinary Tract Infections		Vit. D3		Hormone	
Urinorm		Vita-B12		Musculoskeletal	
Uromitexan		VitA-POS		Zoledronic acid Mylan	
Ursodeoxycholic acid		Vitabdeck		Zopiclone	
Ursosan		Vital		Zopiclone Actavis	141
Utrogestan	86	Vitamin B complex		Zostavax	
- V -		Vitamin B6 25		Zostrix	
Vaccinations		Vitamins		Zostrix HP	
Vaclovir		Vivonex TEN		Zuclopenthixol decanoate	
Valaciclovir		Voltaren		Zuclopenthixol hydrochloride	
Valganciclovir		Voltaren D		Zusdone	
Valganciclovir Mylan		Voltaren Ophtha		Zyban	
Vancomycin		Volumatic		Zypine	136
Vannair		Voriconazole		Zypine ODT	
Varenicline Pfizer		Vosol		Zyprexa Relprevv	
Varenicline tartrate		Votrient		Zytiga	167
Varicella vaccine [Chickenpox		Vttack	101		
vaccine]		- W -			
Varicella zoster virus (Oka str		Warfarin sodium			
attenuated vaccine [shingle		Wart Preparations			
vaccine]		Wasp venom allergy treatmen	t223		
Various		Water			
Varivax		Blood			
Vasodilators		Extemporaneous			
Vasopressin Agonists		Wool fat with mineral oil	70		
Vasorex		- X -			
Vedafil		Xarelto			
Veletri	64	Xifaxan	10		

