May 2021 Volume 28 Number 1

Editors:

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

_	introducing i manimac	
5	General Rules	Section A
6	Alimentary Tract & Metabolism	Section B
37	Blood & Blood Forming Organs	
47	Cardiovascular System	
61	Dermatologicals	
71	Genito Urinary System	
78	Hormone Preparations – Systemic	
89	Infections – Agents For Systemic Use	
110	Musculoskeletal System	
118	Nervous System	
144	Oncology Agents & Immunosuppressants	
220	Respiratory System & Allergies	
229	Sensory Organs	
235	Various	
237	Extemporaneous Compounds (ECPs)	Section C
240	Special Foods	Section D
261	National Immunisation Schedule	Section I
273	Index	

Introducing PHARMAC

Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	✓	Gaviscon Infant
SODIUM ALGINATE Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	✓	Entocort CIR
⇒SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
1 Mild to moderate ileal, ileocaecal or proximal Crohn's dise	ease; and			

0.4 Dishetes as

2 Any of the following:

2.1 Diabetes; or

2.2 Cushingoid habitus; or

2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)			Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

TITUROCORTISONE ACETATE			
Rectal foam 10%, CFC-Free (14 applications)	26.55	21.1 g OP	Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDRO	CHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 19	%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg	49.50	100	✓ Asamax
Tab long-acting 500 mg	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g		100 OP	✓ Pentasa
Enema 1 g per 100 ml	41.30	7	✓ Pentasa
Suppos 500 mg		20	✓ Asacol
Suppos 1 g		28	✓ Pentasa
OLSALAZINE			
Tab 500 mg	93.37	100	✓ Dipentum
Cap 250 mg	53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mgSULFASALAZINE	92.91	100	✓	Nalcrom
* Tab 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g6.35	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg2.66	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy 30 q OP ✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

* Tab 200 mcg - Up to 120 tab available on a PSO41.50 ✓ Cytotec

Subsidy (Manufacturer's Price)			Brand or Generic	
<u> </u>	Per		Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg − Subsidy by endorsement......10.40 14 ✓ Apo-Clarithromycin

- a) Maximum of 14 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

H2 Antagonists

FΑ	MOTIDINE - Only on a prescription			
*	Tab 20 mg	4.91	100	✓ Famotidine Hovid S29
*	Tab 40 mg	8.48	100	✓ Famotidine Hovid \$29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement			•
	Subsidy by endorsement – Subsidised for patients receiving t	reatment as	part of pallia	ıtive care.

RANITIDINE - Subsidy by endorsement

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the
 prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record
 of prior dispensing of ranitidine.

*	Oral liq 150 mg per 10 ml5.14	4 300 ml	•	Peptisoothe
*	Inj 25 mg per ml, 2 ml13.40	5	1	Zantac
(Pa	entisoothe Oral lig 150 mg per 10 ml to be delisted 1 September 2021)			

(Zantac Inj 25 mg per ml, 2 ml to be delisted 1 June 2021)

Proton Pump Inhibitors

LANSOPRAZOLE		
* Cap 15 mg4.58	100	✓ Lanzol Relief
* Cap 30 mg5.41	100	✓ Lanzol Relief
OMEPRAZOLE		
For omeprazole suspension refer Standard Formulae, page 237		
* Cap 10 mg	90	✓ Omeprazole actavis
		10
Omeprazole actavis 10 to be Sole Supply on 1 August 2021		
Cap 20 mg	90	✓ Omeprazole actavis
54p 20 mg	00	20
Omeprazole actavis 20 to be Sole Supply on 1 August 2021		
Cap 40 mg	90	✓ Omeprazole actavis
oup 40 mg	00	40
Omeprazole actavis 40 to be Sole Supply on 1 August 2021		40
* Powder – Only in combination	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspension.	Jy	Wildwest
* Inj 40 mg ampoule with diluent33.98	5	✓ Dr Reddv's
Till 40 filly ampoule with undertained	3	Omeprazole
		Onicpi azole

✓ Ocicure \$29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PANTOPRAZOLE				
₭ Tab EC 20 mg		100		Panzop Relief
★ Tab EC 40 mg	2.85	100	•	Panzop Relief
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE				
Tab 120 mg	14.51	50	1	Gastrodenol S29
SUCRALFATE				
Tab 1 g		120		
	(48.28)			Carafate
Bile and Liver Therapy				
RIFAXIMIN - Special Authority see SA1461 below - Reta	ail pharmacy			
Tab 550 mg	625.00	56	1	Xifaxan
enending from treatment.				
Diabetes				
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret	· ·			
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg	110.00	100		Proglicem \$29
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg	110.00	100	✓	Proglicem \$29
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg	110.00		✓	-
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 ml C e use	DP ✓	Proglicem \$29 Proglycem \$29 reatment of confirmed
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 ml C e use	DP ✓	Proglicem \$29 Proglycem \$29 reatment of confirmed
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 ml C e use	DP d for the tr	Proglicem \$29 Proglycem \$29 reatment of confirmed
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 o ml C e use s noti	DP d for the tr	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 o ml C e use s noti	DP d for the tr	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 o ml C e use s noti	DP d for the tri ffied where	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains Glucagen Hypokit Actrapid
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 o ml C e use s noti	opp of d for the transfer depends on the transfer depe	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains Glucagen Hypokit
Diabetes Hyperglycaemic Agents DIAZOXIDE – Special Authority see SA1320 below – Ret Cap 25 mg		100 ml C e use s noti	opp of d for the transfer depends on the transfer depe	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains Glucagen Hypokit Actrapid Humulin R Actrapid Penfill
Diabetes Hyperglycaemic Agents DIAZOXIDE - Special Authority see SA1320 below - Ret Cap 25 mg		100 ml C e use s noti	opp of d for the transfer depends on the transfer depe	Proglicem \$29 Proglycem \$29 reatment of confirmed the treatment remains Glucagen Hypokit Actrapid Humulin R Actrapid Penfill

	Subsidy	F	ully Brand or
	(Manufacturer's Price	e) Subsidis	,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			
Inj human 100 u per ml	17.68		✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
- , ,			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
•			✓ PenMix 30
			✓ PenMix 40
JOHN N. JORDO WITH INCH IN JORDO DROTAMINE			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml.			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml			
3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml			✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
Inj 100 u per ml, 3 ml			✓ Apidra
Inj 100 u per ml, 3 ml disposable pen			✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml		10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
★ Tab 50 mg		90	✓ Glucobay
₭ Tab 100 mg	10.47 6.40	90	✓ Accarb✓ Glucobay
- 145 155 Hig	20.23	00	✓ Accarb
Oral Hypoglycaemic Agents			
ilibenclamide			
₭ Tab 5 mg	6.00	100	✓ <u>Daonil</u>
			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	-	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
GLICLAZIDE				
* Tab 80 mg	15.18	500	1	<u>Glizide</u>
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000	✓	<u>Apotex</u>
* Tab immediate-release 850 mg	7.04	500	•	Apotex
PIOGLITAZONE				
* Tab 15 mg	1.12	28	✓	<u>Vexazone</u>
	3.47	90	_	Vexazone
* Tab 30 mg		90	_	<u>Vexazone</u>
* Tab 45 mg	7.10	90	•	<u>Vexazone</u>
VILDAGLIPTIN			_	
Tab 50 mg	35.00	60	/	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride		60	_	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	/	Galvumet

SGLT2 Inhibitors

⇒SA2029 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Any of the following:
 - 2.1 Patient is Maaori or any Pacific ethnicity*: or
 - 2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.5 Patient has diabetic kidney disease (see note b)*; and
- 3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months; and
- 4 Treatment will not be used in combination with a funded GLP-1 agonist.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

ΕN	IPAGLIFLOZIN - Special Authority see SA2029 above - Retail	pharmacy		
*	Tab 10 mg	58.56	30	✓ Jardiance
*	Tah 25 mg	58 56	30	✓ lardiance

Fully

Brand or

	(Manufacturer's Price)	Sul Per	bsidised	Generic Manufacturer	
EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Sp	pecial Authority see S		n the pre		ail
* Tab 5 mg with 1,000 mg metformin hydrochloride * Tab 5 mg with 500 mg metformin hydrochloride * Tab 12.5 mg with 1,000 mg metformin hydrochloride * Tab 12.5 mg with 500 mg metformin hydrochloride	58.56 58.56	60 60 60	✓ J.	ardiamet ardiamet ardiamet ardiamet	

Subsidy

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test 1 OP ✓ CareSens N ✓ CareSens N POP 20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

50 test OP CareSens N CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	1	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES	 Maximum of 200 	dev per prescription
---------------------	------------------------------------	----------------------

	00 407	40.50	400	/ D D 14:
*	29 g × 12.7 mm	10.50	100	B-D Micro-Fine
*	31 g × 5 mm	11.75	100	 B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm	10.50	100	✓ B-D Micro-Fine
*			100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200) dev per pre	escription
	Syringe 0.3 ml with 29 g × 12.7 mm needle		100	✓ B-D Ultra Fine
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle	` '	100	✓ B-D Ultra Fine
•	-,···g- ··· ··· g · · · ··· · · · ·	1.30	10	
		(1.99)	. •	B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
•	Symigo do mi mar or give min nocale minimini	1.30	10	
		(1.99)	. •	B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	` '	100	✓ B-D Ultra Fine
•	5)g5 : =5 g × .=	1.30	10	2 2 0 0
		(1.99)	. •	B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
-1-	Syrings This High of g A o Hill Hoodio	1.30	100	- D D OMAT HICH
			10	D D I III Time II
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year p	eriod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
•	Por 🗸	Manufacturer	

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Suk	sidised	Generic	
(Manuacturer 3 i lice)	Out	Joiuiseu	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
(Manufacturer's Frice)	Per 🗸	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol: and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 2 Eithor
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
pump therapy; and				
4 The patient is continuing to derive benefit from pump the			., .	
5 The patient had achieved and is maintaining a HbA1c o 6 The patient has had no increase in severe unexplained				
7 The patient's HbA1c has not deteriorated more than 5 n	,, ,, ,		iii baseiiile, a	allu
8 Either:	into/mor from baseline	, and		
8.1 Applicant is a relevant specialist; or				
8.2 Applicant is a nurse practitioner working within the	neir vocational scope.			
Renewal — (Previous use before 1 September 2012) only for	rom a relevant specialis	st or n	urse practitio	oner. Approvals valid for 2
years for applications meeting the following criteria:				
All of the following:				
1 The patient is continuing to derive benefit according to t than 80 mmol/mol; and	he treatment plan and	has m	naintained a l	HbA1c of equal to or less
2 The patient's HbA1c has not deteriorated more than 5 n	amol/mol from initial an	nlicat	ion: and	
3 The patient has not had an increase in severe unexplain				ne: and
4 Either:				,
4.1 Applicant is a relevant specialist; or				
4.2 Applicant is a nurse practitioner working within the control of the contro	neir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 18 – Retail p	harma	асу	
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 packs of cartridge sets will be funded p		4 00		andan Oastaldaa
Cartridge 300 U, t:lock × 10		1 OP		andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Specia	al Authority see SA198	5 on p	oage 18 – Re	etail pharmacy
a) Maximum of 3 sets per prescription b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T
g				MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	• • • M	liniMed Sure-T
Commented mandles 00 and taking a 40	100.00	4.00		MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	• W	liniMed Sure-T

8 mm steel needle; 80 cm tubing × 10130.00

6 mm steel needle; 29 G; manual insertion; 60 cm tubing x

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

1 OP

1 OP

1 OP

1 OP

MMT-866A

MMT-874A

✓ MiniMed Sure-T

✓ MiniMed Sure-T

MMT-876A

✓ Sure-T MMT-863

✓ Sure-T MMT-873

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 18 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00) 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.00	0 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 18 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- a) Maximum of 12 infusion sate will be funded nor year

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A
			WINI I -OOOA

	Subsidy (Manufacturer's Pr	rice) S	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SERTION WITH	H INSERTI	ON DEVIC	E) - Special Authority see
SA1985 on page 18 – Retail pharmacy				
a) Maximum of 3 sets per prescription b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c	cm			
line × 10 with 10 needles		1 OP	✓ A	utoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cr		4.00		
line × 10 with 10 needles		1 OP		utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	NSERTION) - S	pecial Auth	ority see S	A1985 on page 18 –
Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles; luer lock		1 OP	_	ilhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	IT INSERTION V	VITH INSE	RTION DE	VICE) - Special Authority
see SA1985 on page 18 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 c line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device;	140.00	1 01	• ^	atooon so
110 cm line × 10 with 10 needles	140.00	1 OP	✓ A	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m			
line × 10 with 10 needles		1 OP		utoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	IT INSERTION)	- Special	Authority se	e SA1985 on page 18 -
Retail pharmacy				
a) Maximum of 3 sets per prescription b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wit	:h			
10 needles; luer lock		1 OP	√ Q	uick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wit		1.00		huiale Cat MMT 000
10 needles; luer lock		1 OP		uick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 of	n page 18 – Heta	aii pharmad	У	
a) Maximum of 3 sets per prescription b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per	year.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP		DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓ M	liniMed
				1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ M	liniMed
2		. 0.	- 10	3.0 Reservoir
				MMT-332A

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
· · · ·	Por 🗸	Manufacturor	

Digestives Including Enzymes

PANCREATIC ENZYME

PANCREATIC ENZYME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase,			
1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	Creon Micro
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	v – Retail pha	rmacy	
Cap 250 mg	32.95	100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Normacol Plus

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soin12.20	500 g OP	Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		
* Dry	2 500 g OP	
(17.32	2)	Normacol Plus
2.41	200 a OP	

(8.72)

Faecal Softeners

* Tab 50 mg2.31	100	✓ Coloxyl
* Tab 120 mg3.13		✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES		
* Tab 50 mg with sennosides 8 mg	200	✓ Laxsol
POLOXAMER – Only on a prescription		
Not funded for use in the ear.		
* Oral drops 10%3.98	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority see	SA1691 below - Retail	pharmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
•	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	(Manufacturer's Pric		osidised Generic	
	\$	Per	✓ Manufacturer	_
Osmotic Laxatives				
GLYCEROL				
* Suppos 3.6 g – Only on a prescription	9.25	20	✓ <u>PSM</u>	
LACTULOSE – Only on a prescription				
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ <u>Laevolac</u>	
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI		SODIUM C	CHLORIDE	
Powder for oral soln 13.125 g with potassium chloride 46.6 n		20		
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg6.70	30	✓ <u>Molaxole</u>	
SODIUM ACID PHOSPHATE – Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema	
CODILINA CITRATE MITH CODILINA LALIDVI. CHII DUCACETATE	Only an a muse		Enema	
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE	, ,	inpuon		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml		50	✓ Micolette	
0111	20.00	00	· imoolette	
Stimulant Laxatives				
BISACODYL - Only on a prescription				
* Tab 5 mg	5.99	200	✓ <u>Lax-Tab</u>	
* Suppos 10 mg	3.74	10	✓ <u>Lax-Suppositories</u>	
SENNA - Only on a prescription				
* Tab, standardised		100		
	(8.21)	20	Senokot	
	0.43	20	Senokot	
	(2.06)		Seriokot	

Subsidy

Fully

Brand or

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and

Subsidy	Fully	Brand or
acturer's Price)	Subsidised	Generic
 \$ Pe	er 🗸	

continued...

- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority	see SA2042 below -	- Retail pharmacy
------------------------------	--------------------	-------------------

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 below - Retail pharmacy

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10	 Special Authority see 	SA2039 on the nex	t page – Retail pharm	acy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below	- Retail pharmacy		
Tab 500 mg	CBŚ	30	Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – Re	tail pharmacy		
Tab 100 mg	CBS	100	Country Life
Cap 100 mg	CBS	100	✓ Solgar

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE	- Special Authority see SA1989 below -	- Retail pharmacy	
Tab soluble 100 mg	1,452.70	30 OP	✓ Kuvan

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and

	Subsidy		Fully	Brand or
(I	Manufacturer's Price)	Subsic	lised	Generic
	\$	Per	✓	Manufacturer

continued...

- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1990 below – Retail pharmacy
Grans 483 mg per g.......2,016.00 174 g OP

Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Fully

Subsidy (Manufacturer's Price)

Subsidised Per

1

Brand or Generic Manufacturer

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

✓ Elelyso

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia. thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse: chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose: and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication.

ALIMENTARY TRACT AND METABOLISM	Subsidy (Manufacturer's Price)	Fully Subsidised Per	
continued and/or adjustment of infusion rates; and 5) Patient has not developed another medical condition th ERT; and 6) Patient is compliant with regular treatment and taligluce every other week rounded to the nearest whole vial (20 7) Supporting clinical information including test reports, M investigations are submitted to the Gaucher Panel for a	erase alfa is to be admin 0 units), unless otherwi RI whole body STIR, ha	istered at a dos se agreed by F	e no greater than 30 unit/kg HARMAC; and
Mouth and Throat Agents Used in Mouth Ulceration			
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with Endorsement	(20.31)	500 ml esult of treatme	Difflam ent for cancer, and the

	(20.31)		Difflam
Additional subsidy by endorsement for a patient who ha prescription is endorsed accordingly.	s oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	•	Orabase
	1.52	5 g OP	
	(3.60)	_	Orabase
Powder	8.48	28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
7 tanooro goro /o mar ootamornam omonao o.or/o	(6.00)	10 g 01	Bonjela
TRIAMCINOLONE ACETONIDE	(0.00)		201,010
Paste 0.1%	F 00	F ~ OD	/ Kanalas in Osahaga
Paste 0.1%	5.33	5 g OP	 Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			3
	4.74	40 g OP	✓ Decozol
Oral gel 20 mg per g	4.74	40 g OF	<u>Decozoi</u>
NYSTATIN			4
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>
Other Oral Agents			

For folinic mouthwash, pilocarpine oral liquid or saliva substitute formula refe	r Standard Formulae, page 237
THYMOL GLYCERIN	
* Compound, BPC9.15	5 500 ml ✓ PSM

Fully

Brand or

Subsidy

	(Manufacturer's Price)		sed Generic Manufacturer
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	SO1.89	3	✓ <u>Neo-B12</u> ✓ Vita-B12
	3.15	5	 Hydroxocobalamin Mercury Pharma
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription			
* Tab 25 mg – No patient co-payment payable * Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Apo-Pyridoxine
THIAMINE HYDROCHLORIDE - Only on a prescription			
* Tab 50 mg VITAMIN B COMPLEX	7.09	100	✓ Max Health
* Tab, strong, BPC	7.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription	0.00	500	. Outh
* Tab 100 mg	9.90	500	✓ <u>Cvite</u>
Vitamin D			
ALFACALCIDOL * Cap 0.25 mcg * Cap 1 mcg * Oral drops 2 mcg per ml	87.98	100 100 0 ml OP	✓ One-Alpha ✓ One-Alpha ✓ One-Alpha
CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg		100 100	✓ <u>Calcitriol-AFT</u> ✓ <u>Calcitriol-AFT</u>
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) — Maximum of 12 cap per prescripti * Oral liq 188 mcg per ml (7,500 iu per ml)		12 .8 ml OP	✓ <u>Vit.D3</u> ✓ Puria
Multivitamin Preparations			
MULTIVITAMIN RENAL – Special Authority see SA1546 below – * Cap		30	✓ Clinicians Renal Vit
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either:			
1 The patient has chronic kidney disease and is receiving ei			dialysis; or

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of <

15 ml/min/1.73 m² body surface area (BSA).

ALIMENTARY TRACT AND METABOLISM				
	Subsidy (Manufacturer's Prio \$	ce) Subs	Fully idised	Brand or Generic Manufacturer
MULTIVITAMINS – Special Authority see SA1036 below – Reta	ail pharmacy			
* Powder	72.00	200 g OP	✓ P	aediatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valinborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid withou				·
approval for multivitamins.			·	·
VITAMINS				
* Tab (BPC cap strength)		1,000	✓ <u>N</u>	<u>lvite</u>
* Cap (fat soluble vitamins A, D, E, K) - Special Authority se SA1720 below - Retail pharmacy		60	✓ ∨	itabdeck
⇒SA1720 Special Authority for Subsidy	20.10	00	•	nabaoon
Initial application from any relevant practitioner. Approvals varithe following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; of 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome.	r	aigwai uiiless	noune	o to applications meeting
Minerals Calcium				
CALCIUM CARBONATE				
* Tab 1.25 g (500 mg elemental)	6.69	250	✓ C	alci-Tab 500
* Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme		76	_	Cacit S29
Subsidy by endorsement – Only when prescribed for paconsidered unsuitable.		5 years) whe	ere calci	ium carbonate oral liquid is
CALCIUM GLUCONATE	00.00	10		lavi lia aliib
* Inj 10%, 10 ml ampoule	32.00	10	• 10	lax Health - Hameln S29
	64.00	20	✓ N	lax Health S29
Fluoride				
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	√ P	SM
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	leuroTabs
Iron				

FERRIC CARBOXYMALTOSE - Special Authority see SA1840 on the next page - Retail pharmacy Inj 50 mg per ml, 10 ml......150.00 ✓ Ferinject

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or egual to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:

MAGNESIUM HYDROXIDE

- 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
- 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
- 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
- 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	100	✓ <u>Ferro-tab</u>
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg4.68	60	✓ <u>Ferro-F-Tabs</u>
FERROUS SULFATE * Oral liq 30 mg (6 mg elemental) per 1 ml12.08	500 ml	✓ Ferodan
FERROUS SULPHATE * Tab long-acting 325 mg (105 mg elemental)2.06	30	✓ <u>Ferrograd</u>
IRON POLYMALTOSE * Inj 50 mg per ml, 2 ml ampoule34.50	5	✓ Ferrosig
Magnesium		
For magnesium hydroxide mixture refer Standard Formulae, page 237		

Suspension 8%	33.60	355 ml	✓ Phillips Milk of
			Magnesia S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MAGNESIUM SULPHATE				
* Inj 2 mmol per ml, 5 ml ampoule	25.53 28.00	10	✓	Martindale DBL DBL S29 S29
Martindale to be Sole Supply on 1 July 2021 (DBL Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 July 2021) (DBL S29 S29 Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 J	July 2021)			
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	/	<u>Zincaps</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per		Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу		
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	•	Binocrit

Megaloblastic

-01	10	40	
-OL	_IC	AC	טו

*	Tab 0.8 mg21.84	1,000	1	Apo-Folic Acid
	Tab 5 mg12.12	500	1	Apo-Folic Acid
	Oral lig 50 mcg per ml	25 ml OP	1	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
lnj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 Wastage claimable	below - Retail pharmacy		

Tab 50 mg3,100.00 **➤ SA1743** Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

continued...

✓ Revolade

✓ Revolade

28 28

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
 \$	Per	✓	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

✓ Hemlibra	1	vial3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	ıl vial	Inj 60 mg in 0.4 ml via
✓ Hemlibra	1	ml vial12,492.00	Inj 105 mg in 0.7 ml vi
✓ Hemlibra	1	vial17,846.00	Inj 150 mg in 1 ml vial

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:

continued...

				
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Su	ıbsidised	Generic
	\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Subject to criteria.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

NONACOG GAMMA. [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

nj 500 iu vial	435.00	1	✓ RIXUBIS
nj 1,000 iu vial	870.00	1	✓ RIXUBIS
nj 2,000 iu vial		1	✓ RIXUBIS
nj 3,000 iu vial	·	1	✓ RIXUBIS
) -/ · · · · · · · · · · · · · · · · ·	,		

	0.1		- ·	
	Subsidy (Manufacturer's Price)	Su	Fully bsidised	
	\$	Per	√	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	[Xpharm]			
For patients with haemophilia. Preferred Brand of short half	-life recombinant factor	r VIII. A	ccess to	o funded treatment is
managed by the Haemophilia Treaters Group in conjunction				
Inj 250 iu vial		1		Advate
Inj 500 iu vial	420.00	1	✓	Advate
Inj 1,000 iu vial	840.00	1	✓	Advate
Inj 1,500 iu vial	1,260.00	1	✓	Advate
Inj 2,000 iu vial		1		Advate
Inj 3,000 iu vial	,	1	/	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the I	Vational	Haemo	philia Management Group,
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1	•	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatm		d treatm	ent is n	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			_	
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial	· ·	1		Adynovate
Inj 2,000 iu vial	2,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Konakion MM
ing to mg por mi, i mi op to o mg available on a room		•		Tronauton iiiii
Antithrombotic Agents				
- Tillian Omsons Algente				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	•	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
				<u>Multichem</u>
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	1	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next pag				-
* Tab 90 mg		56	1	Brilinta

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

				=
Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

10

10

10

10

10

Fully

Brand or

✓ Clexane

✓ Clexane

✓ Clexane

✓ Clexane Forte

✓ Clexane Forte

	(Manufacturer's Price) \$	Sub Per		Generic Manufacturer	
Heparin and Antagonist Preparations					
ENOXAPARIN SODIUM - Special Authority see SA1646 below	, ,				
Inj 20 mg in 0.2 ml syringe	27.93	10	✓ Cle	xane	
Inj 40 mg in 0.4 ml syringe	37.27	10	✓ Cle	xane	

Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Low molecular weight heparin treatment is required during a patients pregnancy; or

Inj 60 mg in 0.6 ml syringe......56.18

Inj 80 mg in 0.8 ml syringe......74.90

Inj 120 mg in 0.8 ml syringe.......116.55

- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery: or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	✓ Hospira
	42.40		✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg	76.36	60	✓	Pradaxa
Cap 150 mg		60	1	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	✓	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	1	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
WARFARIN SODIUM Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	1	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg		100	1	Marevan
* Tab 5 mg		50	✓	Coumadin
•	11.48	100	1	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail p	harmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe		10	1	Nivestim
OA4050 On a stat Authority for Outside.				

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC $< 0.5 \times 10^9$ /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy	,	Fully Brand or
	(Manufacturer's Price \$	e) Per	Subsidised Generic Manufacturer
	Ψ	1 01	Wandadard
Fluids and Electrolytes			
Intravenous Administration			
GLUCOSE [DEXTROSE]			
* Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO	30.65	5	✓ Biomed
★ Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO	15.00	1	✓ <u>Biomed</u>
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	55.00	50	✓ AstraZeneca
			Potassium Chloride
			Aguettant S29
	65.00		✓ Juno
AstraZeneca Inj 75 mg per ml, 10 ml to be delisted 1 November	r 2021)		
Potassium Chloride Aguettant S29 Inj 75 mg per ml, 10 ml to l	be delisted 1 Novemb	ber 202	21)
SODIUM BICARBONATE			,
Inj 8.4%, 50 ml	19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO		-	
b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebulise	er use except when u	sed in o	conjunction with an antibiotic intenden
for nebuliser use.			,
Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.23	500 m	l ✓ Baxter
	1.26	1,000 n	ml ✓ Baxter
Only if prescribed on a prescription for renal dialysis, m for emergency use. (500 ml and 1,000 ml packs)	aternity or post-natal	care in	n the home of the patient, or on a PS
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standa			
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi
OTAL PARENTERAL NUTRITION (TPN)			
Infusion	CBS	1 OP	✓ TPN
VATER			
1) On a prescription or Practitioner's Supply Order only v	when on the same for	m as a	n injection listed in the Pharmaceuti
Schedule requiring a solvent or diluent; or			
On a bulk supply order; or			
3) When used in the extemporaneous compounding of e			
4) When used for the dilution of sodium chloride soln 7%	tor cystic tibrosis pa	tients o	only.
lai E ml amagula . Un to E ini available an a BOO	7.00	F0	✓ InterPharma
Inj 5 ml ampoule – Up to 5 inj available on a PSOInj 10 ml ampoule – Up to 5 inj available on a PSO		50 50	✓ InterPharma ✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO		20	✓ Frizer ✓ Fresenius Kabi
ing 20 mil ampoule – op to 3 mil avaliable on a F30		20	✓ Multichem
	7.50	30	✓ InterPharma
(InterPharma Inj 5 ml ampoule to be delisted 1 June 2021)		50	
InterPharma Inj 20 ml ampoule to be delisted 1 June 2021)			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully Brand or idised Generic Manufacturer	
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ Calcium Resoniur	m
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	9.77	50	✓ <u>Electral</u>	
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)	•	1,000 ml OP	✓ <u>Pedialyte -</u> Bubblegum	
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓ Phosphate Phebra	а
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (11.85)	60	Chlorvescent	
* Tab long-acting 600 mg (8 mmol)	8.90 [′]	200	✓ Span-K	
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	✓ Sodibic✓ Sodibic	
SODIUM POLYSTYRENE SULPHONATE				

✓ Resonium-A

454 g OP

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised ✓	Brand or Generic Manufacturer	
Alpha-Adrenoceptor Blockers	·				
Alpha Adrenoceptor Blockers					
DOXAZOSIN * Tab 2 mg * Tab 4 mg PHENOXYBENZAMINE HYDROCHLORIDE		500 500		po-Doxazosin po-Doxazosin	
* Cap 10 mg	65.00 216.67	30 100		NM S29 ibenzyline S29	

TERAZOSIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking terazosin prior to 1 October 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of terazosin.

Tab 2 mg	500
14.20	28
Tab 5 mg10.90	500
24.80	28

✓ Apo-Terazosin

✓ Apo-Prazosin

✓ Apo-Prazosin✓ Apo-Prazosin

✓ Teva S29

✓ 7anril

100

100

100

✓ Apo-Terazosin
✓ Teva S29

(Apo-Terazosin Tab 2 mg to be delisted 1 August 2021)

(Teva S29 Tab 2 mg to be delisted 1 August 2021)

(Apo-Terazosin Tab 5 mg to be delisted 1 August 2021)

(Teva S29) Tab 5 mg to be delisted 1 August 2021)

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRII			

Tah 0.5 mg

D

PRAZOSIN

*	Oral liq 5 mg per ml94.99	95 ml OP	✓ Capoten
	135.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

~	1ab 0.5 mg	.00	•	Zapili
*	Tab 2.5 mg4	.80	90 🗸	Zapril
	Tab 5 mg8		90	Zapril
ΕN	ALAPRIL MALEATE			
*	Tab 5 mg1	.82 1	00 🗸	Acetec
	Tab 10 mg2		00 🗸	Acetec
	Tab 20 mg2		00 🗸	Acetec

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

=					
		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
LIS	INOPRIL				
_	Tab 5 mg	2.07	90	1	Ethics Lisinopril
	Tab 10 mg		90		Ethics Lisinopril
-	Tab 20 mg		90		Ethics Lisinopril
	v		00		<u> </u>
PΕ	RINDOPRIL	0.75	~~	,	An a Bankada adl
	Tab 2 mg		30		Apo-Perindopril
	Tab. Assess	4.95	~~		Coversyl
	Tab 4 mg		30		Apo-Perindopril
		6.30		•	Coversyl
QU	INAPRIL				
	Tab 5 mg	6.01	90	✓	Arrow-Quinapril 5
	Tab 10 mg	3.16	90	✓	Arrow-Quinapril 10
	Tab 20 mg	4.89	90	✓	Arrow-Quinapril 20
Α	CE Inhibitors with Diuretics				
QU	INAPRIL WITH HYDROCHLOROTHIAZIDE				
	Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	✓	Accuretic
		3.83	30	✓	Accuretic 10
*	Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	✓	Accuretic 20
A	ngiotensin II Antagonists				
CA	NDESARTAN CILEXETIL				
*	Tab 4 mg	1.90	90	✓	Candestar
*	Tab 8 mg		90		Candestar
*	Tab 16 mg		90		Candestar
	Tab 32 mg		90		Candestar
	SARTAN POTASSIUM				
_		1 56	84	./	Lacartan Actovia
	Tab 12.5 mg		84		Losartan Actavia
	Tab 25 mg		84		Losartan Actavia
	Tab 50 mg				Losartan Actavis
*	Tab 100 mg	3.50	84	•	Losartan Actavis
A	ngiotensin II Antagonists with Diuretics				
10	SARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
LO	Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	1	Arrow-Losartan &
	rab 50 mg with hydrochlorothazide 12.5 mg	1.00	00	•	Hydrochlorothiazide
					<u> </u>
Α	ngiotensin II Antagonists with Neprilysin Inhib	oitors			
-	g				
SA	CUBITRIL WITH VALSARTAN - Special Authority see SA190				
	Note: Due to the angiotensin II receptor blocking activity of s	sacubitril with valsarta	n it s	hould not	be co-administered with an
	ACE inhibitor or another ARB.				
	Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓	Entresto 24/26
	Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓	Entresto 49/51
	Tab 97.2 mg with valsartan 102.8 mg		56	✓	Entresto 97/103
	•				

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOLIS SYSTEM. Anaesthetics, Local, page 118

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaestnetics, Local,	page 118	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSO	10	✓ Max Health
	10	wax nealth
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		
PSO12.07	10	✓ Hameln S29
		✓ Martindale
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	240	✓ Lanoxin
* Oral lig 50 mcg per ml	60 ml	✓ Lanoxin
* Oral liq 50 micg per mi10.00	00 1111	
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		
▲ Cap 100 mg23.87	100	Rythmodan
FLECAINIDE ACETATE		
▲ Tab 50 mg	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide
— oup ong coming too mg	00	Controlled
		Release Teva
▲ Cap long-acting 200 mg61.06	90	✓ Flecainide
Cap long-acting 200 mg01.00	30	Controlled
		Release Teva
Ini 10 ma nov ml 15 ml amnovila	-	
Inj 10 mg per ml, 15 ml ampoule100.00	5	Tambocor

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100		ANI \$29 Mexiletine Hydrochloride USP \$29
▲ Cap 250 mg	202.00	100	✓	Teva ©29 Mexiletine Hydrochloride USP ©29
				Teva S29
PROPAFENONE HYDROCHLORIDE Tab 150 mg	40.90	50	•	Rytmonorm
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – Retail pha Tab 2.5 mg Tab 5 mg	53.00	100 100	_	Gutron Gutron

Tab 5 mg SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

АТ	ENOLOL	
*	Tab 50 mg4.26	500
	Tab 100 mg7.30	500
*	Oral liq 25 mg per 5 ml21.25	300 ml OP

✓ Mylan Atenolol✓ Mylan Atenolol

✓ Atenolol AFT

✓ Atenolol AFT S29 S29

Restricted to children under 12 years of age.

BISOPROLOL FUMARATE - Brand switch fee payable (Pha	rmacode 2607034) - :	see page 23	5 for details
* Tab 2.5 mg			 Bisoprolol Mylan
* Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
* Tab 10 mg		90	✓ Bisoprolol Mylan
CARVEDILOL			
* Tab 6.25 mg	2.24	60	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	Carvedilol Sandoz
* Tab 25 mg		60	Carvedilol Sandoz

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
LABETALOL				
* Tab 100 mg	14.50	100	1	Trandate
* Tab 200 mg	27.00	100	✓	Trandate
* Inj 5 mg per ml, 20 ml ampoule	59.06	5		
	(88.60)			Trandate
* inj 5 mg per ml, 20 ml vial	42.29	1		
	(48.20)			Alvogen S29
METOPROLOL SUCCINATE	(/			- 3-
* Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
* Tab long-acting 25.75 mg		30		Betaloc CR
* Tab long-acting 95 mg		30		Betaloc CR
* Tab long-acting 190 mg		30		Betaloc CR
		00	-	Dotailo Oil
METOPROLOL TARTRATE	F 00	100		Ana Matannalal
* Tab 50 mg		100		Apo-Metoprolol
* Tab 100 mg		60		Apo-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	•	Metoprolol IV Mylan
NADOLOL				
* Tab 40 mg	16.69	100		Apo-Nadolol
* Tab 80 mg	26.43	100	•	Apo-Nadolol
PINDOLOL				
* Tab 5 mg	13.22	100	✓	Apo-Pindolol
* Tab 10 mg	23.12	100	✓	Apo-Pindolol
* Tab 15 mg	33.31	100	✓	Apo-Pindolol
PROPRANOLOL				
* Tab 10 mg	4 64	100	1	Apo-Propranolol
* Tab 40 mg		100		Apo-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 l				
Retail pharmacy		500 m	· •	Roxane-
				Propranolol \$29
				i iopianoloi

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

*	Tab 80 mg	32.58	500	✓ Mylan
*	Tab 160 mg	10.98	100	✓ Mylan

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	•	Manufacturer

TIMOLOL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking timolol prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol.

Calcium Channel Blockers

AMI ODIDINE

Dihydropyridine Calcium Channel Blockers

ΑN	ILODIPINE			
*	Tab 2.5 mg	08 90	1	Vasorex
	1.7	72 100	1	Apo-Amlodipine
	16.3	20 28	1	Bristol S29
	Vasorex to be Sole Supply on 1 June 2021			
*	Tab 5 mg0.	96 90	1	Vasorex
	1.9	56 28	1	Sandoz S29
			1	Teva S29
	3.5	33 250		Apo-Amlodipine
	Vasorex to be Sole Supply on 1 June 2021			
*	Tab 10 mg	19 90	1	Vasorex
	· ·	66 28	1	Sandoz S29
	· · · · · · · · · · · · · · · · · · ·	40 250		Apo-Amlodipine
	Vasorex to be Sole Supply on 1 June 2021			
(Aı	po-Amlodipine Tab 2.5 mg to be delisted 1 June 2021)			
٠,	po-Amlodipine Tab 5 mg to be delisted 1 June 2021)			
٠,	po-Amlodipine Tab 10 mg to be delisted 1 June 2021)			
٠,	LODIPINE			
. –	Tab long-acting 2.5 mg1.	45 30	1	Plendil ER
	Tab long-acting 5 mg			Felo 5 ER
*	Tab long-acting 10 mg4.			Felo 10 ER
		0 <u>2</u> 30		I CIO TO LIT
	FEDIPINE	00 00		Adalat 10
*	Tab long-acting 10 mg10.	63 60		Adalat 10
				Adefin S29
	18.	80 56		Tensipine MR10 S29
*	Tab long-acting 20 mg9.	12 50	✓	Mylan S29
	17.		1	Nyefax Retard
*	Tab long-acting 30 mg3.	14 30	1	Adalat Oros
	34.	10 100	1	Mylan S29
*	Tab long-acting 60 mg5.	67 30	1	Adalat Oros
	- · ·		✓	Adefin XL
	52.	81 100	1	Mylan S29
(A	dalat 10 Tab long-acting 10 mg to be delisted 1 August 2021)			•

(Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021)

(Adefin \$29) Tab long-acting 10 mg to be delisted 1 August 2021)

(Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021)

(Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021)

(Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021)

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
★ Tab 30 mg	4.60	100	✓	Dilzem
★ Tab 60 mg		100		Dilzem
Cap long-acting 120 mg	33.42	500	✓	Apo-Diltiazem CD
Cap long-acting 180 mg	50.05	500	✓	Apo-Diltiazem CD
Cap long-acting 240 mg	66.76	500	✓	Apo-Diltiazem CD
Dilzem Tab 30 mg to be delisted 1 June 2021)				
Dilzem Tab 60 mg to be delisted 1 January 2022)				
PERHEXILINE MALEATE				
₭ Tab 100 mg	62.90	100	1	Pexsig
· ·		. 55	-	<u></u>
(ERAPAMIL HYDROCHLORIDE	7.04	100	.1	loontin
≮ Tab 40 mg ≮ Tab 80 mg		100 100		Isoptin
				Isoptin
Fab long-acting 120 mg	36.02	100		Isoptin Retard \$29
h T.I.I	45.40			Isoptin SR
Fab long-acting 240 mg	15.12	30	•	Isoptin SR
★ Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a		_	_	
PSO	25.00	5	•	Isoptin
Centrally-Acting Agents				
Centrally-Acting Agents				
CLONIDINE				
Fatch 2.5 mg, 100 mcg per day - Only on a prescription	10.34	4	1	<u>Mylan</u>
Patch 5 mg, 200 mcg per day - Only on a prescription		4		Mylan
Patch 7.5 mg, 300 mcg per day - Only on a prescription		4		Mylan
CLONIDINE HYDROCHLORIDE				
	8 75	110	J	Cloniding RNM
★ Tab 25 mcg		112		Clonidine BNM Catapres
★ Tab 25 mcg★ Tab 150 mcg	34.32	100	/	Catapres
 ★ Tab 25 mcg ★ Tab 150 mcg ★ Inj 150 mcg per ml, 1 ml ampoule 	34.32	–	/	
₭ Tab 25 mcg ₭ Tab 150 mcg ₭ Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96	100	✓ ✓	Catapres Medsurge
 ★ Tab 25 mcg ★ Tab 150 mcg ★ Inj 150 mcg per ml, 1 ml ampoule 	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres Medsurge Methyldopa Mylan
₭ Tab 25 mcg ₭ Tab 150 mcg ₭ Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96	100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
₭ Tab 25 mcg ₭ Tab 150 mcg ₭ Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres Medsurge Methyldopa Mylan
K Tab 25 mcg K Tab 150 mcg K Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
₭ Tab 25 mcg ₭ Tab 150 mcg ₭ Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
K Tab 25 mcg K Tab 150 mcg K Inj 150 mcg per ml, 1 ml ampoule METHYLDOPA K Tab 250 mg	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
K Tab 25 mcg K Tab 150 mcg K Inj 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
K Tab 25 mcg K Tab 150 mcg K Inj 150 mcg per ml, 1 ml ampoule METHYLDOPA K Tab 250 mg	34.32 25.96 15.10	100 10 100	<i>y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan
* Tab 25 mcg * Tab 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10 52.85	100 10 100 500	<i>y y y y</i>	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan S29 S29
* Tab 25 mcg * Tab 150 mcg per ml, 1 ml ampoule	34.32 25.96 15.10 52.85	100 10 100	\(\frac{1}{2}\)	Catapres <u>Medsurge</u> Methyldopa Mylan Methyldopa Mylan

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	
UROSEMIDE [FRUSEMIDE]				
€ Tab 40 mg – Up to 30 tab available on a PSO	7.24	1,000	1	Apo-Furosemide
← Tab 500 mg	25.00	50	1	Urex Forte
	89.48		1	Furosemid-
				Ratiopharm S29
	169.96	100	✓	Furosemid-
				Ratiopharm S29
Oral liq 10 mg per ml	11.20	30 ml OP	1	Lasix
Inj 10 mg per ml, 25 ml ampoule	60.65	6	✓	Lasix
Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available or	a PSO1.15	5	•	Furosemide-Baxter
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	30.00	25 ml OP	1	Biomed
PLERENONE - Special Authority see SA1728 below - Ret	ail pharmacy			
Tab 50 mg		30	1	Inspra
Tab 25 mg	11.87	30	✓	Inspra
 ▶SA1728 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals ne following criteria: oth: Patient has heart failure with ejection fraction less that 2 Either: Patient is intolerant to optimal dosing of spiron. Patient has experienced a clinically significant 	n 40%; and			
IETOLAZONE			_	
Tab 5 mg	CBS	1		Metolazone S29
		50	1	Zaroxolyn S29
PIRONOLACTONE			_	
€ Tab 25 mg		100		Spiractin
F Tab 100 mg		100		Spiractin
Oral liq 5 mg per ml	30.60	25 ml OP	•	Biomed
Potassium Sparing Combination Diuretics				
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE				

AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE

28

50

✓ Frumil

✓ Moduretic

	Subsidy	:\	Fully Brand or	
	(Manufacturer's Pr	ice) S Per	Subsidised Generic Manufacture	r
Thiazide and Related Diuretics	•			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]	00.00	500		
* Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	✓ <u>Arrow-</u> Bendroflua	-ido
			Dendronda	<u>ziue</u>
May be supplied on a PSO for reasons other than em	nergency.			
* Tab 5 mg	34.55	500	✓ Arrow-	
			Bendroflua	<u>zide</u>
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	26.00	25 ml OF	✓ Biomed	
	20.00	20 1111 01	- Biomicu	
CHLORTALIDONE [CHLORTHALIDONE]	2.00	20	/ laroton coo	
Tab 25 mg	6.50	30 50	✓ Igroton S29 ✓ Hygroton	
INDAPAMIDE	0.50	50	- <u>riygroton</u>	
₩ Tab 2.5 mg	10.45	90	✓ Dapa-Tabs	
74 Tub 2.0 mg	10.40	00	• <u>Bupu 1ubo</u>	
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE				
* Tab 200 mg	19.01	90	✓ Bezalip	
* Tab long-acting 400 mg		30	✓ Bezalip Retar	d
				_
Other Lipid-Modifying Agents				
ACIPIMOX				
* Cap 250 mg	21.56	30	✓ Olbetam	
			✓ Olbetam S29	S29
Resins				
COLECTION HYDDOCHLODIDE				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	✓ Colestid	
Grano for ordering org			- Odiodilu	
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg		500	✓ Lorstat	
* Tab 20 mg		500	Lorstat	
* Tab 40 mg		500	Lorstat	
* Tab 80 mg	27.19	500	✓ <u>Lorstat</u>	
PRAVASTATIN	0.11	00	/ Du	ll.a
* Tab 20 mg* Tab 40 mg		28 28	✓ <u>Pravastatin M</u>✓ Pravastatin M	
•		20	▼ <u>FIAVASIAIII IV</u>	iyiaii
SIMVASTATIN * Tab 10 mg	1 22	90	✓ Simvastatin N	/lylan
* Tab 10 mg		90	✓ Simvastatin N	
* Tab 40 mg		90	✓ Simvastatin N	
* Tab 80 mg		90	✓ Simvastatin N	
•				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

Selective Cholesterol Absorption Inhibitors

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than $10 \times \text{normal}$) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg		30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg		30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy		Fully Brand or
	(Manufacturer's		
	\$	Per	✓ Manufacturer
Nitrates			
Milates			
GLYCERYL TRINITRATE			
* Oral pump spray, 400 mcg per dose - Up to 250 dose			
available on a PSO	6.09	250 dose OP	Nitrolingual Pump
			Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS
ISOSORBIDE MONONITRATE			
* Tab 20 mg	19.55	100	✓ <u>Ismo 20</u>
* Tab long-acting 40 mg		30	✓ Ismo 40 Retard
* Tab long-acting 60 mg	9.25	90	✓ <u>Duride</u>
O more alleged by a street			
Sympathomimetics			
ADRENALINE			
Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a F	SO4.98	5	✓ Aspen Adrenaline
, ,, ,	10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a	a PSO27.00	5	✓ Hospira
	49.00	10	Aspen Adrenaline
Vasodilators			
HYDRALAZINE HYDROCHLORIDE			
* Tab 25 mg - Special Authority see SA1321 below - Reta	iil		
pharmacy		1	✓ Hydralazine
,		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink \$29
* Inj 20 mg ampoule	25.90	5	✓ Apresoline
⇒SA1321 Special Authority for Subsidy			•
Initial application from any relevant practitioner. Approvals v	alid without furthe	r renewal unless	notified for applications meeting
the following criteria:			3
Either:			
1 For the treatment of refractory hypertension; or			
2 For the treatment of heart failure in combination with a	nitrate, in patients	who are intolera	nt or have not responded to ACE
inhibitors and/or angiotensin receptor blockers.			
MINOXIDIL			
▲ Tab 10 mg	70.00	100	✓ Loniten
NICORANDIL			
▲ Tab 10 mg		60	✓ <u>Ikorel</u>
▲ Tab 20 mg	32.28	60	✓ <u>Ikorel</u>
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42.26	50	✓ Trental 400
-			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

Endothelin Receptor Antagonists

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1991 below - Retail pharmacy

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV: and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV: or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

continued...

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail ph	armacy		
Tab 25 mg	0.64	4	✓ Vedafil
Tab 50 mg	0.64	4	✓ Vedafil
Tab 100 mg	6.60	12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dvn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retail pharmacy		
Inj 500 mcg vial36.61	1	✓ Veletri
lnj 1.5 mg vial73.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC. PO Box 10-254. WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

Differin
Differin
Oratane
Oratane
Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription13.90 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

DERMATOLOGICALS

	Subsidy	luino) O: I	Fully	Brand or
	(Manufacturer's P \$	rice) Subs Per	idised •	Generic Manufacturer
MUPIROCIN				
Oint 2%	6.60	15 g OP		
	(10.50)		В	actroban
a) Only on a prescription Not in a sorbination				
b) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.50	5 g OP	√ E	oban
a) Maximum of 5 g per prescription	1.39	3 y OF	• [<u>DDall</u>
b) Only on a prescription				
c) Not in combination				
Oint 2%	1.59	5 g OP	✓ <u>F</u>	<u>oban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER Crm 1%	10.90	50 g OP	√ =	amazine
a) Up to 250 g available on a PSO	10.60	50 y OF	• [amazme
b) Not in combination				
,				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, page	96			
AMOROLFINE				
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	14.93	5 ml OP	✓ <u>M</u>	<u>ycoNail</u>
CICLOPIROX OLAMINE				
a) Only on a prescription				
b) Not in combination Nail-soln 8%	E 70	7 ml OP	./ A	na Cialaniray
	3.72	7 ml OP	▼ <u>A</u>	po-Ciclopirox
CLOTRIMAZOLE * Crm 1%	0.77	20 g OP	√ ∩	lomazol
a) Only on a prescription		20 y Oi	• 0	ΙσπαΣσι
b) Not in combination				
* Soln 1%	4.36	20 ml OP		
	(7.55)		С	anesten
a) Only on a prescription				
b) Not in combination				
ECONAZOLE NITRATE	4.00	00 - 00		
Crm 1%		20 g OP	D	evend
a) Only on a prescription	(7.48)		Γ	evaryl
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
-	(17.23)		Р	evaryl
a) Only on a prescription				
b) Not in combination				

			DEIW	IATOLOGICALS
	Subsidy (Manufacturer's Pr \$	rice) Per	Fully Subsidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g O)P 🗸	<u>Multichem</u>
a) Only on a prescriptionb) Not in combination				
* Lotn 2%	4.36	30 ml C	OP	
	(10.03)			Daktarin
a) Only on a prescriptionb) Not in combination				
* Tinct 2%	4.36	30 ml C	OP	
	(12.10)			Daktarin
a) Only on a prescriptionb) Not in combination				
Antipruritic Preparations				
CALAMINE a) Only on a prescription				
b) Not in combination Crm, aqueous, BP	1.26	100 დ	g 🗸	healthE Calamine Aqueous Cream

CROTAMITON

- a) Only on a prescription
- b) Not in combination

....3.29 20 g OP

✓ Itch-Soothe

ΒP

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Crystals6.92	25 g
29.60	100 g

✓ MidWest

✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE		
Crm 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE		
* Crm 0.1%	50 g OP	✓ Beta Cream
* Oint 0.1%3.45	50 g OP	✓ Beta Ointment
* Lotn 0.1%	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE		
* Crm 0.05%2.18	30 g OP	✓ Dermol
* Oint 0.05%2.12	30 g OP	✓ Dermol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Pr \$	ice) Subs Per	sidised	Generic Manufacturer
I ODETAÇONE DI ITVDATE	Ψ	1 61		Manadalo
LOBETASONE BUTYRATE Crm 0.05%	E 20	30 g OP		
GIII 0.05%	(10.00)	30 g OF		Eumovate
	(10.00)			Lumovale
IFLUCORTOLONE VALERATE	0.07	50 OD		
Fatty oint 0.1%		50 g OP		dedesse
Navianna Fattu sint 0.40/ to be delicted 4. Assesset 0.004)	(15.86)		ľ	Nerisone
Nerisone Fatty oint 0.1% to be delisted 1 August 2021)				
YDROCORTISONE				
F Crm 1% – Only on a prescription	3.70	100 g OP	✓ [Hydrocortisone
				(PSM)
	17.15	500 g	✓ <u>I</u>	-tydrocortisone
				(PSM)
Powder – Only in combination		25 g		ABM
Up to 5% in a dermatological base (not proprietary Top	pical Corticosteriod	– Plain) with o	or witho	out other dermatological
galenicals				
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	y on			
a prescription	10.57	250 ml	✓ [OP Lotn HC
YDROCORTISONE BUTYRATE				
Lipocream 0.1%	6.85	100 g OP	√ I	ocoid Lipocream
Oint 0.1%		100 g OP		_ocoid
Milky emul 0.1%		100 ml OP	-	ocoid Crelo
IETHYLPREDNISOLONE ACEPONATE			_	
Crm 0.1%	4.46	15 g OP	1	Advantan
Oint 0.1%		15 g OP	_	Advantan
	4.40	13 9 01	• •	-uvantan
IOMETASONE FUROATE	4.54	45 OD	, ,	Alb-1
Crm 0.1%		15 g OP	_	Elocon Alcohol Free
O:-+ 0.40/	2.50	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
Lata 0.19/	2.90	50 g OP		Elocon
Lotn 0.1%	0.30	30 ml OP	• i	Elocon
RIAMCINOLONE ACETONIDE			_	
Crm 0.02%		100 g OP		Aristocort
Oint 0.02%	6.35	100 g OP		<u>Aristocort</u>
Corticosteroids - Combination				
Our neosterolus - Compination				
ETAMETHASONE VALERATE WITH CLIOQUINOL - Only of				
Crm 0.1% with clioquinol 3%	3.49	15 g OP		
	(4.90)		E	Betnovate-C
Betnovate-C Crm 0.1% with clioquinol 3% to be delisted 1 Jun	ne 2021)			
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [F	FUSIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
,	(10.45)	•	F	ucicort
a) Maximum of 15 g per prescription	` '			
b) Only on a prescription				
	rintian			
VDROCORTISONE WITH MICONIAZOLE - Only on a proce				
YDROCORTISONE WITH MICONAZOLE – Only on a presc Crm 1% with miconazole nitrate 2%		15 g OP	. / 1	Micreme H

	Subsidy (Manufacturer's F \$		Fully Brand or dised Generic Manufacturer
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — C Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	otion 15 g OP 15 g OP	✓ Pimafucort ✓ Pimafucort
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m and gramicidin 250 mcg per g - Only on a prescription.	g	ГIN 15 g ОР	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 10%</u>
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm	1.92	500 g	✓ Basic AquaCream ✓ Boucher ✓ Medco
(Topiderm Crm to be delisted 1 September 2021)	5.75		✓ Topiderm
CETOMACROGOL * Crm BP CETOMACROGOL WITH GLYCEROL	2.48	500 g	✓ <u>healthE</u>
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ Boucher
	3.10	1,000 ml OP	✓ Kenkay Sorbolene✓ ADE✓ Boucher
EMULSIFYING OINTMENT * Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION * Crm	2.19	500 g	✓ <u>O/W Fatty Emulsion</u> <u>Cream</u>
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50% UREA	5.35	500 ml OP	✓ <u>healthE</u>
* Crm 10%	1.37	100 g OP	✓ healthE Urea Cream

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

Subsidy (Manufacturer's F \$		Fully dised	Brand or Generic Manufacturer
5.60	1,000 ml		
(11.95)			P Lotion
1.40	250 ml OP		
(4.53)			P Lotion
5.60	1,000 ml		
(20.53)		Д	Alpha-Keri Lotion
(23.91)		В	SK Lotion
1.40	250 ml OP		
(7.73)		Е	BK Lotion
	\$	\$ Per 5.60 1,000 ml (11.95) 1.40 250 ml OP (4.53) 5.60 1,000 ml (20.53) (23.91) 1.40 250 ml OP	\$ Per

Other Dermatological Bases

PARAFFIN

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIN			

		Lotion	
IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy			
Tab 3 mg - Up to 100 tab available on a PSO 17.20	4	✓ Stromectol	

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

200 ml OP

healthE

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or

dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5% 5.75 Lotn 5% 3.99	30 g OP 30 ml OP	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN	000 OD	/ Daysaidaas
Shampoo 0.5%	200 MI OP	Parasidose

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail pharmacy			
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment;
- 2 Patient is not of child bearing potential.

RETAMETHASONE DIDRODIONATE WITH CALCIDOTRIO

DETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	52.24	60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ <u>Midwest</u>

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

				ATOLOGICALO
	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP	HUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%		75 g OP 30 g OP		gopsoryl TA gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR				
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP		Coco-Scalp Coco-Scalp
a) Maximum of 15 g per prescription b) Note: a maximum of 15 g per prescription and no more the Cream 1%	nan one prescript28.50 almologist or any alid without further	15 g OP y relevant pracer renewal unl	ctitioner ess noti	ified for applications dermatitis, rosacea,
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium		n a prescription 500 ml		Pinetarsol
SALICYLIC ACID Powder – Only in combination	18.88	250 g	✓ N ✓ P	// Aidwest PSM
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	proprietary Topic	al Corticostero	oid – Pla	ain or collodion flexible
SULPHUR				
Precipitated – Only in combination	6.35	100 g	✓ N	lidwest

1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain

2) With or without other dermatological galenicals.

Scalp Preparations		
BETAMETHASONE VALERATE * Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%3.23	100 ml OP	✓ <u>Sebizole</u>
a) Maximum of 100 ml per prescription b) Only on a prescription		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

DERMATOLOGICALS

Subsidy (Manufacturer's Price) Per

Fully Subsidised Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP

✓ Marine Blue Lotion SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 ✓ Perrigo

PODOPHYLLOTOXIN

3.5 ml OP ✓ Condyline

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

20 g OP ✓ Efudix

GENITO-URINARY SYSTEM

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	•	Manufacturer	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Contraceptives - Non-hormonal

Condoms

СО	NDOMS			
*	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
*	53 mm	0.95	10	✓ Moments
		11.64	144	✓ <u>Moments</u>
	 a) Maximum of 60 dev per prescription 			
	b) Up to 60 dev available on a PSO			_
*	53 mm, 0.05 mm thickness		10	✓ <u>Moments</u>
		11.42	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
ŧ	53 mm, chocolate, brown		10	✓ <u>Moments</u>
) II	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.05	40	
ĸ	53 mm, strawberry, red		10	✓ Moments
	a) He to 00 day and lake are a DOO	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.07	10	/ Namenta
6	56 mm	0.97	10 144	✓ Moments
	a) Marrian and CO days and agree education	11.04	144	✓ Moments
	a) Maximum of 60 dev per prescription			
E	b) Up to 60 dev available on a PSO 56 mm, 0.05 mm thickness	1 20	12	Cold Knight
~	50 mm, 0.05 mm unckness	15.57	144	✓ Gold Knight✓ Gold Knight
	a). Un to 60 day available on a DCO	13.37	144	♥ Gold Kilight
a) Up to 60 dev available on a PS0	b) Maximum of 60 dev per prescription			
F	56 mm, 0.05mm thickness (bulk pack)	1/161	144	✓ Gold Knight
•	a) Maximum of 60 dev per prescription	14.01	144	• Gold Killgill
	b) Up to 60 dev available on a PSO			
k	56 mm, 0.08 mm thickness	0.97	10	✓ Moments
•	30 mm, 0.00 mm uncaress	11.64	144	✓ Moments ✓
	a) Up to 60 dev available on a PSO	11.04	177	· momento
	b) Maximum of 60 dev per prescription			
F	56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
•	55, 5.55 mm anomicso, 100	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.01		
	b) Maximum of 60 dev per prescription			
k	56 mm, chocolate	1.30	12	✓ Gold Knight
•	, 5553440	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			***************************************
	b) Maximum of 60 dev per prescription			
*	56 mm, strawberry	1.30	12	✓ Gold Knight
	, , ,	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
K	60 mm	1.42	12	✓ Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		✓ Gold Knight XL

a) Maximumosidisedev per prescription b) substitutions a validable on a PSO

S29 Unapproved medicine supplied under Section 29

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
* 60 mm (bulk pack)	14.87	144	✓ <u>G</u>	old Knight XL

Contraceptive Devices

INTRA-UTERINE DEVICE

a) Up to 40 dev available on a PSO

b) Up to 60 dev available on a PSO

b) Only on a PSO

	b) Only on a 1 00		
*	IUD 29.1 mm length × 23.2 mm width	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	1	✓ Choice
			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to		
	84 tab available on a PSO10.00	84	✓ Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab6.62	84	
	(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

(Marvelon 28 Tab 30 mcg with desogestrel 150 mcg and 7 inert tab to be delisted 1 November 2021)

GENITO-URINARY SYSTEM

) ED
D
) ED
D
(

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO	.16.50 22.00	84 112	✓ <u>Microlut</u> ✓ <u>Microlut</u>
*	Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO	106.92	1	✓ Jadelle

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a P NORETHISTERONE		1	✓ <u>D</u>	epo-Provera	
Tab 350 mcg - Up to 84 tab available on a PSO Emergency Contraceptives	6.25	84	✓ <u>N</u>	loriday 28	
LEVONORGESTREL * Tab 1.5 mg		1 Part I c	-	ostinor-1	

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.00)	9	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	✓ Micreme
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

Myometrial and Vaginal Hormone Preparations

	,				
ER	RGOMETRINE MALEATE				
	Inj 500 mcg per ml, 1 ml ampoule - Up	to 5 inj available on a			
	PSO	160.00	5	 DBL Ergometrine 	
OE	ESTRIOL				
*	Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin	
	Pessaries 500 mcg		15	✓ Ovestin	
ОХ	(YTOCIN - Up to 5 inj available on a PS	0			
	Inj 5 iu per ml, 1 ml ampoule	3.98	5	✓ Oxytocin BNM	
	Inj 10 iu per ml, 1 ml ampoule		5	✓ Oxytocin BNM	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer	
OXYTOCIN WITH ERGOMETRINE MALEATE — Up to 5 inj avai Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	✓ <u>s</u> y	/ntometrine	

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

✓ David One Step Cassette Pregnancy Test

✓ Smith BioMed Rapid Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

★ Tab 5 mg4.81 100 ✓ Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE – Special Authority see SA1032 below – Retail pharmacy

<u>* Cap 400 mcg17.73 100 ✓ Tamsulosin-Rex</u>

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

 ★ Tab 5 mg
 11.70
 500
 ✓ Apo-Oxybutynin

 ★ Oral liq 5 mg per 5 ml
 60.40
 473 ml
 ✓ Apo-Oxybutynin

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 on the

next page − Retail pharmacy......31.80 200 ml OP ✓ Biomed

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	Ful Subsidise	•	
\$	Per •	 Manufacturer 	

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	3.00	30	✓ Solifenacin Mylan
Tab 10 mg	5.50	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE

* Compound diagnostic sticks	7.50	50 test OP	
, , , , , , , , , , , , , , , , , , ,	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
•	(13.92)		Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

✓ Mifegyne	1	Tab 200 mg60.00
✓ Mifegyne	3	180.00

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Calcium Homeostasis

CALCITONIN		
* Inj 100 iu per ml, 1 ml ampoule121.0	00 5	✓ Miacalcic
CINACALCET - Special Authority see SA1618 below - Retail pharmacy		
Tab 30 mg - Wastage claimable210.	30 28	Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g., painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial - Special Authority see SA2031 below -1 ✓ Zoledronic acid Retail pharmacy......38.03 Mylan

⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

	IAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATI Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5 5	
	(36.96)		Celestone
DE	VAMETUACONE		Chronodose
*	XAMETHASONE Tab 0.5 mg - Up to 60 tab available on a PSO	30	✓ Dexmethsone
	Tab 4 mg — Up to 30 tab available on a PSO	30	✓ Dexmethsone
	Oral liq 1 mg per ml45.00	25 ml OP	✓ Biomed
DE	XAMETHASONE PHOSPHATE		
	Dexamethasone phosphate injection will not be funded for oral use.		
*	Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
*	Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
	JDROCORTISONE ACETATE		
*	Tab 100 mcg14.32	100	✓ Florinef
	DROCORTISONE		
	Tab 5 mg	100	✓ <u>Douglas</u>
	Tab 20 mg	100	 ✓ <u>Douglas</u> ✓ Solu-Cortef
不	Inj 100 mg vial	1	▼ Solu-Cortei
	a) Up to 5 inj available on a PSO b) Only on a PSO		
ME	THYLPREDNISOLONE		
	Tab 4 mg	100	✓ Medrol
	Tab 100 mg	20	✓ Medrol
	THYLPREDNISOLONE (AS SODIUM SUCCINATE)	_0	<u></u>
IVIL	Inj 40 mg vial18.90	1	✓ Solu-Medrol-Act-
	11, 10 mg 11d	•	O-Vial
	Inj 125 mg vial	1	✓ <u>Solu-Medrol-Act-</u> <u>O-Vial</u>
	Inj 500 mg vial	1	✓ <u>Solu-Medrol-Act-O-Vial</u>
	Inj 1 g vial27.83	1	✓ Solu-Medrol
ME	THYLPREDNISOLONE ACETATE		- Join monto
IVIL	Inj 40 mg per ml, 1 ml vial44.40	5	✓ Depo-Medrol
DD		J	- Depo-Medioi
	EDNISOLONE Oral lig 5 mg per ml – Up to 30 ml available on a PSO6.00	30 ml OP	✓ Redipred
~	Restricted to children under 12 years of age.	JU IIII UF	• <u>Ileuipieu</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

^{*}Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PREDNISONE				
* Tab 1 mg	10.68	500	1	Apo-Prednisone
* Tab 2.5 mg		500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO		500	1	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	29.03	500	✓	Apo-Prednisone
FETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen \$29
			1	AU Synacthen
			1	Synacthen
★ Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Depot
			✓	Synacthene
				Retard S29
FRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
, -, -,	26.62		1	Adcortyl \$29
Inj 40 mg per ml, 1 ml ampoule	11.30	1		Triaver S29
,	51.10	5	_	Kenacort-A 40
	70.62			Kenalog S29

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CYPROTERONE ACETATE			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	85.00	1	 Depo-Testosterone
TESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	Sustanon Ampoules
TESTOSTERONE UNDECANOATE			
Cap 40 mg	21.00	60	✓ Andriol Testocaps
Inj 250 mg per ml, 4 ml vial	86.00	1	✓ Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	ice) Sub	sidised	Generic
	\$	Per	1	Manufacturer
Oestrogens				
Cestrogens				
DESTRADIOL – See prescribing guideline on the previous page				
₭ Tab 1 mg	4.12	28 OP		
	(11.10)		E	strofem
₭ Tab 2 mg		28 OP		
	(11.10)		_	strofem
Real Patch 100 mcg per 24 hours	7.91	4	✓ C	limara
 a) No more than 1 patch per week 				
b) Only on a prescription				
Patch 50 mcg per 24 hours	7.04	4	✓ C	limara
 a) No more than 1 patch per week 				
b) Only on a prescription				
Patch 25 mcg per day	6.12	8	✓ E	stradot
• • •	7.85		✓ E	stradiol TDP
				Mylan S29
a) No more than 2 patch per week				•
b) Only on a prescription				
Patch 50 mcg per day	7 04	8	√ F	stradot 50 mcg
T dion oo mog por day	9.22	O		stradiol TDP
	5.22		· _	Mylan S29
a). No many than O match man would				IVI y I a I I 329
a) No more than 2 patch per week				
b) Only on a prescription	7.04	•		
Patch 75 mcg per day		8		stradot
	10.60		▼ E	stradiol TDP
				Mylan S29
 a) No more than 2 patch per week 				
b) Only on a prescription				
Patch 100 mcg per day	7.91	8	√ E	stradot
 a) No more than 2 patch per week 				
b) Only on a prescription				
Climara Patch 100 mcg per 24 hours to be delisted 1 August 2	021)			
Climara Patch 50 mcg per 24 hours to be delisted 1 August 202	1)			
DESTRADIOL VALERATE - See prescribing guideline on the p	,			
★ Tab 1 mg	, ,	84	√ D	rogynova
⊁ Tab 2 mg		84	_	rogynova
· ·		U 1	, <u>r</u>	i ogynova
DESTROGENS – See prescribing guideline on the previous page		00		
★ Conjugated, equine tab 300 mcg		28	_	
h 0 :	(17.50)	20	Р	remarin
★ Conjugated, equine tab 625 mcg		28	_	
	(17.50)		Р	remarin
Duamantamana				
Progestogens				
MEDROXYPROGESTERONE ACETATE - See prescribing gui	deline on the prev	ious page		
★ Tab 2.5 mg		30	✓ P	rovera
* Tab 5 mg		100		rovera
⊁ Tab 10 mg		30		rovera
•			-	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		Subsidy (Manufacturer's Price	,	Fully Subsidised	Brand or Generic				
P	\$ Per ✓ Manufacturer Progestogen and Oestrogen Combined Preparations								
OE	OESTRADIOL WITH NORETHISTERONE - See prescribing guideline on page 80								
*	Tab 1 mg with 0.5 mg norethisterone acetate	, ,	28 OF		Kliovance				
*	Tab 2 mg with 1 mg norethisterone acetate	` '	28 OF	•	Kliogest				
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	,			Kilogest				
	oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40 (18.10)	28 OF		Trisequens				
0	ther Oestrogen Preparations								
ETI *	HINYLOESTRADIOL Tab 10 mcg	17.60	100	1	NZ Medical and Scientific				
-	STRIOL Tab 2 mg	7.00	30	•	Ovestin				
0	ther Progestogen Preparations								
* *	/ONORGESTREL Intra-uterine device 52 mg Intra-uterine device 13.5 mg		1		<u>Mirena</u> Jaydess				
	DROXYPROGESTERONE ACETATE Tab 100 mg RETHISTERONE	116.15	100	✓	Provera HD				
*	Tab 5 mg - Up to 30 tab available on a PSO OGESTERONE	5.49	30	✓	Primolut N				
	Cap 100 mg - Special Authority see SA1609 below - Retail			_					

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

pharmacy......16.50

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

30

✓ Utrogestan

_		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
T	hyroid and Antithyroid Agents				
CA *	RBIMAZOLE Tab 5 mg	10.80	100		Neo-Mercazole Neo-Mercazole S29 S29
LE'	VOTHYROXINE				
*	Tab 25 mcg	5.55	90	1	Synthroid
*	Tab 50 mcg	1.71	28	✓	Mercury Pharma
		5.79	90	✓	Synthroid
		64.28	1,000	/	Eltroxin
*	Tab 100 mcg	1.78	28	✓	Mercury Pharma
		6.01	90	✓	Synthroid
		66.78	1,000	/	Eltroxin
PR	OPYLTHIOURACIL – Special Authority see SA1199 below – Propylthiouracil is not recommended for patients under the a treatments are contraindicated.		the	patient is pı	regnant and other
	Tab 50 mg	35.00	100	1	PTU \$29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SC	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	v – Retail p	harmacy	
*	Inj 5 mg cartridge	34.88	1	 Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	.104.63	1	✓ Omnitrope

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

(Manufacturer's Price) Subsidised Generic	
\$ Per 🗸 Manufacturer	

continued...

- children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

Subsidy (Manufacturer's Po	rice) Per	Fully Subsidised	Brand or Generic Manufacturer	
_				

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price	,	Subsidised	Generic	
\$	Per		Manutacturer	

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and

(Manufacturer's Price)	Subsidised	Generic
\$ Pe	er 🗸	Manufacturer

continued...

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH	Ana	logues
------	-----	--------

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe		1	✓ Teva
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a chil	ld or adolescent a	and is unable	e to tolerate ad

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

(591.68)

Vasopressin Agonists

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

Lucrin Depot 3-month

	Subsidy (Manufacturer's		Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	linirin
Tab 200 mcg	54.45	30	✓ N	linirin
▲ Nasal drops 100 mcg per ml	39.03	2.5 ml OP	✓ N	linirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	√ <u>D</u>	esmopressin- PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	✓ N	linirin

Other Endocrine Agents

CABERGOI INF

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA1370 below3.75
✓ Dostinex	8	15.20

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

\cap	\cap	/ 11□			\cap	TD/	ΔTF	
l /	()I\	/!![-1	15	l d	ı m	4 I F	

Tab 50 mg	29.84	10	✓ Mylan
,			Clomiphen S29
METYRAPONE			
Cap 250 mg	558.00	50	✓ Metopirone

INFECTIONS - AGENTS FOR SYSTEMIC USE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 Fskazole S29 ⇒SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription 6 ✓ Vermox 15 ml (7.53)Vermox PRAZIQUANTFI 8 Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 61 b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 229 Cephalosporins and Cephamycins CEFACLOR MONOHYDRATE Cap 250 mg......24.70 100 ✓ Ranbaxy-Cefaclor Grans for oral lig 125 mg per 5 ml - Wastage claimable......3.53 100 ml Ranbaxy-Cefaclor **CEFALEXIN** 20 Cephalexin ABM ✓ Cephalexin ABM 20 Grans for oral lig 25 mg per ml - Wastage claimable.....8.75 100 ml ✓ Cefalexin Sandoz Grans for oral lig 50 mg per ml - Wastage claimable......11.75 100 ml ✓ Cefalexin Sandoz CEFAZOLIN - Subsidy by endorsement Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly. ✓ AFT ✓ AFT CEFTRIAXONE - Subsidy by endorsement a) Up to 10 inj available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

CEFUROXIME AXETIL - Subsidy by endorsement

✓ Ceftriaxone-AFT

✓ Ceftriaxone-AFT

Zinnat

5

Subsidy	Fı	ılly	Brand or	۰
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	.8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable1	14.38	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

1 Atypical mycobacterial infection; or

ERYTHROMYCIN (AS LACTORIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO			•
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
 a) Up to 200 ml available on a PSO 			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg	29.90	100	
	(44.58)		ERA
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-
			<u>Roxithromycin</u>
Tab 300 mg	16 33	50	✓ Arrow-
Tab 500 Hig	10.33	50	Roxithromycin
			1 to Aittii Oili yolii

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		idised	
	\$	Per	✓	Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500	1	<u>Alphamox</u>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500	1	<u>Alphamox</u>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	•	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable	4.70	4001	,	Al-, b 050
Grans for oral liq 250 mg per 5 ml	1./3	100 ml	•	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable Inj 250 mg vial	15.07	10	1	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial — Up to 5 inj available on a PSO		10		Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID		. •		
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab available on a PSO	0.90	10	1	Curam Duo 500/125
available on a FSO	5.00	20		Augmentin
Curam Duo 500/125 to be Sole Supply on 1 July 2021	3.00	20	•	Augmentin
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25	ma			
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5	ma			
per ml – Up to 200 ml available on a PSO	•	100 ml OP	1	Curam
(Augmentin Tab 500 mg with clavulanic acid 125 mg to be delisted				
BENZATHINE BENZYLPENICILLIN	, ,			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	344.93	10	1	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10	1	Sandoz
, , , , , ,	00 11.03	10	•	<u>Januoz</u>
FLUCLOXACILLIN Cap 250 mg – Up to 30 cap available on a PSO	16 00	250	./	Staphlex
Cap 500 mg - Up to 30 cap available on a PSO		500		Staphlex
Grans for oral liq 25 mg per ml		100 ml		AFT
a) Up to 200 ml available on a PSO	2.23	100 1111	٠	<u>AL 1</u>
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	1	AFT
a) Up to 200 ml available on a PSO			-	<u>-</u>
b) Wastage claimable				
Inj 250 mg vial	17.56	10	1	Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial - Up to 5 inj available on a PSO		5	✓	Flucil

	Subsidy		Fully	
	(Manufacturer's Price)			
	\$	Per		Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	2.59	50		Cilicaine VK
Cap 500 mg	4.26	50	/	Cilicaine VK
 a) Up to 20 cap available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	1	<u>AFT</u>
 a) Up to 200 ml available on a PSO 				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	/	<u>AFT</u>
 a) Up to 300 ml available on a PSO 				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	/	Cilicaine
Tetracyclines				
DOVVOVO INF				
DOXYCYCLINE * Tab 100 mg - Up to 30 tab available on a PSO	64.49	500	/	Doxine
	04.43	300	•	DOXIIIE
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg – Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy		60		Martin
W Con 100 mm	(12.05)	100		Mino-tabs
* Cap 100 mg		100		Minancia
CA40FF Consist Authority for Manufacturers Bries	(52.04)			Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 $\label{temperature} \textbf{TETRACYCLINE} \ - \ \textbf{Special Authority see} \ \frac{\textbf{SA1332 below}}{\textbf{SA1332 below}} - \ \textbf{Retail pharmacy}$

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully	
	(Manufacturer's Price)	D	Subsidised	
	\$	Per		Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg		24	✓	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	•	Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and th	e prescription is endo	sed	accordingly	у.
Inj 150 mg	65.00	1	1	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement.	25.00	5	✓	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement.	182.00	10	1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection	and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	17.50	10	✓	Pfizer
	87.50	50	1	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection a	and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Reta No patient co-payment payable	l pharmacy			
Tab 400 mg	42.00	5	1	Avelox
⇒SA1740 Special Authority for Subsidy				

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications:
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

	Subsidy (Manufacturer's Price) \$	Per	•	Brand or Generic Manufacturer
PAROMOMYCIN - Special Authority see SA1689 below - Retai	*			- Indianalara
Cap 250 mg		16	√ Hu	matin \$29
⇒SA1689 Special Authority for Subsidy				
Initial application only from an infectious disease specialist, clin month for applications meeting the following criteria: Either:	ical microbiologist or	gastro	penterologist.	Approvals valid for 1
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
Renewal only from an infectious disease specialist, clinical micro applications meeting the following criteria: Either:	biologist or gastroent	teroloç	gist. Approva	ls valid for 1 month for
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE – Special Authority see SA1328 below – Retain Tab 25 mg		30	✔ Da	raprim S29
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months	r a period of 3 months		nless notified	for applications meetin
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg	34.50	12	√ Fu	cidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below				
Tab 500 mg		56	✓ Wo	ockhardt \$29
⇒SA1331 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:	d without further rene	wal ur	nless notified	for applications meetin
For the treatment of toxoplasmosis in patients with HIV for pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months	•	s; or		
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient an Solution for inhalation 60 mg per ml, 5 ml – Subsidy by		5 endors		<mark>bramycin Mylan</mark> ly.
endorsementa) Wastage claimable	395.00 5	66 dos	e ✓ <u>To</u> l	bramycin BNM
b) Only if prescribed for a cystic fibrosis patient and the	prescription is endor	sed a	ccordingly.	
TRIMETHOPRIM			4	_
* Tab 300 mg - Up to 30 tab available on a PSO		50	✓ <u>TM</u>	<u>P</u>
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX				
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - U to 30 tab available on a PSO		500	✓ Tri	sul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200 available on a PSO		100 m	✓ De	prim

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

VANCOMYCIN - Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 62
- b) For topical antifungals refer to GENITO URINARY, page 75

FLUCONAZOLE

Cap 50 mg2.75	28	Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg12.89	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy	35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 4 Detient as action
 - 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
 - 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOI F

TO COTT LEGEL		
Cap 100 mg4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e)	Subsidised	Generic
	\$	Per	•	Manufacturer
KETOCONAZOLE				
Tab 200 mg - PCT	CBS	30	✓	Link Healthcare S29
· ·			/	Nizoral S29
		100	✓	Strides Shasun S29
NYSTATIN				
Tab 500,000 u	14.16	50		
	(17.09)			Nilstat
Cap 500,000 u	12.81 [′]	50		
•	(15.47)			Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Ret	ail pharmacy			
Tab modified-release 100 mg	869.86	24	✓	Noxafil
Oral liq 40 mg per ml	761.13	105 ml	OP 🗸	Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	8.15	84	Deolate
Deolate to be Sole Supply on 1 August 2021			
VORICONAZOLE - Special Authority see SA1273 below -	Retail pharmacy		
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,437.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 Patient has proven or probable invasive aspergillus infection; or
- 3.2 Patient has possible invasive aspergillus infection; or
- 3.3 Patient has fluconazole resistant candidiasis: or
- 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Reta	il pharmacy		
Tab 7.5 mg	117.00	56	✓ Primacin S29
Tab 15 mg	400.00	100	✓ Sanofi
			Primaquine S29

(Primacin \$29) Tab 7.5 mg to be delisted 1 June 2021)

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antiparasitics

Antiprotozoals

QUININE SULPHATE		
* Tab 300 mg61.91	500	✓ Q 300
(Q 300 Tab 300 mg to be delisted 1 July 2021)		

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl

		INFECTIONS - AC			
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
	AZOLE				
Ta	b 500 mg	32.95	10	/	Arrow-Ornidazole
Anti	tuberculotics and Antileprotics				
lote:	There is no co-payment charge for all pharmaceuticals li	sted in the Antitubercul	otics	and Antile	protics group regardless
	ation status.				F
LOF	ZIMINE - Retail pharmacy-Specialist				
	No patient co-payment payable				
b)	Prescriptions must be written by, or on the recommendate dermatologist.		iseas	se physicia	n, clinical microbiologist o
€ Ca	p 50 mg	442.00	100	1	Lamprene S29
YCLO	SERINE - Retail pharmacy-Specialist				
	No patient co-payment payable				
b)	Prescriptions must be written by, or on the recommendar respiratory physician.		iseas	se physicia	n, clinical microbiologist c
Ca	p 250 mg	344.00	60	1	Cyclorin S29
APS	DNE - Retail pharmacy-Specialist				
	No patient co-payment payable				
b)	Prescriptions must be written by, or on the recommenda	ation of, an infectious d	iseas	se physicia	n, clinical microbiologist o
	dermatologist				
	b 25 mg		100		Dapsone
	b 100 mg		100	/	Dapsone
THAN	IBUTOL HYDROCHLORIDE - Retail pharmacy-Special	list			
	No patient co-payment payable				
	Prescriptions must be written by, or on the recommendarespiratory physician				-
Ta	b 100 mg	85.73	100	/	EMB Fatol S29
Ta	b 400 mg	49.34	56	/	Myambutol S29
ONIA	AZID - Retail pharmacy-Specialist				
	No patient co-payment payable				
b)	Prescriptions must be written by, or on the recommendation	ation of, an internal med	dicine	e physician	, paediatrician, clinical
_	microbiologist, dermatologist or public health physician				
· Ia	b 100 mg	22.00	100	•	<u>PSM</u>
	AZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
	No patient co-payment payable				
b)	Prescriptions must be written by, or on the recommendation	ation of, an internal med	dicine	e physician	, paediatrician, clinical
т.	microbiologist, dermatologist or public health physician	05.54	400		Differen
	b 100 mg with rifampicin 150 mgb 150 mg with rifampicin 300 mg		100 100	_	Rifinah Difinah
			100	•	Rifinah
	AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
	No patient co-payment payable		:	! . ! ! .	4 - aliusia al musia malai al a mia 4 -
D)	Prescriptions must be written by, or on the recommendarespiratory physician	allon of, an infectious d	iseas	se specialis	a, cimical microbiologist c
Gr	ans for oral liq 4 g sachet	280.00	30	1	Paser S29
		200.00	50	•	. 4301
	ONAMIDE – Retail pharmacy-Specialist				
	No patient co-payment payable Prescriptions must be written by, or on the recommenda	ation of an infactious d	icon	e enocialia	t clinical microbiologist s
h١		anon or, an intectious u	iseas	oc specialis	i, cimicai microbiologist C
b)	resniratory physician				
·	respiratory physician b 250 mg	305.00	100	/	Peteha \$29

		Subsidy (Manufacturer's Price) Subsi	Fully	Brand or Generic
		(Manufacturer's Frice)	Per	uiseu ✓	Manufacturer
PY	RAZINAMIDE - Retail pharmacy-Specialist				
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation	on of, an infectious	disease phy	sician,	clinical microbiologist or
*	respiratory physician Tab 500 mg	59 00	100	✓ A	FT-Pyrazinamide
	ABUTIN – Retail pharmacy-Specialist		100		ar i i yrazmamac
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation	on of, an infectious	disease phy	sician,	respiratory physician or
	gastroenterologist				
	Cap 150 mg	299.75	30	✓ N	lycobutin
RIF	AMPICIN – Subsidy by endorsement				
	a) No patient co-payment payableb) For confirmed recurrent Staphylococcus aureus infection	in combination with	other effecti	ve anti	i-etanhylococcal
	antimicrobial based on susceptibilities and the prescription				
	Retail pharmacy - Specialist. Specialist must be an intern				
.1.	paediatrician, or public health physician.	50.54	400		N
*	Cap 150 mg		100 100	_	<u>Rifadin</u> Rifadin
	Oral liq 100 mg per 5 ml		60 ml	_	Rifadin
				_	
Α	ntivirals				
For	eye preparations refer to Eye Preparations, Anti-Infective Pre	parations, page 229)		
	epatitis B Treatment	paranono, pago 220			
	TECAVIR Tab 0.5 mg	52.00	30	/ =	ntecavir Sandoz
	MIVUDINE – Special Authority see SA1685 below – Retail pha		30	• -	intecavii Sandoz
LAI	Tab 100 mg		28	✓ Z	etlam
	Oral liq 5 mg per ml		40 ml OP	_	effix
	SA1685 Special Authority for Subsidy				
	ial application only from a relevant specialist or medical pract		nmendation	of a re	levant specialist.
	provals valid for 1 year where used for the treatment or preven newal from any relevant practitioner. Approvals valid for 2 year	•	ha traatman	t or pro	avention of henatitic B
	NOFOVIR DISOPROXIL	ars wriere asea for the	ne neannen	t or pre	evention of nepatitis b.
	Tenofovir disoproxil prescribed under endorsement for the tre	eatment of HIV is inc	cluded in the	count	of up to 4 subsidised
	antiretrovirals for the purposes of Special Authority SA1651.,				·
*	Tab 245 mg (300.6 mg as a succinate)	38.10	30	✓ <u>T</u>	enofovir Disoproxil
					<u>Teva</u>
Н	erpesvirus Treatments				
	CLOVIR				
*	Tab dispersible 200 mg	1.60	25	✓ L	ovir
*	Tab dispersible 400 mg	5.38	56	✓ L	.ovir
*	Tab dispersible 800 mg	5.98	35	✓ L	<u>.ovir</u>
VA	_ACICLOVIR		00	٠. م	ta al andre
	Tab 1,000 mg		30 30		<u>′aclovir</u> ′aclovir
	TWO 1,000 Hig		00	- <u>v</u>	40.0111

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
VALGANCICLOVIR – Special Authority see SA1993 below – Tab 450 mg	'	60	_	alganciclovir Mvlan	

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Roth:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

⇒SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and

Subsidy		Fully	Brand or
(Manufacturer's Price) Subs		sidised	Generic
\$	Per	✓	

continued...

- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Subsi	idy F	ully	Brand or
(Manufacture	er's Price) Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

EFAVIRENZ - Special Authority see SA1651 on the previous	s page – Retail phari	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	ıs page – Retail pha	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	us page – Retail pha	rmacy	
Tab 200 mg	60.00	60	✓ Nevirapine Alphapharm
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune Suspension

ABACAVIR SULPHATE - Special Authority see SA1651 on the previous page - Retail pharmacy

Nucleosides Reverse Transcriptase Inhibitors

Tab 300 mg Oral liq 20 mg per ml		60 240 ml OP	✓ <u>Ziagen</u>✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Author Note: abacavir with lamivudine (combination tablets) coun anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISO Retail pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil		•	
anti-retroviral Special Authority	counts as timee a	iiu-ieuoviiai iiie	dications for the purposes of the
Tab 600 mg with emtricitabine 200 mg and tenofovir disop 245 mg (300 mg as a maleate)		30	✓ <u>Mylan</u>
EMTRICITABINE – Special Authority see SA1651 on the prev Cap 200 mg		pharmacy 30	✓ Emtriva
LAMIVUDINE - Special Authority see SA1651 on the previous	page – Retail pha	armacy	
Tab 150 mg	84.50	60	✓ <u>Lamivudine</u> Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] - Special Authority see SA1651 on the pr			
Cap 100 mg Oral liq 10 mg per ml		100 200 ml OP	✓ Retrovir✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority s Note: zidovudine [AZT] with lamivudine (combination table the anti-retroviral Special Authority.			
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ Alphapharm

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg	141.68	rmac 60 60	/	<u>Teva</u> Teva
DARUNAVIR – Special Authority see SA1651 on page 104 – Re Brand switch fee payable (Pharmacode 2607026) - see page Tab 400 mg Tab 600 mg	235 for details 132.00	60 60		Darunavir Mylan Darunavir Mylan
LOPINAVIR WITH RITONAVIR — Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg Tab 200 mg with ritonavir 50 mg Oral liq 80 mg with ritonavir 20 mg per ml	183.75 463.00	l phar 60 120 0 ml	/	Kaletra Kaletra Kaletra
RITONAVIR – Special Authority see SA1651 on page 104 – Reta Tab 100 mg		30	✓	<u>Norvir</u>
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 Tab 50 mg	1,090.00	30		Tivicay
Tab 400 mg	1,090.00	60 60	′ 🗸	Isentress Isentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test: or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

✓ Pegasys

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

00.00 4

SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed: or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

Subsidy	Fully Subsidised		Brand or	
(Manufacturer's Price)			Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Wastage claimable	86.40	100	✓ Macrobid
Macrobid to be Sole Supply on 1 August 2021			
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	Arrow-Norfloxacin
Only if prescribed for a nationt with an uncomplicated u		n that is unr	cononcius to a first line agent

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	19.60	10		Juno S29
	29.40		✓	Max Health
	98.00	50	✓	AstraZeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45 79	100	1	Mestinon
_ 145 00 mg		100		MCGUITOII
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg		50		Diclofenac Sandoz
* Tab 50 mg dispersible	1.50	20	1	Voltaren D
* Tab EC 50 mg	1.23	50	✓	Diclofenac Sandoz
* Tab long-acting 75 mg	22.80	500	✓	Apo-Diclo SR
* Tab long-acting 100 mg	25.15	500	1	Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a l		5		Voltaren
* Suppos 12.5 mg		10		Voltaren
* Suppos 25 mg		10		Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
* Suppos 100 mg		10		Voltaren
11	7.00	10	•	Voltaren
IBUPROFEN				
* Tab 200 mg		1,000		Relieve
* Tab long-acting 800 mg		30		Ibuprofen SR BNM
* Oral liq 20 mg per ml	1.88 2	200 m	· •	<u>Ethics</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	1	Oruvail SR
MEFENAMIC ACID				
* Cap 250 mg	1.05	EΟ		
* Cap 250 mg		50		Danatan
	(9.16)	00		Ponstan
	0.50	20		
	(5.60)			Ponstan
NAPROXEN				
* Tab 250 mg	32.69	500	1	Noflam 250
* Tab 500 mg	22.19	250	1	Noflam 500
* Tab long-acting 750 mg		28	1	Naprosyn SR 750
* Tab long-acting 1 g		28		Naprosyn SR 1000
SULINDAC				
	0.57	- -0		Mulan 200
* Tab 100 mg		56		Mylan S29
* Tab 200 mg		50		Aclin
	16.91	56	•	Sulindac Mylan S29
TENOXICAM				
* Tab 20 mg	9.15	100	1	Tilcotil
* Inj 20 mg vial		1		AFT
		•	•	•

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	I Generic
NSAIDs Other				
ELECOXIB				
Cap 100 mg	5.80	60		Celecoxib Pfizer
Cap 200 mg	2.30	30		Celebrex
	3.30		✓	Celecoxib Pfizer
Topical Products for Joint and Muscular Pain				
CAPSAICIN				
Crm 0.025% - Special Authority see SA1289 below - Retail				
pharmacy	9.75	45 g C)P 🗸	Zostrix
Antirheumatoid Agents				
IYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systemic suppression, relevant dermatological conditions (cutaneous for mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmor Pharmacists may annotate the prescription as endorsed wher hydroxychloroquine. Note: Indication marked with a * is an u * Tab 200 mg	orms of lupus and li nary)*, and the pres te there exists a rec anapproved indicati	chen p cription cord of	olanus, cut n is endor prior disp	aneous vasculitides and sed accordingly.
EFLUNOMIDE				<u> </u>
Tab 10 mg	6.00	30	1	Arava
Tab 20 mg		30		Arava
PENICILLAMINE				
Tab 125 mg	67.23	100	/	D-Penamine
Tab 250 mg		100		D-Penamine
- a.5 _ 5 0 mg				
Drugs Affecting Bone Metabolism				
Alendronate for Osteoporosis				

ALENDRONATE SODIUM

* Tab 70 mg	.2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5 600 iu	1.51	4	✓ Fosamax Pli

Other Treatments

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
	Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
	Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RA	LOXIFENE HYDROCHLORIDE - Special Authority see SA1779 or	the next page	- Retail pha	armacy
*	Tab 60 mg	53.76	28	✓ Evista

Subsidy (Manufacturer's Price) \$ Pe

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily;

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per •	Manufacturer

continued...

zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

(Ma	Subsidy anufacturer's Price)	Subsidi Per	ully	Brand or Generic Manufacturer
	<u></u>	Per		Manufacturer

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene: and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA19	63 below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	 Benzbromaron AL
			100 829

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	9.58	100	Colgout
FEBUXOSTAT - Special Authority see SA1996 below - Reta	ail pharmacy		
Tab 80 mg	39.50	28	Adenuric
Tab 120 mg	39.50	28	✓ Adenuric

⇒SA1996 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout: and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

\$ Per ✔ Manufacturer	Subsidy (Manufacturer's Price)	Fully Subsidised	
	\$	Per 🗸	Manufacturer

continued...

2 Patient has a documented history of allopurinol intolerance.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

Muscle Relaxants

BACLOFE	

*	Tab 10 mg4.20	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients where oral ar	ntispastic ag	ents have been ineffective or have
	caused intolerable side effects and the prescription is endorsed according	ly.	

Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement............372.98 5 Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

DANTDOLENE

Cap 25 mg97.50	100	✓ Dantrium
Cap 50 mg77.00	100	✓ Dantrium S29 S29✓ Dantrium
ORPHENADRINE CITRATE		
Tab 100 mg18.54	100	✓ Norflex

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE ▲ Cap 100 mg38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	5	✓ Movapo

BROMOCRIPTINE MESYLATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of bromocriptine mesylate.

	prior dispensing of bromocriptine mesylate.			
*	Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ΕN	TACAPONE			
\blacktriangle	Tab 200 mg	22.00	100	✓ Entapone
LE	VODOPA WITH BENSERAZIDE			
*	Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
*	Cap 50 mg with benserazide 12.5 mg	13.75	100	✓ Madopar 62.5
*	Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
*	Cap long-acting 100 mg with benserazide 25 mg	22.85	100	✓ Madopar HBS
*	Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
LE	VODOPA WITH CARBIDOPA			
*	Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
*	Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
	Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
	AMIPEXOLE HYDROCHLORIDE			
\blacktriangle	Tab 0.25 mg	6.12	100	✓ Ramipex
\blacktriangle	Tab 1 mg		100	✓ Ramipex
RΩ	PINIROLE HYDROCHLORIDE			
A	Tab 0.25 mg	2.85	84	✓ Ropin
	•	3.39	100	✓ Mylan S29
\blacktriangle	Tab 1 mg	3.95	84	✓ Ropin
	v	4.70	100	✓ Mylan \$29
\blacktriangle	Tab 2 mg	5.48	84	✓ Ropin
\blacktriangle	Tab 5 mg		84	✓ Ropin
SF	LEGILINE HYDROCHLORIDE			 _
	Tab 5 mg	22.00	100	✓ Apo-Selegiline S29 S29
ТО	LCAPONE			
A	Tab 100 mg	152.38	100	✓ Tasmar

Anticholinergics

RENZ∆TROPINE ME	CVIATE

Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Un to 10 ini queilable on a DCO			

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	•	Kemadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	-	56	√	Rilutek
►SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:				
1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.				e initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow.	onths for applications	s mee	ting the fo	llowing criteria:
TETRABENAZINE Tab 25 mg	91.10	112	✓	<u>Motetis</u>
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement	dministration and the	30 ml prese 10	cription is	Xylocaine 2% Jelly endorsed accordingly. Instillagel Lido

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

a) Up to 5 each available on a PSO

accordingly.

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 m	ı 🗸	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	✓	Lidocaine-Claris
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓	Lidocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓	Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓	Lidocaine-Claris
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	103.32	10	1	Pfizer
a) Up to 5 each available on a PSO		. •		
b) On heiding a heid and a firm and the standard and a second and a				

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see	SA0906 above – Retail pharn	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - S	Special Authority see SA0906	above – Reta	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

VCDIDINI

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

ASPININ			
* Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabe	etic periphera	ıl neuropathy a	nd the prescription is endorsed
accordingly.			
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	Rugby Capsaicin
			Topical
			Cream S29
NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
	\$	Per	/	Manufacturer
PARACETAMOL				
Tab 500 mg - blister pack	0.50	20	✓ I	Medco
			-	Paracare
				Pharmacy Health
	1.12		✓ [Ethics Paracetamol Classic
	2.48	100	✓ F	Paracare
			✓ F	Pharmacy Health
	11.75	96	✓ [Panadol Mini Caps
	24.82	1,000	√ [Paracetamol Pharmacare
			✓ F	Pharmacare

- b) Up to 30 tab available on a PSO

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

Tab 500 mg - bottle pack - Maximum of 300 tab per prescription; can be waived by endorsement24.82 1.000 ✓ Paracetamol **Pharmacare**

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

*	a) Up to 200 ml available on a PSO	5.45	1,000 ml	✓ Paracare
	b) Not in combination			
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ <u>Paracare Double</u> <u>Strength</u>
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3.29	10	✓ Gacet
*	Suppos 250 mg	3.79	10	✓ Gacet
*	Suppos 500 mg	12.40	50	✓ Gacet
(Pa	aracare Tab 500 mg - blister pack to be delisted 1 June 2021)			
(Pa	aracare Tab 500 mg - blister pack to be delisted 1 June 2021)			

Opioid Analgesics

CODEINE PHOSPHATE - Safety medicine; prescriber may dete	rmine dispensing	frequency	
Tab 15 mg		100	✓ PSM
Tab 30 mg	7.45	100	✓ PSM
Tab 60 mg	14.25	100	✓ PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ DHC Continus

121

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	equency			
Inj 50 mcg per ml, 2 ml ampoule		10		Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
Patch 12.5 mcg per hour	2.95	5	✓	Fentanyl Sandoz
Patch 25 mcg per hour	3.66	5	✓	Fentanyl Sandoz
Patch 50 mcg per hour	6.65	5		Fentanyl Sandoz
Patch 75 mcg per hour	9.25	5	✓	Fentanyl Sandoz
Patch 100 mcg per hour	11.40	5	✓	Fentanyl Sandoz
 c) Safety medicine; prescriber may determine dispensing fr d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablets). e) For methadone hydrochloride oral liquid refer Standard F 	reimbursed at the rate	e of th	ne cheapes	st form available
Tab 5 mg		10	1	Methatabs
Oral lig 2 mg per ml		200 n		Biodone
Oral lig 5 mg per ml		200 n		Biodone Forte
Oral liq 10 mg per ml		200 n		Biodone Extra Forte
Inj 10 mg per ml, 1 ml		10		AFT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug formb) No patient co-payment payablec) Safety medicine; prescriber may determine dispensing fr				
Oral liq 1 mg per ml		200 n		RA-Morph
Oral liq 2 mg per ml		200 n		RA-Morph
Oral liq 5 mg per ml	19.44	200 n	nl 🗸	Ordine S29

Oral liq 10 mg per ml27.74

✓ RA-Morph

✓ Ordine S29

✓ RA-Morph

200 ml

(Manufacturer's Price) Subsidised Ger	and or
	nufacturer
· · · · · · · · · · · · · · · · · · ·	- Iuidotaroi
MORPHINE SULPHATE	
a) Only on a controlled drug form	
b) No patient co-payment payable	
c) Safety medicine; prescriber may determine dispensing frequency	امام
Tab immediate-release 10 mg	
Tab immediate-release 20 mg	
	/-Morphine LA
Cap long-acting 30 mg	
Cap long-acting 100 mg	
	Morphine
, 01	phate
· · · · · · · · · · · · · · · · · · ·	
, 01	Morphine
·	phate
Inj 15 mg per ml, 1 ml ampoule − Up to 5 inj available on a PSO7.08 5 ✓ DBL N	•
· · · · · · · · · · · · · · · · · · ·	phate
Inj 30 mg per ml, 1 ml ampoule − Up to 5 inj available on a PSO7.28 5 ✓ DBL N	•
Sulp	phate
(Arrow-Morphine LA Tab long-acting 30 mg to be delisted 1 June 2021)	
OXYCODONE HYDROCHLORIDE	
a) Only on a controlled drug form	
b) No patient co-payment payable	
c) Safety medicine; prescriber may determine dispensing frequency	
	odone Sandoz
	odone Sandoz
\$29	
Tab controlled-release 10 mg2.15 20 ✓ Oxycc	odone Sandoz
	odone Sandoz
S29	
	odone Sandoz
· · · · · · · · · · · · · · · · · · ·	
\$29	
•	odone Sandoz
S29	
	odone Sandoz
5.38 50 C Oxyco	odone Sandoz
S29	\$29
10.75 100 ✓ Oxycc	odone Sandoz
S29	\$29
Tab controlled-release 40 mg	odone Sandoz
•	odone Sandoz
Cap immediate-release 5 mg	
Cap immediate-release 10 mg	
Cap immediate-release 20 mg	
Oral lig 5 mg per 5 ml	
Inj 10 mg per ml, 1 ml ampoule	
Inj 10 mg per ml, 2 ml ampoule	
Inj 50 mg per ml, 1 ml ampoule	

	Subsidy	Ful	
((Manufacturer's Price) \$	Subsidise Per •	d Generic Manufacturer
PARACETAMOL WITH CODEINE – Safety medicine; prescriber r			
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000	⊂y [′] Paracetamol +
Tab paracetamer doe mg war doddine phospitate o mg	20.01	1,000	Codeine (Relieve)
PETHIDINE HYDROCHLORIDE			(
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing free	quency		
Tab 50 mg			' <u>PSM</u>
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO 29.88	5	DBL Pethidine
		_	Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a PS	SO 30.72	5	DBL Pethidine
			Hydrochloride
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg			Tramal SR 100
Tab sustained-release 150 mg			Tramal SR 150
Tab sustained-release 200 mg Cap 50 mg			
Cap 50 mg	2.00	100	ATTOW-TTAINIAUOI
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE - Safety medicine; prescriber may determine dis	enoncina froguency		
Tab 10 mg		100	Arrow-Amitriptyline
Tab 25 mg			Arrow-Amitriptyline
Tab 50 mg			Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescrib	er may determine d	lispensina frea	uency
Tab 10 mg			Apo-Clomipramine
Tab 25 mg			Apo-Clomipramine
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by end			
Safety medicine; prescriber may determine dispensing frec			
b) Subsidy by endorsement – Subsidised for patients who we		[dothiepin] hyd	Irochloride prior to 1 June
2019 and the prescription is endorsed accordingly. Pharm			
exists a record of prior dispensing of dosulepin [dothiepin]	hydrochloride.		_
Tab 75 mg			Dosulepin Mylan
Cap 25 mg	7.83	50	Dosulepin
			Mylan S29
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber n		ensing frequen	СУ
Tab 10 mg			Tofranil
T-1: 05	10.96		Tofranil
Tab 25 mg	8.80	50	Tofranil
MAPROTILINE HYDROCHLORIDE – Subsidy by endorsement			
Safety medicine; prescriber may determine dispensing fred Subsidiary by and arranged to the dispension of the profit of the dispension of the		المارة المارية المارية	andanta d Cantandra
b) Subsidy by endorsement – Subsidised for patients who we			
2020 and the prescription is endorsed accordingly. Pharm		e ine prescripti	on as endorsed where there
exists a record of prior dispensing of maprotiline hydrochlo Tab 75 mg		20	′ Ludiomil
rab ro my	21.01		Ludiomil
(Ludiomil Tab 75 mg to be delisted 1 August 2021)			
(Ludiomil Tab 75 mg to be delisted 1 August 2021)			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; presc	riber may determine o	dispe		
Tab 10 mg		100		Norpress Norpress
Tab 25 mg	5.98	180	✓ [Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective			
TRANYLCYPROMINE SULPHATE				
Tab 10 mg	12.85	28	✓	Parnate S29 S29
	22.94	50	✓	Parnate
	45.88	100	✓	Parnate S29 S29
	96.00		✓	Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg	6.40	60	✓.	Aurorix
* Tab 300 mg	9.80	60	1	Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	1.52	84	✓	PSM Citalopram
ESCITALOPRAM			•	
* Tab 10 mg	1.07	28	✓	Escitalopram
				(Ethics)
	1.40		•	Escitalopram- Apotex
* Tab 20 mg	1.92	28	✓	Escitalopram
	0.40			(Ethics)
	2.49		•	Escitalopram- Apotex
(Escitalopram-Apotex Tab 10 mg to be delisted 1 October 2021) (Escitalopram-Apotex Tab 20 mg to be delisted 1 October 2021)				
FLUOXETINE HYDROCHLORIDE	4.00	00		Fl
* Tab dispersible 20 mg, scored - Subsidy by endorsement Subsidised by endorsement	1.98	30	•	Fluox
When prescribed for a patient who cannot swallow accordingly; or	whole tablets or caps	sules	and the pre	escription is endorsed
When prescribed in a daily dose that is not a multiple endorsed. Note: Tablets should be combined with a state of the combined with t	•			•
Cap 20 mg	2 91	84	✓	Fluox
PAROXETINE		0-7		11447
* Tab 20 mg	1 20	30	1	Paxtine
7- Tub 20 Hig	3.61	90		Loxamine
	3.01	90	•	LOXUIIIIIC

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
SERTRALINE				
* Tab 50 mg	0.92	30		Setrona Setrona AU
	3.05	90	✓	Arrow-Sertraline
* Tab 100 mg	1.61	30		Setrona Setrona AU
	5.25	90		Arrow-Sertraline
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.63	30	1	Apo-Mirtazapine
Tab 45 mg	3.48	30	/	Apo-Mirtazapine
/ENLAFAXINE				
* Cap 37.5 mg	6.38	84	1	Enlafax XR
* Cap 75 mg	8.11	84	1	Enlafax XR
* Cap 150 mg	11.16	84	✓	Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM - Safety medicine; prescriber may determ	nine dispensing frequency			
Inj 1 mg per ml, 1 ml		5	✓	Rivotril
(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)				
DIAZEPAM - Safety medicine; prescriber may determine	dispensing frequency			
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsen	nent23.66	5	✓	Hospira
a) Up to 5 inj available on a PSO				
b) Only on a PSO				
 c) PSO must be endorsed "not for anaesthetic pr 				
Rectal tubes 5 mg - Up to 5 tube available on a PSO	43.50	5	•	Stesolid
PARALDEHYDE				
* Inj 5 ml	1,500.00	5	1	AFT S29
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available	on a PSO88.63	5	1	Hospira
* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available				
PSO		5	1	Hospira
Control of Epilepsy				
CARBAMAZEPINE				
* Tab 200 mg		100	✓	Tegretol
* Tab long-acting 200 mg		100	✓	Tegretol CR
* Tab 400 mg	34.58	100	1	Tegretol
* Tab long-acting 400 mg	39.17	100	1	Tegretol CR
* Oral liq 20 mg per ml	26.37	250 n	nl 🗸	Tegretol
CLOBAZAM - Safety medicine; prescriber may determine	dispensing frequency			
Tab 10 mg		50	✓	Frisium
CLONAZEPAM - Safety medicine; prescriber may determ	nine dispensina freauency			

	Subsidy		Fully	Brand or
	(Manufacturer's P	rice) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
ETHOSUXIMIDE				
Cap 250 mg	140.88	100	✓ Z	arontin
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Z	arontin
GABAPENTIN				
Note: Not subsidised in combination with subsidised prega	abalin			
* Cap 100 mg	2.65	100	✓ <u>A</u>	po-Gabapentin
* Cap 300 mg	4.07	100	✓ Ā	po-Gabapentin
* Cap 400 mg		100	✓ A	po-Gabapentin
LACOSAMIDE - Special Authority see SA1125 below - Retail	pharmacy			
▲ Tab 50 mg	25.04	14	✓ V	'impat
▲ Tab 100 mg		14	✓ V	impat
ŭ	200.24	56	✓ ∨	'impat
▲ Tab 150 mg	75.10	14	✓ ∨	'impat
ŭ	300.40	56	✓ V	'impat
▲ Tab 200 mg	400.55	56	✓ V	'impat

⇒SA1125 Special Authority for Subsidy

LAMOTRIGINE

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

L/ \IV	OTTIGINE			
A .	Tab dispersible 2 mg5	5.00	30	✓ Lamictal
A .	Tab dispersible 5 mg	0.00	30	✓ Lamictal
*	Tab dispersible 25 mg	2.76	56	✓ Logem
*	Tab dispersible 50 mg	3.31	56	✓ Logem
*	Tab dispersible 100 mg	4.40	56	✓ Logem
LEV	ETIRACETAM			
	Tab 250 mg	4.99	60	✓ Everet
	Tab 500 mg		60	✓ Everet
	Tab 750 mg1		60	✓ Everet
	Tab 1,000 mg1		60	✓ Everet
	Oral liq 100 mg per ml4		0 ml OP	✓ Levetiracetam-AFT
	NOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae, page 237			
*	Tab 15 mg	0.00	500	✓ PSM
*	Tab 30 mg4	0.00	500	✓ PSM
PHE	NYTOIN SODIUM			
*	Tab 50 mg	5.00	200	✓ Dilantin Infatab
	Cap 30 mg	4.00	200	✓ Dilantin
	Cap 100 mg	37.00	200	✓ Dilantin
*	Oral liq 30 mg per 5 ml2	2.03 5	600 ml	✓ Dilantin

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
PREGABALIN				
Note: Not subsidised in combination with subsidised gabap	entin			
* Cap 25 mg	2.25	56	1	Pregabalin Pfizer
* Cap 75 mg	2.65	56	✓	Pregabalin Pfizer
* Cap 150 mg	4.01	56	✓	Lyrica
			✓	Pregabalin Pfizer
* Cap 300 mg	7.38	56	✓	Pregabalin Pfizer
PRIMIDONE				
* Tab 250 mg	17.25	100	1	Apo-Primidone
	62.00	200		Mysoline S29 S29
(Mysoline S29 S29 Tab 250 mg to be delisted 1 July 2021)	02.00			,
SODIUM VALPROATE	40.05	400	,	Fulling Owner habits
Tab 100 mg		100		Epilim Crushable
Tab 200 mg EC		100		Epilim
Tab 500 mg EC	52.24	100	•	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 m	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail p	nharmacy			-
Cap 250 mg	•	60	1	Diacomit \$29
Powder for oral liq 250 mg sachet	509.29	60	•	Diacomit S29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE	07	00	.
▲ Tab 25 mg11	.07	60	✓ Arrow-Topiramate
			✓ Topiramate Actavis
26	.04		✓ Topamax
▲ Tab 50 mg18	.81	60	 Arrow-Topiramate
			✓ Topiramate Actavis
44	.26		✓ Topamax
▲ Tab 100 mg31	.99	60	✓ Arrow-Topiramate
-			✓ Topiramate Actavis
75	.25		✓ Topamax
▲ Tab 200 mg55	.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
129	.85		✓ Topamax
▲ Sprinkle cap 15 mg20	.84	60	✓ Topamax
Sprinkle cap 25 mg26		60	✓ Topamax
/IGABATRIN – Special Authority see SA1997 on the next page – Retail p	harmacy		
▲ Tab 500 mg119	.30	100	✓ Sabril

Subsidy		Fully
(Manufacturer's Price)		Subsidised
\$	Per	/

sidised Ger Mar

Brand or Generic Manufacturer

⇒SA1997 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Fither:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Fither:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

RIZATRIPTAN Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	24.44	100	✓ Apo-Sumatriptan
Tab 100 mg	46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per			
prescription	34.00	2 OP	✓ Imigran

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50 PIZOTIFEN

1 12	OTH EN		
*	Tab 500 mcg23.21	100	✓ Sandomigran

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy
Cap 2 × 80 mg and 1 × 125 mg......84.00 3 OP ✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

* Tab To mg	84	vergo 16
CYCLIZINE HYDROCHLORIDE Tab 50 mg	10	✓ Nausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml	10	✓ Hameln
DOMPERIDONE		
* Tab 10 mg	100	✓ Pharmacy Health
* Inj 400 mcg per ml, 1 ml ampoule	10	✓ Martindale \$29
pharmacy14.11	2	✓ Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

* Tab 10 mg – Up to 30 tab available on a PSO1.30	100	✓ <u>Metoclopramide</u> Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50 ONDANSETRON	10	✓ <u>Pfizer</u>
* Tab 4 mg2.68	50	✓ Onrex
* Tab disp 4 mg – Up to 10 tab available on a PSO	10	✓ Ondansetron ODT-DRLA
* Tab 8 mg4.57	50	✓ Onrex
* Tab disp 8 mg – Up to 10 tab available on a PSO1.13	10	✓ Ondansetron ODT-DRLA
PROCHLORPERAZINE		
* Tab 3 mg buccal	50	
(30.00)		Buccastem
* Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
* Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
` \$	Per	1	Manufacture

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dispe	ensing frequenc	СУ	
Tab 100 mg		30	✓ Sulprix
	17.16	100	✓ Amisulpride
			Mylan S29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix
ARIPIPRAZOLE – Safety medicine; prescriber may determine disp		ncv	
Tab 5 mg		30	✓ Aripiprazole Sandoz
	28.58	49	✓ Aripiprazole 1A
			Pharma S29
Tab 10 mg	17.50	30	✓ Aripiprazole Sandoz
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	✓ Aripiprazole Sandoz
(Aripiprazole 1A Pharma S29 Tab 5 mg to be delisted 1 June 202		00	7 III pipi azolo canacz
	*		. ,
CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; pres			
Tab 10 mg - Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Tab 25 mg — Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Tab 100 mg – Up to 30 tab available on a PSO		100	✓ <u>Largactil</u>
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	30.79	10	✓ <u>Largactil</u>
CLOZAPINE – Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequen	•		
Tab 25 mg		50	✓ Clozaril
	6.69		✓ Clopine
	11.36	100	✓ Clozaril
T-h 50	13.37	50	✓ Clopine
Tab 50 mg		50	✓ Clopine
Toh 100 mg	17.33	100 50	✓ Clopine ✓ Clozaril
Tab 100 mg	17.33	50	✓ Clozarii ✓ Clopine
	29.45	100	✓ Clopine ✓ Clozaril
	34.65	100	✓ Clopine
Tab 200 mg		50	✓ Clopine
1 ab 200 mg	69.30	100	✓ Clopine
Suspension 50 mg per ml		100 ml	✓ Clopine
Casponsion of mg per minimum.	67.62	1001111	✓ Versacloz
HALORERIDOL Cofety medicines are suited as a second at	*****		701000102
HALOPERIDOL – Safety medicine; prescriber may determine disp Tab 500 mcg – Up to 30 tab available on a PSO		•	./ Coveres
5 1		100	✓ <u>Serenace</u>✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO Tab 5 mg - Up to 30 tab available on a PSO		100 50	✓ <u>Serenace</u> ✓ Serenace
1 ab 3 mg - 0p to 30 tab available on a F30	29.72	100	✓ Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSC		100 1111	✓ Serenace
my o my por mi, i mi ampoulo op to o my avallable on a i oc	, <u>L</u> 1.00	10	- Jerenace

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
LEVOMEPROMAZINE - Safety medicine; prescriber may dete	rmine dispensing frequ	uency		
Tab 25 mg (33.8 mg as a maleate)	16.10	100	1	Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100	•	Nozinan
Tab 100 mg (135 mg as a maleate)	41.75	100	•	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	•	<u>Nozinan</u>
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may determ	nine di	spensing	frequency
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
LITHIUM CARBONATE – Safety medicine; prescriber may dete		uonov		
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100		Douglas
, ,		100	•	Douglas
OLANZAPINE – Safety medicine; prescriber may determine dis			_	
Tab 2.5 mg		28	_	Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28	_	Zypine
Tab orodispersible 10 mg	2.38	28	•	Zypine ODT
PERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg	10.49	84	✓	Neulactil
•	12.49	100	✓	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
	44.45	100	✓	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	_	Quetapel
v				
RISPERIDONE – Safety medicine; prescriber may determine di		60	./	Risperidone (Teva)
Tab 0.5 mg Tab 1 mg		60		Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 3 mg		60	_	Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral lig 1 mg per ml		30 ml		Risperon
		00 1111	•	порстоп
ZIPRASIDONE – Safety medicine; prescriber may determine di			,	
Cap 20 mg		60	_	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60		Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	escriber may determin	e disp	•	' '
Tab 10 mg	31.45	100	•	Clopixol
Depot Injections				
FILIDENTINO DECANOATE Cofety mandicine and the same		da a fo		
FLUPENTHIXOL DECANOATE – Safety medicine; prescriber n	•	-		Fluencel
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	•	Fluanxol

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	ay determine dispensir	ng freque	ency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	,	5 '		laldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO	55.90	5		laldol Concentrate laldol
				Decanoas S29
OLANZAPINE – Special Authority see SA1428 below – Retail pl Safety medicine; prescriber may determine dispensing frequ	•			
Inj 210 mg vial	252.00	1	✓ Z	yprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Z	yprexa Relprevv
Inj 405 mg vial	504.00	1	✓ <u>Z</u>	yprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing fr	requency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe		1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see SA1427 on the next page – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency			
Inj 25 mg vial	135.98	1	✓ Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Anx	iol۱	/tics

BUSPIRONE HYDROCHLORIDE * Tab 5 mg		100	Orion
* Tab 10 mg	13.16	100	✓ <u>Orion</u>
CLONAZEPAM - Safety medicine; prescriber may determine	dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg		500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine di	spensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ <u>Ativan</u>
OXAZEPAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 10 mg	6.17	100	✓ Ox-Pam
Tab 15 mg	8.53	100	✓ Ox-Pam

Multiple Sclerosis Treatments

⇒SA2026 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

past 24 months; and

- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2026 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

 Cap 120 mg
 520.00
 14
 ✓ Tecfidera

 Cap 240 mg
 2,000.00
 56
 ✓ Tecfidera

FINGOLIMOD - Special Authority see SA2026 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Cap 0.5 mg......2,200.00 28 Gilenya

GLATIRAMER ACETATE - Special Authority see SA2026 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

lnj 40 mg prefilled syringe.................2,275.00 12 **✓ Copaxone**

INTERFERON BETA-1-ALPHA - Special Authority see SA2026 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer \$ INTERFERON BETA-1-BETA - Special Authority see SA2026 on page 134 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. NATALIZUMAB - Special Authority see SA2026 on page 134 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. OCRELIZUMAB - Special Authority see SA2026 on page 134 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Inj 30 mg per ml, 10 ml vial......9,346.00 ✓ Ocrevus TERIFLUNOMIDE - Special Authority see SA2026 on page 134 - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 28 ✓ Aubagio Aubagio to be Sole Supply on 1 June 2021 Sedatives and Hypnotics MELATONIN - Special Authority see SA1666 below - Retail pharmacy Tab modified-release 2 mg - No more than 5 tab per day28.22 30 ✓ Circadin ⇒SA1666 Special Authority for Subsidy Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and 4 Patient is aged 18 years or under*. Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient is aged 18 years or under*; and

- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine dispe	ensing frequency		
Inj 1 mg per ml, 5 ml ampoule	2.98	10	Mylan Midazolam
, 01	4.30		✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available)		
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for state	tus epilepticu	ıs use only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available	on		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for state	tus epilepticu	ıs use onlv.

			NEF	RVOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PHENOBARBITONE SODIUM – Special Authority see SA1386 b		acy 10	✓	Max Health \$29
■ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both: 1 For the treatment of terminal agitation that is unresponsive	to other agents; and		nless notifi	ed for applications meeting
The applicant is part of a multidisciplinary team working in TEMAZEPAM – Safety medicine; prescriber may determine dispertite to 10 mg	ensing frequency	25	✓ !	<u>Normison</u>
TRIAZOLAM – Safety medicine; prescriber may determine disper Tab 125 mcg	5.10 (9.85) 4.10	100		Нурат
ZOPICLONE – Safety medicine; prescriber may determine dispertable 7.5 mg		500		⊣ypam <mark>Zopiclone Actavis</mark>
Stimulants/ADHD Treatments				
ATOMOXETINE Cap 10 mg	18.41 107.03	28		Generic Partners Strattera

ATOMOXETINE			
Cap 10 mg	18.41	28	 Generic Partners
•	107.03		✓ Strattera
Cap 18 mg	27.06	28	✓ Generic Partners
Cap 25 mg	29.22	28	✓ Generic Partners
Cap 40 mg	29.22	28	✓ Generic Partners
	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ Generic Partners
Cap 80 mg	56.45	28	✓ Generic Partners
Cap 100 mg	58.48	28	✓ Generic Partners
DEXAMFETAMINE SULFATE - Special Authority	see SA1149 below – Retail pha	ırmacy	
a) Only on a controlled drug form		•	
b) Safety medicine; prescriber may determine	dispensing frequency		
Tab 5 mg	1 0 1 7	100	✓ PSM

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:



	Subsidy	Full	/ Brand or
(Manu	facturer's Price)	Subsidise	d Generic
	\$	Per 🗸	Manufacturer

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency Tab immediate-release 5 mg.......3.20 30 ✓ Rubifen ✓ Ritalin 30 ✓ Rubifen Tab extended-release 18 mg......7.75 ✓ Methylphenidate ER 30 - Teva ✓ Rubifen 30 ✓ Rubifen SR 30 ✓ Ritalin SR 100 ✓ Methylphenidate ER 30 - Teva ✓ Methylphenidate ER 30 - Teva Tab extended-release 54 mg......22.25 ✓ Methylphenidate ER 30 - Teva

(Ritalin SR Tab sustained-release 20 mg to be delisted 1 June 2021)

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 27 mg65.4430✓ ConcertaTab extended-release 36 mg71.9330✓ Concerta
Tab extended-release 54 mg
Cap modified-release 10 mg
Cap modified-release 20 mg20.40 30 ✓ Ritalin LA
Cap modified-release 30 mg25.52 30 ✓ Ritalin LA
Cap modified-release 40 mg30.60 30 ✔ Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or



Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 ` \$	Per	✓	Manufacturer

4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia DONEPEZIL HYDROCHLORIDE ★ Tab 5 mg 4.34 90 ✓ Donepezil-Rex ★ Tab 10 mg 6.64 90 ✓ Donepezil-Rex RIVASTIGMINE – Special Authority see SA1488 below – Retail pharmacy 48.75 30 ✓ Generic Partners Patch 4.6 mg per 24 hour 48.75 30 ✓ Generic Partners Patch 9.5 mg per 24 hour 48.75 30 ✓ Generic Partners

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

28

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ <u>Buprenorphine</u> Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg.......11.00 30 ✓ Zyban

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
DISULFIRAM Tab 200 mg	250.00	100	✓ A	ıntabuse	
NALTREXONE HYDROCHLORIDE – Special Authority see SA1 Tab 50 mg	'	harmacy 30	_	laltraccord	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

 b) Note: Direct Provision by a pharmacist permitted under the p 	rovisions in	Part I of Sectio	n A.
Patch 7 mg - Up to 28 patch available on a PSO	18.14	28	Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	Habitrol
Patch 14 mg - Up to 28 patch available on a PSO	19.95	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	22.86	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	21.02	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	38.21	384	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	44.17	384	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.01	96	Habitrol
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	44.17	384	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 422	5.64	53 OP	✓ Champix
·			✓ Varenicline Pfizer
Tab 1 mg	7.10	56	✓ Champix
·			✓ Varenicline Pfizer

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see \$A2046 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

(Ma	Subsidy	F	ully	Brand or
	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	1	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		· ·	
Inj 100 mg vial	1 387 00	1	✓ BiCNU
ing 100 mg viai	1,507.00	'	✓ Bicnu Heritage \$29
Ini 100 ma for ECD	1 207 00	100 mg OP	✓ Baxter
Inj 100 mg for ECP	1,307.00	100 mg OF	Daxiei
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	 Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	19.70	1	✓ DBL Cisplatin
	21.00		 Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
rab oo mg	158.00	100	✓ Procytox S29
Wastage claimable	156.00	100	V Procytox 329
Inj 1 g vial – PCT – Retail pharmacy-Specialist	25.65	1	✓ Endoxan
inj i g viai – POT – Hetali phamiacy-Specialist	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist		1	✓ Endoxan
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
	0.04	ring	Daxiei
IFOSFAMIDE – PCT only – Specialist	22.22	i i	4 11 1
lnj 1 g		1	✓ Holoxan
Inj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
	Ψ	rei		Manuacturei
_OMUSTINE - PCT - Retail pharmacy-Specialist	100.50		,	
Cap 10 mg		20		CeeNU
Cap 40 mg	399.15	20	•	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist		25		Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	•	Alkeran
			1	Alkeran S29 S29
	420.00		1	Tillomed S29
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
,		•		100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg		Baxter
THIOTEPA – PCT only – Specialist		9		
, ,	000		,	D - 46 1 000
Inj 15 mg vial	BS	1		Bedford S29
				THIO-TEPA S29
			/	Tepadina S29
Inj 100 mg vial	CBS	1	1	Tepadina S29
A malino actacle a lista a				
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see S	A1467 below			
Inj 100 mg vial	139.00	1	1	Azacitidine Dr

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or

605.00

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

✓ Vidaza

✓ Baxter

	Subsidy (Manufacturer's Price)	20) C··	Fully Brand or
	(Manufacturer's Prior	ce) Su Per	bsidised Generic Manufacturer
CALCIUM FOLINATE			
Tab 15 mg – PCT – Retail pharmacy-Specialist	114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia		1	✓ <u>Calcium Folinate</u> Sandoz
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓ Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓ Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Calcium Folinate Sandoz
Inj 1 g - PCT only - Specialist	67.51	1	✓ Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter
CAPECITABINE – Retail pharmacy-Specialist	40.00	00	()
Tab 500 mg		60 120	✓ <u>Capercit</u> ✓ Capercit
Tab 500 mg	49.00	120	Capercit
CLADRIBINE - PCT only - Specialist Inj 1 mg per ml, 10 ml	7/0.06	1	✓ Leustatin
Inj 10 mg for ECP		10 mg OP	✓ Baxter
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	alist400.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail		•	
pharmacy-Specialist	41.36	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	alist80.00	100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg OP	✓ Baxter
FLUOROURACIL			4
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	✓ Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1 100 mg	✓ Fluorouracil Ebewe✓ Baxter
Inj 1 mg for ECP – PCT only – Specialist	0.00	100 mg	Daxiei
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist	60.50	1	✓ DPI Compitabine
Inj 1 g, 26.3 ml vial		1	✓ DBL Gemcitabine ✓ Gemcitabine Flower
Inj 1 mg for ECP		1 mg	 ✓ Gemcitabine Ebewe ✓ Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist	0.02	illy	· DUALGI
Inj 20 mg per ml, 5 ml vial	71 44	1	✓ Irinotecan
ing 20 mg por mi, o mi viai		I	Accord \$29
			✓ Irinotecan Actavis
			100
lei 4 mm for FOR	100.00	4.	✓ Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Prio \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
MERCAPTOPURINE Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	√ P	Puri-nethol	
Oral suspension 20 mg per ml – Retail pharmacy-Specialist Special Authority see SA1725 below		100 ml (OP ✓ A	Allmercap	

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ <u>Trexate</u>
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
	, , , ,		Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
•	,gp	·	Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
~	ing 10 mg promised syringe14.77	'	Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
不	ing 20 mg premied symble14.00	1	Sandoz
	1.05		
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
			Sandoz
*	Inj 30 mg prefilled syringe15.09	1	Methotrexate
			Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL
			Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
			Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
~	pharmacy-Specialist	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist4.73	5 mg OP	✓ Baxter
-		5 mg Oi	Daxiei
PE	METREXED - PCT only - Specialist - Special Authority see SA1679 below		
	Inj 100 mg vial60.89	1	Juno Pemetrexed
	Inj 500 mg vial217.77	1	Juno Pemetrexed
	Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE – PCT – Retail pharmacy-Specialist			
Tab 40 mg	126.31	25	Lanvis

Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	161.01	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see	SA1889 on the n	ext page	
Inj 3.5 mg vial	105.00	1	✓ Bortezomib Dr-Reddy's
Inj 1 mg for ECP	31.20	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.			
DACARBAZINE – PCT only – Specialist	60.70	1	✓ DBL Dacarbazine
Inj 200 mg vial	52.70 580.60	10	✓ DBL Dacarbazine ✓ Dacarbazine
	360.00	10	APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		Ü	
Inj 2 mg per ml, 10 ml	149 50	1	✓ Pfizer
Inj 20 mg for ECP		20 mg OP	✓ Baxter
	170.00	20 mg Oi	Daxiei
DOCETAXEL - PCT only - Specialist	10.40		A DDI Deceteval
Inj 10 mg per ml, 2 ml vial		1	✓ DBL Docetaxel
Inj 20 mg	48.75	1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial	46.89	1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg		1	Docetaxel Sandoz
Inj 1 mg for ECP		1 mg	✓ Baxter
(DBL Docetaxel Inj 10 mg per ml, 2 ml vial to be delisted 1 June	2021)		
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebewe
, ,	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	56.15	1	 Doxorubicin Ebewe
	65.00		✓ Arrow-Doxorubicin
Inj 1 mg for ECP	0.29	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP	0.43	1 mg	✓ Baxter
ETOPOSIDE		Ü	
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	✓ Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓ Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special		10	✓ Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
		i ilig	- BUNG!
ETOPOSIDE PHOSPHATE – PCT only – Specialist	40.00	4	/ Flamenhaa
Inj 100 mg (of etoposide base)		1	✓ Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	rmacy-Specialist			
Cap 500 mg	23.82	100	✓ [<u>Devatis</u>
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial - PCT only - Specialist	93.00	1	1	Zavedos
Inj 10 mg vial - PCT only - Specialist		1	✓ 7	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	21.84	1 mg	✓ [Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Author Wastage claimable	ity see SA2047 below	1		
Cap 5 mg	5.122.76	28	✓ [Revlimid
Cap 10 mg		21	√ [Revlimid
	6,207.00	28	✓ F	Revlimid
Cap 15 mg	5,429.39	21	✓ [Revlimid
	7,239.18	28	√ [Revlimid
Cap 25 mg	7,627.00	21	✓ [Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 2.1 Long
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

	Subsidy	.\ 0l-	Fully	Brand or
	(Manufacturer's Price \$	Per	sidised •	Generic Manufacturer
MESNA				
Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	1	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	1	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	1	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	1	Baxter
MITOMYCIN C - PCT only - Specialist		•		
Inj 5 mg vial	851.37	1	1	Teva
Inj 20 mg vial		1	1	Omegapharm S29
11) 20 119 1141		•		Teva
Inj 1 mg for ECP	288.09	1 mg		Baxter
(Teva Inj 5 mg vial to be delisted 1 June 2021)	200.00	9		
MITOZANTRONE - PCT only - Specialist				
Inj 2 mg per ml, 10 ml vial	97.50	1	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	_	Baxter
		illig	•	Daxlei
DLAPARIB – Retail pharmacy-Specialist – Special Authority se				
Tab 100 mg		56		Lynparza
Tab 150 mg		56	_	Lynparza
Cap 50 mg - Wastage claimable	7,402.00	448	1	Lynparza
Lynparza Cap 50 mg to be delisted 1 July 2021)				

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

	Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
PACLITAXEL - PCT only - Specialist				
Inj 30 mg	47.30	5	✓	Paclitaxel Ebewe
Inj 100 mg		1	✓	Paclitaxel Ebewe
•	91.67		✓	Paclitaxel Actavis
Inj 150 mg	26.69	1	✓	Paclitaxel Ebewe
	137.50		✓	Anzatax
			✓	Paclitaxel Actavis
Inj 300 mg	44.00	1	✓	Paclitaxel Ebewe
	275.00		✓	Anzatax
			✓	Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓	Baxter
PEGASPARGASE - PCT only - Special Authority see SA1979 be	low			
Inj 750 iu per ml, 5 ml vial		1	✓	Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Speci	alist		
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharma	acy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the ne	xt page – Retail phar	macy	
Cap 5 mg	9.13	5	✓ <u>Temaccord</u>
Cap 20 mg - No patient co-payment payable	16.38	5	✓ <u>Temaccord</u>
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord \$29
Cap 100 mg	35.98	5	✓ <u>Temaccord</u>
	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord \$29
Cap 140 mg	50.12	5	✓ <u>Temaccord</u>
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord \$29
Cap 250 mg	86.34	5	✓ <u>Temaccord</u>
	688.00		✓ Amneal S29

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

Subsidy (Manufacturer's F	rice) Su	Fully	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

- Fither:
 - 1 The patient has multiple myeloma; or
 - 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 belo)W	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		14 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8.209.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax: and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VINBLASTINE SULPHATE			
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓	DBL Vinblastine S29
		•	Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	•	Baxter
VINCRISTINE SULPHATE			
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	1	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	·	Baxter
VINORELBINE - PCT only - Specialist			
Inj 10 mg per ml, 1 ml vial12.00	1	✓	Navelbine
42.00		✓	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓	Navelbine
210.00		✓	Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	·	Baxter

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable 224 ✓ Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

vvastage ciaimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7.692.58	60	✓ Sprycel
	,		-1. 7

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day: or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB – Retail pharmacy-Specialist – Special Authority see SA2000 below

Tab 100 mg	764.00	30	Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

⇒SA2000 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
 - 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
 - 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
 - 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2001 below

Tab 250 mg1,700.00 30 ✓ Iressa

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and

- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460

*	Cap 100 mg	58.23	60	Imatinib-Rex
	,	98.00		✓ Imatinib-AFT
	Imatinib-Rex to be Sole Supply on 1 June 2021			
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex
		197.50		✓ Imatinib-AFT

Imatinib-Rex to be Sole Supply on 1 June 2021

(Imatinib-AFT Cap 100 mg to be delisted 1 June 2021) (Imatinib-AFT Cap 400 mg to be delisted 1 June 2021)

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 **PHARMAC** Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

70 ✓ Tykerb

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

continued...

✓ Glivec

Subsidy	Fı	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	•	Manufacturer

continued...

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable			
Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CMI treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
 - 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
 - 3 Maximum nilotinib dose of 800 mg/day; and
 - 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable	•		
Cap 75 mg	4,000.00	21	✓ Ibrance
Cap 100 mg	4,000.00	21	✓ Ibrance
Cap 125 mg	4,000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and

4.2.2 Either:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
- 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB	- Special Authority see	SA1190 below -	Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
RUXOLITINIB – Special Authority see SA1890 below – Retail pl	narmacy			
Wastage claimable				
Tab 5 mg	2,500.00	56	•	Jakavi
Tab 15 mg	5,000.00	56	✓	Jakavi
Tab 20 mg	5,000.00	56	✓	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2002 below - Retail pharmacy

Cap 12.5 mg	2,315.38	28	Sutent
Cap 25 mg	· · · · · · · · · · · · · · · · · · ·	28	✓ Sutent
Cap 50 mg	9,261.54	28	✓ Sutent

⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see SA2003 on the next page Wastage claimable

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline: and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone: and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg	1.36	10	✓ Calutide-50 S29
-	4.07	30	✓ Binarex
	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Sp	ecial Authority see SA1895 below		
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MEGESTROL ACETATE	00.50			A M
Tab 160 mg OCTREOTIDE	63.53	30	•	Apo-Megestrol
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	1	Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	/	Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	1	Octreotide
				MaxRx S29
	56.87		✓	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓	DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule		5	/	Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓	DBL Octreotide
	222.00		✓	Octreotide
				(Sun) \$29
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Special	Authority see SA2004	belo	w – Retail	pharmacv
Inj LAR 10 mg prefilled syringe		1		Sandostatin LAR
Inj LAR 20 mg prefilled syringe		1	✓	Sandostatin LAR
Inj LAR 30 mg prefilled syringe		1	✓	Sandostatin LAR

⇒SA2004 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application - (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued...

relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Fither:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

IΑ	WOXIFEN CITATE		
*	Tab 10 mg15.00	60	✓ Tamoxifen Sandoz
*	Tab 20 mg	60	✓ Tamoxifen Sandoz

Aromatase Inhibitors

TAMOVICENI CITRATE

ANASTROZOLE	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE	30	✓ <u>Letrole</u>

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg	5 60	✓ Azamun
* Tab 50 mg	0 100	 Azamun
* Inj 50 mg vial199.00	0 1	✓ Imuran
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	0 50	 Cellcept
Cap 250 mg35.90	0 100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187.25	5 165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

Fusion Proteins

ETANERCEPT	- Special Authority	see SA2048 below	 Retail pharmacy
------------	---------------------	------------------	-------------------------------------

Inj 25 mg	, ,	4	Enbrel
Inj 25 mg autoinjector		4	✓ Enbrel
Inj 50 mg autoinjector		4	✓ Enbrel
lnj 50 mg prefilled syringe	·	4	✓ Enbrel

⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
- 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment: and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
 - 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plague psoriasis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

. -...

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.1 Both:

- 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

2.2 Both:

- 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	ılist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG \$29 Inj 40 mg per ml, vial to be delisted 1 April	2022)		

Monoclonal Antibodies

		below – Retail pharmacy	ADALIMUMAB – Special Authority see SA2049 be
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
Humira	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2049 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992:19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes): and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy	Fully	/ Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

continued...

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab: or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 Patient has pyoderma gangrenosum*; and

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

2.6 Fither:

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints:
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Fither:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al. J Rheumatol. 2004;31:931-7.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:

 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
 - 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 212 Fither
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or
 - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
 - 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

1 Both:

- 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

1 ✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or

Subsidy	Fully Subsidised		Brand or
(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
Ψ	rei		Manufacturei

continued...

- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

		AB - PCT only - Specialist - Special Authority see SA1697 below	CETUXIMAB - PCT only - Specia
Erbitux	1	g per ml, 20 ml vial364.00	Inj 5 mg per ml, 20 ml vial
Erbitux	1	g per ml, 100 ml vial	Inj 5 mg per ml, 100 ml vial
✓ Baxter	1 ma	g for FCP 3.82	Ini 1 mg for FCP

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see SA2050	on the next page		
Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2050 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation: or

2 Roth:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss: and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose: or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	,	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Roth:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- Litner
 - 1 A withdrawal period has been tried and the patient has relapsed; or
 - 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:

Subs (Manufactur		
\$	Per	Manufacturer

continued...

- 2.3.1 There has been an improvement in MRI appearances; or
- 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Fither:

- 1 Roth:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plague psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

1 Fither

Both:

- 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values: or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plaque psoriasis: or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fullminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - Retail p	oharmacy		
Inj 100 mg prefilled pen	1,638.00	1	✓ Nucala
Ini 100 mg vial	1.638.00	1	✓ Nucala

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10°9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

years for applications meeting the following criteria:

Roth

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail phar	macy		
Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

continued...

exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and

- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	

continued...

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

Inj 100 mg per 10 ml vial		2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold: or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Fither:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

than 5 mg per day and has done so for more than three months; and

- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and

continued...

- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
 - 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2028 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2028 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or

(manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer		Subsidy (Manufacturer's Price) \$		Fully bsidised	Brand or Generic Manufacturer	
--	--	---	--	----------------	-------------------------------------	--

continued...

2.2.2 Both:

- 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
- 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
- 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per ✓	Manufacturer

continued...

- 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:

	Subsidy		Fully	Brand or
(Ma	nufacturer's Price)	Subsic	lised	Generic
	\$	Per	•	Manufacturer

continued...

- 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
- 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective:
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued...

applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles: or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — **(haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*: and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Manuacturer's Frice)		oubsidised		
\$	Per	/	Manufacturer	

continued...

practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	bsidised	Generic
 \$	Per	/	Manufacturer

continued...

- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*: and
- 2 Either
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

⇒SA2044 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	✓	Manufacturer

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.
 SILTUXIMAB Special Authority see SA1596 on the next page Retail pharmacy

Note	: Siltuximab is to	be administered at do	ses no greater than 11	l mg/kg eve	ry 3 weeks.	
lnj 1	00 mg vial			770.57	1	Sylvant
Ini /	10 ma vial		3	U82 33	1	✓ Sylvant

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1977 below

Inj 20 mg per ml, 4 ml vial220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial	1	✓ Actemra
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1977 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis: or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or

Subsidy	F	ılly	Brand or
(Manufacturer's	s Price) Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule: and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	 Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib: and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
 - 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
 - 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadiuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 on the next page

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	23.20	1 mg	✓ Baxter

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
· •	Por 🗸	Manufacturer

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

		NIVOLUMAB – PCT only – Specialist – Special Authority see SA2006 below	
Opdivo	1	Inj 10 mg per ml, 4 ml vial1,051.98	
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96	
✓ Baxter	1 mg	Inj 1 mg for ECP27.62	

⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- ılııeı.
 - 1 All of the following:
 - 1.1 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
- 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
- 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 1.2 Either:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Author	ority see SA2007 below		
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	49.14	1 mg	✓ Baxter

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither

Subsidy (Manufacturer's Price)	S	Fully subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 Patient has not received funded nivolumab; or
- 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Fither:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy		Fully	Brand or
(Manufacturer's Price)) :	Subsidised	Generic
\$	Per	•	Manufacturer

continued...

 Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg	. 177.81	50	✓ Neoral
Oral liq 100 mg per ml	. 198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pharm	асу		
Wastage claimable	F40.00	00	/ Att-11
Tab 10 mg6		•	✓ Afinitor
Tab 5 mg4	,555./6	30	✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml		60 ml OP	✓ Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy: or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has severe non-malignant lymphovascular malformation*; and

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
 - 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
 - 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009:45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*: and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam,

\$29 Unapproved medicine supplied under Section 29

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and

- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS	 Special Authority 	/ see SA1745	below – Retail pl	narmacv

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe..................2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above	ve – Retail pharmacy	
Initiation kit - 5 vials freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		
diluent285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		
9 ml, 3 diluent 1.8 ml305.00	0 1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA1367 at	oove – Retail pharmac	у
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	100	✓ Alle and
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	0 1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze	1.00	/ Vanamil coo
dried venom, with diluent305.00	0 1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Pr	ice) Subs	sidised Generic
	\$	Per	✓ Manufacturer
	<u>_</u>		
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
			4
* Tab 10 mg		100	✓ Zista
* Oral liq 1 mg per ml	3.37	200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE			
	0.27	500 ml	✓ Histafen
* Oral liq 2 mg per 5 ml	9.37	500 1111	▼ nistaleli
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
· · · · · · · · · · · · · · · · · · ·	(8.40)		Polaramine
	1.01	20	rolaramino
		20	Delementos
	(5.99)		Polaramine
* Oral liq 2 mg per 5 ml	1.77	100 ml	
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
	4.04	00	
* Tab 60 mg		20	-
	(8.23)		Telfast
* Tab 120 mg	4.74	10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
	(20.11)		Tondot
LORATADINE			
* Tab 10 mg	1.69	100	✓ Lorafix
* Oral liq 1 mg per ml	1.43	100 ml	Haylor syrup
	2.95	120 ml	✓ Lorfast
(Lorfast Oral liq 1 mg per ml to be delisted 1 September 2021)			
PROMETHAZINE HYDROCHLORIDE			_
* Tab 10 mg	1.68	50	✓ Allersoothe
* Tab 25 mg	1.89	50	✓ Allersoothe
* Oral lig 1 mg per 1 ml	2.69	100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a F	PSO 17.87	5	✓ Hospira
		•	
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	9.30	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
, 01			
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250
BUDESONIDE			
	17.00	200 dose OP	✓ Pulmicort
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
• • • • • • • • • • • • • • • • • • • •			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
i owder for illitalation, 400 filey per dose	02.00	LUU UUSE UF	
			Turbuhaler

	Subsidy		Fully Brand	
	(Manufacturer's F	Price) Sub Per	sidised Gener	ric facturer
	Ψ	1 61	• Manu	lacturer
LUTICASONE			4 -	
Aerosol inhaler, 50 mcg per dose		120 dose OP		
Powder for inhalation, 50 mcg per dose		60 dose OP		e Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP		e Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP		_
Aerosol inhaler, 250 mcg per dose		120 dose OP		_
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide	e Accuhaler
nhaled Long-acting Beta-adrenoceptor Agonis	ts			
FORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose devi		60 dose		
	(35.80)		Foradil	
FORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose	e)10.32	60 dose OP		
(oquirus in to ordinate or language or mag motores upon	(16.90)		Oxis Tur	buhaler
IDACATEROL	(10.00)		Oxio rui	Dariaioi
IDACATEROL	04.00	00 de e OD		D
Powder for inhalation 150 mcg		30 dose OP	✓ Onbrez	
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez	Breezhaler
ALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	Serever	ıt
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	✓ Serever	t Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocepto	or Agonists	;	
UDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide w	iith			
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoRes	p Spiromax
,		120 00se OF	Duones	p Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumar				
per dose (equivalent to 400 mcg budesonide with 12 mc	g			
eformoterol fumarate metered dose) - No more than 2	00.50	400 00		
dose per day		120 dose OP		p Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP		
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OP	-,	
				haler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP		
Powder for inhalation 200 mcg with eformoterol fumarate 6 n	ncg44.08	120 dose OP	- ,	
			Turbu	haler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg - No more than 2 dose per day	44.08	60 dose OP	✓ Symbice	ort
, ,			•	haler 400/12
LUTICASONE FUROATE WITH VILANTEROL				
	44.00	30 dose OP	√ Bros Ell	into
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.00	30 dose OP	✓ Breo Ell	ihra

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer	
FLUTICASONE WITH SALMETEROL	*			
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 79	120 dose OP	✓ Seretide	
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide	
Powder for inhalation 100 mcg with salmeterol 50 mcg - No				
more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler	r
Powder for inhalation 250 mcg with salmeterol 50 mcg - No)			
more than 2 dose per day	44.08	60 dose OP	 Seretide Accuhale 	r
Beta-Adrenoceptor Agonists				
SAL PLITAMOL				
SALBUTAMOL Oral lig 400 mcg per ml	20.00	150 ml	✓ Ventolin	
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin	
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Ventolin	
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	✓ Respigen✓ SalAir	
	(6.00)		Ventolin	
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb	` '		VEHIOIIII	
available on a PSO		20	✓ Asthalin	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO		20	✓ <u>Asthalin</u>	
FERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhale	er
Autich clinovnic Augusta				
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	✓ Atrovent	
a) Up to 400 dose available on a PSO				
b) No patient co-payment payable				
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 n		20		
available on a PSO	11./3	20	✓ <u>Univent</u>	
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg	per			
dose CFC-free	•	200 dose OP	✓ Duolin HFA	
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial 2.5 ml amnoula. Un to 20 nob available on a DSC	5 20	20	✓ Duolin	

vial, 2.5 ml ampoule – Up to 20 neb available on a PSO5.20

✓ Duolin

20

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Seebri Breezhaler

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose......50.37 30 dose ✓ Spiriva Soln for inhalation 2.5 mcg per dose......50.37 60 dose OP ✓ Spiriva Respimat

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Incruse Ellipta

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg.....81.00 ✓ Ultibro Breezhaler 30 dose OP

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg.....81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP ✓ Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg......2,554.00 60 OP ✓ Ofev 60 OP ✓ Ofev

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib

Tab 801 mg	3,645.00	90	✓ Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

Leukotriene Receptor Antagonists

МС	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
*	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	28	✓ Montelukast Mylan

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO18	0.00	5	✓ DBL Aminophylline
THE	EOPHYLLINE			
*	Tab long-acting 250 mg2	3.02	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml1	6.60	500 ml	✓ <u>Nuelin</u>

Mucolytics

		8 below – Retail pharmacy	DORNASE ALFA – Special Authority see SA1978
✓ Pulmozyme	6	250.00	Nebuliser soln, 2.5 mg per 2.5 ml ampoule

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	S	Subsidised	Generic	
	\$	Per	1	Manufacturer	
IVACAFTOR - PCT only - Specialist - Special Authority see SA	2017 below				
Tab 150 mg	29,386.00	56	✓ K	Calydeco	
Oral granules 50 mg, sachet	29,386.00	56	✓ K	alydeco	
Oral granules 75 mg, sachet	29,386.00	56	✓ K	Calydeco	

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHI ORIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE	5 23	15 ml OP	✓ Univent

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

Small 2.20 1 ✓ e-chamber Mask

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
	\$	rei		Manuacturei
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	1	Mini-Wright AFS
-				Low Range
Normal range	9.54	1	1	Mini-Wright
Č				Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	2.95	1	1	e-chamber Turbo
510 ml (single patient)	5.12	1	_	e-chamber La
o . o (eg.o pano)				Grande
800 ml	6.50	1	1	Volumatic
000 111				Volumento
Respiratory Stimulants				
nespiratory Stillidiants				
CAFFEINE CITRATE				
Oral liq 20 mg per ml (10 mg base per ml)	15.10 25	ml C)P 🗸	Biomed
The state of the s				

	Subsidy		Fully Brand or
	(Manufacturer's F \$	Price) Subs Per	sidised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1. 2- PROPANEDIOL DIACETATE AND BI	ENZETHONIUM		
For Vosol ear drops with hydrocortisone powder refer Stand	ard Formulae, pa	age 237	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE		00 1111 01	7 70001
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's ✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	TIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate	5.40	75 100	
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50	8 ml OP	
gramicium 50 mcg per mi	(9.27)	o IIII OF	Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
- B	()		, , , , , , , , , , , , , , , , , , ,
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expl	citly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%	1.54	10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * a CIPROFLOXACIN	e unapproved in	dications.	
Eye drops 0.3% – Subsidy by endorsement		5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis of the second line treatment of chronic suppurative otiting the second line treatment of chronic suppurative otiting the second line is an unapproved indication marked with a * is an un	s media (CSOM)		
GENTAMICIN SULPHATE			

5 ml OP

10 ml OP

5 g OP

(14.55)

✓ Genoptic

Brolene

✓ Fucithalmic

PROPAMIDINE ISETHIONATE

SODIUM FUSIDATE [FUSIDIC ACID]

* Eye drops 0.1%......2.97

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F	Price) S	Fully ubsidised	Brand or Generic
	\$	Per	1	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	✓ 1	Tobrex
Eye drops 0.3%	11.48	5 ml OP	✓ 1	Tobrex
Corticosteroids and Other Anti-Inflammatory Pr	eparations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ I	Maxidex
* Eye drops 0.1%		5 ml OP		Maxidex
Ocular implant 700 mcg - Special Authority see SA1680 belo				

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not vet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	sulphate 6,000 u per g5.3	39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml4.5		5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM Eye drops 0.1%13.8	80	5 ml OP	✓ Voltaren Ophtha

✓ Ozurdex

	Subsidy (Manufacturer's F	Price) Sub	Fully sidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
FLUOROMETHOLONE					
* Eye drops 0.1%	3.09	5 ml OP	✓ F	ML	
	5.20		√ F	lucon	
KETOROLAC TROMETAMOL - Special Authority see SA1981	below - Retail ph	armacy			
Eye drops 0.5%	9.50 ·	5 ml OP	√ p	Acular	
OA4004 Out a dad Anatha with a fam Ondra late.					

⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71	4 ml OP	
, , , , , , , , , , , , , , , , , , , ,	(10.34)		Livostin
	(10.01)		21100111
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eve drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
	г оо	40 ml OD	/ Duadwisslams AFT
Eye drops 1%		10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	- Retail pharn	nacv
Eve drops 0.5%, single dose (preservative free)		20 dose	✓ Minims
2,0 diopo 0.0/0, oliigio dobo (proborvativo 1100)		_0 0000	
			Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

5 ml OP

2.5 ml OP

✓ Rexacrom

✓ Arrow-Timolol

✓ Timoptol XE

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers		
BETAXOLOL		
* Eye drops 0.25%	5 ml OP	✓ Betoptic S
* Eye drops 0.5%	5 ml OP	✓ Betoptic
TIMOLOL		
* Eve drops 0.25%	5 ml OP	✓ Arrow-Timolol

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer
Glaucoma Preparations - Carbonic Anhydrase I	nhibitors		
ACETAZOLAMIDE	47.00	100	(D)
* Tab 250 mg	17.03	100	✓ Diamox
BRINZOLAMIDE * Eye drops 1%	7.30	5 ml OP	✓ Azopt
DORZOLAMIDE HYDROCHLORIDE			
* Eye drops 2%	9.77	5 ml OP	
	(17.44)		Trusopt
DORZOLAMIDE WITH TIMOLOL	0.07	E ml OB	✓ Dortiment
* Eye drops 2% with timolol 0.5%	2.87	5 ml OP	✓ <u>Dortimopt</u>
Glaucoma Preparations - Prostaglandin Analog	ues		
BIMATOPROST			
* Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost
ATANOPPOOT			Multichem
_ATANOPROST * Eve drops 0.005%	1.57	2.5 ml OP	✓ Teva
TRAVOPROST		2.0 0.	
* Eye drops 0.004%	7.30	5 ml OP	✓ Travopt
	10.50		✓ Mylan S29
	19.50	2.5 ml OP	✓ Travatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE			
* Eye drops 0.2%	12.25	5 ml OP	✓ Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE	40.50	5l OD	. Ozwitiwa
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL Eye drops 0.005% with timolol 0.5%	2 49	2.5 ml OP	✓ Arrow - Lattim
PILOCARPINE HYDROCHLORIDE			
* Eye drops 1%		15 ml OP	✓ Isopto Carpine
* Eye drops 2%		15 ml OP	✓ Isopto Carpine
Eye drops 4% Subsidised for oral use pursuant to the Standard Formul		15 ml OP	✓ Isopto Carpine
 Eye drops 2% single dose – Special Authority see SA0895 	iuo.		
below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine
⇒SA0895 Special Authority for Subsidy			

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Pri \$	ice) Subsi Per	Fully dised	Brand or Generic Manufacturer
Mydriatics and Cycloplegics				
ATROPINE SULPHATE * Eye drops 1% CYCLOPENTOLATE HYDROCHLORIDE	17.36	15 ml OP	✓ <u>A</u>	tropt
* Eye drops 1% * Eye drops 1%, single dose (preservative free) – Only on a	8.76	15 ml OP	✓ C	yclogyl
prescription	52.86	20 dose		inims Cyclopentolate
TROPICAMIDE * Eye drops 0.5% * Eye drops 1%		15 ml OP 15 ml OP		ydriacyl ydriacyl
Preparations for Tear Deficiency				
For acetylcysteine eye drops refer Standard Formulae, page 237 HYPROMELLOSE	,			
* Eye drops 0.5%	19.50	15 ml OP	✓ M	ethopt
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓ P	oly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Other Eve Preparations

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail phar	macy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authorit	y see SA1388	above – Retail	pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Author	rity see SA138	8 above – Reta	il pharmacy
Eye drops 1 mg per ml	22.00	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Phar	macy Procedu	res Manual res	triction allowing one bottle per
month is not relevant and therefore only the prescribed do	sage to the nea	arest OP may b	oe claimed.

Other Lyc i reparations		
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.20	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
	\$	Per	·	Manufacturer	
Variana			•		

Various

PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee4.50
1 fee

- ✓ BSF Ambrisentan Mylan
 - ✓ BSF Bisoprolol Mylan
 - ✓ BSF Darunavir Mylan
- a) The Pharmacode for BSF Ambrisentan Mylan is 2605309 see also page 58
- b) The Pharmacode for BSF Darunavir Mylan is 2607026 see also page 106
- c) The Pharmacode for BSF Bisoprolol Mylan is 2607034 see also page 50

(BSF Ambrisentan Mylan Brand switch fee to be delisted 1 June 2021)
(BSF Bisoprolol Mylan Brand switch fee to be delisted 1 July 2021)

(BSF Darunavir Mylan Brand switch fee to be delisted 1 July 2021)

Agents Used in the Treatment of Poisonings

Antidotes

ACF	$\Gamma V I$	\sim	/CT	N	ᆮ

Pharma \$29

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

Removal and Elimination

CHARCOAL

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

 Tab 125 mg dispersible
 276.00
 28
 Exjade

 Tab 250 mg dispersible
 552.00
 28
 Exjade

 Tab 500 mg dispersible
 1.105.00
 28
 Exjade

 Exjade
 Exjade
 Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 (Manufacturer's Frice)	Per	Jubsiuiseu ✓	

continued...

- 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis: or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - F	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> <u>Desferrioxamine</u> <u>Mesylate for Inj</u> <u>BP</u>
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

Standard i Orindiae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE ORAL LIQUID	4
Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	Phenobarbitone Sodium Glycerol BP	1 g 70 ml
Sultable eye drop base	Чэ	Water	to 100 ml
CODEINE LINCTUS (3 mg per 5 ml)			
Codeine phosphate	60 mg	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Glycerol Preservative	40 ml	mg per ml) Phenobarbitone Sodium	400 ma
Water	qs to 100 ml	Glycerol BP	400 mg 4 ml
Water	10 100 1111	Water	to 40 ml
CODEINE LINCTUS (15 mg per 5 ml)			
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol Preservative	40 ml gs	Pilocarpine 4% eye drops Preservative	qs qs
Water	to 100 ml	Water	to 500 ml
	10 100 1111	(Preservative should be used if quantity supplied is for	
FOLINIC MOUTHWASH		than 5 days.)	
Calcium folinate 15 mg tab	1 tab	SALIVA SUBSTITUTE FORMULA	
Preservative Water	qs to 500 ml	Methylcellulose	5 g
(Preservative should be used if quantity supplied is f		Preservative	qs
than 5 days. Maximum 500 ml per prescription.)		Water	to 500 ml
MAGNESIUM HYDROXIDE 8% MIXTURE		(Preservative should be used if quantity supplied is for	or more
Magnesium hydroxide paste 29%	275 g	than 5 days. Maximum 500 ml per prescription.)	
Methyl hydroxybenzoate	1.5 q	SODIUM CHLORIDE ORAL LIQUID	
Water	to 1,000 ml	Sodium chloride inj 23.4%, 20 ml	qs
METHADONE MIXTURE		Water	qs
METHADONE MIXTURE Methadone powder	ac	(Only funded if prescribed for treatment of hyponatra	ıemia)
Glycerol	qs qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection	10 vials
		Glycerol BP	40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION	10 ~	Water	to 100 ml
Methyl hydroxybenzoate Propylene glycol	10 g to 100 ml	(Only funded if prescribed for treatment of Clostridiur following metronidazole failure)	n difficile
(Use 1 ml of the 10% solution per 100 ml of oral liqui		following metrorildazole failure)	
	,	VOSOL EAR DROPS	
OMEPRAZOLE SUSPENSION	~~	WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder Sodium bicarbonate powder BP	qs 8.4 g	Hydrocortisone powder	1%
Water	to 100 ml	Vosol Ear Drops	to 35 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy

Fully

Brand or

	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully Brand or sidised Generic Manufacturer
Extemporaneously Compounded Preparations	and Galenica	İs	
CODEINE PHOSPHATE – Safety medicine; prescriber may det			
Powder – Only in combination		25 q	
, , , , , , , , , , , , , , , , , , , ,	(90.09)	- 3	Douglas
Only in extemporaneously compounded codeine linctus			
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the determined.	supplier and will be	e delisted fror	n the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination		100 1111	- 1 Om
Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ <u>Midwest</u>
GLYCERIN WITH SODIUM SACCHARIN - Only in combination	1		
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination			
Only in combination with Ora-Plus. Suspension	20.05	473 ml	✓ Ora-Sweet
	50.95	4/3 1111	▼ <u>Ora-Sweet</u>
GLYCEROL * Liquid – Only in combination	3 23	500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparation		000 1111	inculting dryocron bi
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing fr			and the second second second
 d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablets). 	reimbursed at the	rate of the ch	eapest form available
Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE		3	
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE			
Powder		100 g	✓ <u>MidWest</u>
Suspension – Only in combination		473 ml	✓ <u>Ora-Plus</u>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH	•		4.6 51 1.65
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On	,	470	A Ove Bland
Suspension	30.95	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM Powder – Only in combination	52.50	10 g	✓ MidWest
Fowder - Only in combination	325.00	10 g	✓ MidWest
Only in children up to 12 years	0_0.00	.00 9	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxyben:			
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE	10.05	F00	✓ Midweet

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

✓ Midwest

500 g

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>M</u>	<u>lidwest</u>	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 241

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	200 ml OP	✓ Calogen
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml114.92	4 OP	✓ Liquigen ´

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	 Special Authority see SA1524 above – Hospital p 	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufactu

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML	 Special Authority see SA1095 above 	– Hospital pharn	nacy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
	7.50	1,000 ml OP	Diason RTH
			✓ Glucerna Select
			RTH

(Glucerna Select RTH Liquid to be delisted 1 September 2021)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Sustagen Diabetic
	(2.10)	200 ml OP	Nutren Diabetes

(Glucerna Select Liquid (vanilla) to be delisted 1 September 2021) (Sustagen Diabetic Liquid (vanilla) to be delisted 1 October 2021)

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 243



Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

 Liquid
 54.00
 400 g OP
 ✓ Kindergen

 Powder
 54.00
 400 g OP
 ✓ Kindergen

(Kindergen Liquid to be delisted 1 August 2021)

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2.3 faltering growth in an infant/child; or
- 2.4 increased nutritional requirements; or
- 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see \$ Liquid		he previous pag 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA Liquid		previous page 500 ml OP	− Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Apharmacy [HP3]	Authority see	SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	379 on the r	revious page -	Hospital pharmacy [HP3]
Liquid (strawberry)		200 ml OP	✓ Fortini
Liquid (vanilla)		200 ml OP	✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA137	9 on the pre	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)		200 ml OP	✓ Pediasure
Liquid (strawberry)		200 ml OP	✓ Pediasure
Liquid (vanilla)		200 ml OP	✓ Pediasure
, ,	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authorharmacy [HP3]	ority see SA	1379 on the pre	evious page - Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)		200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the	e previous r	age - Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

245 ✓ fully subsidised

	Subsidy (Manufacturer's Pr \$	rice) Subsi Per	Fully idised	Brand or Generic Manufacturer
RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid		evious page – 500 ml OP		al pharmacy [HP3] lepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		us page – Hos 220 ml OP	✓ N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110 Liquid	2.88	page – Hospi 237 ml OP		,
Liquid (apricot) 125 ml Liquid (caramel) 125 ml		4 OP 4 OP	✓ R	lovaSource Renal Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Liquid	•		
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see	e SA1377 above	– Hospital phari	macy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above –	Hospital pharma	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Aut	hority see SA137	7 above – Hosp	ital pharmacy [HP3]
l iquid	12.04	1.000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per 🗸 Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 247

Subs (Manufactur		
\$	Per	Manufacturer

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per	· •	Manufacturer	

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube: or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum: or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3] Liquid......7.00 1.000 ml OP ✓ Nutrison Energy

✓ fully subsidised 249

	Subsidy		Fully Brand or
	(Manufacturer's		idised Generic
	\$	Per	✓ Manufacturer
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on Liquid		spital pharmacy 250 ml OP 1,000 ml OP	[HP3] ✓ Isosource Standard ✓ Nutrison Standard RTH ✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit Liquid		on page 247 – F 1,000 ml OP	
ENTERAL FEED WITH FIBRE 1 KCAL/ML — Special Authority so Liquid		0age 247 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority Liquid		page 247 – Ho 250 ml OP 1,000 ml OP	spital pharmacy [HP3] Finsure Plus HN Ensure Plus RTH Jevity HiCal RTH Nutrison Energy Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on page Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription. Powder (chocolate) – Higher subsidy of up to \$26.00 per 850	be reimbursed	for patients wit	h both a valid Special Authority
with Endorsement	26.00 9.54	850 g OP 840 g OP	✓ Ensure
	(26.00)	3	Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patier prescription must be endorsed accordingly. Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g	its with fat mala	bsorption, fat in	tolerance or chyle leak. The
with Endorsement	26.00 9.54	857 g OP 850 g OP 840 g OP	✓ Fortisip ✓ Ensure
	(26.00)		Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patier	its with fat mala	ibsorption, fat in	tolerance or chyle leak. The

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

(Fortisip Powder (vanilla) to be delisted 1 August 2021)

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26)	200 0.	Ensure Plus
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with	(1.26)		Fortisip
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	- D
	(1.26)		Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with	0.70	000 ml OD	
Endorsement	0.72 (1.26)	200 ml OP	Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with	(1.20)		rorusip
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 247 — Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.26)	200 0.	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

continued...

✓ fully subsidised 251

Subsidy (Manufacturer's	Ful Price) Subsidise	·
	Per •	Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
FOOD THICKENER - Special Authority see SA1106 on	1 0 1	•	,		
Powder		00 g OP		lutilis	
	7.25 3	80 g OP		eed Thickener Karicare Aptamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX — Special Authority see SA1729 above — Hospit Powder	al pharmacy [HP3] 1,000 g OP	
(5.15)		Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 above - Hospita	al pharmacy [HP3]	
Powder	1,000 g OP	
(7.32)		NZB Low Gluten Bread Mix
3.51		
(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above - Hospital pha	armacy [HP3]	
Powder	2,000 g OP	
(18.10)		Horleys Flour

✓ fully subsidised 253

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		sidised	Generic
	\$	Per		Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – Ho	ospital pharr	nacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)		C	Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		C	Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		C	Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)		C	Orgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)		C	Orgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)		C	Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet		30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton		18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX − Special Authority see SA1108 on the previous page − Hospital pharmacy [HP3]

Powder8.22 500 g OP

Loprofin Mix

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] Animal shapes11.91 500 q OP ✓ Loprofin ✓ Loprofin 250 g OP 500 g OP ✓ Loprofin 250 q OP ✓ Loprofin Macaroni 5.95 500 g OP ✓ Loprofin 500 g OP ✓ Loprofin 500 g OP ✓ Loprofin

✓ fully subsidised

255

Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see SA1940 below Powder	,. ,.	✓ Alfamino Junior
Powder (unflavoured)	53.00 400 g OP	✓ Elecare
,	ū	✓ Elecare LCP
		✓ Neocate Gold
		Neocate Junior Unflavoured
		✓ Neocate SYNEO
Powder (vanilla)	53.00 400 g OP	✓ Elecare
. ,	· ·	 Neocate Junior Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short aut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IqE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

257 ✓ fully subsidised

SPECIAL FOODS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut: or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	- Special Authority see SA1953 below	- Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption: or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see \$A1557 below - Hospital pharmacy [HP3]

Powder	15.21	450 g OP	 Aptamil Gold+ Pepti Junior
	30.42	900 g OP	 Aptamil AllerPro SYNEO 1
			 Aptamil AllerPro SYNEO 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

✓ fully subsidised

259



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.35 125 ml OP ✓ Infatrini

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > 0 requal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
 Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE			
(1	Subsidy Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	D HAEMOPHILUS I	NFLUENZAE TY	PE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria:			
1) Up to four doses for children up to and under the age of 1			- d d - u d - u - d - d
2) An additional four doses (as appropriate) are funded for (
10 who are patients post haematopoietic stem cell transp			
post solid organ transplant, renal dialysis and other sever 3) Up to five doses for children up to and under the age of 1	, , , , , , , , , , , , , , , , , , , ,	•	
Note: A course of up-to four vaccines is funded for catch up pi		, ,	
to complete full primary immunisation. Please refer to the Imm			
programmes.	iumsation nanuboor	tion the approprie	ate sofication for eaten up
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10 🗸 In	fanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]		_	
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)imm	nunisation for patien	ts post haematop	oietic stem cell
transplantation, or chemotherapy; functional asplenic; pre			
or post cochlear implants, renal dialysis and other severe			3
3) For use in testing for primary immunodeficiency diseases	, ,,	•	al medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg;			
prefilled syringe plus vial 0.5 ml	0.00	1 ✓ H	iberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
1) Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver dis-	ease; or		
3) One dose of vaccine for close contacts of known hepatitis	s A cases.		

✓ Havrix

✓ Havrix Junior

1

	NATIONAL	IIVIIVI	UNISATI	JN SCHEDOLL
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe		1	√ E	ngerix-B
Funded for patients meeting any of the following criteria 1) for household or sexual contacts of known acute h 2) for children born to mothers who are hepatitis B su 3) for children up to and under the age of 18 years in serology and require additional vaccination or requ 4) for HIV positive patients; or 5) for hepatitis C positive patients; or 6) for patients following non-consensual sexual interce 7) for patients following immunosuppression; or 8) for solid organ transplant patients; or 9) for post-haematopoietic stem cell transplant (HSC) 10) following needle stick injury.	repatitis B patients or harface antigen (HBsAg clusive who are considure a primary course course; or) posi dered	tive; or not to have	
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following criteria		1	√ <u>E</u>	ngerix-B
1) for household or sexual contacts of known acute he go for children born to mothers who are hepatitis B such for children up to and under the age of 18 years in serology and require additional vaccination or request for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual interest for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC) following needle stick injury; or for dialysis patients; or for liver or kidney transplant patients.	nepatitis B patients or hurface antigen (HBsAg clusive who are considure a primary course course; or) posi dered	tive; or not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 8 Any of the following: 1) Maximum of two doses for children aged 14 years and	under; or	- [Xph	arm]	
2) Maximum of three doses for patients meeting any of th 1) People aged 15 to 26 years inclusive; or 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: o	r			
3) Maximum of four doses for people aged 9 to 26 years	·			and all o
Inj 270 mcg in 0.5 ml syringe	0.00	10	✓ <u>G</u>	ardasil 9

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Sul	Fully osidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine)90.00

✓ Fluad Quad (2021 Formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

- INFLUENZA VACCINE people 65 years and over
 is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccin	e) –		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	√	Manufacturer	

A) INFLUENZA VACCINE – people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

MEASI ES, MUMPS AND BUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

${\sf MENINGOCOCCAL} \ ({\sf GROUPS} \ {\sf A}, \ {\sf C}, \ {\sf Y} \ {\sf AND} \ {\sf W-135}) \ {\sf CONJUGATE} \ {\sf VACCINE} \ -[{\sf Xpharm}]$

Fither:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases; or
 - 3) A maximum of two doses for bone marrow transplant patients; or
 - 4) A maximum of two doses for patients following immunosuppression*; or
- B) Both
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Ini 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

✓ Synflorix

10

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	\$	Per 🗸	Manufacturer
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]			
Both:			
1) The child is under 9 months of age; and			
Any of the following:			
 Up to three doses for patients pre- and post spler HIV, complement deficiency (acquired or inherited Two doses for close contacts of meningococcal c A maximum of two doses for bone marrow transp A maximum of two doses for patients pre- and po 	d), or pre or post solid ases; or lant patients; or	organ transplant	
Note: children under nine months of age require two d booster schedules with meningococcal ACWY vaccine.		Refer to the Imm	unisation Handbook for
*Immunosuppression due to steroid or other immunosu	ippressive therapy mu	ist be for a period	d of greater than 28 days.
Inj 10 mcg in 0.5 ml syringePNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm		1 ~ N	leisvac-C
A primary course of three doses for previously unvaccing	•	the age of 59 m	onths inclusive
Note: please refer to the Immunisation Handbook for the ap	propriate schedule for	catch up progra	mmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously
 received two doses of the primary course of PCV10: or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISAT	ION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE	– [Xpharm]		
Either:			
 Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with funcomplement deficiency (acquired or inherited), coch All of the following: 	ctional asplenia, pre- or p	oost-solid organ	transplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)immur b) Treatment is for a maximum of two doses; and c) Any of the following: 	· ·		
 i) on immunosuppressive therapy or radiati immune response; or 	ion therapy, vaccinate wl	nen there is expe	ected to be a sufficient
ii) with primary immune deficiencies; oriii) with HIV infection; or			
iv) with renal failure, or nephrotic syndrome;	or		
v) who are immune-suppressed following or		uding haematop	oietic stem cell transplant);
or vi) with cochlear implants or intracranial shu	inte: or		
vii) with cerebrospinal fluid leaks; or	into, or		
viii) receiving corticosteroid therapy for more			
prednisone of 2 mg/kg per day or greater	r, or children who weigh	more than 10 kg	on a total daily dosage of
20 mg or greater; or ix) with chronic pulmonary disease (includin	g asthma treated with hig	ah-dose corticos	teroid therapy): or
x) pre term infants, born before 28 weeks g		g., acco cocc	10.0.u 1o. up///, 0.
xi) with cardiac disease, with cyanosis or fai	lure; or		
xii) with diabetes; or			
xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	functional asplenia		
xiv) who are pre or poor opionocionity, or with	Turiotional aspionia.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each			
23 pneumococcal serotype)	0.00	1 🗸	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]			
Up to three doses for patients meeting either of the follow			
 For partially vaccinated or previously unvaccinated i For revaccination following immunosuppression. 	ndividuals; or		
Note: Please refer to the Immunisation Handbook for app	propriate schedule for ca	tch-up programn	100
Inj 80D antigen units in 0.5 ml syringe			POL
ROTAVIRUS ORAL VACCINE - [Xpharm]		-	
Maximum of two doses for patients meeting the following:			
 first dose to be administered in infants aged under 1 no vaccination being administered to children aged 2 			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

		Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
VARICELLA V Either:	ACCINE [CHICKENPOX VACCINE] - [Xpharm]				
1) Maxi	imum of one dose for primary vaccination for eithe	er:			
,	Any infant born on or after 1 April 2016; or For previously unvaccinated children turning 11 varicella infection (chickenpox), or	years old on or after 1	July 201	7, who h	nave not previously had a
	imum of two doses for any of the following:				
a)	Any of the following for non-immune patients:				
	ii) with chronic liver disease who may in future iii) with deteriorating renal function before tran iii) prior to solid organ transplant; or	splantation; or	nsplanta	ition; or	
	iv) prior to any elective immunosuppression*,				
L١	v) for post exposure prophylaxis who are imm			_:::	-liet
,	For patients at least 2 years after bone marrow to For patients at least 6 months after completion of				·
	For HIV positive non immune to varicella with mi				
	For patients with inborn errors of metabolism at r varicella, or				
,	For household contacts of paediatric patients wh immune compromise where the household conta For household contacts of adult patients who have	ct has no clinical histo	ry of var	ricella, or	
9/	immunocompromised, or undergoing a procedur has no clinical history of varicella.				
28 days	suppression due to steroid or other immunosuppre	.,	for a tre	eatment p	period of greater than
Inj 1350 P	FU prefilled syringe	0.00	1	✓ \	/arivax
			10	✓ \	<u>/arivax</u>
	OSTER VIRUS (OKA STRAIN) LIVE ATTENUAT r patients meeting either of the following criteria:	ED VACCINE [SHING	LES VA	CCINE]	– [Xpharm]
1) One	dose for all people aged 65 years; or				
	dose for all people aged between 66 and 80 year	s inclusive from 1 Apri	l 2018 a	nd 31 De	ecember 2021.
Ini 19.400	PFU prefilled syringe plus vial	0.00	1	√ 2	ostavax .
,,	- F		10		ostavax
Diagnosti	c Agents				
	PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU po	er 0.1 ml, 1 ml vial	0.00	1	✓]	<u>ubersol</u>

- Symbols -	Afluria Quad Junior	Amoxicillin with clavulanic acid9
UK Synacthen80	(2021 Formulation) 264	Amphotericin B3
3TC105	AFT-Pyrazinamide100	Amsacrine14
- A -	Agents Affecting the	AmsaLyo14
A-Scabies68	Renin-Angiotensin System 47	Amsidine14
Abacavir sulphate105	Agents for Parkinsonism and Related	Amzoate3
Abacavir sulphate with	Disorders 118	Anaesthetics11
lamivudine 105	Agents Used in the Treatment of	Anagrelide hydrochloride14
Abiraterone acetate162	Poisonings235	Analgesics12
Acarbose11	Agrylin149	Anastrozole16
Accarb11	Albendazole89	Anatrole16
Accuretic48	Albey220	Andriol Testocaps8
Accuretic 1048	Albustix77	Androderm8
Accuretic 2048	Aldurazyme28	ANI5
Acetazolamide232	Alecensa156	Anoro Ellipta22
Acetec47	Alectinib156	Antabuse14
Acetic acid with 1, 2- propanediol	Alendronate sodium111	Antacids and Antiflatulents
diacetate and	Alendronate sodium with	Anthelmintics8
benzethonium229	colecalciferol111	Antiacne Preparations6
Acetic acid with hydroxyquinoline and	Alfacalcidol33	Antiallergy Preparations22
ricinoleic acid75	Alfamino Junior256	Antianaemics3
Acetylcysteine235	Alginic acid6	Antiandrogen Oral
Aci-Jel75	Alglucosidase alfa26	Contraceptives7
Aciclovir	Alkeran146	Antiarrhythmics4
Infection100	Alkeran S29146	Antibacterials8
Sensory229	Allersoothe221	Antibacterials Topical6
Acidex6	Allmercap148	Anticholinergic Agents22
Acipimox55	Allopurinol116	Anticholinesterases11
Acitretin68	Alpha-Adrenoceptor Blockers47	Antidepressants12
Aclasta114	Alpha-Keri Lotion66	Antidiarrhoeals
Aclin110	Alphamox92	Antiepilepsy Drugs12
Actemra209	Alphamox 12592	Antifibrinolytics, Haemostatics and
Actinomycin D150	Alphamox 25092	Local Sclerosants3
Actrapid10	Alprolix38	Antifibrotics22
Actrapid Penfill10	Alu-Tab6	Antifungals9
Acular231	Aluminium hydroxide6	Antifungals Topical6
Acupan120	Alvogen51	Antihistamines22
Adalat 1052	Amantadine hydrochloride118	Antihypotensives5
Adalat Oros52	Ambrisentan58	Antimalarials9
Adalimumab173	Ambrisentan Mylan58	Antimigraine Preparations12
Adapalene61	Amiloride hydrochloride54	Antinausea and Vertigo Agents 13
Adcortyl80	Amiloride hydrochloride with	Antiparasitics9
Adefin52	furosemide54	Antipruritic Preparations6
Adefin XL52	Amiloride hydrochloride with	Antipsychotics13
Adenuric116	hydrochlorothiazide 54	Antiretrovirals10
ADR Cartridge 1.823	Aminophylline226	Antirheumatoid Agents11
Adrenaline57	Amiodarone hydrochloride49	Antispasmodics and Other Agents
Advantan64	Amisulpride131	Altering Gut Motility
Advate41	Amisulpride Mylan131	Antithrombotic Agents4
Adynovate41	Amitriptyline124	Antithymocyte globulin
Afinitor217	Amlodipine52	(equine) 17
Aflibercept183	Amneal153	Antitrichomonal Agents9
Afluria Quad	Amorolfine62	Antituberculotics and
(2021 Formulation) 265	Amoxicillin92	Antileprotics9

Antiulcerants	8	Arrow-Doxorubicin	150	Bacillus Calmette-Guerin (BCG)	
Antivirals	100	Arrow-Losartan &		vaccine	173
Anxiolytics	134	Hydrochlorothiazide	48	Bacillus Calmette-Guerin	
Anzatax	153	Arrow-Morphine LA	123	vaccine	26
Apidra	11	Arrow-Norfloxacin	109	Baclofen	11
Apidra SoloStar	11	Arrow-Ornidazole	99	Bactroban	62
Apo-Amlodipine	52	Arrow-Quinapril 10	48	Balance	29
Apo-Azithromycin		Arrow-Quinapril 20		Barrier Creams and Emollients	6
Apo-Bromocriptine		Arrow-Quinapril 5	48	Basic AquaCream	6
Apo-Ciclopirox	62	Arrow-Roxithromycin		BCG Vaccine	26
Apo-Clarithromycin		Arrow-Sertraline		Beclazone 100	22
Alimentary	9	Arrow-Timolol	231	Beclazone 250	22
Infection	90	Arrow-Topiramate	128	Beclazone 50	22
Apo-Clomipramine	124	Arrow-Tramadol	124	Beclomethasone dipropionate	22
Apo-Diclo SR		Arsenic trioxide	149	Bee venom allergy treatment	
Apo-Diltiazem CD		Asacol		Bendamustine hydrochloride	
Apo-Doxazosin		Asamax	7	Bendrofluazide	
Apo-Folic Acid		Ascorbic acid	33	Bendroflumethiazide	
Apo-Furosemide		Aspen Adrenaline		[Bendrofluazide]	5
Apo-Gabapentin		Aspirin		Benzathine benzylpenicillin	
Apo-Megestrol		Blood	41	Benzatropine mesylate	
Apo-Metoprolol		Nervous		Benzbromaron AL 100	
Apo-Mirtazapine		Asthalin		Benzbromarone	
Apo-Nadolol		Atazanavir sulphate		Benztrop	
Apo-Oxybutynin		Atenolol		Benzydamine hydrochloride	
Apo-Perindopril		Atenolol AFT		Benzylpenicillin sodium [Penicillin	
Apo-Pindolol		Atenolol AFT S29		G]	
Apo-Prazosin		ATGAM		Beta Cream	
Apo-Prednisone		Ativan		Beta Ointment	
Apo-Primidone		Atomoxetine		Beta Scalp	
Apo-Propranolol	51	Atorvastatin		Beta-Adrenoceptor Agonists	
Apo-Pyridoxine		Atropine sulphate		Beta-Adrenoceptor Blockers	
Apo-Selegiline S29		Cardiovascular	40	Betadine	
Apo-Sumatriptan		Sensory		Betadine Skin Prep	
Apo-Temozolomide		Atropt		Betaferon	
		Atrovent		Betahistine dihydrochloride	
Apo-Terazosin Apo-Timol		All Synacthen		Betaine	
Apomorphine hydrochloride		Aubagio		Betaloc CR	
Aprending		Augmentin		Betamethasone dipropionate	
Apresoline		AutoCoft 20		Betamethasone dipropionate with	
Aptamil AllerPro SYNEO 1		AutoSoft 30		calcipotriol	
Aptamil AllerPro SYNEO 2		AutoSoft 90		Betamethasone sodium phosphat	
Aptamil Gold+ Pepti Junior				with betamethasone acetate Betamethasone valerate	
Aqueous cream		Avonex		Betamethasone valerate with	. 03, 0
Arava		Azacitidine		clioquinol	
Arginine		Azacitidine Dr Reddy's		Betamethasone valerate with sod	
Aripiprazole		Azamun		fusidate [fusidic acid]	64
Aripiprazole 1A Pharma		Azathioprine		Betaxolol	23
Aripiprazole Sandoz		Azithromycin		Betnovate	
Aristocort		Azopt		Betnovate-C	
Arrow - Lattim		AZT	105	Betoptic	
Arrow-Amitriptyline		-B-		Betoptic S	
Arrow-Bendrofluazide		B-D Micro-Fine		Bezafibrate	
Arrow-Brimonidine		B-D Ultra Fine		Bezalip	5
Arrow-Diazepam	134	B-D Ultra Fine II	15	Bezalip Retard	5

Bicalutamide	163	Buprenorphine with naloxone	141	Carmustine	145
Bicillin LA	92	Bupropion hydrochloride		Carvedilol	50
BiCNU	145	Burinex	53	Carvedilol Sandoz	50
Bicnu Heritage	145	Burinex S29	53	Catapres	53
Bile and Liver Therapy		Buscopan	8	CeeNU	
Biltricide		Buspirone hydrochloride		Cefaclor monohydrate	
Bimatoprost		Busulfan		Cefalexin	
Bimatoprost Multichem		- C -		Cefalexin Sandoz	
Binarex		Cabergoline	88	Cefazolin	
Binocrit	38	Cacit		Ceftriaxone	
Biodone	122	Caffeine citrate	228	Ceftriaxone-AFT	89
Biodone Extra Forte	122	Calamine	63	Cefuroxime axetil	89
Biodone Forte	122	Calci-Tab 500	34	Celebrex	
Bisacodyl		Calcipotriol	68	Celecoxib	111
Bisoprolol fumarate		Calcitonin		Celecoxib Pfizer	111
Bisoprolol Mylan		Calcitriol		Celestone Chronodose	
BK Lotion		Calcitriol-AFT		Cellcept	
Bleomycin sulphate		Calcium carbonate		Centrally-Acting Agents	
Blood Colony-stimulating		Calcium Channel Blockers		Cephalexin ABM	
Factors	44	Calcium Disodium Versenate		Cetirizine hydrochloride	
Blood glucose diagnostic test		Calcium folinate		Cetomacrogol	
meter	14	Calcium Folinate Ebewe		Cetomacrogol with glycerol	
Blood glucose diagnostic test		Calcium Folinate Sandoz		Cetuximab	
strip	14	Calcium gluconate		Champix	
Blood glucose test strips (visua		Calcium Homeostasis		Charcoal	
impaired)		Calcium polystyrene sulphonate.		Chemotherapeutic Agents	
Blood Ketone Diagnostic Test		Calcium Resonium		Chickenpox vaccine	
Strip	13	Calogen		Chlorafast	
Bonjela		Calutide-50		Chlorambucil	
Boostrix		Candesartan cilexetil		Chloramphenicol	
Bortezomib		Candestar		Chlorothiazide	
Bortezomib Dr-Reddy's		Canesten		Chlorpheniramine maleate	
Bosentan		Capecitabine		Chlorpromazine hydrochloride	
Bosentan Dr Reddy's		Capercit		Chlortalidone [Chlorthalidone]	
Bplex		Capoten		Chlorthalidone	
Breo Ellipta		Capsaicin		Chlorvescent	
Brevinor 1/28		Musculoskeletal	111	Choice Load 375	
Bricanyl Turbuhaler		Nervous		Choice TT380 Short	
Brilinta		Captopril		Choice TT380 Standard	
Brimonidine tartrate		Captopril-Mylan		Choline salicylate with cetalkoniur	
Brimonidine tartrate with timolo		Carafate		chloride	
maleate		Carbaccord		Ciclopirox olamine	
Brinzolamide		Carbamazepine		Ciclosporin	
Bristol		Carbimazole		Cilazapril	
Brolene		Carbomer		Cilicaine	
Bromocriptine mesylate		Carboplatin		Cilicaine VK	
BSF Ambrisentan Mylan		Carboplatin Ebewe		Cinacalcet	
BSF Bisoprolol Mylan		Carbosorb-X		Cipflox	
BSF Darunavir Mylan	235	Cardinol LA		Ciprofloxacin	
Buccastem		CareSens Dual		Infection	93
Budesonide		CareSens N		Sensory	
Alimentary	6	CareSens N POP		Ciprofloxacin Teva	229
Respiratory	.221, 227	CareSens N Premier		Circadin	
Budesonide with eformoterol	222	CareSens PRO		Cisplatin	
Bumetanide		Carmellose sodium with gelatin a		Cisplatin Ebewe	
Buprenorphine Naloxone BNM		pectin		Citalopram hydrobromide	
- r r		r			

Cladribine 147	Combigan232	Dapsone9
Clarithromycin	Compound electrolytes46	Daraprim9
Alimentary9	Compound electrolytes with glucose	Darunavir10
Infection90	[Dextrose]46	Darunavir Mylan10
Clexane43	Compound hydroxybenzoate238	Dasatinib15
Clexane Forte43	Concerta139	Daunorubicin15
Climara81	Condoms72	David One Step Cassette Pregnancy
Climara81	Condyline70	Test7
Clindamycin94	Contraceptives - Hormonal73	DBL Acetylcysteine23
Clinicians27	Contraceptives - Non-hormonal71	DBL Adrenaline5
Clinicians Renal Vit33	Copaxone 135	DBL Aminophylline22
Clobazam 126	Corticosteroids and Related Agents	DBL Bleomycin Sulfate14
Clobetasol propionate63, 69	for Systemic Use79	DBL Carboplatin14
Clobetasone butyrate64	Corticosteroids Topical63	DBL Cisplatin14
Clofazimine99	Cosentyx206	DBL Dacarbazine15
Clomazol	Cosmegen 150	DBL Desferrioxamine Mesylate for Inj
Dermatological62	Coumadin44	BP23
Genito-Urinary75	Country Life29	DBL Docetaxel15
Clomifene citrate88	Coversyl48	DBL Ergometrine7
Clomipramine hydrochloride124	Creon 1000024	DBL Gemcitabine14
Clonazepam 126, 134	Creon 2500024	DBL Gentamicin9
Clonidine53	Creon Micro24	DBL Heparin Sodium4
Clonidine BNM53	Crotamiton63	DBL Leucovorin Calcium14
Clonidine hydrochloride53	Crystaderm61	DBL Methotrexate Onco-Vial14
Clopidogrel41	Curam92	DBL Morphine Sulphate12
Clopidogrel Multichem41	Curam Duo 500/12592	DBL Naloxone Hydrochloride23
Clopine131	Cvite33	DBL Octreotide
Clopixol	Cyclizine hydrochloride130	DBL Pethidine Hydrochloride 12
Clotrimazole	Cyclizine lactate130	DBL Vinblastine
Dermatological62	Cyclogyl233	DBL Vincristine Sulfate15
Genito-Urinary75	Cyclopentolate hydrochloride 233	Decozol3
Clozapine131	Cyclophosphamide145	Deferasirox23
Clozaril131	Cyclorin99	Deferiprone23
Co-trimoxazole95	Cycloserine99	Denosumab11
Coal tar68	Cyproterone acetate80	Deolate9
Coal tar with allantoin, menthol,	Cyproterone acetate with	Deoxycoformycin15
phenol and sulphur69	ethinyloestradiol75	Depo-Medrol7
Coal tar with salicylic acid and	Cystadane27	Depo-Provera7
sulphur69	Cytarabine147	Depo-Testosterone8
Coco-Scalp69	Cytotec8	Deprim9
Codeine phosphate	Cytoxan145	Dermol
Extemporaneous238	- D -	Desferrioxamine mesilate23
Nervous121	D-Penamine111	Desmopressin8
Coenzyme Q1027	Dabigatran44	Desmopressin acetate8
Colchicine116	Dacarbazine150	Desmopressin-PH&T8
Colecalciferol33	Dacarbazine APP150	Desuric11
Colestid55	Dactinomycin [Actinomycin D]150	Detection of Substances in
Colestipol hydrochloride55	Daivobet68	Urine7
Colgout 116	Daivonex68	Dexamethasone
Colifoam7	Daktarin63	Hormone7
Colistin sulphomethate94	Dalacin C94	Sensory23
Colistin-Link94	Dantrium117	Dexamethasone phosphate7
Collodion flexible238	Dantrium S29117	Dexamethasone Phosphate
Colloidal bismuth subcitrate10	Dantrolene117	Panpharma7
Colofac8	Daonil11	Dexamethasone with framycetin and
Coloxyl25	Dapa-Tabs <u>55</u>	gramicidin 22
•	· · · · · · · · · · · · · · · · · · ·	-

sennosides	. 25		
Dolutegravir	106		
Donepezil hydrochloride	140	Eltrombopag	38
Donepezil-Rex	140		
Dornase alfa	226		
Dorzolamide hydrochloride	232		
Dostinex	88		12
Dosulepin [Dothiepin]		Empagliflozin with metformin	
		Emtricitabine	105
Dothiepin	124	Emtricitabine with tenofovir	
Doxazosin	47	disoproxil	102
Doxine	93	Emtriva	105
Doxorubicin Ebewe	150		
Doxorubicin hydrochloride	150	Emulsifying Ointment ADE	65
Doxycycline	93		
DP Lotn HC	64		
DP-Allopurinol	116		
		Engerix-B	263
Drugs Affecting Bone		Enlafax XR	126
Metabolism	111	Enoxaparin sodium	43
		Enstilar	68
diagnostic test meter	. 13	Ensure	250
		Ensure Plus	251
Duolin	223	Ensure Plus HN	250
Duolin HFA	223	Ensure Plus RTH	250
DuoResp Spiromax	222	Entacapone	118
Duride	. 57		
-E-		Entecavir	100
e-chamber La Grande	228		
e-chamber Mask	227	Entocort CIR	6
e-chamber Turbo	228	Entresto 24/26	48
E-Mycin	91	Entresto 49/51	48
Ear Preparations	229		
		Epilim	128
Econazole nitrate	62		
Efavirenz	105	Epilim S/F Liquid	128
Efavirenz with emtricitabine and			
tenofovir disoproxil	105		
Eformoterol fumarate	222		
Eformoterol fumarate dihydrate	222		
Eftrenonacog alfa [Recombinant		Epoetin alfa	38
	. 38	Epoprostenol	60
		VIIa]	40
		Erbitux	184
	Dolutegravir	hydrochloride 124 Dosulepin Mylan 124 Dothiepin 124 Doxazosin 47 Doxine 93 Doxorubicin Ebewe 150 Doxorubicin hydrochloride 150 Doxycycline 93 DD Lotion 66 DP Lotion HC 64 DP-Allopurinol 116 Dr Reddy's Omeprazole 9 Drugs Affecting Bone 9 Metabolism 111 Dual blood glucose and blood ketone diagnostic test meter 13 Ducal Super Soluble Powder 241 Duolin 223 Duolin HFA 223 Duolin HFA 223 Duoride 57 - E - e-chamber La Grande 228 e-chamber Mask 227 e-chamber Turbo 228 E-Mycin 91 Ear Preparations 229 Earlyee Preparations 229 Easiphen Liquid 255 Econazole nitrat	Dolutegravir

Erythromycin ethyl succinate	91	Felodipine	52	Fluoxetine hydrochloride	12
Erythromycin stearate	91	Femme-Tab ED	74	Flupenthixol decanoate	132
Esbriet	225	Fentanyl	122	Flutamide	163
Escitalopram	125	Fentanyl Sandoz		Flutamin	160
Escitalopram (Ethics)		Ferinject		Fluticasone	222
Escitalopram-Apotex		Ferodan		Fluticasone furoate with	
Eskazole		Ferric carboxymaltose	34	vilanterol	222
Essential Prednisolone	8	Ferriprox	236	Fluticasone propionate	22
Estradiol TDP Mylan	81	Ferro-F-Tabs	35	Fluticasone with salmeterol	223
Estradot		Ferro-tab	35	FML	23 ⁻
Estradot 50 mcg	81	Ferrograd	35	Foban	62
Estrofem	81	Ferrosig	35	Folic acid	38
Etanercept	166	Ferrous fumarate	35	Food Thickeners	252
Ethambutol hydrochloride	99	Ferrous fumarate with folic acid.	35	Foods And Supplements For Int	orn
Ethics Aspirin	120	Ferrous sulfate	35	Errors Of Metabolism	254
Ethics Aspirin EC		Ferrous sulphate	35	Foradil	22
Ethics Lisinopril		Fexofenadine hydrochloride	221	Forteo	113
Ethics Paracetamol Classic		Fibro-vein	41	Fortini	24
Ethinyloestradiol	82	Filgrastim		Fortini Multi Fibre	
Ethinyloestradiol with		Finasteride	76	Fortisip	250-25 ⁻
desogestrel	73	Fingolimod	135	Fortisip Multi Fibre	
Ethinyloestradiol with		Firazyr		Fosamax	11
levonorgestrel	74	Flagyl	98	Fosamax Plus	
Ethinyloestradiol with		Flagyl-S		Framycetin sulphate	229
norethisterone	74	Flamazine		Frisium	
Ethosuximide	127	Flecainide acetate	49	Frumil	54
Etopophos		Flecainide BNM	49	Frusemide	54
Etoposide		Flecainide Controlled Release		Fucicort	
Etoposide phosphate		Teva	49	Fucidin	
Etravirine		Fleet Phosphate Enema	26	Fucithalmic	229
Eumovate		Flixonase Hayfever & Allergy		Fulvestrant	163
Everet	127	Flixotide		Fungilin	
Everolimus		Flixotide Accuhaler		Furosemid-Ratiopharm	
Evista		Florinef		Furosemide [Frusemide]	
Exemestane		Fluad Quad		Furosemide-Baxter	
Exjade		(2021 Formulation)	265	fusidic acid	•
Extemporaneously Compounded		Fluanxol		Dermatological	62. 64
Preparations and	-	Flucil		Infection	
Galenicals	238	Flucloxacillin		Sensory	
Eye Preparations		Flucloxin		- G -	
Eylea		Flucon		Gabapentin	12
Ezetimibe		Fluconazole		Gacet	12
Ezetimibe Sandoz		Fludara Oral		Galsulfase	
Ezetimibe with simvastatin		Fludarabine Ebewe		Galvumet	
- F -		Fludarabine phosphate		Galvus	
Factor eight inhibitor bypassing		Fludrocortisone acetate		Gardasil 9	
fraction	40	Fluids and Electrolytes		Gastrodenol	
Famotidine		Flumetasone pivalate		Gaviscon Double Strength	
Famotidine Hovid		Fluocortolone caproate with		Gaviscon Infant	
Faslodex		fluocortolone pivalate and		Gazyva	19
Febuxostat		cinchocaine	8	Gefitinib	
Feed Thickener Karicare		Fluorometholone		Gemcitabine Ebewe	
Aptamil	253	Fluorouracil		Gemcitabine hydrochloride	
FEIBA NF		Fluorouracil Ebewe		Genoptic	
Felo 10 ER		Fluorouracil sodium		Gentamicin sulphate	
Felo 5 ER		Fluox		Infection	Q
	· · · · · · · · · · · · · · · · · · ·				· · · · · · · · · · · · · · · · · · ·

Sensory229	Hemastix77	Hygroton5
Gilenya135	Hemlibra39	Hylo-Fresh23
Ginet75	Heparin sodium43	Hymenoptera22
Glatiramer acetate 135	Heparinised saline43	Hyoscine butylbromide
Glecaprevir with pibrentasvir 102	Heparon Junior244	Hyoscine hydrobromide13
Glibenclamide11	Hepatitis A vaccine262	Hypam13
Gliclazide12	Hepatitis B recombinant	Hyperuricaemia and Antigout11
Glipizide12	vaccine263	Hypromellose23
Glivec158	Herceptin212	Hypromellose with dextran23
Glizide12	Hiberix262	-1-
Glucagen Hypokit10	Hiprex109	Ibiamox9
Glucagon hydrochloride10	Histaclear221	Ibrance15
Glucerna Select243	Histafen221	Ibuprofen11
Glucerna Select RTH243	Holoxan145	Ibuprofen SR BNM11
Glucobay11	Horleys Bread Mix253	Icatibant22
Glucose [Dextrose]45	Horleys Flour253	Idarubicin hydrochloride15
Gluten Free Foods253	Hormone Replacement Therapy -	Idursulfase2
Glycerin with sodium saccharin238	Systemic 80	Ifosfamide14
Glycerin with sucrose238	HPV263	Igroton5
Glycerol	Humalog11	Ikorel5
Alimentary26	Humalog Mix 2511	Ilevro23
Extemporaneous238	Humalog Mix 5011	Iloprost6
Glyceryl trinitrate	Human papillomavirus (6, 11, 16, 18,	Imatinib mesilate15
Alimentary8	31, 33, 45, 52 and 58) vaccine	Imatinib-AFT15
Cardiovascular57	[HPV]263	Imatinib-Rex15
Glycopyrronium224	Humatin95	Imigran12
Glycopyrronium bromide8	Humira173	Imipramine hydrochloride12
Glycopyrronium with	HumiraPen173	Imiquimod7
indacaterol224	Humulin 30/7011	Immune Modulators10
Go Healthy27	Humulin NPH11	Immunosuppressants16
Gold Knight72	Humulin R10	Imuran16
Gold Knight XL72–73	Hyaluronic acid233	Incruse Ellipta22
Goserelin87	Hydralazine57	Indacaterol22
Gutron50	Hydralazine hydrochloride57	Indapamide5
Gynaecological Anti-infectives75	Hydrocortisone	Infanrix IPV26
- H -	Dermatological64	Infanrix-hexa26
Habitrol142	Hormone79	Infant Formulae25
Haemophilus influenzae type B	Hydrocortisone (PSM)64	Infatrini26
vaccine 262	Hydrocortisone acetate7	Infliximab18
Haldol133	Hydrocortisone acetate with	Influenza vaccine26
Haldol Concentrate133	pramoxine hydrochloride 7	Influvac Tetra
Haldol Decanoas133	Hydrocortisone and paraffin liquid	(2021 Formulation) 26
Haloperidol131	and lanolin64	Inhaled Corticosteroids22
Haloperidol decanoate133	Hydrocortisone butyrate64, 69	Inhaled Long-acting
Harvoni102	Hydrocortisone with cinchocaine8	Beta-adrenoceptor Agonists 22
Havrix262	Hydrocortisone with miconazole64	Inspra5
Havrix Junior262	Hydrocortisone with natamycin and	Instillagel Lido11
Haylor syrup221	neomycin 65	Insulin aspart1
healthE Calamine Aqueous Cream	Hydrogen peroxide61	Insulin aspart with insulin aspart
BP63	Hydroxocobalamin33	protamine1
healthE Dimethicone 10%65	Hydroxocobalamin Mercury	Insulin glargine1
healthE Dimethicone 4% Lotion 66	Pharma33	Insulin glulisine1
healthE Dimethicone 5%65	hydroxycarbamide151	Insulin isophane1
healthE Glycerol BP238	Hydroxychloroquine111	Insulin isophane with insulin
healthE Urea Cream65	Hydroxyurea	neutral1
Healtheries Simple Baking Mix253	[hydroxycarbamide] 151	Insulin lispro1

Insulin lispro with insulin lispro	Ivacaftor	227	Lantus SoloStar	
protamine11	Ivermectin	66	Lanvis	149
Insulin neutral10	- J -		Lanzol Relief	
Insulin pen needles15	Jadelle	74	Lapatinib ditosylate	158
Insulin pump15	Jakavi	161	Largactil	13 ⁻
Insulin pump cartridge20		13	Laronidase	28
Insulin pump infusion set (steel	Jardiance	12	Lasix	54
cannula)20	Jaydess	82	Latanoprost	232
Insulin pump infusion set (steel	Jevity HiCal RTH		Latanoprost with timolol	
cannula, straight insertion) 21			Lax-Suppositories	
Insulin pump infusion set (teflon	Juno Pemetrexed		Lax-Tab	
cannula)22			Laxatives	
Insulin pump infusion set (teflon	Kadcyla	213	Laxsol	2
cannula, angle insertion with	Kaletra		Ledipasvir with sofosbuvir	
insertion device)23			Leflunomide	
Insulin pump infusion set (teflon	Kemadrin		Lenalidomide	15
cannula, angle insertion) 23			Letrole	
Insulin pump infusion set (teflon	Kenacort-A 10		Letrozole	
cannula, straight insertion with	Kenacort-A 40		Leukeran FC	
insertion device)23			Leukotriene Receptor	
Insulin pump infusion set (teflon	Kenalog in Orabase		Antagonists	226
cannula, straight insertion) 23			Leuprorelin	
Insulin pump reservoir23			Leustatin	
	KetoCal 4:1		Levetiracetam	
Insulin syringes, disposable with attached needle		200	Levetiracetam-AFT	
Intal Forte CFC Free		60	Levlen ED	
	•			
Intelence 105			Levocabastine	
Interferon beta-1-alpha			Levocarnitine	
Interferon beta-1-beta136			Levodopa with benserazide	
Intra-uterine device			Levodopa with carbidopa	118
Invega Sustenna133			Levomepromazine	132
IPOL271	•		Levomepromazine	
lpratropium bromide223, 227			hydrochloride	132
Iressa157			Levonorgestrel	
Irinotecan Accord147			Genito-Urinary	
Irinotecan Actavis 100147			Hormone	
Irinotecan hydrochloride147			Levothyroxine	
Irinotecan-Rex147	· ·		Lidocaine [Lignocaine]	.119–120
Iron polymaltose35			Lidocaine [Lignocaine]	
Isentress106	-		hydrochloride	120
Isentress HD106		29	Lidocaine [Lignocaine] with	
Ismo 2057			chlorhexidine	120
Ismo 40 Retard57	Labetalol	51	Lidocaine [Lignocaine] with	
Isoniazid99		127	prilocaine	
Isoniazid with rifampicin99			Lidocaine-Claris	120
Isoptin53	Laevolac	26	Life Extension	
Isoptin Retard53	Lamictal	127	Lignocaine	.119-120
Isoptin SR53	Lamivudine	100, 105	Lioresal Intrathecal	117
Isopto Carpine232	Lamivudine Alphapharm	105	Lipid-Modifying Agents	5
Isosorbide mononitrate57			Liquigen	242
Isosource Standard250			Lisinopril	48
Isotretinoin61			Lithium carbonate	
Ispaghula (psyllium) husk25	Lanoxin PG	49	Livostin	
Itch-Soothe63			LMX4	
Itraconazole96			Locacorten-Viaform ED's	
Itrazole96	•		Local preparations for Anal and	
			1 1	

Rectal Disorders	8	Maviret	102	Methylphenidate hydrochloride	
Locasol	256	Maxidex	230	extended-release	139
Locoid	64, 69	Maxitrol		Methylprednisolone	79
Locoid Crelo	64	MCT oil (Nutricia)	242	Methylprednisolone (as sodium	
Locoid Lipocream	64	Measles, mumps and rubella		succinate)	79
Locorten-Vioform	229	vaccine	268	Methylprednisolone aceponate	64
Lodoxamide	231	Mebendazole	89	Methylprednisolone acetate	79
Logem	127	Mebeverine hydrochloride	8	Methylxanthines	226
Lomide	231	Medco		Metoclopramide Actavis 10	130
Lomustine	146	Dermatological	65	Metoclopramide hydrochloride	130
Loniten	57	Nervous	121	Metolazone	
Loperamide hydrochloride	6	Medrol	79	Metopirone	88
Lopinavir with ritonavir	106	Medroxyprogesterone acetate		Metoprolol IV Mylan	51
Loprofin		Genito-Urinary	75	Metoprolol succinate	
Loprofin Mix		Hormone		Metoprolol tartrate	
Lorafix		Mefenamic acid	110	Metrogyl	
Loratadine		Megestrol acetate	164	Metronidazole	
Lorazepam	134	Melatonin		Metyrapone	88
Lorfast		Melphalan		Mexiletine hydrochloride	
Lorstat		Menactra		Mexiletine Hydrochloride USP	
Losartan Actavis	48	Meningococcal (groups A, C, Y an		Miacalcic	
Losartan potassium	48	W-135) conjugate vaccine		Micolette	
Losartan potassium with		Meningococcal C conjugate		Miconazole	
hydrochlorothiazide	48	vaccine	269	Miconazole nitrate	
Lovir		Menthol		Dermatological	63
Loxamine		Mepolizumab		Genito-Urinary	
Lucrin Depot 1-month		Mercaptopurine		Micreme	
Lucrin Depot 3-month		Mercilon 28		Micreme H	
Ludiomil		Mesalazine		Microgynon 20 ED	
Lyderm		Mesna		Microgynon 30	
Lynparza		Mestinon		Microgynon 50 ED	
Lyrica		Metabolic Disorder Agents		Microlut	
- M -		Metformin hydrochloride		Midazolam	
m-Eslon	123	Methadone hydrochloride		Midazolam-Baxter	
Mabthera		Extemporaneous	238	Midodrine	
Macrobid		Nervous		Mifegyne	
Macrogol 3350 with potassium		Methatabs		Mifepristone	
chloride, sodium bicarbonate	and	Methenamine (hexamine)		Minerals	
sodium chloride		hippurate	109	Mini-Wright AFS Low Range	
Macrogol 400 and propylene		Methopt		Mini-Wright Standard	
glycol	233	Methotrexate		Minidiab	
Madopar 125		Methotrexate DBL Onco-Vial		MiniMed 1.8 Reservoir	
Madopar 250		Methotrexate Ebewe		MMT-326A	23
Madopar 62.5		Methotrexate Sandoz		MiniMed 3.0 Reservoir	20
Madopar HBS		Methyl hydroxybenzoate		MMT-332A	23
Madopar Rapid		Methylcellulose		MiniMed 640G	
Magnesium hydroxide		Methylcellulose with glycerin and	200	MiniMed Mio MMT-921A	
Magnesium sulphate		sodium saccharin	238	MiniMed Mio MMT-923A	
Mantoux		Methylcellulose with glycerin and	200	MiniMed Mio MMT-925A	
Maprotiline hydrochloride		SUCTOSE	238	MiniMed Mio MMT-941A	
Marevan		Methyldopa		MiniMed Mio MMT-943A	
Marine Blue Lotion SPF 50+		Methyldopa Mylan		MiniMed Mio MMT-945A	
Martindale Pharma		Methyldopa Mylan S29	50	MiniMed Mio MMT-965A	
Marvelon 28		Methylnaltrexone bromide		MiniMed Mio MMT-975A	
Mask for spacer device		Methylphenidate ER - Teva		MiniMed Quick-Set MMT-386A	
Mast Cell Stabilisers		Methylphenidate hydrochloride		MiniMed Quick-Set MMT-387A	
made dell'olabilidolo		month of the state		TITLE CONTROL COLLINIAL TOTAL	

MiniMed Quick-Set MMT-396A.	22	Mucolytics	226	Nepro HP RTH	24
MiniMed Quick-Set MMT-397A.		Mucosoothe		Nerisone	6
MiniMed Quick-Set MMT-398A.		Multiple Sclerosis Treatments.		Neulactil	
MiniMed Quick-Set MMT-399A.	22	Multivitamin renal	33	Neulastim	4
MiniMed Silhouette MMT-368A.		Multivitamins		NeuroTabs	
MiniMed Silhouette MMT-377A.		Mupirocin		Nevirapine	
MiniMed Silhouette MMT-378A.		Muscle Relaxants		Nevirapine Alphapharm	
MiniMed Silhouette MMT-381A.		Mvite		Nicorandil	5
MiniMed Silhouette MMT-382A.		Myambutol		Nicotine	
MiniMed Silhouette MMT-383A.		Mycobutin		Nifedipine	
MiniMed Silhouette MMT-384A.		MycoNail		Nifuran	
MiniMed Sure-T MMT-864A		Mycophenolate mofetil		Nilotinib	
MiniMed Sure-T MMT-866A		Mydriacyl		Nilstat	
MiniMed Sure-T MMT-874A		Mylan Atenolol		Alimentary	3
MiniMed Sure-T MMT-876A		Mylan Clomiphen		Genito-Urinary	
MiniMed Sure-T MMT-884A		Mylan Midazolam		Infection	
MiniMed Sure-T MMT-886A		Myleran		Nintedanib	
Minims Cyclopentolate		Myometrial and Vaginal Hormo		Nipent	
Minims Pilocarpine		Preparations		Nitrates	
Minims Prednisolone		Myozyme		Nitroderm TTS	
Minirin		Mysoline S29	128	Nitrofurantoin	
Minirin Melt		- N -		Nitrolingual Pump Spray	
Mino-tabs		Nadolol		Nivestim	
Minocycline hydrochloride		Naglazyme		Nivolumab	
Minomycin		Nalcrom		Nizoral	
Minor Skin Infections		Naloxone hydrochloride		Nodia	
Minoxidil		Naltraccord		Noflam 250	
Mirena		Naltrexone hydrochloride		Noflam 500	11
Mirtazapine		Naphazoline hydrochloride		Non-Steroidal Anti-Inflammatory	
Misoprostol		Naphcon Forte		Drugs	11
Mitomycin C	152	Naprosyn SR 1000	110	Nonacog gamma, [Recombinant	
Mitozantrone	152	Naprosyn SR 750	110	Factor IX]	4
Mitozantrone Ebewe	152	Naproxen	110	Norethisterone	
Mixtard 30	11	Narcaricin mite	116	Genito-Urinary	7
MMR II	268	Nasal Preparations	227	Hormone	8
Moclobemide	125	Natalizumab		Norflex	11
Modafinil	140	Natulan	153	Norfloxacin	10
Modavigil	140	Nausafix	130	Noriday 28	7
Moduretic		Nausicalm	130	Norimin	
Molaxole	26	Navelbine	156	Normacol Plus	2
Moments	72	Necon	74	Normison	
Mometasone furoate		Nedocromil		Norpress	
Monogen		Nefopam hydrochloride	120	Nortriptyline hydrochloride	
Montelukast		Neisvac-C		Norvir	
Montelukast Mylan		Neo-B12		NovaSource Renal	
Moroctocog alfa [Recombinant fa		Neo-Mercazole		Novatretin	
VIII]		Neo-Mercazole S29		NovoMix 30 FlexPen	
Morphine hydrochloride		Neocate Gold		NovoRapid	
Morphine sulphate		Neocate Junior Unflavoured		NovoRapid FlexPen	
Motetis		Neocate Junior Vanilla		NovoRapid Penfill	
Mouth and Throat		Neocate SYNEO		NovoSeven RT	
Movapo		Neoral		Noxafil	
Moxifloxacin		Neostigmine metilsulfate		Nozinan	
MSUD Maxamum				Nozinan (Swiss)	ران م
	254	Nepafenac	231		
Mucilaginous laxatives with	0.5	Nepro HP (strawberry)		Nucala	
stimulants	25	Nepro HP (vanilla)	246	Nuelin	22

Nuelin-SR226	Omalizumab193	Pacifen	11
Nutilis253	Omeprazole9	Paclitaxel	
Nutren Diabetes243	Omeprazole actavis 109	Paclitaxel Actavis	
Nutrient Modules240	Omeprazole actavis 209	Paclitaxel Ebewe	15
Nutrini Energy Multi Fibre245	Omeprazole actavis 409	Paediatric Seravit	
Nutrini Energy RTH245	Omnitrope83	Palbociclib	15
Nutrini Low Energy Multi Fibre247	Onbrez Breezhaler222	Paliperidone	
Nutrini Peptisorb258	Oncaspar LYO153	Pamidronate disodium	112
Nutrini Peptisorb Energy258	OncoTICE173	Pamisol	
Nutrini RTH245	Ondansetron130	Panadol Mini Caps	12
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA130	Pancreatic enzyme	
Fibre 250	One-Alpha33	Pantoprazole	10
Nutrison Concentrated252	Onrex130	Panzop Relief	10
Nutrison Energy249	Opdivo214	Panzytrat	2
Nutrison Energy Multi Fibre250	Ora-Blend238	Papaverine hydrochloride	5
Nutrison Multi Fibre250	Ora-Blend SF238	Para-amino salicylic acid	
Nutrison Standard RTH250	Ora-Plus238	Paracare	12
Nyefax Retard52	Ora-Sweet238	Paracare Double Strength	12
Nystatin	Ora-Sweet SF238	Paracetamol	12
Alimentary32	Orabase32	Paracetamol + Codeine	
Genito-Urinary75	Oral and Enteral Feeds243	(Relieve)	124
Infection97	Oratane61	Paracetamol Pharmacare	12
NZB Low Gluten Bread Mix253	Ordine122	Paracetamol with codeine	124
-0-	Orgran254	Paraffin	65-6
O/W Fatty Emulsion Cream65	Ornidazole99	Paraffin liquid with wool fat	23
Obinutuzumab193	Orphenadrine citrate117	Paraldehyde	12
Obstetric Preparations77	Ortho-tolidine77	Parasidose	6
Ocicure9	Oruvail SR110	Parasiticidal Preparations	6
Ocrelizumab136	Osmolite RTH250	Parnate	12
Ocrevus136	Other Endocrine Agents88	Parnate S29	12
Octocog alfa [Recombinant factor	Other Oestrogen Preparations82	Paromomycin	9
VIII] (Advate)41	Other Progestogen	Paroxetine	12
Octocog alfa [Recombinant factor	Preparations82	Paser	9
VIII] (Kogenate FS)41	Other Skin Preparations70	Paxam	13
Octreotide	Ovestin	Paxtine	12
Octreotide (Sun) 164	Genito-Urinary75	Pazopanib	160
Octreotide GH164	Hormone82	Peak flow meter	
Octreotide LAR (somatostatin	Ox-Pam134	Pedialyte - Bubblegum	4
analogue) 164	Oxaliplatin146	Pediasure	24
Octreotide MaxRx164	Oxaliplatin Accord146	Pediasure RTH	24
Oestradiol81	Oxaliplatin Actavis 100146	Pegaspargase	15
Oestradiol valerate81	Oxaliplatin Ebewe146	Pegasys	
Oestradiol with norethisterone82	Oxazepam 134	Pegfilgrastim	4
Oestriol	Oxis Turbuhaler222	Pegylated interferon alfa-2a	10
Genito-Urinary75	Oxpentifylline57	Pembrolizumab	
Hormone82	Oxybutynin	Pemetrexed	14
Oestrogens81	Oxycodone hydrochloride123	Penicillamine	
Ofev224	Oxycodone Sandoz123	Penicillin G	9
Oil in water emulsion65	Oxycodone Sandoz S29123	PenMix 30	1
Olanzapine 132–133	OxyNorm123	PenMix 40	1
Olaparib	Oxytocin	PenMix 50	
Olbetam55	Oxytocin BNM75	Pentasa	
Olbetam S2955	Oxytocin with ergometrine	Pentostatin [Deoxycoformycin]	
Olopatadine233	maleate	Pentoxifylline [Oxpentifylline]	
Olopatadine Teva233	Ozurdex230	Peptamen Junior	
Olsalazine7	- P -	Peptisoothe	

Peptisorb	246	vaccine	270	Proglycem	1
Perhexiline maleate		Pneumococcal (PPV23)		Progynova	
Pericyazine		polysaccharide vaccine	271	Prolia	
Perindopril		Pneumovax 23		Promethazine hydrochloride	
Perjeta	194	Podophyllotoxin		Propafenone hydrochloride	
Permethrin		Polaramine		Propamidine isethionate	
Perrigo		Poliomyelitis vaccine		Propranolol	
Pertuzumab		Poloxamer		Propylene glycol	
Peteha		Poly-Gel		Propylthiouracil	
Pethidine hydrochloride		Poly-Tears		Protaphane	
Pevaryl		Poly-Visc		Protaphane Penfill	
Pexsig		Polycal		Protifar	
Pfizer Exemestane		Ponstan		Protionamide	
Pharmacy Services		Posaconazole		Provera	
Pheburane		Postinor-1		Provera HD	
Phenasen		Potassium chloride		PSM Citalopram	
Phenobarbitone		Potassium Chloride Aguettant		Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate		Preparations	6
Extemporaneous	238	Potassium iodate		PTU	
Nervous		Povidone iodine		Pulmicort Turbuhaler	
Phenothrin		Pradaxa		Pulmozyme	
Phenoxybenzamine	00	Pramipexole hydrochloride		Puri-nethol	
hydrochloride	47			Puria	
		Prayastatin		Pyrazinamide	
Phenoxymethylpenicillin (Penicil		Pravastatin Mylan			
V)		Praziquantel		Pyridostigmine bromide	
		Prazosin Pred Forte		Pyridoxine hydrochloride	
Phillips Milk of Magnesia				Pyrimethamine	
Phlexy 10		Prednisolone Prednisolone acetate		Pytazen SR	4
Phosphate Phebra	40				0
Phosphorus		Prednisolone sodium	0	Q 300	
Phytomenadione		Prednisolone sodium	001	Quetapel	
Pilocarpine hydrochloride		phosphate		Quetiapine	13
Pimafucort		Prednisolone-AFT		Quick-Set MMT-392	
Pimecrolimus		Prednisone		Quick-Set MMT-393	
Pindolol		Pregabalin		Quinapril	4
Pine tar with trolamine laurilsulfa		Pregabalin Pfizer		Quinapril with	
and fluorescein		Pregnancy Tests - hCG Urine		hydrochlorothiazide	
Pinetarsol		Premarin		Quinine sulphate	
Pioglitazone		Prevenar 13		Qvar	22
Pirfenidone		Priadel		-R-	
Pizotifen		Primacin		RA-Morph	
PKU Anamix Infant		Primaquine		Raloxifene hydrochloride	
PKU Anamix Junior		Primidone		Raltegravir potassium	
PKU Anamix Junior Chocolate		Primolut N		Ramipex	
PKU Anamix Junior LQ		Priorix		Ranbaxy-Cefaclor	
PKU Anamix Junior Orange		Probenecid		Ranitidine	
PKU Anamix Junior Vanilla		Probenecid-AFT		Rapamune	
PKU Lophlex LQ 10		Procaine penicillin		Reandron 1000	
PKU Lophlex LQ 20		Procarbazine hydrochloride		Recombinant factor IX	
PKU Lophlex Powder		Prochlorperazine		Recombinant factor VIIa	
PKU Lophlex Sensation 20		Proctofoam		Recombinant factor VIII	
Plaquenil		Proctosedyl		Rectogesic	
Plendil ER		Procyclidine hydrochloride		Redipred	
Pneumococcal (PCV10) conjuga	ate	Procytox		Relieve	
vaccine					
		Progesterone		Relistor	
Pneumococcal (PCV13) conjuga		Progesterone Proglicem		Relistor Remicade	

Renilon 7.5		Ruxolitinib		Sodium acid phosphate	
Resonium-A		Rythmodan		Sodium alginate	
Resource Beneprotein		Rytmonorm	50	Sodium benzoate	30
Respigen		- S -		Sodium bicarbonate	
Respiratory Devices		Sabril		Blood	. 45–46
Respiratory Stimulants	228	Sacubitril with valsartan	48	Extemporaneous	238
Retinol palmitate	<mark>234</mark>	SalAir	223	Sodium calcium edetate	236
ReTrieve	61	Salazopyrin	8	Sodium chloride	
Retrovir	105	Salazopyrin EN	8	Blood	45
Revlimid	151	Salbutamol	223	Respiratory	227
Revolade	38	Salbutamol with ipratropium		Sodium citrate with sodium lauryl	
Rexacrom	231	bromide	223	sulphoacetate	26
Riboflavin	29	Salicylic acid	69	Sodium citro-tartrate	
Ribomustin	144	Salmeterol		Sodium cromoglicate	
Ricit	76	Sandomigran	129	Alimentary	8
Rifabutin	100	Sandostatin LAR		Respiratory	
Rifadin	100	Sanofi Primaquine		Sensory	
Rifampicin		Sapropterin dihydrochloride		Sodium fluoride	34
Rifaximin		Scalp Preparations		Sodium Fusidate [fusidic acid]	
Rifinah		Scopoderm TTS		Dermatological	62
Rilutek		Sebizole		Infection	
Riluzole		Secukinumab		Sensory	
Riodine		Sedatives and Hypnotics		Sodium hyaluronate [Hyaluronic	220
Risedronate Sandoz		Seebri Breezhaler		acid]	220
Risedronate sodium		Selegiline hydrochloride		Sodium phenylbutyrate	
Risperdal Consta		Senna		Sodium polystyrene sulphonate	
		Senokot		. , , .	
Risperidone				Sodium tetradecyl sulphate	
Risperidone (Teva)		Sensipar SensoCard		Sodium valproate Sofradex	
Risperon					
Ritalin		Serenace		Soframycin	
Ritalin LA		Seretide		Solgar	
Ritalin SR		Seretide Accuhaler		Solifenacin Mylan	
Ritonavir		Serevent		Solifenacin succinate	
Rituximab (Mabthera)		Serevent Accuhaler		Solu-Cortef	
Rituximab (Riximyo)		Sertraline		Solu-Medrol	
Rivaroxaban		Setrona		Solu-Medrol-Act-O-Vial	
Rivastigmine		Setrona AU		Somatropin (Omnitrope)	
Rivotril		Sevredol	123	Sotalol	
Riximyo		Sex Hormones Non		Spacer device	
RIXUBIS		Contraceptive		Span-K	46
Rizamelt	129	Shield XL		Spiolto Respimat	224
Rizatriptan		shingles vaccine		Spiractin	54
Ropin	118	SII-Onco-BCG		Spiriva	
Ropinirole hydrochloride	118	Sildenafil	59	Spiriva Respimat	224
Rotarix	<mark>271</mark>	Silhouette MMT-373	<mark>23</mark>	Spironolactone	
Rotavirus oral vaccine	271	Siltuximab	208	Sporanox	96
Roxane		Simvastatin		Sprycel	156
Roxane-Propranolol	51	Simvastatin Mylan	55	Staphlex	92
Roxithromycin		Sinemet		Stemetil	130
Rubifen		Sinemet CR	118	SteroClear	
Rubifen SR		Sirolimus	217	Stesolid	126
Rugby Capsaicin Topical		Siterone		Stimulants/ADHD Treatments	
Cream	120	Slow-Lopresor		Stiripentol	
Rulide D		Smith BioMed Rapid Pregnanc		Stocrin	105
Rurioctocog alfa pegol [Reco		Test		Stomahesive	
factor VIII]		Sodibic		Strattera	

Strides Shasun	97	Temaccord	153	Topiramate	12
Stromectol	66	Temazepam	137	Topiramate Actavis	
Sucralfate	10	Temozolomide	153	Total parenteral nutrition (TPN)	4
Sulfadiazine Silver	62	Tenofovir disoproxil	100	TPN	
Sulfadiazine sodium	95	Tenofovir Disoproxil Teva		Tramadol hydrochloride	12
Sulfasalazine	8	Tenoxicam		Tramal SR 100	
Sulindac	110	Tensipine MR10	52	Tramal SR 150	12
Sulindac Mylan	110	Tepadina		Tramal SR 200	12
Sulphur	69	Terazosin		Trandate	5
Sulprix		Terbinafine	97	Tranexamic acid	
Sumatriptan		Terbutaline sulphate	223	Tranylcypromine sulphate	
Sunitinib		Teriflunomide	136	Trastuzumab	
Sunscreens		Teriparatide		Trastuzumab emtansine	
Sunscreens, proprietary	70	Testosterone		Travatan	23
Sure-T MMT-863		Testosterone cipionate		Travoprost	
Sure-T MMT-873		Testosterone esters		Travopt	
Sustagen Diabetic		Testosterone undecanoate		Treatments for Dementia	
Sustagen Hospital Formula		Tetrabenazine		Treatments for Substance	
Active	250	Tetrabromophenol		Dependence	14
Sustanon Ampoules		Tetracosactrin		Trental 400	
Sutent		Tetracycline		Tretinoin	
Sylvant		Thalidomide		Dermatological	6
Symbicort Turbuhaler 100/6		Thalomid		Oncology	
Symbicort Turbuhaler 200/6		Theophylline		Trexate	
Symbicort Turbuhaler 400/12		Thiamine hydrochloride		Triamcinolone acetonide	
Symmetrel		THIO-TEPA		Alimentary	3
Sympathomimetics		Thioguanine		Dermatological	6
Synacthen		Thiotepa		Hormone	
Synacthen Depot		Thymol glycerin		Triamcinolone acetonide with	
Synacthene Retard		Thyroid and Antithyroid Agents		gramicidin, neomycin and nysi	atin
Synflorix		Ticagrelor		Dermatological	
Synthroid		Tilade		Sensory	
Syntometrine		Tilcotil		Triaver	
Syrup (pharmaceutical grade)		Tillomed		Triazolam	
Systane Unit Dose		Timolol	170	Trimethoprim	
- T -	200	Cardiovascular	52	Trimethoprim with	
Tacrolimus	219	Sensory		sulphamethoxazole	
Tacrolimus Sandoz		Timoptol XE		[Co-trimoxazole]	q
Taliglucerase alfa		Tiotropium bromide		Trisequens	Q
Tambocor		Tiotropium bromide with		Trisul	
Tamoxifen citrate		olodaterol	224	Trophic Hormones	
Tamoxifen Sandoz		Tivicay		Tropicamide	23
Tamsulosin hydrochloride		TMP		Trusopt	
Tamsulosin-Rex		Tobramycin		TruSteel	
Tandem Cartridge		Infection	95	Tuberculin PPD [Mantoux] test	
Tandem t:slim X2 with Basal-IC		Sensory		Tubersol	27
Tap water		Tobramycin BNM		Two Cal HN	
Tarceva		Tobramycin Mylan		Two Cal HN RTH	
Tasigna		Tobrex		Tykerb	
Tasmar		Tocilizumab		Tysabri	13
Taurine		Tofranil		- U -	13
Tecfidera		Tolcapone		Ultibro Breezhaler	22
Tegretol		Topamax		Ultraproct	
Tegretol CR		Topical Products for Joint and	120	Umeclidinium	ეე
Telfast		Muscular Pain	111	Umeclidinium with vilanterol	
Teligent					
religerit	94	Topiderm	00	Univent2	۷۵, ۷۷

Ural	77	Vinorelbine	156	Zista	221
Urea	65	Vinorelbine Ebewe	156	Zithromax	90
Urex Forte	54	Viramune Suspension	105	Zoledronic acid	
Urinary Agents	76	ViruPOS		Hormone	78
Urinary Tract Infections		Vit.D3	33	Musculoskeletal	
Urinorm		Vita-B12	33	Zoledronic acid Mylan	7 <mark>8</mark>
Uromitexan		VitA-POS	234	Zopiclone	
Ursodeoxycholic acid		Vitabdeck		Zopiclone Actavis	137
Ursosan		Vital	246	Zostavax	
Utrogestan	82	Vitamin B complex	33	Zostrix	111
- V -		Vitamin B6 25		Zostrix HP	
Vaccinations	261	Vitamins		Zuclopenthixol decanoate	134
Vaclovir		Vivonex TEN		Zuclopenthixol hydrochloride	132
Valaciclovir		Voltaren		Zusdone	
Valganciclovir		Voltaren D	110	Zyban	141
Valganciclovir Mylan		Voltaren Ophtha		Zypine	
Vancomycin		Volumatic		Zypine ODT	
Vannair		Voriconazole		Zyprexa Relprevv	
Varenicline Pfizer		Vosol		Zytiga	
Varenicline tartrate		Votrient		=,9	
Varicella vaccine [Chickenpox		Vttack			
vaccine]	272	- W -			
Varicella zoster virus (Oka strai		Warfarin sodium	44		
attenuated vaccine [shingles	,	Wart Preparations			
vaccine]		Wasp venom allergy treatme			
Various		Water			
Varivax		Blood	45		
Vasodilators		Extemporaneous			
Vasopressin Agonists		Wool fat with mineral oil			
Vasorex		- X -			
Vedafil		Xarelto	44		
Veletri		Xifaxan			
Venclexta		XMET Maxamum			
Venetoclax		Xolair			
Venlafaxine		XP Maxamum			
Venomil		Xylocaine			
VENOX		Xylocaine 2% Jelly			
Ventavis		Xyntha			
Ventolin		- 7 -			
Vepesid		Zantac	9		
Verapamil hydrochloride		Zapril			
Vergo 16		Zarontin			
Vermox		Zaroxolyn			
Versacloz		Zavedos			
Vesanoid		Zeffix			
Vexazone		Zetlam			
Vfend		Ziagen			
Viaderm KC		Zidovudine [AZT]			
VidazaVidaza		Zidovudine [AZT] with	103		
Vigabatrin		lamivudine	105		
Vildagliptin		Zimvbe			
Vildagliptin with metformin	12	Zinc and castor oil			
hydrochloride	10	Zinc sulphate			
Vimpat		Zincaps			
Vinblastine sulphate		Zinnat			
Vincristine sulphate		Ziprasidone			
virionaline auiphale	100	∠ıµı asıu∪ı ı c	102		

