# June 2021 Volume 28 Number 1

#### **Editors:**

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

#### Circulation

Published each April, August and December. Changes to the contents are published in monthly updates.

Accessible in an electronic format at no cost from the PHARMAC website <a href="https://www.pharmac.govt.nz/schedule">www.pharmac.govt.nz/schedule</a>.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

#### Production

Typeset automatically from XML and T<sub>E</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

#### **Programmers**

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz @Pharmaceutical Management Agency



ISSN 1179-3686 pdf ISSN 1172-9376 print

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to PHARMAC and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to PHARMAC should be in written form and not by reproduction of the PHARMAC logo. While care has been taken in compiling this Schedule, PHARMAC takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	illitioducing Phanimac	L
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	61
	Genito Urinary System	71
	Hormone Preparations – Systemic	78
	Infections – Agents For Systemic Use	89
	Musculoskeletal System	110
	Nervous System	118
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	220
	Sensory Organs	229
	Various	235
Section C	Extemporaneous Compounds (ECPs)	237
Section D	Special Foods	240
Section I	National Immunisation Schedule	261
	Index	273

Introducing PHARMAC

# Introducing PHARMAC

The Pharmaceutical Management Agency (PHARMAC) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. PHARMAC negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

#### PHARMAC's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about PHARMAC and the way we make funding decisions can be found on the PHARMAC website at <a href="https://www.pharmac.govt.nz/about">https://www.pharmac.govt.nz/about</a>.

# **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by PHARMAC.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements PHARMAC has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

# **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

### Example



# Glossary

### **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

# **SECTION B: ALIMENTARY TRACT AND METABOLISM**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	<b>✓</b>	Gaviscon Infant
SODIUM ALGINATE     Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60		Gaviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 m		Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg	12.56	100	✓.	Alu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsementOnly when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according	cium carbonate tablet	500 m s or v		Roxane um carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on  * Tab 2 mg*  * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy	166.50	90	<b>✓</b>	Entocort CIR
<b>⇒SA1886</b> Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant practithe following criteria: Both:	titioner. Approvals va	ılid fo	r 6 months	for applications meeting
1 Mild to moderate ileal, ileocaecal or proximal Crohn's dise	ease; and			

0.4 Dishetes as

2 Any of the following:

2.1 Diabetes; or

2.2 Cushingoid habitus; or

2.3 Osteoporosis where there is significant risk of fracture; or

(N	Subsidy fanufacturer's Price)	Subsi	Fully dised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes; or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture; or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
  - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with \* is an unapproved indication.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

#### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	<ul><li>✓ Cortifoam S29</li><li>✓ Colifoam</li></ul>
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone S29
SODIUM CROMOGLICATE  Cap 100 mgSULFASALAZINE	92.91	100	✓	Nalcrom
* Tab 500 mg		100 100		Salazopyrin Salazopyrin EN

# Local preparations for Anal and Rectal Disorders

### Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g6.35	30 g OP	<ul><li>Ultraproct</li></ul>
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg2.66	12	<ul><li>Ultraproct</li></ul>
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

# **Management of Anal Fissures**

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy 30 q OP ✓ Rectogesic

### ⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

# **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	<ul><li>Buscopan</li></ul>
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

# **Antiulcerants**

# **Antisecretory and Cytoprotective**

#### MISOPROSTOL

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

\* Tab 200 mcg - Up to 120 tab available on a PSO ......41.50 ✓ Cytotec

		ALIMENTAR	Y TR	ACT AND	METABOLISM
		Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Brand or Generic Manufacturer
Helicol	pacter Pylori Eradication				
Tab 5	ROMYCIN 00 mg - Subsidy by endorsement  Maximum of 14 tab per prescription Subsidised only if prescribed for helicobacter pylori endote: the prescription is considered endorsed if clarify inhibitor and either amoxicillin or metronidazole.	radication and preso		n is endorse	
H2 Ant	agonists				
	INE – Only on a prescription 0 mg	4.91	100	<b>✓</b> F	Famotidine Hovid §29
⊁ Tab 4	0 mg	8.48	100	<b>✓</b> F	Famotidine Hovid §29
	mg per ml, 4 ml - Subsidy by endorsementubsidy by endorsement - Subsidised for patients receiv		10 rt of p		Mylan S29 e.
a) Or b) Su pre of & Oral li	NE – Subsidy by endorsement only on a prescription besidy by endorsement – Subsidised for patients who we escription is endorsed accordingly. Pharmacists may a prior dispensing of ranitidine. only 150 mg per 10 ml	nnotate the prescrip		s endorsed v	
	Pump Inhibitors	:1 2021)			
ANSOPF	•				
k Cap 1 k Cap 3	5 mg 0 mg		100 100	_	anzol Relief anzol Relief
	POLE meprazole suspension refer Standard Formulae, page 2 0 mg		90	✓ (	Omeprazole actavis
	meprazole actavis 10 to be Sole Supply on 1 August 20 0 mg		90	<b>√</b> (	Omeprazole actavis
	meprazole actavis 20 to be Sole Supply on 1 August 20 mg		90	✓ (	Omeprazole actavis
0	meprazole actavis 40 to be Sole Supply on 1 August 20	)21	<b>.</b>		Al along a A

5 g

100

100

✓ Midwest

✓ <u>Dr Reddy's</u>

<u>Omeprazole</u>
✓ Ocicure (\$29)

✓ Panzop Relief

✓ Panzop Relief

**PANTOPRAZOLE** 

Powder – Only in combination......42.50

Only in extemporaneously compounded omeprazole suspension. Inj 40 mg ampoule with diluent .......33.98

\* Tab EC 20 mg ......2.02

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE  Tab 120 mgSUCRALFATE	14.51	50	<b>√</b> (	Gastrodenol 629
Tab 1 g	35.50 (48.28)	120	(	Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Reta	ail pharmacy			
Tab 550 mg	625.00	56	<b>✓</b> <u>&gt;</u>	<u> (ifaxan</u>
➤SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatol epatologist. Approvals valid for 6 months where the patiolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Piepatologist. Approvals valid without further renewal unleasenefiting from treatment.	ent has hepatic encephalop Practitioner on the recomme	athy d	espite an a	dequate trial of maximun penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Ref Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 0 ml 0	<b>✓</b> F	Proglicem \$29 Proglicem \$29 Proglycem \$29
■ SA1320 Special Authority for Subsidy nitial application from any relevant practitioner. Approvypoglycaemia caused by hyperinsulinism.	als valid for 12 months whe	re use	d for the tre	atment of confirmed
Renewal from any relevant practitioner. Approvals valid of ppropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE				
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓ (	Glucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL Inj human 100 u per ml	25.26 1	0 ml C		Actrapid Humulin R
Inj human 100 u per ml, 3 ml	42.66	5	<b>√</b>	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE				
Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	<b>✓</b> 1	NovoMix 30 FlexPen

	Subsidy	F	ully Brand or
	(Manufacturer's Price	e) Subsidis	,
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE			<b></b>
Inj human 100 u per ml	17.68		<ul><li>✓ Humulin NPH</li><li>✓ Protaphane</li></ul>
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
- , ,			✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	<ul><li>✓ Humulin 30/70</li><li>✓ Mixtard 30</li></ul>
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
•			✓ PenMix 30
			✓ PenMix 40
JOHN N. JORDO WITH INCH IN JORDO DROTAMINE			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per ml.			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml			
3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen		5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
Inj 100 u per ml, 10 ml			✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE Inj 100 u per ml, 10 ml	27.02	1	✓ Apidra
Inj 100 u per ml, 3 ml			✓ Apidra
Inj 100 u per ml, 3 ml disposable pen			✓ Apidra SoloStar
NSULIN LISPRO			
Inj 100 u per ml, 10 ml		10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
CARBOSE			
★ Tab 50 mg		90	✓ Glucobay
₭ Tab 100 mg	10.47 6.40	90	<ul><li>✓ Accarb</li><li>✓ Glucobay</li></ul>
- 145 155 Hig	20.23	00	✓ Accarb
Oral Hypoglycaemic Agents			
ilibenclamide			
₭ Tab 5 mg	6.00	100	✓ <u>Daonil</u>
			<del></del>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	-	Fully Subsidised	Brand or Generic
	\$	Per	1	Manufacturer
GLICLAZIDE				
* Tab 80 mg	15.18	500	1	<u>Glizide</u>
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓	<u>Minidiab</u>
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000	✓	<u>Apotex</u>
* Tab immediate-release 850 mg	7.04	500	•	Apotex
PIOGLITAZONE				
* Tab 15 mg	1.12	28	✓	<u>Vexazone</u>
	3.47	90	_	Vexazone
* Tab 30 mg		90	_	<u>Vexazone</u>
* Tab 45 mg	7.10	90	•	<u>Vexazone</u>
VILDAGLIPTIN			_	
Tab 50 mg	35.00	60	/	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride		60	_	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	/	Galvumet

### **SGLT2 Inhibitors**

### ⇒SA2029 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Any of the following:
  - 2.1 Patient is Maaori or any Pacific ethnicity\*: or
  - 2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
  - 2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*: or
  - 2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
  - 2.5 Patient has diabetic kidney disease (see note b)\*; and
- 3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months; and
- 4 Treatment will not be used in combination with a funded GLP-1 agonist.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

ΕN	IPAGLIFLOZIN - Special Authority see SA2029 above - Retail	pharmacy		
*	Tab 10 mg	58.56	30	✓ Jardiance
*	Tah 25 mg	58 56	30	✓ lardiance

Fully

Brand or

	(Manufacturer's Price)	Sul Per	bsidised	Generic Manufacturer	
EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – pharmacy	Special Authority see S	SA2029 o	n the pre	vious page - Reta	ail
* Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ J	ardiamet	
* Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	✓ J	ardiamet	
* Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ J	ardiamet	
* Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	✓ J	ardiamet	

Subsidy

# **Diabetes Management**

### **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

# **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

13

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes; or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

### BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood	l glucose	test s	trips26	5.20 50	o test OP	SensoCard

			_
Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

# **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES	<ul> <li>Maximum of 200</li> </ul>	dev per prescription
---------------------	------------------------------------	----------------------

	00 407	40.50	400	/ D D 14:
*	29 g × 12.7 mm	10.50	100	B-D Micro-Fine
*	31 g × 5 mm	11.75	100	<ul> <li>B-D Micro-Fine</li> </ul>
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm	10.50	100	✓ B-D Micro-Fine
*			100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200	) dev per pre	escription
	Syringe 0.3 ml with 29 g × 12.7 mm needle		100	✓ B-D Ultra Fine
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle	` '	100	✓ B-D Ultra Fine
	-,···g- ··· ··· g · · · ··· · · · ·	1.30	10	
		(1.99)	. •	B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
•	Symigo do mi mar or give min nocale minimini	1.30	10	
		(1.99)	. •	B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	` '	100	✓ B-D Ultra Fine
•	5)g5 : =5 g % :=1	1.30	10	2 2 0 0
		(1.99)	. •	B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
-1-	Syrings This High of g A o Hill Hoodio	1.30	100	- D D ORIGINICII
			10	D D I Illiana Einea II
		(1.99)		B-D Ultra Fine II

# **Insulin Pumps**

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year	period.		
Min basal rate 0.025 U/h	8,800.00	1	MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

### ⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Por 🗸	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Fither:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Suk	sidised	Generic	
(Manuacturer 3 i lice)	Out	Joiuiseu	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
(Manufacturer's Frice)	Per 🗸	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol: and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Either:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.

### **Insulin Pump Consumables**

### ⇒SA1985 Special Authority for Subsidy

**Initial application** — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 2 Eithor
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
    - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
pump therapy; and				
4 The patient is continuing to derive benefit from pump the			., .	
5 The patient had achieved and is maintaining a HbA1c o 6 The patient has had no increase in severe unexplained				
7 The patient's HbA1c has not deteriorated more than 5 n	,, ,, ,		iii baseiiile, a	allu
8 Either:	into/mor from baseline	, and		
8.1 Applicant is a relevant specialist; or				
8.2 Applicant is a nurse practitioner working within the	neir vocational scope.			
Renewal — (Previous use before 1 September 2012) only for	rom a relevant specialis	st or n	urse practitio	oner. Approvals valid for 2
years for applications meeting the following criteria:				
All of the following:				
1 The patient is continuing to derive benefit according to t than 80 mmol/mol; and	he treatment plan and	has m	naintained a l	HbA1c of equal to or less
2 The patient's HbA1c has not deteriorated more than 5 n	amol/mol from initial an	nlicat	ion: and	
3 The patient has not had an increase in severe unexplain				ne: and
4 Either:				·-, -··-
4.1 Applicant is a relevant specialist; or				
4.2 Applicant is a nurse practitioner working within the control of the contro	neir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 18 – Retail p	harma	асу	
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 packs of cartridge sets will be funded p		4 00		andan Oastaldaa
Cartridge 300 U, t:lock × 10		1 OP		andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Specia	al Authority see SA198	5 on p	oage 18 – Re	etail pharmacy
a) Maximum of 3 sets per prescription     b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T
g				MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ M	liniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	• • • M	liniMed Sure-T
Commented mandles 00 and taking a 40	100.00	4.00		MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	• W	liniMed Sure-T

8 mm steel needle; 80 cm tubing × 10 ......130.00

6 mm steel needle; 29 G; manual insertion; 60 cm tubing x

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

1 OP

1 OP

1 OP

1 OP

MMT-866A

MMT-874A

✓ MiniMed Sure-T

✓ MiniMed Sure-T

MMT-876A

✓ Sure-T MMT-863

✓ Sure-T MMT-873

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 18 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles	1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00	) 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.00	0 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.00	1 OP	✓ TruSteel

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per ✓ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 18 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- a) Maximum of 12 infusion sate will be funded nor year

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette  MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set  MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing x 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A
			WINI I -OOOA

	Subsidy (Manufacturer's Pr	rice) S	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SERTION WITH	H INSERTI	ON DEVIC	E) - Special Authority see
SA1985 on page 18 – Retail pharmacy				
a) Maximum of 3 sets per prescription     b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c	cm			
line × 10 with 10 needles		1 OP	✓ A	utoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cr		4.00		
line × 10 with 10 needles		1 OP		utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	NSERTION) - S	pecial Auth	ority see S	A1985 on page 18 –
Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles; luer lock		1 OP	_	ilhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	IT INSERTION V	VITH INSE	RTION DE	VICE) - Special Authority
see SA1985 on page 18 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles		1 OP	✓ A	utoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 c line × 10 with 10 needles		1 OP	<b>√</b> ∆	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device;	140.00	1 01	• ^	atooon so
110 cm line × 10 with 10 needles	140.00	1 OP	✓ A	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m			
line × 10 with 10 needles		1 OP		utoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	IT INSERTION)	- Special	Authority se	e SA1985 on page 18 -
Retail pharmacy				
a) Maximum of 3 sets per prescription     b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wit	:h			
10 needles; luer lock		1 OP	<b>√</b> Q	uick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wit		1.00		huiale Cat MMT 000
10 needles; luer lock		1 OP		uick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 of	n page 18 – Heta	aii pharmad	У	
a) Maximum of 3 sets per prescription     b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per	year.			
10 × luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP		DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10	50.00	1 OP	✓ M	liniMed
				1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	✓ M	liniMed
2		. 0.	- 10	3.0 Reservoir
				MMT-332A

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
· · · ·	Por 🗸	Manufacturor	

# **Digestives Including Enzymes**

#### PANCREATIC ENZYME

PANCREATIC ENZYME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase,			
1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	<ul><li>Creon Micro</li></ul>
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	v – Retail pha	rmacy	
Cap 250 mg	32.95	100	✓ <u>Ursosan</u>

### ⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

### Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Normacol Plus

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

### Laxatives

# **Bulk-forming Agents**

* Powder for oral soin12.20	500 g OP	Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		
* Dry	2 500 g OP	
(17.32	2)	Normacol Plus
2.41	200 a OP	

(8.72)

### Faecal Softeners

* Tab 50 mg2.31	100	✓ Coloxyl
* Tab 120 mg3.13		✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES		
* Tab 50 mg with sennosides 8 mg	200	✓ Laxsol
POLOXAMER – Only on a prescription		
Not funded for use in the ear.		
* Oral drops 10%	30 ml OP	✓ Coloxyl

# **Opioid Receptor Antagonists - Peripheral**

DOCUSATE SODIUM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority see	SA1691 below - Retail	pharmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	Relistor
•	246.00	7	✓ Relistor

### ⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	(Manufacturer's Pric		osidised Generic	
	\$	Per	✓ Manufacturer	_
Osmotic Laxatives				
GLYCEROL				
* Suppos 3.6 g – Only on a prescription	9.25	20	✓ <u>PSM</u>	
LACTULOSE – Only on a prescription				
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ <u>Laevolac</u>	
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BI		SODIUM C	CHLORIDE	
Powder for oral soln 13.125 g with potassium chloride 46.6 n		20		
sodium bicarbonate 178.5 mg and sodium chloride 350.	7 mg6.70	30	✓ <u>Molaxole</u>	
SODIUM ACID PHOSPHATE – Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate  Enema	
CODILINA CITRATE MITH CODILINA LALIDVI, CHILDHOACETATE	Only an a muse		Enema	
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE	, ,	inpuon		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml		50	✓ Micolette	
0111	20.00	00	· imoolette	
Stimulant Laxatives				
BISACODYL - Only on a prescription				
* Tab 5 mg	5.99	200	✓ <u>Lax-Tab</u>	
* Suppos 10 mg	3.74	10	✓ <u>Lax-Suppositories</u>	
SENNA - Only on a prescription				
* Tab, standardised		100		
	(8.21)	20	Senokot	
	0.43	20	Senokot	
	(2.06)		Seriokot	

Subsidy

Fully

Brand or

# **Metabolic Disorder Agents**

### ⇒SA1986 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and

Subsidy	Fully	Brand or
acturer's Price)	Subsidised	Generic
 \$ Pe	er 🗸	

continued...

- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Special Authority	see SA2042 below -	- Retail pharmacy
------------------------------	--------------------	-------------------

Tab 1,000 mg	CBS	90	<ul><li>Clinicians</li></ul>
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### BETAINE - Special Authority see SA1987 below - Retail pharmacy

# ⇒SA1987 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
  - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
  - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10	<ul> <li>Special Authority see</li> </ul>	SA2039 on the nex	t page – Retail pharm	acy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

### ⇒SA2039 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

### ⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT

# ⇒SA1623 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
  - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

### ⇒SA1695 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts: or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below	- Retail pharmacy		
Tab 500 mg	CBŚ	30	<ul><li>Solgar</li></ul>
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

### ⇒SA2040 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see SA2041 below – Re	tail pharmacy		
Tab 100 mg	CBS	100	<ul><li>Country Life</li></ul>
Cap 100 mg	CBS	100	✓ Solgar

### ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

**Renewal** only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE	- Special Authority see SA1989 below -	- Retail pharmacy	
Tab soluble 100 mg	1,452.70	30 OP	✓ Kuvan

### ⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and

	Subsidy		Fully	Brand or
(I	Manufacturer's Price)	Subsic	lised	Generic
	\$	Per	✓	Manufacturer

continued...

- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

#### 

### ⇒SA1599 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1990 below – Retail pharmacy
Grans 483 mg per g.......2,016.00 174 g OP 

Pheburane

# **⇒SA1990** Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# ⇒SA2043 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Fully

Subsidy (Manufacturer's Price)

Subsidised Per

1

Brand or Generic Manufacturer

### Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy 

✓ Elelyso

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 PHARMAC PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

#### **Access Criteria**

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by PHARMAC; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia. thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
  - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse: chronic bone pain not controlled by other pharmaceuticals; or
  - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
  - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

#### \*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose: and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication.

ALIMENTARY TRACT AND METABOLISM	Subsidy (Manufacturer's Price)	Fully Subsidised Per	
continued and/or adjustment of infusion rates; and 5) Patient has not developed another medical condition th ERT; and 6) Patient is compliant with regular treatment and taligluce every other week rounded to the nearest whole vial (20 7) Supporting clinical information including test reports, M investigations are submitted to the Gaucher Panel for a	erase alfa is to be admin 0 units), unless otherwi RI whole body STIR, ha	istered at a dos se agreed by F	e no greater than 30 unit/kg HARMAC; and
Mouth and Throat  Agents Used in Mouth Ulceration			
BENZYDAMINE HYDROCHLORIDE  Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with Endorsement	(20.31)	500 ml esult of treatme	Difflam ent for cancer, and the

	(20.31)		Difflam
Additional subsidy by endorsement for a patient who ha prescription is endorsed accordingly.	s oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	•	Orabase
	1.52	5 g OP	
	(3.60)	_	Orabase
Powder	8.48	28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
Transcrib got on 70 min obtaine man one need of or 70 min min	(6.00)	. o g o .	Bonjela
TRIAMCINOLONE ACETONIDE	(0.00)		20.,jo.u
Paste 0.1%	E 22	5 a OP	✓ Kenalog in Orabase
Fasie 0.176		5 y OF	Kenalog III Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			•
Oral gel 20 mg per g	171	40 g OP	✓ Decozol
	4.74	70 g Oi	- <u>DCCO201</u>
NYSTATIN	4.70	04 100	( NIII
Oral liq 100,000 u per ml	1./6	24 ml OP	✓ <u>Nilstat</u>
Other Oral Agents			

For folinic mouthwash, pilocarpine oral liquid or saliva substitute formula refe	r Standard Formulae, page 237
THYMOL GLYCERIN	
* Compound, BPC	5 500 ml <b>✓ PSM</b>

Fully

Brand or

Subsidy

	(Manufacturer's Pric	e) Subs Per	Hully Brand or dised Generic  Manufacturer
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN  * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	SO 1.89	3	✓ <u>Neo-B12</u> ✓ Vita-B12
	3.15	5	<ul> <li>Hydroxocobalamin</li> <li>Mercury Pharma</li> </ul>
PYRIDOXINE HYDROCHLORIDE  a) No more than 100 mg per dose b) Only on a prescription			·
* Tab 25 mg – No patient co-payment payable      * Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Apo-Pyridoxine
THIAMINE HYDROCHLORIDE – Only on a prescription	10.00	300	• Apo-i yildoxille
* Tab 50 mg	7.09	100	✓ Max Health
VITAMIN B COMPLEX  * Tab, strong, BPC	7.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID  a) No more than 100 mg per dose b) Only on a prescription  * Tab 100 mg	9.90	500	✓ Cvite
Vitamin D			
ALFACALCIDOL  * Cap 0.25 mcg  * Cap 1 mcg  * Oral drops 2 mcg per ml  CALCITRIOL	87.98	100 100 20 ml OP	✓ One-Alpha ✓ One-Alpha ✓ One-Alpha
* Cap 0.25 mcg      * Cap 0.5 mcg  COLECALCIFEROL		100 100	✓ Calcitriol-AFT ✓ Calcitriol-AFT
* Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescripti * Oral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml OP	✓ <u>Vit.D3</u> ✓ Puria
Multivitamin Preparations			
MULTIVITAMIN RENAL – Special Authority see SA1546 below –  * Cap		30	✓ Clinicians Renal Vit
■ SA1546 Special Authority for Subsidy  Initial application from any relevant practitioner. Approvals valid the following criteria:  Either:  1. The nationt has chronic kidney disease and is receiving all the patient has			

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

ALIMENTARY TRACT AND METABOLISM				
	Subsidy (Manufacturer's Pri	ce) Subs	Fully idised	Brand or Generic Manufacturer
MULTIVITAMINS – Special Authority see SA1036 below – Reta	ail pharmacy			
* Powder	72.00	200 g OP	<b>✓</b> P	aediatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valinborn errors of metabolism.  Renewal from any relevant practitioner. Approvals valid withou				·
approval for multivitamins.			·	·
VITAMINS			_	
* Tab (BPC cap strength)		1,000	✓ <u>N</u>	<u>lvite</u>
* Cap (fat soluble vitamins A, D, E, K) - Special Authority se SA1720 below - Retail pharmacy		60	<b>✓</b> ∨	itabdeck
⇒SA1720 Special Authority for Subsidy	20.10	00	•	nabaoon
Initial application from any relevant practitioner. Approvals varithe following criteria:  Any of the following:  1 Patient has cystic fibrosis with pancreatic insufficiency; of 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome.	r	aigwai uiiless	noune	o to applications meeting
Minerals  Calcium				
CALCIUM CARBONATE				
* Tab 1.25 g (500 mg elemental)	6.69	250	<b>✓</b> C	alci-Tab 500
* Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme		76	_	Cacit S29
Subsidy by endorsement – Only when prescribed for paconsidered unsuitable.		5 years) whe	ere calci	ium carbonate oral liquid is
CALCIUM GLUCONATE	00.00	10		lavi lia aliib
* Inj 10%, 10 ml ampoule	32.00	10	• 10	lax Health - Hameln S29
	64.00	20	✓ N	lax Health S29
Fluoride				
SODIUM FLUORIDE  * Tab 1.1 mg (0.5 mg elemental)	5.75	100	<b>√</b> P	SM
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	leuroTabs
Iron				

FERRIC CARBOXYMALTOSE - Special Authority see SA1840 on the next page - Retail pharmacy Inj 50 mg per ml, 10 ml......150.00 ✓ Ferinject

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
<b>\$</b>	Per	•	Manufacturer	

### ⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- - 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
  - 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

FERROUS FUMARATE  * Tab 200 mg (65 mg elemental)	100	✓ Ferro-tab	
FERROUS FUMARATE WITH FOLIC ACID  * Tab 310 mg (100 mg elemental) with folic acid 350 mcg4.68	60	✓ Ferro-F-Tabs	
FERROUS SULFATE  * Oral liq 30 mg (6 mg elemental) per 1 ml12.08	500 ml	✓ <u>Ferodan</u>	
FERROUS SULPHATE  * Tab long-acting 325 mg (105 mg elemental)2.06	30	✓ Ferrograd	
IRON POLYMALTOSE  * Inj 50 mg per ml, 2 ml ampoule34.50	5	✓ Ferrosig	
Magnesium			
For magnesium hydroxide mixture refer Standard Formulae, page 237			

# For magnesium hydroxide mixture refer Standard Formulae, page 237

MAGNESIUM HYDROXIDE

355 ml ✓ Phillips Milk of Magnesia S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MAGNESIUM SULPHATE				
* Inj 2 mmol per ml, 5 ml ampoule	25.53 28.00	10	✓	Martindale DBL DBL S29 S29
Martindale to be Sole Supply on 1 July 2021 (DBL Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 July 2021) (DBL S29 S29 Inj 2 mmol per ml, 5 ml ampoule to be delisted 1 J	luly 2021)			
Zinc				
ZINC SULPHATE  * Cap 137.4 mg (50 mg elemental)	11.00	100	/	Zincaps

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

## **Antianaemics**

### Hypoplastic and Haemolytic

#### ⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus: and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus: and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal — (chronic renal failure)** from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

	Subsidy (Manufacturer's Price)		Fully	
	\$	Per		Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу		
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	•	Binocrit

### Megaloblastic

-01	10	40	
-OL	_IC	AC	טו

*	Tab 0.8 mg21.84	1,000	1	Apo-Folic Acid
	Tab 5 mg	500	1	Apo-Folic Acid
	Oral lig 50 mcg per ml	25 ml OP	1	Biomed

## Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
lnj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	Alprolix
ELTROMBOPAG – Special Authority see SA1743 Wastage claimable	below - Retail pharmacy		

# Tab 50 mg ......3,100.00 **➤ SA1743** Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding: or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

**Initial application** — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

continued...

✓ Revolade

✓ Revolade

28 28

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
 \$	Per	✓	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

## 

✓ Hemlibra	1	vial3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	ıl vial	Inj 60 mg in 0.4 ml via
✓ Hemlibra	1	ml vial12,492.00	Inj 105 mg in 0.7 ml vi
✓ Hemlibra	1	vial17,846.00	Inj 150 mg in 1 ml vial

⇒SA1969 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:

continued...

<del></del>				
	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Su	ıbsidised	Generic
	\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
  - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
    - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

**Renewal** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

#### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

#### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

#### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Subject to criteria.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

#### NONACOG GAMMA. [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

nj 500 iu vial	435.00	1	✓ RIXUBIS
nj 1,000 iu vial	870.00	1	✓ RIXUBIS
nj 2,000 iu vial		1	✓ RIXUBIS
nj 3,000 iu vial	·	1	✓ RIXUBIS
) -/ · · · · · · · · · · · · · · · · ·	,		

	0.1		- ·	
	Subsidy (Manufacturer's Price)	Su	Fully bsidised	
	\$	Per	<b>√</b>	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	[Xpharm]			
For patients with haemophilia. Preferred Brand of short half	-life recombinant factor	r VIII. A	ccess to	o funded treatment is
managed by the Haemophilia Treaters Group in conjunction				
Inj 250 iu vial		1		Advate
Inj 500 iu vial	420.00	1	✓	Advate
Inj 1,000 iu vial	840.00	1	✓	Advate
Inj 1,500 iu vial	1,260.00	1	✓	Advate
Inj 2,000 iu vial		1		Advate
Inj 3,000 iu vial	,	1	/	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the I	Vational	Haemo	philia Management Group,
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1	•	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatm		d treatm	ent is n	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			_	
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial		1		Adynovate
Inj 2,000 iu vial	2,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Konakion MM
ing to mg por mi, i mi op to o mg available on a room		•		Tronauton iiiii
Antithrombotic Agents				
- Tillian Omsono Algorito				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	•	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
				<u>Multichem</u>
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	1	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next pag				-
* Tab 90 mg		56	1	Brilinta

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

#### ⇒SA1955 Special Authority for Subsidy

**Initial application** — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
- 1.2 Patient is about to have a neurological stenting procedure performed\*; and
- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

**Initial application** — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

10

10

10

10

10

Fully

Brand or

✓ Clexane

✓ Clexane

Clexane

✓ Clexane Forte

✓ Clexane Forte

	(Manufacturer's Price) \$	Sub Per		Generic Manufacturer	
Heparin and Antagonist Preparations					
ENOXAPARIN SODIUM - Special Authority see SA1646 below	, ,				
Inj 20 mg in 0.2 ml syringe	27.93	10	✓ Cle	xane	
Inj 40 mg in 0.4 ml syringe	37.27	10	✓ Cle	xane	

Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Low molecular weight heparin treatment is required during a patients pregnancy; or

Inj 60 mg in 0.6 ml syringe......56.18

Inj 80 mg in 0.8 ml syringe......74.90

Inj 120 mg in 0.8 ml syringe.......116.55

- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery: or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	✓ Hospira
	42.40		✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg	76.36	60	✓	Pradaxa
Cap 150 mg		60	1	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	✓	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	1	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
WARFARIN SODIUM  Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	1	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg		100	1	Marevan
* Tab 5 mg		50	✓	Coumadin
•	11.48	100	1	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail p	harmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe		10	1	Nivestim
OA4050 On a stat Authority for Outside				

#### ⇒SA1259 Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10<sup>9</sup>/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC  $< 0.5 \times 10^9$ /L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

#### ⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*).

Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	DECOD AND			TIMING OTIGANS
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE]  * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO  * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO		5 1		Biomed Biomed
POTASSIUM CHLORIDE  * Inj 75 mg per ml, 10 ml	55.00	50		AstraZeneca Potassium Chloride Aquettant \$29
	65.00		•	Juno
(AstraZeneca Inj 75 mg per ml, 10 ml to be delisted 1 November	,	000	41	
(Potassium Chloride Aguettant S29 Inj 75 mg per ml, 10 ml to be	e aeiistea 1 Novemb	er 202	1)	
SODIUM BICARBONATE Inj 8.4%, 50 ml	19.95	1	/	Biomed
<ul><li>a) Up to 5 inj available on a PSO</li><li>b) Not in combination</li></ul>		1		Diamad
Inj 8.4%, 100 mla) Up to 5 inj available on a PSO b) Not in combination	20.50	ı	•	Biomed
SODIUM CHLORIDE  Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	use except when us	sed in o	conjunctio	on with an antibiotic intended
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml 1,000 m		Baxter Baxter
Only if prescribed on a prescription for renal dialysis, ma	ternity or post-natal	care in	the home	e of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standar		5	•	Biomed
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	2.80	20	•	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50		Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	•	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN) Infusion	CBS	1 OP	/	TPN
WATER				
<ol> <li>On a prescription or Practitioner's Supply Order only when Schedule requiring a solvent or diluent; or</li> <li>On a bulk supply order; or</li> <li>When used in the extemporaneous compounding of eyed) When used for the dilution of sodium chloride soln 7% f</li> </ol>	e drops; or			n listed in the Pharmaceutical
Inj 10 ml ampoule – Up to 5 inj available on a PSO Inj 20 ml ampoule – Up to 5 inj available on a PSO		50 20	✓	Pfizer Fresenius Kabi Multichem
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85 3	00 g O	P 🗸	Calcium Resonium
▲ Th		- "		

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pri \$	ice) Subs Per	Fully idised	Brand or Generic Manufacturer
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO  COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)		50 1.000 ml OP	-	Electral Pedialyte -
PHOSPHORUS Tab eff 500 mg (16 mmol)		100	_	Bubblegum Phosphate Phebra
POTASSIUM CHLORIDE  * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	(11.85)	60		Chlorvescent
* Tab long-acting 600 mg (8 mmol) SODIUM BICARBONATE Cap 840 mg		200		<u>Span-K</u> Sodibic
SODIUM POLYSTYRENE SULPHONATE Powder	84.65	454 g OP		Sodibic Resonium-A

	Subsidy (Manufacturer's Price \$	) Sub Per	Fully osidised	Brand or Generic Manufacturer
Alpha-Adrenoceptor Blockers				
Alpha Adrenoceptor Blockers				
DOXAZOSIN  * Tab 2 mg  * Tab 4 mg  PHENOXYBENZAMINE HYDROCHLORIDE		500 500		po-Doxazosin po-Doxazosin

# \* Tab 5 mg ......11.70 TERAZOSIN – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking terazosin prior to 1 October 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of terazosin.

216.67

1ab 2 mg	500
14.20	28
Tab 5 mg10.90	500
24.80	28

\* Cap 10 mg......65.00

✓ Apo-Terazosin

✓ Teva S29

✓ BNM S29

✓ Dibenzyline S29

✓ Apo-Prazosin

✓ Apo-Prazosin

✓ Apo-Prazosin

30

100

100

100

100

✓ Apo-Terazosin✓ Teva S29

(Apo-Terazosin Tab 2 mg to be delisted 1 August 2021)

(Teva \$29) Tab 2 mg to be delisted 1 August 2021)

(Apo-Terazosin Tab 5 mg to be delisted 1 August 2021)

(Teva \$29 Tab 5 mg to be delisted 1 August 2021)

## Agents Affecting the Renin-Angiotensin System

#### **ACE Inhibitors**

CAPTOPRII

**PRAZOSIN** 

Oral liq 5 mg per ml	95 ml OP	✓ Capoten
135.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

(Captopril-Mylan S29 Oral lig 5 mg per ml to be delisted 1 January 2022)

#### CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

* Tab 0.5 mg	2.09	90	Zapril
* Tab 2.5 mg		90	✓ Zapril
Tab 5 mg		90	✓ Zapril
ENALAPRIL MALEATE			
* Tab 5 mg	1.82	100	✓ Acetec
* Tab 10 mg		100	✓ Acetec
* Tab 20 mg		100	✓ Acetec

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

=					
		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	
_		\$	Per		Manufacturer
LIS	INOPRIL				
_	Tab 5 mg	2.07	90	1	Ethics Lisinopril
	Tab 10 mg		90		Ethics Lisinopril
-	Tab 20 mg		90		Ethics Lisinopril
	v		00		<u> </u>
PΕ	RINDOPRIL	0.75	~~	,	An a Bankada adl
	Tab 2 mg		30		Apo-Perindopril
	Tab. Assess	4.95	~~		Coversyl
	Tab 4 mg		30		Apo-Perindopril
		6.30		•	Coversyl
QU	INAPRIL				
	Tab 5 mg	6.01	90	✓	Arrow-Quinapril 5
	Tab 10 mg	3.16	90	✓	Arrow-Quinapril 10
	Tab 20 mg	4.89	90	✓	Arrow-Quinapril 20
Α	CE Inhibitors with Diuretics				
QU	INAPRIL WITH HYDROCHLOROTHIAZIDE				
	Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	✓	Accuretic
		3.83	30	✓	Accuretic 10
*	Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	✓	Accuretic 20
A	ngiotensin II Antagonists				
CA	NDESARTAN CILEXETIL				
*	Tab 4 mg	1.90	90	✓	Candestar
*	Tab 8 mg		90		Candestar
*	Tab 16 mg		90		Candestar
	Tab 32 mg		90		Candestar
	SARTAN POTASSIUM				
_		1 56	84	./	Lacartan Actovia
	Tab 12.5 mg		84		Losartan Actavia
	Tab 25 mg		84		Losartan Actavia
	Tab 50 mg				Losartan Actavis
*	Tab 100 mg	3.50	84	•	Losartan Actavis
A	ngiotensin II Antagonists with Diuretics				
10	SARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
LO	Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	1	Arrow-Losartan &
	rab 50 mg with hydrochlorothazide 12.5 mg	1.00	00	•	Hydrochlorothiazide
					<u> </u>
Α	ngiotensin II Antagonists with Neprilysin Inhib	oitors			
-	g				
SA	CUBITRIL WITH VALSARTAN - Special Authority see SA190				
	Note: Due to the angiotensin II receptor blocking activity of s	sacubitril with valsarta	n it s	hould not	be co-administered with an
	ACE inhibitor or another ARB.				
	Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓	Entresto 24/26
	Tab 48.6 mg with valsartan 51.4 mg	190.00	56	1	Entresto 49/51
	Tab 97.2 mg with valsartan 102.8 mg		56	✓	Entresto 97/103
	•				

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	•	Manufacturer

#### **⇒SA1905** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III: or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 118

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

### **Antiarrhythmics**

	INCOMPONE LINGUOCINOTINE TELET TO MENVOUS STOTEM, ATTAESHIN	elics, Local, po	aye 110	
	IODARONE HYDROCHLORIDE  Tab 100 mg	2 90	30	✓ Aratac
	Tab 200 mg		30	✓ Aratac
_	Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a	5.25	30	Midlac
	PSOPSO	16 27	10	✓ Max Health
		10.37	10	wax ricalui
	ROPINE SULPHATE			
*	Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a			
	PSO	12.07	10	✓ Hameln S29
				✓ <u>Martindale</u>
DIC	GOXIN			
*	Tab 62.5 mcg - Up to 30 tab available on a PSO		240	✓ Lanoxin PG
*	Tab 250 mcg - Up to 30 tab available on a PSO		240	✓ <u>Lanoxin</u>
*	Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
				Lanoxin Paediatric
				Elixir S29
				✓ Lanoxin S29 S29
DIS	SOPYRAMIDE PHOSPHATE			
$\blacktriangle$	Cap 100 mg	23.87	100	✓ Rythmodan
FLF	ECAINIDE ACETATE			•
<b>A</b>	Tab 50 mg	19.95	60	✓ Flecainide BNM
$\blacktriangle$	Cap long-acting 100 mg		90	✓ Flecainide
				Controlled
				Release Teva
$\blacktriangle$	Cap long-acting 200 mg	61.06	90	✓ Flecainide
				Controlled
				Release Teva
	Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100		ANI \$29 Mexiletine Hydrochloride USP \$29
▲ Cap 250 mg	202.00	100	✓	Teva ©29 Mexiletine Hydrochloride USP ©29
				Teva S29
PROPAFENONE HYDROCHLORIDE  ▲ Tab 150 mg	40.90	50	•	Rytmonorm
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – Retail pha Tab 2.5 mg Tab 5 mg	53.00	100 100	_	Gutron Gutron

# Tab 5 mg ...... SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## **Beta-Adrenoceptor Blockers**

## **Beta Adrenoceptor Blockers**

АТ	ENOLOL	
*	Tab 50 mg4.26	500
	Tab 100 mg7.30	500
*	Oral liq 25 mg per 5 ml21.25	300 ml OP

✓ Mylan Atenolol✓ Mylan Atenolol

✓ Atenolol AFT

✓ Atenolol AFT S29 S29

Restricted to children under 12 years of age.

BISOPROLOL FUMARATE - Brand switch fee payable (Pha	rmacode 2607034) - :	see page 23	5 for details
* Tab 2.5 mg			<ul> <li>Bisoprolol Mylan</li> </ul>
* Tab 5 mg	2.55	90	✓ Bisoprolol Mylan
* Tab 10 mg		90	✓ Bisoprolol Mylan
CARVEDILOL			
* Tab 6.25 mg	2.24	60	<ul><li>Carvedilol Sandoz</li></ul>
* Tab 12.5 mg	2.30	60	<ul><li>Carvedilol Sandoz</li></ul>
* Tab 25 mg		60	<ul><li>Carvedilol Sandoz</li></ul>

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
LABETALOL				
* Tab 100 mg	14.50	100	1	Trandate
* Tab 200 mg	27.00	100	✓	Trandate
* Inj 5 mg per ml, 20 ml ampoule	59.06	5		
	(88.60)			Trandate
* inj 5 mg per ml, 20 ml vial	42.29	1		
	(48.20)			Alvogen S29
METOPROLOL SUCCINATE	( /			- 3-
* Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
* Tab long-acting 25.75 mg		30		Betaloc CR
* Tab long-acting 95 mg		30		Betaloc CR
* Tab long-acting 190 mg		30		Betaloc CR
		00	-	Dotailo Oil
METOPROLOL TARTRATE	F 00	100		Ana Matannalal
* Tab 50 mg		100		Apo-Metoprolol
* Tab 100 mg		60		Apo-Metoprolol
* Tab long-acting 200 mg		28		Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	•	Metoprolol IV Mylan
NADOLOL				
* Tab 40 mg	16.69	100		Apo-Nadolol
* Tab 80 mg	26.43	100	•	Apo-Nadolol
PINDOLOL				
* Tab 5 mg	13.22	100	✓	Apo-Pindolol
* Tab 10 mg	23.12	100	✓	Apo-Pindolol
* Tab 15 mg	33.31	100	✓	Apo-Pindolol
PROPRANOLOL				
* Tab 10 mg	4 64	100	1	Apo-Propranolol
* Tab 40 mg		100		Apo-Propranolol
* Cap long-acting 160 mg		100		Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 l				
Retail pharmacy		500 m	· •	Roxane-
				Propranolol \$29
				i iopianoloi

#### ⇒SA1327 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

#### SOTALOL

*	Tab 80 mg	32.58	500	✓ Mylan
*	Tab 160 mg	10.98	100	✓ Mylan

MoLOL — Subsidy by endorsement		Subsidy (Manufacturer's Price)		Fully bsidised	Brand or Generic
Subsidy by endorsement – Subsidised for patients who were taking timolol prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of timolol.  ₹ Tab 10 mg		\$	Per		Manufacturer
Calcium Channel Blockers	Subsidy by endorsement – Subsidised for patients who endorsed accordingly. Pharmacists may annotate the dispensing of timolol.	prescription as endorsed wh			
MILODIPINE		10.55	100	✓ A	po-Timol
MLODIPINE	Calcium Channel Blockers				
Tab 2.5 mg	Dihydropyridine Calcium Channel Blockers	<b>;</b>			
## Tab 5 mg	MLODIPINE				
ELODIPINE  E Tab long-acting 2.5 mg	F Tab 2.5 mg	1.08	90	✓ V	<u>asorex</u>
ELODIPINE  Tab long-acting 2.5 mg	÷ Tab 5 mg	0.96	90	✓ <u>V</u>	asorex as a second
Tab long-acting 2.5 mg	v	1.19	90	✓ <u>v</u>	<u>asorex</u>
## Tab long-acting 5 mg		1.45	30	<b>√</b> □	lendil FD
## Tab long-acting 10 mg					
## Tab long-acting 10 mg	Tab long-acting 10 mg			_	
18.80   56					
## Tab long-acting 20 mg ## 17.72	Fab long-acting 10 mg	10.63	60		
## Tab long-acting 20 mg				<b>✓</b> A	defin S29
17.72		18.80	56	<b>√</b> T	ensipine MR10 S29
17.72	Fab long-acting 20 mg	9.12	50	✓ N	lylan S29
34.10 100 ✓ Mylan 529 ✓ Adalat Oros ✓ Adefin XL  52.81 100 ✓ Mylan 529 ✓ Adalat Oros ✓ Adefin XL  52.81 100 ✓ Mylan 529 ✓ Adalat Oros ✓ Adefin XL  52.81 100 ✓ Mylan 529 ✓ Adalat Oros ✓ Adefin XL  Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021) ✓ Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021) ✓ Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) ✓ Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021) ✓ Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021) ✓ Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021)  Other Calcium Channel Blockers  □ LTIAZEM HYDROCHLORIDE ✓ Tab 60 mg			100	✓ N	yefax Retard
## Tab long-acting 60 mg	Fab long-acting 30 mg	3.14	30	<b>✓</b> A	dalat Oros
Adalat Oros Adefin XL  52.81 100 Mylan  Adalat Oros Adefin XL  Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 10 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021)  Other Calcium Channel Blockers  MILTIAZEM HYDROCHLORIDE Tab 60 mg Sales Cap long-acting 120 mg Sales Apo-Diltiazem CD Apo-Diltiazem CD Cap long-acting 240 mg Sales Apo-Diltiazem CD Apo-Diltiazem CD			100	✓ N	Ivlan S29
Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021) Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021) Adefin XL Tab long-acting 60 mg to be delisted 1 August 2021)  Other Calcium Channel Blockers  ILTIAZEM HYDROCHLORIDE  E Tab 60 mg	Tab long-acting 60 mg	5.67	30		•
Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021)  Adelin	0 0			<b>✓</b> A	defin XL
Adalat 10 Tab long-acting 10 mg to be delisted 1 August 2021)  Adelin © Tab long-acting 10 mg to be delisted 1 August 2021)  Adalat Oros Tab long-acting 30 mg to be delisted 1 August 2021)  Adalat Oros Tab long-acting 60 mg to be delisted 1 August 2021)  Adelin XL Tab long-acting 60 mg to be delisted 1 August 2021)  Other Calcium Channel Blockers  ILITIAZEM HYDROCHLORIDE  © Tab 60 mg		52.81	100	✓ N	Ivlan \$29
ILTIAZEM HYDROCHLORIDE	Adefin \$29 Tab long-acting 10 mg to be delisted 1 Augus Adalat Oros Tab long-acting 30 mg to be delisted 1 Augus Adalat Oros Tab long-acting 60 mg to be delisted 1 Augus	t 2021) t 2021) t 2021)			
€ Tab 60 mg	Other Calcium Channel Blockers				
€ Tab 60 mg	II TIAZEM HYDROCHI ORIDE				
€ Cap long-acting 120 mg		8.50	100	<b>√</b> ⊓	)ilzem
€ Cap long-acting 180 mg				_	
€ Cap long-acting 240 mg	Cap long-acting 180 mg	50.05			
· · · · · · · · · · · · · · · · · · ·					
	Dilzem Tab 60 mg to be delisted 1 January 2022)				

\* Tab 100 mg ......62.90

PERHEXILINE MALEATE

✓ Pexsig

100

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	I Generic
	\$	Per		Manufacturer
ERAPAMIL HYDROCHLORIDE				
★ Tab 40 mg		100	•	Isoptin
<b>₭</b> Tab 80 mg	11.74	100	•	Isoptin
★ Tab long-acting 120 mg	36.02	100	✓	Isoptin Retard S29
			✓	Isoptin SR
★ Tab long-acting 240 mg	15.12	30	✓	Isoptin SR
k Inj 2.5 mg per ml, 2 ml ampoule − Up to 5 inj available on a				
PSO	25.00	5	•	Isoptin
Centrally-Acting Agents				
CLONIDINE				
א Patch 2.5 mg, 100 mcg per day − Only on a prescription	10.34	4		Mylan
Patch 5 mg, 200 mcg per day — Only on a prescription		4	_	Mylan
Patch 7.5 mg, 300 mcg per day — Only on a prescription		4	_	Mylan
		7	•	<u>, 1411</u>
CLONIDINE HYDROCHLORIDE	0.75	110		Clanidina DNM
k Tab 150 mag		112		Clonidine BNM
* Tab 150 mcg		100	_	Catapres
k Inj 150 mcg per ml, 1 ml ampoule	∠5.96	10	•	<u>Medsurge</u>
METHYLDOPA			_	
* Tab 250 mg		100		Methyldopa Mylan
	52.85	500	•	Methyldopa Mylan
				<b>S29</b> S29
<b>5</b> 1 1				
Diuretics				
Loop Diuretics				
BUMETANIDE				
·	/ O1	30	./	Burinex S29 S29
k Tab 1 mg	16.36	100		Burinex 529 529
₭ Inj 500 mcg per ml, 4 ml vial		5		Burinex
	1.30	5	•	Dulliex
FUROSEMIDE [FRUSEMIDE]	7.04	4 000		A F
★ Tab 40 mg – Up to 30 tab available on a PSO		1,000		Apo-Furosemide
★ Tab 500 mg		50		Urex Forte
	89.48		•	Furosemid-
				Ratiopharm S29
	160.06	100		Eurocomid
	169.96	100	•	Furosemid-
				Ratiopharm S29
K Oral lig 10 mg per ml	11.00	30 ml (	חם . <b>/</b>	Lasix
Oral liq 10 mg per ml  In liq 10 mg per ml .25 ml ampoule.				
<ul><li>Inj 10 mg per ml, 25 ml ampoule</li><li>Inj 10 mg per ml, 2 ml ampoule − Up to 5 inj available on a</li></ul>		6 5		<u>Lasix</u> Furosemide-Baxter
inj to my per mi, z mi ampoule – op to 5 mj avallable on a	1 50 1.15	5		utosettiide-Daxter
Potassium Sparing Diuretics				
i otaooiani opannig bialetios				
MILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	30.00	25 ml (	OP 🗸	Biomed
PLERENONE - Special Authority see SA1728 on the next pa	ge – Retail pharmacy			
Tab 50 mg		30	/	Inspra
Tab 25 mg		30		Inspra
··· - ··· •				

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic	
	\$	Per	1	Manufacturer	
⇒SA1728 Special Authority for Subsidy					

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has heart failure with ejection fraction less than 40%; and
- - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
  - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

METOLAZONE		o opa. aoo	g or opinoriolization.
Tab 5 mg	CBS	1	✓ Metolazone S29
		50	✓ Zaroxolyn S29
SPIRONOLACTONE  ** Tab 05 mg	4.00	100	./ Cuivantin
* Tab 25 mg * Tab 100 mg		100	<ul><li>✓ Spiractin</li><li>✓ Spiractin</li></ul>
Oral liq 5 mg per ml		25 ml OP	✓ <u>Biomed</u>
Potassium Sparing Combination Diuretics			
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
* Tab 5 mg with furosemide 40 mg	8.63	28	✓ Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE			
* Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ Moduretic
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
* Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ Arrow-
			<u>Bendrofluazide</u>
May be supplied on a PSO for reasons other than emergency	•		
* Tab 5 mg	34.55	500	✓ <u>Arrow-</u> Bendrofluazide
			<u> Deliurolluaziue</u>
CHLOROTHIAZIDE			_
Oral liq 50 mg per ml	26.00	25 ml OP	✓ Biomed
CHLORTALIDONE [CHLORTHALIDONE]			41
Tab 25 mg	3.90 6.50	30 50	✓ Igroton S29 ✓ Hygroton
INDAPAMIDE	0.50	30	- <u>riygioton</u>
* Tab 2.5 mg	10.45	90	✓ Dapa-Tabs
Lipid-Modifying Agents			
Fibrates			

✓	fully subsidised
Sol	le Subsidised Supply

✓ Bezalip ✓ Bezalip Retard

90

30

**BEZAFIBRATE** 

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Lipid-Modifying Agents				
ACIPIMOX * Cap 250 mg	21.56	30		Olbetam Olbetam S29 S29
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	1	Colestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN  * Tab 10 mg	9.99 	500 500 500 500 28 28 90 90 90	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Lorstat Lorstat Lorstat Lorstat Pravastatin Mylan Pravastatin Mylan Simvastatin Mylan Simvastatin Mylan Simvastatin Mylan Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors				
EZETIMIBE — Special Authority see SA1045 below — Retail pha  ★ Tab 10 mg	•	30	1	Ezetimibe Sandoz

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
EZETIMIBE WITH SIMVASTATIN - Special Authority see SA104	6 below – Retail pha	rmacy		
Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zi	imybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zi	imybe
Tab 10 mg with simvastatin 40 mg		30	✓ Zi	imybe
Tab 10 mg with simvastatin 80 mg		30	<b>✓</b> Zi	imybe

#### ⇒SA1046 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

\* Oral pump spray, 400 mcg per dose - Up to 250 dose

3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

#### **Nitrates**

|--|

available on a PSO	250 dose OP	✓ Nitrolingual Pump Spray
* Patch 25 mg, 5 mg per day	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day18.62	30	✓ Nitroderm TTS
ISOSORBIDE MONONITRATE		
* Tab 20 mg19.55	100	✓ Ismo 20

K	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
K	Tab long-acting 60 mg	9.25	90	✓ <u>Duride</u>

## **Sympathomimetics**

ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

#### Vasodilators

#### HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 on the next page -		
	Retail pharmacyCBS	1	✓ Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM S29
		100	✓ Onelink S29
*	Inj 20 mg ampoule25.90	5	✓ Apresoline

Subsidy	Fu	lly Brand or	
(Manufacturer's P	Price) Subsidise	ed Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

#### ⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

#### MINOXIDIL

▲ Tab 10 mg	70.00	100	✓ Loniten
NICORANDIL			
▲ Tab 10 mg	25.57	60	✓ <u>Ikorel</u>
▲ Tab 20 mg	32.28	60	✓ Ikorel
PAPAVERINE HYDROCHLORIDE			
* Inj 12 mg per ml, 10 ml ampoule	.217.90	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]			
Tab 400 mg	42.26	50	✓ Trental 400

## **Endothelin Receptor Antagonists**

AMBRISENTAN - Special Authority see SA1702 below - Retail pharmacy		
Tab 5 mg1,550.00	30	✓ Ambrisentan Mylan
Tab 10 mg1,550.00	30	✓ Ambrisentan Mylan

#### ⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ROSENTAN	<ul> <li>Special Authority see</li> </ul>	SA1991 he	low – Retail pharmacy

Tab 62.5 mg	141.00	60	✓ Bosentan Dr
			Reddy's
Tab 125 mg	141.00	60	✓ Bosentan Dr
			Reddy's

#### ⇒SA1991 Special Authority for Subsidy

**Initial application** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 Bosentan is to be used as PAH monotherapy; and
    - 4.1.2 Either:
      - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
      - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or

continued...

<del>-</del>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	` ¢	Por 🗸	Manufacturor

continued...

- 4.2 Both:
  - 4.2.1 Bosentan is to be used as PAH dual therapy; and
  - 4.2.2 Either:
    - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
    - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
- 4.3 Both:
  - 4.3.1 Bosentan is to be used as PAH triple therapy; and
    - 4.3.2 Any of the following:
      - 4.3.2.1 Patient is on the lung transplant list; or
      - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
      - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
      - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

**Renewal** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Bosentan is to be used as PAH monotherapy; and
  - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
  - 2.1 Bosentan is to be used as PAH dual therapy; and
  - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
  - 3.1 Bosentan is to be used as PAH triple therapy; and
  - 3.2 Any of the following:
    - 3.2.1 Patient is on the lung transplant list; or
    - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
    - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
    - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

## **Phosphodiesterase Type 5 Inhibitors**

SILDENAFIL - Special Authority see SA1992 below - Retail pharmacy		
Tab 25 mg	4 4	✓ Vedafil
Tab 50 mg	4 4	✓ Vedafil
Tab 100 mg	0 12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

1 Patient has Raynaud's Phenomenon\*; and

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	

- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 Any of the following:
  - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
  - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
  - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
  - 3.1 PAH is in NYHA/WHO functional class II; or
  - 3.2 PAH is in NYHA/WHO functional class III: or
  - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.2 Fither:
      - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
      - 4.1.2.2 Patient is peri Fontan repair; and
    - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
  - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with \* are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## Prostacvclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retail pharmacy		
Inj 500 mcg vial36.61	1	✓ Veletri
Inj 1.5 mg vial73.21	1	✓ Veletri
SA1606 Chaoial Authority for Subcidy		

#### SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 on the next page - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml .......740.10 30

✓ Ventavis

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from PHARMAC's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

PHARMAC, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

#### **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Fer ✓ Manufacturer

## **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a procomption			
Crm 0.1%	22.89	30 g OP	<ul><li>Differin</li></ul>
Gel 0.1%	22.89	30 g OP	<ul><li>Differin</li></ul>
ISOTRETINOIN - Special Authority see SA2023 below -	- Retail pharmacy	•	
Cap 5 mg	8.14	60	<ul><li>Oratane</li></ul>
Cap 10 mg	13.34	120	✓ Oratane
Cap 20 mg	20.49	120	✓ Oratane

#### ⇒SA2023 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
  - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

**TRFTINOIN** 

Crm 0.5 mg per q − Maximum of 50 g per prescription......13.90 50 g OP ✓ ReTrieve

## **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

## **DERMATOLOGICALS**

	Subsidy	luino) O: I	Fully	Brand or
	(Manufacturer's P \$	rice) Subs Per	idised •	Generic Manufacturer
MUPIROCIN				
Oint 2%	6.60	15 g OP		
	(10.50)		В	actroban
a) Only on a prescription     Not in a sorbination				
b) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID]  Crm 2%	1.50	5 g OP	<b>√</b> E	oban
a) Maximum of 5 g per prescription	1.39	3 y OF	• [	<u>DDall</u>
b) Only on a prescription				
c) Not in combination				
Oint 2%	1.59	5 g OP	✓ <u>F</u>	<u>oban</u>
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER  Crm 1%	10.90	50 g OP	<b>√</b> =	amazine
a) Up to 250 g available on a PSO	10.60	50 y OF	• [	amazme
b) Not in combination				
,				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, page	96			
AMOROLFINE				
a) Only on a prescription				
b) Not in combination				
Nail soln 5%	14.93	5 ml OP	✓ <u>M</u>	<u>ycoNail</u>
CICLOPIROX OLAMINE				
a) Only on a prescription				
b) Not in combination Nail-soln 8%	E 70	7 ml OP	./ A	na Cialaniray
	3.72	7 ml OP	▼ <u>A</u>	po-Ciclopirox
CLOTRIMAZOLE  * Crm 1%	0.77	20 g OP	<b>√</b> ∩	lomazol
a) Only on a prescription		20 y Oi	• 0	ΙσπαΣσι
b) Not in combination				
* Soln 1%	4.36	20 ml OP		
	(7.55)		С	anesten
a) Only on a prescription				
b) Not in combination				
ECONAZOLE NITRATE	4.00	00 - 00		
Crm 1%		20 g OP	D	evend
a) Only on a prescription	(7.48)		Γ	evaryl
b) Not in combination				
Foaming soln 1%, 10 ml sachets	9.89	3		
-	(17.23)		Р	evaryl
a) Only on a prescription				
b) Not in combination				

			DEIW	IATOLOGICALS
	Subsidy (Manufacturer's Pr \$	rice) Per	Fully Subsidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g O	)P 🗸	<u>Multichem</u>
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
* Lotn 2%	4.36	30 ml C	OP	
	(10.03)			Daktarin
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
* Tinct 2%	4.36	30 ml C	OP	
	(12.10)			Daktarin
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>				
Antipruritic Preparations				
CALAMINE  a) Only on a prescription				
b) Not in combination Crm, aqueous, BP	1.26	100 დ	g 🗸	healthE Calamine Aqueous Cream

#### CROTAMITON

- a) Only on a prescription
- b) Not in combination

Crm 10%......3.29

....3.29 20 g OP

✓ Itch-Soothe

ΒP

#### MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Crystals6.92	25 g
29.60	100 g

✓ MidWest

✓ MidWest

## **Corticosteroids Topical**

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

#### Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE		
Crm 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05%2.96	15 g OP	✓ Diprosone
36.00	50 g OP	✓ Diprosone
Oint 0.05% in propylene glycol base4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERATE		
* Crm 0.1%	50 g OP	✓ Beta Cream
* Oint 0.1%3.45	50 g OP	✓ Beta Ointment
* Lotn 0.1%	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE		
* Crm 0.05%	30 g OP	✓ Dermol
* Oint 0.05%2.12	30 g OP	✓ Dermol

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
CLOBETASONE BUTYRATE			
Crm 0.05%	5.38	30 g OP	
	(10.00)		Eumovate
DIFLUCORTOLONE VALERATE			
Fatty oint 0.1%	8 97	50 g OP	
Tady one or 70	(15.86)	00 g 0.	Nerisone
Nerisone Fatty oint 0.1% to be delisted 1 August 2021)	(10.00)		1101100110
,			
HYDROCORTISONE	0.70	100 = OD	. I lively a continuo
★ Crm 1% – Only on a prescription	3.70	100 g OP	✓ <u>Hydrocortisone</u>
			(PSM)
	17.15	500 g	✓ <u>Hydrocortisone</u>
			(PSM)
★ Powder – Only in combination		25 g	✓ ABM
Up to 5% in a dermatological base (not proprietary Top	oical Corticosterio	d – Plain) with c	or without other dermatologica
galenicals			
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only	/ on		
a prescription		250 ml	✓ DP Lotn HC
		200 1111	<u> </u>
HYDROCORTISONE BUTYRATE	0.05	400 00	
Lipocream 0.1%		100 g OP	✓ Locoid Lipocream
Oint 0.1%		100 g OP	Locoid
Milky emul 0.1%	13.70	100 ml OP	✓ Locoid Crelo
METHYLPREDNISOLONE ACEPONATE			
Crm 0.1%	4.46	15 g OP	✓ Advantan
Oint 0.1%	4.46	15 g OP	✓ Advantan
MOMETASONE FUROATE			
Crm 0.1%	1 51	15 g OP	✓ Elocon Alcohol Free
0111 0.170	2.50	50 g OP	✓ Elocon Alcohol Free
Oint 0.1%		15 g OP	✓ Elocon
OII t 0.170	2.90	50 g OP	✓ Elocon
Lotn 0.1%		30 ml OP	✓ Elocon
	0.50	30 1111 01	LIOCOII
TRIAMCINOLONE ACETONIDE			
Crm 0.02%		100 g OP	✓ <u>Aristocort</u>
Oint 0.02%	6.35	100 g OP	✓ Aristocort
Corticosteroids - Combination			
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE (F	TIGIDIC ACIDI		
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP	
Citi 0.1% with souldin idsidate (idsidic acid) 2%		15 y OF	Fusiont
A Mariana of 45 amount of the	(10.45)		Fucicort
a) Maximum of 15 g per prescription			
b) Only on a prescription			
HYDROCORTISONE WITH MICONAZOLE - Only on a presci	ription		
★ Crm 1% with miconazole nitrate 2%	2.00	15 g OP	✓ Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN -	Only on a prescrip	otion	
TO TO CONTINUOUS WITH INVAINABLE OF AND INCOME OF THE	, , ,		/ Discontinuous
Crm 1% with natamyoin 1% and neomyoin sulphate 0.5%	7 75	15 (I L)P	♥ PIMATIICORT
Crm 1% with natamycin 1% and neomycin sulphate 0.5%. Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP 15 g OP	<ul><li>✓ Pimafucort</li><li>✓ Pimafucort</li></ul>

		U	ENWATOLOGICALS
	Subsidy (Manufacturer's F	Price) Subsi Per	Fully Brand or dised Generic  Manufacturer
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	N AND NYSTA	ΓIN	
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g - Only on a prescription		15 g OP	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE  * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u>
* Crm 10% pump bottle	4.52	500 ml OP	Dimethicone 5%  ✓ healthE  Dimethicone 10%
ZINC AND CASTOR OIL  * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM  * Crm		500 g	✓ Basic AquaCream ✓ Boucher ✓ Medco
(Topiderm Crm to be delisted 1 September 2021) CETOMACROGOL	5.75		✓ Topiderm
* Crm BP CETOMACROGOL WITH GLYCEROL		500 g	✓ <u>healthE</u>
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ Boucher ✓ Kenkay Sorbolene ✓ Pharmacy Health Sorbolene with Glycerin
	3.10	1,000 ml OP	✓ ADE ✓ Boucher
# Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION  * Crm	2.19	500 g	✓ <u>O/W Fatty Emulsion</u> <u>Cream</u>
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ <u>healthE</u>
UREA	4.07	400 · OD	/ h = - Wh = 1 l O

100 g OP

✓ healthE Urea Cream

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

#### **DERMATOLOGICALS**

Subsidy (Manufacturer's F \$		Fully dised	Brand or Generic Manufacturer
5.60	1,000 ml		
(11.95)			P Lotion
1.40	250 ml OP		
(4.53)			P Lotion
5.60	1,000 ml		
(20.53)		Д	Alpha-Keri Lotion
(23.91)		В	SK Lotion
1.40	250 ml OP		
(7.73)		Е	BK Lotion
	\$	\$ Per 5.60 1,000 ml (11.95) 1.40 250 ml OP (4.53) 5.60 1,000 ml (20.53) (23.91) 1.40 250 ml OP	\$ Per

#### Other Dermatological Bases

#### **PARAFFIN**

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

#### **Minor Skin Infections**

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

## **Parasiticidal Preparations**

ME.		

		Lotion	
IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy			
Tab 3 mg - Up to 100 tab available on a PSO 17.20	4	✓ Stromectol	

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

200 ml OP

healthE

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

#### ⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or

dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	1	Manufacturer	

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

#### PERMETHRIN

Crm 5%       5.75         Lotn 5%       3.99	30 g OP 30 ml OP	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN	000   OD	/ Daysaidaas
Shampoo 0.5%	200 MI OP	Parasidose

## **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail pharmacy			
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

#### ⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment;
- 2 Patient is not of child bearing potential.

#### RETAMETHASONE DIDRODIONATE WITH CALCIDOTRIO

DETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	52.24	60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	19.95	30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ <u>Midwest</u>

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

				ATOLOGICALO
	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULP	HUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%		75 g OP 30 g OP		gopsoryl TA gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR				
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP		Coco-Scalp Coco-Scalp
a) Maximum of 15 g per prescription b) Note: a maximum of 15 g per prescription and no more the Cream 1%	nan one prescript28.50 almologist or any alid without further	15 g OP y relevant pracer renewal unl	ctitioner ess noti	ified for applications dermatitis, rosacea,
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORES  * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium		n a prescription 500 ml		Pinetarsol
SALICYLIC ACID Powder – Only in combination	18.88	250 g	✓ N ✓ P	Aidwest PSM
<ol> <li>Only in combination with a dermatological base or p</li> <li>With or without other dermatological galenicals.</li> </ol>	proprietary Topic	al Corticostero	oid – Pla	ain or collodion flexible
SULPHUR				
Precipitated – Only in combination	6.35	100 g	✓ N	lidwest

1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain

2) With or without other dermatological galenicals.

Scalp Preparations		
BETAMETHASONE VALERATE  * Scalp app 0.1%	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE  * Scalp app 0.05%	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%3.23	100 ml OP	✓ <u>Sebizole</u>
a) Maximum of 100 ml per prescription     b) Only on a prescription		

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

#### **DERMATOLOGICALS**

Subsidy (Manufacturer's Price) Per

Fully Subsidised Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP

✓ Marine Blue Lotion SPF 50+

**Wart Preparations** 

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

**IMIQUIMOD** 

Crm 5%, 250 mg sachet......21.72 24 ✓ Perrigo

**PODOPHYLLOTOXIN** 

3.5 ml OP ✓ Condyline

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

**Other Skin Preparations** 

Antineoplastics

FLUOROURACIL SODIUM

20 g OP ✓ Efudix

## **GENITO-URINARY SYSTEM**

Sub	bsidy	Fully Brand or
(Manufactu	turer's Price) Subsic	lised Generic
•	\$ Per	<ul> <li>Manufacturer</li> </ul>

Subsidy (Manufacturer's Price)

Fully Subsidised

Brand or Generic

Per Manufacturer

## **Contraceptives - Non-hormonal**

## **Condoms**

	NDOMS			
	49 mm - Up to 144 dev available on a PSO		144	✓ Moments
*	53 mm	0.95	10	✓ <u>Moments</u>
		11.64	144	✓ Moments
	<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>			
	b) Up to 60 dev available on a PSO			
*	53 mm, 0.05 mm thickness	0.95	10	✓ <u>Moments</u>
		11.42	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	53 mm, chocolate, brown	0.95	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	53 mm, strawberry, red	0.95	10	✓ Moments
	•	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
*	56 mm, 0.05 mm thickness	1.30	12	✓ Gold Knight
	· · · · · · · · · · · · · · · · · · ·	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
•	a) Maximum of 60 dev per prescription			<u></u>
	b) Up to 60 dev available on a PSO			
*	56 mm, 0.08 mm thickness	0.97	10	✓ Moments
•	oo min, o.oo min ullowlood	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.01		<u> </u>
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
•	30 mm, 0.00 mm unokilo33, red	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.04	177	Momenta
	b) Maximum of 60 dev per prescription			
*	56 mm, chocolate	1 30	12	✓ Gold Knight
~	50 mm, chocolate	15.57	144	✓ Gold Knight
	a) Un to 60 day available on a DCO	10.07	144	4 dola Kliight
	<ul><li>a) Up to 60 dev available on a PSO</li><li>b) Maximum of 60 dev per prescription</li></ul>			
×	56 mm, strawberry	1 20	12	✓ Cold Knight
*	56 mm, strawberry	15.57	144	✓ Gold Knight
	a) Un to CO day available on a BCO	10.07	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
•	b) Maximum of 60 dev per prescription	1 40	10	. Cald Knight VI
*	60 mm		12 144	✓ Gold Knight XL
		14.87	144	✓ Shield XL
_		17.02		✓ Gold Knight XL

a) Maximumosidisedev per prescription b) substantification a PSO

S29 Unapproved medicine supplied under Section 29

### **GENITO-URINARY SYSTEM**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
* 60 mm (bulk pack)	14.87	144	<b>✓</b> <u>G</u>	old Knight XL

### **Contraceptive Devices**

#### INTRA-UTERINE DEVICE

a) Up to 40 dev available on a PSO

b) Up to 60 dev available on a PSO

b) Only on a PSO

	b) Only on a 1 00		
*	IUD 29.1 mm length × 23.2 mm width	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	1	✓ Choice
			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	1	✓ Choice Load 375

## **Contraceptives - Hormonal**

### **Combined Oral Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to		
	84 tab available on a PSO10.00	84	✓ Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab6.62	84	
	(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

(Marvelon 28 Tab 30 mcg with desogestrel 150 mcg and 7 inert tab to be delisted 1 November 2021)

### **GENITO-URINARY SYSTEM**

) ED
D
) ED
D
(

# **Progestogen-only Contraceptives**

### **⇒SA0500** Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO	.16.50 22.00	84 112	✓ <u>Microlut</u> ✓ <u>Microlut</u>
*	Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO	106.92	1	✓ Jadelle

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a P NORETHISTERONE		1	<b>✓</b> <u>D</u>	epo-Provera	
Tab 350 mcg - Up to 84 tab available on a PSO  Emergency Contraceptives	6.25	84	✓ <u>N</u>	loriday 28	
LEVONORGESTREL  * Tab 1.5 mg		1 Part I c	-	ostinor-1	

# **Antiandrogen Oral Contraceptives**

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

# **Gynaecological Anti-infectives**

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID		
Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.00)	9	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	✓ Micreme
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

# Myometrial and Vaginal Hormone Preparations

	,				
ER	RGOMETRINE MALEATE				
	Inj 500 mcg per ml, 1 ml ampoule - Up	to 5 inj available on a			
	PSO	160.00	5	<ul> <li>DBL Ergometrine</li> </ul>	
OE	ESTRIOL				
*	Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin	
	Pessaries 500 mcg		15	✓ Ovestin	
ОХ	(YTOCIN - Up to 5 inj available on a PS	0			
	Inj 5 iu per ml, 1 ml ampoule	3.98	5	✓ Oxytocin BNM	
	Inj 10 iu per ml, 1 ml ampoule		5	✓ Oxytocin BNM	

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

### GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price)	_	Fully sidised	Brand or Generic	
	\$	Per		Manufacturer	
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj av	ailable on a PSO				
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml	15.00	5	✓ Sy	<u>yntometrine</u>	

## Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

40 test OP ✓ David One Step Cassette

> **Pregnancy Test** ✓ Smith BioMed Rapid **Pregnancy Test**

# **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 107

### 5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy \* Tab 5 mg .......4.81 100 ✓ Ricit

#### ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

# Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

✓ Tamsulosin-Rex

## ⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

# Other Urinary Agents

OXYBUTYNIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking oxybutynin prior to 1 June 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of oxybutynin.

*	Tab 5 mg11./0	500	✓ Apo-Oxybutynin
*	Oral liq 5 mg per 5 ml60.40	473 ml	Apo-Oxybutynin

## **GENITO-URINARY SYSTEM**

	Subsidy	-	ully	Brand or
(N	Manufacturer's Price	) Subsid	ised	Generic
	\$	Per	✓	Manufacturer
POTASSIUM CITRATE				
Oral liq 3 mmol per ml - Special Authority see SA1083 below -	-			
Retail pharmacy	31.80 20	00 ml OP	<b>✓</b> B	iomed

### ⇒SA1083 Special Authority for Subsidy

CODULINA OUTDO TARTERATE

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22	28	✓ Ural
SOLIFENACIN SUCCINATE			_
Tab 5 mg	3.00	30	<ul> <li>Solifenacin Mylan</li> </ul>
Tab 10 mg	5.50	30	✓ Solifenacin Mylan

# **Detection of Substances in Urine**

ORTHO-TOLIDINE	7.50	50 test OP	
* Compound diagnostic sticks	(8.25)	50 lesi OF	Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
	(13.92)		Albustix

# **Obstetric Preparations**

# **Antiprogesterones**

#### **MIFEPRISTONE**

Subsidised on a PSO only if from a Family Planning New Zealand Clinic or an abortion service provider with a DHB contract and the PSO is endorsed with the name of the institution for which the PSO is required.

✓ Mifegyne	1	Tab 200 mg60.00
✓ Mifegyne	3	180 00

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

# Calcium Homeostasis

CALCITONIN		
* Inj 100 iu per ml, 1 ml ampoule121.0	00 5	✓ Miacalcic
CINACALCET - Special Authority see SA1618 below - Retail pharmacy		
Tab 30 mg - Wastage claimable210.	30 28	<ul><li>Sensipar</li></ul>

### ⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g., painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

#### ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial - Special Authority see SA2031 below -1 ✓ Zoledronic acid Retail pharmacy......38.03 Mylan

#### ⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
  - 2.1 Patient has bone metastases or involvement; and
  - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
  - 3.1 Patient has bone metastases or involvement; and
  - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

Subsidy	Fully	y Brand or	
(Manufacturer's Price)	Subsidised	d Generic	
\$	Per 🗸	Manufacturer	

#### continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

# Corticosteroids and Related Agents for Systemic Use

PETAMETHA COME CODII IM DUOCDHATE WITH PETAMETHA COME ACETATE

BE	TAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASO	ONE ACETAT	Έ	
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20	5	
	, ,	(36.96)		Celestone
		, ,		Chronodose
חר	XAMETHASONE			
		0.00	00	/ Daymetheens
*	Tab 0.5 mg – Up to 60 tab available on a PSO		30	✓ <u>Dexmethsone</u>
*	Tab 4 mg – Up to 30 tab available on a PSO		30	✓ <u>Dexmethsone</u>
	Oral liq 1 mg per ml	45.00	25 ml OP	✓ Biomed
DE	XAMETHASONE PHOSPHATE			
	Dexamethasone phosphate injection will not be funded for oral	use.		
*	Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO		10	✓ Dexamethasone
	, 34. ,			Phosphate
				Panpharma
*	Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO	16 27	10	✓ Dexamethasone
*	inj 4 mg per mi, 2 mi ampoule – op to 5 mj avallable on a F30	10.37	10	Phosphate
				<u>Panpharma</u>
	JDROCORTISONE ACETATE			
*	Tab 100 mcg	14.32	100	✓ Florinef
НΥ	DROCORTISONE			
	Tab 5 mg	8 10	100	✓ Douglas
	Tab 20 mg		100	✓ Douglas
	Inj 100 mg vial		1	✓ Solu-Cortef
~	, ,		•	• Join-Corter
	a) Up to 5 inj available on a PSO			
	b) Only on a PSO			
ME	THYLPREDNISOLONE			
*	Tab 4 mg	112.00	100	✓ <u>Medrol</u>
*	Tab 100 mg	194.00	20	✓ Medrol
ME	THYLPREDNISOLONE (AS SODIUM SUCCINATE)			
	Inj 40 mg vial	18 90	1	✓ Solu-Medrol-Act-
	iiij 40 iiig viai	10.00	•	0-Vial
				<u>O-viai</u>
	Inj 125 mg vial	28 90	1	✓ Solu-Medrol-Act-
	111 120 111g viai	20.00	•	0-Vial
				<u>O-Viai</u>
	Inj 500 mg vial	22 78	1	✓ Solu-Medrol-Act-
	11, 555 mg via		•	0-Vial
				<u>O Viui</u>
	Inj 1 g vial	27.83	1	✓ Solu-Medrol
		27.00	•	<u>ooia moaror</u>
IVI	THYLPREDNISOLONE ACETATE	44.40	_	<b>45</b>
	Inj 40 mg per ml, 1 ml vial	44.40	5	✓ Depo-Medrol
PR	EDNISOLONE			
*	Oral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00	30 ml OP	✓ Redipred
	Restricted to children under 12 years of age.			
	• •			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PREDNISONE				
* Tab 1 mg	10.68	500	1	Apo-Prednisone
* Tab 2.5 mg		500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO		500	1	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	29.03	500	✓	Apo-Prednisone
FETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen \$29
			1	AU Synacthen
			1	Synacthen
★ Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Depot
			✓	Synacthene
				Retard S29
FRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	1	Kenacort-A 10
, -, -,	26.62		1	Adcortyl \$29
Inj 40 mg per ml, 1 ml ampoule	11.30	1		Triaver S29
,	51.10	5	_	Kenacort-A 40
	70.62			Kenalog S29

# **Sex Hormones Non Contraceptive**

# **Androgen Agonists and Antagonists**

CYPROTERONE ACETATE			
Tab 50 mg	13.17	50	✓ Siterone
Tab 100 mg	26.75	50	✓ Siterone
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	85.00	1	<ul><li>Depo-Testosterone</li></ul>
TESTOSTERONE ESTERS			
Inj 250 mg per ml, 1 ml	12.98	1	<ul><li>Sustanon Ampoules</li></ul>
TESTOSTERONE UNDECANOATE			
Cap 40 mg	21.00	60	✓ Andriol Testocaps
Inj 250 mg per ml, 4 ml vial	86.00	1	✓ Reandron 1000

# Hormone Replacement Therapy - Systemic

### **Prescribing Guideline**

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

	Subsidy		Fully	Brand or
	(Manufacturer's Pr	ice) Sub	sidised	Generic
	\$	Per	1	Manufacturer
Oestrogens				
Cestrogens				
DESTRADIOL – See prescribing guideline on the previous page				
<b>₭</b> Tab 1 mg	4.12	28 OP		
	(11.10)		E	strofem
<b>₭</b> Tab 2 mg		28 OP		
	(11.10)		_	strofem
Real Patch 100 mcg per 24 hours	7.91	4	✓ C	limara
<ul> <li>a) No more than 1 patch per week</li> </ul>				
b) Only on a prescription				
Patch 50 mcg per 24 hours	7.04	4	✓ C	limara
<ul> <li>a) No more than 1 patch per week</li> </ul>				
b) Only on a prescription				
Patch 25 mcg per day	6.12	8	<b>✓</b> E	stradot
• • •	7.85		<b>✓</b> E	stradiol TDP
				Mylan S29
a) No more than 2 patch per week				•
b) Only on a prescription				
Patch 50 mcg per day	7 04	8	<b>√</b> F	stradot 50 mcg
T dion oo mog por day	9.22	O		stradiol TDP
	5.22		· _	Mylan S29
a). No many than O match man would				IVI y I a I I 329
a) No more than 2 patch per week				
b) Only on a prescription	7.04	•		
Patch 75 mcg per day		8		stradot
	10.60		<b>▼</b> E	stradiol TDP
				Mylan S29
<ul> <li>a) No more than 2 patch per week</li> </ul>				
b) Only on a prescription				
Patch 100 mcg per day	7.91	8	<b>√</b> E	stradot
<ul> <li>a) No more than 2 patch per week</li> </ul>				
b) Only on a prescription				
Climara Patch 100 mcg per 24 hours to be delisted 1 August 2	021)			
Climara Patch 50 mcg per 24 hours to be delisted 1 August 202	1)			
DESTRADIOL VALERATE - See prescribing guideline on the p	,			
★ Tab 1 mg	, ,	84	<b>√</b> D	rogynova
⊁ Tab 2 mg		84	_	rogynova
· ·		U <del>1</del>	, <u>r</u>	i ogynova
DESTROGENS – See prescribing guideline on the previous page		00		
★ Conjugated, equine tab 300 mcg		28	_	
h 0 :	(17.50)	20	Р	remarin
★ Conjugated, equine tab 625 mcg		28	_	
	(17.50)		Р	remarin
Duamantamana				
Progestogens				
MEDROXYPROGESTERONE ACETATE - See prescribing gui	deline on the prev	ious page		
<b>★</b> Tab 2.5 mg		30	<b>✓</b> P	rovera
* Tab 5 mg		100		rovera
⊁ Tab 10 mg		30		rovera
•			-	

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		Subsidy (Manufacturer's Price	) Per	Fully Subsidised	Brand or Generic Manufacturer		
Progestogen and Oestrogen Combined Preparations  Manufacturer  Manufacturer							
OE	STRADIOL WITH NORETHISTERONE - See prescribing gui	ideline on page 80					
*	Tab 1 mg with 0.5 mg norethisterone acetate	, ,	28 OF		Kliovance		
*	Tab 2 mg with 1 mg norethisterone acetate	` '	28 OF	•	Kliogest		
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	,			Kilogest		
	oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40 (18.10)	28 OF		Trisequens		
0	ther Oestrogen Preparations						
ETI *	HINYLOESTRADIOL Tab 10 mcg	17.60	100	1	NZ Medical and Scientific		
-	STRIOL Tab 2 mg	7.00	30	•	Ovestin		
0	ther Progestogen Preparations						
* *	/ONORGESTREL Intra-uterine device 52 mg Intra-uterine device 13.5 mg		1		<u>Mirena</u> Jaydess		
	DROXYPROGESTERONE ACETATE Tab 100 mg RETHISTERONE	116.15	100	✓	Provera HD		
*	Tab 5 mg - Up to 30 tab available on a PSO OGESTERONE	5.49	30	✓	Primolut N		
	Cap 100 mg - Special Authority see SA1609 below - Retail			_			

### ⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 For the prevention of pre-term labour\*; and
- - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

pharmacy......16.50

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour\*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
  - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are unapproved indications.

30

✓ Utrogestan

_		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
T	hyroid and Antithyroid Agents				
CA *	RBIMAZOLE Tab 5 mg	10.80	100		Neo-Mercazole Neo-Mercazole S29 S29
LE'	VOTHYROXINE				
*	Tab 25 mcg	5.55	90	1	Synthroid
*	Tab 50 mcg	1.71	28	✓	Mercury Pharma
		5.79	90	✓	Synthroid
		64.28	1,000	<b>/</b>	Eltroxin
*	Tab 100 mcg	1.78	28	✓	Mercury Pharma
		6.01	90	✓	Synthroid
		66.78	1,000	<b>/</b>	Eltroxin
PR	OPYLTHIOURACIL – Special Authority see SA1199 below – Propylthiouracil is not recommended for patients under the a treatments are contraindicated.		the	patient is pı	regnant and other
	Tab 50 mg	35.00	100	1	PTU \$29

### ⇒SA1199 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

# **Trophic Hormones**

#### **Growth Hormones**

SC	MATROPIN (OMNITROPE) - Special Authority see SA2032 below	v – Retail p	harmacy	
*	Inj 5 mg cartridge	34.88	1	<ul> <li>Omnitrope</li> </ul>
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	.104.63	1	✓ Omnitrope

### ⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

(Manufacturer's Price) Subsidised Generic	
\$ Per 🗸 Manufacturer	

continued...

- children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

**Initial application — (Turner syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

Subsidy (Manufacturer's Po	rice) Per	Fully Subsidised	Brand or Generic Manufacturer	
<del>_</del>				

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

**Initial application** — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price	(Manufacturer's Price) Subsidised		Generic	
\$	Per		Manutacturer	

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and

(Manufacturer's Price)	Subsidised	Generic
\$ Pe	er 🗸	Manufacturer

continued...

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH	Ana	logues
------	-----	--------

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe		1	✓ Teva
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a chil	ld or adolescent a	and is unable	e to tolerate ad

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

(591.68)

Vasopressin Agonists

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

Lucrin Depot 3-month

		Subsidy (Manufacturer's P	rice) Subs	Fully	Brand or Generic
		\$	Per	1	Manufacturer
DE	SMOPRESSIN ACETATE				
	Tab 100 mcg	25.00	30	✓ N	linirin
	Tab 200 mcg	54.45	30	✓ N	linirin
$\blacktriangle$	Nasal drops 100 mcg per ml	39.03	2.5 ml OP	✓ N	linirin
•	Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ □	esmopressin- PH&T
	Inj 4 mcg per ml, 1 ml	67.18	10	✓ N	linirin

# **Other Endocrine Agents**

#### CABERGOI INF

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
<ul><li>Dostinex</li></ul>	2	waived by Special Authority see SA1370 below
✓ Dostinex	8	15.20

### ⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly\*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

$\cap$	$\cap M$	IIFFN	10	$\cap$ IT	DV.	TE
l d	UNV	יוחחוו	v =	L /I I	na.	16

Tab 50 mg	29.84	10	✓ Mylan Clomiphen ©29
METYRAPONE			
Cap 250 mg	558.00	50	Metopirone

**INFECTIONS - AGENTS FOR SYSTEMIC USE** Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 Fskazole S29 ⇒SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription 6 ✓ Vermox 15 ml (7.53)Vermox PRAZIQUANTFI 8 Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 61 b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 229 Cephalosporins and Cephamycins CEFACLOR MONOHYDRATE Cap 250 mg......24.70 100 ✓ Ranbaxy-Cefaclor Grans for oral lig 125 mg per 5 ml - Wastage claimable......3.53 100 ml Ranbaxy-Cefaclor **CEFALEXIN** 20 Cephalexin ABM ✓ Cephalexin ABM 20 Grans for oral lig 25 mg per ml - Wastage claimable.....8.75 100 ml ✓ Cefalexin Sandoz Grans for oral lig 50 mg per ml - Wastage claimable......11.75 100 ml ✓ Cefalexin Sandoz CEFAZOLIN - Subsidy by endorsement Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly. ✓ AFT ✓ AFT CEFTRIAXONE - Subsidy by endorsement a) Up to 10 inj available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

CEFUROXIME AXETIL - Subsidy by endorsement

✓ Ceftriaxone-AFT

✓ Ceftriaxone-AFT

Zinnat

5

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
	Por 🗸	Manufacturor	

#### **Macrolides**

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO	0.93	2	✓ Apo-Azithromycin
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	14.38	15 ml	✓ Zithromax

### **⇒SA1683** Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

#### CLARITHROMYCIN

- a) Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 on the next page
- b) Wastage claimable

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

#### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

#### Fither:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

#### Fither:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

**Initial application — (Helicobacter pylori eradication)** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.

Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral lig 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
a) Up to 300 ml available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP     c) Wastage claimable			,
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg	29.90	100	
	(44.58)		ERA

	Subsidy (Manufacturer's Price \$	e) ; Per	Fully Brand or Subsidised Generic Manufacturer	
BOXITHROMYCIN				
Tab disp 50 mg	8.29	10	✓ Rulide D	
Tab 150 mg	8.28	50	✓ <u>Arrow-</u> Roxithromy	<u>cin</u>
Tab 300 mg	16.33	50	✓ <u>Arrow-</u> <u>Roxithromy</u>	<u>cin</u>
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500	✓ Alphamox	
<ul><li>a) Up to 30 cap available on a PSO</li><li>b) Up to 10 x the maximum PSO quantity for RFPP</li></ul>			<del></del>	
Cap 500 mg	36.98	500	✓ <u>Alphamox</u>	
<ul><li>a) Up to 30 cap available on a PSO</li><li>b) Up to 10 x the maximum PSO quantity for RFPP</li></ul>				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	✓ Alphamox 125	į
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	✓ Alphamox 250	)
a) Up to 300 ml available on a PSO			_	
b) Up to 10 x the maximum PSO quantity for RFPP     Wastage claimable				
Inj 250 mg vial		10	✓ Ibiamox	
Inj 500 mg vial		10	✓ Ibiamox	
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	✓ Ibiamox	
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO		10	✓ Curam Duo 50	0/125
O D 500/405 to be Oale O 4 lede 0004	5.00	20	<ul><li>Augmentin</li></ul>	
Curam Duo 500/125 to be Sole Supply on 1 July 2021				
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r	-	100 ml		
per mla) Up to 200 ml available on a PSO	5.00	100 ml	✓ Augmentin	
b) Wastage claimable				
Grans for oral lig amoxicillin 50 mg with clavulanic acid 12.5 r	ma			
per ml – Up to 200 ml available on a PSO	•	100 ml C	P <b>✓ Curam</b>	
(Augmentin Tab 500 mg with clavulanic acid 125 mg to be deliste				
BENZATHINE BENZYLPENICILLIN	, ,			
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	344.93	10	✓ Bicillin LA	
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.09	10	✓ Sandoz	
, 110g (1				

	Subsidy (Manufacturer's P	rice) Sub	Fully Brand or sidised Generic
	\$	Per	✓ Manufacturer
LUCLOXACILLIN			
Cap 250 mg - Up to 30 cap available on a PSO		250	✓ Staphlex
Cap 500 mg - Up to 30 cap available on a PSO	56.61	500	✓ Staphlex
Grans for oral liq 25 mg per ml	2.29	100 ml	✓ <u>AFT</u>
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>			
b) Wastage claimable			
Grans for oral liq 50 mg per ml	3.68	100 ml	✓ <u>AFT</u>
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
Inj 250 mg vial	17.56	10	✓ Flucloxin
Inj 500 mg vial		10	✓ Flucloxin
Inj 1 g vial - Up to 5 inj available on a PSO	5.70	5	✓ Flucil
HENOXYMETHYLPENICILLIN (PENICILLIN V)			
Cap 250 mg - Up to 30 cap available on a PSO	2.59	50	✓ Cilicaine VK
Cap 500 mg		50	✓ Cilicaine VK
a) Up to 20 cap available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	✓ AFT
a) Up to 200 ml available on a PSO			<del></del>
b) Wastage claimable			
Grans for oral lig 250 mg per 5 ml	3.99	100 ml	✓ AFT
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
ROCAINE PENICILLIN			
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	122.50	5	✓ Cilicaine
iiij 1.5 g iii 5.4 iiii syriiige — Op to 5 iiij avallable oii a F5O	123.30	J	Cilicalile
etracyclines			
DXYCYCLINE			_
Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓ Doxine
INOCYCLINE HYDROCHLORIDE			
Tab 50 mg - Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	5.79	60	
	(12.05)		Mino-tabs
Cap 100 mg	19.32	100	
	(52.04)		Minomycin
SA1355 Special Authority for Manufacturers Price			
itial application from any relevant practitioner. Approvals val	id without further	renewal unles	s notified where the patient ha
sacea.			
ETRACYCLINE - Special Authority see SA1332 below - Reta	il pharmacy		
Tab 250 mg	' '	28	✓ Accord S29
Special Authority for Subsidy		20	- 7100014

### ⇒SA1332 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

-					
	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Su	bsidised	Generic	
	\$	Per	/	Manufacturer	

#### Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

#### **CIPROFLOXACIN**

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

, g			
Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	✓ Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule		10	✓ Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and the Inj 150 mg	ne prescription is er		ordingly. <b>✓ Colistin-Link</b>
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement.	25.00	5	✓ DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urin	nary tract inf	fection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement.	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urin	nary tract inf	fection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.	17.50	10	✓ Pfizer

Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.

87.50

50

✓ Pfizer

MOXIFLOXACIN - Special Authority see SA1740 below - Retail pharmacy

No patient co-payment payable

Tab 400 mg ......42.00 5 ✓ Avelox

### ⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*; and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications: or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
    - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
    - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
    - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;

INFECTI	ONS - A	GENTS F	OR S	SYSTEMIC USE
	bsidy urer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
continued				
2 Mycobacterium avium-intracellulare complex not responding to othe 3 Patient is under five years of age and has had close contact with a Note: Indications marked with * are unapproved indications. Renewal only from a respiratory specialist or infectious disease specialist.	confirmed n	nulti-drug re	esistan	t tuberculosis case.
remains appropriate and the patient is benefiting from treatment.	Appiovais	valiu ioi i	year w	nere the treatment
Initial application — (Mycoplasma genitalium) only from a sexual health specialist. Approvals valid for 1 month for applications meeti All of the following:				the recommendation of a
<ol> <li>Has nucleic acid amplification test (NAAT) confirmed Mycoplasma g</li> <li>Either:</li> </ol>	enitalium* a	and is sym	ptomat	ic; and
<ul><li>2.1 Has tried and failed to clear infection using azithromycin; or</li><li>2.2 Has laboratory confirmed azithromycin resistance; and</li></ul>				
3 Treatment is only for 7 days.				
Initial application — (Penetrating eye injury) only from an ophthalmolog requires prophylaxis following a penetrating eye injury and treatment is for Note: Indications marked with * are unapproved indications.			or 1 mo	nth where the patient
PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmac	v			
Cap 250 mg126		16	<b>✓</b> H	umatin S29
Initial application only from an infectious disease specialist, clinical microl month for applications meeting the following criteria:  Either:  1 Patient has confirmed cryptosporidium infection; or		gastroente	rologis	t. Approvals valid for 1
2 For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microbiologist of applications meeting the following criteria: Either:	r gastroent	erologist.	Approv	rals valid for 1 month for
Patient has confirmed cryptosporidium infection; or     For the eradication of Entamoeba histolyica carriage.				
PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharma	су			
Tab 25 mg48	.00	30	<b>✓</b> D	araprim \$29
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without the following criteria: Any of the following:	urther rene	wal unless	notifie	d for applications meeting

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
  3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]			
Tab 250 mg	34.50	12	✓ Fucidin
SULFADIAZINE SODIUM - Special Authority see SA13	331 on the next page – Retail	pharmacy	
Tab 500 mg	543.20	56	✓ Wockhardt S29

s	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsic	dised	Generic
	\$ Per	✓	Manufacturer

### **⇒SA1331** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

#### **TOBRAMYCIN**

Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	15.00	5 <b>/</b>	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient an	d the prescription is end	dorsed accord	dingly.
* · · · * · · · · · · · · · · · · · · ·			

Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement.......395.00

a) Wastage claimableb) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

#### **TRIMETHOPRIM**

*	Tab 300 mg - Up to 30 tab available on a PSO	16.50	50	✓ TMP
不	1 ab 300 iiiu = 0b io 30 iab available 011 a F30	10.30	50	▼ IIVIF

#### TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]

*	Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - Up	

to 30 tab available on a PSO.......53.96 500 **Trisul** 

100 ml ✓ Deprim

56 dose

✓ Tobramycin BNM

#### VANCOMYCIN - Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

# **Antifungals**

- a) For topical antifungals refer to DERMATOLOGICALS, page 62
- b) For topical antifungals refer to GENITO URINARY, page 75

### **FLUCONAZOLE**

00011112022		
Cap 50 mg2.75	28	Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg12.89	28	✓ Mylan
Powder for oral suspension 10 mg per ml — Special Authority		_ <del>-</del>
see SA1359 below – Retail pharmacy109.34	35 ml	Diflucan
Wastage claimable		

#### ⇒SA1359 Special Authority for Subsidy

**Initial application — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

(Ma	Subsidy	F	ully	Brand or
	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	1	Manufacturer

#### continued...

- 1 Patient is immunocompromised; and
  - 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

#### ITRACONAZOI F

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy	141.80	150 ml OP	Sporanox

#### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

# KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare \$29 ✓ Nizoral \$29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Reta	il pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml		105 ml OP	✓ Noxafil

#### ⇒SA1285 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the

	Subsidy		Fully	Brand or
(Man	ufacturer's Price)	Subsi	dised	Generic
	\$	Per	•	Manufacturer

continued...

following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

#### **TERRINAFINE**

* Tab 250 mg8.15	84	✓ Deolate
Deolate to be Sole Supply on 1 August 2021		
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy		
Tab 50 mg91.00	56	✓ Vttack
Tab 200 mg350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,437.00	70 ml	✓ Vfend

#### ⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is immunocompromised: and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

## **Antimalarials**

	Subsidy	F	ully	Brand or
(Ma	nufacturer's Price)	Subsidis	sed	Generic
	\$	Per	•	Manufacturer

### ⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

### **Antiparasitics**

### **Antiprotozoals**

QUININE	SULPHATE	Ξ

*	Tab 300 mg	61.91	500	✓ Q 300
---	------------	-------	-----	---------

(Q 300 Tab 300 mg to be delisted 1 July 2021)

### **Antitrichomonal Agents**

MF	$\Gamma \Omega C$	ואר	IΠΔ	70	ΙF
IVIE	וחו	ועוכ	IUA	/ ( )	1 -

Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg		10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	32.95	10	✓ Arrow-Ornidazole

## **Antituberculotics and Antileprotics**

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

#### CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

#### CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

### DAPSONE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Tab 25 mg	268.50	100	Dapsone
Tab 100 mg	329.50	100	Dapsone

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
FT.	HAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist	<u> </u>			
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation respiratory physician		iseas	e physicia	n, clinical microbiologist or
	Tab 100 mg	85.73	100	✓	EMB Fatol S29
	Tab 400 mg	49.34	56	✓	Myambutol S29
ISC	DNIAZID - Retail pharmacy-Specialist				
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation microbiologist, dermatologist or public health physician				
*	Tab 100 mg	22.00	100	/	<u>PSM</u>
ISC	ONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist  a) No patient co-payment payable  b) Prescriptions must be written by, or on the recommendation	on of, an internal med	dicine	physician	, paediatrician, clinical
	microbiologist, dermatologist or public health physician	05.54	400		B.//
	Tab 100 mg with rifampicin 150 mg		100 100		Rifinah Rifinah
	Tab 150 mg with rifampicin 300 mg	170.60	100	•	niiiiaii
ΓA	RA-AMINO SALICYLIC ACID — Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation respiratory physician Grans for oral liq 4 g sachet		isease 30		t, clinical microbiologist or
PR	OTIONAMIDE - Retail pharmacy-Specialist				
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation respiratory physician  Tab 250 mg		isease	·	t, clinical microbiologist or
DΥ	RAZINAMIDE – Retail pharmacy-Specialist				
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation respiratory physician			. ,	•
*	Tab 500 mg	59.00	100	✓	AFT-Pyrazinamide
RIF	FABUTIN - Retail pharmacy-Specialist				
	a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommendation gastroenterologist				
*	Cap 150 mg	299.75	30	•	Mycobutin
RIF	FAMPICIN - Subsidy by endorsement				
	<ul> <li>a) No patient co-payment payable</li> <li>b) For confirmed recurrent Staphylococcus aureus infection i antimicrobial based on susceptibilities and the prescriptior Retail pharmacy - Specialist. Specialist must be an intern paediatrician, or public health physician.</li> </ul>	n is endorsed accord	ingly;	can be wa	aived by endorsement -
	Cap 150 mg		100		Rifadin
	Cap 300 mg Oral liq 100 mg per 5 ml		100 60 ml		Rifadin Rifadin

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

## Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 229

### **Hepatitis B Treatment**

ENTECAVIR  * Tab 0.5 mg52.	00 30	✓ Entecavir Sandoz
LAMIVUDINE - Special Authority see SA1685 below - Retail pharmacy		
Tab 100 mg6.	95 28	✓ Zetlam
Oral liq 5 mg per ml270.	00 240 ml 0	OP ✓ Zeffix

### ⇒SA1685 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 104

# **Herpesvirus Treatments**

4 CICL OV/IE

ACICLOVIA			
* Tab dispersible 200 mg	1.60	25	✓ Lovir
* Tab dispersible 400 mg		56	✓ Lovir
* Tab dispersible 800 mg		35	✓ Lovir
VALACICLOVIR			
Tab 500 mg	5.75	30	✓ Vaclovir
Tab 1,000 mg	11.35	30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1993 bele	ow – Retail pharmacy		
Tab 450 mg		60	✓ Valganciclovir
•			Mylan

### **⇒SA1993** Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

#### 1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis: and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or

#### 2 Both:

- 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
- 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist.

Subsidy		ully Brand or	
(Manufacturer's \$	Price) Subsidis Per	sed Generic  Manufacturer	

continued

Approvals valid for 3 months for applications meeting the following criteria:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient is immunocompromised: and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- - 1 Patient is immunocompromised; and
  - 2 Any of the following:
    - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
    - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
    - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

# **Hepatitis C Treatment**

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via PHARMAC's approved direct distribution supply. Further details can be found on PHARMAC's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg .......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable Tab 90 mg with sofosbuvir 400 mg......24.363.46 28 ✓ Harvoni

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per ✔ Manufacturer

continued...

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from PHARMAC's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

PHARMAC, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

### **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL — Subsidy by endorsement; can be waived by Special Authority see SA1994 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the PHARMAC website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate) .......61.15

30 **✓ Teva** 

### **⇒SA1994** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

**Renewal** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and

- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

### **Antiretrovirals**

#### ⇒SA1651 Special Authority for Subsidy

**Initial application** — (**Confirmed HIV**) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsid	ised	Generic
	\$	Per	1	Manufacturer

continued

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

# Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous		•	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	us page – Retail pha	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previo	us page – Retail pha	rmacy	
Tab 200 mg	60.00	60	✓ Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

# Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA16	51 on the previous page - Retail	pharmacy
Tab 300 mg	180.00	60 ✓ Ziagen
Oral liq 20 mg per ml	256.31 240	ml OP ✓ Ziagen

	Cub aids		Fully Prond or
	Subsidy (Manufacturer's Price		Fully Brand or dised Generic
	\$	Per	✓ Manufacturer
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPF	ROXIL - Special Au	thority see S	SA1651 on page 104 – Retail
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox 245 mg (300 mg as a maleate)	ounts as three anti-re	·	
EMTRICITABINE – Special Authority see SA1651 on page 104 – Cap 200 mg	- Retail pharmacy	30	✓ Emtriva
LAMIVUDINE – Special Authority see SA1651 on page 104 – Re Tab 150 mg	etail pharmacy	60	✓ <u>Lamivudine</u> Alphapharm
Oral liq 10 mg per ml	102.50 2	40 ml OP	✓ 3TC
ZIDOVUDINE [AZT] - Special Authority see SA1651 on page 10		/	
Cap 100 mg		100	✓ Retrovir
Oral liq 10 mg per ml		00 ml OP	✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg	) counts as two anti		,
Protease Inhibitors			
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p. Cap 150 mg Cap 200 mg	141.68	armacy 60 60	✓ <u>Teva</u> ✓ <u>Teva</u>
DARUNAVIR – Special Authority see SA1651 on page 104 – Re Brand switch fee payable (Pharmacode 2607026) - see page Tab 400 mg	235 for details	60	✓ Darunavir Mylan
Tab 600 mg	196.65	60	✓ Darunavir Mylan
LOPINAVIR WITH RITONAVIR — Special Authority see SA1651  Tab 100 mg with ritonavir 25 mg  Tab 200 mg with ritonavir 50 mg  Oral liq 80 mg with ritonavir 20 mg per ml	183.75 463.00	ail pharmacy 60 120 00 ml OP	<ul><li>✓ Kaletra</li><li>✓ Kaletra</li><li>✓ Kaletra</li></ul>
RITONAVIR - Special Authority see SA1651 on page 104 - Reta Tab 100 mg		30	✓ <u>Norvir</u>
Strand Transfer Inhibitors			
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 - Tab 50 mg		30	✓ Tivicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 o Tab 400 mg Tab 600 mg	1,090.00	pharmacy 60 60	✓ Isentress ✓ Isentress HD

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

## **Immune Modulators**

#### Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

#### **Criteria for Treatment**

- 1) Diagnosis
  - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
  - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
  - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

#### **Exclusion Criteria**

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10<sup>9</sup>) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

#### Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

#### **Exit Criteria**

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: PHARMAC will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at PHARMAC on 0800-023-588 option 4.

#### ✓ Pegasys

#### ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV: or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

#### Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	•	Manufacturer

continued...

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C. genotype 1: and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed: or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*: or
- 2 All of the following:

### **INFECTIONS - AGENTS FOR SYSTEMIC USE**

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Patient has a myeloproliferative disorder\*; and
- 2.2 Patient is intolerant of hydroxyurea; and
- 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Fither:
  - 3.1 Patient has a cutaneous T cell lymphoma\*: or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Fither:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with \* are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

**Initial application** — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

**Renewal — (post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

## **Urinary Tract Infections**

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg - Wastage claimable Macrobid to be Sole Supply on 1 August 2021	86.40	100	✓ Macrobid
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	✓ Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated uri	inary tract infectio	on that is unre	esponsive to a first line agent or

with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	Jupolaloca J	Manufacturer
	Ψ	1 61		Manuacturei
And the Literature of the Control of				
Anticholinesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	19.60	10	1	Juno S29
, , , ,	29.40		1	Max Health
	98.00	EΩ		AstraZeneca
	98.00	50	V	Astrazeneca
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.70	100	1	Mestinon
- Tab oo mg		100	•	Mestinon
New Observation And the General Laws December				
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM				
* Tab EC 25 mg	1.23	50	✓	Diclofenac Sandoz
* Tab 50 mg dispersible		20	1	Voltaren D
* Tab EC 50 mg		50	1	Diclofenac Sandoz
* Tab long-acting 75 mg		500		Apo-Diclo SR
* Tab long-acting 100 mg		500		Apo-Diclo SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a	ı PSO 13.20	5	✓	Voltaren
* Suppos 12.5 mg	2.04	10	1	Voltaren
* Suppos 25 mg		10	1	Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
•				
* Suppos 100 mg	7.00	10	•	Voltaren
IBUPROFEN				
* Tab 200 mg	21.40	1.000	1	Relieve
•		,		
* Tab long-acting 800 mg		30		Ibuprofen SR BNM
* Oral liq 20 mg per ml	1.88	200 m	/	<u>Ethics</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	1	Oruvail SR
* Cap long-acting 200 mg	12.07	20	•	Oruvan Sh
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
· · · · · · · · · · · · · · · · · · ·	(9.16)			Ponstan
	0.50	20		Tonotan
		20		5 .
	(5.60)			Ponstan
NAPROXEN				
* Tab 250 mg	32.60	500	1	Noflam 250
•				
* Tab 500 mg		250		Noflam 500
* Tab long-acting 750 mg	6.16	28	•	Naprosyn SR 750
* Tab long-acting 1 g	8.21	28	•	Naprosyn SR 1000
SULINDAC				
	a			
* Tab 100 mg		56		Mylan S29
* Tab 200 mg	15.10	50	✓	Aclin
- -	16.91	56	/	Sulindac Mylan S29
(Adin Tah 200 mg to be delicted 1 January 2022)		-	_	
(Aclin Tab 200 mg to be delisted 1 January 2022)				
(Sulindac Mylan S29) Tab 200 mg to be delisted 1 January 202	22)			
TENOXICAM				
	0.15	100		Tileatil
* Tab 20 mg		100		Tilcotil
* Inj 20 mg vial	9.95	1	•	AFT

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	I Generic
NSAIDs Other				
ELECOXIB				
Cap 100 mg	5.80	60		Celecoxib Pfizer
Cap 200 mg	2.30	30		Celebrex
	3.30		✓	Celecoxib Pfizer
Topical Products for Joint and Muscular Pain				
CAPSAICIN				
Crm 0.025% - Special Authority see SA1289 below - Retail				
pharmacy	9.75	45 g C	)P 🗸	Zostrix
Antirheumatoid Agents				
IYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systemic suppression, relevant dermatological conditions (cutaneous for mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmor Pharmacists may annotate the prescription as endorsed wher hydroxychloroquine. Note: Indication marked with a * is an u  * Tab 200 mg	orms of lupus and li nary)*, and the pres te there exists a rec napproved indicati	chen p cription cord of	olanus, cut n is endor prior disp	aneous vasculitides and sed accordingly.
EFLUNOMIDE				<u> </u>
Tab 10 mg	6.00	30	1	Arava
Tab 20 mg		30		Arava
PENICILLAMINE				
Tab 125 mg	67.23	100	/	D-Penamine
Tab 250 mg		100		D-Penamine
- a.5 _ 5 0 mg				
Drugs Affecting Bone Metabolism				
Alendronate for Osteoporosis				

# ALENDRONATE SODIUM

* Tab 70 mg	.2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5 600 iu	1.51	4	✓ Fosamax Pli

### **Other Treatments**

**⇒SA1777** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

	Subsidy	F	ully	Brand or
(Ma	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	✓	Manufacturer

continued...

#### All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

	Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
	Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
	Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RA	LOXIFENE HYDROCHLORIDE - Special Authority see SA1779 or	the next page	- Retail pha	armacy
*	Tab 60 mg	53.76	28	✓ Evista

Subsidy (Manufacturer's Price) \$ Pe

Subsidised Per

Fully

Brand or Generic Manufacturer

### ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### RISEDRONATE SODIUM

### **⇒SA1139** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily;

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per •	Manufacturer

continued...

zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

#### ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

#### ⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:
Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

**Initial application — (Underlying cause - glucocorticosteroid therapy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
  - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

(Ma	Subsidy anufacturer's Price)	Subsidi Per	ully	Brand or Generic Manufacturer
	<u></u>	Per		Manufacturer

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
  - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
  - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene: and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	<ul> <li>Manufac</li> </ul>	turer

#### continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

## Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA19	63 below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	<ul> <li>Benzbromaron AL</li> </ul>
			100 829

#### ⇒SA1963 Special Authority for Subsidy

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

### COLCHICINE

* Tab 500 mcg	9.58	100	Colgout
FEBUXOSTAT - Special Authority see SA1996 below - Reta	ail pharmacy		
Tab 80 mg	39.50	28	✓ Adenuric
Tab 120 mg	39.50	28	✓ Adenuric

#### ⇒SA1996 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with gout: and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
  - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Both:

1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and

\$ Per ✔ Manufacturer	Subsidy (Manufacturer's Price)	Fully Subsidised	
	\$	Per 🗸	Manufacturer

continued...

2 Patient has a documented history of allopurinol intolerance.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

#### PROBENECID

### Muscle Relaxants

BACLOFE	

*	Tab 10 mg4.20	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patients where oral ar	ntispastic ag	ents have been ineffective or have
	caused intolerable side effects and the prescription is endorsed according	ly.	

Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement............372.98 5 Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

#### DANTDOLENE

Cap 25 mg97.50	100	✓ Dantrium
Cap 50 mg77.00	100	<ul><li>✓ Dantrium S29 S29</li><li>✓ Dantrium</li></ul>
ORPHENADRINE CITRATE		
Tab 100 mg18.54	100	✓ Norflex

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

## **Agents for Parkinsonism and Related Disorders**

## **Dopamine Agonists and Related Agents**

AMANTADINE HYDROCHLORIDE  ▲ Cap 100 mg38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE		
▲ Inj 10 mg per ml, 2 ml ampoule59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	5	✓ Movapo

#### BROMOCRIPTINE MESYLATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of bromocriptine mesylate.

	prior dispensing of bromocriptine mesylate.			
*	Tab 2.5 mg	32.08	100	✓ Apo-Bromocriptine
ΕN	TACAPONE			
$\blacktriangle$	Tab 200 mg	22.00	100	✓ Entapone
LE	VODOPA WITH BENSERAZIDE			
*	Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
*	Cap 50 mg with benserazide 12.5 mg	13.75	100	✓ Madopar 62.5
*	Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
*	Cap long-acting 100 mg with benserazide 25 mg	22.85	100	✓ Madopar HBS
*	Cap 200 mg with benserazide 50 mg		100	✓ Madopar 250
LE	VODOPA WITH CARBIDOPA			
*	Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
*	Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
	Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
	AMIPEXOLE HYDROCHLORIDE			
$\blacktriangle$	Tab 0.25 mg	6.12	100	✓ Ramipex
$\blacktriangle$	Tab 1 mg		100	✓ Ramipex
RΩ	PINIROLE HYDROCHLORIDE			<del></del>
<b>A</b>	Tab 0.25 mg	2.85	84	✓ Ropin
	•	3.39	100	✓ Mylan S29
$\blacktriangle$	Tab 1 mg	3.95	84	✓ Ropin
	<b>v</b>	4.70	100	✓ Mylan \$29
$\blacktriangle$	Tab 2 mg	5.48	84	✓ Ropin
$\blacktriangle$	Tab 5 mg		84	✓ Ropin
SF	LEGILINE HYDROCHLORIDE			<del></del> _
	Tab 5 mg	22.00	100	✓ Apo-Selegiline S29 S29
TO	LCAPONE			
<b>A</b>	Tab 100 mg	152.38	100	✓ Tasmar

## **Anticholinergics**

RENZ∆TROPINE ME	CVIATE

Tab 2 mg	9.59	60	✓ Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓ Phebra
a) Un to 10 ini queilable on a DCO			

- a) Up to 10 inj available on a PSO
- b) Only on a PSO

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	•	Kemadrin
Agents for Essential Tremor, Chorea and Relate	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	-	56	<b>√</b>	Rilutek
►SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialis following criteria: All of the following:				
1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following:  5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow.				e initial application; and
Renewal from any relevant practitioner. Approvals valid for 18 m All of the following:  1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following:  3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow.	onths for applications	s mee	ting the fo	llowing criteria:
TETRABENAZINE Tab 25 mg	91.10	112	✓	<u>Motetis</u>
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE]  Gel 2%, tube - Subsidy by endorsement	dministration and the	30 ml prese 10	cription is	Xylocaine 2% Jelly endorsed accordingly. Instillagel Lido

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

a) Up to 5 each available on a PSO

accordingly.

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price	e) Subs	Fully idised	
LIDOGANIE II JONGGANIET LIVERDOGUI ORIDE	Ψ	Геі		iviariuiacturei
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%		200 ml	•	Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO	8.75	25	•	Lidocaine-Baxter
			1	Lidocaine-Claris
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	` ,	25	1	Lidocaine-Claris
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO		5		
ing 176, 20 mi ampoule— Op to 0 mj available on a 1 00	(20.00)	Ü		Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO	\ /	5	./	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO		5	•	<u>Lidocaine-Claris</u>
(Lidocaine-Claris Inj 1%, 5 ml ampoule to be delisted 1 January 2	2022)			
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	102 22	10	1	Pfizer
	103.32	10	•	FIIZEI
<ul> <li>a) Up to 5 each available on a PSO</li> </ul>				

## **Topical Local Anaesthetics**

### ⇒SA0906 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 above –	Retail phar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Retai	I pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

## **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

## **Non-opioid Analgesics**

ASPIRIN  * Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diab accordingly.	oetic periphera	l neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price)	Fully Subsidised		,	
	(Manufacturer 3 i fice)	Per	√ v	Manufacturer	
PARACETAMOL					
Tab 500 mg - blister pack	0.50	20	<b>✓</b> N	/ledco	
•			<b>✓</b> F	harmacy Health	
	1.12		<b>√</b> E	thics Paracetamol	
				Classic	
	2.48	100	<b>√</b> F	harmacy Health	
	11.75	96	<b>√</b> F	Panadol Mini Caps	
	24.82	1,000	<b>√</b> F	aracetamol .	
				Pharmacare	
			<b>✓</b> F	harmacare	

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
  regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
  annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

*	Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓ Paracare
	a) Up to 200 ml available on a PSO			
	b) Not in combination			
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double
				<u>Strength</u>
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3.29	10	✓ Gacet
*	Suppos 250 mg	3.79	10	✓ Gacet
*	Suppos 500 mg	12.40	50	✓ Gacet
(Pa	aracetamol Pharmacare Tab 500 mg - bottle pack to be o	delisted 1 December 20	021)	

## **Opioid Analgesics**

CODEINE PHOSPHATE - Safety medicine; prescriber may dete	rmine dispensing	frequency	
Tab 15 mg		100	✓ PSM
Tab 30 mg	7.45	100	✓ PSM
Tab 60 mg	14.25	100	✓ PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ DHC Continus

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic  Manufacturer
ENTANYL		1 01	- Manadalai
a) Only on a controlled drug form			
b) No patient co-payment payable			
<ul><li>c) Safety medicine; prescriber may determine dispensin</li></ul>	a frequency		
Inj 50 mcg per ml, 2 ml ampoule		10	✓ Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	✓ Boucher and Muir
Patch 12.5 mcg per hour		5	✓ Fentanyl Sandoz
Patch 25 mcg per hour		5	✓ Fentanyl Sandoz
Patch 50 mcg per hour		5	✓ Fentanyl Sandoz
Patch 75 mcg per hour		5	✓ Fentanyl Sandoz
Patch 100 mcg per hour		5	✓ Fentanyl Sandoz
IETHADONE HYDROCHLORIDE		·	
a) Only on a controlled drug form     No notice to a new root to be a least of the controlled drug form			
b) No patient co-payment payable	a fraguenou		
c) Safety medicine; prescriber may determine dispensin		o of th	a channet form available
d) Extemporaneously compounded methadone will only	ne reminursed at the fate	ะ บเ เก	e cheapest form available
(methadone powder, not methadone tablets).	rd Formulae name 007		
e) For methadone hydrochloride oral liquid refer Standar		10	✓ Methatabs
Tab 5 mg		10	
Oral liq 2 mg per ml		200 m	
Oral lig 5 mg per ml		200 m	
Oral liq 10 mg per ml		200 m	
Inj 10 mg per ml, 1 ml	61.00	10	✓ AFT
ORPHINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
<ul> <li>Safety medicine; prescriber may determine dispensin</li> </ul>	g frequency		
Oral liq 1 mg per ml	9.28	200 m	✓ <u>RA-Morph</u>
Oral liq 2 mg per ml	16.24	200 m	l ✓ <u>RA-Morph</u>
Oral liq 5 mg per ml	19.44	200 m	✓ Ordine S29
			✓ RA-Morph
Oral liq 10 mg per ml	27.74 2	200 m	
Oral liq 10 mg per ml	27.74 2	200 m	
	27.74	200 m	✓ Ordine \$29
IORPHINE SULPHATE	27.74 2	200 m	✓ Ordine \$29
IORPHINE SULPHATE  a) Only on a controlled drug form	27.74 2	200 m	✓ Ordine \$29
IORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable		200 m	✓ Ordine \$29
IORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin	g frequency		✓ Ordine \$29 ✓ <u>RA-Morph</u>
IORPHINE SULPHATE  a) Only on a controlled drug form  b) No patient co-payment payable  c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency 2.80	10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency 2.80 5.52	10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg Tab immediate-release 20 mg Cap long-acting 10 mg	g frequency 2.80 5.52 2.05	10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg Tab immediate-release 20 mg Cap long-acting 10 mg Cap long-acting 30 mg	g frequency2.805.522.053.00	10 10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg Tab immediate-release 20 mg Cap long-acting 10 mg Cap long-acting 30 mg Cap long-acting 60 mg	g frequency2.805.522.053.006.12	10 10 10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency2.805.522.053.006.127.13	10 10 10 10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg Tab immediate-release 20 mg Cap long-acting 10 mg Cap long-acting 30 mg Cap long-acting 60 mg	g frequency2.805.522.053.006.127.13	10 10 10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ DBL Morphine
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency	10 10 10 10 10 10 5	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ DBL Morphine Sulphate
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency	10 10 10 10 10 10	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ DBL Morphine Sulphate ✓ DBL Morphine
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency	10 10 10 10 10 10 5	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ DBL Morphine Sulphate ✓ DBL Morphine Sulphate
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency	10 10 10 10 10 10 5	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Esion ✓ m-Esion ✓ m-Esion ✓ DBL Morphine Sulphate ✓ DBL Morphine Sulphate ✓ DBL Morphine
AORPHINE SULPHATE  a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensin Tab immediate-release 10 mg	g frequency	10 10 10 10 10 10 5	✓ Ordine S29 ✓ RA-Morph  ✓ Sevredol ✓ Sevredol ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ m-Eslon ✓ DBL Morphine Sulphate ✓ DBL Morphine Sulphate

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	
OXYCODONE HYDROCHLORIDE	<u> </u>			
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency	uency			
Tab controlled-release 5 mg	2.15	20	1	Oxycodone Sandoz
	3.01	28	✓	Oxycodone Sandoz
				<b>S29</b> S29
Tab controlled-release 10 mg	2.15	20	✓	Oxycodone Sandoz
•	3.23	30	✓	Oxycodone Sandoz
				S29 S29
	5.38	50	/	Oxycodone Sandoz
	0.00	00		S29 S29
	10.75	100	1	Oxycodone Sandoz
	10.75	100	•	•
T	0.45		,	S29 S29
Tab controlled-release 20 mg		20		Oxycodone Sandoz
	5.38	50	•	Oxycodone Sandoz
				<b>S29</b> S29
	10.75	100	✓	Oxycodone Sandoz
				<b>S29</b> S29
Tab controlled-release 40 mg	3.20	20	✓	Oxycodone Sandoz
Tab controlled-release 80 mg		20	✓	Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg	3.32	20	✓	OxyNorm
Cap immediate-release 20 mg		20	✓	OxyNorm
Oral liq 5 mg per 5 ml		250 m	nl 🗸	OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5	✓	OxyNorm
Inj 10 mg per ml, 2 ml ampoule	14.36	5	1	OxyNorm
Inj 50 mg per ml, 1 ml ampoule	30.60	5	1	OxyNorm
PARACETAMOL WITH CODEINE - Safety medicine; prescriber m	av determine disn	ensin	a frequenc	·V
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1.000		Paracetamol +
Tab paradotanio dos nig mar dodonio priospriato e niginimimi	20.0	.,000		Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				(10.000)
a) Only on a controlled drug form     b) No period on payment payable				
b) No patient co-payment payable	longy			
c) Safety medicine; prescriber may determine dispensing frequency	•	10	./	PSM
Tab 50 mgInj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS		5		DBL Pethidine
ing 50 mg per mi, i mi ampoule – op to 5 mg available on a F5	J29.00	5	•	Hydrochloride
Ini E0 ma nor ml. 0 ml amnoula. Lin to E ini quailable on a DC	0 20 70	_	./	
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS	J3U.12	5	•	DBL Pethidine
				Hydrochloride
TRAMADOL HYDROCHLORIDE	. =-		_	
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.80	100	✓	Arrow-Tramadol

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Subsidi Per	Fully Brand or ised Generic  Manufacturer
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE – Safety medicine; prescriber may determine d Tab 10 mg Tab 25 mg Tab 50 mg  CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescri Tab 10 mg Tab 25 mg  DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by en	2.491.512.51 ber may determine di13.999.46 dorsement	100 100 100 ispensing fre 100 100	Arrow-Amitriptyline Arrow-Amitriptyline Arrow-Amitriptyline Arrow-Amitriptyline equency Apo-Clomipramine Apo-Clomipramine
Safety medicine; prescriber may determine dispensing freb) Subsidy by endorsement – Subsidised for patients who we 2019 and the prescription is endorsed accordingly. Pharmal exists a record of prior dispensing of dosulepin [dothiepin] Tab 75 mg	ere taking dosulepin   macists may annotate   hydrochloride. 3.85		
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber Tab 10 mg Tab 25 mg	5.48 10.96	nsing freque 50 100 50	ency  Tofranil  Tofranil  Tofranil
MAPROTILINE HYDROCHLORIDE – Subsidy by endorsement  a) Safety medicine; prescriber may determine dispensing fre  b) Subsidy by endorsement – Subsidised for patients who w  2020 and the prescription is endorsed accordingly. Pharr  exists a record of prior dispensing of maprotiline hydrochl  Tab 75 mg	equency ere taking maprotiline nacists may annotate oride.		
(Ludiomil Tab 75 mg to be delisted 1 August 2021)	21.01	30	✓ Ludiomil
(Ludiomil Tab 75 mg to be delisted 1 August 2021)  NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; prescriptab 10 mg	2.44	lispensing fr 100 180	requency  Norpress Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective		
TRANYLCYPROMINE SULPHATE Tab 10 mg	12.85 22.94 45.88 96.00	28 50 100	✓ Parnate S29 S29 ✓ Parnate ✓ Parnate S29 S29 ✓ Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE  * Tab 150 mg  * Tab 300 mg		60 60	✓ <u>Aurorix</u> ✓ <u>Aurorix</u>

			IVEII V	7000 OTOTEW
	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE  * Tab 20 mg	1.52	84	<b>√</b> <u>P</u> \$	SM Citalopram
* Tab 10 mg	1.07	28		scitalopram (Ethics)
	1.40		✓ Es	scitalopram- Apotex
* Tab 20 mg	1.92	28		scitalopram (Ethics)
	2.49			scitalopram- Apotex
(Escitalopram-Apotex Tab 10 mg to be delisted 1 October 2021) (Escitalopram-Apotex Tab 20 mg to be delisted 1 October 2021)				
FLUOXETINE HYDROCHLORIDE  * Tab dispersible 20 mg, scored – Subsidy by endorsement  Subsidised by endorsement	1.98	30	<b>✓</b> FI	uox
<ol> <li>When prescribed for a patient who cannot swallow accordingly; or</li> <li>When prescribed in a daily dose that is not a multipendorsed. Note: Tablets should be combined with</li> </ol>	ple of 20 mg in which	case th	e prescripti	ion is deemed to be
Cap 20 mg	2.91	84	<b>✓</b> Fl	uox
PAROXETINE				
* Tab 20 mg		30		axtine
(Paxtine Tab 20 mg to be delisted 1 January 2022)	3.61	90	✓ Lo	oxamine
SERTRALINE				
* Tab 50 mg	0.92	30		etrona etrona AU
	3.05	90		row-Sertraline
* Tab 100 mg	1.61	30		etrona etrona AU
	5.25	90		rrow-Sertraline
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg		30 30	_	oo-Mirtazapine oo-Mirtazapine
Tab 45 mg VENLAFAXINE	3.40	30	▼ A	оо-тип тагарите
* Cap 37.5 mg	6.38	84		nlafax XR
* Cap 75 mg	8.11	84	✓ Er	ılafax XR

84

✓ Enlafax XR

\* Cap 150 mg......11.16

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM – Safety medicine; prescriber may determine dis Inj 1 mg per ml, 1 ml(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)		5	<b>✓</b> R	ivotril
DIAZEPAM – Safety medicine; prescriber may determine dispen Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO b) Only on a PSO	23.66	5	<b>✓</b> H	ospira
<ul> <li>c) PSO must be endorsed "not for anaesthetic procedur Rectal tubes 5 mg - Up to 5 tube available on a PSO</li> <li>PARALDEHYDE</li> </ul>		5	✓ S	tesolid

## **Control of Epilepsy**

PHENYTOIN SODIUM

\* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO ....88.63

\* Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on a

control of =phopo,			
CARBAMAZEPINE			
* Tab 200 mg	14.53	100	✓ Tegretol
* Tab long-acting 200 mg	16.98	100	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg		100	Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CLOBAZAM - Safety medicine; prescriber may determine dis	spensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM - Safety medicine; prescriber may determine	e dispensing frequer	ncy	
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml		200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pre	gabalin		
* Cap 100 mg	2.65	100	✓ Apo-Gabapentin
* Cap 300 mg	4.07	100	✓ Apo-Gabapentin
* Cap 400 mg	5.64	100	✓ Apo-Gabapentin
LACOSAMIDE - Special Authority see SA1125 on the next p	age - Retail pharm	acy	
▲ Tab 50 mg	25.04	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
	200.24	56	✓ Vimpat
▲ Tab 150 mg		14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

✓ AFT S29

✓ Hospira

✓ Hospira

5

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

#### ⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE			
▲ Tab dispersible 2 mg	55.00	30	✓ Lamictal
▲ Tab dispersible 5 mg	50.00	30	✓ Lamictal
* Tab dispersible 25 mg	2.76	56	✓ Logem
* Tab dispersible 50 mg	3.31	56	✓ Logem
* Tab dispersible 100 mg	4.40	56	✓ Logem
LEVETIRACETAM			
Tab 250 mg	4.99	60	✓ Everet
Tab 500 mg		60	✓ Everet
Tab 750 mg		60	✓ Everet
Tab 1,000 mg		60	✓ Everet
Oral liq 100 mg per ml		300 ml OP	✓ Levetiracetam-AFT
PHENOBARBITONE			
For phenobarbitone oral liquid refer Standard Formulae, pag	ie 237		
* Tab 15 mg		500	✓ PSM
* Tab 30 mg		500	✓ PSM
PHENYTOIN SODIUM			<del></del>
* Tab 50 mg	75.00	200	✓ Dilantin Infatab
9		200	✓ Dilantin
Cap 30 mg Cap 100 mg		200	✓ Dilantin
* Oral lig 30 mg per 5 ml		500 ml	✓ Dilantin
	22.03	300 IIII	Dilanun
PREGABALIN			
Note: Not subsidised in combination with subsidised gabape			4 D
* Cap 25 mg		56	✓ Pregabalin Pfizer
* Cap 75 mg		56	✓ Pregabalin Pfizer
* Cap 150 mg	4.01	56	✓ Lyrica
ut. 0 000	7.00		✓ Pregabalin Pfizer
* Cap 300 mg	7.38	56	Pregabalin Pfizer
PRIMIDONE			
* Tab 250 mg	17.25	100	Apo-Primidone
	62.00	200	✓ Mysoline S29 S29
(Mysoline S29 S29 Tab 250 mg to be delisted 1 July 2021)			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

### **NERVOUS SYSTEM**

	Subsidy		Fully	Brand or	
	(Manufacturer's Pric	e)	Subsidised	Generic	
	\$	Per	1	Manufacturer	
SODIUM VALPROATE					
Tab 100 mg	13.65	100	✓	Epilim Crushable	
Tab 200 mg EC	27.44	100	✓	Epilim	
Tab 500 mg EC	52.24	100	✓	Epilim	
* Oral lig 200 mg per 5 ml	20.48	300 m	<b>/</b>	Epilim S/F Liquid	
, ,,			✓	Epilim Syrup	
* Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV	
STIRIPENTOL - Special Authority see SA1330 below - Retail p	harmacy				
Cap 250 mg	509.29	60	/	Diacomit S29	
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29	

#### ⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

## TOPIRAMATE

▲ Tab 25 mg	11.07	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	26.04		✓ Topamax
▲ Tab 50 mg	18.81	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	44.26		✓ Topamax
▲ Tab 100 mg	31.99	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	75.25		✓ Topamax
▲ Tab 200 mg	55.19	60	✓ Arrow-Topiramate
·			✓ Topiramate Actavis
	129.85		✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg	26.04	60	✓ Topamax
VIGABATRIN - Special Authority see SA1997 below - F	Retail pharmacy		
▲ Tab 500 mg		100	✓ Sabril

### ⇒SA1997 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and
    - 1.2.2 Either:
      - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
      - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

## **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

A	B # ! !	
ACIITE	Midrain	e Treatment
,,,,,,,,,	migrani	o iloutilloit

RIZATRIPTAN Tab orodispersible 10 mg3.65	30	✓ Rizamelt
SUMATRIPTAN Tab 50 mg24.44	100	✓ Apo-Sumatriptan
Tab 100 mg	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription34.00	2 OP	✓ <u>Imigran</u>

### **Prophylaxis of Migraine**

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50 PIZOTIFEN

**※** Tab 500 mcg......23.21 100 **✓ Sandomigran** 

# Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT – Special Authority see SA0987 below – Retail pharmacy

**⇒SA0987** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

### **NERVOUS SYSTEM**

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
BETAHISTINE DIHYDROCHLORIDE  * Tab 16 mg	3.88	84	<b>✓</b> V	/ergo 16
CYCLIZINE HYDROCHLORIDE Tab 50 mg	0.55	10	<b>✓</b> N	lausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml		10	- ✔ H	
DOMPERIDONE  * Tab 10 mg	2.25	100	- ✓ P	Pharmacy Health
HYOSCINE HYDROBROMIDE  * Inj 400 mcg per ml, 1 ml ampoule		10	- -	Martindale \$29
Patch 1.5 mg – Special Authority see SA1998 below – Retai pharmacy	il	2	_	Scopoderm TTS

### ⇒SA1998 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

#### METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO	.1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO	.9.50	10	✓ Pfizer
ON	DANSETRON			
*	Tab 4 mg	.2.68	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO	.0.76	10	✓ Ondansetron ODT-DRLA
*	Tab 8 mg	1.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO	.1.13	10	✓ Ondansetron ODT-DRLA
PR	OCHLORPERAZINE			
*	Tab 3 mg buccal	.5.97	50	
		30.00)		Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO	.8.00	250	✓ Nausafix
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO	25.81	10	✓ Stemetil

## **Antipsychotics**

#### General

AMISULPRIDE - Safety medicine; prescriber may dete	ermine dispensing frequency	/	
Tab 100 mg	5.15	30	✓ Sulprix
·	17.16	100	✓ Amisulpride
			Mylan \$29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	1	Manufacturer
ARIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 5 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 10 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 15 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 20 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 30 mg	17.50	30	✓	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may determi	ne dis	spensina fr	eguency
Tab 10 mg – Up to 30 tab available on a PSO		100		Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100		Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10		Largactil
CLOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing freq	Hanov			
Tab 25 mg	•	50	1	Clozaril
1 ab 25 mg	6.69	50		Clopine
	11.36	100		Clozaril
	13.37	100		Clopine
Tab 50 mg		50		Clopine
1 ab 50 mg	17.33	100		Clopine
Tab 100 mg		50		Clozaril
Tab 100 mg	17.33	50		Clopine
	29.45	100		Clozaril
		100		Clopine
Tab 200 mg	34.65	EΩ		Clopine
1 ab 200 mg	69.30	50 100		Clopine
Suspension 50 mg per ml		100 m		Clopine
Suspension so my per mi	67.62	10011		Versacloz
			•	VEISACIOZ
HALOPERIDOL – Safety medicine; prescriber may determine				
Tab 500 mcg – Up to 30 tab available on a PSO		100		<u>Serenace</u>
Tab 1.5 mg — Up to 30 tab available on a PSO		100		<u>Serenace</u>
Tab 5 mg - Up to 30 tab available on a PSO		50		<u>Serenace</u>
0 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	29.72	100		<u>Serenace</u>
Oral liq 2 mg per ml – Up to 200 ml available on a PSO		100 m		<u>Serenace</u>
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a l	2SO21.55	10	•	<u>Serenace</u>
LEVOMEPROMAZINE - Safety medicine; prescriber may dete	ermine dispensing fred	uency		
Tab 25 mg (33.8 mg as a maleate)	16.10	100	✓	Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100	✓	<u>Nozinan</u>
Tab 100 mg (135 mg as a maleate)	41.75	100	✓	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	✓	<u>Nozinan</u>
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may deterr	nine d	lispensing	frequency
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
LITHIUM CARBONATE - Safety medicine; prescriber may det				
				Priadel
Tab long-acting 400 mg		100	_	
Cap 250 mg		100	•	Douglas
OLANZAPINE - Safety medicine; prescriber may determine dis		_	_	
Tab 2.5 mg		28		Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28		Zypine
Tab orodispersible 10 mg	2.38	28	✓	Zypine ODT

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg	10.49	84	1	Neulactil
•	12.49	100	1	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
•	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensina frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90	✓	Quetapel
Tab 200 mg		90	✓	Quetapel
Tab 300 mg		90	1	Quetapel
RISPERIDONE – Safety medicine; prescriber may determine di	spensing frequency			
Tab 0.5 mg	, , ,	60	1	Risperidone (Teva)
Tab 1 mg	2.06	60	1	Risperidone (Teva)
Tab 2 mg		60	✓	Risperidone (Teva)
Tab 3 mg	2.50	60	1	Risperidone (Teva)
Tab 4 mg	3.42	60	1	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 m	· •	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine dis	spensing frequency			
Cap 20 mg		60	1	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60	_	Zusdone
Cap 80 mg		60	1	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre		e disi	nensina fra	Pallency
Tab 10 mg	•	100		Clopixol
100 10 mg		.00	•	Ciopixoi

## **Depot Injections**

FLUPENTHIXOL DECANOATE – Safety medicine; prescribe Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO	13.14	nsing freq 5 5 5	uency  Fluanxol  Fluanxol  Fluanxol  Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber	mav determine dispen	sina freau	encv
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	, ,	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO.	55.90	5	<ul><li>✓ Haldol Concentrate</li><li>✓ Haldol</li></ul>
			Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retai	pharmacy		
Safety medicine; prescriber may determine dispensing fre	quency		
Inj 210 mg vial	252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv

## **⇒SA1428** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and

#### **NERVOUS SYSTEM**

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
- 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

#### ⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see SA1427 below – Retail pharmacy

Risperdal Consta
Risperdal Consta
Risperdal Consta
•

#### ⇒SA1427 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.



	\$	Per	Manutacturer
ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber m	nay determine	dispensing fr	equency
Inj 200 mg per ml, 1 ml - Up to 5 inj available on a PSO	19.80	5	✓ Clopixol
Anxiolytics			
BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine dispe	nsing frequenc	су	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispensin	g frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispens	sing frequency		
Tab 1 mg	9.72	250	✓ Ativan

Subsidy

(Manufacturer's Price)

Fully

Subsidised

100

100

100

Ativan

✓ Ox-Pam

✓ Ox-Pam

Brand or

Generic

### **Multiple Sclerosis Treatments**

### **⇒SA2051** Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and

OXAZEPAM - Safety medicine: prescriber may determine dispensing frequency

Tab 15 mg ......8.53

- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
  - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
  - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
  - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
  - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
  - 5.5 Either:
    - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
    - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and

		N	NER\	OUS SYSTEM
	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully sed	Brand or Generic Manufacturer
continued				
Evidence of new inflammatory activity on an MR scan with     Any of the following:     7.1 A sign of that new inflammatory activity is a gadoli	•			
7.2 A sign of that new inflammatory activity is a lesion				
<ul><li>7.3 A sign of that new inflammatory is a T2 lesion with</li><li>7.4 A sign of that new inflammatory activity is a promit</li></ul>			nsible	for the clinical features of
a recent relapse that occurred within the last 2 year 7.5 A sign of that new inflammatory activity is new T2		a previous	MR c	nan
Note: Natalizumab can only be dispensed from a pharmacy regi				
operated by the supplier. Treatment on two or more funded mult	iple sclerosis treatmer	nts simultan	eously	is not permitted.
Renewal — (Multiple sclerosis) only from a neurologist or gen had an EDSS score of 0 to 6.0 (inclusive) with or without the use				
(i.e. the patient has walked 100 metres or more with or without a	ids in the last six mon	ths).		
Note: Natalizumab can only be dispensed from a pharmacy region operated by the supplier. Treatment on two or more funded multi-				
DIMETHYL FUMARATE – Special Authority see SA2051 on the	•		-	io not pomittou.
a) Wastage claimable				
b) Note: Treatment on two or more funded multiple scleros Cap 120 mg		eously is no 14		nitted. e <b>cfidera</b>
Cap 240 mg		56		ecfidera
FINGOLIMOD - Special Authority see SA2051 on the previous	o <mark>age</mark> – Retail pharmad	су		
a) Wastage claimable     b) Note: Treatment on two or more funded multiple scleros Cap 0.5 mg	is treatments simultan	eously is no 28		nitted. Ienya
GLATIRAMER ACETATE - Special Authority see SA2051 on th	e previous page – Ret	ail pharmac	у	•
Note: Treatment on two or more funded multiple sclerosis tr Inj 40 mg prefilled syringe		ısly is not pe 12		d. <b>opaxone</b>
INTERFERON BETA-1-ALPHA - Special Authority see SA2051				
Note: Treatment on two or more funded multiple sclerosis tr Inj 6 million iu prefilled syringe		isly is not pe 4		d. /onex
Injection 6 million iu per 0.5 ml pen injector		4		onex Pen
INTERFERON BETA-1-BETA - Special Authority see SA2051 c				
Note: Treatment on two or more funded multiple sclerosis tr		isly is not pe 15		d. e <b>taferon</b>
NATALIZUMAB - Special Authority see SA2051 on the previous				
Note: Treatment on two or more funded multiple sclerosis tr Inj 20 mg per ml, 15 ml vial		isly is not pe		d. <b>rsabri</b>
OCRELIZUMAB – Special Authority see SA2051 on the previou Note: Treatment on two or more funded multiple sclerosis tr Inj 30 mg per ml, 10 ml vial	eatments simultaneou			d. crevus
TERIFLUNOMIDE – Special Authority see SA2051 on the previo			<b>.</b>	· · · · · · · · · · · · · · · · · · ·

## **Sedatives and Hypnotics**

MELATONIN - Special Authority see SA1666 on the next page - Retail pharmacy 30 ✓ Circadin Tab modified-release 2 mg - No more than 5 tab per day ......28.22

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

28

✓ Aubagio



Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

#### ⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

Trote: maladione manea min are unapproved maladione.			
MIDAZOLAM - Safety medicine; prescriber may determine of	dispensing frequency		
Inj 1 mg per ml, 5 ml ampoule	2.98	10	Mylan Midazolam
	4.30		✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj avail	able		
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO mus	t be endorsed for stat	us epileptici	us use only.
Inj 5 mg per ml, 3 ml ampoule	2.50	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj availa	able on		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO mus	t be endorsed for stat	us epileptici	us use only.
PHENOBARBITONE SODIUM - Special Authority see SA13	886 below – Retail pha	armacy	
Inj 200 mg per ml, 1 ml ampoule	78.20	10	✓ Max Health S29
⇒SA1386 Special Authority for Subsidy			

#### SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determine	dispensing frequency		
Tab 10 mg	1.33	25	✓ Normison
TRIAZOLAM - Safety medicine; prescriber may determine d	ispensing frequency		
Tab 125 mcg	5.10	100	
•	(9.85)		Hypam
Tab 250 mcg	4.10	100	
•	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine d	ispensing frequency		
Tab 7.5 mg	9.56	500	Zopiclone Actavis

✓ PSM

	Subsidy (Manufacturer's Price		Fully Subsidised	Brand or Generic
	\$	Per		Manufacturer
Stimulants/ADHD Treatments				
TOMOXETINE				
Cap 10 mg	18.41	28	1	Generic Partners
•	107.03		1	Strattera
Cap 18 mg	27.06	28	1	Generic Partners
Cap 25 mg	29.22	28	1	Generic Partners
Cap 40 mg	29.22	28	1	Generic Partners
, -	107.03		<b>✓</b> 9	Strattera
Cap 60 mg	46.51	28	✓ (	Generic Partners
Cap 80 mg	56.45	28	1	Generic Partners
	58.48	28	1	Generic Partners

### ⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or

b) Safety medicine; prescriber may determine dispensing frequency

3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.



	\$	Per	✓ Manufacturer
METHYLPHENIDATE HYDROCHLORIDE - Special Authority s	ee SA1964 below	– Retail pha	ırmacy
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing fr	equency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg	7.75	30	<ul><li>Methylphenidate ER</li></ul>
			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg	11.45	30	<ul><li>Methylphenidate ER</li></ul>
			- Teva
Tab extended-release 36 mg	15.50	30	Methylphenidate ER
•			- Teva
Tab extended-release 54 mg	22.25	30	<ul> <li>Methylphenidate ER</li> </ul>
<b>3</b>			- Teva

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

#### ⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy\*)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

#### **NERVOUS SYSTEM**

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
(Manufacturer's Frice)	_	Subsidised	Generic
\$	Per	✓	Manufacturer

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) Salety medicine, prescriber may determine dispensing in	equency		
Tab extended-release 18 mg	58.96	30	<ul><li>Concerta</li></ul>
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg	71.93	30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg	86.24	30	<ul><li>Concerta</li></ul>
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

### ⇒SA1965 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

#### All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

**Renewal** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

#### ⇒SA1999 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or

### **NERVOUS SYSTEM**

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

## **Treatments for Dementia**

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below - F	Retail pharmacy		
Patch 4.6 mg per 24 hour	48.75	30	✓ Generic Partners
Patch 9.5 mg per 24 hour	48.75	30	✓ Generic Partners

#### ⇒SA1488 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

### **Treatments for Substance Dependence**

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

	aloxone 0.5 mg			

Tab sublingual 8 mg with naloxone 2 mg ......53.12

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine Naloxone BNM

#### ⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is opioid dependent: and

continued...

- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone);
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE			
Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see SA	A1408 below – Reta	il pharmacy	
Tab 50 mg	133.33	30	✓ Naltraccord

#### ⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.



Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Patch 7 mg - Up to 28 patch available on a PSO		28	<ul><li>Habitrol</li></ul>
Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	<ul><li>Habitrol</li></ul>
Patch 14 mg - Up to 28 patch available on a PSO	19.95	28	<ul><li>Habitrol</li></ul>
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	<ul><li>Habitrol</li></ul>
Patch 21 mg - Up to 28 patch available on a PSO	22.86	28	<ul><li>Habitrol</li></ul>
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	<ul><li>Habitrol</li></ul>
Lozenge 1 mg - Up to 216 loz available on a PSO	19.18	216	<ul><li>Habitrol</li></ul>
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	<ul><li>Habitrol</li></ul>
Lozenge 2 mg - Up to 216 loz available on a PSO	21.02	216	<ul><li>Habitrol</li></ul>
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	<ul><li>Habitrol</li></ul>
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO	38.21	384	<ul><li>Habitrol</li></ul>
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	8.64	96	<ul><li>Habitrol</li></ul>
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	38.21	384	<ul><li>Habitrol</li></ul>
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	8.64	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO	44.17	384	<ul><li>Habitrol</li></ul>
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	10.01	96	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) - Up to 384 piece available on a PSO	44.17	384	<ul><li>Habitrol</li></ul>
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	10.01	96	<ul><li>Habitrol</li></ul>

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4225.6	4 53 OP	Champix
		✓ Varenicline Pfizer
Tab 1 mg27.1	0 56	✓ Champix
•		✓ Varenicline Pfizer

#### ⇒SA1845 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

### NERVOUS SYSTEM

Subsidy (Manufacturer's Price)	Subsid	Fully lised	Brand or Generic
\$	Per	•	Manufacturer

continued...

#### All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

#### ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

## **Chemotherapeutic Agents**

### **Alkylating Agents**

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see \$A2046 below

Inj 25 mg vial77.00	1	<ul> <li>Ribomustin</li> </ul>
Inj 100 mg vial308.00	1	✓ Ribomustin
Inj 1 mg for ECP	1 mg	✓ Baxter

#### ⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 All of the following:
    - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
    - 3.2.2 The patient has not received prior bendamustine therapy; and
    - 3.2.3 Fither:
      - 3.2.3.1 Both:
        - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
        - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
      - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
  - 2.1 Both:

(Ma	Subsidy	F	ully	Brand or
	anufacturer's Price)	Subsidi	sed	Generic
	\$	Per	1	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		· ·	
Inj 100 mg vial	1 387 00	1	✓ BiCNU
ing 100 mg viai	1,507.00	'	✓ Bicnu Heritage \$29
Ini 100 ma for ECD	1 207 00	100 mg OP	✓ Baxter
Inj 100 mg for ECP	1,307.00	100 mg OF	Daxiei
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	<ul><li>Leukeran FC</li></ul>
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	<ul> <li>Cisplatin Ebewe</li> </ul>
Inj 1 mg per ml, 100 ml vial	19.70	1	✓ DBL Cisplatin
	21.00		<ul> <li>Cisplatin Ebewe</li> </ul>
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
rab oo mg	158.00	100	✓ Procytox S29
Wastage claimable	156.00	100	V Procytox 329
Inj 1 g vial – PCT – Retail pharmacy-Specialist	25.65	1	✓ Endoxan
inj i g viai – POT – Hetali phamiacy-Specialist	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist		1	✓ Endoxan
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
	0.04	ring	Daxiei
IFOSFAMIDE – PCT only – Specialist	22.22	i i	<b>4</b> 11 1
lnj 1 g		1	✓ Holoxan
Inj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
	Ψ	rei		Manuacturei
_OMUSTINE - PCT - Retail pharmacy-Specialist	100.50		,	
Cap 10 mg		20		CeeNU
Cap 40 mg	399.15	20	•	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist		25		Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	•	Alkeran
			1	Alkeran S29 S29
	420.00		1	Tillomed S29
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
,		•		100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg		Baxter
THIOTEPA – PCT only – Specialist		9		
, ,	000		,	D - 46 1 000
Inj 15 mg vial	BS	1		Bedford S29
				THIO-TEPA S29
			/	Tepadina S29
Inj 100 mg vial	CBS	1	1	Tepadina S29
A malion and a little a				
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see S	A1467 below			
Inj 100 mg vial	139.00	1	1	Azacitidine Dr

# 

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
  - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or

605.00

- 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

✓ Vidaza

✓ Baxter

	Out side		F. II.	. December
	Subsidy (Manufacturer's Price	s) S	Fully ubsidised	
	\$	Per	<b>√</b>	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg – PCT – Retail pharmacy-Specialist	114.69	10	✓	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia		1		Calcium Folinate Sandoz
			•	Calcium Folinate Sandoz S29 S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	•	Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	•	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	•	Calcium Folinate Sandoz
			•	Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	✓	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	•	Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓	Baxter
CAPECITABINE - Retail pharmacy-Specialist		•		
Tab 150 mg	10.00	60	1	Capercit
Tab 500 mg		120	✓	Capercit
CLADRIBINE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml	749.96	1	/	Leustatin
Inj 10 mg for ECP		0 mg OF	•	Baxter
CYTARABINE		_		
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 20 ml vial – PCT – Retail	alist400.00	5	•	Pfizer
pharmacy-Specialist	41.36	1	/	Pfizer
Inj 1 mg for ECP - PCT only - Specialist	0.25	10 mg	1	Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	alist80.00 1	00 mg O	P 🗸	Baxter
FLUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	✓	Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		i0 mg OF	· •	Baxter
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	12.00	1	✓	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	30.00	1	✓	Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist	0.66	100 mg		Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1		DBL Gemcitabine
lnj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	✓	Baxter

	Subsidy (Manufacturer's Pri \$	ce) Per	Fully Subsidised	I Generic
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist Inj 20 mg per ml, 5 ml vial	71.44	1	/	Irinotecan
			•	Accord S29 Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	•	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist Oral suspension 20 mg per ml - Retail pharmacy-Specialist		25	•	Puri-nethol
Special Authority see SA1725 below		100 ml	OP 🗸	Allmercap

# **⇒SA1725** Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

**Renewal** only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

# METHOTREXATE

	1110111270112		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist8.05	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist31.75	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
	, 0, , 0		Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
•	, <del>-</del> -,	-	Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
•••	Trip 20 mg promised syrings	•	Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
~	ing 20 mg promited syringe14.00	•	Sandoz
*	Inj 30 mg prefilled syringe	1	✓ Methotrexate
~	ing of mg premied syninge	•	Sandoz
*	Ini 05 ma nor ml. 0 ml.viol. BCT. Botoil pharmany Chaniclist. 20.00	5	✓ Methotrexate DBL
不	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	Onco-Vial
	Ini OF man you and OO and visal DOT Detail about 200 Consciolist 45 00	4	*****
*	Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist45.00	I	✓ DBL Methotrexate
	1:400 L40 L BOT B L " L		Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ <u>Methotrexate Ebewe</u>
*	, 3	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
PΕ	METREXED – PCT only – Specialist – Special Authority see SA1679 on the n	ext page	
	Inj 100 mg vial60.89	1	<ul> <li>Juno Pemetrexed</li> </ul>
	Inj 500 mg vial217.77	1	<ul> <li>Juno Pemetrexed</li> </ul>
	Inj 1 mg for ECP	1 mg	✓ Baxter

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

### ⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient has chemotherapy-naïve disease; and
    - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cvcles: or
  - 2.2 All of the following:
    - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
    - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
    - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m<sup>2</sup> every 21 days.

Tab 40 mg	126.31	25	Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

	Subsidy (Manufacturer's Pr	rica) S	Fully	
	\$	Per	Jubsiuiseu ✓	Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist				
Inj 15,000 iu, vial	161.01	1	✓	DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓	Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see S.	A1889 below			
Inj 3.5 mg vial	105.00	1	✓	Bortezomib Dr-Reddy's
Inj 1 mg for ECP	31.20	1 mg	•	Baxter

# ⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis \*.

Note: Indications marked with \* are unapproved indications

62.70	1	✓ DBL Dacarbazine
580.60	10	✓ Dacarbazine
		APP S29
62.70	200 mg OP	✓ Baxter
255.00	1	✓ Cosmegen
	0.5 mg OP	✓ Baxter
	-	
149.50	1	✓ Pfizer
	20 mg OP	✓ Baxter
	ŭ	
48.75	1	✓ Docetaxel Sandoz
46.89	i	✓ DBL Docetaxel
	1	✓ Docetaxel
		Accord \$29
195.00	1	✓ Docetaxel Sandoz
	1 mg	✓ Baxter
	-	
10.00	1	Doxorubicin Ebewe
11.50	1	Doxorubicin Ebewe
17.00		✓ Arrow-Doxorubicin
23.00	1	Doxorubicin Ebewe
56.15	1	<ul><li>Doxorubicin Ebewe</li></ul>
65.00		Arrow-Doxorubicin
0.29	1 mg	✓ Baxter
25.00	1	Epirubicin Ebewe
	1	✓ Epirubicin Ebewe
	1	<ul> <li>Epirubicin Ebewe</li> </ul>
0.43	1 mg	✓ Baxter
	62.70255.00149.50149.5048.7546.8926.95195.0006510.0011.50 17.0023.0056.15	580.60 1062.70 200 mg OP255.00 1255.00 0.5 mg OP149.50 1149.50 20 mg OP48.75 146.89 126.95 1195.00 10.65 1 mg150 10.65 1 mg

	Subsidy	Ful	ly Brand or
	(Manufacturer's Price)	Subsidise	
	\$	Per •	Manufacturer
ETOPOSIDE			
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali		1 •	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist		1 mg	Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist		3	
, ,	40.00	1	/ Etonophoo
Inj 100 mg (of etoposide base)			✓ Etopophos ✓ Baxter
Inj 1 mg (of etoposide base) for ECP		1 mg •	Daxier
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail phan	macy-Specialist		
Cap 500 mg	23.82	100	<u>Devatis</u>
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial - PCT only - Specialist	93.00	1 •	Zavedos
Inj 10 mg vial - PCT only - Specialist		· 1 •	Zavedos
Inj 1 mg for ECP – PCT only – Specialist		· ·	Baxter
, ,		9	Durio
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority Wastage claimable	y see SA2047 below		
Cap 5 mg	5,122.76	28	Revlimid
Cap 10 mg		21	Revlimid
•	6,207.00	28	Revlimid
Cap 15 mg	,	21	/ Revlimid
	7.239.18		Revlimid
Cap 25 mg	,		Revlimid
	,0200		

# **⇒SA2047** Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

**Initial application** — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

continued...

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

#### **MESNA**

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Speciali	ist177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specia	alist407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 20 mg vial	3,275.00	1	✓ Omegapharm \$29
, ,	•		✓ Teva
Inj 1 mg for ECP	288.09	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist		· ·	
Inj 2 mg per ml, 10 ml vial	97.50	1	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authorit	ty see SA1883 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg	3,701.00	56	✓ Lynparza
Cap 50 mg - Wastage claimable	7,402.00	448	✓ Lynparza
(Lynparza Cap 50 mg to be delisted 1 July 2021)			

### **⇒SA1883** Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued

for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg	24.00	1	Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see	e SA1979 below		

### ⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only -	Specialist		
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail p	harmacy-Specialist		
Can 50 mg	980.00	50	✓ Natulan S29

✓ Oncaspar LYO S29

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Brand or Generic Manufacturer
TEMOZOLOMIDE - Special Authority see SA	1741 below – Retail pharmacy			
Cap 5 mg		5	1	Temaccord
Cap 20 mg		5	<b>✓</b>	Temaccord
, ,	18.30		✓	Apo-Temozolomide
	136.00	14	✓.	Accord S29
Cap 100 mg	35.98	5	1	Temaccord
	40.20		1	Apo-Temozolomide
	532.00	14	✓.	Accord S29
Cap 140 mg	50.12	5	1	Temaccord
	400.00		1	Amneal S29
Cap 180 mg	620.00	14	1	Accord \$29
Cap 250 mg		5	1	Temaccord
	688.00		1	Amneal \$29

# ⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

**Initial application — (neuroendocrine tumours)** only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has glioblastoma multiforme; and
  - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
  - 2.1 Patient has anaplastic astrocytoma\*; and
  - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
  - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Aut	hority see SA1124 below	I	
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

### ⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an unapproved indication.

#### **TRFTINOIN**

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 bel	OW	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	✓ Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	120	✓ Venclexta

### ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

#### VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓ DBL Vinblastine S29
		✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓ Navelbine
210.00		✓ Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	✓ Baxter

# Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

continued...

Alecensa

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

for 6 months for applications meeting the following criteria:

#### Roth:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

# DASATINIB - Special Authority see SA1805 below - Retail pharmacy

### Wastage claimable

Tab 20 mg3,774.06	60	✓ Sprycel
Tab 50 mg6,214.20	60	✓ Sprycel
Tab 70 mg	60	✓ Sprycel

### ⇒SA1805 Special Authority for Subsidy

**Initial application** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

# Any of the following:

- 1 Both:
  - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
  - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
  - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
  - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Maximum dose of 100 mg/day; and
  - 3.3 Any of the following:
    - 3.3.1 Patient has documented treatment failure\* with imatinib; or
    - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
    - 3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Lack of treatment failure while on dasatinib\*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA2000 below

✓ Tarceva	30	Tab 100 mg764.00
✓ Tarceva	30	Tab 150 mg

## ⇒SA2000 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 3 Either:
  - 3.1 Patient is treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued defitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
  - 4 Erlotinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2001 below

Tab 250 mg .......1,700.00 30 ✓ Iressa

# ⇒SA2001 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fitha
  - 2.1 Patient is treatment naive; or
    - 2.2 Both:
      - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
      - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

# **IMATINIB MESILATE**

Note: Imatinib-AFT is not a registered for the treatment of Gastro Intestinal Stromal Tumours (GIST). The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg - [Xpharm] - Special Authority see SA1460 on the

	next page	2,400.00	60	✓ Glivec
*	Cap 100 mg		60	✓ Imatinib-Rex
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex

Subsidy (Manufacturer's Price) Per

Fully Subsidised

Brand or Generic Manufacturer

# ⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from PHARMAC's website schedule pharmac govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 **PHARMAC** Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

#### Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

### LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

70 Tykerb

# ⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology):
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

# NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

# ⇒SA1489 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
  - 2.1 Patient has documented CML treatment failure\* with imatinib; or
  - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer	
PALBOCICLIB – Retail pharmacy-Specialist – Special Authority Wastage claimable	see SA1894 below				
Cap 75 mg	4,000.00	21	<b>✓</b> Ib	rance	
Cap 100 mg	4,000.00	21	✓ Ib	rance	
Cap 125 mg	4,000.00	21	<b>✓</b> Ib	rance	

### ⇒SA1894 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
  - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
  - 4.2.2.2 All of the following:
    - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
    - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
    - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

# PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

✓ Votrient	30	1,334.70	·······	Tab 200 mg
✓ Votrient	30	2.669.40		Tab 400 mg

## ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and

Subsidy	e)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

#### continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

vvastage ciaimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5.000.00	56	✓ Jakavi

# ⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
    - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Aut	thority see SA2002	on the next page -	- Retail pharmacy

Cap 12.5 mg	2,315.38	28	Sutent
Cap 25 mg	· ·	28	✓ Sutent
Cap 50 mg	· ·	28	✓ Sutent

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

# **⇒SA2002** Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

**Renewal — (RCC)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
 \$	Per	1	Manufacturer

continued...

- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

# **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

Tab 250 mg ......4,276.19 120 **✓ Zytiga** 

# ⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline: and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

#### BICAL LITAMIDE

BIOALOTAMIDE			
Tab 50 mg	1.36	10	✓ Calutide-50 S29
·	4.07	30	✓ Binarex
	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	119.50	100	✓ Flutamin

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer	
FULVESTRANT – Retail pharmacy-Specialist – Special Authority Inj 50 mg per ml, 5 ml prefilled syringe		2	✓ Fa	aslodex	

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease: and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

#### MEGESTROL ACETATE

1ab 160 mg	63.53	30	Apo-Megestroi
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	✓ Octreotide GH \$29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	<ul> <li>Octreotide</li> </ul>
			MaxRx S29
	56.87		✓ DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule	72.50	5	✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓ DBL Octreotide
	222.00		<ul> <li>Octreotide</li> </ul>
			(Sun) \$29

#### OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Special Authority see \$A2004 below - Retail pharmacy

Inj LAR 10 mg prefilled syringe1,772	2.50 1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe2,358		Sandostatin LAR
Inj LAR 30 mg prefilled syringe2,951	1.25 1	Sandostatin LAR

### ⇒SA2004 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed: and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued

specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma: and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas: and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

**Renewal — (Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

*	Tab 10 mg	60 60	✓ <u>Tamoxifen Sandoz</u> ✓ <u>Tamoxifen Sandoz</u>

Aromatase Inhibitors		
ANASTROZOLE	30	✓ Anatrole
EXEMESTANE		
* Tab 25 mg14.50	30	✓ Pfizer Exemestane

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LETROZOLE  * Tab 2.5 mg	4.68	30	✓	<u>Letrole</u>
Immunosuppressants				
Cytotoxic Immunosuppressants				

# A7ATHIOPRINE

60 100 1	✓ <u>Azamun</u> ✓ <u>Azamun</u> ✓ Imuran
50	✓ Cellcept
100	✓ Cellcept
165 ml OP	<ul> <li>Cellcept</li> </ul>
	100 1 50 100

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

# **Fusion Proteins**

ETANERCEPT - Special Authority see SA2048 below - F	Retail pharmacy		
Inj 25 mg	690.00	4	<ul><li>Enbrel</li></ul>
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

## ⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Either:

1 Both:

- 1.1 Fither:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Fither:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application** — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

and

- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
  - 2 Either:
    - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
    - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
  - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pvoderma gangrenosum\*: and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:
    - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application** — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

# Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

 Subsidy	Fully		Brand or
 Manufacturer's Price)	Subsidis	sea	Generic
\$	Per	✓	Manufacturer

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (severe chronic plague psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Fither:
      - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 2.2 Both:
    - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
    - 2.2.2 Fither:
      - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip: and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose): and
- 5 Any of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
<u> </u>	Per	1	Manufacturer

continued...

- 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or

ANTITUVMOCYTE OLODIJI IN /FOLIME\ DOT only Choololist

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

### **Immune Modulators**

ANTITITINOCTTE GLOBULIN (EQUINE) - POT ONLY - S	pecialist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT	only - Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG S29 Inj 40 mg per ml, vial to be delisted 1	April 2022)		

# Monoclonal Antibodies

		9 below – Retail pharmacy	ADALIMUMAB – Special Authority see SA2049 be
<ul><li>Humira</li></ul>	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
<ul><li>Humira</li></ul>	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

### ⇒SA2049 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg. non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis: or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sacittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
    - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
    - 2.1.2 CDAI score is 150 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (Crohn's disease - children)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a gastroenterologist; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
    - 2.1.2 PCDAI score is 15 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (fistulising Crohn's disease)** only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

**Renewal — (hidradenitis suppurativa)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

<u></u>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
  - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Roth:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:
    - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
  - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plague psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
    - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from. at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
- 2.2 Both:
  - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2.2 Fither:
    - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
    - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation: or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
  - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## ⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

**Initial application — (diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	<ul><li>Erbitux</li></ul>
Inj 1 mg for ECP	3.82	1 mg	Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### ⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see SA2050 below

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	Baxter

#### ⇒SA2050 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation: or
- 2 Roth:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
<b>\$</b>	Per	1	Manufacturer	

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — **(neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcojosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Either:
    - 2.3.1 There has been an improvement in MRI appearances; or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
  - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Fither:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Either:
    - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis: or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis: or
  - 2.4 Severe ocular inflammation: or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults): or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease; or
  - 2.9 Severe fulminant ulcerative colitis: or
  - 2.10 Severe ulcerative colitis: or
  - 2.11 Plague psoriasis; or
  - 2.12 Neurosarcoidosis: or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

#### Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

**Initial application — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application** — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
  - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

**Renewal — (ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

		6 below - Retail pharmacy	MEPOLIZUMAB - Special Authority see SA189
✓ Nucala	1	1,638.00	Inj 100 mg prefilled pen
✓ Nucala	1	1.638.00	Ini 100 mg vial

#### ⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB – PCT only – Specialist – Special Author	ority see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

### ⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

OMALIZUMAB – Special Authority see SA1744 below – Retail	pharmacy		
Inj 150 mg prefilled syringe	450.00	1	Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

#### ⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and

- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
  - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
  - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
  - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal** — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjet	а
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxte	r

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✔ Manufacturer

## ⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist	pecial Authority see SA197	'6 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

#### ⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy (Manufacturer's Price)	Fu Subsidise	,	
 \$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

#### ⇒SA2028 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

**Initial application** — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	•	Por 🗸	Manufacturer

continued...

- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive; or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
  - 3 The patient has good performance status; and
  - 4 Either:
    - 4.1 The patient does not have chromosome 17p deletion CLL; or
    - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
  - 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
  - 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Either:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

375 mg/m2 administered weekly for four weeks; and

- 2 Either
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD: and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
  - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
  - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Steroid resistant nephrotic syndrome (SRNS))** only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles: or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal** — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of  $2 \times 1,000$ mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

(Manufacturer <sup>'</sup> s Price) Subsidised Generic \$ Per ✔ Manufacturer
---

continued...

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles: or
  - 2 Both:
    - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
    - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Fither:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

#### Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

SECUKINUMAB – Special Authority see SA2044 on the n	iext page – Retail pharmac	:y	
Inj 150 mg per ml, 1 ml prefilled syringe	799.50	1	✓ Cosentyx
	1,599.00	2	✓ Cosentyx

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

### ⇒SA2044 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
  - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Fu Subsidis	,	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fithe
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

**Initial application — (psoriatic arthritis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for psoriatic arthritis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

	Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic			
	\$	Per	1	Manufacturer			
SILTUXIMAB – Special Authority see SA1596 below – Retail pharmacy							
Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.							
Inj 100 mg vial	770.57	1		Sylvant			
Inj 400 mg vial	3,082.33	1	1	Sylvant			

## ⇒SA1596 Special Authority for Subsidy

**Initial application** only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1977 below

Inj 20 mg per ml, 4 ml vial220.00	1	Actemra
Inj 20 mg per ml, 10 ml vial550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial	1	✓ Actemra
Inj 1 mg for ECP	1 mg	✓ Baxter

## ⇒SA1977 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
  - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial: and
  - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis: or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease: or
  - 2.4 polyarticular juvenile idiopathic arthritis; or
  - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Sub	Fully sidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Both:
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Bules of the Pharmaceutical Schedule: and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	<ul> <li>Herceptin</li> </ul>
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

#### ⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 Both:
    - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and

Subsidy (Manufacturer's Price)	Ful Subsidise	,	
\$	Per •	Manufacturer	

continued...

- 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
  - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 3.2 Both:
    - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 3.2.2 The cancer did not progress whilst on lapatinib; or
  - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
  - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 4.2 All of the following:
    - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
    - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully bsidised	Brand or Generic Manufacturer	
TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special	Authority see SA187	'1 below			
Inj 100 mg vial	2,320.00	1	✓ K	(adcyla	
Inj 160 mg vial	3,712.00	1	✓ K	(adcyla	
Inj 1 mg for ECP	23.20	1 mg	<b>✓</b> B	Baxter	

### **⇒SA1871** Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
  - 3.1 The patient has received prior therapy for metastatic disease\*: or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: \*Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

# Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB – PCT only – Specialist – Special Autho	rity see SA2006 below		
Inj 10 mg per ml, 4 ml vial	1,051.98	1	<ul><li>Opdivo</li></ul>
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

## ⇒SA2006 Special Authority for Subsidy

**Initial application** only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

for 4 months for applications meeting the following criteria:

#### Fither

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Either:
    - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
    - 1.2.2 Both:
      - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
      - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm.</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

		PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2007 below
✓ Keytruda	1	Inj 25 mg per ml, 4 ml vial
✓ Baxter	1 mg	Inj 1 mg for ECP49.14

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Either:
    - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
    - 1.2.2 Both:
      - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
      - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm.</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest

### ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

ıbsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
\$ Per	1	

#### continued...

sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).

 Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

## Other Immunosuppressants

CICLOSPORIN		
Cap 25 mg	50	✓ Neoral
Cap 50 mg88.91	50	✓ Neoral
Cap 100 mg177.81	50	✓ Neoral
Oral liq 100 mg per ml198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pharmacy		
Wastage claimable		
Tab 10 mg6,512.29	30	✓ Afinitor
Tab 5 mg4,555.76	30	✓ Afinitor

#### ⇒SA2008 Special Authority for Subsidy

**Initial application** only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months: and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

### SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

### ⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

continued...

## ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

(1	Subsidy Manufacturer's Price)	F Subsid	ully	Brand or Generic
· ·	\$	Per	•	Manufacturer

continued...

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note): or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

 $\textbf{Renewal--(renal angiomyolipoma(s) associated with tuberous sclerosis complex*)} \ \ \text{from any relevant practitioner}.$ 

Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and

continued...

## ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

	operation of the second of the		
Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

### ⇒SA1745 Special Authority for Subsidy

**Initial application** — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with \* are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

# **Antiallergy Preparations**

## Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe..................2,668.00 1 ✓ Firazyr

#### ⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## Allergy Desensitisation

## **⇒SA1367** Special Authority for Subsidy

**Initial application** only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above	ve – Retail pharmacy	
Initiation kit - 5 vials freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with		
diluent285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent		
9 ml, 3 diluent 1.8 ml305.00	0 1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA1367 at	oove – Retail pharmac	у
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze		
dried venom, with diluent305.00	0 1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	100	✓ Alle and
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	0 1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze	1.00	/ Vanamil coo
dried venom, with diluent305.00	0 1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Pr	rice) Subsid	dised Generic
	\$	Per	✓ Manufacturer
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral lig 1 mg per ml		200 ml	✓ Histaclear
		200 1111	riistacieai
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
本 Tau 2 IIIy		40	Delevenine
	(8.40)		Polaramine
	1.01	20	
	(5.99)		Polaramine
* Oral liq 2 mg per 5 ml	1.77	100 ml	
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE	, ,		
	4.04	00	
* Tab 60 mg		20	T-161
	(8.23)		Telfast
* Tab 120 mg		10	
	(8.23)		Telfast
	14.22	30	
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg	1.60	100	✓ Lorafix
ŭ			
* Oral liq 1 mg per ml		100 ml	✓ Haylor syrup
	2.95	120 ml	✓ Lorfast
Haylor syrup to be Sole Supply on 1 September 2021			
(Lorfast Oral liq 1 mg per ml to be delisted 1 September 2021)			
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1 68	50	✓ Allersoothe
* Tab 25 mg		50	✓ Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	✓ Allersoothe
, 0,		5	
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	-30 17.07	5	✓ Hospira
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	9.30	200 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
. 01		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose			
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.6/	200 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort
1 311301 101 Hillianation, 200 Hog per 4030		_00 0000 OI	Turbuhaler
Decodes for inhelation, 400 construction	00.00	000 4 00	
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy		Fully Brand	
	(Manufacturer's F	Price) Sub Per	sidised Gener	ric facturer
	Ψ	1 61	• Manu	lacturer
LUTICASONE			<b>4 -</b>	
Aerosol inhaler, 50 mcg per dose		120 dose OP		
Powder for inhalation, 50 mcg per dose		60 dose OP		e Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP		e Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP		_
Aerosol inhaler, 250 mcg per dose		120 dose OP		_
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide	e Accuhaler
nhaled Long-acting Beta-adrenoceptor Agonis	ts			
FORMOTEROL FUMARATE				
Powder for inhalation, 12 mcg per dose, and monodose devi		60 dose		
	(35.80)		Foradil	
FORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose	e)10.32	60 dose OP		
(oquirus in to ordinate or language or mag motores upon	(16.90)		Oxis Tur	buhaler
IDACATEROL	(10.00)		Oxio rui	Dariaioi
IDACATEROL	04.00	00 de e OD		D
Powder for inhalation 150 mcg		30 dose OP	✓ Onbrez	
Powder for inhalation 300 mcg	61.00	30 dose OP	<ul><li>Onbrez</li></ul>	Breezhaler
ALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	<ul><li>Serever</li></ul>	ıt
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	✓ Serever	t Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocepto	or Agonists	;	
UDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide w	<i>ii</i> th			
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoRes	p Spiromax
,		120 00se OF	Duones	p Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumar				
per dose (equivalent to 400 mcg budesonide with 12 mc	g			
eformoterol fumarate metered dose) - No more than 2	00.50	100   00		
dose per day		120 dose OP		p Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP		
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OP	-,	
				haler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP		
Powder for inhalation 200 mcg with eformoterol fumarate 6 n	ncg44.08	120 dose OP	- ,	
			Turbu	haler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate				
12 mcg - No more than 2 dose per day	44.08	60 dose OP	✓ Symbice	ort
, ,			•	haler 400/12
LUTICASONE FUROATE WITH VILANTEROL				
	44.00	30 dose OP	√ Bros Ell	into
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.00	30 dose OP	✓ Breo Ell	ihra

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or sidised Generic  Manufacturer	
FLUTICASONE WITH SALMETEROL	*			
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25 79	120 dose OP	✓ Seretide	
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide	
Powder for inhalation 100 mcg with salmeterol 50 mcg - No				
more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler	r
Powder for inhalation 250 mcg with salmeterol 50 mcg - No	)			
more than 2 dose per day	44.08	60 dose OP	<ul> <li>Seretide Accuhale</li> </ul>	r
Beta-Adrenoceptor Agonists				
SAL PLITAMOL				
SALBUTAMOL Oral lig 400 mcg per ml	20.00	150 ml	✓ Ventolin	
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin	
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO		5	✓ Ventolin	
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	<ul><li>✓ Respigen</li><li>✓ SalAir</li></ul>	
	(6.00)		Ventolin	
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb	` '		VEHIOIIII	
available on a PSO		20	✓ Asthalin	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule - Up to 30 neb				
available on a PSO		20	✓ <u>Asthalin</u>	
FERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhale	er
Autich clinovnic Augusta				
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	✓ Atrovent	
a) Up to 400 dose available on a PSO				
b) No patient co-payment payable				
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 n		20		
available on a PSO	11./3	20	✓ <u>Univent</u>	
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg	per			
dose CFC-free	•	200 dose OP	✓ Duolin HFA	
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial 2.5 ml amnoula. Un to 20 nob available on a DSC	5 20	20	✓ Duolin	

vial, 2.5 ml ampoule – Up to 20 neb available on a PSO .............5.20

✓ Duolin

20

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

## **Long-Acting Muscarinic Antagonists**

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Seebri Breezhaler

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose......50.37 30 dose ✓ Spiriva Soln for inhalation 2.5 mcg per dose......50.37 60 dose OP ✓ Spiriva Respimat

#### UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

30 dose OP ✓ Incruse Ellipta

# Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

### ⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg.....81.00 ✓ Ultibro Breezhaler 30 dose OP

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg.....81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg ......77.00 30 dose OP ✓ Anoro Ellipta

### **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg......2,554.00 60 OP ✓ Ofev 60 OP ✓ Ofev 

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

### ⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib

Tab 801 mg	3,645.00	90	✓ Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	<ul><li>Esbriet</li></ul>

### ⇒SA2013 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	1	Manufacturer

# Leukotriene Receptor Antagonists

МС	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
*	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	28	✓ Montelukast Mylan

## **Mast Cell Stabilisers**

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

## Methylxanthines

#### **AMINOPHYLLINE**

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO18	0.00	5	✓ DBL Aminophylline
THE	EOPHYLLINE			
*	Tab long-acting 250 mg2	3.02	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml1	6.60	500 ml	✓ <u>Nuelin</u>

# **Mucolytics**

		8 below – Retail pharmacy	DORNASE ALFA – Special Authority see SA1978
✓ Pulmozyme	6	250.00	Nebuliser soln, 2.5 mg per 2.5 ml ampoule

### ⇒SA1978 Special Authority for Subsidy

**Initial application — (cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	S	Subsidised	Generic	
	\$	Per	1	Manufacturer	
IVACAFTOR - PCT only - Specialist - Special Authority see SA	2017 below				
Tab 150 mg	29,386.00	56	✓ K	Calydeco	
Oral granules 50 mg, sachet	29,386.00	56	✓ K	alydeco	
Oral granules 75 mg, sachet	29,386.00	56	✓ K	Calydeco	

### **⇒SA2017** Special Authority for Subsidy

**Initial application** only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

#### SODIUM CHI ORIDE

Not funded for use as a nasal drop.

## **Nasal Preparations**

# Allergy Prophylactics

BUDESONIDE  Metered aqueous nasal spray, 50 mcg per dose  Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
FLUTICASONE PROPIONATE  Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE	5 23	15 ml OP	✓ Univent

## **Respiratory Devices**

#### MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

Small 2.20 1 ✓ e-chamber Mask

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer
	\$	rei		Manuacturei
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	1	Mini-Wright AFS
				Low Range
Normal range	9.54	1	1	Mini-Wright
Č				Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	2.95	1	1	e-chamber Turbo
510 ml (single patient)	5.12	1	_	e-chamber La
o . o (eg.o pano)				Grande
800 ml	6.50	1	1	Volumatic
000 111				Volumento
Respiratory Stimulants				
nespiratory Stillidiants				
CAFFEINE CITRATE				
Oral liq 20 mg per ml (10 mg base per ml)	15.10 25	ml C	)P 🗸	Biomed
The state of the s				

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or idised Generic ✓ Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE		ngo 027	
For Vosol ear drops with hydrocortisone powder refer Stand	aru Formulae, pa	ige 237	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	ΓΙΝ	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
3	(9.27)		Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%	4.13	8 ml OP	
	(8.65)		Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL			
Eye oint 1%		5 g OP	✓ <u>Devatis</u>
Eye drops 0.5%	1.54 re unapproved in	10 ml OP dications.	✓ <u>Chlorafast</u>
CIPROFLOXACIN			
Eye drops 0.3% – Subsidy by endorsement	or severe bacteria		
Note: Indication marked with a * is an unapproved indic		, and the presi	supplier to endersed accordingly.

✓ Genoptic

Brolene

✓ Fucithalmic

5 ml OP

10 ml OP

5 g OP

(14.55)

**GENTAMICIN SULPHATE** 

PROPAMIDINE ISETHIONATE

SODIUM FUSIDATE [FUSIDIC ACID]

	Subsidy (Manufacturer's Price)		Full Subsidise	
	\$	Pe	r 🗸	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g	OP 🗸	Tobrex
Eye drops 0.3%	11.48	5 ml (	OP 🗸	Tobrex
Corticosteroids and Other Anti-Inflammatory Pre	parations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g	OP 🗸	Maxidex
* Eye drops 0.1%	4.50	5 ml (	OP 🗸	Maxidex
Ocular implant 700 mcg - Special Authority see SA1680 belo	)W			

### ⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

**Initial application — (Women of child bearing age with diabetic macular oedema)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

## DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 a OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6.000 u per ml		5 ml OP	✓ Maxitrol
חור	CLOFENAC SODIUM	4.50	3 IIII OF	• Maxill OI
Dic	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha

✓ Ozurdex

	Subsidy (Manufacturer's F	Price) Sub	Fully sidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
FLUOROMETHOLONE					
* Eye drops 0.1%	3.09	5 ml OP	<b>✓</b> F	ML	
	5.20		<b>√</b> F	lucon	
KETOROLAC TROMETAMOL - Special Authority see SA1981	below - Retail ph	armacy			
Eye drops 0.5%	9.50 ·	5 ml OP	<b>√</b> p	Acular	
OA4004 Out a dad Anatha with a fam Ondra late.					

#### ⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
  - 2.1 The patient is at risk of postoperative macular oedema; and
  - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

### LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71	4 ml OP	
, , , , , , , , , , , , , , , , , , , ,	(10.34)		Livostin
	(10.01)		21100111
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eve drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
	г оо	40 ml OD	/ Duadwisslams AFT
Eye drops 1%		10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	- Retail pharn	nacv
Eve drops 0.5%, single dose (preservative free)		20 dose	✓ Minims
2,0 diopo 0.0/0, oliigio dobo (proborvativo 1100)		_0 0000	
			Prednisolone

## **⇒SA1715** Special Authority for Subsidy

**Initial application** only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

5 ml OP

2.5 ml OP

✓ Rexacrom

✓ Arrow-Timolol

✓ Timoptol XE

### SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers		
BETAXOLOL		
* Eye drops 0.25%	5 ml OP	✓ Betoptic S
* Eye drops 0.5%	5 ml OP	✓ Betoptic
TIMOLOL		
* Eve drops 0.25%	5 ml OP	✓ Arrow-Timolol

	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully Brand or sidised Generic Manufacturer
Glaucoma Preparations - Carbonic Anhydrase I	nhibitors		
ACETAZOLAMIDE	17.02	100	✓ Diamox
* Tab 250 mg	17.03	100	Diamox
BRINZOLAMIDE  * Eye drops 1%	7.30	5 ml OP	✓ Azopt
DORZOLAMIDE HYDROCHLORIDE		0 1111 01	- Azopt
* Eye drops 2%	9 77	5 ml OP	
	(17.44)	0 1111 01	Trusopt
DORZOLAMIDE WITH TIMOLOL	, ,		
* Eye drops 2% with timolol 0.5%	2.87	5 ml OP	✓ Dortimopt
Glaucoma Preparations - Prostaglandin Analog	ues		
BIMATOPROST			
* Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost
			Multichem
LATANOPROST			
* Eye drops 0.005%	1.57	2.5 ml OP	✓ <u>Teva</u>
TRAVOPROST			
* Eye drops 0.004%		5 ml OP	✓ Travopt
	10.50	2.5 ml OP	✓ Mylan <sup>©29</sup> ✓ Travatan
	19.50	2.5 IIII OP	ravatan
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE			
* Eye drops 0.2%	12.25	5 ml OP	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	<ul><li>Combigan</li></ul>
LATANOPROST WITH TIMOLOL			
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	Arrow - Lattim
Arrow - Lattim to be Sole Supply on 1 September 2021			
PILOCARPINE HYDROCHLORIDE			
* Eye drops 1%		15 ml OP	✓ Isopto Carpine
* Eye drops 2%  * Eye drops 4%		15 ml OP 15 ml OP	<ul><li>✓ Isopto Carpine</li><li>✓ Isopto Carpine</li></ul>
Subsidised for oral use pursuant to the Standard Formul		13 1111 01	• ISOPIO CAI PIIIE
* Eye drops 2% single dose – Special Authority see SA0895	~~·		
below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine
- CARROLL Created Authority for Cubaidy			•

### ⇒SA0895 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Pri \$	ice) Subsi Per	Fully dised	Brand or Generic Manufacturer
Mydriatics and Cycloplegics				
ATROPINE SULPHATE  * Eye drops 1%  CYCLOPENTOLATE HYDROCHLORIDE	17.36	15 ml OP	✓ <u>A</u>	tropt
* Eye drops 1%      * Eye drops 1%, single dose (preservative free) – Only on a	8.76	15 ml OP	<b>✓</b> C	yclogyl
prescription	52.86	20 dose		inims Cyclopentolate
TROPICAMIDE  * Eye drops 0.5%  * Eye drops 1%		15 ml OP 15 ml OP		ydriacyl ydriacyl
Preparations for Tear Deficiency				
For acetylcysteine eye drops refer Standard Formulae, page 237 HYPROMELLOSE	,			
* Eye drops 0.5%	19.50	15 ml OP	✓ M	ethopt
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	<b>✓</b> P	oly-Tears

### **Preservative Free Ocular Lubricants**

## **⇒SA1388** Special Authority for Subsidy

Other Eve Preparations

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail phar	macy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authorit	y see SA1388	above – Retail	pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Author	rity see SA138	8 above – Reta	il pharmacy
Eye drops 1 mg per ml	22.00	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Phar	macy Procedu	res Manual res	triction allowing one bottle per
month is not relevant and therefore only the prescribed do	sage to the nea	arest OP may b	oe claimed.

Other Lyc i reparations		
NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.20	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT  * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

# **SENSORY ORGANS**

Pharma S29

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
Various					
PHARMACY SERVICES  May only be claimed once per patient.					
* Brand switch fee	4.50	1 fee		SF Bisoprolol Mylan	
a) The Dharmonde for DCC Davinguis Mulan is 000700	20	•	_	SF Darunavir Mylan	

a) The Pharmacode for BSF Darunavir Mylan is 2607026 - see also page 106

b) The Pharmacode for BSF Bisoprolol Mylan is 2607034 - see also page 50

(BSF Bisoprolol Mylan Brand switch fee to be delisted 1 July 2021)

(BSF Darunavir Mylan Brand switch fee to be delisted 1 July 2021)

# **Agents Used in the Treatment of Poisonings**

### **Antidotes**

ACETVI CVCTEINI	_

#### NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

### Removal and Elimination

### CHARCOAL

*	Oral lig 50 g per 250 n	ıl	250 ml OP	✓ Carbosorb-X

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

## DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable Tab 125 mg dispers

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

## ⇒SA1492 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis; or

continued...



Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE - Special Authority see SA1480 below - Retail pl	harmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

### ⇒SA1480 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

## DESFERRIOXAMINE MESILATE

* Inj 500 mg vial	84.53	10	✓ <u>DBL</u> <u>Desferrioxamine</u> <u>Mesylate for Inj</u> <u>BP</u>
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

# **Standard Formulae**

Standard i Orindiae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Acetylcysteine inj 200 mg per ml, 10 ml	qs	mg per ml)	
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
CODEINE LINOTHO (C		Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)	00	Water	to 40 ml
Codeine phosphate	60 mg	DII OOADDINE ODAL LIQUID	
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg nor 5 ml)		Water	to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)	200 ma	(Preservative should be used if quantity supplied is	for more
Codeine phosphate Glycerol	300 mg 40 ml	than 5 days.)	
Preservative		SALIVA SUBSTITUTE FORMULA	
Water	qs to 100 ml	Methylcellulose	5 g
water	10 100 1111	Preservative	gs qs
FOLINIC MOUTHWASH		Water	to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	ioi iliole
Water	to 500 ml	than 5 days. Maximum 500 mi per prescription.)	
(Preservative should be used if quantity supplied is		SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)	101 111010	Sodium chloride inj 23.4%, 20 ml	qs
man o dayo. Maximum ooo mi por procomption.		Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatr	aemia)
Methadone powder	qs		,
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection	10 vials
		Glycerol BP	40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridiu	um difficile
Propylene glycol	to 100 ml	following metronidazole failure)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	uid mixture)	VOSOL EAR DROPS	
OMEPRAZOLE SUSPENSION		WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder	qs		1%
Sodium bicarbonate powder BP	43 8.4 g	Hydrocortisone powder	to 35 ml
Water	to 100 ml	Vosol Ear Drops	10 35 1111
Water	10 100 1111		
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Water	to 100 ml		

## EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy

Fully

Brand or

	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully Brand or sidised Generic  Manufacturer
Extemporaneously Compounded Preparations	and Galenica	İs	
CODEINE PHOSPHATE – Safety medicine; prescriber may det			
Powder – Only in combination		25 q	
, , , , , , , , , , , , , , , , , , , ,	(90.09)	- 3	Douglas
Only in extemporaneously compounded codeine linctus			
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the determined.	supplier and will be	e delisted fror	n the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination		100 1111	- 1 Om
Only in extemporaneously compounded oral mixtures.			
Soln	30.00	100 ml	✓ <u>Midwest</u>
GLYCERIN WITH SODIUM SACCHARIN - Only in combination	1		
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination			
Only in combination with Ora-Plus. Suspension	20.05	473 ml	✓ Ora-Sweet
	50.95	4/3 1111	▼ <u>Ora-Sweet</u>
GLYCEROL  * Liquid – Only in combination	3 23	500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparation		000 1111	inculting dryocron bi
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing fr			and the second second second
<ul> <li>d) Extemporaneously compounded methadone will only be (methadone powder, not methadone tablets).</li> </ul>	reimbursed at the	rate of the ch	eapest form available
Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE		3	
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE			
Powder		100 g	✓ <u>MidWest</u>
Suspension – Only in combination		473 ml	✓ <u>Ora-Plus</u>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH	•		4.6 51 1.65
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On	,	470	A Ove Bland
Suspension	30.95	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM Powder – Only in combination	52.50	10 g	✓ MidWest
Fowder - Only in combination	325.00	10 g	✓ MidWest
Only in children up to 12 years	0_0.00	.00 9	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxyben:			
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE	10.05	F00	✓ Midweet

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

✓ Midwest

500 g

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>M</u>	<u>lidwest</u>	
WATER Tap - Only in combination	0.00	1 ml	<b>✓</b> Ta	ap water	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

# **Nutrient Modules**

## Carbohydrate

## ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal 

# Carbohydrate And Fat

# ⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Fat

## **⇒SA1523** Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 241

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	200 ml OP	✓ Calogen
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml114.92	4 OP	✓ Liquigen ´

## **Protein**

### ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	<ul> <li>Special Authority see SA1524 above – Hospital p</li> </ul>	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacture

## **Oral and Enteral Feeds**

#### **Diabetic Products**

### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Sp	pecial Authority see SA1095 above – F	Hospital pharm	acy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
	7.50	1,000 ml OP	<ul><li>Diason RTH</li></ul>
			✓ Glucerna Select
			RTH

(Glucerna Select RTH Liquid to be delisted 1 September 2021)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Sustagen Diabetic
	(2.10)	200 ml OP	Nutren Diabetes

(Glucerna Select Liquid (vanilla) to be delisted 1 September 2021) (Sustagen Diabetic Liquid (vanilla) to be delisted 1 October 2021)

### Fat Modified Products

### ⇒SA1525 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 243



Subsidy (Manufacturer's Price) Su

Fully Subsidised

Brand or Generic Manufacturer

## **Paediatric Products For Children Awaiting Liver Transplant**

#### ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

## Paediatric Products For Children With Chronic Renal Failure

### ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

 Liquid
 54.00
 400 g OP
 ✓ Kindergen

 Powder
 54.00
 400 g OP
 ✓ Kindergen

(Kindergen Liquid to be delisted 1 August 2021)

### Paediatric Products

### ⇒SA1379 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2.3 faltering growth in an infant/child; or
- 2.4 increased nutritional requirements; or
- 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see \$ Liquid		he previous pag 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA Liquid		previous page 500 ml OP	<ul><li>− Hospital pharmacy [HP3]</li><li>✓ Nutrini RTH</li><li>✓ Pediasure RTH</li></ul>
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Apharmacy [HP3]	Authority see	SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13	379 on the r	revious page -	Hospital pharmacy [HP3]
Liquid (strawberry)		200 ml OP	✓ Fortini
Liquid (vanilla)		200 ml OP	✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA137	9 on the pre	evious page – H	ospital pharmacy [HP3]
Liquid (chocolate)		200 ml OP	✓ Pediasure
Liquid (strawberry)		200 ml OP	✓ Pediasure
Liquid (vanilla)		200 ml OP	✓ Pediasure
, ,	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authorharmacy [HP3]	ority see SA	1379 on the pre	evious page - Hospital
Liquid (unflavoured)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry)		200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla)		200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the	e previous r	age - Hospital	pharmacy [HP3]
Powder		400 g OP	✓ Peptamen Junior

### **Renal Products**

## ⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

245 ✓ fully subsidised

	Subsidy (Manufacturer's Pri	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see Liquid		evious page – 500 ml OP		al pharmacy [HP3] lepro HP RTH
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		s page – Hos 220 ml OP	· <b>√</b> N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA11(		page – Hospi 237 ml OP	ital pha	rmacy [HP3]
Liquid (apricot) 125 ml Liquid (caramel) 125 ml	(3.31) 11.52	4 OP 4 OP	<b>✓</b> R	lovaSource Renal Renilon 7.5 Renilon 7.5

## **Specialised And Elemental Products**

### ⇒SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LiquidLiquid	•					
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see	SA1377 above	- Hospital phari	macy [HP3]			
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see \$	SA1377 above -	Hospital pharma	acy [HP3]			
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN			
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]						
l iquid '	•		, ,, ,			

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per 🗸 Brand or Generic Manufacturer

# Paediatric Products For Children With Low Energy Requirements

## ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

## Standard Supplements

### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 247

Subs (Manufactur		
\$	Per	Manufacturer

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m<sup>2</sup> and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3]

Liquid.......7.00 1,000 ml OP ✓ Nutrison Energy

✓ fully subsidised 249

	Subsidy		Fully Brand or
	(Manufacturer's		idised Generic
	\$	Per	✓ Manufacturer
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on Liquid		spital pharmacy 250 ml OP 1,000 ml OP	[HP3]  ✓ Isosource Standard  ✓ Nutrison Standard  RTH  ✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit Liquid		on page 247 – F 1,000 ml OP	
ENTERAL FEED WITH FIBRE 1 KCAL/ML — Special Authority so Liquid		0age 247 – Hos 1,000 ml OP	pital pharmacy [HP3]  ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML — Special Authority Liquid		page 247 – Ho 250 ml OP 1,000 ml OP	spital pharmacy [HP3]  Finsure Plus HN  Ensure Plus RTH  Jevity HiCal RTH  Nutrison Energy  Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on page Note: Higher subsidy for Sustagen Hospital Formula will only number and an appropriately endorsed prescription.  Powder (chocolate) – Higher subsidy of up to \$26.00 per 850	be reimbursed	for patients wit	h both a valid Special Authority
with Endorsement	26.00 9.54	850 g OP 840 g OP	✓ Ensure
	(26.00)	3	Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patier prescription must be endorsed accordingly.  Powder (vanilla) – Higher subsidy of up to \$26.00 per 850 g	its with fat mala	bsorption, fat in	tolerance or chyle leak. The
with Endorsement	26.00 9.54	857 g OP 850 g OP 840 g OP	✓ Fortisip ✓ Ensure
	(26.00)		Sustagen Hospital Formula Active
Additional subsidy by endorsement is available for patier	its with fat mala	ibsorption, fat in	tolerance or chyle leak. The

Additional subsidy by endorsement is available for patients with fat malabsorption, fat intolerance or chyle leak. The prescription must be endorsed accordingly.

(Fortisip Powder (vanilla) to be delisted 1 August 2021)

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	- 5
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	Francis Dive
1: :1/	(1.26)		Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
Lituoisement	(1.26)	200 IIII OF	Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3] Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.26)	200 0.	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			·
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

# **High Calorie Products**

### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

continued...

251 ✓ fully subsidised

Subsidy		Fully	Brand or
(Manufacturer's Price)	_ 8	Subsidised	Generic
\$	Per		Manufacturer

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

harmacy [HP3]	ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]			
✓ Nutrison				
Concentrated				
✓ Two Cal HN RTH	1.000 ml OP	11.00		

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

## Food Thickeners

### ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
FOOD THICKENER - Special Authority see SA1106 on	1 0 1	•	,		
Powder		00 g OP		lutilis	
	7.25 3	80 g OP		eed Thickener Karicare Aptamil	

## **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by PHARMAC from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX — Special Authority see SA1729 above — Hospit Powder2.81	al pharmacy [HP3] 1,000 g OP	
(5.15)		Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 above - Hospita	al pharmacy [HP3]	
Powder	1,000 g OP	
(7.32)		NZB Low Gluten Bread Mix
3.51		
(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above - Hospital pha	armacy [HP3]	
Powder	2,000 g OP	
(18.10)		Horleys Flour

✓ fully subsidised 253

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		sidised	Generic
	\$	Per		Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – Ho	ospital pharr	nacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)		C	Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)		C	Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)		C	Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)		C	Orgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)		C	Orgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)		C	Orgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)		C	Orgran

# Foods And Supplements For Inborn Errors Of Metabolism

## ⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

# **Supplements For Homocystinuria**

## **Supplements For MSUD**

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturer

## **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet		30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton		18 OP	<ul> <li>Easiphen Liquid</li> </ul>
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

## Foods

LOW PROTEIN BAKING MIX − Special Authority see SA1108 on the previous page − Hospital pharmacy [HP3]

Powder .......8.22 500 g OP 

Loprofin Mix

LOW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] Animal shapes .......11.91 500 q OP ✓ Loprofin ✓ Loprofin 250 g OP 500 g OP ✓ Loprofin 250 q OP ✓ Loprofin Macaroni 5.95 500 g OP ✓ Loprofin 500 g OP ✓ Loprofin 500 g OP ✓ Loprofin 

✓ fully subsidised

255

Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

## Infant Formulae

## For Williams Syndrome

### ⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder .......44.40 400 g OP ✓ Locasol

## **Gastrointestinal and Other Malabsorptive Problems**

AMINO ACID FORMULA – Special Authority see SA1940 below Powder	,. ,.	✓ Alfamino Junior
Powder (unflavoured)	53.00 400 g OP	✓ Elecare
,	ū	✓ Elecare LCP
		✓ Neocate Gold
		<ul><li>Neocate Junior Unflavoured</li></ul>
		✓ Neocate SYNEO
Powder (vanilla)	53.00 400 g OP	✓ Elecare
. ,	· ·	<ul><li>Neocate Junior Vanilla</li></ul>

#### ⇒SA1940 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short aut: or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Fither:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

**Initial application** — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency: or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Fither:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
      - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
  - 2.1 Patient has non IqE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
  - 2.2 All of the following:
    - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
    - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 2.2.3 Amino acid formula is required for a nutritional deficit; and
    - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

257 ✓ fully subsidised

# SPECIAL FOODS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut: or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
      - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	- Special Authority see SA1953 below	- Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

### ⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea: or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption: or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see \$A1557 below - Hospital pharmacy [HP3]

Powder	15.21	450 g OP	<ul> <li>Aptamil Gold+ Pepti Junior</li> </ul>
	30.42	900 g OP	<ul> <li>Aptamil AllerPro SYNEO 1</li> </ul>
			<ul> <li>Aptamil AllerPro SYNEO 2</li> </ul>

## ⇒SA1557 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula; and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

✓ fully subsidised

259



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3] Liquid.......2.35 125 ml OP ✓ Infatrini

#### ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

# **Ketogenic Diet**

### ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

### **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

# **Vaccinations**

#### BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > 0 requal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

## DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
  Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE			
(1	Subsidy Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	D HAEMOPHILUS I	NFLUENZAE TY	PE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria:			
1) Up to four doses for children up to and under the age of 1			- d d - u d - u - d - d
2) An additional four doses (as appropriate) are funded for (			
10 who are patients post haematopoietic stem cell transp			
post solid organ transplant, renal dialysis and other sever 3) Up to five doses for children up to and under the age of 1	, , , , , , ,	•	
Note: A course of up-to four vaccines is funded for catch up pi		,	
to complete full primary immunisation. Please refer to the Imm			
programmes.	iumsation nanuboor	tion the approprie	ate sofication for eaten up
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10 🗸 In	fanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]		_	
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
<ul><li>2) An additional dose (as appropriate) is funded for (re-)imm</li></ul>	nunisation for patien	ts post haematop	oietic stem cell
transplantation, or chemotherapy; functional asplenic; pre			
or post cochlear implants, renal dialysis and other severe			3
3) For use in testing for primary immunodeficiency diseases	, ,,	•	al medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg;			
prefilled syringe plus vial 0.5 ml	0.00	1 🗸 H	iberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
1) Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver dis-	ease; or		
3) One dose of vaccine for close contacts of known hepatitis	s A cases.		

✓ Havrix

✓ Havrix Junior

1

	NATIONAL	IIVIIVI	UNISATI	JN SCHEDOLL
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe		1	<b>√</b> E	ngerix-B
Funded for patients meeting any of the following criteria  1) for household or sexual contacts of known acute h  2) for children born to mothers who are hepatitis B su  3) for children up to and under the age of 18 years in serology and require additional vaccination or requ  4) for HIV positive patients; or  5) for hepatitis C positive patients; or  6) for patients following non-consensual sexual interce  7) for patients following immunosuppression; or  8) for solid organ transplant patients; or  9) for post-haematopoietic stem cell transplant (HSC)  10) following needle stick injury.	repatitis B patients or harface antigen (HBsAg clusive who are considure a primary course course; or	) posi dered	tive; or not to have	
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following criteria		1	<b>√</b> <u>E</u>	ngerix-B
1) for household or sexual contacts of known acute he go for children born to mothers who are hepatitis B such for children up to and under the age of 18 years in serology and require additional vaccination or request. If the form the f	nepatitis B patients or hurface antigen (HBsAg clusive who are considure a primary course course; or	) posi dered	tive; or not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 8 Any of the following: 1) Maximum of two doses for children aged 14 years and	under; or	- [Xph	arm]	
2) Maximum of three doses for patients meeting any of th  1) People aged 15 to 26 years inclusive; or  2) Either: People aged 9 to 26 years inclusive  1) Confirmed HIV infection; or  2) Transplant (including stem cell) patients: o	r			
3) Maximum of four doses for people aged 9 to 26 years	·			and all o
Inj 270 mcg in 0.5 ml syringe	0.00	10	✓ <u>G</u>	ardasil 9

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

#### INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

#### A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
  - a) ischaemic heart disease, or
  - b) congestive heart failure, or
  - c) rheumatic heart disease, or
  - d) congenital heart disease, or
  - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
  - a) asthma, if on a regular preventative therapy, or
  - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
  - a) autoimmune disease, or
  - b) immune suppression or immune deficiency, or
  - c) HIV, or
  - d) transplant recipients, or
  - e) neuromuscular and CNS diseases/disorders, or
  - f) haemoglobinopathies, or
  - g) on long term aspirin, or
  - h) have a cochlear implant, or
  - i) errors of metabolism at risk of major metabolic decompensation, or
  - i) pre and post splenectomy, or
  - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Sul	Fully osidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable
- C)

#### A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by PHARMAC:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes: or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV. or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - j) pre and post splenectomy, or
    - k) down syndrome, or
  - vii) are pregnant;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine) .......90.00

✓ Fluad Quad (2021 Formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

С

- INFLUENZA VACCINE people 65 years and over
  is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccin	e) –		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic	
\$	Per	<b>√</b>	Manufacturer	

#### A) INFLUENZA VACCINE – people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by PHARMAC:

- i) have any of the following cardiovascular diseases
  - a) ischaemic heart disease, or
  - b) congestive heart failure, or
  - c) rheumatic heart disease, or
  - d) congenital heart disease, or
  - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
  - a) asthma, if on a regular preventative therapy, or
  - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
  - a) autoimmune disease, or
  - b) immune suppression or immune deficiency, or
  - c) HIV, or
  - d) transplant recipients, or
  - e) neuromuscular and CNS diseases/disorders, or
  - f) haemoglobinopathies, or
  - g) are children on long term aspirin, or
  - h) have a cochlear implant, or
  - i) errors of metabolism at risk of major metabolic decompensation, or
  - i) pre and post splenectomy, or
  - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

#### MEASI ES, MUMPS AND BUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

#### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

# ${\sf MENINGOCOCCAL} \ ({\sf GROUPS} \ {\sf A}, \ {\sf C}, \ {\sf Y} \ {\sf AND} \ {\sf W-135}) \ {\sf CONJUGATE} \ {\sf VACCINE} \ -[{\sf Xpharm}]$

Fither:

- A) Any of the following:
  - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
  - 2) One dose for close contacts of meningococcal cases; or
  - 3) A maximum of two doses for bone marrow transplant patients; or
  - 4) A maximum of two doses for patients following immunosuppression\*; or
- B) Both
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either:
    - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
    - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Ini 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

✓ Synflorix

10

	Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
	\$	Per 🗸	Manufacturer
MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm]			
Both:			
1) The child is under 9 months of age; and			
<ol><li>Any of the following:</li></ol>			
<ol> <li>Up to three doses for patients pre- and post spler HIV, complement deficiency (acquired or inherited</li> <li>Two doses for close contacts of meningococcal c</li> <li>A maximum of two doses for bone marrow transp</li> <li>A maximum of two doses for patients pre- and po</li> </ol>	d), or pre or post solid ases; or lant patients; or	organ transplant	
Note: children under nine months of age require two d booster schedules with meningococcal ACWY vaccine.		Refer to the Imm	unisation Handbook for
*Immunosuppression due to steroid or other immunosu	ippressive therapy mu	ist be for a period	d of greater than 28 days.
Inj 10 mcg in 0.5 ml syringePNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm		1 <b>~</b> N	leisvac-C
A primary course of three doses for previously unvaccing	•	the age of 59 m	onths inclusive
Note: please refer to the Immunisation Handbook for the ap	propriate schedule for	catch up progra	mmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B, 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer

### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously
  received two doses of the primary course of PCV10: or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) with primary immune deficiencies; or
  - c) with HIV infection; or
  - d) with renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) with cochlear implants or intracranial shunts; or
  - g) with cerebrospinal fluid leaks; or
  - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) with cardiac disease, with cyanosis or failure; or
  - I) with diabetes; or
  - m) with Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISAT	ION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE	– [Xpharm]		
Either:			
<ol> <li>Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with funcomplement deficiency (acquired or inherited), coch</li> <li>All of the following:</li> </ol>	ctional asplenia, pre- or p	oost-solid organ	transplant, renal dialysis,
<ul> <li>a) Patient is a child under 18 years for (re-)immur</li> <li>b) Treatment is for a maximum of two doses; and</li> <li>c) Any of the following:</li> </ul>	· ·		
<ul> <li>i) on immunosuppressive therapy or radiati immune response; or</li> </ul>	ion therapy, vaccinate wl	nen there is expe	ected to be a sufficient
<ul><li>ii) with primary immune deficiencies; or</li><li>iii) with HIV infection; or</li></ul>			
iv) with renal failure, or nephrotic syndrome;	or		
v) who are immune-suppressed following or		uding haematop	oietic stem cell transplant);
or vi) with cochlear implants or intracranial shu	inte: or		
vii) with cerebrospinal fluid leaks; or	into, or		
viii) receiving corticosteroid therapy for more			
prednisone of 2 mg/kg per day or greater	r, or children who weigh	more than 10 kg	on a total daily dosage of
20 mg or greater; or ix) with chronic pulmonary disease (includin	g asthma treated with hig	ah-dose corticos	teroid therapy): or
x) pre term infants, born before 28 weeks g		g., acco cocc	10.0.u 1o. up///, 0.
xi) with cardiac disease, with cyanosis or fai	lure; or		
xii) with diabetes; or			
xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	functional asplenia		
xiv) who are pre or poor opionocionity, or with	Turiotional aspionia.		
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each			
23 pneumococcal serotype)	0.00	1 🗸	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm]			
Up to three doses for patients meeting either of the follow			
<ol> <li>For partially vaccinated or previously unvaccinated i</li> <li>For revaccination following immunosuppression.</li> </ol>	ndividuals; or		
Note: Please refer to the Immunisation Handbook for app	propriate schedule for ca	tch-up programn	100
Inj 80D antigen units in 0.5 ml syringe			POL
ROTAVIRUS ORAL VACCINE - [Xpharm]		-	<del></del>
Maximum of two doses for patients meeting the following:			
<ol> <li>first dose to be administered in infants aged under 1</li> <li>no vaccination being administered to children aged 2</li> </ol>			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

		Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
VARICELLA V Either:	ACCINE [CHICKENPOX VACCINE] - [Xpharm]				
1) Maxi	imum of one dose for primary vaccination for eithe	er:			
,	Any infant born on or after 1 April 2016; or For previously unvaccinated children turning 11 varicella infection (chickenpox), or	years old on or after 1	July 201	7, who h	nave not previously had a
	imum of two doses for any of the following:				
a)	Any of the following for non-immune patients:				
	ii) with chronic liver disease who may in future     iii) with deteriorating renal function before tran     iii) prior to solid organ transplant; or	splantation; or	nsplanta	ition; or	
	iv) prior to any elective immunosuppression*,				
L١	v) for post exposure prophylaxis who are imm			_:::	-liet
,	For patients at least 2 years after bone marrow to For patients at least 6 months after completion of				·
	For HIV positive non immune to varicella with mi				
	For patients with inborn errors of metabolism at r varicella, or				
,	For household contacts of paediatric patients wh immune compromise where the household conta For household contacts of adult patients who have	ct has no clinical histo	ry of var	ricella, or	
9/	immunocompromised, or undergoing a procedur has no clinical history of varicella.				
28 days	suppression due to steroid or other immunosuppre	.,	for a tre	eatment p	period of greater than
Inj 1350 P	FU prefilled syringe	0.00	1	<b>✓</b> \	/arivax
			10	✓ \	<u>/arivax</u>
	OSTER VIRUS (OKA STRAIN) LIVE ATTENUAT r patients meeting either of the following criteria:	ED VACCINE [SHING	LES VA	CCINE]	– [Xpharm]
1) One	dose for all people aged 65 years; or				
	dose for all people aged between 66 and 80 year	s inclusive from 1 Apri	l 2018 a	nd 31 De	ecember 2021.
Ini 19.400	PFU prefilled syringe plus vial	0.00	1	<b>√</b> 2	ostavax
,,	- F		10		ostavax
Diagnosti	c Agents				
	PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU po	er 0.1 ml, 1 ml vial	0.00	1	✓ ]	<u>ubersol</u>

- Symbols -	Afluria Quad Junior	Amoxicillin with clavulanic acid92
UK Synacthen80	(2021 Formulation) 264	Amphotericin B32
3TC106	AFT-Pyrazinamide100	Amsacrine149
- A -	Agents Affecting the	AmsaLyo149
A-Scabies68	Renin-Angiotensin System 47	Amsidine149
Abacavir sulphate 105	Agents for Parkinsonism and Related	Amzoate30
Abacavir sulphate with	Disorders 118	Anaesthetics119
lamivudine 106	Agents Used in the Treatment of	Anagrelide hydrochloride149
Abiraterone acetate163	Poisonings235	Analgesics120
Acarbose11	Agrylin149	Anastrozole165
Accarb11	Albendazole89	Anatrole
Accuretic48	Albey220	Andriol Testocaps80
Accuretic 1048	Albustix77	Androderm80
Accuretic 2048	Aldurazyme28	ANI50
Acetazolamide232	Alecensa156	Anoro Ellipta224
Acetec47	Alectinib156	Antabuse141
Acetic acid with 1, 2- propanediol	Alendronate sodium111	Antacids and Antiflatulents6
diacetate and	Alendronate sodium with	Anthelmintics89
benzethonium229	colecalciferol111	Antiacne Preparations61
Acetic acid with hydroxyquinoline and	Alfacalcidol33	Antiallergy Preparations220
ricinoleic acid	Alfamino Junior256	Antianaemics37
Acetylcysteine235	Alginic acid6	Antiandrogen Oral
Aci-Jel75	Alglucosidase alfa26	Contraceptives75
Aciclovir	Alkeran146	Antiarrhythmics49
Infection101	Alkeran S29146	Antibacterials89
Sensory229	Allersoothe221	Antibacterials Topical61
Acidex6	Allmercap148	Anticholinergic Agents
Acipimox55	Allopurinol116	Anticholinesterases
Acitretin	Alpha-Adrenoceptor Blockers47	Antidepressants
Aclasta114	Alpha-Keri Lotion	Antidiarrhoeals6
Aclin	Alphamox92	Antiepilepsy Drugs
Actemra209	Alphamox 12592	Antifibrinolytics, Haemostatics and
Actinomycin D150	Alphamox 25092	Local Sclerosants38
Actrapid10	Alprolix38	Antifibrotics224
Actrapid Penfill10	Alu-Tab6	Antifungals96
Acular231	Aluminium hydroxide6	Antifungals Topical62
Acupan	Alvogen51	Antihistamines221
Adalat 1052	Amantadine hydrochloride118	Antihypotensives50
Adalat Oros52	Ambrisentan57	Antimalarials98
Adalimumab173	Ambrisentan Mylan57	Antimigraine Preparations129
Adapalene61	Amiloride hydrochloride53	Antinausea and Vertigo Agents 129
Adcortyl80	Amiloride hydrochloride with	Antiparasitics99
Adefin52	furosemide54	Antipruritic Preparations63
Adefin XL52	Amiloride hydrochloride with	Antipsychotics130
Adenuric	hydrochlorothiazide54	Antiretrovirals104
ADR Cartridge 1.823	Aminophylline226	Antirheumatoid Agents111
Adrenaline56	Amiodarone hydrochloride49	Antispasmodics and Other Agents
Advantan64	Amisulpride130	Altering Gut Motility8
Advate41	Amisulpride Mylan130	Antithrombotic Agents41
Adynovate41	Amitriptyline124	Antithymocyte globulin
Afinitor217	Amlodipine52	(equine)
Aflibercept	Amneal	Antitrichomonal Agents99
Afluria Quad	Amorolfine	Antituberculotics and
(2021 Formulation)	Amoxicillin92	Antileprotics
,		•

Antiulcerants	8	Hydrochlorothiazide	48	vaccine	26
Antivirals	101	Arrow-Norfloxacin	109	Baclofen	11
Anxiolytics	134	Arrow-Ornidazole		Bactroban	6
Anzatax		Arrow-Quinapril 10	48	Balance	2
Apidra	11	Arrow-Quinapril 20		Barrier Creams and Emollients	6
Apidra SoloStar		Arrow-Quinapril 5		Basic AquaCream	6
Apo-Azithromycin		Arrow-Roxithromycin		BCG Vaccine	
Apo-Bromocriptine		Arrow-Sertraline		Beclazone 100	
Apo-Ciclopirox		Arrow-Timolol		Beclazone 250	22
Apo-Clarithromycin		Arrow-Topiramate		Beclazone 50	
Alimentary	9	Arrow-Tramadol		Beclomethasone dipropionate	
Infection		Arsenic trioxide		Bee venom allergy treatment	
Apo-Clomipramine		Asacol		Bendamustine hydrochloride	
Apo-Diclo SR		Asamax		Bendrofluazide	
Apo-Diltiazem CD		Ascorbic acid		Bendroflumethiazide	
Apo-Doxazosin		Aspen Adrenaline		[Bendrofluazide]	5
Apo-Folic Acid		Aspirin		Benzathine benzylpenicillin	
Apo-Furosemide		Blood	41	Benzatropine mesylate	
Apo-Gabapentin		Nervous		Benzbromaron AL 100	
		Asthalin		Benzbromarone	
Apo-Megestrol		Atazanavir sulphate		Benztrop	
				•	
Apo-Mirtazapine		Atenolol		Benzydamine hydrochloride	
Apo-Nadolol		Atenolol AFT		Benzylpenicillin sodium [Penicillin	
Apo-Oxybutynin		Atenolol AFT S29		G]	
Apo-Perindopril		ATGAM		Beta Cream	
Apo-Pindolol		Ativan		Beta Ointment	
Apo-Prazosin		Atomoxetine		Beta Scalp	
Apo-Prednisone		Atorvastatin	55	Beta-Adrenoceptor Agonists	
Apo-Primidone		Atropine sulphate		Beta-Adrenoceptor Blockers	
Apo-Propranolol		Cardiovascular		Betadine	
Apo-Pyridoxine		Sensory		Betadine Skin Prep	
Apo-Selegiline S29		Atropt		Betaferon	
Apo-Sumatriptan		Atrovent		Betahistine dihydrochloride	13
Apo-Temozolomide		AU Synacthen		Betaine	2
Apo-Terazosin		Aubagio	135	Betaloc CR	5
Apo-Timol	52	Augmentin	92	Betamethasone dipropionate	6
Apomorphine hydrochloride	118	Aurorix	124	Betamethasone dipropionate with	1
Aprepitant	129	AutoSoft 30	<mark>23</mark>	calcipotriol	6
Apresoline	56	AutoSoft 90	<mark>23</mark>	Betamethasone sodium phospha	ıte
Aptamil AllerPro SYNEO 1	259	Avelox	94	with betamethasone acetate	<mark>7</mark>
Aptamil AllerPro SYNEO 2	259	Avonex	135	Betamethasone valerate	63, 6
Aptamil Gold+ Pepti Junior	259	Avonex Pen	135	Betamethasone valerate with soc	muit
Aqueous cream	65	Azacitidine	146	fusidate [fusidic acid]	6
Aratac	49	Azacitidine Dr Reddy's	146	Betaxolol	23
Arava	111	Azamun		Betnovate	6
Arginine	27	Azathioprine		Betoptic	23
Aripiprazole		Azithromycin	90	Betoptic S	
Aripiprazole Sandoz		Azopt		Bezafibrate	5
Aristocort		AZT		Bezalip	
Arrow - Lattim		-B-		Bezalip Retard	5
Arrow-Amitriptyline		B-D Micro-Fine	15	Bicalutamide	
Arrow-Bendrofluazide		B-D Ultra Fine		Bicillin LA	Q
Arrow-Brimonidine		B-D Ultra Fine II		BiCNU	
Arrow-Diazepam		Bacillus Calmette-Guerin (I		Bicnu Heritage	
Arrow-Doxorubicin		vaccine		Bile and Liver Therapy	
Arrow-Losartan &	100	Bacillus Calmette-Guerin	173	Biltricide	
AITOW-LOSAITAIT &		Dacinus Cannette-Guellii		Ditt ICIUE	0

				_
Bimatoprost	232	Cabergoline	kin Sandoz	. 89
Bimatoprost Multichem	232	Cacit34 Cefazo	lin	. 89
Binarex	163	Caffeine citrate228 Ceftriax	xone	. 89
Binocrit	38	Calamine63 Ceftriax	xone-AFT	. 89
Biodone	122		xime axetil	
Biodone Extra Forte	122		ex1	
Biodone Forte	122		xib1	
Bisacodyl			xib Pfizer1	
Bisoprolol fumarate			one Chronodose	.79
Bisoprolol Mylan			pt1	
BK Lotion			lly-Acting Agents	
Bleomycin sulphate			exin ABM	
Blood Colony-stimulating		•	ne hydrochloride2	
Factors	44		acrogol	.65
Blood glucose diagnostic test			acrogol with glycerol	
meter	14		nab1	
Blood glucose diagnostic test			oix1	
strip	14		oal2	
Blood glucose test strips (visually			therapeutic Agents1	
impaired)	14		npox vaccine2	
Blood Ketone Diagnostic Test	14		fast	
Strip	13		mbucil1	
Bonjela			mphenicol	
Boostrix			thiazide	
Bortezomib			heniramine maleate2	
Bortezomib Dr-Reddy's			romazine hydrochloride1	
Bosentan			llidone [Chlorthalidone]	
Bosentan Dr Reddy's		·	ialidone	
Bplex			escent	
Breo Ellipta			Load 375	
Brevinor 1/28			TT380 Short	
Bricanyl Turbuhaler			TT380 Standard	
Brilinta			e salicylate with cetalkonium	
Brimonidine tartrate			ride	20
Brimonidine tartrate with timolol	202		rox olamine	
maleate	222		porin2	
Brinzolamide			oril	
Brolene			ie	
Bromocriptine mesylate			ie VK	
BSF Bisoprolol Mylan		•	lcet	
BSF Darunavir Mylan				
Buccastem		Cardinol LA		. 5-
Budesonide	100		ction	۵/
Alimentary	6		sory2	
			oxacin Teva2	
Respiratory22			n1	
Budesonide with eformoterol  Bumetanide			in1	
Buprenorphine Naloxone BNM			in Ebewe1	
Buprenorphine with naloxone			ram hydrobromide	
Bupropion hydrochloride			ine1	147
Burinex		Carvedilol		,
Burinex S29			entary	
Buscopan	ŏ		ction	
Buspirone hydrochloride			ee Forte	
Busulfan	145	•		
- 6 -		Cefalexin89 Climara	a	۰٥۱

Climara	81	Condyline	70	David One Step Cassette Pregna	ancy
Clindamycin	94	Contraceptives - Hormonal		Test	
Clinicians	27	Contraceptives - Non-hormonal.	71	DBL Acetylcysteine	
Clinicians Renal Vit	33	Copaxone	135	DBL Adrenaline	56
Clobazam	126	Corticosteroids and Related Age		DBL Aminophylline	
Clobetasol propionate	63, 69	for Systemic Use		DBL Bleomycin Sulfate	
Clobetasone butyrate		Corticosteroids Topical		DBL Carboplatin	
Clofazimine		Cortifoam		DBL Cisplatin	14
Clomazol		Cosentyx		DBL Dacarbazine	
Dermatological	62	Cosmegen		DBL Desferrioxamine Mesylate f	
Genito-Urinary		Coumadin		BP	
Clomifene citrate		Country Life	29	DBL Docetaxel	150
Clomipramine hydrochloride	124	Coversyl		DBL Ergometrine	7
Clonazepam		Creon 10000		DBL Gemcitabine	
Clonidine		Creon 25000	24	DBL Gentamicin	
Clonidine BNM		Creon Micro		DBL Heparin Sodium	
Clonidine hydrochloride		Crotamiton		DBL Leucovorin Calcium	
Clopidogrel		Crystaderm		DBL Methotrexate Onco-Vial	
Clopidogrel Multichem		Curam		DBL Morphine Sulphate	
Clopine		Curam Duo 500/125		DBL Naloxone Hydrochloride	
Clopixol		Cvite		DBL Octreotide	
Clotrimazole	, , , , , ,	Cyclizine hydrochloride		DBL Pethidine Hydrochloride	
Dermatological	62	Cyclizine lactate		DBL Vinblastine	
Genito-Urinary		Cyclogyl		DBL Vincristine Sulfate	
Clozapine		Cyclopentolate hydrochloride		Decozol	
Clozaril		Cyclophosphamide		Deferasirox	
Co-trimoxazole		Cyclorin		Deferiprone	
Coal tar		Cycloserine		Denosumab	11
Coal tar with allantoin, menthol		Cyproterone acetate		Deolate	
phenol and sulphur		Cyproterone acetate with		Deoxycoformycin	
Coal tar with salicylic acid and		ethinyloestradiol	75	Depo-Medrol	
sulphur	69	Cystadane		Depo-Provera	
Coco-Scalp		Cytarabine		Depo-Testosterone	
Codeine phosphate		Cytotec		Deprim	
Extemporaneous	238	Cytoxan		Dermol	
Nervous		- D -		Desferrioxamine mesilate	,
Coenzyme Q10		D-Penamine	111	Desmopressin	
Colchicine		Dabigatran		Desmopressin acetate	
Colecalciferol		Dacarbazine		Desmopressin-PH&T	
Colestid		Dacarbazine APP		Desuric	
Colestipol hydrochloride		Dactinomycin [Actinomycin D]		Detection of Substances in	
Colgout		Daivobet		Urine	7
Colifoam		Daivonex		Dexamethasone	/ 1
Colistin sulphomethate		Daktarin		Hormone	70
Colistin-Link		Dalacin C		Sensory	
Collodion flexible		Dantrium		Dexamethasone phosphate	
Colloidal bismuth subcitrate		Dantrium S29		Dexamethasone Phosphate	
Colofac		Dantrolene		Panpharma	70
Coloxyl		Daonil		Dexamethasone with framycetin	
Combigan		Dapa-Tabs		gramicidin	
Compound electrolytes		Dapsone		Dexamethasone with neomycin	44
Compound electrolytes with glu		Daraprim		sulphate and polymyxin B	
[Dextrose]		DaraprimDaraprim		sulphate	221
Compound hydroxybenzoate		Darunavir Mylan		Dexamfetamine sulfate	ادے
Concerta		Dasatinib		Dexamethsone	
Condoms		Dasatinib		Dextrochlorpheniramine	/
OUTIQUITIS	/ _	Dauliulubiciii	150	Dexilocillorpheniranine	

maleate	221	Dortimopt		Emend Tri-Pack	129
Dextrose	45–46	Dorzolamide hydrochloride		Emicizumab	
DHC Continus	121	Dorzolamide with timolol	232	EMLA	
Diabetes	10	Dostinex	88	Empagliflozin	12
Diabetes Management		Dosulepin [Dothiepin]		Empagliflozin with metformin	
Diacomit	128	hydrochloride	124	hydrochloride	
Diagnostic Agents		Dosulepin Mylan		Emtricitabine	106
Diamide Relief	6	Dothiepin	124	Emtricitabine with tenofovir	
Diamox		Doxazosin	47	disoproxil	103
Diasip	243	Doxine		Emtriva	
Diason RTH	243	Doxorubicin Ebewe	150	Emulsifying ointment	65
Diazepam 120	6, 134	Doxorubicin hydrochloride	150	Emulsifying Ointment ADE	65
Diazoxide	10	Doxycycline	93	Enalapril maleate	47
Dibenzyline	47	DP Lotion	66	Enbrel	
Diclofenac Sandoz	110	DP Lotn HC	64	Endocrine Therapy	163
Diclofenac sodium		DP-Allopurinol	116	Endoxan	
Musculoskeletal		Dr Reddy's Omeprazole	9	Engerix-B	263
Sensory	230	Drugs Affecting Bone		Enlafax XR	125
Differin		Metabolism	111	Enoxaparin sodium	43
Difflam	32	Dual blood glucose and blood ket	one	Enstilar	68
Diflucan	96	diagnostic test meter	13	Ensure	250
Diflucortolone valerate	64	Duocal Super Soluble Powder		Ensure Plus	251
Digestives Including Enzymes	24	Duolin	223	Ensure Plus HN	250
Digoxin		Duolin HFA	223	Ensure Plus RTH	250
Dihydrocodeine tartrate	121	DuoResp Spiromax	222	Entacapone	118
Dilantin	127	Duride	56	Entapone	
Dilantin Infatab	127	-E-		Entecavir	101
Diltiazem hydrochloride	52	e-chamber La Grande	228	Entecavir Sandoz	101
Dilzem		e-chamber Mask	227	Entocort CIR	6
Dimethicone	65–66	e-chamber Turbo	228	Entresto 24/26	
Dimethyl fumarate	135	E-Mycin	91	Entresto 49/51	
Dipentum		Ear Preparations		Entresto 97/103	
Diphtheria, tetanus and pertussis		Ear/Eye Preparations		Epilim	128
vaccine	261	Easiphen Liquid		Epilim Crushable	128
Diphtheria, tetanus, pertussis and		Econazole nitrate		Epilim IV	
polio vaccine	261	Efavirenz		Epilim S/F Liquid	
Diphtheria, tetanus, pertussis, polici		Efavirenz with emtricitabine and		Epilim Syrup	
hepatitis B and haemophilus	-,	tenofovir disoproxil	106	Epirubicin Ebewe	
influenzae type B vaccine	262	Eformoterol fumarate		Epirubicin hydrochloride	
Diprosone		Eformoterol fumarate dihydrate		Eplerenone	
Diprosone OV		Eftrenonacog alfa [Recombinant		Epoetin alfa	
Dipyridamole		factor IX]	38	Epoprostenol	59
Disopyramide phosphate		Efudix		Eptacog alfa [Recombinant factor	
Disulfiram		Egopsoryl TA		VIIa]	40
Diuretics		Elaprase		ERA	
Docetaxel		Elecare		Erbitux	
Docetaxel Accord		Elecare LCP		Ergometrine maleate	
Docetaxel Sandoz		Electral		Erlotinib	157
Docusate sodium		Elelyso		Erythrocin IV	91
Docusate sodium with		Elemental 028 Extra		Erythromycin (as lactobionate)	
sennosides	25	Elidel		Erythromycin ethyl succinate	
Dolutegravir		Elocon		Erythromycin stearate	
Domperidone		Elocon Alcohol Free		Esbriet	22
Donepezil hydrochloride		Eltrombopag		Escitalopram	
Donepezil-Rex		Eltroxin		Escitalopram (Ethics)	
Dornase alfa		EMB Fatol		Escitalopram-Apotex	
Domado ana	۲۲0	LIND I alvi	100	Lookalopiani Apolex	140

Eskazole	89	Ferric carboxymaltose	34	vilanterol	22
Essential Prednisolone	8	Ferriprox	236	Fluticasone propionate	22
Estradiol TDP Mylan	81	Ferro-F-Tabs	35	Fluticasone with salmeterol	
Estradot	81	Ferro-tab	35	FML	23
Estradot 50 mcg	81	Ferrograd	35	Foban	
Estrofem	81	Ferrosig	35	Folic acid	3
Etanercept	166	Ferrous fumarate	35	Food Thickeners	
Ethambutol hydrochloride	100	Ferrous fumarate with folic acid	35	Foods And Supplements For Inbor	rn
Ethics Aspirin	120	Ferrous sulfate	35	Errors Of Metabolism	25
Ethics Aspirin EC	41	Ferrous sulphate	35	Foradil	22
Ethics Lisinopril		Fexofenadine hydrochloride	221	Forteo	11
Ethics Paracetamol Classic	121	Fibro-vein	41	Fortini	24
Ethinyloestradiol	82	Filgrastim	44	Fortini Multi Fibre	24
Ethinyloestradiol with		Finasteride		Fortisip25	50-25
desogestrel	73	Fingolimod	135	Fortisip Multi Fibre	25
Ethinyloestradiol with		Firazyr		Fosamax	
levonorgestrel	74	Flagyl	99	Fosamax Plus	
Ethinyloestradiol with		Flagyl-S		Framycetin sulphate	22
norethisterone	74	Flamazine		Frisium	
Ethosuximide		Flecainide acetate		Frumil	
Etopophos		Flecainide BNM	49	Frusemide	5
Etoposide		Flecainide Controlled Release		Fucicort	6
Etoposide phosphate		Teva	49	Fucidin	9
Etravirine		Fleet Phosphate Enema	26	Fucithalmic	
Eumovate		Flixonase Hayfever & Allergy		Fulvestrant	
Everet	127	Flixotide		Fungilin	3
Everolimus	217	Flixotide Accuhaler		Furosemid-Ratiopharm	5
Evista		Florinef	79	Furosemide [Frusemide]	
Exemestane	165	Fluad Quad		Furosemide-Baxter	
Exjade	235	(2021 Formulation)	265	fusidic acid	
Extemporaneously Compounded		Fluanxol		Dermatological	62, 6
Preparations and		Flucil	93	Infection	9
Galenicals	238	Flucloxacillin	93	Sensory	22
Eye Preparations		Flucloxin		´ - G -	
Eylea		Flucon	231	Gabapentin	12
Ezetimibe		Fluconazole	96	Gacet	
Ezetimibe Sandoz	55	Fludara Oral	147	Galsulfase	2
Ezetimibe with simvastatin	56	Fludarabine Ebewe	147	Galvumet	1
-F-		Fludarabine phosphate		Galvus	1
Factor eight inhibitor bypassing		Fludrocortisone acetate		Gardasil 9	
fraction	40	Fluids and Electrolytes		Gastrodenol	1
Famotidine	9	Flumetasone pivalate		Gaviscon Double Strength	
Famotidine Hovid	9	Fluocortolone caproate with		Gaviscon Infant	
Faslodex	164	fluocortolone pivalate and		Gazyva	19
Febuxostat	116	cinchocaine	8	Gefitinib	
Feed Thickener Karicare		Fluorometholone	231	Gemcitabine Ebewe	14
Aptamil	253	Fluorouracil	147	Gemcitabine hydrochloride	
FEIBA NF	40	Fluorouracil Ebewe	147	Genoptic	
Felo 10 ER		Fluorouracil sodium		Gentamicin sulphate	
Felo 5 ER		Fluox		Infection	9
Felodipine		Fluoxetine hydrochloride		Sensory	
Femme-Tab ED		Flupenthixol decanoate		Gilenya	
Fentanyl		Flutamide	163	Ginet	7
Fentanyl Sandoz		Flutamin		Glatiramer acetate	
Ferinject		Fluticasone		Glecaprevir with pibrentasvir	
Ferodan		Fluticasone furoate with		Glibenclamide	1

Gliclazide12	Hepatitis B recombinant	Hyperuricaemia and Antigout	
Glipizide12	vaccine 263	Hypromellose	
Glivec158	Herceptin212	Hypromellose with dextran	23
Glizide12	Hiberix262	-1-	
Glucagen Hypokit10	Hiprex109	Ibiamox	9
Glucagon hydrochloride10	Histaclear221	Ibrance	160
Glucerna Select243	Histafen221	Ibuprofen	110
Glucerna Select RTH243	Holoxan145	Ibuprofen SR BNM	110
Glucobay11	Horleys Bread Mix253	Icatibant	22
Glucose [Dextrose]45	Horleys Flour253	Idarubicin hydrochloride	15
Gluten Free Foods253	Hormone Replacement Therapy -	Idursulfase	2
Glycerin with sodium saccharin238	Systemic 80	Ifosfamide	14
Glycerin with sucrose238	HPV263	Igroton	5
Glycerol	Humalog11	lkorel	5
Alimentary26	Humalog Mix 2511	llevro	23
Extemporaneous238	Humalog Mix 5011	lloprost	5
Glyceryl trinitrate	Human papillomavirus (6, 11, 16, 18,	Imatinib mesilate	15
Alimentary8	31, 33, 45, 52 and 58) vaccine	Imatinib-Rex	15
Cardiovascular56	[HPV]263	Imigran	12
Glycopyrronium224	Humatin95	Imipramine hydrochloride	12
Glycopyrronium bromide8	Humira173	Imiquimod	
Glycopyrronium with	HumiraPen173	Immune Modulators	
indacaterol224	Humulin 30/7011	Immunosuppressants	
Go Healthy27	Humulin NPH11	Imuran	
Gold Knight72	Humulin R10	Incruse Ellipta	
Gold Knight XL72–73	Hyaluronic acid233	Indacaterol	
Goserelin87	Hydralazine56	Indapamide	
Gutron50	Hydralazine hydrochloride56	Infanrix IPV	
Gynaecological Anti-infectives	Hydrocortisone	Infanrix-hexa	26
́ -Н-	Dermatological64	Infant Formulae	25
Habitrol142	Hormone79	Infatrini	26
Haemophilus influenzae type B	Hydrocortisone (PSM)64	Infliximab	18
vaccine	Hydrocortisone acetate7	Influenza vaccine	
Haldol132	Hydrocortisone acetate with	Influvac Tetra	
Haldol Concentrate132	pramoxine hydrochloride	(2021 Formulation)	26
Haldol Decanoas	Hydrocortisone and paraffin liquid	Inhaled Corticosteroids	
Haloperidol131	and lanolin64	Inhaled Long-acting	
Haloperidol decanoate132	Hydrocortisone butyrate64, 69	Beta-adrenoceptor Agonists	22
Harvoni102	Hydrocortisone with cinchocaine8	Inspra	
Havrix262	Hydrocortisone with miconazole64	Instillagel Lido	
Havrix Junior262	Hydrocortisone with natamycin and	Insulin aspart	
Haylor syrup221	neomycin 64	Insulin aspart with insulin aspart	
healthE Calamine Aqueous Cream	Hydrogen peroxide61	protamine	10
BP63	Hydroxocobalamin33	Insulin glargine	
healthE Dimethicone 10%65	Hydroxocobalamin Mercury	Insulin glulisine	
healthE Dimethicone 4% Lotion 66	Pharma33	Insulin isophane	
healthE Dimethicone 5%65	hydroxycarbamide151	Insulin isophane with insulin	
healthE Glycerol BP238	Hydroxychloroquine111	neutral	1:
healthE Urea Cream	Hydroxyurea	Insulin lispro	
Healtheries Simple Baking Mix253	[hydroxycarbamide]	Insulin lispro with insulin lispro	
Hemastix	Hygroton54	protamine	1:
Hemlibra	Hylo-Fresh233	Insulin neutral	10
Heparin sodium43	Hymenoptera220	Insulin pen needles	
Heparinised saline43	Hyoscine butylbromide8	Insulin pump	
Heparon Junior244	Hyoscine hydrobromide130	Insulin pump cartridge	
	,		
Hepatitis A vaccine262	Hypam 136	Insulin pump infusion set (steel	

cannula)	20	Jaydess	82	Lasix	53
Insulin pump infusion set (steel		Jevity HiCal RTH	250	Latanoprost	232
cannula, straight insertion)	21	Jevity RTH	250	Latanoprost with timolol	232
Insulin pump infusion set (teflon		Juno Pemetrexed		Lax-Suppositories	26
cannula)	22	- K -		Lax-Tab	
Insulin pump infusion set (teflon		Kadcyla	214	Laxatives	25
cannula, angle insertion with		Kaletra		Laxsol	25
insertion device)	23	Kalydeco		Ledipasvir with sofosbuvir	102
Insulin pump infusion set (teflon		Kemadrin		Leflunomide	
cannula, angle insertion)	23	Kenacomb		Lenalidomide	
Insulin pump infusion set (teflon		Kenacort-A 10		Letrole	
cannula, straight insertion with		Kenacort-A 40		Letrozole	
insertion device)	23	Kenalog		Leukeran FC	
Insulin pump infusion set (teflon	20	Kenalog in Orabase		Leukotriene Receptor	170
cannula, straight insertion)	23	Kenkay Sorbolene			226
				Antagonists	
Insulin pump reservoir	23	Ketocal 3:1		Leuprorelin	
Insulin syringes, disposable with	45	KetoCal 4:1	200	Leustatin	
attached needle		Ketoconazole	00	Levetiracetam	
Intal Forte CFC Free		Dermatological		Levetiracetam-AFT	
Intelence		Infection		Levlen ED	
Interferon beta-1-alpha		Ketogenic Diet		Levocabastine	
Interferon beta-1-beta		Ketoprofen		Levocarnitine	
Intra-uterine device		Ketorolac trometamol		Levodopa with benserazide	
Invega Sustenna		KetoSens	13	Levodopa with carbidopa	
IPOL	271	Keytruda	215	Levomepromazine	131
Ipratropium bromide22	3, 227	Kindergen	244	Levomepromazine	
Iressa	158	Kivexa		hydrochloride	131
Irinotecan Accord	148	Klacid	90	Levonorgestrel	
Irinotecan Actavis 100	148	Kliogest	82	Genito-Urinary	74–75
Irinotecan hydrochloride	148	Kliovance	82	Hormone	82
Irinotecan-Rex	148	Kogenate FS	41	Levothyroxine	83
Iron polymaltose	35	Konakion MM	41	Lidocaine [Lignocaine]	119-120
Isentress	106	Konsyl-D	25	Lidocaine [Lignocaine]	
Isentress HD	106	Kuvan		hydrochloride	120
Ismo 20	56	-L-		Lidocaine [Lignocaine] with	
Ismo 40 Retard	56	Labetalol	51	chlorhexidine	120
Isoniazid		Lacosamide		Lidocaine [Lignocaine] with	
Isoniazid with rifampicin		Lactulose		prilocaine	120
Isoptin		Laevolac		Lidocaine-Baxter	
Isoptin Retard		Lamictal		Lidocaine-Claris	
Isoptin SR		Lamivudine		Life Extension	
Isopto Carpine		Lamivudine Alphapharm		Lignocaine	
Isosorbide mononitrate		Lamotrigine		Lioresal Intrathecal	
Isosource Standard		Lamprene		Lipid-Modifying Agents	
Isotretinoin		Lanoxin		Liquigen	
Ispaghula (psyllium) husk		Lanoxin Paediatric Elixir		Lisinopril	
Itch-Soothe		Lanoxin PG		Lithium carbonate	
Itraconazole		Lanoxin S29		Livostin	
ltrazole		Lansoprazole		LMX4	
lvacaftor		Lantus Colo Stor		Locacorten-Viaform ED's	
lvermectin	66	Lantus SoloStar		Local preparations for Anal a	
- <b>J</b> -		Lanvis		Rectal Disorders	
Jadelle		Lanzol Relief		Locasol	
Jakavi		Lapatinib ditosylate		Locoid	
Jardiamet		Largactil		Locoid Crelo	
Jardiance	12	Laronidase	28	Locoid Lipocream	64

Locorten-Vioform	229	vaccine	Methylprednisolone aceponate	6
Lodoxamide	231	Mebendazole89	Methylprednisolone acetate	7
Logem	127	Mebeverine hydrochloride8	Methylxanthines	22
Lomide	231	Medco	Metoclopramide Actavis 10	130
Lomustine	146	Dermatological65	Metoclopramide hydrochloride	130
Loniten	57	Nervous121	Metolazone	5
Loperamide hydrochloride	6	Medrol79	Metopirone	8
Lopinavir with ritonavir		Medroxyprogesterone acetate	Metoprolol IV Mylan	5
Loprofin		Genito-Urinary75	Metoprolol succinate	
Loprofin Mix		Hormone81–82	Metoprolol tartrate	
Lorafix	221	Mefenamic acid110	Metrogyl	9
Loratadine		Megestrol acetate164	Metronidazole	
Lorazepam	134	Melatonin135	Metyrapone	8
Lorfast		Melphalan146	Mexiletine hydrochloride	5
Lorstat	<u>55</u>	Menactra268	Mexiletine Hydrochloride USP	
Losartan Actavis	48	Meningococcal (groups A, C, Y and	Miacalcic	
Losartan potassium	48	W-135) conjugate vaccine 268	Micolette	2
Losartan potassium with		Meningococcal C conjugate	Miconazole	3
hydrochlorothiazide	48	vaccine	Miconazole nitrate	
Lovir		Menthol63	Dermatological	6
Loxamine	125	Mepolizumab192	Genito-Urinary	
Lucrin Depot 1-month	87	Mercaptopurine148	Micreme	
Lucrin Depot 3-month		Mercilon 2873	Micreme H	6
Ludiomil		Mesalazine7	Microgynon 20 ED	74
Lyderm	68	Mesna152	Microgynon 30	
Lynparza		Mestinon110	Microgynon 50 ED	
Lyrica		Metabolic Disorder Agents26	Microlut	
- M -		Metformin hydrochloride12	Midazolam	
m-Eslon	122	Methadone hydrochloride	Midazolam-Baxter	
Mabthera	195	Extemporaneous238	Midodrine	5
Macrobid	109	Nervous122	Mifegyne	
Macrogol 3350 with potassium		Methatabs122	Mifepristone	
chloride, sodium bicarbonat		Methenamine (hexamine)	Minerals	
sodium chloride		hippurate109	Mini-Wright AFS Low Range	
Macrogol 400 and propylene		Methopt233	Mini-Wright Standard	
glycol	233	Methotrexate148	Minidiab	
Madopar 125	118	Methotrexate DBL Onco-Vial148	MiniMed 1.8 Reservoir	
Madopar 250		Methotrexate Ebewe148	MMT-326A	2
Madopar 62.5		Methotrexate Sandoz148	MiniMed 3.0 Reservoir	
Madopar HBS		Methyl hydroxybenzoate238	MMT-332A	2
Madopar Rapid		Methylcellulose238	MiniMed 640G	
Magnesium hydroxide	35	Methylcellulose with glycerin and	MiniMed Mio MMT-921A	
Magnesium sulphate		sodium saccharin238	MiniMed Mio MMT-923A	
Mantoux		Methylcellulose with glycerin and	MiniMed Mio MMT-925A	
Maprotiline hydrochloride		sucrose238	MiniMed Mio MMT-941A	
Marevan		Methyldopa53	MiniMed Mio MMT-943A	
Marine Blue Lotion SPF 50+		Methyldopa Mylan53	MiniMed Mio MMT-945A	2
Martindale Pharma	235	Methyldopa Mylan S2953	MiniMed Mio MMT-965A	2
Marvelon 28		Methylnaltrexone bromide25	MiniMed Mio MMT-975A	
Mask for spacer device		Methylphenidate ER - Teva138	MiniMed Quick-Set MMT-386A	
Mast Cell Stabilisers		Methylphenidate hydrochloride 138	MiniMed Quick-Set MMT-387A	
Maviret		Methylphenidate hydrochloride	MiniMed Quick-Set MMT-396A	
Maxidex		extended-release	MiniMed Quick-Set MMT-397A	2
Maxitrol		Methylprednisolone79	MiniMed Quick-Set MMT-398A	
MCT oil (Nutricia)		Methylprednisolone (as sodium	MiniMed Quick-Set MMT-399A	
Measles, mumps and rubella	_	succinate) 79	MiniMed Silhouette MMT-368A	
		•		

MiniMed Silhouette MMT-377A	22	Mupirocin	62	Nevirapine	10
MiniMed Silhouette MMT-378A	22	Muscle Relaxants		Nevirapine Alphapharm	
MiniMed Silhouette MMT-381A	22	Mvite	34	Nicorandil	
MiniMed Silhouette MMT-382A	22	Myambutol	100	Nicotine	14
MiniMed Silhouette MMT-383A		Mycobutin	100	Nifedipine	5
MiniMed Silhouette MMT-384A		MycoNail		Nifuran	
MiniMed Sure-T MMT-864A	20	Mycophenolate mofetil		Nilotinib	
MiniMed Sure-T MMT-866A	20	Mydriacyl		Nilstat	
MiniMed Sure-T MMT-874A	20	Mylan Atenolol		Alimentary	3
MiniMed Sure-T MMT-876A	20	Mylan Clomiphen		Genito-Urinary	
MiniMed Sure-T MMT-884A	20	Mylan Midazolam		Infection	
MiniMed Sure-T MMT-886A	20	Myleran		Nintedanib	
Minims Cyclopentolate	233	Myometrial and Vaginal Hormor		Nipent	
Minims Pilocarpine		Preparations		Nitrates	
Minims Prednisolone		Myozyme	26	Nitroderm TTS	
Minirin		Mysoline S29		Nitrofurantoin	10
Minirin Melt		- N -		Nitrolingual Pump Spray	
Mino-tabs	93	Nadolol	51	Nivestim	
Minocycline hydrochloride		Naglazyme		Nivolumab	
Minomycin		Nalcrom		Nizoral	
Minor Skin Infections		Naloxone hydrochloride		Nodia	
Minoxidil		Naltraccord		Noflam 250	
Mirena		Naltrexone hydrochloride		Noflam 500	
Mirtazapine		Naphazoline hydrochloride		Non-Steroidal Anti-Inflammatory	
Misoprostol		Naphcon Forte		Drugs	11
Mitomycin C		Naprosyn SR 1000		Nonacog gamma, [Recombinant	
Mitozantrone		Naprosyn SR 750		Factor IX]	4
Mitozantrone Ebewe		Naproxen		Norethisterone	
Mixtard 30		Narcaricin mite		Genito-Urinary	7
MMR II		Nasal Preparations		Hormone	
Moclobemide		Natalizumab		Norflex	
Modafinil		Natulan		Norfloxacin	
Modavigil		Nausafix		Noriday 28	
Moduretic		Nausicalm		Norimin	
Molaxole		Navelbine		Normacol Plus	
Moments		Necon		Normison	
Mometasone furoate		Nedocromil		Norpress	
Monogen		Nefopam hydrochloride		Nortriptyline hydrochloride	
Montelukast		Neisvac-C		Norvir	
Montelukast Mylan		Neo-B12		Noumed Paracetamol	
Moroctocog alfa [Recombinant fa		Neo-Mercazole		NovaSource Renal	
VIII]		Neo-Mercazole S29		Novatretin	
Morphine hydrochloride		Neocate Gold		NovoMix 30 FlexPen	
Morphine sulphate		Neocate Junior Unflavoured		NovoRapid	
Motetis		Neocate Junior Vanilla		NovoRapid FlexPen	
Mouth and Throat		Neocate SYNEO		NovoRapid Penfill	
Movapo		Neoral		NovoSeven RT	
Moxifloxacin		Neostigmine metilsulfate		Noxafil	
MSUD Maxamum		Nepafenac		Nozinan	
Mucilaginous laxatives with	204	Nepro HP (strawberry)		Nozinan (Swiss)	
stimulants	25	Nepro HP (vanilla)		Nucala	
Mucolytics		Nepro HP RTH		Nuelin	
Mucosoothe		Nerisone		Nuelin-SR	
Multiple Sclerosis Treatments		Neulactil		Nutilis	
Multivitamin renal		Neulastim		Nutren Diabetes	
Multivitamins		NeuroTabs		Nutrient Modules	
141GHU VIIGITIII IS		1400101000		TAGGICTIC INICOGUICS	44

Nutrini Energy Multi Fibre245	Omeprazole actavis 40	9	Paediatric Seravit	34
Nutrini Energy RTH245	Omnitrope	83	Palbociclib	160
Nutrini Low Energy Multi Fibre247	Onbrez Breezhaler	222	Paliperidone	133
Nutrini Peptisorb258	Oncaspar LYO	153	Pamidronate disodium	112
Nutrini Peptisorb Energy258	OncoTICE	173	Pamisol	112
Nutrini RTH245	Ondansetron	130	Panadol Mini Caps	
Nutrison 800 Complete Multi	Ondansetron ODT-DRLA	130	Pancreatic enzyme	24
Fibre 250	One-Alpha	33	Pantoprazole	9
Nutrison Concentrated252	Onrex		Panzop Relief	§
Nutrison Energy249	Opdivo	214	Panzytrat	24
Nutrison Energy Multi Fibre250	Ora-Blend	238	Papaverine hydrochloride	57
Nutrison Multi Fibre250	Ora-Blend SF	238	Para-amino salicylic acid	100
Nutrison Standard RTH250	Ora-Plus	238	Paracare	121
Nyefax Retard52	Ora-Sweet	238	Paracare Double Strength	
Nystatin	Ora-Sweet SF	238	Paracetamol	121
Alimentary32	Orabase	32	Paracetamol + Codeine	
Genito-Urinary75	Oral and Enteral Feeds	243	(Relieve)	123
Infection97	Oratane	61	Paracetamol Pharmacare	121
NZB Low Gluten Bread Mix253	Ordine	122	Paracetamol with codeine	123
-0-	Orgran	254	Paraffin	65-66
O/W Fatty Emulsion Cream65	Ornidazole	99	Paraffin liquid with wool fat	233
Obinutuzumab193	Orphenadrine citrate	117	Paraldehyde	126
Obstetric Preparations77	Ortho-tolidine	77	Parasidose	68
Ocicure9	Oruvail SR	110	Parasiticidal Preparations	66
Ocrelizumab 135	Osmolite RTH	250	Parnate	124
Ocrevus135	Other Endocrine Agents	88	Parnate S29	124
Octocog alfa [Recombinant factor	Other Oestrogen Preparations	82	Paromomycin	95
VIII] (Advate)41	Other Progestogen		Paroxetine	
Octocog alfa [Recombinant factor	Preparations	82	Paser	100
VIII] (Kogenate FS)41	Other Skin Preparations	70	Paxam	134
Octreotide164	Ovestin		Paxtine	
Octreotide (Sun) 164	Genito-Urinary	75	Pazopanib	160
Octreotide GH164	Hormone	82	Peak flow meter	228
Octreotide LAR (somatostatin	Ox-Pam	134	Pedialyte - Bubblegum	46
analogue) 164	Oxaliplatin	146	Pediasure	245
Octreotide MaxRx164	Oxaliplatin Accord	146	Pediasure RTH	245
Oestradiol81	Oxaliplatin Actavis 100	146	Pegaspargase	153
Oestradiol valerate81	Oxaliplatin Ebewe	146	Pegasys	107
Oestradiol with norethisterone82	Oxazepam	134	Pegfilgrastim	44
Oestriol	Oxis Turbuhaler	222	Pegylated interferon alfa-2a	107
Genito-Urinary75	Oxpentifylline	57	Pembrolizumab	
Hormone82	Oxybutynin	76	Pemetrexed	148
Oestrogens81	Oxycodone hydrochloride		Penicillamine	
Ofev224	Oxycodone Sandoz	123	Penicillin G	
Oil in water emulsion65	Oxycodone Sandoz S29		PenMix 30	
Olanzapine131–132	OxyNorm	123	PenMix 40	11
Olaparib152	Oxytocin		PenMix 50	11
Olbetam55	Oxytocin BNM	75	Pentasa	7
Olbetam S2955	Oxytocin with ergometrine		Pentostatin [Deoxycoformycin]	153
Olopatadine233	maleate	<del>76</del>	Pentoxifylline [Oxpentifylline]	57
Olopatadine Teva233	Ozurdex	230	Peptamen Junior	24
Olsalazine7	- P -		Peptisoothe	9
Omalizumab193	Pacifen		Peptisorb	246
Omeprazole9	Paclitaxel		Perhexiline maleate	52
Omeprazole actavis 109	Paclitaxel Actavis	153	Pericyazine	132
Omeprazole actavis 209	Paclitaxel Ebewe	153	Perindopril	48

Perjeta	194	polysaccharide vaccine	271	Promethazine hydrochloride	22
Permethrin	68	Pneumovax 23	271	Propafenone hydrochloride	5
Perrigo		Podophyllotoxin		Propamidine isethionate	22
Pertuzumab		Polaramine	221	Propranolol	5
Peteha	100	Poliomyelitis vaccine		Propylene glycol	
Pethidine hydrochloride	123	Poloxamer		Propylthiouracil	
Pevaryl	62	Poly-Gel		Protaphane	
Pexsig		Poly-Tears	233	Protaphane Penfill	
Pfizer Exemestane		Poly-Visc		Protifar	
Pharmacy Health Sorbolene with		Polycal		Protionamide	
Glycerin		Ponstan		Provera	
Pharmacy Services		Posaconazole		Provera HD	
Pheburane		Postinor-1		PSM Citalopram	
Phenasen		Potassium chloride		Psoriasis and Eczema	
Phenobarbitone		Potassium Chloride Aguettant		Preparations	6
Phenobarbitone sodium		Potassium citrate		PTU	8
Extemporaneous	238	Potassium iodate		Pulmicort Turbuhaler	
Nervous		Povidone iodine		Pulmozyme	
Phenothrin		Pradaxa		Puri-nethol	
Phenoxybenzamine		Pramipexole hydrochloride		Puria	
hydrochloride	47	Pravastatin		Pyrazinamide	
Phenoxymethylpenicillin (Penicill		Pravastatin Mylan		Pyridostigmine bromide	
V)		Praziquantel		Pyridoxine hydrochloride	را ا
		Prazosin			
Phenytoin sodium1 Phillips Milk of Magnesia1		Pred Forte		Pyrimethamine	
Phlexy 10	255	Prednisolone		Pytazen SR	4
		Prednisolone acetate		Q 300	0
Phospharus				Quetapel	
Phosphorus		Prednisolone sodium	0		
Phytomenadione		Prednisolone sodium	004	Quetiapine	
Pilocarpine hydrochloride		phosphate		Quick-Set MMT-392	
Pimafucort		Prednisolone-AFT		Quick-Set MMT-393	
Pimecrolimus		Prednisone		Quinapril	4
Pindolol		Pregabalin		Quinapril with	
Pine tar with trolamine laurilsulfa		Pregabalin Pfizer		hydrochlorothiazide	4
and fluorescein		Pregnancy Tests - hCG Urine		Quinine sulphate	
Pinetarsol		Premarin		Qvar	22
Pioglitazone		Prevenar 13		-R-	
Pirfenidone		Priadel		RA-Morph	12
Pizotifen		Primaquine		Raloxifene hydrochloride	
PKU Anamix Infant		Primidone		Raltegravir potassium	
PKU Anamix Junior		Primolut N		Ramipex	
PKU Anamix Junior Chocolate		Priorix		Ranbaxy-Cefaclor	
PKU Anamix Junior LQ		Probenecid		Ranitidine	
PKU Anamix Junior Orange		Probenecid-AFT		Rapamune	21
PKU Anamix Junior Vanilla		Procaine penicillin		Reandron 1000	
PKU Lophlex LQ 10		Procarbazine hydrochloride	153	Recombinant factor IX	
PKU Lophlex LQ 20	255	Prochlorperazine	130	Recombinant factor VIIa	4
PKU Lophlex Powder	255	Proctofoam	<mark>7</mark>	Recombinant factor VIII	40–4
PKU Lophlex Sensation 20		Proctosedyl		Rectogesic	
Plaquenil		Procyclidine hydrochloride		Redipred	<mark>7</mark>
Plendil ER		Procytox		Relieve	11
Pneumococcal (PCV10) conjuga		Progesterone	82	Relistor	2
vaccine		Proglicem		Remicade	18
Pneumococcal (PCV13) conjuga		Proglycem		Renilon 7.5	
vaccine		Progynova		Resonium-A	
Pneumococcal (PPV23)		Prolia		Resource Beneprotein	24
, ,				•	

Respigen		Sabril		Sodium bicarbonate	
Respiratory Devices		Sacubitril with valsartan		Blood	45–46
Respiratory Stimulants		SalAir	223	Extemporaneous	238
Retinol palmitate	234	Salazopyrin	8	Sodium calcium edetate	236
ReTrieve	61	Salazopyrin EN	8	Sodium chloride	
Retrovir	106	Salbutamol	223	Blood	45
Revlimid	151	Salbutamol with ipratropium		Respiratory	
Revolade	38	bromide	223	Sodium citrate with sodium laury	/l
Rexacrom	231	Salicylic acid	69	sulphoacetate	26
Riboflavin		Salmeterol	222	Sodium citro-tartrate	
Ribomustin	144	Sandomigran		Sodium cromoglicate	
Ricit	<mark>76</mark>	Sandostatin LAR		Alimentary	8
Rifabutin		Sanofi Primaquine		Respiratory	
Rifadin	100	Sapropterin dihydrochloride		Sensory	
Rifampicin	100	Scalp Preparations		Sodium fluoride	
Rifaximin		Scopoderm TTS		Sodium Fusidate [fusidic acid]	
Rifinah		Sebizole		Dermatological	62
Rilutek		Secukinumab		Infection	
Riluzole		Sedatives and Hypnotics		Sensory	
Riodine		Seebri Breezhaler		Sodium hyaluronate [Hyaluronic	
Risedronate Sandoz		Selegiline hydrochloride		acid]	
Risedronate sodium		Senna		Sodium phenylbutyrate	
Risperdal Consta		Senokot		Sodium polystyrene sulphonate	
Risperidone		Sensipar		Sodium tetradecyl sulphate	
Risperidone (Teva)		SensoCard		Sodium valproate	
Risperon		Serenace		Sofradex	
Ritalin		Seretide		Soframycin	
Ritalin LA		Seretide Accuhaler		Solgar	
Ritonavir		Serevent		Solifenacin Mylan	
Rituximab (Mabthera)		Serevent Accuhaler		Solifenacin succinate	
Rituximab (Riximyo)		Sertraline		Solu-Cortef	
Rivaroxaban		Setrona		Solu-Medrol	
Rivastigmine		Setrona AU		Solu-Medrol-Act-O-Vial	
Rivotril		Sevredol		Somatropin (Omnitrope)	
Riximyo		Sex Hormones Non	122	Sotalol	
			90		
RIXUBIS		Contraceptive		Spacer device	
Rizamelt		Shield XL		Span-K	
Rizatriptan		shingles vaccine		Spiolto Respimat	
Ropin		SII-Onco-BCG		Spiractin	
Ropinirole hydrochloride		Sildenafil		Spiriva Daggimet	
Rotarix		Silhouette MMT-373		Spiriva Respimat	
Rotavirus oral vaccine		Siltuximab		Spironolactone	
Roxane		Simvastatin		Sporanox	
Roxane-Propranolol		Simvastatin Mylan		Sprycel	
Roxithromycin		Sinemet		Staphlex	
Rubifen		Sinemet CR		Stemetil	
Rubifen SR	138	Sirolimus	217	SteroClear	
Rugby Capsaicin Topical	400	Siterone	80	Stesolid	
Cream		Slow-Lopresor		Stimulants/ADHD Treatments	
Rulide D		Smith BioMed Rapid Pregnancy		Stiripentol	
Rurioctocog alfa pegol [Reco		Test		Stocrin	
factor VIII]		Sodibic		Stomahesive	
Ruxolitinib		Sodium acid phosphate	26	Strattera	137
Rythmodan		Sodium alginate		Strides Shasun	
Rytmonorm	50	Sodium benzoate	30	Stromectol	66
- S -					

Sucralfate	10	Temozolomide	154	Total parenteral nutrition (TPN)	4
Sulfadiazine Silver	62	Tenofovir disoproxil	101	TPN	
Sulfadiazine sodium	95	Tenofovir Disoproxil Teva	101	Tramadol hydrochloride	12
Sulfasalazine	8	Tenoxicam	110	Tramal SR 100	12
Sulindac	110	Tensipine MR10	52	Tramal SR 150	
Sulindac Mylan	110	Tepadina	146	Tramal SR 200	12
Sulphur	69	Terazosin	47	Trandate	5
Sulprix		Terbinafine	98	Tranexamic acid	4
Sumatriptan	129	Terbutaline sulphate	223	Tranylcypromine sulphate	12
Sunitinib		Teriflunomide		Trastuzumab	21
Sunscreens	70	Teriparatide	113	Trastuzumab emtansine	
Sunscreens, proprietary	70	Testosterone	80	Travatan	23
Sure-T MMT-863	20	Testosterone cipionate	80	Travoprost	
Sure-T MMT-873	20	Testosterone esters		Travopt	
Sustagen Diabetic	243	Testosterone undecanoate	80	Treatments for Dementia	
Sustagen Hospital Formula		Tetrabenazine	119	Treatments for Substance	
Active	250	Tetrabromophenol	77	Dependence	14
Sustanon Ampoules	80	Tetracosactrin	80	Trental 400	
Sutent		Tetracycline		Tretinoin	
Sylvant		Thalidomide		Dermatological	6
Symbicort Turbuhaler 100/6		Thalomid		Oncology	
Symbicort Turbuhaler 200/6		Theophylline	226	Trexate	
Symbicort Turbuhaler 400/12		Thiamine hydrochloride		Triamcinolone acetonide	
Symmetrel		THIO-TEPÁ		Alimentary	3
Sympathomimetics		Thioguanine		Dermatological	6
Synacthen		Thiotepa		Hormone	
Synacthen Depot		Thymol glycerin		Triamcinolone acetonide with	
Synacthene Retard		Thyroid and Antithyroid Agent		gramicidin, neomycin and nysta	atin
Synflorix		Ticagrelor		Dermatological	
Synthroid	83	Tilade		Sensory	
Syntometrine		Tilcotil	110	Triaver	
Syrup (pharmaceutical grade)		Tillomed	146	Triazolam	
Systane Unit Dose		Timolol		Trimethoprim	
-T-		Cardiovascular	52	Trimethoprim with	
Tacrolimus	219	Sensory		sulphamethoxazole	
Tacrolimus Sandoz		Timoptol XE		[Co-trimoxazole]	9
Taliglucerase alfa		Tiotropium bromide		Trisequens	
Tambocor		Tiotropium bromide with		Trisul	
Tamoxifen citrate		olodaterol	224	Trophic Hormones	
Tamoxifen Sandoz		Tivicay		Tropicamide	
Tamsulosin hydrochloride		TMP		Trusopt	23
Tamsulosin-Rex		Tobramycin		TruSteel	
Tandem Cartridge		Infection	96	Tuberculin PPD [Mantoux] test	
Tandem t:slim X2 with Basal-IQ		Sensory		Tubersol	
Tap water		Tobramycin BNM		Two Cal HN	
Tarceva		Tobramycin Mylan		Two Cal HN RTH	
Tasigna		Tobrex		Tykerb	
Tasmar		Tocilizumab		Tysabri	
Taurine		Tofranil		- U -	
Tecfidera		Tolcapone		Ultibro Breezhaler	22
Tegretol		Topamax		Ultraproct	
Tegretol CR		Topical Products for Joint and		Umeclidinium	22
Telfast		Muscular Pain		Umeclidinium with vilanterol	
Teligent		Topiderm		Univent22	
Temaccord		Topiramate		Ural	
Temazepam		Topiramate Actavis		Urea	
10111azopa111	100	i opiiamate Actavio	120	O104	

Urex Forte		Viramune Suspension	105	Zoledronic acid	
Urinary Agents		ViruPOS		Hormone	
Urinary Tract Infections	109	Vit.D3	33	Musculoskeletal	
Urinorm	116	Vita-B12		Zoledronic acid Mylan	
Uromitexan		VitA-POS	234	Zopiclone	
Ursodeoxycholic acid	24	Vitabdeck	34	Zopiclone Actavis	136
Ursosan	24	Vital	246	Zostavax	272
Utrogestan	82	Vitamin B complex		Zostrix	111
- V -		Vitamin B6 25	33	Zostrix HP	
Vaccinations	261	Vitamins	33–34	Zuclopenthixol decanoate	134
Vaclovir	101	Vivonex TEN	246	Zuclopenthixol hydrochloride	132
Valaciclovir	101	Voltaren	110	Zusdone	132
Valganciclovir		Voltaren D		Zyban	141
Valganciclovir Mylan	101	Voltaren Ophtha	230	Zypine	131
Vancomycin	96	Volumatic	228	Zypine ODT	131
Vannair	222	Voriconazole	98	Zyprexa Relprevv	132
Varenicline Pfizer	142	Vosol	229	Zytiga	163
Varenicline tartrate	142	Votrient	160		
Varicella vaccine [Chickenpox		Vttack	98		
vaccine]	272	- W -			
Varicella zoster virus (Oka stra	in) live	Warfarin sodium	44		
attenuated vaccine [shingles		Wart Preparations	70		
vaccine]	272	Wasp venom allergy treatme	nt220		
Various	235	Water			
Varivax	272	Blood	45		
Vasodilators	<mark>56</mark>	Extemporaneous	239		
Vasopressin Agonists	87	Wool fat with mineral oil	66		
Vasorex	52	- X -			
Vedafil	58	Xarelto	44		
Veletri	<u>59</u>	Xifaxan	10		
Venclexta	155	XMET Maxamum	254		
Venetoclax	155	Xolair	193		
Venlafaxine	125	XP Maxamum	255		
Venomil	220	Xylocaine	120		
VENOX	220	Xylocaine 2% Jelly	119		
Ventavis	59	Xyntha			
Ventolin	223	- Z -			
Vepesid	151	Zapril	47		
Verapamil hydrochloride	53	Zarontin	126		
Vergo 16	130	Zaroxolyn	54		
Vermox		Zavedos	151		
Versacloz	131	Zeffix	101		
Vesanoid	155	Zetlam	101		
Vexazone	12	Ziagen	105		
Vfend	98	Zidovudine [AZT]	106		
Viaderm KC	65	Zidovudine [AZT] with			
Vidaza	146	lamivudine	106		
Vigabatrin	128	Zimybe	56		
Vildagliptin		Zinc and castor oil			
Vildagliptin with metformin		Zinc sulphate	36		
hydrochloride	12	Zincaps	36		
Vimpat		Zinnat			
Vinblastine sulphate		Ziprasidone			
Vincristine sulphate		Zista			
Vinorelbine		Zithromax	90		
Vinorelbine Ebewe					

