August 2021 Volume 28 Number 2

Editors:

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at no cost from the Pharmac website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

		L
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	60
	Genito Urinary System	70
	Hormone Preparations – Systemic	77
	Infections – Agents For Systemic Use	88
	Musculoskeletal System	109
	Nervous System	117
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	220
	Sensory Organs	229
	Various	235
Section C	Extemporaneous Compounds (ECPs)	237
Section D	Special Foods	240
Section I	National Immunisation Schedule	260
	Index	272

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 n sachet	0.	30	✓ Ga	viscon Infant
GODIUM ALGINATE ★ Tab 500 mg with sodium bicarbonate 267 mg and calciur carbonate 160 mg - peppermint flavour		60		viscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and cal carbonate 160 mg per 10 ml		500 ml	Ac	idex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE		100	✓ Alı	u-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) Subsidy by endorsement Only when prescribed for patients unable to swallow inappropriate and the prescription is endorsed accor	39.00	500 ml ts or whe	✓ Ro re calcium	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available Tab 2 mg Cap 2 mg	10.75	400 400	✓ No ✓ <u>Dia</u>	dia amide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Reta pharmacy	166.50	90 alid for 6		tocort CIR r applications meeting

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

(N	Subsidy fanufacturer's Price)	Subsi	Fully dised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	1	Essential
				Prednisolone S29
SODIUM CROMOGLICATE				
Cap 100 mg	92.91	100	✓	Nalcrom
SULFASALAZINE				
* Tab 500 mg	14.00	100	1	Salazopyrin
Tab EC 500 mg	15.53	100	1	Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and	00 × OD	✓ Illituanus at		
cinchocaine hydrochloride 5 mg per g	30 g OP	✓ Ultraproct		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and	40	/ 1111		
cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct		
HYDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl		
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl		

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a			
PSO	65.45	10	Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

IVII	SOPROSTOL		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	✓ Cytotec

		AL IMENTA DV	TD	ACT AND) METABOLISM
		ALIMENTANT	חו	ACT ANL	VIVIETABOLISIVI
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Н	elicobacter Pylori Eradication				
CLA	ARITHROMYCIN Tab 500 mg — Subsidy by endorsement	radication and prescri		is endorsed	
H	2 Antagonists				
FAN	MOTIDINE - Only on a prescription				
*	Tab 20 mg	4.91	100	√ F	amotidine Hovid S29
*	Tab 40 mg	8.48	100	√ F	amotidine Hovid S29
*	Inj 10 mg per ml, 4 ml — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients receive		10 of p		lylan S29
RAI	NITIDINE - Subsidy by endorsement	-	·		

- a) Only on a prescription
- b) Subsidy by endorsement Subsidised for patients who were taking ranitidine prior to 1 November 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ranitidine.
- 300 ml ✓ Peptisoothe (Peptisoothe Oral lig 150 mg per 10 ml to be delisted 1 September 2021)

(,		
Proton Pump Inhibitors		
LANSOPRAZOLE		
* Cap 15 mg4.20	100	✓ Lanzol Relief
* Cap 30 mg5.26	100	✓ Lanzol Relief
OMEPRAZOLE		
For omeprazole suspension refer Standard Formulae, page 237		
* Cap 10 mg1.94	90	✓ Omeprazole actavis 10
* Cap 20 mg	90	✓ <u>Omeprazole actavis</u> 20
* Cap 40 mg3.11	90	✓ Omeprazole actavis 40
* Powder – Only in combination42.50	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspension.	-	
* Inj 40 mg ampoule with diluent	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
		✓ Ocicure S29
(Ocicure 1992) Inj 40 mg ampoule with diluent to be delisted 1 October 2021) PANTOPRAZOLE		
* Tab EC 20 mg2.02	100	✓ Panzop Relief
* Tab EC 40 mg2.85	100	✓ Panzop Relief

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Brand or Generic Manufacturer
Site Protective Agents				
COLLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	✓ (Gastrodenol S29
Tab 1 g	35.50 (48.28)	120	(Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail pha Tab 550 mg SA1461 Special Authority for Subsidy	,	56	✓ <u>)</u>	<u> Kifaxan</u>
Initial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient hat lolerated doses of lactulose.				
Renewal only from a gastroenterologist, hepatologist or Practiti hepatologist. Approvals valid without further renewal unless no benefiting from treatment.			•	•
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Retail ph	armacy			
Cap 25 mg Cap 100 mg		100 100		Proglicem S29 Proglicem S29
Oral liq 50 mg per ml		100 80 ml O		Proglycem S29
⇒SA1320 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals va	alid for 12 months whe	re use	d for the tre	eatment of confirmed
hypoglycaemia caused by hyperinsulinism.				
Renewal from any relevant practitioner. Approvals valid withou appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE	ut turtner renewai unie	SS NOU	nea where	the treatment remains
Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	√ <u>(</u>	Glucagen Hypokit
Insulin - Short-acting Preparations				
NSULIN NEUTRAL				
▲ Inj human 100 u per ml	25.26 1	0 ml O		Actrapid Humulin R
▲ Inj human 100 u per ml, 3 ml	42.66	5	1	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE ▲ Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	V 1	NovoMix 30 FlexPen

	Subsidy (Manufacturer's F \$	Price) Subs	Fully Brand or idised Generic Manufacturer
NSULIN ISOPHANE			
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH✓ Protaphane
Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH✓ Protaphane Penfill
NSULIN ISOPHANE WITH INSULIN NEUTRAL ▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Mixtard 30 ✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 40 ✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE Inj lispro 25% with insulin lispro protamine 75% 100 u per m 3 ml	42.66	5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per m 3 ml		5	✓ Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml Inj 100 u per ml, 3 ml disposable pen	94.50	1 5 5	✓ Lantus✓ Lantus✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART Inj 100 u per ml, 10 ml Inj 100 u per ml, 3 ml Inj 100 u per ml, 3 ml syringe	51.19	1 5 5	✓ NovoRapid✓ NovoRapid Penfill✓ NovoRapid FlexPen
▲ Inj 100 u per ml, 10 ml		1	✓ Apidra✓ Apidra
▲ Inj 100 u per ml, 3 ml		5 5	✓ Apidra SoloStar
N30EIN EISFNO ▲ Inj 100 u per ml, 10 ml ▲ Inj 100 u per ml, 3 ml		10 ml OP 5	✓ Humalog✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE * Tab 50 mg		90	✓ Glucobay✓ Accarb
★ Tab 100 mg	8.95 6.40 15.29	90	✓ Glucobay ✓ Accarb
Glucobay Tab 50 mg to be delisted 1 December 2021) Glucobay Tab 100 mg to be delisted 1 December 2021)	70.20		
Oral Hypoglycaemic Agents			
SLIBENCLAMIDE * Tab 5 mg	7.50	100	✓ Daonil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
GLICLAZIDE				
* Tab 80 mg	15.18	500	•	<u>Glizide</u>
GLIPIZIDE				
* Tab 5 mg	3.27	100	•	Minidiab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg		1,000	/	Apotex
* Tab immediate-release 850 mg	7.04	500	•	Apotex
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓	Vexazone
* Tab 30 mg	7.30	90	✓	Vexazone
* Tab 45 mg	12.25	90	✓	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	1	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	1	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	1	Galvumet

SGLT2 Inhibitors

⇒SA2029 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has type 2 diabetes; and
- 2 Any of the following:
 - 2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*: or
 - 2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.5 Patient has diabetic kidney disease (see note b)*; and
- 3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months; and
- 4 Treatment will not be used in combination with a funded GLP-1 agonist.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

ΕN	IPAGLIFLOZIN - Special Authority see SA2029 above - Retail pharmacy		
*	Tab 10 mg	30	Jardiance
*	Tab 25 mg	30	 Jardiance

Fully

Brand or

	(Manufacturer's Price)	Sul Per	bsidised	Generic Manufacturer	
EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Sp	pecial Authority see S		n the pre		ail
* Tab 5 mg with 1,000 mg metformin hydrochloride * Tab 5 mg with 500 mg metformin hydrochloride * Tab 12.5 mg with 1,000 mg metformin hydrochloride * Tab 12.5 mg with 500 mg metformin hydrochloride	58.56 58.56	60 60 60	✓ J.	ardiamet ardiamet ardiamet ardiamet	

Subsidy

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly: or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	1	Manufacturer	

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES	 Maximum of 200 	dev per prescription
---------------------	------------------------------------	----------------------

	00 407	40.50	400	/ D D 14:
*	29 g × 12.7 mm	10.50	100	B-D Micro-Fine
*	31 g × 5 mm	11.75	100	 B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm	10.50	100	✓ B-D Micro-Fine
*			100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	- Maximum of 200) dev per pre	escription
	Syringe 0.3 ml with 29 g × 12.7 mm needle		100	✓ B-D Ultra Fine
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
	3, 3, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1, 1,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g × 12.7 mm needle	` '	100	✓ B-D Ultra Fine
•	-,···g g · · · · · · · · · · ·	1.30	10	
		(1.99)	. •	B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
•	Symige de im mar er give im necese imministration	1.30	10	
		(1.99)	. •	B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	` '	100	✓ B-D Ultra Fine
•	5)g5 : =5 g × .=	1.30	10	2 2 0 0
		(1.99)	. •	B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	` '	100	✓ B-D Ultra Fine II
-1-	Syrings This High of g A o Hill Hoodio	1.30	100	- D D OMAT HICH
			10	D D I III Time II
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year p	eriod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Suk	sidised	Generic	
(Manuacturer 3 i lice)	Out	Joiuiseu	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	/	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 2 Eithor:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

	Subsidy (Manufacturer's Price) \$	S Per	Fully subsidised	Brand or Generic Manufacturer
continued pump therapy; and 4 The patient is continuing to derive benefit from pump the 5 The patient had achieved and is maintaining a HbA1c o 6 The patient has had no increase in severe unexplained 7 The patient's HbA1c has not deteriorated more than 5 n 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within the second series of the second	f equal to or less than a hypoglycaemic episod nmol/mol from baseline	es from		
Renewal — (Previous use before 1 September 2012) only fi years for applications meeting the following criteria: All of the following:		st or nu	rse practiti	oner. Approvals valid for 2
 The patient is continuing to derive benefit according to than 80 mmol/mol; and The patient's HbA1c has not deteriorated more than 5 not The patient has not had an increase in severe unexplair Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within the 	nmol/mol from initial ap ned hypoglycaemic epi	plication	n; and	·
NSULIN PUMP CARTRIDGE – Special Authority see SA1985 a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded p		harmac	у	
Cartridge 300 U, t:lock × 10		1 OP		andem Cartridge
NSULIN PUMP INFUSION SET (STEEL CANNULA) – Specia a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	al Authority see SA198	5 on pa	ge 18 – Re	etail pharmacy
10 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T

6 mm steel needle; 29 G; manual insertion; 60 cm tubing x

8 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

1 OP

1 OP

MMT-876A

✓ Sure-T MMT-863

✓ Sure-T MMT-873

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 18 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

,	nm steel cannula; straight insertion; 80 cm line × 10 with			
01	10 needles	130.00	1 OP	✓ TruSteel
8 n	nm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
6 n	nm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
6 n	nm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 n	nm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 n	nm steel cannula; straight insertion; 81 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel

(TruSteel 6 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021) (TruSteel 8 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 18 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	✓ MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓ MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Quick-Set MMT-386A

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH IN	SERT	ION DEVIC	CE) - Special Authority see
SA1985 on page 18 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c	m			
line x 10 with 10 needles		1 OP	•	AutoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm				
line x 10 with 10 needles	140.00	1 OP	•	AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION) - Speci	al Au	thority see	SA1985 on page 18 –
Retail pharmacy				
a) Maximum of 3 sets per prescription				
b) Only on a prescriptionc) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with				
10 needles; luer lock	130.00	1 OP		Silhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH		-		
see SA1985 on page 18 – Retail pharmacy	I INSENTION WITE	I IIVO		= VICE) - Special Authority
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles		1 OP	•	AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cr	m		_	
line x 10 with 10 needles	140.00	1 OP	•	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device;	110.00	4 00		A1-0-4-00
110 cm line × 10 with 10 needles		1 OP	•	AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr		1 OP		AutoSoft 90
			-	
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH	I INSERTION) - S	pecial	Authority s	see SA1985 on page 18 –
Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing × 10 with	h			
10 needles; luer lock	130.00	1 OP		Quick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with				
10 needles; luer lock	130.00	1 OP	•	Quick-Set MMT-392
INSULIN PUMP RESERVOIR - Special Authority see SA1985 or	n page 18 – Retail pl	narma	асу	
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 packs of reservoir sets will be funded per		4 00		ADD Contribution 4.0
10 × luer lock conversion cartridges 1.8 ml for Paradigm pump Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		ADR Cartridge 1.8 MiniMed
Carriage for 5 and 7 series pump, 1.6 mil x 10	50.00	I OF	•	1.8 Reservoir
				MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP	•	MiniMed
			'	3.0 Reservoir
				MMT-332A

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	•	Manufacturer	
				-

Digestives Including Enzymes

PANCREATIC FNZYME

PANCREATIC ENZYME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	Creon Micro
JRSODEOXYCHOLIC ACID – Special Authority see SA1739 below	/ – Retail pha	rmacy	
Cap 250 mg	32.95	100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

U

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Normacol Plus

Subsidy		ully	Brand or	_
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	•	Manufacturer	

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soin	500 g OP	Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		
* Dry6.02	500 g OP	
(17.32)	· ·	Normacol Plus
2.41	200 g OP	

(8.72)

Faecal Softeners

* Tab 50 mg		✓ <u>Coloxyl</u> ✓ Coloxyl
* Tab 120 mg3.13 DOCUSATE SODIUM WITH SENNOSIDES	3 100	Coloxyl
* Tab 50 mg with sennosides 8 mg3.10	200	✓ Laxsol
POLOXAMER - Only on a prescription		
Not funded for use in the ear. * Oral drops 10%	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCLISATE SODILIM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority s	see SA1691 below – Retail	pharmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
•	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- - 1 The patient is receiving palliative care; and
 - 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy	, ,	Fully Bran	
	(Manufacturer's Price	ce) Subs Per	sidised Gen	eric ufacturer
	\$	rei	V IVIAII	uiaciurei
Osmotic Laxatives				
GLYCEROL	0.05	00	4 DOM	
* Suppos 3.6 g - Only on a prescription	9.25	20	✓ PSM	
LACTULOSE - Only on a prescription				
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ Laevol	ac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM B	ICARBONATE AND	SODIUM C	HLORIDE	
Powder for oral soln 13.125 g with potassium chloride 46.6				
sodium bicarbonate 178.5 mg and sodium chloride 350		30	✓ Molaxo	le
SODIUM ACID PHOSPHATE - Only on a prescription	•			_
Enema 16% with sodium phosphate 8%	2 50	1	✓ Fleet P	hosphate
= 10.11a 10.75 11a1 0001a1.1 p.100p.1a10 0.75		•	Enen	•
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETAT	F _ Only on a pres	crintion		
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per m	, ,	Сприоп		
5 ml	*	50	✓ Micole	to
J III	29.90	30	• IVIICOIC	<u></u>
Stimulant Laxatives				
Olimaiant Editativoo				
BISACODYL – Only on a prescription				
* Tab 5 mg		200	✓ Lax-Ta	-
* Suppos 10 mg	3.69	10	✓ Lax-Su	ppositories
SENNA - Only on a prescription				
* Tab, standardised	2.17	100		
	(8.21)		Senoko	t
	0.43	20		
	(2.06)		Senoko	t
SODIUM PICOSULFATE - Special Authority see SA2053 below	w – Retail pharmac	/		
Oral soln 7.5 mg per ml	, ,	30 ml OP	✓ Dulcola	ax SP Drop
TO CARDES Changed Authority for Cubaidy				•

Subsidy

Fully

Brand or

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:

Subsidy (Manufacturer's Price)	9	Fully Subsidised	Brand or Generic
 \$	Per	Jubsidised ✓	Manufacturer

continued...

- 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
- 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
- 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
- 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authority see SA2042 below – Retail	pharmacy		
Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 ` \$	Per	✓	Manufacturer

continued...

2.3 A disorder of intracellular cobalamin metabolism; and

3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy	
Cap 120 mgCBS	

30 ✓ Solgar Cap 160 mg.......CBS ✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

patient is benefiting from treatment.

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1988 below - Retail pharmacy ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts: or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to

IDURSULFASE - Special Authority see SA1623 below - Retail pharmacy

✓ Elaprase

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts: or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene, and

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic	
--	--	--

continued...

- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – Special Authority see SA2040 below –	Retail pharmacy		
Tab 500 mg	CBS	30	Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE − Special Authority see SA1599 below − Retail pharmacy
Soln 100 mg per mlCBS 100 ml ✓ Amzoate \$29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 Pharmac PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific
 deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
 - Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

		Subsidy acturer's Price) \$	Sı Per	Fully obsidised	Brand or Generic Manufacturer
	tinued				
All of	of the following:				
1)	 Patient has demonstrated a symptomatic improvement or no dete initiated; and 	rioration in the	main :	symptom	for which therapy was
2)	Patient has demonstrated a clinically objective improvement or no liver and spleen size; and	deterioration	in haer	noglobin l	evels, platelet counts and
3)	 Radiological (MRI) signs of bone activity performed at two years of demonstrate no deterioration shown by the MRI, compared with Nor adjusted dose; and 				
4)	 Patient has not had severe infusion-related adverse reactions wh and/or adjustment of infusion rates; and 	ich were not p	reventa	ble by ap	propriate pre-medication
5)	 Patient has not developed another medical condition that might re ERT: and 	easonably be e	expecte	d to comp	promise a response to
6)	 Patient is compliant with regular treatment and taliglucerase alfa i every other week rounded to the nearest whole vial (200 units), i 				
7)	 Supporting clinical information including test reports, MRI whole be investigations are submitted to the Gaucher Panel for assessmen 	ody STIR, had			

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with			
Endorsement		500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient who ha prescription is endorsed accordingly.	s oral mucositis a	as a result of tr	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder	8.48	28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
•	(6.00)		Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
		- 9	
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			. J
Oral gel 20 mg per g	4 74	40 g OP	✓ Decozol
		40 g Ol	- 5600201
NYSTATIN			
Oral liq 100,000 u per ml	4 70	24 ml OP	✓ Nilstat

	Subsidy (Manufacturer's Price	e) (Fully Subsidised	Brand or Generic
Other Oral Agents	\$	rei		Manufacturer
For folinic mouthwash, pilocarpine oral liquid or saliva substitut	e formula refer Standa	ard Forn	nulae, <mark>pag</mark>	e 237
FHYMOL GLYCERIN * Compound, BPC	9.15	500 ml	✓	PSM
Vitamins				
Vitamin B				
HYDROXOCOBALAMIN	PSO1.89	3		Neo-B12 Vita-B12
	3.15	5	✓	Hydroxocobalamin Mercury Pharma
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription				·
Tab 25 mg - No patient co-payment payable Tab 50 mg		90 500		<u>Vitamin B6 25</u> Apo-Pyridoxine
FHIAMINE HYDROCHLORIDE − Only on a prescription * Tab 50 mg	7.09	100	✓	Max Health
/ITAMIN B COMPLEX ★ Tab, strong, BPC	7.15	500	✓	Bplex
Vitamin C				
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription				
* Tab 100 mg	9.90	500	•	<u>Cvite</u>
Vitamin D				
ALFACALCIDOL ★ Cap 0.25 mcg ★ Cap 1 mcg ★ Oral drops 2 mcg per ml	87.98	100 100 20 ml O	1	One-Alpha One-Alpha One-Alpha
CALCITRIOL * Cap 0.25 mcg * Cap 0.5 mcg		100 100		Calcitriol-AFT Calcitriol-AFT
COLECALCIFEROL Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescri Coral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml O		<u>Vit.D3</u> Puria
Multivitamin Preparations				
MULTIVITAMIN RENAL - Special Authority see SA1546 on th ★ Cap		oharmad 30		Clinicians Renal Vit

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$) S Per	Fully Subsidised	Brand or Generic Manufacturer
■SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Either:	d without further ren	ewal un	less notifie	d for applications meeting
 The patient has chronic kidney disease and is receiving e The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m² body surface area (BSA). 				
MULTIVITAMINS - Special Authority see SA1036 below - Reta * Powder		:00 g OF	⊃ √ P	aediatric Seravit
■ SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali inborn errors of metabolism.	d without further ren	ewal un	less notifie	d where the patient has
Renewal from any relevant practitioner. Approvals valid without approval for multivitamins. VITAMINS	further renewal unle	ss notifi	ed where p	patient has had a previous
* Tab (BPC cap strength)		1,000	✓ <u>N</u>	<u>lvite</u>
* Cap (fat soluble vitamins A, D, E, K) - Special Authority see SA1720 below - Retail pharmacy		60	✓ V	'itabdeck
Initial application from any relevant practitioner. Approvals valithe following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut 3 Patient has severe malabsorption syndrome.		ewai uii	iess notine	u tor applications meeting
Minerals				
Calcium				
CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme Subsidy by endorsement – Only when prescribed for pa considered unsuitable.	nt54.60	250 76 years)	√ 0	Calci-Tab 500 Cacit \$29 ium carbonate oral liquid is
CALCIUM GLUCONATE				
* Inj 10%, 10 ml ampoule		10		lax Health - Hameln (\$29)
	64.00	20	✓ N	Max Health \$29
Fluoride				
SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)	5.75	100	✓ P	PSM

Tab 253 mcg (150 mcg elemental iodine)4.58

✓ NeuroTabs

90

lodine

POTASSIUM IODATE

Fully

Brand or

	(Manufacturer's Price) \$	Sı Per	ubsidised	Generic Manufacturer
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.09	100	✓ Fe	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	✓ Fe	erro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority se Inj 50 mg per ml, 10 ml vial	ee SA1840 below – F		armacy	erinject
SA1940 Special Authority for Subsidy				

Subsidy

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
 - 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON	POL	_YMAL	_TOSE
INUN	PUL	_ Y IVI <i>I</i> AL	-100

Magnesium

MAGNESIUM HYDROXIDE

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓	<u>Martindale</u>
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	•	Zincaps

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу			_
Wastage claimable					
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	/	Binocrit	
Inj 2,000 iu in 1 ml, syringe	100.00	6	✓	Binocrit	
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit	
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	1	Binocrit	
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	1	Binocrit	
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	1	Binocrit	
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	1	Binocrit	
Inj 10,000 iu in 1 ml, syringe		6	✓	Binocrit	
Inj 40,000 iu in 1 ml, syringe		1	✓]	Binocrit	

Megaloblastic

FOLIC ACID			
* Tab 0.8 mg	21.84	1,000	✓ Apo-Folic Acid
* Tab 5 mg		100	✓ Folic Acid Mylan
· ·	12.12	500	✓ Apo-Folic Acid
Oral lig 50 mcg per ml	26.00	25 ml OP	✓ Biomed
(Apo-Folic Acid Tab 5 mg to be delisted 1 December 2021)			

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

ricators aroup in conjunction with the reationary	idomoprilia ividnagomoni gro	up.	
Inj 250 iu vial	612.50	. 1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	-	1	Alprolix
TROMBOPAG - Special Authority see SA1743 be	elow – Retail pharmacy		
Wastage claimable			
Tab 25 mg	1,550.00	28	✓ Revolade
Tab 50 mg	3,100.00	28	✓ Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

EL.

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist.

continued...

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- Fither:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

FMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

✓ Hemlibra	1	3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	7,138.00	Inj 60 mg in 0.4 ml vial
✓ Hemlibra	1	12,492.00	Inj 105 mg in 0.7 ml vial
✓ Hemlibra	1	17.846.00	Ini 150 mg in 1 ml vial

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and

continued...

Subsidy (Manufacturer's		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more: and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe		1	✓ NovoSeven RT
Inj 5 mg syringe		1	✓ NovoSeven RT
Inj 8 mg syringe		1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U) 1	✓ FEIBA NF
Inj 1,000 U) 1	✓ FEIBA NF
Inj 2,500 U6,575.00) 1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

37.50 1	Xyntha
75.00 1	Xyntha
	✓ Xyntha
0.00 1	✓ Xyntha
60.00 1	✓ Xyntha
	5.00 1 0.00 1 0.00 1

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 1,000 iu vial. 870.00 1 ✓ RIXUBIS Inj 2,000 iu vial. 1,740.00 1 ✓ RIXUBIS Inj 3,000 iu vial. 2,610.00 1 ✓ RIXUBIS	Inj 500 iu vial	1	✓ RIXUBIS
, =,000 13 113	Inj 1,000 iu vial870.00	1	✓ RIXUBIS
lni 3 000 iu vial. 2 610 00 1 ✓ RIXUBIS	Inj 2,000 iu vial	1	✓ RIXUBIS
	Inj 3,000 iu vial2,610.00	1	✓ RIXUBIS

	0.1		- ·	
	Subsidy (Manufacturer's Price)	Su	Fully bsidised	
	\$	Per	√	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	[Xpharm]			
For patients with haemophilia. Preferred Brand of short half	-life recombinant factor	r VIII. A	ccess to	o funded treatment is
managed by the Haemophilia Treaters Group in conjunction				
Inj 250 iu vial		1		Advate
Inj 500 iu vial	420.00	1	✓	Advate
Inj 1,000 iu vial	840.00	1	✓	Advate
Inj 1,500 iu vial	1,260.00	1	✓	Advate
Inj 2,000 iu vial		1		Advate
Inj 3,000 iu vial	,	1	/	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the I	Vational	Haemo	philia Management Group,
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1	•	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatm		d treatm	ent is n	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			_	
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial	· ·	1		Adynovate
Inj 2,000 iu vial	2,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Konakion MM
ing to mg por mi, i mi op to o mg available on a room		•		Tronauton iiiii
Antithrombotic Agents				
- Tillian Omsons rigonic				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	•	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
				<u>Multichem</u>
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	1	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next pag				-
* Tab 90 mg		56	1	Brilinta

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA1646	below - Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe	101.30	10	✓ Clexane
Inj 120 mg in 0.8 ml syringe		10	Clexane Forte
Inj 150 mg in 1 ml syringe	143.86	10	Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODILIM

Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	203.68	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	19.00	5	✓ Hospira
	42.40		✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg		60	✓	Pradaxa
Cap 150 mg		60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	1	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	77.56	28	✓	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓	Coumadin
•	6.46	100	✓	Marevan
* Tab 2 mg	4.31	50	1	Coumadin
* Tab 3 mg	10.03	100	✓	Marevan
* Tab 5 mg	5.93	50	✓	Coumadin
	11.48	100	1	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail ph	narmacy			
Inj 300 mcg per 0.5 ml prefilled syringe		10		Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓	Nivestim

FILGRASTIM - Special Authority see SA1259 below - Retail ph	narmacy		
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓ Nivestim

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

✓ Neulastim

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	DECOD AND			TIMING OTIGANS
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	d Generic
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO		5 1		Biomed Biomed
POTASSIUM CHLORIDE * Inj 75 mg per ml, 10 ml	55.00	50		AstraZeneca Potassium Chloride Aquettant \$29
	65.00		•	Juno
(AstraZeneca Inj 75 mg per ml, 10 ml to be delisted 1 November	,	000	41	
(Potassium Chloride Aguettant S29 Inj 75 mg per ml, 10 ml to be	e aeiistea 1 Novemb	er 202	1)	
SODIUM BICARBONATE Inj 8.4%, 50 ml	19.95	1	/	Biomed
a) Up to 5 inj available on a PSOb) Not in combination		1		Diamad
Inj 8.4%, 100 mla) Up to 5 inj available on a PSO b) Not in combination	20.50	ı	·	Biomed
SODIUM CHLORIDE Not funded for use as a nasal drop. Not funded for nebuliser for nebuliser use.	use except when us	sed in o	conjunctio	on with an antibiotic intended
Inj 0.9%, bag – Up to 2000 ml available on a PSO		500 ml 1,000 m		Baxter Baxter
Only if prescribed on a prescription for renal dialysis, ma	ternity or post-natal	care in	the home	e of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs) Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standar		5	•	Biomed
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO	2.80	20	•	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50		Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	•	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN) Infusion	CBS	1 OP	/	TPN
WATER				
 On a prescription or Practitioner's Supply Order only when Schedule requiring a solvent or diluent; or On a bulk supply order; or When used in the extemporaneous compounding of eyed) When used for the dilution of sodium chloride soln 7% f 	e drops; or			n listed in the Pharmaceutical
Inj 10 ml ampoule – Up to 5 inj available on a PSO Inj 20 ml ampoule – Up to 5 inj available on a PSO		50 20	✓	Pfizer Fresenius Kabi Multichem
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85 3	00 g O	P 🗸	Calcium Resonium
▲ Th		- "		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F		Fully dised	Brand or Generic
	\$	Per	✓	Manufacturer
COMPOUND ELECTROLYTES				
Powder for oral soln - Up to 5 sach available on a PSO	9.77	50	✓ [<u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]				
Soln with electrolytes (2 × 500 ml)	6.55	1,000 ml OP	√	Pedialyte - Bubblegum
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓ I	Phosphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60		
	(11.85)		(Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90	200	1	Span-K
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	1	Sodibic
· -			1	Sodibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84.65	454 a OP	✓	Resonium-A

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Series Per Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DO	XAZOSIN	
*	Tab 2 ma	

*	Tab 2 mg17.35	500	✓ Apo-Doxazosin
*	Tab 4 mg	500	✓ Apo-Doxazosin

PHENOXYBENZAMINE HYDROCHLORIDE

*	Cap 10 mg	 65.00	30	✓ BNM S29
		216.67	100	✓ Dibenzyline S29

PRAZOSIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking prazosin prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of prazosin.

*	Tab 1 mg5.53	100	✓ Apo-Prazosin
	Tab 2 mg	100	✓ Apo-Prazosin
	Tab 5 mg11.70	100	✓ Apo-Prazosin

(Apo-Prazosin Tab 1 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 2 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 5 mg to be delisted 1 May 2022)

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

*	Oral liq 5 mg per ml94.99	95 ml OP	✓ Capoten
	135.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

(Captopril-Mylan S29) Oral lig 5 mg per ml to be delisted 1 January 2022)

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

*	Tab 0.5 mg	2.09	90	✓ Zapril
*	Tab 2.5 mg	4.80	90	✓ Zapril
	Tab 5 mg		90	✓ Zapril
ΕN	IALAPRIL MALEATE			
*	Tab 5 mg	1.82	100	✓ Acetec
*	Tab 10 mg	2.02	100	✓ Acetec
*	Tab 20 mg	2.42	100	✓ Acetec
LIS	SINOPRIL			
*	Tab 5 mg	2.07	90	 Ethics Lisinopril
*	Tab 10 mg	2.36	90	✓ Ethics Lisinopril
	Tab 20 mg		90	✓ Ethics Lisinopril

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	_	Subsidised	Generic
	\$	Per		Manufacturer
PERINDOPRIL			_	
Tab 2 mg		30		Apo-Perindopril
	4.95			Coversyl
Tab 4 mg		30		Apo-Perindopril
	6.30		•	Coversyl
Apo-Perindopril Tab 2 mg to be delisted 1 October 2021) Apo-Perindopril Tab 4 mg to be delisted 1 October 2021)				
QUINAPRIL				
Tab 5 mg	6.01	90	✓	Arrow-Quinapril 5
Tab 10 mg	3.16	90	/	Arrow-Quinapril 10
Tab 20 mg	4.89	90	✓	Arrow-Quinapril 20
ACE Inhibitors with Diuretics				
UINAPRIL WITH HYDROCHLOROTHIAZIDE				
Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	1	Accuretic
·- · · · · · · · · · · · · · · · ·	3.83	30		Accuretic 10
* Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	1	Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
₹ Tab 4 mg	2.00	90	1	Candestar
₹ Tab 8 mg		90	1	Candestar
F Tab 16 mg	3.31	90	1	Candestar
€ Tab 32 mg	5.26	90	1	Candestar
OSARTAN POTASSIUM				
€ Tab 12.5 mg	1 56	84	1	Losartan Actavis
€ Tab 25 mg		84		Losartan Actavis
€ Tab 50 mg		84		Losartan Actavis
€ Tab 100 mg		84		Losartan Actavis
Angiotensin II Antagonists with Diuretics				
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
Tab 50 mg with hydrochlorothiazide 12.5 mg	1 88	30	1	Arrow-Losartan &
rab oo mg war nyaroomoroanazido 12.0 mg		00	•	Hydrochlorothiazide
				ny ar comorcanazia
Angiotensin II Antagonists with Neprilysin Inhi	bitors			
ACUBITRIL WITH VALSARTAN - Special Authority see SA19	905 below – Retail pha	rmac	ev	
Note: Due to the angiotensin II receptor blocking activity of ACE inhibitor or another ARB.				be co-administered with
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	1	Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg		56		Entresto 49/51
Tab 07.0 mg with valoaten 100.0 mg		E 6		Entroote 07/102

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

continued...

✓ Entresto 97/103

56

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has heart failure; and
 - 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II: or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV: and
 - 3 Fither:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
 - 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthe	etics, Local, <mark>pa</mark>	ge 117	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg	3.80	30	✓ Aratac
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a			
PSO	16.37	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
PSO	12.07	10	✓ Hameln S29
	15.09		✓ Martindale
(Hameln S29 Inj 600 mcg per ml, 1 ml ampoule to be delisted 1 Jar	nuary 2022)		
DIGOXIN			
* Tab 62.5 mcg - Up to 30 tab available on a PSO	7.00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO		240	✓ Lanoxin
* Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			 Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	✓ Rythmodan
FLECAINIDE ACETATE			•
▲ Tab 50 mg	19.95	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg		90	✓ Flecainide
			Controlled
			Release Teva
▲ Cap long-acting 200 mg	61.06	90	✓ Flecainide
			Controlled
			Release Teva
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor

Restricted to children under 12 years of age.

BISOPROLOL FUMARATE

Principal Supply

CARVEDILOL

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
MEXILETINE HYDROCHLORIDE				
▲ Cap 150 mg	162.00	100		ANI \$29 Mexiletine Hydrochloride USP \$29
▲ Cap 250 mg	202.00	100	✓	Teva \$29 Mexiletine Hydrochloride USP \$29
ANI S29 Cap 150 mg to be delisted 1 January 2022)			•	Teva S29
Mexiletine Hydrochloride USP \$29 Cap 150 mg to be Mexiletine Hydrochloride USP \$29 Cap 250 mg to be PROPAFENONE HYDROCHLORIDE Tab 150 mg	delisted 1 January 2022)	50	•	Rytmonorm
Antihypotensives				
MIDODRINE – Special Authority see SA1474 below – F Tab 2.5 mg	53.00	100 100		Gutron Gutron
SA1474 Special Authority for Subsidy initial application from any relevant practitioner. Appro- not due to drugs. Note: Treatment should be started with small doses and the usual target is a standing systolic blood pressure of Renewal from any relevant practitioner. Approvals valid tenefiting from treatment.	ovals valid for 2 years where p d titrated upwards as necessa 90 mm Hg.	atient	has disab	ling orthostatic hypotension
Beta-Adrenoceptor Blockers				
Beta Adrenoceptor Blockers				
ATENOLOL				
⊀ Tab 50 mg ★ Tab 100 mg ★ Oral liq 25 mg per 5 ml	14.20	500 500 00 ml (OP 🗸	Mylan Atenolol Mylan Atenolol Atenolol AFT Atenolol AFT S29 829

✓ fully subsidised	S29	Unapproved medicine supplied under Section 29

90

90

90

60

60

60

Sole Subsidised Supply

✓ <u>Bisoprolol Mylan</u>✓ Bisoprolol Mylan

✓ Bisoprolol Mylan

✓ Carvedilol Sandoz

✓ Carvedilol Sandoz

✓ Carvedilol Sandoz

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LA	BETALOL				
*	Tab 100 mg	14.50	100		<u>Trandate</u>
*	Tab 200 mg		100	✓	<u>Trandate</u>
*	Inj 5 mg per ml, 20 ml ampoule	59.06	5		
		(88.60)			Trandate
*	inj 5 mg per ml, 20 ml vial	42.29	1		
		(48.20)			Alvogen S29
ME	TOPROLOL SUCCINATE				
*	Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
*	Tab long-acting 47.5 mg		30	/	Betaloc CR
*	Tab long-acting 95 mg		30		Betaloc CR
*	Tab long-acting 190 mg		30	/	Betaloc CR
ME	TOPROLOL TARTRATE				
*	Tab 50 mg	5.66	100	1	Apo-Metoprolol
*	Tab 100 mg		60		Apo-Metoprolol
*	Tab long-acting 200 mg		28		Slow-Lopresor
*	Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
		20.50	J	•	MELOPIOIOI IV MYIAII
	DOLOL Tab 40 mm	10.00	100		Ana Nadalal
	Tab 40 mg		100		Apo-Nadolol
	Tab 80 mg	20.43	100	•	Apo-Nadolol
	NDOLOL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who vendorsed accordingly. Pharmacists may annotate the predispensing of pindolol.	escription as endorsed v	vhere	there exist	s a record of prior
	Tab 5 mg		100		Apo-Pindolol
	Tab 10 mg		100		Apo-Pindolol
*	Tab 15 mg	33.31	100	•	Apo-Pindolol
٠,	po-Pindolol Tab 5 mg to be delisted 1 May 2022)				
٠,	po-Pindolol Tab 10 mg to be delisted 1 May 2022)				
٠,	po-Pindolol Tab 15 mg to be delisted 1 May 2022)				
PR	OPRANOLOL				
*	Tab 10 mg		100		Apo-Propranolol
*	Tab 40 mg	5.72	100		Apo-Propranolol
*	Cap long-acting 160 mg	18.17	100	•	Cardinol LA
*	Oral liq 4 mg per ml - Special Authority see SA1327 bel	ow –			
	Retail pharmacy	CBS	500 m	nl 🗸	Roxane-
	-				

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Propranolol \$29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OTALOL				
₭ Tab 80 mg		500		Mylan
Fab 160 mg	10.98	100	•	<u>Mylan</u>
Calcium Channel Blockers				
Dihydropyridine Calcium Channel Blockers				
MLODIPINE				
★ Tab 2.5 mg		90	•	<u>Vasorex</u>
₭ Tab 5 mg	0.96	90		Vasorex
├ Tab 10 mg	1.19	90	1	Vasorex
ELODIPINE				
★ Tab long-acting 2.5 mg	1.45	30	✓	Plendil ER
₭ Tab long-acting 5 mg		90	✓	Felo 5 ER
₭ Tab long-acting 10 mg		90	_	Felo 10 ER
NIFEDIPINE				
₭ Tab long-acting 10 mg	18.80	56	✓	Tensipine MR10 S29
₭ Tab long-acting 20 mg	0.10	50	./	Mylan (12 hr
* Tab long-acting 20 mg		50	•	Mylan (12 hr
	47.70	465		release) \$29
	17.72	100		Nyefax Retard
* Tab long-acting 30 mg	34.10	100	•	Mylan (24 hr release) \$29
* Tab long-acting 60 mg	52.81	100	•	Mylan (24 hr release) S29
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
★ Tab 60 mg	8.50	100	1	Dilzem
K Cap long-acting 120 mg		500	✓	Apo-Diltiazem CD
K Cap long-acting 180 mg		500		Apo-Diltiazem CD
Cap long-acting 240 mg		500	1	Apo-Diltiazem CD
Dilzem Tab 60 mg to be delisted 1 January 2022)				
PERHEXILINE MALEATE				
★ Tab 100 mg	62.90	100	/	Pexsig
/ERAPAMIL HYDROCHLORIDE				<u>~~a</u>
* Tab 40 mg	7.01	100	J	Isoptin
₭ Tab 80 mg		100		Isoptin
				•
* Tab long-acting 120 mg	36.02	100		Isoptin Retard \$29
K. Tob long acting 040 mg	15 10	20		Isoptin SR
* Tab long-acting 240 mg		30	•	Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5	1	Isoptin
				·
Centrally-Acting Agents				
CLONIDINE	40.51			
★ Patch 2.5 mg, 100 mcg per day — Only on a prescription		4		Mylan
★ Patch 5 mg, 200 mcg per day — Only on a prescription		4		<u>Mylan</u>
Patch 7.5 mg, 300 mcg per day – Only on a prescription	16.93	4	/	Mylan

		•	JANL	IUVAS	CULAR SYSTEM
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
* Tab 150 mcg	OCHLORIDE r ml, 1 ml ampoule	37.07	112 100 10	✓	Clonidine BNM Catapres Medsurge
METHYLDOPA * Tab 250 mg		15.10 52.85	100 500		Methyldopa Mylan Methyldopa Mylan S29 S29
Diuretics					
Loop Diuretic	s				
Ü	r ml, 4 ml vial	16.36	30 100 5	✓	Burinex S29 S29 Burinex
* Tab 40 mg - L	lp to 30 tab available on a PSO		1,000 50	✓	Apo-Furosemide Urex Forte Furosemid- Ratiopharm \$29
		169.96	100	•	Furosemid- Ratiopharm S29
* Inj 10 mg per n	per ml nl, 25 ml ampoule nl, 2 ml ampoule – Up to 5 inj available	60.65	0 ml O 6 5	1	<u>Lasix</u> <u>Lasix</u> <u>Furosemide-Baxter</u>
Potassium Sp	paring Diuretics				
	er ml		5 ml O	P 🗸	Biomed
Tab 50 mg Tab 25 mg	Special Authority see SA1728 below – R	17.00	30 30		Inspra Inspra
Initial application the following criteria Both: 1 Patient has 2 Either: 2.1 Patie	I Authority for Subsidy from any relevant practitioner. Approva a: heart failure with ejection fraction less the series intolerant to optimal dosing of spiront has experienced a clinically significant	an 40%; and			
METOLAZONE Tab 5 mg		CBS	1	/	Metolazone 629

✓ Zaroxolyn S29

50

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy	\ 0::	Fully Brand or
	(Manufacturer's Pric	ce) Subsi Per	idised Generic Manufacturer
SPIRONOLACTONE			
* Tab 25 mg	4.38	100	✓ Spiractin
* Tab 100 mg		100	✓ Spiractin
Oral liq 5 mg per ml		25 ml OP	✓ Biomed
Potassium Sparing Combination Diuretics			
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
* Tab 5 mg with furosemide 40 mg	8.63	28	✓ Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIE			
* Tab 5 mg with hydrochlorothiazide 50 mg		50	✓ Moduretic
- 145 5 mg with hydrodillorodillazide 50 mg		50	- INIOGUI GUO
Thiazide and Related Diuretics			
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]	00.00	F00	/ A
* Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ Arrow-
			<u>Bendrofluazide</u>
May be supplied on a PSO for reasons other than emerge	ency.		
* Tab 5 mg	•	500	✓ Arrow-
·			Bendrofluazide
CHI ODOTHIAZIDE			
CHLOROTHIAZIDE Oral lig 50 mg por ml	26.00	25 ml OP	✓ Biomed
Oral liq 50 mg per ml	∠0.00	20 IIII UP	▼ bioilleu
CHLORTALIDONE [CHLORTHALIDONE]			-
Tab 25 mg		30	✓ Igroton \$29
	6.50	50	✓ <u>Hygroton</u>
NDAPAMIDE			4.5
* Tab 2.5 mg		90	✓ <u>Dapa-Tabs</u>
	11.61	100	✓ Mylan
			Indapamide \$29
Lipid-Modifying Agents			
Fibrates			
BEZAFIBRATE			
* Tab 200 mg		90	✓ Bezalip
* Tab long-acting 400 mg	12.89	30	Bezalip Retard
Other Lipid-Modifying Agents			
ACIPIMOX			
* Cap 250 mg	21.56	30	Olbetam
			✓ Olbetam S29 S29
Resins			
COLESTIPOL HYDROCHLORIDE			
Grans for oral liq 5 g	32.89	30	✓ Colestid
			y

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 80 mg	9.24 14.92	500 500 500 500	✓	Lorstat Lorstat Lorstat Lorstat
PRAVASTATIN * Tab 20 mg * Tab 40 mg	2.11	28 28	✓	Pravastatin Mylan Pravastatin Mylan
SIMVASTATIN * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 80 mg	2.03 3.58	90 90 90 90	√	Simvastatin Mylan Simvastatin Mylan Simvastatin Mylan Simvastatin Mylan
Salective Chalesteral Absorption Inhibitors				

Selective Cholesterol Absorption Inhibitors

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:

2.0 mmol/litre.

- 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
- 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
- 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN – Special Authority see SA1046 below – Retail pharmacy

30 ✓ Zimyb e	30	5.15	Tab 10 mg with simvastatin 10 mg
30 ✓ Zimyb e	30	6.15	Tab 10 mg with simvastatin 20 mg
30 ✓ Zimyb e	30	7.15	
30 ✓ Zimybe	30	8.15	Tab 10 mg with simvastatin 80 mg

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

continued...

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

不	available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISC	OSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
*	Tab long-acting 60 mg	9.25	90	✓ Duride

Sympathomimetics

Α	D	R	FI	N	Α	П	IN	ΙE

Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	Aspen Adrenaline

Vasodilators

HYDRAI AZINE HYDROCHI ORIDE

*	Tab 25 mg - Special Authority see SA1321 below - Retail		
	pharmacyCBS	1	 Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM S29
		100	✓ Onelink S29
*	Inj 20 mg ampoule25.90	5	✓ Apresoline

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

Reddy's

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised	
MINOXIDII	Ψ			manadaror
MINOAIDIE ▲ Tab 10 mg	70.00	100	1	Loniten
· ·	70.00	100	•	Loniten
NICORANDIL	05.53		,	
▲ Tab 10 mg		60		Ikorel
▲ Tab 20 mg	32.28	60	•	<u>lkorel</u>
PAPAVERINE HYDROCHLORIDE			_	
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	•	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	1	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN - Special Authority see SA1702 below - Reta	il pharmacy			
Tab 5 mg		30	1	Ambrisentan Mylan
Tab 10 mg		30		Ambrisentan Mylan
⇒SA1702 Special Authority for Subsidy				
Special Authority approved by the Pulmonary Arterial Hypertens	sion Panel			
Notes: Application details may be obtained from Pharmac's we		c do/	rt nz/SAFo	irms or
The Coordinator. PAH Panel	bono <u>conoccionamia</u>	o.go	11.112/0711	on.
Pharmac, PO Box 10-254, WELLINGTON				
Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharma	ic.govt.nz			
BOSENTAN - Special Authority see SA1991 below - Retail ph				
Tab 62.5 mg	•	60	1	Bosentan Dr
y				Reddy's
Tab 125 mg	440.05	60	,	Bosentan Dr

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*: and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.3.2.1 Patient is on the lung transplant list; or
- 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
- 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
- 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail pharmacy		
Tab 25 mg0.85	4	✓ Vedafil
Tab 50 mg1.70	4	✓ Vedafil
Tab 100 mg10.20	12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*: and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

continued...

Subsidy	Fully	Brand or	
•	. ,		
(Manufacturer's Price)	Subsidised	Generic	
\$	Per ✓	Manufacturer	
Ψ		manadatata	

continued...

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications, and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 412 Fither
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retai	l pharmacy		
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri
- CA1606 Chaolal Authority for Cubaidy			

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WFI LINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Re	etail pharmacy	-	
Cap 5 mg	8.14	60	Oratane
Cap 10 mg	13.34	120	Oratane
Cap 20 mg	20.49	120	Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription15.57 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 88

HYDROGEN PEROXIDE

111	DITOGEN I ETIONIDE		
*	Crm 1%8.56	10 g OP	Crystaderm
		15 g OP	Crystaderm

			LIMAI	OLOGICALO
	Subsidy (Manufacturer's F \$	Price) Subs	idised G	rand or eneric anufacturer
MUPIROCIN Oint 2%	6.60 (10.50)	15 g OP	Bact	oban
a) Only on a prescriptionb) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	✓ Foba	ın
b) Only on a prescription c) Not in combination Oint 2%	1.50	5 a OB	✓ Foba	·n
a) Maximum of 5 g per prescription b) Only on a prescription c) Not in combination	1.59	5 g OP	V FODA	ın
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ Flam	azine
a) Up to 250 g available on a PSOb) Not in combination				
Antifungals Topical				
For systemic antifungals, refer to INFECTIONS, Antifungals, AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ Myc	nNail
CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination		3 m 31	inyo.	/////////////////////////////////////
Nail-soln 8%	5.72	7 ml OP	✓ Apo-	Ciclopirox
* Crm 1% a) Only on a prescription b) Not in combination		20 g OP	✓ Clon	nazol
* Soln 1%	4.36 (7.55)	20 ml OP	Cane	esten
b) Not in combination ECONAZOLE NITRATE				
Crm 1%	1.00 (7.48)	20 g OP	Peva	ryl
a) Only on a prescription b) Not in combination Foaming soln 1%, 10 ml sachets	9.89	3		
a) Only on a prescription b) Not in combination	(17.23)	-	Peva	ryl

DERMATOLOGICALS

	Subsidy (Manufacturer's F	Price) Subs	Fully	Brand or Generic
	(Manufacturers i	Per Subs	siuiseu ✓	Manufacturer
IICONAZOLE NITRATE				
₭ Crm 2%	0.81	15 g OP	✓ <u>N</u>	<u>//ultichem</u>
a) Only on a prescriptionb) Not in combination				
k Lotn 2%	4.36 (10.03)	30 ml OP		Daktarin
a) Only on a prescription				
b) Not in combination	4.00	00 ml OD		
≰ Tinct 2%	4.36	30 ml OP	г	Daktarin
a) Only on a prescription	(12.10)			Januaiii i
b) Not in combination				
,				
Antipruritic Preparations				
CALAMINE			-	
a) Only on a prescription				
b) Not in combination	4.00	400	,.	
Crm, aqueous, BP	1.26	100 g	√ r	nealthE Calamine Aqueous Cream BP
ROTAMITON				
a) Only on a prescription				
b) Not in combination	0.55	00 05		
Crm 10%	3.29	20 g OP	✓	tch-Soothe
MENTHOL – Only in combination				
 Only in combination with a dermatological base With or without other dermatological galenicals. 	or proprietary Topical C	Corticosteriod –	Plain	
Crystals	6.92	25 g	✓ N	/lidWest
,	29.60	100 g		/lidWest
Corticosteroids Topical				
or systemic corticosteroids, refer to CORTICOSTEROID	S AND RELATED AGE	NTS, page 78		
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE				
Crm 0.05%		15 g OP	_	Diprosone
	36.00	50 g OP	-	<u>Diprosone</u>
Oint 0.05%		15 g OP	-	<u>Diprosone</u>
	36.00	50 g OP		Diprosone

	Crm 0.05%	2.96	15 g OP	✓ <u>Diprosone</u>
		36.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BE	TAMETHASONE VALERATE			
*	Crm 0.1%	4.53	50 g OP	✓ Beta Cream
*	Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
*	Lotn 0.1%	18.00	50 ml OP	✓ Betnovate
CL	OBETASOL PROPIONATE			
*	Crm 0.05%	2.18	30 g OP	✓ Dermol
*	Oint 0.05%	2.12	30 g OP	✓ Dermol
			3 -	

	Subsidy		Fully Brand or	
	(Manufacturer's F	Price) Subs Per	idised Generic Manufacturer	
OLODETA COME DUTYDATE	Ψ	1 61	Wallulacturer	
CLOBETASONE BUTYRATE	F 00	00 = OD		
Crm 0.05%		30 g OP	5	
	(10.00)		Eumovate	
HYDROCORTISONE				
* Crm 1% - Only on a prescription	3.70	100 g OP	 Hydrocortisone 	
			(PSM)	
	17.15	500 g	✓ Hydrocortisone	
			(PSM)	
* Powder - Only in combination	49.95	25 g	✓ ABM	
Up to 5% in a dermatological base (not proprietary Topic	al Corticosterio	d – Plain) with o	r without other dermatol	ogical
galenicals		,		J
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only o	n			
a prescription		250 ml	✓ DP Lotn HC	
	10.57	250 1111	DF LOUITIC	
HYDROCORTISONE BUTYRATE	0.05	100 00	4	
Lipocream 0.1%		100 g OP	✓ Locoid Lipocrear	n
Oint 0.1%		100 g OP	Locoid	
Milky emul 0.1%	12.33	100 ml OP	✓ Locoid Crelo	
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	✓ Advantan	
Oint 0.1%	4.46	15 g OP	✓ Advantan	
MOMETASONE FUROATE				
Crm 0.1%	1.51	15 a OP	✓ Elocon Alcohol F	ree
	2.50	50 g OP	✓ Elocon Alcohol F	:ree
Oint 0.1%	1.51	15 g OP	✓ Elocon	
	2.90	50 g OP	✓ Elocon	
Lotn 0.1%	6.30	30 ml OP	✓ Elocon	
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	✓ Aristocort	
Oint 0.02%		100 g OP	✓ Aristocort	
<u> </u>		.00 g 0.		
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS	SIDIC ACIDI			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
((10.45)		Fucicort	
a) Maximum of 15 g per prescription	(10110)			
b) Only on a prescription				
	·:			
HYDROCORTISONE WITH MICONAZOLE — Only on a prescrip		45 = OD	/ Miawawa II	
* Crm 1% with miconazole nitrate 2%		15 g OP	✓ Micreme H	
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O	, ,			
Crm 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	✓ Pimafucort	
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	✓ Pimafucort	
(Pimafucort Crm 1% with natamycin 1% and neomycin sulphate (0.5% to be delis	ted 1 May 2022 _,	")	
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	N AND NYSTA	TIN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg	1			
and gramicidin 250 mcg per g - Only on a prescription	,	15 g OP		
3 31 3 7 1 33 1 3	(9.28)	3	Viaderm KC	
	` '			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Barrier Creams and Emollients

DIMETHICONE		
* Crm 5% pump bottle4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle4.52	500 ml OP	✓ healthE
		Dimethicone 10%
ZINC AND CASTOR OIL * Oint	F00 ~	✓ Boucher
未 OIII	500 g	▼ Boucher
Emollients		
AQUEOUS CREAM		
* Crm	500 g	✓ Basic AquaCream✓ Boucher
		✓ Medco
5.75		✓ Topiderm
(Topiderm Crm to be delisted 1 September 2021)		
CETOMACROGOL * Crm BP	500 g	✓ healthE
CETOMACROGOL WITH GLYCEROL	3	
Crm 90% with glycerol 10%2.35	500 ml OP	✓ ADE
		 ✓ Boucher ✓ Kenkay Sorbolene
		✓ Pharmacy Health
		Sorbolene with
3.10	1,000 ml OP	Glycerin ✓ ADE
	.,	✓ Boucher
EMULSIFYING OINTMENT		4
* Oint BP	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION		<u></u>
* Crm2.19	500 g	✓ O/W Fatty Emulsion
DADAFIN		Cream
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%5.35	500 ml OP	✓ healthE
UREA		
		_

100 g OP

✓ healthE Urea Cream

	Subsidy		Fully Brand or
	(Manufacturer's P		sidised Generic
	\$	Per	✓ Manufacturer
WOOL FAT WITH MINERAL OIL - Only on a prescription			
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
	(11.95)		DP Lotion
	1.40	250 ml OP	
	(4.53)		DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft - Only in combination	4.99	450 g	✓ healthE
•	19.99	2,500 g	✓ healthE
Only in combination with a dermatological galenical or	as a diluent for a p		ical Corticosteroid – Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
•	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIMETHICONE

*	✓ <u>healthE</u> <u>Dimethicone 4%</u> Lotion
IVE	✓ Stromectol
	1

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...

DERMATOLOGICALS

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — **(Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5% 5.75 Lotn 5% 3.99	0 -	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN		
Shampoo 0.5%11.36	200 ml OP	Parasidose
(Parasidose Shampoo 0.5% to be delisted 1 January 2022)		

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail p	harmacy		
Cap 10 mg	17.86	60	Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g	35 60 g OP	✓ Enstilar✓ Daivobet✓ Daivobet
CALCIPOTRIOL Oint 50 mcg per g40.0	00 120 g OP	✓ Daivonex

	Subsidy		Fully Brand or
	(Manufacturer's Pri		sidised Generic
OOAL TAD	\$	Per	✓ Manufacturer
COAL TAR Soln BP - Only in combination	36.25	200 ml	✓ Midwest
 Up to 10% only in combination with a dermatologi With or without other dermatological galenicals. 		tary Topical (
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% ar allantoin crm 2.5%		75 g OP	
and to the Lie /e	(8.00)	70 g 01	Egopsoryl TA
	3.43	30 g OP	
	(4.35)		Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	4.07	05 - OD	(Occas Occale
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP	✓ Coco-Scalp✓ Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Reta		40 g Oi	· Ooco-ocaip
a) Maximum of 15 g per prescription	iii priarriacy		
b) Note: a maximum of 15 g per prescription and no more	than one prescripti	on per 12 we	eks.
Cream 1%		15 g OP	✓ <u>Elidel</u>
neeting the following criteria: 3oth: 1 Patient has atopic dermatitis on the eyelid; and 2 Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure.	ppical corticosteroid	ds, cataracts,	glaucoma, or raised intraocula
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE Soln 2.3% with trolamine laurilsulfate and fluorescein sodiur		a prescription 500 ml	n ✓ <u>Pinetarsol</u>
SALICYLIC ACID	40.00	0.50	
Powder - Only in combination	18.88	250 g	✓ Midwest ✓ PSM
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topica	al Corticostero	
SULPHUR			
Precipitated – Only in combination		100 g	✓ Midwest
 Only in combination with a dermatological base or With or without other dermatological galenicals. 	proprietary Topica	ii Corticosterd	oid – Piain
Scalp Preparations			
BETAMETHASONE VALERATE ★ Scalp app 0.1%	0.84	100 ml OP	✓ Beta Scalp
	5.04	100 IIII OP	- Deta Stalp
CLOBETASOL PROPIONATE ★ Scalp app 0.05%	5 69	30 ml OP	✓ Dermol
IYDROCORTISONE BUTYRATE		30 1111 01	- <u>Bonnor</u>
Scalp lotn 0.1%	6.57	100 ml OP	✓ Locoid

DERMATOLOGICALS

	Subsidy (Manufacturer's Pri \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	√ <u>S</u>	sebizole

Sunscreens

Wart Preparations

Other Skin Preparations

Antineoplastics

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

Contraceptives - Non-hormonal

Condoms

-	DOMS 49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments
	53 mm		10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
: 5	53 mm, 0.05 mm thickness	0.95	10	✓ <u>Moments</u>
		11.42	144	✓ Moments
	 a) Up to 60 dev available on a PSO 			
	b) Maximum of 60 dev per prescription			
: {	53 mm, chocolate, brown		10	✓ <u>Moments</u>
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.05	40	
• 5	53 mm, strawberry, red		10	✓ Moments
	a) He to 00 downwilled 1 200	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.07	10	✓ Moments
• 5	56 mm	0.97	10 144	✓ <u>Moments</u> ✓ Moments
	a) Maximum of 60 day per prescription	11.04	144	• WOMENIS
	a) Maximum of 60 dev per prescriptionb) Up to 60 dev available on a PSO			
:	56 mm, 0.05 mm thickness	1 20	12	✓ Gold Knight
• •	70 IIIII, 0.00 IIIII UII0NIIG93	15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO	10.07	177	- Gold Kingill
	b) Maximum of 60 dev per prescription			
÷	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
•	a) Maximum of 60 dev per prescription			<u> </u>
	b) Up to 60 dev available on a PSO			
+ 5	56 mm, 0.08 mm thickness	0.97	10	✓ Moments
•	,	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
+ 5	56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
	•	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
+ 5	56 mm, chocolate	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
: 5	56 mm, strawberry		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
• (60 mm		12	✓ Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

GENITO-URINARY SYSTEM

b) Up to 60 dev available on a PSO

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	√ <u>G</u>	old Knight XL
	a) Maximum of 60 dev per prescription				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IÚD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓ Choice
				TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to		
	84 tab available on a PSO10.00	84	✓ Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	84	
	(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

(Marvelon 28 Tab 30 mcg with desogestrel 150 mcg and 7 inert tab to be delisted 1 November 2021)

_					_
		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	I Generic
		\$	Per	•	Manufacturer
ET	HINYLOESTRADIOL WITH LEVONORGESTREL				
*	Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -	_			
	Up to 112 tab available on a PSO	2.18	84	1	Microgynon 20 ED
	•	6.45	112	✓	Femme-Tab ED
*	Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - Up)			
	to 84 tab available on a PSO		84	1	Microgynon 50 ED
*	Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		•
		(16.50)			Microgynon 30
	a) Higher subsidy of \$15.00 per 63 tab with Special Auth	nority see SA0500 or	the p	revious p	age
	b) Up to 63 tab available on a PSO	,			
*	Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets -	_			
	Up to 112 tab available on a PSO		84	1	Levlen ED
	- F	6.45	112	1	Femme-Tab ED
ET	HINYLOESTRADIOL WITH NORETHISTERONE				
	Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to				
	84 tab available on a PSO		84	1	Brevinor 1/28
	Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U		04	•	Dicvillor 1/20
	to 84 tab available on a PSO		84	./	Necon
	10 04 Idu avallable oli a F50		04	_	
		21.99		•	Norimin

(Necon Tab 35 mcg with norethisterone 500 mcg and 7 inert tab to be delisted 1 September 2021)

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	✓ Microlut
		22.00	112	✓ Microlut
*	Subdermal implant (2 × 75 mg rods) - Up to 3 pack available			
	on a PSO	106.92	1	✓ <u>Jadelle</u>

GENITO-URINARY SYSTEM

(1	Subsidy Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSI NORETHISTERONE	O7.98	1	✓ [Depo-Provera
Tab 350 mcg - Up to 84 tab available on a PSO	6.25	84	√	Noriday 28
Emergency Contraceptives				
LEVONODOFOTDEL				

LEVONORGESTREL

- - a) Maximum of 2 tab per prescription
 - b) Up to 5 tab available on a PSO
 - c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

ACETIC ACID WITH HADDOAAOHINOLINE AND DICINOLEIC VOID

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	✓ Micreme
NYSTATIN	ū	
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat
raginal of the roots of port of their applicator(o)	, o g o.	- Itilotat

Myometrial and Vaginal Hormone Preparations

. Ergometrine
stin
stin
tocin BNM
tocin BNM

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic	
	\$	Per	1	Manufacturer	
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj av	ailable on a PSO				
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml	30.00	5	✓ Sy	yntometrine	

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

✓ David One Step Cassette Pregnancy Test

✓ Smith BioMed Rapid Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking oxybutynin prior to 1 June 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of oxybutynin.

 ★ Tab 5 mg
 11.70
 500
 ✓ Apo-Oxybutynin

 ★ Oral liq 5 mg per 5 ml
 60.40
 473 ml
 ✓ Apo-Oxybutynin

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 below -

Retail pharmacy......31.80 200 ml OP ✓ Biomed

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

*	Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SO	LIFENACIN SUCCINATE			
	Tab 5 mg	2.05	30	✓ Solifenacin Mylan
	Tab 10 mg	3.72	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE

*	Compound diagnostic sticks	7.50	50 test OP	
	(1	8.25)		Hemastix

TETRABROMOPHENOL

*	Blue diagnostic strips7.02	100 test OP	
	(13 92)		Alhustiy

Obstetric Preparations

Antiprogesterones

MI	FΕ	PR	IST	$\Gamma \cap I$	ΝF	
IVII		гΠ	IO I	v	VL.	

✓ Mifegyne	1	Tab 200 mg60.00
✓ Mifegyne	3	180.00

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Por 🗸	Manufacturer	

Calcium Homeostasis

$C\Delta$		

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable210.30 28 ✓ Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA2031 below −
Retail pharmacy.......18.00 1 ✓ Zoledronic acid
Mylan

⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

Subsid	dy Full	/ Brand or
(Manufacture	r's Price) Subsidise	d Generic
\$	Per 💌	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASO	NE ACETATE	Ī	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20	5	
	(36.96)		Celestone
			Chronodose
DEXAMETHASONE			
* Tab 0.5 mg - Up to 60 tab available on a PSO	1.50	30	✓ Dexmethsone
* Tab 4 mg - Up to 30 tab available on a PSO		30	✓ Dexmethsone
Oral liq 1 mg per ml		25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE			
Dexamethasone phosphate injection will not be funded for oral u	ise		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO.		10	✓ Dexamethasone
and the state of t		. •	Phosphate
			Panpharma
* Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO.	16.37	10	✓ Dexamethasone
and the state of t		. •	Phosphate
			Panpharma
FLUDROCORTISONE ACETATE			
* Tab 100 mcg	1/132	100	✓ Florinef
•	14.02	100	• Homie
HYDROCORTISONE	0.40	100	(December
* Tab 5 mg		100	✓ Douglas✓ Douglas
* Tab 20 mg * Inj 100 mg vial		100 1	✓ Douglas ✓ Solu-Cortef
, ,	4.38	1	▼ Solu-Cortei
a) Up to 5 inj available on a PSO			
b) Only on a PSO a) Solu Cost of to be Principal Supply on 1 Nevember 2021			
c) Solu-Cortef to be Principal Supply on 1 November 2021			
METHYLPREDNISOLONE			
* Tab 4 mg		100	✓ Medrol
* Tab 100 mg	194.00	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial	18.90	1	✓ Solu-Medrol-Act-
			O-Vial
Let 405 monetal	00.00	_	Code Made Lat
Inj 125 mg vial	28.90	1	✓ Solu-Medrol-Act-
			O-Vial
Inj 500 mg vial	22 78	1	✓ Solu-Medrol-Act-
inj ood ing via	22.70	•	O-Vial
			O Tidi
Inj 1 g vial	27.83	1	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE			
Inj 40 mg per ml, 1 ml vial	44.40	5	✓ Depo-Medrol
		Ū	Sobo modioi

	Subsidy		Fully	Brand or
(Manufacturer's Pi		idised	Generic
	\$	Per		Manufacturer
PREDNISOLONE				
Yoral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00	30 ml OP	√ F	Redipred
PREDNISONE				
* Tab 1 mg	18.58	500	√	Apo-Prednisone
* Tab 2.5 mg		500	√	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	√	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	√	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen S29
				AU Synacthen
			✓ 9	Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	√ 9	Synacthen Depot
			✓ 9	Synacthene
				Retard \$29
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓	Cenacort-A 10
	26.62		✓ <u>F</u>	Adcortyl \$29
Inj 40 mg per ml, 1 ml ampoule	11.30	1		riaver S29
.,g p,p	51.10	5	_	Cenacort-A 40
	70.62		_	Kenalog S29
(Adcortyl \$29 Inj 10 mg per ml, 1 ml ampoule to be delisted 1 Oct				tonalog
(Triaver \$29 Inj 40 mg per ml, 1 ml ampoule to be delisted 1 Octo	,			
, , , ,	,			
(Kenalog S29 Inj 40 mg per ml, 1 ml ampoule to be delisted 1 Oct	ober 2021)			

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

50	Rex S29
•	Siterone
50	Siterone
30	Androderm
1	Depo-Testosterone
1 1	Sustanon Ampoules
	oustation Ampoules
00	Audulul Tankanan
	Andriol Testocaps
1	Reandron 1000
	50

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

		Subsidy (Manufacturer's Price	e) Sub	Fully sidised	Brand or Generic
		\$	Per	√	Manufacturer
Oe	estrogens				
DES	TRADIOL - See prescribing guideline on the previous page				
*	Tab 1 mg	4.12	28 OP		
		(11.10)		Е	strofem
*	Tab 2 mg		28 OP	_	
		(11.10)			strofem
*	Patch 100 mcg per 24 hours	7.91	4	✓ (Climara
	No more than 1 patch per week				
	b) Only on a prescription				
*	Patch 50 mcg per 24 hours	7.04	4	✓ (Climara
	a) No more than 1 patch per week				
	b) Only on a prescription				
	Patch 25 mcg per day		8		stradot
		7.85		✓ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 50 mcg per day	7.04	8	√ E	stradot 50 mcg
		9.22		✓ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 75 mcg per day	7.91	8	√ E	stradot
		10.60		✓ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				•
	b) Only on a prescription				
	Patch 100 mcg per day	7.91	8	√ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
Clin	nara Patch 100 mcg per 24 hours to be delisted 1 January 20	122)			
	nara Patch 50 mcg per 24 hours to be delisted 1 January 202				
	TRADIOL VALERATE – See prescribing guideline on the pro-				
	Tab 1 mg		84	√ □	rogynova
	Tab 1 mg		84		rogynova Progynova
	· ·		04	• -	Togyttova
	TROGENS – See prescribing guideline on the previous page				
*	Conjugated, equine tab 300 mcg		28	_	
	Operior and a service state OOF areas	(17.50)	00	۲	remarin
*	Conjugated, equine tab 625 mcg		28	_	
		(17.50)		۲	Premarin
Pr	ogestogens				
ΛΕΓ	PROXYPROGESTERONE ACETATE - See prescribing guid	eline on the previo	ous page		
	Tab 2.5 mg		30	√ P	rovera
	Tab 5 mg		100	-	rovera
	Tab 10 mg		30		rovera
••	144 14 mg		00	• 1	

	Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer							
Progestogen and Oestrogen Combined Preparations								
OESTRADIOL WITH NORETHISTERONE - See presc	eribing guideline on page 79							
* Tab 1 mg with 0.5 mg norethisterone acetate								
Tab Ting Mar 6.5 mg Noroanotorono acotato	(18.10) Kliovance							
* Tab 2 mg with 1 mg norethisterone acetate	,							
Tub 2 mg war i mg nordanotorone doctate	(18.10) Kliogest							
* Tab 2 mg with 1 mg norethisterone acetate (10), and	, ,							
· /·	· ·							
oestradiol tab (12) and 1 mg oestradiol tab (6)								
	(18.10) Trisequens							
Other Oestrogen Preparations								
Other Destroyen Freparations								
ETHINYLOESTRADIOL								
* Tab 10 mcg	17.60 100 V NZ Medical and							
· · · · · · · · · · · · · · · · · · ·	Scientific							
OESTRIOL								
* Tab 2 mg	7.00 30 V Ovestin							
* Tab 2 Hig	7.00 30 • Ovestin							
Other Progestogen Preparations								
Other i rogestogen i reparations								
LEVONORGESTREL								
* Intra-uterine device 52 mg	269.50 1 Mirena							
* Intra-uterine device 13.5 mg								
MEDROXYPROGESTERONE ACETATE								
Tab 100 mg	116.15 100 ✓ Provera HD							
· ·	- 11070112							
NORETHISTERONE	5.40 00 A Drive alist N							
* Tab 5 mg - Up to 30 tab available on a PSO	5.49 30 •• Primolut N							
PROGESTERONE								
Cap 100 mg - Special Authority see SA1609 below								
pharmacy	16.50 30 Vtrogestan							

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Cubaidy

Eully.

Drand or

		Subsidy (Manufacturer's Price) \$	Per	Subsidised	
Th	nyroid and Antithyroid Agents				
CAF	RBIMAZOLE				
*	Tab 5 mg	10.80	100		Neo-Mercazole Neo-Mercazole S29 S29
LEV	OTHYROXINE				
*	Tab 25 mcg	5.55	90	✓	Synthroid
	Tab 50 mcg		28	✓	Mercury Pharma
		5.79	90	✓	Synthroid
		64.28	1,000	/	Eltroxin
*	Tab 100 mcg	1.78	28	✓	Mercury Pharma
	-	6.01	90	✓	Synthroid
		66.78	1,000	/	Eltroxin
	DPYLTHIOURACIL – Special Authority see SA1199 below Propylthiouracil is not recommended for patients under the treatments are contraindicated.	, ,	the	patient is p	regnant and other
	Tab 50 mg	35.00	100	✓	PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see S	A2032 below – Retail pharn	nacy	
*	Inj 5 mg cartridge	69.75	í	Omnitrope
	Inj 10 mg cartridge		1	✓ Omnitrope
	Inj 15 mg cartridge		1	✓ Omnitrope
_				•

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

continued...

- children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate: and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and

Subsidy	,	Fully	Brand or	
(Manufacturer's		ubsidised	Generic	
<u> </u>	Per		Manufacturer	_
				_

continued...

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

Implant 3.6 mg, syringe65.68

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues	
----------------	--

GOSFREI IN

Implant 10.8 mg, syringe	122.37	1	✓ <u>Teva</u>
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a child of goserelin and the prescription is endorsed accordingly.	or adolescent a	nd is unable	to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
	(591.68)		Lucrin Depot 3-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy	(221.60) 177.50	1	·

Wach	pressi	\mathbf{n} \mathbf{A} \mathbf{a}	ANI	210
	4.4		чш	-1
		7		

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

1

Teva

	Subsidy		Fully	Brand or
	(Manufacturer's	,	sidised	Generic
	\$	Per		Manufacturer
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	Minirin
Tab 200 mcg	54.45	30	✓ N	Minirin
▲ Nasal drops 100 mcg per ml	39.03	2.5 ml OP	✓ N	Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ [esmopressin-
				PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	✓ N	Minirin
ing 4 mag per mi, 1 mi	07.10	10	J 10	

CABERGOLINE

Tab 0.5 mg - Maximum of 2 tab per prescription; can be		
waived by Special Authority see SA1370 below3.75	2	Dostinex
15.20	8	Dostinex

⇒SA1370 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 pathological hyperprolactinemia; or
- 2 acromegaly*.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

\cap	\bigcirc N	ΛIΕ		CIT	DΛ.	TE
l d	(JI	/11 [гілг	(7)	\neg A	1 -

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE			
Cap 250 mg5	58.00	50	✓ <u>Metopirone</u>

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

Anthelmintics

ALBENDAZOLE – Special Authority see SA1318 b	pelow – Retail pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	7.97	6	✓ Vermox
Oral lig 100 mg per 5 ml		15 ml	
	(7.53)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 60
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 229

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ Ranbaxy-Cefaclor
CEFALEXIN			
Cap 250 mg	3.33	20	✓ Cephalexin ABM
Cap 500 mg		20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable	8.75	100 ml	✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable	11.75	100 ml	✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement			
Only if prescribed for dialysis or cellulitis in accordance with accordingly.	a DHB approved	protocol and	he prescription is endorsed
Inj 500 mg vial	3.39	5	✓ <u>AFT</u>
Inj 1 g vial	3.49	5	✓ <u>AFT</u>
CEFTRIAXONE - Subsidy by endorsement			
a) Up to 10 inj available on a PSO			
 Subsidised only if prescribed for a dialysis or cystic fibros pelvic inflammatory disease, or the treatment of suspecte endorsed accordingly. 			
Inj 500 mg vial	0.89	1	✓ Ceftriaxone-AFT
Inj 1 g vial		5	✓ Ceftriaxone-AFT
CEFUROXIME AXETIL – Subsidy by endorsement			

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.

Tab 250 mg45.93

✓ Zinnat

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
• ,	2.57		✓ Zithromax
Grans for oral lig 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	14.38	15 ml	✓ Zithromax

(Apo-Azithromycin Tab 250 mg to be delisted 1 May 2022)

(Apo-Azithromycin Tab 500 mg to be delisted 1 December 2021)

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*: or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN

Tab 250 mg − Maximum of 28 tab per prescription; can be waived by Special Authority see SA1857 on the next page3.98

Grans for oral liq 250 mg per 5 ml192.00

Table 250 mg − Maximum of 28 tab per prescription; can be waived by Special Authority see SA1857 on the next page3.98

Klacid

- a) Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 on the next page
- b) Wastage claimable

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician.

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

1 Atypical mycobacterial infection; or

FRYTHROMYCIN (AS LACTORIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LITTITION (AO LAOTODIONATE)	40.00		
Inj 1 g vial	10.00	1	 Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	E-Mycin
a) Up to 20 tab available on a PSO			•
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			•
b) Wastage claimable			
, ,			
ERYTHROMYCIN STEARATE			
Tab 250 mg - Up to 30 tab available on a PSO	14.95	100	
	(22.29)		ERA
Tab 500 mg	29.90	100	
-	(44.58)		ERA

	Subsidy (Manufacturer's Price \$	e) ; Per	Fully Subsidised	Brand or Generic Manufacturer
ROXITHROMYCIN				
Tab disp 50 mg	8.29	10	✓ F	Rulide D
Restricted to children under 12 years of age.				
Tab 150 mg	8.28	50	✓ <u>I</u>	Arrow-
				Roxithromycin
Tab 300 mg	16.33	50	✓ <u>I</u>	Arrow- Roxithromycin
Penicillins				
AMOXICILLIN	00.50	500		
Cap 250 mg	22.50	500	✓ <u>I</u>	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	36.08	500	11	Alphamox
a) Up to 30 cap available on a PSO		300	• •	прпашох
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	✓	Alphamox 125
a) Up to 200 ml available on a PSO			_	
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	✓ <u>I</u>	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable Inj 250 mg vial	15.07	10	./ 1	biamox
Inj 500 mg vial		10	-	biamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		biamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO	0.89	10	✓ (Curam Duo 500/125
Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25 r			_	
per ml	5.00	100 ml	✓	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	•			_
per ml – Up to 200 ml available on a PSO	2.20 1	100 ml C)P 🗸 (Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	344.93	10	✓ E	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.09	10	✓ <u>§</u>	<u>Sandoz</u>

91

UCLOXACILLIN Cap 250 mg – Up to 30 cap available on a PSO	56.61 3.29	250 500 100 ml	1	Generic Manufacturer Staphlex Staphlex AFT
Cap 250 mg — Up to 30 cap available on a PSO	56.61 3.29	500 100 ml	1	Staphlex
Cap 500 mg — Up to 30 cap available on a PSO	56.61 3.29	500 100 ml	1	Staphlex
Cap 500 mg — Up to 30 cap available on a PSO	56.61 3.29	100 ml	1	Staphlex
Grans for oral liq 25 mg per ml	3.29		•	
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 50 mg per ml		100 ml	,	
Grans for oral liq 50 mg per ml	3.68	100 ml		
a) Up to 200 ml available on a PSOb) Wastage claimable		100 1111	•	AFT
			•	ALI
	17.56	10	1	Flucloxin
Inj 250 mg vial		10		Flucioxin
Inj 1 g vial – Up to 5 inj available on a PSO		5	_	
	5.70	5	•	Flucil
ENOXYMETHYLPENICILLIN (PENICILLIN V)	0.04	50	,	Olli I MK
Cap 250 mg – Up to 30 cap available on a PSO		50		Cilicaine VK
Cap 500 mg	6.86	50	•	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2.99	100 ml		<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	✓	<u>AFT</u>
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
OCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe - Up to 5 inj available on a PSO.	123.50	5	1	Cilicaine
etracyclines				
XYCYCLINE				
Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	1	Doxine
NOCYCLINE HYDROCHLORIDE				
Tab 50 mg – Additional subsidy by Special Authority see	F 70	00		
SA1355 below – Retail pharmacy		60		Mine teles
0	(12.05)	400		Mino-tabs
Cap 100 mg		100		Minamonata
	(52.04)			Minomycin
SA1355 Special Authority for Manufacturers Price				
tial application from any relevant practitioner. Approvals va acea.	alid without further	renewal unle	ss notifi	ed where the patient h
TRACYCLINE - Special Authority see SA1332 below - Reta	ail pharmacy			
Tab 250 mg	21.42	28	1	Accord S29
SA1332 Special Authority for Subsidy				

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ıbsidised	Generic	
\$	Per	✓	Manufacturer	

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 60

CIPROFLOXACIN

Recommended for patients with any of the following:

Tah 250 mg - Up to 5 tah available on a PSO

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

1 ab 250 mg - Op to 5 tab available on a F50	2.42	20	▼ <u>Cipilox</u>
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	✓ Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓ Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – Su Only if prescribed for dialysis or cystic fibrosis patient and the			ordingly.
Inj 150 mg	65.00	1	✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.			✓ DBL Gentamicin ection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	complicated uri	nary tract inf	ection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓ Pfizer
	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient or endorsed accordingly.	complicated uri	nary tract inf	ection and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 below - Retail p	harmacy		

2 42

28

5

✓ Cinflov

✓ Avelox

⇒SA1740 Special Authority for Subsidy

No patient co-payment payable

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*; and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or

Tab 400 mg42.00

- 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
- 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
- 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
- 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;

INFECTIONS - AGENTS FOR SYSTEMIC US	E		
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued 2 Mycobacterium avium-intracellulare complex not respond 3 Patient is under five years of age and has had close conta Note: Indications marked with * are unapproved indications. Renewal only from a respiratory specialist or infectious diseases remains appropriate and the patient is benefiting from treatment. Initial application — (Mycoplasma genitalium) only from a se sexual health specialist. Approvals valid for 1 month for application.	act with a confirmed management of specialist. Approvals xual health specialist	nulti-drug resistar valid for 1 year w or Practitioner or	nt tuberculosis case. where the treatment
All of the following: 1 Has nucleic acid amplification test (NAAT) confirmed Myc 2 Either: 2.1 Has tried and failed to clear infection using azithro 2.2 Has laboratory confirmed azithromycin resistance; 3 Treatment is only for 7 days.	mycin; or and	, ,	
Initial application — (Penetrating eye injury) only from an opt requires prophylaxis following a penetrating eye injury and treatm Note: Indications marked with * are unapproved indications. PAROMOMYCIN — Special Authority see SA1689 below — Retail	nent is for 5 days only		,
Cap 250 mg			Humatin \$29 st. Approvals valid for 1
For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microapplications meeting the following criteria: Either: 1 Patient has confirmed cryptosporidium infection; or	biologist or gastroent	erologist. Appro	vals valid for 1 month for
For the eradication of Entamoeba histolyica carriage. PYRIMETHAMINE – Special Authority see SA1328 below – Ret Tab 25 mg		30	Daraprim ©29
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valithe following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 2 For infection with coversite toxoplasmosis in patients with 110 preplacements.	r a period of 3 months		ed for applications meeting

SODIUM FUSIDATE [FUSIDIC ACID]

3 For infants with congenital toxoplasmosis until 12 months of age.

Tab 250 mg34.50

Tab 500 mg543.20

SULFADIAZINE SODIUM - Special Authority see SA1331 on the next page - Retail pharmacy

12

56

✓ Fucidin

✓ Wockhardt S29

56 dose

✓ Tobramycin BNM

✓ Trisul

(N	Subsidy fanufacturer's Price)	F Subsid	ully	Brand or Generic
,,,	\$	Per	✓	Manufacturer

⇒SA1331 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

TOBRAMYCIN

Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement	18.50	5	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient and	d the prescription is er	ndorsed accor	rdingly.
Calchian for inhalation CO man you and Fund Conhaids by			

Solution for innalation 60 mg p	er mi, 5 mi	- Subsidy by

endorsement......395.00

a) Wastage claimable

b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.

TRIMETHOPRIM

*	Tab 300 mg - Up to 30 tab available on a PSO	18.55	50	✓ TMP
---	--	-------	----	-------

TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]

*	Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – Up	
	3 -1	

to 30 tab available on a PSO......64.80 500

VANCOMYCIN - Subsidy by endorsement

Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 61
- b) For topical antifungals refer to GENITO URINARY, page 74

FLUCONAZOLE

Cap 50 mg2.7	5 28	Mylan
Cap 150 mg	5 1	✓ Mylan
Cap 200 mg12.8		✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy	4 35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's P	Price) Sub-	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOI F

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral lig 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy	141.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare \$29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
,	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Reta	il pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml		105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

continued...

following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

8/1	✓ Deolate
04	Declate
56	✓ Vttack
56	✓ Vttack
70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA16	34 on the next page – Retail pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaquine S29

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml		100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceuticals list immigration status.	ed in the Antitube	erculotics and	Antileprotics group regardless of
· ·			
CLOFAZIMINE – Retail pharmacy-Specialist			
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendati	on of an infaction	uc dicasca ph	veisian elipical microbiologist or
dermatologist.	on oi, an imedioi	us disease pri	rysician, clinical microbiologist of
* Cap 50 mg	442.00	100	✓ Lamprene S29
CYCLOSERINE - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendati	on of an infection	ie dieasea nh	veician clinical microhiologist or
respiratory physician.	on oi, an imedio	us discase pii	rysician, clinical microbiologist of
Cap 250 mg	344.00	60	✓ Cyclorin \$29
DAPSONE – Retail pharmacy-Specialist			Cyc.c
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendati	on of an infaction	uc dicasca ph	veisian elipical microbiologist or
dermatologist	on oi, an imedioi	us disease pri	rysician, clinical microbiologist of
Tab 25 mg	268 50	100	✓ Dapsone
Tab 100 mg		100	✓ Dapsone
ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialis			Бироспо
	,L		
No patient co-payment payable Prescriptions must be written by or on the recommendation.	on of an infaction	ua diagona nh	voicion, clinical microbiologist or
 b) Prescriptions must be written by, or on the recommendati respiratory physician 	on or, an intection	us disease pri	iyəlcian, ciinicai microbiologist or
Tab 100 mg	85 73	100	✓ EMB Fatol S29
Tab 400 mg		56	✓ Myambutol \$29
Tab 700 mg		50	· wyambutor

_		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
ISC	NIAZID - Retail pharmacy-Specialist				
	a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation microbiologist, dermatologist or public health physician	on of, an internal me	dicine	physician	, paediatrician, clinical
*	Tab 100 mg	23.00	100	1	PSM
ISC	NIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
	 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation microbiologist, dermatologist or public health physician 	on of, an internal me	dicine	physician	, paediatrician, clinical
*	Tab 100 mg with rifampicin 150 mg	89.82	100	1	Rifinah
*	Tab 150 mg with rifampicin 300 mg	179.13	100	/	Rifinah
PA	RA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation	on of, an infectious o	liseas	e specialis	t, clinical microbiologist or
	respiratory physician				
	Grans for oral liq 4 g sachet	280.00	30	•	Paser S29
PR	OTIONAMIDE – Retail pharmacy-Specialist				
	 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation respiratory physician 	on of, an infectious c	liseas	e specialis	t, clinical microbiologist or
	Tab 250 mg	305.00	100	1	Peteha S29
PΥ	RAZINAMIDE – Retail pharmacy-Specialist				
	 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation respiratory physician 	on of, an infectious o	liseas	e physicia	n, clinical microbiologist or
*	Tab 500 mg	59.00	100	1	AFT-Pyrazinamide
RIF	ABUTIN - Retail pharmacy-Specialist				
	a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation gastroenterologist	on of, an infectious c	liseas	e physicia	n, respiratory physician or
*	Cap 150 mg	299.75	30	1	Mycobutin
RIF	AMPICIN - Subsidy by endorsement				
	a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection i antimicrobial based on susceptibilities and the prescription Retail pharmacy - Specialist. Specialist must be an intern	n is endorsed accord	lingly;	can be wa	aived by endorsement -
*	paediatrician, or public health physician. Cap 150 mg	58 54	100	1	Rifadin
	Cap 300 mg		100		Rifadin
	Oral liq 100 mg per 5 ml		60 m	✓	Rifadin
Α	ntivirals				
For	eye preparations refer to Eye Preparations, Anti-Infective Prep	parations, page 229			
	epatitis B Treatment				
	TECAVIR Tab 0.5 mg	52.00	30	/	Entecavir Sandoz

	Subsidy (Manufacturer's Prio \$	ce) S	Fully Subsidised	Brand or Generic Manufacturer	
LAMIVUDINE - Special Authority see SA1685 below - Retail ph	narmacy				
Tab 100 mg	6.95	28	✓ Z	etlam	
Oral liq 5 mg per ml	270.00	240 ml O	P ✓Z	effix	

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXII

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651... page 103

Herpesvirus Treatments

ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ Lovir
* Tab dispersible 400 mg		56	✓ Lovir
* Tab dispersible 800 mg	5.98	35	✓ Lovir
VALACICLOVIR			
Tab 500 mg	6.50	30	✓ Vaclovir
Tab 1,000 mg		30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1993 be	low - Retail pharmacy		
Tab 450 mg		60	✓ Valganciclovir Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

Subsidy (Manufacturer's Price) Fully Subsidised Per Brand or Generic Manufacturer

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

below
Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is
co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated
accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 103 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

30

✓ Teva

⇒SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

those risks; and

- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)		ubsidised	Generic
	\$	Per		Manufacturer

continued...

- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA1651 on the previous	s page – Retail phari	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	s page – Retail pha	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	ıs page – Retail pha	rmacy	
Tab 200 mg	84.00	60	Nevirapine Alphapharm
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE - Special Authority see SA1651 on th	e previous page – i	Retali pharmac	cy
Tab 300 mg	180.00	60	✓ Ziagen
Oral liq 20 mg per ml	256.31	240 ml OP	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE - Special Author			
Note: abacavir with lamivudine (combination tablets) coun	ts as two anti-retro	viral medication	ns for the purposes of the
anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa

	Subsidy		Fully	Brand or
	(Manufacturer's		,	Generic
	\$	Per	✓	Manufacturer
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPI	BOYII - Specia	al Authority soo S	Δ1651	on nage 103 – Retail
	HONIL - Specia	al Authority See S	AIOSI	on page 105 - netali
Note: Efavirenz with emtricitabine and tenofovir disoproxil c	ounts as three a	ınti-retroviral med	lications	s for the purposes of the
anti-retroviral Special Authority				
Tab 600 mg with emtricitabine 200 mg and tenofovir disopro 245 mg (300 mg as a maleate)		30	✓ <u>M</u> y	<u>ylan</u>
EMTRICITABINE - Special Authority see SA1651 on page 103	– Retail pharma	CV		
Cap 200 mg	307.20	30	✓ <u>En</u>	ntriva
LAMIVUDINE - Special Authority see SA1651 on page 103 - Re	etail pharmacy			
Tab 150 mg	84.50	60		<u>mivudine</u> Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3T	
ZIDOVUDINE [AZT] - Special Authority see SA1651 on page 10	03 – Retail phari	macy		
Cap 100 mg		100	✓ Re	etrovir
Oral liq 10 mg per ml	30.45	200 ml OP	✓ Re	etrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority ser Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.				
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ Al	phapharm
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA1651 on p	age 103 – Reta	il pharmacy		
Cap 150 mg	•	60	✓ Te	va
Cap 200 mg		60	✓ Te	eva
DARUNAVIR - Special Authority see SA1651 on page 103 - Re				
Tab 400 mg		60	✓ Da	arunavir Mylan
Tab 600 mg		60		arunavir Mylan
· ·			· <u>D</u>	a unavii iiiyiun
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651			/ V-	.l
Tab 100 mg with ritonavir 25 mg		60	✓ Ka	
Tab 200 mg with ritonavir 50 mg		120	✓ Ka	
Oral liq 80 mg with ritonavir 20 mg per ml		300 ml OP	✓ Ka	netra
RITONAVIR - Special Authority see SA1651 on page 103 - Ret	, ,			
Tab 100 mg	43.31	30	✓ <u>No</u>	<u>orvir</u>
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 103 Tab 50 mg		acy 30	✓ Tiv	vicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 c Tab 400 mg Tab 600 mg	1,090.00	etail pharmacy 60 60		entress entress HD

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test: or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4. ✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV: or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*: or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1 Patient has a myeloproliferative disorder*; and
- 2.2 Patient is intolerant of hydroxyurea; and
- 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE * Tab 1 g40.01	100	✓ Hiprex
NITROFURANTOIN		·
* Tab 50 mg - Up to 30 tab available on a PSO	100	✓ Nifuran
* Tab 100 mg	100	✓ Nifuran
* Cap modified-release 100 mg - Wastage claimable86.40	100	✓ Macrobid
NORFLOXACIN		
Tab 400 mg — Subsidy by endorsement 135.00) 100	✓ Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's F		sidised	Generic
	\$	Per		Manufacturer
Anticholinesterases				
Antichonnesterases				
EOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	19.60	10	✓ J	uno S29
	29.40		✓ M	lax Health
	98.00	50	✓ A	straZeneca
RIDOSTIGMINE BROMIDE				
Tab 60 mg	45.79	100	✓ M	lestinon
lan Charaidal Anti Inflammatam Duran				
Non-Steroidal Anti-Inflammatory Drugs				
CLOFENAC SODIUM				
Tab EC 25 mg		50	✓ D	iclofenac Sandoz
Tab 50 mg dispersible	1.50	20	√ V	oltaren D
Tab EC 50 mg		50	_	iclofenac Sandoz
Tab long-acting 75 mg		500		po-Diclo SR
Tab long-acting 100 mg		500		po-Diclo SR
Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a P		5		oltaren
Suppos 12.5 mg		10		oltaren
Suppos 25 mg		10	_	oltaren
Suppos 50 mg — Up to 10 supp available on a PSOSuppos 100 mg		10 10		oltaren oltaren
Suppos 100 mgoo-Diclo SR Tab long-acting 100 mg to be delisted 1 May 2022		10	• •	Ollarell
)			
JPROFEN	04.40	4 000		
Tab 200 mg		1,000		<u>elieve</u> rufen SR
Tab long-acting 800 mg	5.99	30		rulen SK Juprofen SR BNM
Oral liq 20 mg per ml		200 ml		thics
uprofen SR BNM Tab long-acting 800 mg to be delisted 1 Janu		200 1111	٠ ـ	unos
TOPROFEN	ally 2022)			
Cap long-acting 200 mg	12.07	28	√ ∩	ruvail SR
	12.07	20	• 0	iuvali Sh
FENAMIC ACID	4.05	50		
Cap 250 mg	4	50	р	anatan
	(9.16) 0.50	20	Р	onstan
	(5.60)	20	Р	onstan
DDOVEN	(5.55)		ı	onotari
PROXEN Tab 250 mg	32.60	500	⊿ M	oflam 250
Tab 250 mg Tab 500 mg		500 250		oflam 250 oflam 500
Tab long-acting 750 mg		28	_	aprosyn SR 750
Tab long-acting 1 g		28		aprosyn SR 1000
LINDAC		_0		
	0.57	FC	./ 14	lylon con
Tab 100 mg		56 50	✓ IVI	lylan (\$29) clin
Tab 200 mg		50 56		
alia Tah 200 man ta ha daliatad 4 January 2000)	16.91	56	• 5	ulindac Mylan S29
clin Tab 200 mg to be delisted 1 January 2022)	,			
ulindac Mylan 👀 Tab 200 mg to be delisted 1 January 2022)			
NOXICAM			_	
Tab 20 mg		100		ilcotil
Inj 20 mg vial	9.95	1	✓ A	FT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
NSAIDs Other				
CELECOXIB Cap 100 mg Cap 200 mg		60 30	1	Celecoxib Pfizer Celebrex Celecoxib Pfizer
Topical Products for Joint and Muscular Pain				
CAPSAICIN Crm 0.025% - Special Authority see SA1289 below - Retail pharmacy	9.75 4	5 g C	np 🗸	Zostrix
➤SA1289 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid osteoarthritis that is not responsive to paracetamol and oral non-substitution.				
Antirheumatoid Agents HYDROXYCHLOROQUINE – Subsidy by endorsement Subsidised only if prescribed for rheumatoid arthritis, systemi				
suppression, relevant dermatological conditions (cutaneous for mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmon Pharmacists may annotate the prescription as endorsed when hydroxychloroguine. Note: Indication marked with a * is an u	nary)*, and the presc re there exists a reco	riptio rd of	n is endor	sed accordingly.
* Tab 200 mg		100	✓	Plaquenil
LEFLUNOMIDE Tab 10 mg Tab 20 mg		30 30		<u>Arava</u> <u>Arava</u>
PENICILLAMINE Tab 125 mg Tab 250 mg		100 100		D-Penamine D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

* Tab 70 mg	2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			
* Tab 70 mg with colecalciferol 5 600 iu	1 51	1	✓ Fosamay Plus

Other Treatments

ALENDRONATE SODIUM

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy	Fı	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	•	Manufacturer

continued...

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	27.53	1	Pamisol
Inj 6 mg per ml, 10 ml vial		1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	75.95	1	Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see \$	SA1779 on the next page	- Retai	l pharmacy
* Tab 60 mg	53.76	28	✓ Evista

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or egual to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has guantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg3.10 Risedronate Sandoz TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy Forteo

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily;

(Ma	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
	\$	Per	1	Manufacturer

continued...

zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain: or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically: or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

Subsidy (Manufacturer's Price)	Ş	Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg		500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA19	963 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	✓ Benzbromaron AL
			100 S29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

(Adenuric Tab 80 mg to be delisted 1 January 2022) (Adenuric Tab 120 mg to be delisted 1 January 2022)

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

	Muscle Relaxants				
Ī	BACLOFEN				
+	* Tab 10 mg	4.20	100	✓ Pacifen	
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsem	ent11.55	1	Lioresal Intrathecal	
	Subsidised only for use in a programmable pump in p caused intolerable side effects and the prescription is			ents have been ineffective or ha	ıve
	Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	306.82	5	✓ Medsurge	
	Subsidised only for use in a programmable pump in p caused intolerable side effects and the prescription is			ents have been ineffective or ha	ive
I	DANTROLENE				
	Cap 25 mg	97.50	100	✓ Dantrium	
				✓ Dantrium S29 S29	
	Cap 50 mg	77.00	100	✓ Dantrium	

ORPHENADRINE CITRATE

100

✓ Norflex

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE A Cap 100 mg	.38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			•
▲ Inj 10 mg per ml, 2 ml ampoule	.59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo

BROMOCRIPTINE MESYLATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of bromocriptine mesylate.

* Tab 2.5 mg	11.70	30	Parlodel S29
•	32.08	100	Apo-Bromocriptine
ENTACAPONE			
▲ Tab 200 mg	22.00	100	Entapone
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg	13.75	100	Madopar 62.5
* Cap 100 mg with benserazide 25 mg	15.80		Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg	22.85	100	Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25		Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg		100	Sinemet CR
* Tab 250 mg with carbidopa 25 mg		100	Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	6.12	100	Ramipex
▲ Tab 1 mg	20.73	100	Ramipex
RASAGILINE			
* Tab 1 mg	53.50	30	Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.85	84	Ropin
· ·		100	Mylan S29
▲ Tab 1 mg	3.95		Ropin
·		100	Mylan S29
▲ Tab 2 mg	5.48		Ropin
▲ Tab 5 mg			Ropin
-			

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

*	Tab 5 mg22.0	00 10) 🗸	Apo-Selegiline S29 S29
ТО	LCAPONE			
\blacktriangle	Tab 100 mg	8 10	0	Tasmar

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Anticholinergics				
BENZATROPINE MESYLATE Tab 2 mg		60 5		Benztrop <u>Phebra</u>
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ed Disorders			
RILUZOLE - Special Authority see SA1403 below - Retail pharm Wastage claimable Tab 50 mg	-	56	✓	Rilutek
Initial application only from a neurologist or respiratory specialist following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2 The patient has at least 60 percent of predicted forced vita 3 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. Renewal from any relevant practitioner. Approvals valid for 18 m All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and	duration of 5 years o	or less onths	s; and prior to th	e initial application; and
3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. TETRABENAZINE Tab 25 mg	91.10	112	•	<u>Motetis</u>
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube — Subsidy by endorsement	dministration and the	10	cription is	Instillagel Lido

b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed

accordingly.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) Sub	osidised	Generic
	\$	Per		Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	✓ N	lucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	✓ L	idocaine-Baxter
			√ L	idocaine-Claris
	17.50	50		
	(35.00)		Х	ylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25 [°]	25	√ L	idocaine-Baxter
			√ L	idocaine-Claris
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)		Х	ylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20 [°]	5	√ L	idocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO		5	√ L	idocaine-Baxter
• • •			√ L	idocaine-Claris
(Lidocaine-Claris Inj 1%, 5 ml ampoule to be delisted 1 January 2	2022)			
(Lidocaine-Claris Inj 2%, 5 ml ampoule to be delisted 1 January 2	,			
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE	- /			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –		40		e
Subsidy by endorsement	103.32	10	✓ P	fizer
a) Up to 5 each available on a PSO				
 b) Subsidised only if prescribed for urethral or cervical a 	administration and th	ne prescrip	otion is er	ndorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA0906 above - Retail pharmacy

Crm 4%	5.40 27.00	5 g OP 30 g OP	✓ LMX4 ✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA090	6 above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 109

Non-opioid Analgesics			
ASPIRIN			4- 00 - 00
★ Tab dispersible 300 mg - Up to 30 tab availab	ole on a PSO4.50	100	Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic accordingly.	neuralgia or diabetic periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream \$29
NEFOPAM HYDROCHLORIDE			
Tah 30 mg	23.40	an	✓ Acunan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PARACETAMOL				
Tab 500 mg - blister pack	0.50	20	1	Medco
• ,			✓	Pharmacy Health
	1.12		✓	Ethics Paracetamol
				Classic
	2.48	100	✓	Pharmacy Health
	5.01	50	✓	Panadol
	11.75	96	✓	Panadol Mini Caps
	19.75	1,000	/	Pacimol
	24.82		1	Paracetamol
				Pharmacare
			•	Pharmacare

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
 regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
 annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

*	Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓ <u>Paracare</u>
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double Strength
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3.29	10	✓ Gacet
*	Suppos 250 mg	3.79	10	✓ Gacet
*	Suppos 500 mg	12.40	50	✓ Gacet
(Me	edco Tab 500 mg - blister pack to be delisted 1 February 2022)			
(Ph	armacy Health Tab 500 mg - blister pack to be delisted 1 February 2	2022)		
(Etl	nics Paracetamol Classic Tab 500 mg - blister pack to be delisted 1	February 2	2022)	
(Ph	armacy Health Tab 500 mg - blister pack to be delisted 1 February 2	2022)		
(Pa	nadol Mini Caps Tab 500 mg - blister pack to be delisted 1 February	y 2022)		
(Pa	racetamol Pharmacare Tab 500 mg - blister pack to be delisted 1 Fo	ebruary 20.	22)	
(Pa	racetamol Pharmacare Tab 500 mg - bottle pack to be delisted 1 De	ecember 20	021)	
	· ·			

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing fre	quency		
Tab 15 mg	6.25	100	√ <u>F</u>	<u>PSM</u>
Tab 30 mg		100	✓ <u>F</u>	
Tab 60 mg	14.25	100	✓ <u>F</u>	<u>PSM</u>
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	✓ [OHC Continus
FENTANYL				
 a) Only on a controlled drug form 				
b) No patient co-payment payable				
Safety medicine; prescriber may determine dispensing fr Inj 50 mcg per ml, 2 ml ampoule		10	./ 5	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
Patch 12.5 mcg per hour		5		entanyl Sandoz
Patch 25 mcg per hour		5		entanyi Sandoz
Patch 50 mcg per hour		5		entanyl Sandoz
Patch 75 mcg per hour	17.99	5	√ F	entanyl Sandoz
Patch 100 mcg per hour	18.59	5	√ F	entanyl Sandoz
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr				
d) Extemporaneously compounded methadone will only be	reimbursed at the rate	e of the ch	neapest	form available
(methadone powder, not methadone tablets).e) For methadone hydrochloride oral liquid refer Standard F	Formulae page 227			
Tab 5 mg		10	✓ 1	/lethatabs
Oral lig 2 mg per ml		200 ml		Biodone
Oral liq 5 mg per ml		200 ml	_	Biodone Forte
Oral liq 10 mg per ml		200 ml	√ E	Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	✓ Þ	AFT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing fr 				
Oral liq 1 mg per ml		200 ml		RA-Morph
Oral liq 2 mg per ml		200 ml		RA-Morph
Oral liq 5 mg per ml	19.44	200 ml		Ordine S29
Oral lia 10 ma nor ml	07.74	200 ml		RA-Morph
Oral liq 10 mg per ml	27.74	200 ml		Ordine S29

✓ RA-Morph

	Subsidy (Manufacturer's Price)	S	Fully	Brand or Generic
	\$	Per	√	Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f	requency			
Tab immediate-release 10 mg	2.80	10	1	Sevredol
Tab immediate-release 20 mg	5.52	10		Sevredol
Cap long-acting 10 mg	2.05	10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg	6.12	10	✓	m-Eslon
Cap long-acting 100 mg		10		m-Eslon
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	SO6.99	5	✓	DBL Morphine
				Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO5.61	5	✓	DBL Morphine
				Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO7.08	5	1	DBL Morphine
,g p, ppp				Sulphate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 7.28	5	1	DBL Morphine
inj oo nig per nii, 1 nii ampoule — op to 5 inj available on a	1 00 7.20	J	•	Sulphate
OVIVOODONE LIVEDOOLII ODIDE				ouipilate
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 c) Safety medicine; prescriber may determine dispensing f 			_	
Tab controlled-release 5 mg		20		Oxycodone Sandoz
	3.01	28	•	Oxycodone Sandoz
				S29 S29
Tab controlled-release 10 mg	2.15	20	1	Oxycodone Sandoz
	3.23	30	1	Oxycodone Sandoz
				S29 S29
	5.38	50	1	Oxycodone Sandoz
				S29 S29
	10.75	100	1	Oxycodone Sandoz
	10.75	100	•	•
				S29 S29
	11.50	28		OxyContin
Tab controlled-release 20 mg		20		Oxycodone Sandoz
	5.38	50	•	Oxycodone Sandoz
				S29 S29
	10.75	100	1	Oxycodone Sandoz
				S29 S29
	13.25	28	1	OxyContin
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 20 mg		20		OxyNorm
Oral lig 5 mg per 5 ml		250 ml		OxyNorm
OxyNorm to be Principal Supply on 1 September 2021	-	- "		•
Inj 10 mg per ml, 1 ml ampoule	7.28	5	1	OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm
Inj 50 mg per ml, 1 ml ampoule		5		OxyNorm
)a L		-		- ,

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
PARACETAMOL WITH CODEINE – Safety medicine; prescriber * Tab paracetamol 500 mg with codeine phosphate 8 mg PETHIDINE HYDROCHLORIDE		ensing fred 1,000		Paracetamol + Codeine (Relieve)
a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab 50 mg		10	✓	PSM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a P	SO 29.88	5	1	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO30.72	5	•	DBL Pethidine Hydrochloride
TRAMADOL HYDROCHLORIDE Tab sustained-release 100 mg	1.52	20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.80	100	•	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg		100		Arrow-Amitriptyline
Tab 50 mg		100		Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescri				
Tab 10 mg		100		Apo-Clomipramine
Tab 25 mg		100	•	Apo-Clomipramine
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by en				
Safety medicine; prescriber may determine dispensing fre				
 Subsidy by endorsement – Subsidised for patients who w 2019 and the prescription is endorsed accordingly. Pharr exists a record of prior dispensing of dosulepin [dothiepin] 	nacists may annotate			
Tab 75 mg	•	30	1	Dosulepin Mylan
Cap 25 mg		50		Dosulepin, Ida
				Mylan S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine disne	neina froa	IIAncv	•
Tab 10 mg		50		Tofranil
	10.96	100		Tofranil
Tab 25 mg		50		Tofranil
NORTRIPTYLINE HYDROCHLORIDE - Safety medicine; prescr		dispensing	frequ	encv
Tab 10 mg	,	100		Norpress
Tab 25 mg		180		Norpress
-				- —

	Subsidy (Manufacturer's Price)	Cuba	Fully	Brand or Generic
	(Manufacturer's Frice)	Per	luiseu ✓	Manufacturer
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
TRANYLCYPROMINE SULPHATE Tab 10 mg	12.85 22.94 45.88 96.00	28 50 100	✓ P	arnate S29 S29 arnate arnate S29 S29 arnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE * Tab 150 mg * Tab 300 mg		60 60		urorix urorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE * Tab 20 mg ESCITALOPRAM	1.52	84	✓ P	SM Citalopram
* Tab 10 mg	1.07	28	√ E	scitalopram (Ethics)
	1.40		✓ E	scitalopram- Apotex
Escitalopram (Ethics) to be Sole Supply on 1 October 2 * Tab 20 mg	1.92	28		scitalopram (Ethics)
	2.49		✓ E	scitalopram- Apotex
Escitalopram (Ethics) to be Sole Supply on 1 October 2 (Escitalopram-Apotex Tab 10 mg to be delisted 1 October 2021, (Escitalopram-Apotex Tab 20 mg to be delisted 1 October 2021))			
FLUOXETINE HYDROCHLORIDE * Tab dispersible 20 mg, scored – Subsidy by endorsement. Subsidised by endorsement	1.98	30	✓ F	luox
When prescribed for a patient who cannot swallow accordingly; or	v whole tablets or caps	sules and t	he pres	scription is endorsed
When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi				
Cap 20 mg	2.91	84	✓ F	luox
PAROXETINE * Tab 20 mg		30		axtine
(Paxtine Tab 20 mg to be delisted 1 January 2022)	3.61	90	✓ <u>L</u>	oxamine

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
SERTRALINE				
* Tab 50 mg	0.92	30		Setrona
	3.05	90		Setrona AU Arrow-Sertraline
* Tab 100 mg		30		Setrona
- 140 100 mg		00		Setrona AU
	5.25	90	1	Arrow-Sertraline
(Setrona AU Tab 50 mg to be delisted 1 October 2021) (Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021) (Setrona AU Tab 100 mg to be delisted 1 October 2021) (Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)				
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg	2.60	28	1	Noumed S29
v	2.63	30	/	Apo-Mirtazapine
Tab 45 mg		28		Noumed S29
(A AK)	3.48	30	•	Apo-Mirtazapine
(Apo-Mirtazapine Tab 30 mg to be delisted 1 January 2022) (Apo-Mirtazapine Tab 45 mg to be delisted 1 January 2022)				
VENLAFAXINE			_	
* Cap 37.5 mg * Cap 75 mg		84 84		Enlafax XR Enlafax XR
* Cap 75 mg*		84		Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM - Safety medicine; prescriber may determine disp	pensing frequency			
Inj 1 mg per ml, 1 ml		5	✓	Rivotril
(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)				
DIAZEPAM - Safety medicine; prescriber may determine dispens		_		
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement	23.66	5	•	Hospira
a) Up to 5 inj available on a PSOb) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic procedure	es".			
Rectal tubes 5 mg – Up to 5 tube available on a PSO		5	1	Stesolid
PARALDEHYDE				
* Inj 5 ml	1,500.00	5	•	AFT S29
PHENYTOIN SODIUM				
* Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a P	SO88.63	5	•	Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a		_	_	
PSO	133.92	5	/	Hospira

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		(Manufacturer's Fir	Per	✓ Manufactu	ıror
_		Ψ	r ei	- Manufactu	II CI
Co	ontrol of Epilepsy				
CAF	RBAMAZEPINE				
*	Tab 200 mg	14.53	100	✓ Tegretol	
*	Tab long-acting 200 mg	16.98	100	✓ Tegretol CR	}
*	Tab 400 mg	34.58	100	✓ Tegretol	
*	Tab long-acting 400 mg	39.17	100	✓ Tegretol CR	}
*	Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol	
CLC	DBAZAM - Safety medicine; prescriber may determine disper	nsina freauency			
	Tab 10 mg	. ,	50	✓ Frisium	
CLC	DNAZEPAM - Safety medicine; prescriber may determine dis		v		
OLC	Oral drops 2.5 mg per ml		10 ml OP	✓ Rivotril	
СТІ			10 1111 01	- 111701111	
EIF	IOSUXIMIDE	140.00	400	. Zavantin	
	Cap 250 mg		100	✓ Zarontin	
	Oral liq 250 mg per 5 ml	50.35	200 ml	Zarontin	
GAE	BAPENTIN				
	Note: Not subsidised in combination with subsidised pregab				
	Cap 100 mg		100	✓ Apo-Gabap	
	Cap 300 mg		100	✓ Apo-Gabap	
*	Cap 400 mg	5.64	100	Apo-Gabap	entin
LAC	COSAMIDE - Special Authority see SA1125 below - Retail pl	narmacy			
\blacktriangle	Tab 50 mg	25.04	14	✓ Vimpat	
\blacktriangle	Tab 100 mg		14	✓ Vimpat	
		200.24	56	✓ Vimpat	

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Patient has partial-onset epilepsy; and

2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

300.40

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

WOTHIGHT			
Tab dispersible 2 mg	55.00	30	✓ Lamictal
		30	✓ Lamictal
		56	✓ Logem
		56	✓ Logem
		56	✓ Logem
	Tab dispersible 5 mg	Tab dispersible 2 mg .55.00 Tab dispersible 5 mg .50.00 Tab dispersible 25 mg .2.76 Tab dispersible 50 mg .3.31 Tab dispersible 100 mg .4.40	Tab dispersible 2 mg 55.00 30 Tab dispersible 5 mg 50.00 30 Tab dispersible 25 mg 2.76 56 Tab dispersible 50 mg 3.31 56

14

56

✓ Vimpat
✓ Vimpat

✓ Vimpat

	Subsidy		Fully	Brand or
	(Manufacturer's Price		sidised	Generic
	\$	Per		Manufacturer
LEVETIRACETAM				
Tab 250 mg	4.99	60	✓ E	Everet
Tab 500 mg	8.79	60	✓ E	veret
Tab 750 mg	14.39	60	✓ E	veret
Tab 1,000 mg	18.59	60	✓ E	veret
Oral liq 100 mg per ml	44.78	300 ml OP	✓ L	evetiracetam-AFT
PHENOBARBITONE				
For phenobarbitone oral liquid refer Standard Formulae, pa	ge 237			
* Tab 15 mg	•	500	✓ F	PSM
* Tab 30 mg		500	✓ F	PSM
PHENYTOIN SODIUM				
* Tab 50 mg	75.00	200	√ [Dilantin Infatab
Cap 30 mg		200		Dilantin
Cap 100 mg		200	_	Dilantin
* Oral lig 30 mg per 5 ml		500 ml	_	Dilantin
		000 1111	•	Jiidiidii
PREGABALIN	a antin			
Note: Not subsidised in combination with subsidised gabat * Cap 25 mg		56	./ [Pregabalin Pfizer
* Cap 25 mg*		56 56		Pregabalin Pfizer
* Сар 75 mg		56 56		rregabalili Pilzer .vrica
ж Сар 130 mg	4.01	50		Pregabalin Pfizer
* Cap 300 mg	7 38	56		Pregabalin Pfizer
	7.50	30	• 1	regaballii Filzei
PRIMIDONE				
* Tab 250 mg	17.25	100	•	Apo-Primidone
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓ E	Epilim Crushable
Tab 200 mg EC	27.44	100	✓ E	Epilim
Tab 500 mg EC	52.24	100	✓ E	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 ml	✓ E	Epilim S/F Liquid
				Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓ E	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail	pharmacy			
Cap 250 mg		60	√ [Diacomit S29
		00	✓ [

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

(Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
TOPIRAMATE				
▲ Tab 25 mg	11.07	60	_	Arrow-Topiramate
	00.04			Topiramate Actavis
	26.04			Topamax
▲ Tab 50 mg	18.81	60	_	Arrow-Topiramate
				Topiramate Actavis
	44.26		•	Topamax
▲ Tab 100 mg	31.99	60		Arrow-Topiramate
			1	Topiramate Actavis
	75.25		✓.	Topamax
▲ Tab 200 mg	55.19	60	1	Arrow-Topiramate
-			✓.	Topiramate Actavis
	129.85		1	Topamax
Sprinkle cap 15 mg	20.84	60	1	Topamax
Sprinkle cap 25 mg		60	1	Topamax
/IGABATRIN - Special Authority see SA1997 below - Retail phar				-
▲ Tab 500 mg		100	✓ :	Sabril

⇒SA1997 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Fither:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - of treatment with vigabatrin; or

 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...
- Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Por	/	Manufacturer	

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 109

Acute Migraine Treatment		
RIZATRIPTAN		
Tab orodispersible 10 mg3.65	30	✓ Rizamelt
SUMATRIPTAN		
Tab 50 mg24.44	100	✓ Apo-Sumatriptan
Tab 100 mg46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per		
prescription34.00	2 OP	✓ <u>Imigran</u>

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

PIZOTIFFN

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT − Special Authority see SA0987 below − Retail pharmacy
Cap 2 × 80 mg and 1 × 125 mg......30.00 3 OP

✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

# Tab 16 mg3.88	84	✓ Vergo 16
CYCLIZINE HYDROCHLORIDE Tab 50 mg0.49	10	✓ Nausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml21.53	10	✓ Hameln
DOMPERIDONE * Tab 10 mg2.25	100	✓ Pharmacy Health
HYOSCINE HYDROBROMIDE		
* Inj 400 mcg per ml, 1 ml ampoule	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1998 below - Retail pharmacy14.11	2	✓ Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
 ` \$	Per	✓	Manufacturer

continued...

where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or

2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

ME	TOCLOPRAMIDE HYDROCHLORIDE		
*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50	10	✓ <u>Pfizer</u>
ON	DANSETRON		
*	Tab 4 mg	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO	10	Ondansetron ODT-DRLA
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	Ondansetron ODT-DRLA
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal	50	_
	(30.00)		Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dispensing frequency	y	
Tab 100 mg5.15	30	✓ Sulprix
17.16	100	✓ Amisulpride
		Mylan S29
Tab 200 mg14.96	60	✓ Sulprix
Tab 400 mg29.78	60	✓ Sulprix
ARIPIPRAZOLE - Safety medicine; prescriber may determine dispensing frequency	су	
Tab 5 mg17.50	30	Aripiprazole Sandoz
Tab 10 mg 17.50	30	✓ Aripiprazole Sandoz
Tab 15 mg 17.50	30	✓ Aripiprazole Sandoz
Tab 20 mg17.50	30	Aripiprazole Sandoz
Tab 30 mg17.50	30	✓ Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; prescriber may deter	rmine disper	nsing frequency
Tab 10 mg - Up to 30 tab available on a PSO14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO30.79	10	✓ Largactil

	Subsidy		Fully	Brand or
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
LOZAPINE – Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing freq	uency			
Tab 25 mg	5.69	50		Clozaril
	6.69		✓	Clopine
	11.36	100	✓	Clozaril
	13.37		✓	Clopine
Tab 50 mg	8.67	50	✓	Clopine
	17.33	100	✓	Clopine
Tab 100 mg	14.73	50	✓	Clozaril
	17.33		✓	Clopine
	29.45	100		Clozaril
	34.65			Clopine
Tab 200 mg		50		Clopine
1 ab 200 mg	69.30	100		Clopine
Suspension 50 mg per ml		100 m		Clopine
ouspension so mg per mi	67.62	100 111		Versacloz
			•	VEISACIOZ
ALOPERIDOL – Safety medicine; prescriber may determine of			_	_
Tab 500 mcg - Up to 30 tab available on a PSO		100	/	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO	9.43	100	✓	<u>Serenace</u>
Tab 5 mg - Up to 30 tab available on a PSO	14.86	50		<u>Serenace</u>
	29.72	100	✓	<u>Serenace</u>
Oral liq 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a F		10	✓	Serenace
VOMEPROMAZINE - Safety medicine; prescriber may dete		aduanav		
		100		Nozinan (Swiss)
Tab 25 mg (33.8 mg as a maleate)				
Tab 25 mg as a maleate		100		Nozinan (Curios)
Tab 100 mg (135 mg as a maleate)		100	•	Nozinan (Swiss)
Tab 100 mg as a maleate	41./5	100	•	<u>Nozinan</u>
VOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may dete	ermine di	ispensing	frequency
Inj 25 mg per ml, 1 ml ampoule	33.50	10	1	Nozinan
THIUM CARBONATE - Safety medicine; prescriber may dete		ranuancy	,	
Tab long-acting 400 mg		100		Priadel
Priadel to be Principal Supply on 1 September 2021	72.00	100	•	i ilaaci
	0.40	100	./	Davelas
Cap 250 mg		100	•	Douglas
_ANZAPINE - Safety medicine; prescriber may determine dis			_	
		00	/	Zypine
Tab 2.5 mg	1.35	28		
Tab 2.5 mg Tab 5 mg		28	_	Zypine
•	1.58		1	Zypine ODT
Tab 5 mg	1.58 1.81	28	✓	
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg	1.58 1.81 2.01	28 28	√ √	Zypine ODT
Tab 5 mg	1.58 1.81 2.01 2.38	28 28 28 28	√ √	Zypine ODT Zypine
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine d	1.58 1.81 2.01 2.38 ispensing frequency	28 28 28 28	<i>y y y</i>	Zypine ODT Zypine Zypine ODT
Tab 5 mg		28 28 28 28 28	\ \ \	Zypine ODT Zypine Zypine ODT Neulactil
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine d Tab 2.5 mg		28 28 28 28 28 , 84 100	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine d		28 28 28 28 28 4 100 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE – Safety medicine; prescriber may determine d Tab 2.5 mg		28 28 28 28 28 , 84 100	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE — Safety medicine; prescriber may determine d Tab 2.5 mg Tab 10 mg		28 28 28 28 28 4 100 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil
Tab 5 mg Tab orodispersible 5 mg Tab 10 mg Tab orodispersible 10 mg ERICYAZINE — Safety medicine; prescriber may determine d Tab 2.5 mg Tab 10 mg JETIAPINE — Safety medicine; prescriber may determine dis		28 28 28 28 28 4 100 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil Neulactil
Tab 5 mg		28 28 28 28 28 , , , , , , , , , , , , ,	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil Neulactil Neulactil
Tab 5 mg		28 28 28 28 28 4 100 84 100		Zypine ODT Zypine Zypine ODT Neulactil Neulactil Neulactil Neulactil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
RISPERIDONE – Safety medicine; prescriber may determine dis	spensing frequency			
Tab 0.5 mg	1.86	60	✓	Risperidone (Teva)
Tab 1 mg	2.06	60	✓	Risperidone (Teva)
Tab 2 mg	2.29	60	✓	Risperidone (Teva)
Tab 3 mg	2.50	60	✓	Risperidone (Teva)
Tab 4 mg	3.42	60	✓	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 m	✓	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine dis	spensing frequency			
Cap 20 mg	14.50	60	✓	Zusdone
Cap 40 mg	24.70	60	✓	Zusdone
Cap 60 mg	33.80	60	✓	Zusdone
Cap 80 mg	39.70	60	1	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre	scriber may determin	e disp	ensing fre	equency
Tab 10 mg		100		Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO13.14	5	Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO20.90	5	Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO40.87	5	✓ Fluanxol

HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO55.90	5	✓ Haldol Concentrate
		✓ Haldol

Decanoas \$29

OLANZAPINE - Special Authority see SA1428 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 210 mg vial	252.00	1	Zyprexa Relpre	٧V
Inj 300 mg vial	414.00	1	✓ Zyprexa Relpre	٧v
Inj 405 mg vial	504.00	1	✓ Zyprexa Relpre	٧V

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	0 , ,	1	✓ Invega Sustenna
Inj 50 mg syringe		1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

(M	Subsidy fanufacturer's Price)	Subsi	Fully dised	Brand or Generic
,	\$	Per	✓	Manufacturer

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency	uency		
Inj 25 mg vial	135.98	1	 Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

Ini 200 ma nor ml 1 ml	 Un to 5 ini available on a P 	PSO 19.80	5 / CI	loxido

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may de	termine dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determ	nine dispensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
•			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
LORAZEPAM – Safety medicine; prescriber may determine dispe Tab 1 mg Tab 2.5 mg	9.72	250 100		Ativan Ativan
OXAZEPAM – Safety medicine; prescriber may determine disper Tab 10 mg Tab 15 mg	6.17	100 100		Ox-Pam Ox-Pam

Multiple Sclerosis Treatments

⇒SA2051 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. **Renewal — (Multiple sclerosis)** only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

	0.1.11		
	Subsidy (Manufacturer's Price)		fully Brand or sed Generic
	(Manufacturer's Price)	Per	✓ Manufacturer
DIMETLINI FLIMADATE Chariel Authority and CARREL on the	nraviava naga. Data	il nharmaar	
DIMETHYL FUMARATE – Special Authority see SA2051 on the	previous page – Reia	ш рпаппасу	
a) Wastage claimable	ia traatmanta aimultan	aauahi ia na	t narmittad
b) Note: Treatment on two or more funded multiple scleros Cap 120 mg		14	√ Tecfidera
Cap 240 mg		56	✓ Tecfidera
		• •	reciliaera
FINGOLIMOD – Special Authority see SA2051 on the previous	page – Retail pharmac	СУ	
a) Wastage claimable	to the other suite of an item.		A construction of
b) Note: Treatment on two or more funded multiple scleros		•	•
Cap 0.5 mg	•	28	✓ Gilenya
GLATIRAMER ACETATE – Special Authority see SA2051 on th			
Note: Treatment on two or more funded multiple sclerosis tr		, ,	
Inj 40 mg prefilled syringe		12	Copaxone
INTERFERON BETA-1-ALPHA - Special Authority see SA2051			
Note: Treatment on two or more funded multiple sclerosis tr			
Inj 6 million iu prefilled syringe	,	4	✓ Avonex
Injection 6 million iu per 0.5 ml pen injector	•	4	✓ Avonex Pen
INTERFERON BETA-1-BETA - Special Authority see SA2051 of			
Note: Treatment on two or more funded multiple sclerosis tr		, ,	
Inj 8 million iu per 1 ml	1,322.89	15	✓ Betaferon
NATALIZUMAB - Special Authority see SA2051 on the previous			
Note: Treatment on two or more funded multiple sclerosis tr		ısly is not pe	
Inj 20 mg per ml, 15 ml vial	1,750.00	1	✓ Tysabri
OCRELIZUMAB - Special Authority see SA2051 on the previou	s page - Retail pharm	acy	
Note: Treatment on two or more funded multiple sclerosis tr	reatments simultaneou	ısly is not pe	
Inj 30 mg per ml, 10 ml vial	9,346.00	1	Ocrevus
TERIFLUNOMIDE - Special Authority see SA2051 on the previous	ous page – Retail pha	rmacy	
a) Wastage claimable		•	
b) Note: Treatment on two or more funded multiple scleros	is treatments simultan	eously is no	t permitted.
Tab 14 mg	659.90	28	✓ Aubagio

Sedatives and Hypnotics

MELATONIN − Special Authority see SA1666 below − Retail pharmacy
Tab modified-release 2 mg − No more than 5 tab per day......28.22 30 ✓ Circadin

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
continued				
All of the following:				
1 Patient is aged 18 years or under*; and				
2 Patient has demonstrated clinically meaningful benefit from3 Patient has had a trial of funded modified-release melatoning	i funded modified-re	lease	melatonir	(clinician determined); and
recurrence of persistent and distressing insomnia; and	n discontinuation wit	n min	ne past 12	months and has had a
4 Funded modified-release melatonin is to be given at doses	no greater than 10 i	ma ne	er dav	
Note: Indications marked with * are unapproved indications.	no groater than to	9 P	or day.	
MIDAZOLAM – Safety medicine; prescriber may determine disper	neina fraguency			
Inj 1 mg per ml, 5 ml ampoule		10	/	Mylan Midazolam
ing it mg por mi, o mi ampodio	5.50	10		Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available	0.00			
on a PSO	14.90	10	/	Pfizer
On a PSO for status epilepticus use only. PSO must be e		epilep	oticus use	only.
Inj 5 mg per ml, 3 ml ampoule		5		Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available of	on			
a PSO		5		Pfizer
On a PSO for status epilepticus use only. PSO must be e	endorsed for status e	epilep	oticus use	only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 b	elow – Retail pharm	acy		
Inj 200 mg per ml, 1 ml ampoule	78.20	10	•	Max Health S29
⇒SA1386 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	without further rene	wal u	ınless noti	fied for applications meeting
the following criteria:				
Both:				
1 For the treatment of terminal agitation that is unresponsive	-	t		
2 The applicant is part of a multidisciplinary team working in				
TEMAZEPAM - Safety medicine; prescriber may determine dispe				
Tab 10 mg	1.33	25	•	<u>Normison</u>
TRIAZOLAM - Safety medicine; prescriber may determine disper	sing frequency			
Tab 125 mcg		100		
T 050	(9.85)	400		Hypam
Tab 250 mcg		100		Llunam
	(11.20)			Hypam
ZOPICLONE – Safety medicine; prescriber may determine disper	. ,		,	Zantalana Astauta
Tab 7.5 mg	9.56	500	•	Zopiclone Actavis
Stimulants/ADHD Treatments				
Odindiants/ADTID Treatments				
ATOMOXETINE				
Cap 10 mg		28		Generic Partners
Can 10 mg	107.03	00		Strattera
Cap 18 mg	27.06 107.03	28		Generic Partners Strattera
Cap 25 mg		28		Generic Partners
Cap 40 mg		28		Generic Partners
r ·- ··· · · · · · · · · · · · · · ·				<u></u>

Cap 60 mg.......46.51

28 28

28

✓ Strattera

✓ Generic Partners

✓ Generic Partners ✓ Generic Partners

107.03

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

DEXAMFETAMINE SULFATE - Special Authority see SA1149 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — **(Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 on the next page - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency 30 ✓ Rubifen 30 ✓ Ritalin ✓ Rubifen Tab extended-release 18 mg......7.75 30 ✓ Methylphenidate ER - Teva ✓ Rubifen 30 ✓ Rubifen SR 30 ✓ Methylphenidate ER 30 - Teva Tab extended-release 36 mg......15.50 ✓ Methylphenidate ER - Teva Tab extended-release 54 mg.......22.25 30 ✓ Methylphenidate ER

- Teva



Subsidy		Fully	Brand or	
(Manufacturer's P	rice)	Subsidised	Generic	
\$	Per	•	Manufacturer	

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

a) Only on a controlled drug form

 b) Safety medicine; prescriber may determine dispensin 	g trequency		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg	15.60	30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Reta	ail pharmacy		
Tab 100 mg		60	Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia DONEPEZIL HYDROCHLORIDE ★ Tab 5 mg ...4.34 90 ✓ Donepezil-Rex ★ Tab 10 mg .6.64 90 ✓ Donepezil-Rex RIVASTIGMINE – Special Authority see SA1488 on the next page – Retail pharmacy Patch 4.6 mg per 24 hour .48.75 30 ✓ Generic Partners Patch 9.5 mg per 24 hour .48.75 30 ✓ Generic Partners



Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per •	Manufacturer

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	,	28	✓ <u>Buprenorphine</u> Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	53.12	28	✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and

(Mar	Subsidy	Fully	Brand or
	nufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
 - 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	Antabuse
Antabuse to be Principal Supply on 1 November 2021			

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
 - 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.



ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. ✓ Habitrol Patch 7 mg - Up to 28 patch available on a PSO18.14 28 Patch 7 mg for direct distribution only - [Xpharm]......3.94 7 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO19.95 ✓ Habitrol 28 Patch 14 mg for direct distribution only - [Xpharm]......4.52 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO22.86 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......5.18 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......19.18 216 ✓ Habitrol 36 ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......21.02 216 ✓ Habitrol ✓ Habitrol 36 Gum 2 mg (Fruit) - Up to 384 piece available on a PSO38.21 384 ✓ Habitrol Gum 2 mg (Fruit) for direct distribution only - [Xpharm].......8.64 96 ✓ Habitrol

Gum 4 mg (Mint) for direct distribution only – [Xpharm]...............10.01

VARENICLINE TARTRATE – Special Authority see SA1845 below – Retail pharmacy

Gum 2 mg (Mint) - Up to 384 piece available on a PSO......38.21

Gum 2 mg (Mint) for direct distribution only - [Xpharm]......8.64

Gum 4 mg (Fruit) - Up to 384 piece available on a PSO44.17

Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17

a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack

384

96

384

96

384

96

✓ Habitrol

✓ Habitrol

✓ Habitrol

✓ Habitrol

✓ Habitrol

✓ Habitrol

- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4216.67	53 OP	✓ Varenicline Pfizer
25.64		✓ Champix
Tab 1 mg17.62	56	✓ Varenicline Pfizer
27.10		✓ Champix

(Champix Tab 0.5 mg \times 11 and 1 mg \times 42 to be delisted 1 January 2022) (Champix Tab 1 mg to be delisted 1 January 2022)

SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see \$A2046 below

Inj 25 mg vial77.00	1	Ribomustin
Inj 100 mg vial308.00	1	✓ Ribomustin
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

BUSULFAN - PUT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, , , , , , , , , , , , , , , , , , , ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1.387.00	1	✓ BiCNU
, 3	,		✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist	•	Ü	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist		20	- Louitorum i o
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ DBL Cisplatin
iiij i iiig pei iiii, 100 iiii viai	21.00	'	✓ Cisplatin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
, •	0.20	ring	Daxiei
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist		50	✓ Endoxan S29
	145.00		Cyclonex
	158.00	100	✓ Procytox S29
Inj 1 g vial – PCT – Retail pharmacy-Specialist		1	✓ Endoxan
	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist		1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
(Endoxan S29 Tab 50 mg to be delisted 1 January 2022)			
(Procytox S29 Tab 50 mg to be delisted 1 January 2022)			
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price)	_	Subsidised	
	\$	Per		Manufacturer
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	✓	CeeNU
Cap 40 mg	399.15	20	1	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	✓	Alkeran
Inj 50 mg - PCT only - Specialist		1	1	Alkeran
			✓	Alkeran S29 S29
	420.00		1	Tillomed S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
", 100 mg vai		•		100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP		1 mg		Baxter
THIOTEPA - PCT only - Specialist			,	
Inj 15 mg vial	CBS	1	1	Bedford S29
iiij 10 iiig viai		•		THIO-TEPA \$29
lai 400 associal	CDC			Tepadina S29
Inj 100 mg vial	CBS	1	•	Tepadina S29
Antimatabalitas				
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority se	e SA1467 below			
Ini 100 ma viol	75.06	4	./	Amonitidina Du

tidine Dr
dy's
а
r

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

				_
	Subsidy		Fully	Brand or
	(Manufacturer's Price) ;	Subsidised	Generic
	\$	Per	•	Manufacturer
CALCIUM FOLINATE				
Tab 15 mg - PCT - Retail pharmacy-Specialist	114 69	10	1	DBL Leucovorin
rab to mg 1 of Trotal pharmacy openation			-	Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialst		1		Calcium Folinate
ing to mg per mi, 5 mi viai – 1 of – Hetaii phaimacy-Specia	1131 7 .20	'	•	Sandoz
			./	
			•	Calcium Folinate
			_	Sandoz S29 S29
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist	9.49	1	/	Calcium Folinate
				Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓	Calcium Folinate
				Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	1	Calcium Folinate
				Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	/	Calcium Folinate
, , ,				Sandoz
			1	Calcium Folinate
			_	Sandoz S29 S29
laid a DOT ank Considist	67.54	4		
Inj 1 g - PCT only - Specialist		1	•	Calcium Folinate
				Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist	72.00	1	•	Calcium Folinate
			_	Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	/	Baxter
CAPECITABINE - Retail pharmacy-Specialist				
Tab 150 mg	10.00	60	1	Capercit
Tab 500 mg		120	1	Capercit
CLADRIBINE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml	740.06	1	1	Leustatin
Inj 10 mg for ECP		0 mg O		Baxter
, ,		o nig O	. •	Daxiei
CYTARABINE		_		
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	list400.00	5	•	Pfizer
Inj 100 mg per ml, 20 ml vial - PCT - Retail			_	
pharmacy-Specialist		1		Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	_	Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	list80.00 10	00 mg ()P	Baxter
FLUDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	1	Fludara Oral
Inj 50 mg vial - PCT only - Specialist		5	1	Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29 5	0 mg O	P 🗸	Baxter
FLUOROURACIL		-		
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	12 00	1	1	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1		Fluorouracil Ebewe
Inj 1 mg for ECP – PCT only – Specialist		100 mg	_	Baxter
			, -	
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist	00.50		,	DDI Ossaskala
Inj 1 g, 26.3 ml vial		1		DBL Gemcitabine
Inj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	•	Baxter

	Subsidy (Manufacturer's Prio \$	ce) Per	Fully Subsidised	d Generic
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	71.44	1	•	Irinotecan Accord \$29
			•	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	✓	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	✓	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist	_			
Special Authority see SA1725 below	428.00	100 ml (OP 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

IVIL	MOTIENATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
	, , , ,		Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
	, - 3, , 3-		Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
	, ··· · · · · · · · · · · · · · · ·		Sandoz
*	Inj 25 mg prefilled syringe14.99	1	✓ Methotrexate
	, ··· g p· - ···, ··· g -		Sandoz
*	Inj 30 mg prefilled syringe	1	✓ Methotrexate
	, ···g p· - ····, ···g - ·····		Sandoz
*	Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL
	,,,	-	Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
•	- 1, 20 mg por mi, 20 mi mai - 1 0 mi mai priamiady opodanomii motod	·	Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail	•	monionoxato Ebono
~	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*		1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist4.73	5 mg OP	✓ Baxter
-		•	- DUNIOI
rE	METREXED - PCT only - Specialist - Special Authority see SA1679 on the n		✓ Juno Pemetrexed
	Inj 100 mg vial	1	✓ Juno Pemetrexed ✓ Juno Pemetrexed
	Inj 500 mg vial	1 ma	✓ Baxter
	Inj 1 mg for ECP	1 mg	▼ Daxter

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cvcles: or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-S	Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

	Subsidy (Manufacturer's Pr	ice) Sub	Fully	Brand or Generic	
	\$	Per	•	Manufacturer	
BLEOMYCIN SULPHATE - PCT only - Specialist					
Inj 15,000 iu, vial	161.01	1	✓ [DBL Bleomycin	
				Sulfate	
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ E	Baxter	
BORTEZOMIB - PCT only - Specialist - Special Authority see S	SA1889 below				
Inj 3.5 mg vial	105.00	1	✓ E	Bortezomib	
				Dr-Reddy's	
Inj 1 mg for ECP	31.20	1 mg	✓ E	Baxter	

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

62.70	1	✓ DBL Dacarbazine
580.60	10	✓ Dacarbazine
		APP S29
62.70	200 mg OP	✓ Baxter
255.00	1	✓ Cosmegen
	0.5 mg OP	✓ Baxter
	-	
149.50	1	✓ Pfizer
	20 mg OP	✓ Baxter
	Ü	
48.75	1	✓ Docetaxel Sandoz
46.89	1	✓ DBL Docetaxel
	1	✓ Docetaxel
		Accord \$29
195.00	1	✓ Docetaxel Sandoz
	1 mg	✓ Baxter
	-	
10.00	1	Doxorubicin Ebewe
11.50	1	Doxorubicin Ebewe
17.00		✓ Arrow-Doxorubicin
23.00	1	Doxorubicin Ebewe
65.00	1	Arrow-Doxorubicin
69.99		Doxorubicin Ebewe
0.35	1 mg	✓ Baxter
25.00	1	Epirubicin Ebewe
	1	✓ Epirubicin Ebewe
	1	 Epirubicin Ebewe
0.50	1 mg	✓ Baxter
	62.70255.00149.50149.5048.7546.8926.95195.000.6510.0011.50 17.0023.0065.00	580.60 1062.70 200 mg OP255.00 1255.00 0.5 mg OP149.50 1149.50 20 mg OP48.75 146.89 126.95 1195.00 1150 1065 1 mg150 117.00 117.00 117.00 1

	Subsidy	Ful	ly Brand or
	(Manufacturer's Price)	Subsidise	
	\$	Per •	Manufacturer
ETOPOSIDE			
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali		1 •	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist		1 mg	Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist		3	
, ,	40.00	1	/ Etonophoo
Inj 100 mg (of etoposide base)			✓ Etopophos ✓ Baxter
Inj 1 mg (of etoposide base) for ECP		1 mg •	Daxier
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail phan	macy-Specialist		
Cap 500 mg	23.82	100	<u>Devatis</u>
IDARUBICIN HYDROCHLORIDE			
Inj 5 mg vial - PCT only - Specialist	93.00	1 •	Zavedos
Inj 10 mg vial - PCT only - Specialist		1 v	Zavedos
Inj 1 mg for ECP – PCT only – Specialist		· ·	Baxter
, ,		9	Durio
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority Wastage claimable	y see SA2047 below		
Cap 5 mg	5,122.76	28	Revlimid
Cap 10 mg		21	Revlimid
•	6,207.00	28	Revlimid
Cap 15 mg	,	21	/ Revlimid
	7.239.18		Revlimid
Cap 25 mg	,		Revlimid
	,0200		

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.0	0 50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.5	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.4	5 15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.4	0 15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.9	6 100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 20 mg vial	0 1	✓ Omegapharm S29
, ,		✓ Teva
Inj 1 mg for ECP470.7	5 1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial97.5	0 1	 Mitozantrone Ebewe
Inj 1 mg for ECP5.5	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1883 be	elow	
Tab 100 mg3,701.0	0 56	✓ Lynparza
Tab 150 mg3,701.0	0 56	✓ Lynparza
- CA1000 Chariel Authority for Cubaids		

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's P	rice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL – PCT only – Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg		1	✓ Paclitaxel Ebewe
•	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see	SA1979 below		
Inj 750 iu per ml, 5 ml vial		1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 The patient has relapsed acute lymphoblastic leukaemia; and

DENTOCTATINI (DECOVO/COECDAN/ONE) DOT I O : II :

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Sp	pecialist		
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pha			
Cap 50 mg	980.00	50	✓ Natulan S29

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per	•	Manufacturer
TEMOZOLOMIDE - Special Authority see SA1741 belo	v – Retail pharmacy			
Cap 5 mg	9.13	5	✓	Temaccord
Cap 20 mg	16.38	5	✓	Temaccord
	18.30		✓	Apo-Temozolomide
	136.00	14	✓	Accord S29
Cap 100 mg	35.98	5	✓	Temaccord
	40.20		✓	Apo-Temozolomide
	532.00	14	✓	Accord S29
Cap 140 mg	50.12	5	1	Temaccord
	400.00		✓.	Amneal S29
Cap 180 mg	620.00	14	1	Accord S29
Cap 250 mg		5	1	Temaccord
•	688.00		1	Amneal S29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Aut	hority see SA1124 below	1	
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRFTINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist479.50	100	✓ Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA1868 bel	OW	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	42 OP	✓ Venclexta
Tab 10 mg95.78	14 OP	✓ Venclexta
Tab 50 mg239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Specialist270.37	5	✓ DBL Vinblastine S29
		✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓ Navelbine
210.00		✓ Vinorelbine Ebewe
Inj 1 mg for ECP1.25	1 mg	✓ Baxter

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable

224 Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

for 6 months for applications meeting the following criteria:

Roth:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

Tab 20 mg3,774.06	60	✓ Sprycel
Tab 50 mg6,214.20	60	✓ Sprycel
Tab 70 mg	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see SA2000 below

✓ Tarceva	30	Tab 100 mg764.00
✓ Tarceva	30	Tab 150 mg

⇒SA2000 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and

continued...

- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued defitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
 - 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2001 below

Tab 250 mg1,700.00 30 **✓ Iressa**

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Fither
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	rad 100 mg - [xpnarm] - Special Authority see SAT	460 on the		
	next page	2,400.00	60	✓ Glivec
*	Cap 100 mg		60	✓ Imatinib-Rex
	Cap 400 mg		30	✓ Imatinib-Rex

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	✓ Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised		
PALBOCICLIB - Retail pharmacy-Specialist - Special Authority	see SA1894 below				
Wastage claimable					
Cap 75 mg	4,000.00	21	✓	Ibrance	
Cap 100 mg	4,000.00	21	✓	Ibrance	
Cap 125 mg	4,000.00	21	✓	Ibrance	

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail ph	armacy
Tah 200 mg	1.3

✓ Votrient	30	1,334.70	······································	Tab 200 mg
✓ Votrient	30	2.669.40		Tab 400 mg

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and

Subsidy (Manufacturer's Pric	e)	Fully Brand or Subsidised Generic	
\$	Per	1	Manufacturer

continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

vvastage ciaimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5.000.00	56	✓ Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Aut	thority see SA2002	on the next page -	- Retail pharmacy

Cap 12.5 mg	2,315.38	28	Sutent
Cap 25 mg	· ·	28	✓ Sutent
Cap 50 mg	· ·	28	✓ Sutent

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 82

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

Tab 250 mg4,276.19 120 **✓ Zytiga**

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline: and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

DICALOTAMIDE			
Tab 50 mg	1.36	10	✓ Calutide-50 S29
	4.21	28	✓ Binarex
(Calutide-50 S29	Tab 50 mg to be delisted 1 January 2022)		
FLUTAMIDE			
Tab 250 mg.	107.55	90	✓ Prostacur S29
_	119 50	100	✓ Flutamin

	Subsidy (Manufacturer's Price)	Su Per	Fully bsidised	Brand or Generic Manufacturer	
FULLYEOTRANT - Date if the consequence Operation is the Consequence of	OA4005 balana	1 61		Wallulacturei	
FULVESTRANT – Retail pharmacy-Specialist – Special Authority Inj 50 mg per ml, 5 ml prefilled syringe		2	√ Fa	aslodex	

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

Tab 160 mg	63.53	30	✓ Apo-Megestrol
OCTREOTIDE			
Inj 100 mcg per ml, 1 ml ampoule	18.69	5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule		5	✓ Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	✓ Octreotide
, , ,			MaxRx S29
	56.87		✓ DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓ DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule	72.50	5	✓ Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial		5	✓ DBL Octreotide
	222.00		Octreotide
			(Sun) \$29
OCTREOTIDE LAR (SOMATOSTATIN ANALOGUE) - Special Aut	hority see SA2004	4 below – F	Retail pharmacy
Inj LAR 10 mg prefilled syringe		1	✓ Sandostatin LAR
Inj LAR 20 mg prefilled syringe	.2,358.75	1	 Sandostatin LAR
Inj LAR 30 mg prefilled syringe	.2,951.25	1	 Sandostatin LAR

⇒SA2004 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

TAMOXIFEN CITRATE

*	Tab 10 mg	60	✓ <u>Tamoxifen Sandoz</u>
*	Tab 20 mg6.65	60	✓ <u>Tamoxifen Sandoz</u>

Aromatase Inhibitors

AN	ASTROZOLE			
*	Tab 1 mg	.4.55	30	✓ Anatrole

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
EXEMESTANE * Tab 25 mg	14.50	30	√ P	fizer Exemestane
LETROZOLE * Tab 2.5 mg	5.84	30	✓ L	etrole

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE		
* Tab 25 mg7.35	60	Azamun
* Tab 50 mg7.60	100	✓ Azamun
* Inj 50 mg vial199.00	1	✓ Imuran
MYCOPHENOLATE MOFETIL		
Tab 500 mg35.90	50	Cellcept
Cap 250 mg35.90	100	 Cellcept
Powder for oral lig 1 g per 5 ml - Subsidy by endorsement187.25	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2048 below - Retail	pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate: and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm: Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm: Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment: and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate,

Subsidy	,	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

ciclosporin, or acitretin; and

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following: 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
 - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and

Subsidy		Fully	Brand or
(Manufacturer's	Price) Subs	sidised	Generic
\$	Per	•	Manufacturer

continued...

- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application: or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specia	alist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG §29
(SII-Onco-BCG S29 Inj 40 mg per ml, vial to be delisted 1 April	1 2022)		

Monoclonal Antibodies

ADALIMUMAB - Special Authority see SA2049 below -	Retail pharmacy		
Inj 20 mg per 0.4 ml prefilled syringe	1,599.96	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA2049 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either: 1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application. Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

continued...

25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Fither

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less: or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — **(hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per •	Manufacturer

continued...

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plague psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plague or plagues have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from. at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Subsidy (Manufacturer's Price)	Fully Subsidised		. ,		Brand or Generic
 \$	Per	√	Manufacturer		

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see	SA1697 on the next	page	
Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see SA2050 below

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	Baxter

⇒SA2050 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

ilei.

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcojosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plague psoriasis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plague psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot. as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

		nority see SA1896 below - Retail pharmacy	MEPOLIZUMAB - Special Authority see SA189
✓ Nucala	1	1,638.00	Inj 100 mg prefilled pen
✓ Nucala	1		Ini 100 mg vial

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Author	ority see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB – Special Authority see SA1744 below – Retail	pharmacy		
Inj 150 mg prefilled syringe	450.00	1	Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and

- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
 - 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✔ Manufacturer

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist	pecial Authority see SA197	'6 below	
Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2061 below

Inj 100 mg per 10 ml vial	275.33	2	✓ <u>Riximyo</u>
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2061 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
 - 3 The patient has good performance status; and
 - 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
 - 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
 - 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

375 mg/m2 administered weekly for four weeks; and

- 2 Either
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD: and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:

<u> </u>				
	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects: and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles: or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of $2 \times 1,000$ mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

0.1.1		
Subsidy	Fully	Brand or
(Manufacturer's F	Price) Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
 - 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2044 below - Retail pharmacy

⇒SA2044 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for psoriatic arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for psoriatic arthritis; or

2 All of the following:

- 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
 - 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA1977 below

Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

⇒SA1977 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2 systemic juvenile idiopathic arthritis; or
- 2.3 adult-onset Still's disease; or
- 2.4 polyarticular juvenile idiopathic arthritis; or
- 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules: and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule: and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

		TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below	ı
✓ Herceptin	1	Inj 150 mg vial	
✓ Herceptin	1	Inj 440 mg vial	
✓ Baxter	1 mg	Inj 1 mg for ECP	

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned: or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and

Subsidy	Fu	lly Brand	or
(Manufacturer's Price)	Subsidis	ed Generi	ic
\$	Per	 Manufa 	acturer

continued...

- 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
- 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial	2,320.00	1	✓ Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	23.20	1 mg	✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
 - 5 Baseline measurement of overall tumour burden is documented (see Note); and
 - 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Fither:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2007 below

✓ Keytruda 1 mg ✓ Baxter

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and

1.2 Either:

- 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
- 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression: and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
	Per	1	Manufacturer	

continued...

per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg		50	✓ Neoral
Oral liq 100 mg per ml		50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail p	harmacy		
Wastage claimable	-		
Tab 10 mg	6,512.29	30	Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor
•			

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg749.99	100	Rapamune
Tab 2 mg1,499.99	100	✓ Rapamune
Oral lig 1 mg per ml449.99	60 ml OP	✓ Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy (Manufacturer's Price)	Fı Subsidis	ılly ed	Brand or Generic	_
\$	Per	✓	Manufacturer	

continued...

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy	Fully	Brand or
(Manufacturer's Price)	(Manufacturer's Price) Subsidised	
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA	1367 above – I	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see S	SA1367 above	– Retail pharr	macy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	205.00	1 OD	√ Albay
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent	205.00	1 OP	✓ Venomil \$29
uneu venom, wim unuerit	303.00	I OF	A ACHOUNII 258

	Subsidy		Fully Brand or
	(Manufacturer's Price		,
	\$	Per	✓ Manufacturer
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1 12	100	✓ Zista
* Oral liq 1 mg per ml		200 ml	✓ Histaclear
	2.0+	200 1111	- Inotablear
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen
DEXTROCHLORPHENIRAMINE MALEATE			
* Tab 2 mg	2.02	40	
· · · · g	(8.40)		Polaramine
	1.01	20	1 olaramine
	(5.99)	20	Polaramine
* Oral liq 2 mg per 5 ml		100 ml	Foldialillile
* Oral liq 2 mg per 5 ml		100 1111	Doloromino
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg	4.34	20	
·	(8.23)		Telfast
* Tab 120 mg		10	
· · · · · · · · · · · · · · · · · · ·	(8.23)		Telfast
	14.22	30	
	(26.44)	00	Telfast
	(20.44)		Tellast
LORATADINE			
* Tab 10 mg	1.69	100	✓ Lorafix
* Oral liq 1 mg per ml	1.43	100 ml	Haylor syrup
	2.95	120 ml	✓ Lorfast
Haylor syrup to be Sole Supply on 1 September 2021			
(Lorfast Oral lig 1 mg per ml to be delisted 1 September 2021)			
PROMETHAZINE HYDROCHLORIDE			
	1.60	F0	✓ Allersoothe
* Tab 10 mg		50	
* Tab 25 mg		50	✓ Allersoothe
* Oral liq 1 mg per 1 ml		100 ml	✓ Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a f	PSO 17.87	5	✓ Hospira
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
	0.00	00 doos 00	✓ Qvar
Aerosol inhaler, 50 mcg per dose		200 dose OP	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	00 dose OP	✓ Beclazone 250
BUDESONIDE			
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort Turbuhaler
D 1 () 1 1 1 2 200 1	46.55		
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's		dised Generic
	\$	Per	✓ Manufacturer
LUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose	7.50	60 dose OP	 Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose	13.60	120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	sts		
FORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose dev	rice20.64	60 dose	
,,,,,,	(35.80)		Foradil
FORMOTEROL FUMARATE DIHYDRATE	()		
Powder for inhalation 4.5 mcg per dose, breath activated	۵) ۱۵ ۵۵	60 dose OP	
(equivalent to eformoterol fumarate 6 mcg metered dos	,	60 dose OP	Oxis Turbuhaler
	(16.90)		Oxis Turbunaler
NDACATEROL			_
Powder for inhalation 150 mcg		30 dose OP	Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	 Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta	-Adrenocept	or Agonists	
NUDECONUDE WITH EFORMOTEROL			
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol	. 241.		
fumarate per dose (equivalent to 200 mcg budesonide		100 daaa OD	/ Dua Daam Culinaman
6 mcg eformoterol fumarate metered dose)		120 dose OP	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma			
per dose (equivalent to 400 mcg budesonide with 12 mc	cg		
eformoterol fumarate metered dose) - No more than 2	20.50	100 00	(D.D. 0)
dose per day		120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6	mcg33.74	120 dose OP	✓ Symbicort
			Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6	mcg44.08	120 dose OP	✓ Symbicort
			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	44.08	60 dose OP	Symbicort
			Turbuhaler 400/12
LUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta
			F

	_			
	Subsidy (Manufacturer's \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer	
FLUTICASONE WITH SALMETEROL Aerosol inhaler 50 mcg with salmeterol 25 mcg Aerosol inhaler 125 mcg with salmeterol 25 mcg Powder for inhalation 100 mcg with salmeterol 50 mcg – No		120 dose OP 120 dose OP	✓ <u>Seretide</u> ✓ <u>Seretide</u>	
more than 2 dose per day		60 dose OP	 ✓ Seretide Accuhaler ✓ Seretide Accuhaler 	
Beta-Adrenoceptor Agonists		00 4000 01		
SALBUTAMOL Oral liq 400 mcg per ml Infusion 1 mg per ml, 5 ml Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO	118.38	150 ml 10 5	✓ Ventolin✓ Ventolin✓ Ventolin	
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	3.80	200 dose OP	✓ Respigen✓ SalAirVentolin	
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	8.96	20	✓ Asthalin	
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	9.43	20	✓ Asthalin	
TERBUTALINE SULPHATE Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler	
Anticholinergic Agents				
IPRATROPIUM BROMIDE Aerosol inhaler, 20 mcg per dose CFC-free		200 dose OP	✓ Atrovent	
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 nei available on a PSO		20	✓ <u>Univent</u>	
Inhaled Beta-Adrenoceptor Agonists with Anticl	holinergic A	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free		200 dose OP	✓ Duolin HFA	
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	11.04	20	✓ Duolin	

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet
Cap 267 mg - Wastage claimable	3,645.00	270	Esbriet

(Esbriet Cap 267 mg to be delisted 1 January 2022)

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

Leukotriene Receptor Antagonists

MON	ITELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
*	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	28	✓ Montelukast Mylan

Mast Cell Stabilisers

NEDOCROMIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking nedocromil prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of nedocromil.

(Thade Nerosof Inhalor, 2 mg per dose of a free to be deficited if deplember

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a	100.00	-	A DDI Andrea bellin
	PSO	180.00	5	✓ DBL Aminophylline
TH	EOPHYLLINE			
*	Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml	16.60	500 ml	✓ Nuelin

Mucolytics

		8 below – Retail pharmacy	DORNASE ALFA – Special Authority see SA1978 b
✓ Pulmozyme	6	250.00	Nebuliser soln, 2.5 mg per 2.5 ml ampoule

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

e

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	S	Subsidised	Generic	
	\$	Per	1	Manufacturer	
IVACAFTOR - PCT only - Specialist - Special Authority see SA	2017 below				
Tab 150 mg	29,386.00	56	✓ K	Calydeco	
Oral granules 50 mg, sachet	29,386.00	56	✓ K	alydeco	
Oral granules 75 mg, sachet	29,386.00	56	✓ K	Calydeco	

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

BUDESONIDE		
Metered aqueous nasal spray, 50 mcg per dose2.54	200 dose OP	✓ SteroClear
Metered aqueous nasal spray, 100 mcg per dose2.84	200 dose OP	✓ SteroClear
FLUTICASONE PROPIONATE		
Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ Flixonase Hayfever & Allergy
IPRATROPIUM BROMIDE	15 ml OD	Ilmirramt
Aqueous nasal spray, 0.03%	15 ml OP	✓ Univent

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
PEAK FLOW METER				
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	//////////////////////////////////////
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	2.95	1	√ e	-chamber Turbo
510 ml (single patient)	5.12	1	√ e	-chamber La Grande
800 ml	6.50	1	✓ V	/olumatic
Respiratory Stimulants				
CAFFEINE CITRATE				
Oral liq 20 mg per ml (10 mg base per ml)	15.10 2	5 ml OP	✓ <u>E</u>	Biomed

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BI	ENZETHONIUM		
For Vosol ear drops with hydrocortisone powder refer Stand	lard Formulae, <mark>pa</mark>	age 237	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and			
benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYO	IN AND NYSTAT	ΓIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g		7.5 ml OP	✓ Kenacomb
3 3 31 3			
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and	4.50	0 OD	
gramicidin 50 mcg per ml		8 ml OP	Sofradex
	(9.27)		Sorradex
FRAMYCETIN SULPHATE	4.40	0 OD	
Ear/Eye drops 0.5%		8 ml OP	Caframusin
	(8.65)		Soframycin
Eye Preparations			
Lyc i reparations			
Eye preparations are only funded for use in the eye, unless expl	icitly stated other	wise.	
Auti Infantiva Dunamentiana			
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
ViruPOS to be Principal Supply on 1 September 2021		- 3 -	
CHLORAMPHENICOL			
Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%		10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * a		dications.	
CIPROFLOXACIN			
Eye drops 0.3% – Subsidy by endorsement	9.73	5 ml OP	✓ Ciprofloxacin Teva
a) When prescribed for the treatment of bacterial kerati		terial conjunctiv	•
or for the second line treatment of chronic suppurati			
accordingly. Note: Indication marked with a * is an	unapproved indic	cation.	
b) Ciprofloxacin Teva to be Principal Supply on 1 Nove	mber 2021		
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE			•
* Eye drops 0.1%	2.97	10 ml OP	
jyy	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID]	/		
Eve drone 1%	5 20	5 a OP	✓ Fucithalmic

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

5 g OP

✓ Fucithalmic

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer	
TOBRAMYCIN Eye oint 0.3% Eye drops 0.3%		3.5 g OP 5 ml OP		obrex obrex	
Corticosteroids and Other Anti-Inflammatory Pr	reparations				
DEXAMETHASONE # Eye oint 0.1%	4.50	3.5 g OP 5 ml OP		Maxidex Maxidex	
- Retail pharmacy		1	✓ 0)zurdex	

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not vet completed a family: and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eve per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
	Voltaren Ophtha to be Principal Supply on 1 November 2021			

	Subsidy (Manufacturer's F	Price) Sub	Fully sidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
FLUOROMETHOLONE					
* Eye drops 0.1%	3.09	5 ml OP	✓ F	ML	
	5.20		√ F	lucon	
KETOROLAC TROMETAMOL - Special Authority see SA1981	below - Retail ph	armacy			
Eye drops 0.5%	9.50 ·	5 ml OP	√ p	Acular	
OA4004 Out a dal Austraultur fam Outral de					

⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71	4 ml OP	
, , , , , , , , , , , , , , , , , , , ,	(10.34)		Livostin
	(10.01)		21100111
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eve drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
	г оо	40 ml OD	/ Duadwisslams AFT
Eye drops 1%		10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	- Retail pharn	nacv
Eve drops 0.5%, single dose (preservative free)		20 dose	✓ Minims
2,0 diopo 0.0/0, oliigio dobo (proborvativo 1100)		_0 0000	
			Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

5 ml OP

2.5 ml OP

✓ Rexacrom

✓ Arrow-Timolol

✓ Timoptol XE

SODIUM CROMOGLICATE

Glaucoma Preparations - Beta Blockers		
BETAXOLOL		
* Eye drops 0.25%	5 ml OP	✓ Betoptic S
* Eye drops 0.5%	5 ml OP	✓ Betoptic
TIMOLOL		
* Eve drops 0.25%	5 ml OP	✓ Arrow-Timolol



	Subsidy (Manufacturer's Pi \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
Glaucoma Preparations - Carbonic Anhydrase In	hibitors			
CETAZOLAMIDE	17.00	100	./ -	Diamox
F Tab 250 mg	17.03	100	V L	латтох
RINZOLAMIDE Eye drops 1%	7 20	5 ml OP	1	zont
Azopt to be Principal Supply on 1 September 2021	7.30	3 IIII OF	• •	zopt
ORZOLAMIDE HYDROCHLORIDE				
Eye drops 2%	9 77	5 ml OP		
- Lyo diopo 270	(17.44)	0 1111 01	Т	rusopt
ORZOLAMIDE WITH TIMOLOL	()			
Eye drops 2% with timolol 0.5%	2.73	5 ml OP	√ [Oortimopt
				·
Glaucoma Preparations - Prostaglandin Analogu	es			
IMATOPROST				
F Eye drops 0.03%	3.30	3 ml OP	✓ E	Bimatoprost
				Multichem
ATANOPROST				
Fye drops 0.005%	1.57	2.5 ml OP	✓ T	eva
RAVOPROST	7.00	- 100		
Eye drops 0.004%		5 ml OP 2.5 ml OP		ravopt ravatan
	9.75	-		
Fravent Fire drang 0.0040/ to be delicted 1. December 2021)	10.50	5 ml OP	• 1	lylan S29
Travopt Eye drops 0.004% to be delisted 1 December 2021)				
Mylan S29 Eye drops 0.004% to be delisted 1 December 2021)				
Glaucoma Preparations - Other				
RIMONIDINE TARTRATE				
F Eye drops 0.2%	4.29	5 ml OP	✓ A	rrow-Brimonidine
RIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
Fye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	√ (Combigan
ATANOPROST WITH TIMOLOL				
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ A	rrow - Lattim
Arrow - Lattim to be Sole Supply on 1 September 2021				
ILOCARPINE HYDROCHLORIDE				
F Eye drops 1%		15 ml OP		sopto Carpine
Eye drops 2%		15 ml OP		sopto Carpine
Eye drops 4%		15 ml OP	✓	sopto Carpine
Subsidised for oral use pursuant to the Standard Formula	е.			
Eye drops 2% single dose – Special Authority see SA0895	21.05	20 dose	./ 1	linime Dilacerniae
below – Retail pharmacy	31.95	∠u dose	▼ \	linims Pilocarpine

Either:

Subsidy	Ful	ly Brand or	
(Manufacturer's Pr	ice) Subsidise	d Generic	
\$	Per	 Manufacturer 	

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Mudriation and Cycloplagian

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cyclopiegics		
ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a prescription	20 dose	✓ Minims Cyclopentolate
TROPICAMIDE		• •
* Eye drops 0.5% 7.15 ★ Eye drops 1% 8.66	15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 237 HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	✓ Methopt
* Eye drops 0.3% with dextran 0.1%2.30	15 ml OP	✓ Poly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eve drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail	pharmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Au	thority see SA1388 abov	e – Reta	il pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special A	Authority see SA1388 ab	ove – Re	tail pharmacy
Eye drops 1 mg per ml	13.85	ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The	Pharmacy Procedures M	/lanual re	striction allowing one bottle per
month is not relevant and therefore only the prescribe	ed dosage to the neares	OP may	be claimed

SENSORY ORGANS

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully dised	Brand or Generic Manufacturer
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	4.15	15 ml OP	✓ N	laphcon Forte
OLOPATADINE Eye drops 0.1%	2.20	5 ml OP	√ <u>0</u>	Nopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.63	3.5 g OP	✓ P	oly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	3.80	5 g OP	✓ V	itA-POS

Subsidy (Manufacturer's Price) Per

Subsidised

Fully

Brand or Generic Manufacturer

Agents Used in the Treatment of Poisonings

Antidotes

ACETYL CYSTEINE

Inj 200 mg per ml, 10 ml ampoule58.76 10 ✓ DBL Acetylcysteine ✓ Martindale Pharma \$29

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO
- ✓ DBL Naloxone 5 Hydrochloride

Removal and Elimination

CHARCOAL

✓ Carbosorb-X 250 ml OP

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible		28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels. liver or cardiac MRI T2*: or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis: or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE -	 Special Authority see SA1480 on the next page 	- Retail pharmacy	1
Tah 500 mg		533 17	10

Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE * Inj 500 mg vial	84.53	10	✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	. LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium Glycerol BP	400 mg 4 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate	60 mg	Water	to 40 ml
Glycerol Preservative	40 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)		Water (Preservative should be used if quantity supplied is	to 500 ml for more
Codeine phosphate Glycerol	300 mg 40 ml	than 5 days.)	
Preservative Water	qs to 100 ml	SALIVA SUBSTITUTE FORMULA Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative Water	qs to 500 ml
Calcium folinate 15 mg tab Preservative	1 tab qs	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more
Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	to 500 ml for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
METHADONE MIXTURE		Water (Only funded if prescribed for treatment of hyponatra	qs aemia)
Methadone powder Glycerol	qs qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection Glycerol BP	10 vials 40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate	10 g	Water (Only funded if prescribed for treatment of Clostridiu	to 100 ml
Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	to 100 ml	following metronidazole failure)	im dillicile
OMEPRAZOLE SUSPENSION	ila mixture)	VOSOL EAR DROPS	
Omeprazole adules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder Vosol Ear Drops	1% to 35 ml
	10 100 1111		
PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully sidised	Brand or Generic Manufacturer
Extemporaneously Compounded Preparations a	and Galenicals			
CODEINE PHOSPHATE - Safety medicine; prescriber may dete	rmine dispensing fre	equency		
Powder – Only in combination		25 g	0	ouglas
Only in extemporaneously compounded codeine linctus.				
COLLODION FLEXIBLE Note: This product is no longer being manufactured by the s	upplior and will be d	aliated from	a tha Ci	shadula at a data ta ba
determined.	uppliel allu will be u	elisteu ii oli	i iiie oi	inedule at a date to be
Collodion flexible	19.30	100 ml	✓ P	SM
COMPOUND HYDROXYBENZOATE — Only in combination Only in extemporaneously compounded oral mixtures. Soln	30.00	100 ml	✓ N	lidwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination Only in combination with Ora-Plus.				
Suspension	30.95	473 ml	√ 0	ra-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination Only in combination with Ora-Plus.			_	
Suspension	30.95	473 ml	√ <u>C</u>	ra-Sweet
GLYCEROL				
* Liquid – Only in combination Only in extemporaneously compounded oral liquid prepa		500 ml	✓ <u>h</u>	ealthE Glycerol BP
METHADONE HYDROCHLORIDE a) Only on a controlled drug form b) No patient co-payment payable c) Safety medicine; prescriber may determine dispensing fred) Extemporaneously compounded methadone will only be a (methadone powder, not methadone tablets).	eimbursed at the rat			
Powder	7.04	1 g	✓ A	AF I
METHYL HYDROXYBENZOATE Powder	8 98	25 g	✓ N	lidwest
METHYLCELLULOSE		20 g		<u></u>
Powder	36.95	100 g	✓ N	lidWest
Suspension - Only in combination		473 ml		ra-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHA	ARIN - Only in com	bination		
Suspension	30.95	473 ml	√ <u>C</u>	ra-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Onl	,			
Suspension	30.95	473 ml	√ <u>C</u>	ra-Blend
PHENOBARBITONE SODIUM	50.50	40		
Powder – Only in combination	52.50 325.00	10 g 100 g		lidWest lidWest
Only in children up to 12 years	323.00	100 g	V IV	iiuwesi
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxybenz	oate 10% solution.			
Liq		500 ml	✓ N	lidwest
SODIUM BICARBONATE				
Powder BP — Only in combination		500 g	✓ <u>N</u>	lidwest

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation Liq		500 ml	✓ <u>M</u>	lidwest
WATER Tap – Only in combination	0.00	1 ml	✓ T	ap water

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- 1 Infant or child aged four years or under; and
 - 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 241

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
	Per	✓	Manufacturer	

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacture

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Sp	pecial Authority see SA1095 above – F	Hospital pharm	acy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
	7.50	1,000 ml OP	Diason RTH
			✓ Glucerna Select
			RTH

(Glucerna Select RTH Liquid to be delisted 1 September 2021)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	1.88	250 ml OP	✓ Glucerna Select
	1.78	237 ml OP	
	(2.10)		Sustagen Diabetic
	(2.10)	200 ml OP	Nutren Diabetes

(Glucerna Select Liquid (vanilla) to be delisted 1 September 2021) (Sustagen Diabetic Liquid (vanilla) to be delisted 1 February 2022)

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism: or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 243

Subsidy (Manufacturer's Price)

Subsidised

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or

continued...

Sub	bsidy F	ully Br	and or
(Manufactu	urer's Price) Subsidis	sed Ge	eneric
•	\$ Per	✓ Ma	anufacturer

2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1379 on the previous Liquid	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1379 on the previous Liquid	
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1379 pharmacy [HP3]	on the previous page - Hospital
Liquid	OP ✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379 on the previous page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page 1.5KCAL/ML - Special Authority see SA1379 on the page	age – Hospital pharmacy [HP3]
Liquid (strawberry)	
Liquid (vanilla)	
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 on the previous page	e – Hospital pharmacy [HP3]
Liquid (chocolate)	
Liquid (strawberry)1.07 200 ml C	OP ✓ Pediasure
Liquid (vanilla)	OP ✓ Pediasure
1.34 250 ml C	OP ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority see SA1379 on t pharmacy [HP3]	he previous page – Hospital
Liquid (unflavoured)	OP ✓ Fortini Multi Fibre
Liquid (chocolate)	OP ✓ Fortini Multi Fibre
Liquid (strawberry)1.60 200 ml C	OP ✓ Fortini Multi Fibre
Liquid (vanilla)	OP ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED - Special Authority see SA1379 on the previous page - Ho	spital pharmacy [HP3]
Powder	

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised

245

	Subsidy (Manufacturer's Pr \$	rice) Subsi Per	Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		us page – Hos 220 ml OP	•	oharmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110 Liquid	2.88 (3.31)	s page – Hospit 237 ml OP 4 OP		armacy [HP3] NovaSource Renal Renilon 7.5
Liquid (caramel) 125 ml	11.52	4 OP	✓	Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease: or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LiquidLiquid		1,000 ml OP	
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority s	ee SA1377 above	– Hospital phar	macy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	e SA1377 above –	Hospital pharm	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special A	uthority see SA137	7 above – Hosp	oital pharmacy [HP3]
Liquid	12.04	1,000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per 🗸 Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

✓ fully subsidised 247

Subs (Manufactur		
\$	Per	Manufacturer

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy	e)	Fully	Brand or	
(Manufacturer's Pric		Subsidised	Generic	
\$	Per	· •	Manufacturer	

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube: or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant: and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum: or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure: or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia: or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3] Liquid......7.00 1.000 ml OP ✓ Nutrison Energy

✓ fully subsidised 249

	Subsidy		Fully Brand or
	(Manufacturer's		,
	\$	Per	✓ Manufacturer
	Ψ	1 61	Wallulaciulei
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	page 247 - Ho	spital pharmacy [HP3]
Liquid		250 ml OP	✓ Isosource Standard
Liquid			
	5.29	1,000 ml OP	✓ Nutrison Standard
			RTH
			✓ Osmolite RTH
ENTERNI EEER MITH EIRRE 0.00 KOM (MI On a del Andrea)	044050		
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority			
Liquid	5.29	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority se	e SA1859 on ⊧	page 247 – Hospi	tal pharmacy [HP3]
Liquid	5.29	1,000 ml OP	✓ Jevity RTH
1		,	✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s	see SA1859 on	n page 247 – Hosp	oital pharmacy [HP3]
Liquid	1.75	250 ml OP	✓ Ensure Plus HN
	7.00	1,000 ml OP	✓ Ensure Plus RTH
	7.00	1,000 1111 01	
			Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
ORAL EEER (ROWDER) Creatial Authority and CA10EO or rea	047 Haanii		21
ORAL FEED (POWDER) - Special Authority see SA1859 on page			-
Powder (chocolate)	14.00	840 g OP	 Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
Develor (verille)			
Powder (vanilla)	14.00	840 g OP	 Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
		ŭ	
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa			
Additional subsidy by endorsement is available for patients be	eing bolus fed t	through a feeding	tube, who have severe
epidermolysis bullosa, or as exclusive enteral nutrition in child	tren under the	age of 18 years for	or the treatment of Crohn's
disease, or for patients with COPD and hypercapnia, defined			
	as COZ value	exceeding 55mm	rig. The prescription must be
endorsed accordingly.			
Liquid (banana) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	٠,		
	(1.26)		Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)	-	Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 r	nl		
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Lieuwiel (atmosphanes) — Lliebaur authairth, af \$4 00 mar 000 mal with			
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml wi	, ,		•
		007 ml OD	
Endorsement		237 ml OP	E 5
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
	(1.20)		ι στασιρ

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 247 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Liquid (chocolate) — Higher Subsidy of \$1.20 per 200 Hil With			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 251

SPECIAL FOODS

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 c		age – Hospital p 500 ml OP		cy [HP3] lutrison Concentrated
	11.00	1,000 ml OP	_	insure Two Cal HN RTH wo Cal HN RTH
(Two Cal HN RTH Liquid to be delisted 1 February 2022)			-	
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients epidermolysis bullosa. The prescription must be endorsed a Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with	being bolus fed t		, .	•
Endorsement	0.96 (1.90)	200 ml OP	Т	wo Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA1106 above - Hospital pharmacy [HP3]				
Powder	6.53	300 g OP	✓ Nutilis	
	7.25	380 g OP	Feed Thickener	
			Karicare Antamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

(Manufacturer's Price) Subsidised Manufacturer's Price) Per ✓ Manufacturer Manufacturer's Price) Per ✓ Manufacturer SELUTEN FREE BAKING MIX – Special Authority see SA1729 on the previous page – Hospital pharmacy [HP3] Powder		Subsidy	F	ully Brand or
SELUTEN FREE BAKING MIX - Special Authority see SA1729 on the previous page - Hospital pharmacy [HP3] Powder		(Manufacturer's P	rice) Subsidi	sed Generic
Powder		\$	Per	✓ Manufacturer
Powder	SLUTEN FREE BAKING MIX - Special Authority see SA172	9 on the previous pa	ige – Hospital ph	armacy [HP3]
Baking Mix BLUTEN FREE BREAD MIX - Special Authority see SA1729 on the previous page - Hospital pharmacy [HP3] Powder				,. ,
SELUTEN FREE BREAD MIX - Special Authority see SA1729 on the previous page - Hospital pharmacy [HP3] Powder		(5.15)		Healtheries Simple
SELUTEN FREE BREAD MIX - Special Authority see SA1729 on the previous page - Hospital pharmacy [HP3] Powder		, ,		Baking Mix
Powder	SLUTEN EREE RREAD MIX - Special Authority see SA170	on the previous pag	no – Hospital ph	· ·
(7.32) NZB Low Gluten Bread Mix 3.51 (10.87) Horleys Bread Mix 3.51 (10.87) Horleys Bread Mix 3.51 (10.87) Horleys Bread Mix SEUTEN FREE FLOUR - Special Authority see SA1729 on the previous page - Hospital pharmacy [HP3] Powder	·			aillacy [i ir oj
Bread Mix 3.51 (10.87) Horleys Bread Mix 3.51 (10.87) Horleys Bread Mix 3.51 (10.87) Horleys Bread Mix SILUTEN FREE FLOUR – Special Authority see SA1729 on the previous page – Hospital pharmacy [HP3] Powder	1 OWGG1		1,000 g OI	NZR Low Gluten
3.51 (10.87)		(1.52)		
(10.87)		3.51		DICAU MIX
Corn and Vegetable Spirals Corn and Corn Lasagne Sheets Corn and Corn Macaroni Corn and Corn Penne Corn and Corn Spage Corn and Corn				Horleys Bread Mix
Powder	NUTEN EDEE ELOUD. Consid Authority and CA4700	, ,		•
(18.10)	·			cy [HP3]
BLUTEN FREE PASTA — Special Authority see SA1729 on the previous page — Hospital pharmacy [HP3] Buckwheat Spirals	Powder		2,000 g OP	Harlana Flance
Buckwheat Spirals 2.00 250 g OP (3.11) Orgran Corn and Vegetable Shells 2.00 250 g OP (2.92) Orgran Corn and Vegetable Spirals 2.00 250 g OP (2.92) Orgran Rice and Corn Lasagne Sheets 1.60 200 g OP (3.82) Orgran Rice and Corn Macaroni 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP		()		,
Corn and Vegetable Shells				cy [HP3]
Corn and Vegetable Shells 2.00 250 g OP (2.92) Orgran Corn and Vegetable Spirals 2.00 250 g OP (2.92) Orgran Rice and Corn Lasagne Sheets 1.60 200 g OP (3.82) Orgran Rice and Corn Macaroni 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Buckwheat Spirals	2.00	250 g OP	
Corn and Vegetable Spirals (2.92) Orgran Rice and Corn Lasagne Sheets 1.60 200 g OP (3.82) Orgran Rice and Corn Macaroni 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals 2.00 250 g OP Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 250 g OP		` ,		Orgran
Corn and Vegetable Spirals. 2.00 250 g OP (2.92) Orgran Rice and Corn Lasagne Sheets. 1.60 200 g OP (3.82) Orgran Rice and Corn Macaroni. 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals. 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Corn and Vegetable Shells		250 g OP	
(2.92) Orgran		` ,		Orgran
Rice and Corn Lasagne Sheets. 1.60 200 g OP (3.82) Orgran Rice and Corn Macaroni. 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals. 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Corn and Vegetable Spirals		250 g OP	
Rice and Corn Macaroni		` ,		Orgran
Rice and Corn Macaroni 2.00 250 g OP (2.92) Orgran Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and Corn Lasagne Sheets		200 g OP	_
Rice and Corn Penne		\ /		Orgran
Rice and Corn Penne 2.00 250 g OP (2.92) Orgran Rice and Maize Pasta Spirals 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and Corn Macaroni		250 g OP	_
(2.92) Orgran		\ /		Orgran
Rice and Maize Pasta Spirals. 2.00 250 g OP (2.92) Orgran Rice and Millet Spirals. 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles. 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals. 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and Corn Penne		250 g OP	_
(2.92) Orgran	B	\ /		Orgran
Rice and Millet Spirals 2.00 250 g OP (3.11) Orgran Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and Maize Pasta Spirals		250 g OP	
(3.11) Orgran	T	, ,		Orgran
Rice and corn spaghetti noodles 2.00 375 g OP (2.92) Orgran Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and Millet Spirals		250 g OP	_
Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran (2.92) Orgran Italian long style spaghetti 2.00 220 g OP		\ /		Orgran
Vegetable and Rice Spirals 2.00 250 g OP (2.92) Orgran Italian long style spaghetti 2.00 220 g OP	Rice and corn spaghetti noodles		375 g OP	
(2.92) Orgran Italian long style spaghetti2.00 220 g OP		\ /		Orgran
Italian long style spaghetti	Vegetable and Rice Spirals		250 g OP	
		` ,		Orgran
(3.11) Orgran	Italian long style spaghetti		220 g OP	•
		(3.11)		Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

✓ fully subsidised 253

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer Supplements For Homocystinuria AMINOACID FORMULA WITHOUT METHIONINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 500 a OP ✓ XMET Maxamum Powder 461.94 Supplements For MSUD AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] ✓ MSUD Maxamum 500 q OP Supplements For PKU AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 75 OP ✓ Phlexv 10 30 PKU Anamix Junior Orange Powder (chocolate) 36 g sachet......393.00 ✓ PKU Anamix Junior 30 Chocolate Powder (unflavoured) 28 g sachets......936.00 30 ✓ PKU Lophlex Powder ✓ PKU Anamix Junior 30 ✓ PKU Anamix Junior Powder (vanilla) 36 g sachet393.00 30 Vanilla ✓ PKU Anamix Infant 400 g OP ✓ XP Maxamum 500 a OP 500 g OP ✓ XP Maxamum 125 ml OP ✓ PKU Anamix Junior LQ 125 ml OP ✓ PKU Anamix Junior ✓ PKU Anamix Junior 125 ml OP Liquid (forest berries), 250 ml carton......540.00 18 OP ✓ Easiphen Liquid Liquid (juicy tropical) 125 ml......936.00 ✓ PKU Lophlex LQ 20 30 OP

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Powder8.22 500 g OP ✓ Loprofin Mix

36 OP

60 OP

60 OP

60 OP

30 OP

30 OP

✓ PKU Lophlex Sensation 20

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 20

✓ PKU Lophlex LQ 20

Liquid (juicy berries) 62.5 ml......939.00

Liquid (juicy orange) 125 ml936.00

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
LOW PROTEIN PASTA - Special Authority see SA1108 on page	e 253 – Hospital	pharmacy [HP:	3]	
Animal shapes	11.91	500 g OP	✓ L	.oprofin
Lasagne	5.95	250 g OP	√ L	.oprofin
Low protein rice pasta	11.91	500 g OP	√ L	.oprofin
Macaroni	5.95	250 g OP	√ L	.oprofin
Penne	11.91	500 g OP	√ L	.oprofin
Spaghetti	11.91	500 g OP	√ L	.oprofin
Spirals	11.91	500 g OP	√ L	oprofin

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FORMULA – Special Authority see SA1940 below – Hospital phar Powder43.60	400 g OP	✓ Alfamino
Powder (unflavoured)53.00	400 g OP	✓ Alfamino Junior ✓ Elecare ✓ Elecare LCP
		✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	 ✓ Neocate SYNEC ✓ Elecare ✓ Neocate Junior ✓ Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or

continued...

✓ fully subsidised 255

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Por 🗸	Manufacturer	

- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	✓	Manufacturer	

- 2.2.3 Amino acid formula is required for a nutritional deficit; and
- 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis: or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Roth
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	- Special Authority see SA1953 below	- Hospital phan	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

continued...

✓ fully subsidised 257



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	lised	Generic
\$	Per	1	Manufacturer

- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

Powder	450 g OP	✓ Aptamil Gold+ Pepti Junior
30.42	900 g OP	 Aptamil AllerPro SYNEO 1
		✓ Aptamil AllerPro SYNEO 2

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Fither:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Sov milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

✓ fully subsidised 259

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Vaccinations** BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm] For infants at increased risk of tuberculosis. Increased risk is defined as: 1) living in a house or family with a person with current or past history of TB; or 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm] Funded for any of the following criteria: 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or 5) A single dose for vaccination of patients aged from 65 years old; or 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or 7) For vaccination of previously unimmunised or partially immunised patients: or 8) For revaccination following immunosuppression; or 9) For boosting of patients with tetanus-prone wounds. Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous 10 **Boostrix Boostrix** DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm] Funded for any of the following: 1) A single dose for children up to the age of 7 who have completed primary immunisation; or 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive 4) Five doses will be funded for children requiring solid organ transplantation. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

Infanrix IPV

Subsidised

Per

Fully

Brand or

Generic

✓ <u>Havrix</u>✓ Havrix Junior

Manufacturer

Subsidy

(Manufacturer's Price)

φ rei ♥ ivialiulaciulei
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE -
[Xpharm]
Funded for patients meeting any of the following criteria:
 Up to four doses for children up to and under the age of 10 for primary immunisation; or
2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of
10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or
post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.
Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years)
to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up
programmes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg
pertussis toxoid, 25 mcg pertussis filamentous
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,
10 mcg hepatitis B surface antigen in 0.5 ml syringe
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]
One dose for patients meeting any of the following:
1) For primary vaccination in children; or
2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell
transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre-
or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
3) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or
paediatrician.
p-0-0-0-0-0-0-0-0-0-0-0-0-0-0-0-0-0-0-0
Haemophilus Influenzae type B polysaccharide 10 mcg
conjugated to tetanus toxoid as carrier protein 20-40 mcg;
prefilled syringe plus vial 0.5 ml
promot of the transfer from the first transfer from th
HEPATITIS A VACCINE – [Xpharm]
Funded for patients meeting any of the following criteria:
Two vaccinations for use in transplant patients; or
Two vaccinations for use in children with chronic liver disease; or
One dose of vaccine for close contacts of known hepatitis A cases.

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Inj 10 mo	B RECOMBINANT VACCINE - [Xpharm] g per 0.5 ml prefilled syringe Jed for patients meeting any of the following criteria		1	√ E	ingerix-B
1) 2) 3) 4) 5) 6) 7) 8) 9)	for household or sexual contacts of known acute for children born to mothers who are hepatitis B s for children up to and under the age of 18 years in serology and require additional vaccination or rector HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual interfor patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC following needle stick injury.	hepatitis B patients or h urface antigen (HBsAg nclusive who are considuire a primary course of recourse; or) pos derec	sitive; or I not to have	
	cg per 1 ml prefilled syringeded for patients meeting any of the following criteria		1	√ <u>E</u>	ingerix-B
2) 3) 4) 5) 6) 7) 8) 9) 10) 11)	for household or sexual contacts of known acute for children born to mothers who are hepatitis B is for children up to and under the age of 18 years in serology and require additional vaccination or record for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual interfor patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC following needle stick injury; or for dialysis patients; or for liver or kidney transplant patients.	urface antigen (HBsAg nclusive who are consic juire a primary course o rcourse; or) pos derec	sitive; or I not to have	
Any of th	PILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND e following:	,	· [Xpl	harm]	
2) Ma 1 2	ximum of two doses for children aged 14 years and ximum of three doses for patients meeting any of to 1 yeople aged 15 to 26 years inclusive; or 1 yeither: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: of ximum of four doses for people aged 9 to 26 years	he following criteria: or	nerap	ру	
Inj 270 m	ncg in 0.5 ml syringe	0.00	10	√ <u>G</u>	ardasil 9

Subsidy (Manufacturer's Price)	,		Brand or Generic
 \$	Per	✓	Manufacturer

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

- [Xpharm]......9.00 1 ✓ Afluria Quad Junior (2021 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)		Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable

C)

A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine).......90.00 10 ✓ Fluad Quad (2021 Formulation)

Subsidy (Manufacturer's Price)			Brand or Generic
\$	Per	✓	Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

C)

- INFLUENZA VACCINE people 65 years and over
 is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine)	_		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturor

A) INFLUENZA VACCINE - people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	✓	

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

NATIONAL IMMUNISATION SCHEDULE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant: or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C 1 PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00

✓ Synflorix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Immunisation Handbook for the app	propriate schedule for catch up programmes
Inj 30.	8 mcg of pneumococcal polysaccharide serotypes 1, 3,	, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe	10	✓ Prevenar 13
	1	✓ Prevenar 13

Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. Inj 80D antigen units in 0.5 ml syringe			Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or 2) All of the following: a) Patient is a child under 18 years for (re-)immunisation; and b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with oral failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplator or vi) with corelar implants or intracranial shunts; or viii) with corebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xii with diabetes; or xiii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)		PV23) POLYSACCHARIDE VACCINE -	[Xpharm]			
a) Patient is a child under 18 years for (re-)immunisation; and b) Treatment is for a maximum of two doses; and c) Any of the following: i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transpla or vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	Up to three contemplement	py; pre- or post-splenectomy or with functi deficiency (acquired or inherited), cochle	onal asplenia, pre- or p	ost-so	lid organ t	ransplant, renal dialysis,
in mmunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or ii) with HIV infection; or iv) with primary immune deficiencies; or iii) with HIV infection; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transpla or vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	a) Patient b) Treatm	t is a child under 18 years for (re-)immunis nent is for a maximum of two doses; and	sation; and			
immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; or v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplator or vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with Down syndrome; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	, ,	•	n therapy, vaccinate wh	nen the	re is expe	cted to be a sufficient
iv) with renal failure, or nephrotic syndrome; or y) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplator vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with Down syndrome; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ir ii) v	mmune response; or vith primary immune deficiencies; or			. o io onpo	
vi) with cochlear implants or intracranial shunts; or vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	iv) w	vith renal failure, or nephrotic syndrome; o		uding h	naematopo	vietic stem cell transplant);
viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage 20 mg or greater; or ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	vi) v	vith cochlear implants or intracranial shunt	s; or			
ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	viii) r	eceiving corticosteroid therapy for more th				
x) pre term infants, born before 28 weeks gestation; or xi) with cardiac disease, with cyanosis or failure; or xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)		5 5 7	asthma treated with hig	ah-dose	e corticoste	eroid therapy); or
xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	x) p	re term infants, born before 28 weeks ges	tation; or	,		13/
xiv) who are pre-or post-splenectomy, or with functional asplenia. Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	xii) w	vith diabetes; or	, 01			
23 pneumococcal serotype)	,		unctional asplenia.			
POLIOMYELITIS VACCINE – [Xpharm] Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. Inj 80D antigen units in 0.5 ml syringe					4 -	
Up to three doses for patients meeting either of the following: 1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. Inj 80D antigen units in 0.5 ml syringe			0.00	1	▼ <u>P</u>	neumovax 23
1) For partially vaccinated or previously unvaccinated individuals; or 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. Inj 80D antigen units in 0.5 ml syringe			j :			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes. Inj 80D antigen units in 0.5 ml syringe	•		•			
Inj 80D antigen units in 0.5 ml syringe	,	•				
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 weeks of age; and 2) no vaccination being administered to children aged 24 weeks or over. Oral susp live attenuated human rotavirus					0	
Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 weeks of age; and 2) no vaccination being administered to children aged 24 weeks or over. Oral susp live attenuated human rotavirus	, ,	, ,		'	• 11	OL
no vaccination being administered to children aged 24 weeks or over. Oral susp live attenuated human rotavirus						
Oral susp live attenuated human rotavirus	,	•	•			
·	no vaccination	on being administered to children aged 24	weeks or over.			
1,000,000 CCID50 per dose, prefilled oral applicator	Oral susp live atte	nuated human rotavirus				
	1,000,000 CC	CID50 per dose, prefilled oral applicator	0.00	10	✓ <u>R</u>	<u>otarix</u>

	NATIONAL	IMMUNIS	SATIO	ON SCHEDULE
	Subsidy (Manufacturer's Price)	Subsid Per	Fully lised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:				
1) Maximum of one dose for primary vaccination for either	r:			
 a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 y varicella infection (chickenpox), or 	ears old on or after 1	July 2017, v	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
 a) Any of the following for non-immune patients: 				
i) with chronic liver disease who may in future ii) with deteriorating renal function before tran iii) prior to called a reach transplant or iii)		nsplantation	ı; or	
iii) prior to solid organ transplant; oriv) prior to any elective immunosuppression*,	or			
v) for poet exposure prophylaxic who are imm		nto · or		

- v) for post exposure prophylaxis who are immune competent inpatients.; or
- b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
- c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
- d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
- For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
- f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
- g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* immu	unosuppression due to	steroid or other imr	nunosuppressive th	herapy must be fo	or a treatment perio	od of greater than
28 day	'S					

Inj 1350 PFU prefilled syringe	0.00	1	✓ Varivax
		10	✓ Varivax

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting either of the following criteria:

- 1) One dose for all people aged 65 years; or
- 2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 December 2021.

Inj 19,400 PFU prefilled syringe plus vial	0.00	1	Zostavax
		10	✓ Zostavax

Diagnostic Agents

- Symbols -		Renin-Angiotensin System	47	AmsaLyo	14
UK Synacthen	79	Agents for Parkinsonism and		Amsidine	
3TC		Disorders		Amzoate	
- A -		Agents Used in the Treatmen		Anaesthetics	
A-Scabies	67	Poisonings		Anagrelide hydrochloride	14
Abacavir sulphate		Agrylin		Analgesics	11
Abacavir sulphate with		Albendazole		Anastrozole	
lamivudine	104	Albey		Anatrole	
Abiraterone acetate		Albustix		Andriol Testocaps	7
Acarbose	11	Aldurazyme		Androderm	
Accarb	11	Alecensa	156	ANI	
Accuretic	48	Alectinib	156	Anoro Ellipta	
Accuretic 10	48	Alendronate sodium	110	Antabuse	14
Accuretic 20	48	Alendronate sodium with		Antacids and Antiflatulents	
Acetazolamide	232	colecalciferol	110	Anthelmintics	8
Acetec	47	Alfacalcidol	33	Antiacne Preparations	
Acetic acid with 1, 2- propanediol		Alfamino	255	Antiallergy Preparations	22
diacetate and		Alfamino Junior		Antianaemics	
benzethonium	229	Alginic acid	6	Antiandrogen Oral	
Acetic acid with hydroxyquinoline	and	Alglucosidase alfa	26	Contraceptives	7
ricinoleic acid	74	Alkeran	146	Antiarrhythmics	4
Acetylcysteine		Alkeran S29		Antibacterials	8
Aci-Jel		Allersoothe	221	Antibacterials Topical	6
Aciclovir		Allmercap	148	Anticholinergic Agents	
Infection	100	Allopurinol	115	Anticholinesterases	
Sensory	229	Alpha-Adrenoceptor Blockers	47	Antidepressants	12
Acidex	6	Alpha-Keri Lotion	65	Antidiarrhoeals	
Acipimox	54	Alphamox	91	Antiepilepsy Drugs	
Acitretin	67	Alphamox 125	91	Antifibrinolytics, Haemostatics and	1
Aclasta	113	Alphamox 250	91	Local Sclerosants	3
Aclin	109	Alprolix	38	Antifibrotics	
Actemra	209	Alu-Tab		Antifungals	9
Actinomycin D	150	Aluminium hydroxide	6	Antifungals Topical	6
Actrapid	10	Alvogen	51	Antihistamines	
Actrapid Penfill		Amantadine hydrochloride		Antihypotensives	
Acular		Ambrisentan		Antimalarials	
Acupan		Ambrisentan Mylan		Antimigraine Preparations	12
Adalimumab		Amiloride hydrochloride	53	Antinausea and Vertigo Agents	
Adapalene		Amiloride hydrochloride with		Antipruritic Preparations	
Adcortyl		furosemide	54	Antipsychotics	
Adenuric		Amiloride hydrochloride with		Antiretrovirals	
ADR Cartridge 1.8		hydrochlorothiazide		Antirheumatoid Agents	
Adrenaline		Aminophylline		Antispasmodics and Other Agents	
Advantan		Amiodarone hydrochloride		Altering Gut Motility	
Advate		Amisulpride		Antithrombotic Agents	4
Adynovate		Amisulpride Mylan		Antithymocyte globulin	
Afinitor		Amitriptyline		(equine)	
Aflibercept	183	Amlodipine		Antitrichomonal Agents	9
Afluria Quad	004	Amneal		Antituberculotics and	_
(2021 Formulation)	264	Amorolfine		Antilleprotics	
Afluria Quad Junior	000	Amoxicillin		Antiulcerants	
(2021 Formulation)		Amoxicillin with clavulanic aci		Antivirals	
AFT-Pyrazinamide	99	Amphotericin B		Anxiolytics	
Agents Affecting the		Amsacrine	149	Anzatax	15

Apidra	11	Arrow-Roxithromycin	91	Basic AquaCream	6
Apidra SoloStar		Arrow-Sertraline		BCG Vaccine	
Apo-Azithromycin		Arrow-Timolol		Beclazone 100	
Apo-Bromocriptine		Arrow-Topiramate		Beclazone 250	
Apo-Ciclopirox		Arrow-Tramadol		Beclazone 50	
Apo-Clarithromycin		Arsenic trioxide		Beclomethasone dipropionate	
Alimentary	Q	Asacol		Bee venom allergy treatment	
Infection		Asamax		Bendamustine hydrochloride	
Apo-Clomipramine		Ascorbic acid		Bendrofluazide	
		Aspen Adrenaline		Bendroflumethiazide	
Apo-Diclo SR			50	[Bendrofluazide]	5.
Apo-Diltiazem CD		Aspirin	44		
Apo-Doxazosin		Blood		Benzathine benzylpenicillin	
Apo-Folic Acid		Nervous		Benzatropine mesylate	
Apo-Furosemide		Asthalin		Benzbromaron AL 100	
Apo-Gabapentin		Atazanavir sulphate		Benzbromarone	
Apo-Megestrol		Atenolol		Benztrop	
Apo-Metoprolol		Atenolol AFT		Benzydamine hydrochloride	3
Apo-Mirtazapine		Atenolol AFT S29		Benzylpenicillin sodium [Penicillin	
Apo-Nadolol		ATGAM		G]	
Apo-Oxybutynin		Ativan		Beta Cream	
Apo-Perindopril		Atomoxetine		Beta Ointment	
Apo-Pindolol		Atorvastatin	55	Beta Scalp	
Apo-Prazosin		Atropine sulphate		Beta-Adrenoceptor Agonists	
Apo-Prednisone		Cardiovascular		Beta-Adrenoceptor Blockers	
Apo-Primidone	127	Sensory	233	Betadine	
Apo-Propranolol		Atropt	233	Betadine Skin Prep	
Apo-Pyridoxine		Atrovent	223	Betaferon	
Apo-Selegiline S29	117	AU Synacthen	79	Betahistine dihydrochloride	12
Apo-Sumatriptan		Aubagio	135	Betaine	2
Apo-Temozolomide	154	Augmentin	91	Betaloc CR	5
Apomorphine hydrochloride	117	Aurorix	124	Betamethasone dipropionate	6
Aprepitant	129	AutoSoft 30	23	Betamethasone dipropionate with	
Apresoline	56	AutoSoft 90	23	calcipotriol	6
Aptamil AllerPro SYNEO 1	258	Avelox	93	Betamethasone sodium phosphate	
Aptamil AllerPro SYNEO 2	258	Avonex	135	with betamethasone acetate	78
Aptamil Gold+ Pepti Junior	258	Avonex Pen	135	Betamethasone valerate6	2, 6
Aqueous cream		Azacitidine		Betamethasone valerate with sodiu	m
Aratac		Azacitidine Dr Reddy's	146	fusidate [fusidic acid]	6
Arava	110	Azamun		Betaxolol	
Arginine	27	Azathioprine		Betnovate	6
Aripiprazole		Azilect		Betoptic	
Aripiprazole Sandoz		Azithromycin		Betoptic S	
Aristocort		Azopt	232	Bexsero	
Arrow - Lattim		AZT		Bezafibrate	
Arrow-Amitriptyline		- B -		Bezalip	
Arrow-Bendrofluazide		B-D Micro-Fine	15	Bezalip Retard	
Arrow-Brimonidine		B-D Ultra Fine		Bicalutamide	
Arrow-Diazepam		B-D Ultra Fine II		Bicillin LA	
Arrow-Doxorubicin		Bacillus Calmette-Guerin (BCG)		BiCNU	
Arrow-Losartan &	100	vaccine	173	Bicnu Heritage	
Hydrochlorothiazide	48	Bacillus Calmette-Guerin	170	Bile and Liver Therapy	
Arrow-Norfloxacin		vaccine	260	Biltricide	
Arrow-Ornidazole		Baclofen			
Arrow-Quinapril 10		Bactroban		BimatoprostBimatoprost Multichem	<u>८</u> ८१
		Balance		Binarex	
Arrow Quinapril 5		Barrier Creams and Emollients		Binocrit	
Arrow-Quinapril 5	40	Daniel Cleans and Emonients	04	DITIOGRAM	ن ا

Biodone	121	Calcipotriol	67	Celebrex	11
Biodone Extra Forte		Calcitonin		Celecoxib	
Biodone Forte	121	Calcitriol	33	Celecoxib Pfizer	11
Bisacodyl	26	Calcitriol-AFT	33	Celestone Chronodose	7
Bisoprolol fumarate		Calcium carbonate	6, 34	Cellcept	16
Bisoprolol Mylan		Calcium Channel Blockers	<mark>52</mark>	Centrally-Acting Agents	5
BK Lotion		Calcium Disodium Versenate	236	Cephalexin ABM	
Bleomycin sulphate		Calcium folinate	147	Cetirizine hydrochloride	
Blood Colony-stimulating		Calcium Folinate Ebewe	147	Cetomacrogol	
Factors	44	Calcium Folinate Sandoz	147	Cetomacrogol with glycerol	6
Blood glucose diagnostic test		Calcium Folinate Sandoz S29	9 147	Cetuximab	
meter	14	Calcium gluconate	34	Champix	14
Blood glucose diagnostic test		Calcium Homeostasis	77	Charcoal	23
strip	14	Calcium polystyrene sulphona	ate45	Chemotherapeutic Agents	14
Blood glucose test strips (visually	y	Calcium Resonium		Chickenpox vaccine	
impaired)	14	Calogen	242	Chlorafast	
Blood Ketone Diagnostic Test		Calutide-50	163	Chlorambucil	14
Strip	13	Candesartan cilexetil	48	Chloramphenicol	22
Bonjela	32	Candestar	48	Chlorothiazide	
Boostrix	260	Canesten	61	Chlorpheniramine maleate	22
Bortezomib	150	Capecitabine	147	Chlorpromazine hydrochloride	13
Bortezomib Dr-Reddy's	150	Capercit	147	Chlortalidone [Chlorthalidone]	
Bosentan	57	Capoten		Chlorthalidone	
Bosentan Dr Reddy's	57	Capsaicin		Chlorvescent	4
Bplex		Musculoskeletal	110	Choice Load 375	
Breo Ellipta		Nervous	119	Choice TT380 Short	<mark>7</mark>
Brevinor 1/28	73	Captopril	47	Choice TT380 Standard	
Bricanyl Turbuhaler		Captopril-Mylan	47	Choline salicylate with cetalkonius	m
Brilinta		Carafate		chloride	
Brimonidine tartrate	232	Carbaccord	145	Ciclopirox olamine	6
Brimonidine tartrate with timolol		Carbamazepine	126	Ciclosporin	
maleate	232	Carbimazole	82	Cilazapril	4
Brinzolamide	232	Carbomer	233	Cilicaine	9
Brolene	229	Carboplatin	145	Cilicaine VK	
Bromocriptine mesylate	117	Carboplatin Ebewe	145	Cinacalcet	7
Brufen SR	109	Carbosorb-X	235	Cipflox	9
Buccastem	130	Cardinol LA	51	Ciprofloxacin	
Budesonide		CareSens Dual	13	Infection	9
Alimentary	6	CareSens N	14	Sensory	22
Respiratory2	221, 227	CareSens N POP	14	Ciprofloxacin Teva	
Budesonide with eformoterol	222	CareSens N Premier	14	Circadin	13
Bumetanide	53	CareSens PRO	14	Cisplatin	14
Buprenorphine Naloxone BNM	140	Carmellose sodium with gelat	tin and	Cisplatin Ebewe	
Buprenorphine with naloxone	140	pectin	32	Citalopram hydrobromide	
Bupropion hydrochloride		Carmustine	145	Cladribine	
Burinex	53	Carvedilol	50	Clarithromycin	
Burinex S29	53	Carvedilol Sandoz	50	Alimentary	
Buscopan	8	Catapres		Infection	8
Buspirone hydrochloride		CeeNU		Clexane	4
Busulfan		Cefaclor monohydrate		Clexane Forte	4
- C -		Cefalexin		Climara	
Cabergoline	87	Cefalexin Sandoz	88	Climara	8
Cacit	34	Cefazolin	88	Clindamycin	9
Caffeine citrate		Ceftriaxone	88	Clinicians	2
Calamine		Ceftriaxone-AFT		Clinicians Renal Vit	
Calci-Tab 500		Cefuroxime axetil		Clobazam	12

Clobetasol propionate62, 68	for Systemic Use	78	DBL Aminophylline	226
Clobetasone butyrate63	Corticosteroids Topical		DBL Bleomycin Sulfate	150
Clofazimine98	Cortifoam	7	DBL Carboplatin	145
Clomazol	Cosentyx	207	DBL Cisplatin	145
Dermatological61	Cosmegen		DBL Dacarbazine	150
Genito-Urinary74	Coumadin	44	DBL Desferrioxamine Mesylate for	or Inj
Clomifene citrate87	Country Life	29	BP	236
Clomipramine hydrochloride123	Coversyl		DBL Docetaxel	150
Clonazepam 125–126, 133	Creon 10000	24	DBL Ergometrine	74
Clonidine52	Creon 25000		DBL Gemcitabine	147
Clonidine BNM53	Creon Micro	24	DBL Gentamicin	
Clonidine hydrochloride53	Crotamiton	62	DBL Heparin Sodium	
Clopidogrel41	Crystaderm	60	DBL Leucovorin Calcium	147
Clopidogrel Multichem41	Curam		DBL Methotrexate Onco-Vial	148
Clopine131	Curam Duo 500/125	91	DBL Morphine Sulphate	122
Clopixol132–133	Cvite	33	DBL Naloxone Hydrochloride	
Clotrimazole	Cyclizine hydrochloride	129	DBL Octreotide	164
Dermatological61	Cyclizine lactate	129	DBL Pethidine Hydrochloride	123
Genito-Urinary74	Cyclogyl		DBL Vinblastine	
Clozapine131	Cyclonex	145	DBL Vincristine Sulfate	156
Clozaril131	Cyclopentolate hydrochloride	233	Decozol	32
Co-trimoxazole95	Cyclophosphamide		Deferasirox	235
Coal tar68	Cyclorin		Deferiprone	235
Coal tar with allantoin, menthol,	Cycloserine		Denosumab	110
phenol and sulphur68	Cyproterone acetate		Deolate	97
Coal tar with salicylic acid and	Cyproterone acetate with		Deoxycoformycin	153
sulphur68	ethinyloestradiol	74	Depo-Medrol	78
Coco-Scalp68	Cystadane	27	Depo-Provera	74
Codeine phosphate	Cytarabine		Depo-Testosterone	
Extemporaneous238	Cytotec		Deprim	
Nervous121	Cytoxan		Dermol	
Coenzyme Q1028	- D -		Desferrioxamine mesilate	
Colchicine	D-Penamine	110	Desmopressin	86
Colecalciferol33	Dabigatran		Desmopressin acetate	
Colestid54	Dacarbazine		Desmopressin-PH&T	
Colestipol hydrochloride54	Dacarbazine APP		Desuric	
Colgout115	Dactinomycin [Actinomycin D]		Detection of Substances in	
Colifoam7	Daivobet		Urine	76
Colistin sulphomethate93	Daivonex	67	Dexamethasone	
Colistin-Link93	Daktarin	62	Hormone	78
Collodion flexible238	Dalacin C	93	Sensory	
Colloidal bismuth subcitrate10	Dantrium		Dexamethasone phosphate	
Colofac8	Dantrium S29		Dexamethasone Phosphate	
Coloxyl25	Dantrolene		Panpharma	78
Combigan232	Daonil	11	Dexamethasone with framycetin	
Compound electrolytes46	Dapa-Tabs		gramicidin	
Compound electrolytes with glucose	Dapsone		Dexamethasone with neomycin	
[Dextrose]46	Daraprim		sulphate and polymyxin B	
Compound hydroxybenzoate238	Darunavir		sulphate	230
Concerta	Darunavir Mylan		Dexamfetamine sulfate	
Condoms	Dasatinib		Dexmethsone	
Condyline69	Daunorubicin		Dextrochlorpheniramine	
Contraceptives - Hormonal72	David One Step Cassette Pregr		maleate	221
Contraceptives - Non-hormonal70	Test		Dextrose	
Copaxone	DBL Acetylcysteine		DHC Continus	
Corticosteroids and Related Agents	DBL Adrenaline		Diabetes	

			_
Diabetes Management13	hydrochloride123	Empagliflozin with metformin	
Diacomit127	Dosulepin Mylan123	hydrochloride	1
Diagnostic Agents271	Dothiepin123	Emtricitabine1	10
Diamide Relief6	Doxazosin47	Emtricitabine with tenofovir	
Diamox232	Doxine92	disoproxil1	10
Diasip243	Doxorubicin Ebewe150	Emtriva1	10
Diason RTH243	Doxorubicin hydrochloride150	Emulsifying ointment	
Diazepam 125, 133	Doxycycline92	Emulsifying Ointment ADE	6
Diazoxide10	DP Lotion65	Enalapril maleate	4
Dibenzyline47	DP Lotn HC63	Enbrel1	16
Diclofenac Sandoz109	DP-Allopurinol115	Endocrine Therapy1	16
Diclofenac sodium	Dr Reddy's Omeprazole9	Endoxan1	14
Musculoskeletal109	Drugs Affecting Bone	Engerix-B2	26
Sensory230	Metabolism 110	Enlafax XR1	12
Differin60	Dual blood glucose and blood ketone	Enoxaparin sodium	4
Difflam32	diagnostic test meter 13	Enstilar	
Diflucan95	Dulcolax SP Drop26	Ensure2	
Digestives Including Enzymes24	Duocal Super Soluble Powder241	Ensure Plus2	25
Digoxin49	Duolin	Ensure Plus HN2	
Dihydrocodeine tartrate121	Duolin HFA223	Ensure Plus RTH2	
Dilantin127	DuoResp Spiromax222	Ensure Two Cal HN RTH	
Dilantin Infatab127	Duride56	Entacapone1	
Diltiazem hydrochloride52	-E-	Entapone1	11
Dilzem52	e-chamber La Grande228	Entecavir	
Dimethicone64–65	e-chamber Mask227	Entecavir Sandoz	
Dimethyl fumarate135	e-chamber Turbo228	Entocort CIR	
Dipentum	E-Mycin90	Entresto 24/26	
Diphtheria, tetanus and pertussis	Ear Preparations229	Entresto 49/51	
vaccine260	Ear/Eye Preparations229	Entresto 97/103	
Diphtheria, tetanus, pertussis and	Easiphen Liquid254	Epilim1	
polio vaccine	Econazole nitrate 61	Epilim Crushable1	12
Diphtheria, tetanus, pertussis, polio,	Efavirenz104	Epilim IV1	12
hepatitis B and haemophilus	Efavirenz with emtricitabine and	Epilim S/F Liquid1	
influenzae type B vaccine 261	tenofovir disoproxil	Epilim Syrup1	
	Eformoterol fumarate222	Epirubicin Ebewe1	
Diprosone			
Diprosone OV	Eformoterol fumarate dihydrate222	Epirubicin hydrochloride1	
Dipyridamole41 Disopyramide phosphate49	Eftrenonacog alfa [Recombinant	Eplerenone Epoetin alfa	
Disulfiram141	factor IX]		
Diuretics53	Efudix	Epoprostenol	. ၁
	Egopsoryl TA68	Eptacog alfa [Recombinant factor VIIa]	4
Docetaxel	Elaprase		
	Elecare	ERA	
Docetaxel Sandoz	Elecare LCP255	Erbitux	
Docusate sodium	Electral 46	Ergometrine maleate	
Docusate sodium with	Elelyso	Erlotinib1	
sennosides	Elemental 028 Extra246	Erythrocin IV	
Dolutegravir	Elidel68	Erythromycin (as lactobionate)	
Domperidone	Elocon	Erythromycin ethyl succinate	.9
Donepezil hydrochloride	Elocon Alcohol Free	Erythromycin stearate	.9
Donepezil-Rex139	Eltrombopag38	Esbriet	
Dornase alfa226	Eltroxin82	Escitalopram1	
Dortimopt232	EMB Fatol98	Escitalopram (Ethics)1	12
Dorzolamide hydrochloride232	Emend Tri-Pack129	Escitalopram-Apotex1	
Dorzolamide with timolol232	Emicizumab39	Eskazole	
Dostinex87	EMLA119	Essential Prednisolone	
Dosulepin [Dothiepin]	Empagliflozin12	Estradiol TDP Mylan	.8

Estradot	80	Ferro-tab	35	Foban	6
Estradot 50 mcg	80	Ferrograd	35	Folic acid	
Estrofem	80	Ferrosig	35	Folic Acid Mylan	3
Etanercept	166	Ferrous fumarate		Food Thickeners	25
Ethambutol hydrochloride	98	Ferrous fumarate with folic acid	35	Foods And Supplements For Inb	orn
Ethics Aspirin	119	Ferrous sulfate	35	Errors Of Metabolism	25
Ethics Aspirin EC	41	Fexofenadine hydrochloride	221	Foradil	22
Ethics Lisinopril	47	Fibro-vein	41	Forteo	112
Ethics Paracetamol Classic	120	Filgrastim		Fortini	24
Ethinyloestradiol	81	Finasteride	75	Fortini Multi Fibre	
Ethinyloestradiol with		Fingolimod	135	Fortisip	
desogestrel	72	Firazyr		Fortisip Multi Fibre	25
Ethinyloestradiol with		Flagyl	98	Fosamax	
levonorgestrel	73	Flagyl-S	98	Fosamax Plus	<mark>11</mark> 0
Ethinyloestradiol with		Flamazine	61	Framycetin sulphate	22
norethisterone	73	Flecainide acetate	49	Frisium	
Ethosuximide	126	Flecainide BNM	49	Frumil	
Etopophos	151	Flecainide Controlled Release		Frusemide	5
Etoposide	151	Teva	49	Fucicort	6
Etoposide phosphate	151	Fleet Phosphate Enema	26	Fucidin	94
Etravirine	104	Flixonase Hayfever & Allergy	227	Fucithalmic	22
Eumovate	63	Flixotide	222	Fulvestrant	
Everet	127	Flixotide Accuhaler	222	Fungilin	
Everolimus	217	Florinef	78	Furosemid-Ratiopharm	5
Evista		Fluad Quad		Furosemide [Frusemide]	5
Exemestane	166	(2021 Formulation)	264	Furosemide-Baxter	5
Exjade	235	Fluanxol	132	fusidic acid	
Extemporaneously Compound	ed	Flucil		Dermatological	
Preparations and		Flucloxacillin	92	Infection	94
Galenicals		Flucloxin		Sensory	22
Eye Preparations		Flucon	231	- G -	
Eylea		Fluconazole	95	Gabapentin	120
Ezetimibe		Fludara Oral	147	Gacet	
Ezetimibe Sandoz		Fludarabine Ebewe		Galsulfase	
Ezetimibe with simvastatin	55	Fludarabine phosphate	147	Galvumet	
-F-		Fludrocortisone acetate		Galvus	1
Factor eight inhibitor bypassing		Fluids and Electrolytes		Gardasil 9	
fraction		Flumetasone pivalate	229	Gastrodenol	
Famotidine		Fluocortolone caproate with		Gaviscon Double Strength	
Famotidine Hovid		fluocortolone pivalate and		Gaviscon Infant	
Faslodex		cinchocaine		Gazyva	
Febuxostat		Fluorometholone		Gefitinib	
Febuxostat multichem	115	Fluorouracil		Gemcitabine Ebewe	
Feed Thickener Karicare		Fluorouracil Ebewe		Gemcitabine hydrochloride	
Aptamil		Fluorouracil sodium		Genoptic	22
FEIBA NF		Fluox		Gentamicin sulphate	
Felo 10 ER		Fluoxetine hydrochloride		Infection	
Felo 5 ER		Flupenthixol decanoate		Sensory	22
Felodipine		Flutamide		Gilenya	13
Femme-Tab ED		Flutamin		Ginet	
Fentanyl		Fluticasone	222	Glatiramer acetate	
Fentanyl Sandoz		Fluticasone furoate with		Glecaprevir with pibrentasvir	
Ferinject		vilanterol		Glibenclamide	
Ferodan		Fluticasone propionate		Gliclazide	
Ferriprox		Fluticasone with salmeterol		Glipizide	
Ferro-F-Tabs	35	FML	231	Glivec	15

Glizide	12	Hiberix26	-1-	
Glucagen Hypokit	10	Hiprex10	B Ibiamox	91
Glucagon hydrochloride	10	Histaclear22	Ibrance	160
Glucerna Select	243	Histafen22	Ibuprofen	109
Glucerna Select RTH		Holoxan14	Ibuprofen SR BNM	109
Glucobay	11	Horleys Bread Mix25		
Glucose [Dextrose]		Horleys Flour25	Idarubicin hydrochloride	151
Gluten Free Foods		Hormone Replacement Therapy -	Idursulfase	28
Glycerin with sodium sacchar		Systemic		
Glycerin with sucrose		HPV26		
Glycerol		Humalog1		
Alimentary	<mark>26</mark>	Humalog Mix 251	llevro	231
Extemporaneous	238	Humalog Mix 501		59
Glyceryl trinitrate		Human papillomavirus (6, 11, 16, 18,	Imatinib mesilate	158
Alimentary	8	31, 33, 45, 52 and 58) vaccine	Imatinib-Rex	158
Cardiovascular		[HPV] 26		
Glycopyrronium		Humatin		
Glycopyrronium bromide		Humira17		
Glycopyrronium with	•	HumiraPen17		
indacaterol	224	Humulin 30/701		
Go Healthy		Humulin NPH1	• • • • • • • • • • • • • • • • • • • •	
Gold Knight		Humulin R		
Gold Knight XL		Hyaluronic acid23		
Goserelin		Hydralazine		
Gutron		Hydralazine hydrochloride		
Gynaecological Anti-infective		Hydrocortisone	Infanrix-hexa	
- H -	37	Dermatological		
Habitrol	1/12	Hormone		
Haemophilus influenzae type		Hydrocortisone (PSM)		
vaccine		Hydrocortisone acetate		
Haldol		Hydrocortisone acetate with	Influvac Tetra	200
Haldol Concentrate		pramoxine hydrochloride		265
Haldol Decanoas		Hydrocortisone and paraffin liquid	Inhaled Corticosteroids	
Haloperidol		and lanolin		221
Haloperidol decanoate		Hydrocortisone butyrate		222
Harvoni		Hydrocortisone with cinchocaine		
Havrix		Hydrocortisone with miconazole		
Havrix Junior		Hydrocortisone with natamycin and		
			Insulin aspart with insulin cone	
Haylor syrup		neomycin		
healthE Calamine Aqueous C		Hydrogen peroxide		
BP		Hydroxocobalamin		
healthE Dimethicone 10%		Hydroxocobalamin Mercury	Insulin glulisine	
healthE Dimethicone 4% Lotio		Pharma		11
healthE Dimethicone 5%		hydroxycarbamide15		44
healthE Glycerol BP		Hydroxychloroquine11		
healthE Urea Cream		Hydroxyurea	Insulin lispro	
Healtheries Simple Baking Mi		[hydroxycarbamide] 15		
Hemastix		Hygroton		
Hemlibra		Hylo-Fresh23		
Heparin sodium		Hymenoptera22		
Heparinised saline		Hyoscine butylbromide		
Heparon Junior		Hyoscine hydrobromide12		
Hepatitis A vaccine	261	Hypam13		
Hepatitis B recombinant		Hyperuricaemia and Antigout11		
vaccine		Hypromellose23		
Herceptin	212	Hypromellose with dextran23	cannula, straight insertion)	21

Insulin pump infusion set (teflon		Jevity RTH	250	Latanoprost with timolol	232
cannula)	22	Juno Pemetrexed	148	Lax-Suppositories	26
Insulin pump infusion set (teflon		- K -		Lax-Tab	
cannula, angle insertion with		Kadcyla	214	Laxatives	
insertion device)	. 23	Kaletra	105	Laxsol	25
Insulin pump infusion set (teflon		Kalydeco	227	Ledipasvir with sofosbuvir	10 1
cannula, angle insertion)	. 23	Kemadrin		Leflunomide	
Insulin pump infusion set (teflon		Kenacomb	229	Lenalidomide	
cannula, straight insertion with		Kenacort-A 10	79	Letrole	166
insertion device)	. 23	Kenacort-A 40	79	Letrozole	166
Insulin pump infusion set (teflon		Kenalog		Leukeran FC	
cannula, straight insertion)	. 23	Kenalog in Orabase		Leukotriene Receptor	
Insulin pump reservoir		Kenkay Sorbolene		Antagonists	226
Insulin syringes, disposable with		Ketocal 3:1		Leuprorelin	
attached needle	15	KetoCal 4:1		Leustatin	
Intal Forte CFC Free		Ketoconazole		Levetiracetam	
Intelence		Dermatological	69	Levetiracetam-AFT	
Interferon beta-1-alpha		Infection		Levlen ED	
Interferon beta-1-beta		Ketogenic Diet		Levocabastine	231
Intra-uterine device		Ketoprofen		Levocarnitine	
Invega Sustenna		Ketorolac trometamol		Levodopa with benserazide	
IPOL		KetoSens		Levodopa with carbidopa	
Ipratropium bromide223,		Keytruda		Levomepromazine	
Iressa		Kindergen		Levomepromazine	
Irinotecan Accord		Kivexa		hydrochloride	131
Irinotecan Actavis 100		Klacid		Levonorgestrel	
Irinotecan hydrochloride		Kliogest		Genito-Urinary	73_7/
Irinotecan-Rex		Kliovance		Hormone	
Iron (as ferric carboxymaltose)		Kogenate FS		Levothyroxine	
Iron polymaltose		Konakion MM		Lidocaine [Lignocaine]	
Isentress		Konsyl-D		Lidocaine [Lignocaine]	110 110
Isentress HD		Kuvan		hydrochloride	110
Ismo 20		- L -	20	Lidocaine [Lignocaine] with	
Ismo 40 Retard		Labetalol	51	chlorhexidine	110
Isoniazid		Lacosamide		Lidocaine [Lignocaine] with	116
Isoniazid with rifampicin		Lactulose		prilocaine	110
· · · · · · · · · · · · · · · · · · ·		Laevolac		Lidocaine-Baxter	
IsoptinIsoptin Retard		Lamictal		Lidocaine-Claris	
Isoptin SR		Lamivudine Lamivudine Alphapharm.		Life Extension Lignocaine	
Isopto CarpineIsosorbide mononitrate	56	Lamotrigine		Lioresal Intrathecal	110-118
Isosource Standard		Lamprene		Lipid-Modifying Agents	110
		Lanoxin		Liquigen	040
Isotretinoinbusk		Lanoxin Paediatric Elixir			
Ispaghula (psyllium) husk				Lisinopril Lithium carbonate	41
Itch-Soothe		Lanoxin PG		Livostin	
Itraconazole		Lanoxin S29			
Itrazole		Lansoprazole		LMX4 Locacorten-Viaform ED's	
lvacaftor		Lantus Colo Stor			
Ivermectin	65	Lantus SoloStar		Local preparations for Anal and	
- J - Jadelle	70	Lanvis		Rectal Disorders	
		Lanzol Relief		Locasol	
Jakavi		Lapatinib ditosylate		Locoid	
Jardiamet		Largactil		Locoid Crelo	
Jardiance		Laronidase		Locoid Lipocream	
Jaydess	ช1	Lasix		Locorten-Vioform	
Jevity HiCal RTH	250	Latanoprost	232	Lodoxamide	231

Logem	126	Medco	Methylprednisolone aceponate	6
Lomide	231	Dermatological64	Methylprednisolone acetate	7
Lomustine	146	Nervous120	Methylxanthines	22
Loniten	57	Medrol78	Metoclopramide Actavis 10	13
Loperamide hydrochloride	6	Medroxyprogesterone acetate	Metoclopramide hydrochloride	13
Lopinavir with ritonavir	105	Genito-Urinary74	Metolazone	5
Loprofin	255	Hormone80–81	Metopirone	8
Loprofin Mix	254	Mefenamic acid109	Metoprolol IV Mylan	5
Lorafix		Megestrol acetate164	Metoprolol succinate	5
Loratadine	221	Melatonin135	Metoprolol tartrate	
Lorazepam	134	Melphalan146	Metrogyl	9
Lorfast	221	Menactra267	Metronidazole	
Lorstat		Meningococcal (groups A, C, Y and	Metyrapone	
Losartan Actavis		W-135) conjugate vaccine 267	Mexiletine hydrochloride	
Losartan potassium	48	Meningococcal B multicomponent	Mexiletine Hydrochloride USP	5
Losartan potassium with		vaccine	Miacalcic	
hydrochlorothiazide	48	Meningococcal C conjugate	Micolette	
Lovir	100	vaccine	Miconazole	3
Loxamine	124	Menthol62	Miconazole nitrate	
Lucrin Depot 1-month	86	Mepolizumab192	Dermatological	6
Lucrin Depot 3-month	86	Mercaptopurine148	Genito-Urinary	7
Lyderm		Mercilon 2872	Micreme	
Lynparza	152	Mesalazine7	Micreme H	
Lyrica	127	Mesna152	Microgynon 20 ED	7
- M -		Mestinon109	Microgynon 30	
m-Eslon	122	Metabolic Disorder Agents26	Microgynon 50 ED	
Mabthera	195	Metformin hydrochloride12	Microlut	
Macrobid	108	Methadone hydrochloride	Midazolam	13
Macrogol 3350 with potassium		Extemporaneous238	Midazolam-Baxter	13
chloride, sodium bicarbonate		Nervous121	Midodrine	
sodium chloride	26	Methatabs121	Mifegyne	
Macrogol 400 and propylene		Methenamine (hexamine)	Mifepristone	
glycol	233	hippurate108	Minerals	3
Madopar 125		Methopt233	Mini-Wright AFS Low Range	
Madopar 250	117	Methotrexate148	Mini-Wright Standard	22
Madopar 62.5		Methotrexate DBL Onco-Vial148	Minidiab	1
Madopar HBS		Methotrexate Ebewe148	MiniMed 1.8 Reservoir	
Madopar Rapid		Methotrexate Sandoz148	MMT-326A	2
Magnesium hydroxide		Methyl hydroxybenzoate238	MiniMed 3.0 Reservoir	
Magnesium sulphate	36	Methylcellulose238	MMT-332A	2
Mantoux		Methylcellulose with glycerin and	MiniMed 640G	
Marevan		sodium saccharin238	MiniMed Mio MMT-921A	
Marine Blue Lotion SPF 50+	69	Methylcellulose with glycerin and	MiniMed Mio MMT-923A	
Martindale Pharma		sucrose	MiniMed Mio MMT-925A	
Marvelon 28		Methyldopa53	MiniMed Mio MMT-941A	
Mask for spacer device		Methyldopa Mylan53	MiniMed Mio MMT-943A	
Mast Cell Stabilisers	226	Methyldopa Mylan S2953	MiniMed Mio MMT-945A	2
Maviret	101	Methylnaltrexone bromide25	MiniMed Mio MMT-965A	2
Maxidex		Methylphenidate ER - Teva137	MiniMed Mio MMT-975A	
Maxitrol		Methylphenidate hydrochloride 137	MiniMed Quick-Set MMT-386A	2
MCT oil (Nutricia)	242	Methylphenidate hydrochloride	MiniMed Quick-Set MMT-387A	2
Measles, mumps and rubella		extended-release138	MiniMed Quick-Set MMT-396A	
vaccine		Methylprednisolone78	MiniMed Quick-Set MMT-397A	
Mebendazole		Methylprednisolone (as sodium	MiniMed Quick-Set MMT-398A	
Mebeverine hydrochloride	8	succinate) 78	MiniMed Quick-Set MMT-399A	
			MiniMed Silhouette MMT-368A	2

MiniMed Silhouette MMT-377A		Mupirocin	61	NeuroTabs	
MiniMed Silhouette MMT-378A		Muscle Relaxants		Nevirapine	
MiniMed Silhouette MMT-381A		Mvite		Nevirapine Alphapharm	104
MiniMed Silhouette MMT-382A	22	Myambutol		Nicorandil	
MiniMed Silhouette MMT-383A	22	Mycobutin		Nicotine	
MiniMed Silhouette MMT-384A	22	MycoNail	61	Nifedipine	
MiniMed Sure-T MMT-864A	20	Mycophenolate mofetil	166	Nifuran	10
MiniMed Sure-T MMT-866A	20	Mydriacyl	233	Nilotinib	159
MiniMed Sure-T MMT-874A	20	Mylan (12 hr release)		Nilstat	
MiniMed Sure-T MMT-876A	20	Mylan (24 hr release)	52	Alimentary	3
MiniMed Sure-T MMT-884A	20	Mylan Atenolol		Genito-Urinary	
MiniMed Sure-T MMT-886A	20	Mylan Clomiphen		Infection	9
Minims Cyclopentolate	233	Mylan Indapamide		Nintedanib	
Minims Pilocarpine		Mylan Midazolam		Nipent	15
Minims Prednisolone		Myleran		Nitrates	5
Minirin		Myometrial and Vaginal Hormon		Nitroderm TTS	
Minirin Melt		Preparations		Nitrofurantoin	
Mino-tabs		Myozyme		Nitrolingual Pump Spray	
Minocycline hydrochloride		- N -	20	Nivestim	
		Nadolol	51	Nivolumab	
Minorycin Minor Skin Infections		Naglazyme		Nizoral	
				Nodia	
Minoxidil		Nalcrom			
Mirena		Naloxone hydrochloride		Noflam 250	
Mirtazapine		Naltraccord		Noflam 500	10
Misoprostol	8	Naltrexone hydrochloride		Non-Steroidal Anti-Inflammatory	40
Mitomycin C		Naphazoline hydrochloride		Drugs	10
Mitozantrone		Naphcon Forte		Nonacog gamma, [Recombinant	
Mitozantrone Ebewe		Naprosyn SR 1000		Factor IX]	40
Mixtard 30		Naprosyn SR 750		Norethisterone	
MMR II		Naproxen		Genito-Urinary	
Moclobemide	124	Narcaricin mite	115	Hormone	
Modafinil		Nasal Preparations	227	Norflex	
Modavigil	139	Natalizumab	135	Norfloxacin	
Moduretic		Natulan	153	Noriday 28	
Molaxole	26	Nausafix	130	Norimin	7
Moments	71	Nausicalm	129	Normacol Plus	2
Mometasone furoate		Navelbine	156	Normison	13
Monogen	244	Necon	73	Norpress	
Montelukast		Nedocromil	226	Nortriptyline hydrochloride	
Montelukast Mylan		Nefopam hydrochloride		Norvir	
Moroctocog alfa [Recombinant		Neisvac-C		Noumed	
VIII]		Neo-B12		Noumed Paracetamol	
Morphine hydrochloride		Neo-Mercazole		NovaSource Renal	24
Morphine sulphate		Neo-Mercazole S29		Novatretin	
Motetis	118	Neocate Gold		NovoMix 30 FlexPen	
Mouth and Throat		Neocate Junior Unflavoured		NovoRapid	
Movapo		Neocate Junior Vanilla		NovoRapid FlexPen	
Moxifloxacin		Neocate SYNEO		NovoRapid Penfill	
MSUD Maxamum		Neoral		NovoSeven RT	1 //
Mucilaginous laxatives with	204	Neostigmine metilsulfate		Noxafil	41
	05				
stimulants		Nepafenac		Nozinan	
Mucolytics		Nepro HP (strawberry)		Nozinan (Swiss)	د ان
Mucosoothe		Nepro HP (vanilla)		Nucala	
Multiple Sclerosis Treatments		Nepro HP RTH		Nuelin	
Multivitamin renal		Neulactil		Nuelin-SR	
Multivitamins	34	Neulastim	44	Nutilis	25

Nutren Diabetes	243	Omeprazole actavis 10	9	Pacimol	12
Nutrient Modules	240	Omeprazole actavis 20	9	Paclitaxel	15
Nutrini Energy Multi Fibre	245	Omeprazole actavis 40	9	Paclitaxel Actavis	15
Nutrini Energy RTH	245	Omnitrope		Paclitaxel Ebewe	15
Nutrini Low Energy Multi Fibre	247	Onbrez Breezhaler	222	Paediatric Seravit	
Nutrini Peptisorb	257	Oncaspar LYO	153	Palbociclib	16
Nutrini Peptisorb Energy		OncoTICE	173	Paliperidone	13
Nutrini RTH	245	Ondansetron	130	Pamidronate disodium	11
Nutrison 800 Complete Multi		Ondansetron ODT-DRLA	130	Pamisol	11
Fibre	250	One-Alpha	33	Panadol	
Nutrison Concentrated	252	Onrex	130	Panadol Mini Caps	12
Nutrison Energy		Opdivo		Pancreatic enzyme	
Nutrison Energy Multi Fibre	250	Ora-Blend	238	Pantoprazole	
Nutrison Multi Fibre	250	Ora-Blend SF	238	Panzop Relief	
Nutrison Standard RTH	250	Ora-Plus		Panzytrat	2
Nyefax Retard	52	Ora-Sweet		Papaverine hydrochloride	5
Nystatin		Ora-Sweet SF	238	Para-amino salicylic acid	9
Alimentary	32	Orabase	32	Paracare	12
Genito-Urinary	74	Oral and Enteral Feeds	243	Paracare Double Strength	12
Infection	96	Oratane	60	Paracetamol	12
NZB Low Gluten Bread Mix	253	Ordine	121	Paracetamol + Codeine	
-0-		Orgran	253	(Relieve)	12
O/W Fatty Emulsion Cream	64	Ornidazole	98	Paracetamol Pharmacare	12
Obinutuzumab	193	Orphenadrine citrate	116	Paracetamol with codeine	12
Obstetric Preparations	76	Ortho-tolidine	76	Paraffin	64-6
Ocicure	9	Oruvail SR	109	Paraffin liquid with wool fat	23
Ocrelizumab	135	Osmolite RTH	250	Paraldehyde	12
Ocrevus	135	Other Endocrine Agents	87	Parasidose	6
Octocog alfa [Recombinant facto	or	Other Oestrogen Preparation	s81	Parasiticidal Preparations	6
VIII] (Advate)	41	Other Progestogen		Parlodel	11
Octocog alfa [Recombinant factor		Preparations	81	Parnate	12
VIII] (Kogenate FS)	41	Other Skin Preparations	69	Parnate S29	12
Octreotide	164	Ovestin		Paromomycin	9
Octreotide (Sun)	164	Genito-Urinary	74	Paroxetine	
Octreotide GH	164	Hormone	81	Paser	9
Octreotide LAR (somatostatin		Ox-Pam	134	Paxam	13
analogue)	164	Oxaliplatin	146	Paxtine	12
Octreotide MaxRx	164	Oxaliplatin Accord	146	Pazopanib	16
Oestradiol	80	Oxaliplatin Actavis 100		Peak flow meter	22
Oestradiol valerate	80	Oxaliplatin Ebewe	146	Pedialyte - Bubblegum	4
Oestradiol with norethisterone	81	Oxazepam		Pediasure	24
Oestriol		Oxis Turbuhaler	222	Pediasure RTH	24
Genito-Urinary	74	Oxpentifylline	57	Pegaspargase	15
Hormone	81	Oxybutynin	75	Pegasys	10
Oestrogens	80	Oxycodone hydrochloride	122	Pegfilgrastim	
Ofev		Oxycodone Sandoz	122	Pegylated interferon alfa-2a	
Oil in water emulsion	64	Oxycodone Sandoz S29		Pembrolizumab	
Olanzapine		OxyContin		Pemetrexed	14
Olaparib	152	OxyNorm	122	Penicillamine	11
Olbetam		Oxytocin		Penicillin G	9
Olbetam S29	54	Oxytocin BNM		PenMix 30	1
Olopatadine		Oxytocin with ergometrine		PenMix 40	
Olopatadine Teva	234	maleate	75	PenMix 50	
Olsalazine		Ozurdex		Pentasa	
Omalizumab		.P.		Pentostatin [Deoxycoformycin]	
Omeprazole		Pacifen	116	Pentoxifylline [Oxpentifylline]	

Peptamen Junior	245	Pneumococcal (PCV10) conjuga	ate	Progesterone	81
Peptisoothe	9	vaccine		Proglicem	10
Peptisorb	246	Pneumococcal (PCV13) conjuga	ate	Proglycem	10
Perhexiline maleate	52	vaccine	269	Progynova	80
Pericyazine	131	Pneumococcal (PPV23)		Prolia	110
Perindopril	48	polysaccharide vaccine	270	Promethazine hydrochloride	221
Perjeta		Pneumovax 23	270	Propafenone hydrochloride	50
Permethrin	67	Podophyllotoxin	69	Propamidine isethionate	
Perrigo	69	Polaramine	221	Propranolol	
Pertuzumab		Poliomyelitis vaccine	270	Propylene glycol	238
Peteha	99	Poloxamer	25	Propylthiouracil	
Pethidine hydrochloride	123	Poly-Gel		Prostacur	
Pevaryl	61	Poly-Tears	233	Protaphane	11
Pexsig		Poly-Visc		Protaphane Penfill	11
Pfizer Exemestane		Polycal	240	Protifar	
Pharmacy Health Sorbolene wi	ith	Ponstan		Protionamide	99
Glycerin	64	Posaconazole	96	Provera	80
Pheburane		Postinor-1	74	Provera HD	81
Phenasen		Potassium chloride	45-46	PSM Citalopram	
Phenobarbitone	127	Potassium Chloride Aguettant	45	Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate		Preparations	67
Extemporaneous	238	Potassium iodate		PTU	82
Nervous		Povidone iodine		Pulmicort Turbuhaler	
Phenothrin		Pradaxa		Pulmozyme	
Phenoxybenzamine		Pramipexole hydrochloride	117	Puri-nethol	148
hydrochloride	47	Pravastatin		Puria	
Phenoxymethylpenicillin (Penic		Pravastatin Mylan		Pyrazinamide	
V)		Praziquantel		Pyridostigmine bromide	
Phenytoin sodium		Prazosin		Pyridoxine hydrochloride	
Phillips Milk of Magnesia		Pred Forte		Pyrimethamine	
Phlexy 10		Prednisolone		Pytazen SR	
Phosphate Phebra		Prednisolone acetate		- Q -	
Phosphorus		Prednisolone sodium		Quetapel	131
Phytomenadione		Prednisolone sodium		Quetiapine	
Pilocarpine hydrochloride		phosphate	231	Quick-Set MMT-392	
Pimafucort		Prednisolone-AFT		Quick-Set MMT-393	
Pimecrolimus		Prednisone		Quinapril	
Pindolol		Pregabalin		Quinapril with	
Pine tar with trolamine laurilsul		Pregabalin Pfizer		hydrochlorothiazide	48
and fluorescein		Pregnancy Tests - hCG Urine		Qvar	
Pinetarsol		Premarin		- R -	
Pioglitazone		Prevenar 13		RA-Morph	121
Pirfenidone		Priadel		Raloxifene hydrochloride	
Pizotifen		Primaquine		Raltegravir potassium	
PKU Anamix Infant		Primidone		Ramipex	
PKU Anamix Junior		Primolut N		Ranbaxy-Cefaclor	
PKU Anamix Junior Chocolate		Priorix		Ranitidine	
PKU Anamix Junior LQ		Probenecid		Rapamune	
PKU Anamix Junior Orange		Probenecid-AFT		Rasagiline	
PKU Anamix Junior Vanilla		Procaine penicillin		Reandron 1000	70
PKU Lophlex LQ 10		Procarbazine hydrochloride		Recombinant factor IX	
PKU Lophlex LQ 20		Prochlorperazine		Recombinant factor VIIa	
PKU Lophlex Powder				Recombinant factor VIII	40.41
•		Proctofoam		Rectogesic	
PKU Lophlex Sensation 20		Proctosedyl			
Plaquenil		Procyclidine hydrochloride	110	Redipred	
Plendil ER	52	Procytox	145	Relieve	108

Relistor	25	factor VIII]	41	Sodibic	4
Remicade	185	Ruxolitinib	161	Sodium acid phosphate	20
Renilon 7.5	246	Rythmodan		Sodium alginate	
Resonium-A	46	Rytmonorm		Sodium benzoate	
Resource Beneprotein		· - S -		Sodium bicarbonate	
Respigen		Sabril	128	Blood	45-46
Respiratory Devices		Sacubitril with valsartan		Extemporaneous	
Respiratory Stimulants		SalAir	223	Sodium calcium edetate	
Retinol palmitate		Salazopyrin		Sodium chloride	
ReTrieve		Salazopyrin EN		Blood	4
Retrovir		Salbutamol		Respiratory	
Revlimid		Salbutamol with ipratropium		Sodium citrate with sodium lauryl	
Revolade		bromide	223	sulphoacetate	26
Rexacrom		Salicylic acid	68	Sodium citro-tartrate	
Riboflavin	29	Salmeterol		Sodium cromoglicate	
Ribomustin	144	Sandomigran		Alimentary	8
Ricit	75	Sandostatin LAR		Respiratory	
Rifabutin	99	Sanofi Primaquine		Sensory	
Rifadin		Sapropterin dihydrochloride		Sodium fluoride	
Rifampicin		Scalp Preparations		Sodium Fusidate [fusidic acid]	
Rifaximin		Scopoderm TTS		Dermatological	6·
Rifinah		Sebizole		Infection	
Rilutek		Secukinumab		Sensory	
Riluzole	118	Sedatives and Hypnotics		Sodium hyaluronate [Hyaluronic	
Riodine		Seebri Breezhaler		acid]	233
Risedronate Sandoz	112	Selegiline hydrochloride		Sodium phenylbutyrate	
Risedronate sodium		Senna		Sodium picosulfate	
Risperdal Consta		Senokot		Sodium polystyrene sulphonate	
Risperidone		Sensipar		Sodium tetradecyl sulphate	
Risperidone (Teva)		SensoCard		Sodium valproate	
Risperon		Serenace		Sofradex	
Ritalin		Seretide		Soframycin	
Ritalin LA	138	Seretide Accuhaler	223	Solgar	
Ritonavir		Serevent		Solifenacin Mylan	
Rituximab (Mabthera)		Serevent Accuhaler		Solifenacin succinate	
Rituximab (Riximyo)		Sertraline		Solu-Cortef	
Rivaroxaban		Setrona		Solu-Medrol	
Rivastigmine		Setrona AU		Solu-Medrol-Act-O-Vial	
Rivotril		Sevredol		Somatropin (Omnitrope)	
Riximyo		Sex Hormones Non		Sotalol	
RIXUBIS		Contraceptive	79	Spacer device	
Rizamelt		Shield XL		Span-K	4
Rizatriptan		shingles vaccine		Spiolto Respimat	22
Ropin		SII-Onco-BCG		Spiractin	54
Ropinirole hydrochloride		Sildenafil	58	Spiriva	
Rotarix		Silhouette MMT-373		Spiriva Respimat	224
Rotavirus oral vaccine	270	Siltuximab		Spironolactone	
Roxane	6	Simvastatin	55	Sporanox	
Roxane-Propranolol		Simvastatin Mylan		Sprycel	15
Roxithromycin		Sinemet		Staphlex	92
Rubifen		Sinemet CR		Stemetil	
Rubifen SR		Sirolimus		SteroClear	
Rugby Capsaicin Topical		Siterone		Stesolid	
Cream	119	Slow-Lopresor	51	Stimulants/ADHD Treatments	
Rulide D		Smith BioMed Rapid Pregnand		Stiripentol	
Rurioctocog alfa pegol [Reco		Test		Stocrin	
J					

Stomahesive	32	Telfast	221	Topiramate Actavis	128
Strattera	136	Teligent	93	Total parenteral nutrition (TPN)	
Strides Shasun	96	Temaccord		TPN	45
Stromectol	65	Temazepam	136	Tramadol hydrochloride	123
Sucralfate	10	Temozolomide	154	Tramal SR 100	
Sulfadiazine Silver	61	Tenofovir disoproxil	100	Tramal SR 150	
Sulfadiazine sodium	94	Tenofovir Disoproxil Teva		Tramal SR 200	
Sulfasalazine		Tenoxicam		Trandate	
Sulindac		Tensipine MR10		Tranexamic acid	
Sulindac Mylan	109	Tepadina		Tranylcypromine sulphate	124
Sulphur		Terbinafine		Trastuzumab	
Sulprix		Terbutaline sulphate		Trastuzumab emtansine	
Sumatriptan		Teriflunomide		Travatan	
Sunitinib		Teriparatide		Travoprost	
Sunscreens		Testosterone		Travopt	
Sunscreens, proprietary		Testosterone cipionate		Treatments for Dementia	
Sure-T MMT-863		Testosterone esters		Treatments for Substance	
Sure-T MMT-873		Testosterone undecanoate		Dependence	140
Sustagen Diabetic		Tetrabenazine		Trental 400	
Sustagen Hospital Formula		Tetrabromophenol		Tretinoin	
Active	250	Tetracosactrin		Dermatological	60
Sustanon Ampoules		Tetracycline		Oncology	
Sutent		Thalidomide		Trexate	
Sylvant		Thalomid		Triamcinolone acetonide	
Symbicort Turbuhaler 100/6		Theophylline		Alimentary	32
Symbicort Turbuhaler 200/6		Thiamine hydrochloride		Dermatological	
Symbicort Turbuhaler 400/12		THIO-TEPA		Hormone	
Symmetrel		Thioguanine		Triamcinolone acetonide with	
Sympathomimetics		Thiotepa		gramicidin, neomycin and nysta	atin
Synacthen		Thymol glycerin		Dermatological	
Synacthen Depot		Thyroid and Antithyroid Agents .		Sensory	
Synacthene Retard		Ticagrelor		Triaver	
Synflorix		Tilade	226	Triazolam	
Synthroid		Tilcotil		Trimethoprim	
Syntometrine		Tillomed		Trimethoprim with	
Syrup (pharmaceutical grade)		Timolol		sulphamethoxazole	
Systane Unit Dose	233	Timoptol XE		[Co-trimoxazole]	95
- T -		Tiotropium bromide		Trisequens	
Tacrolimus	219	Tiotropium bromide with		Trisul	
Tacrolimus Sandoz		olodaterol	224	Trophic Hormones	
Taliglucerase alfa		Tivicay		Tropicamide	
Tambocor		TMP		Trusopt	
Tamoxifen citrate		Tobramycin		TruSteel	
Tamoxifen Sandoz		Infection	95	Tuberculin PPD [Mantoux] test	271
Tamsulosin hydrochloride		Sensory		Tubersol	271
Tamsulosin-Rex		Tobramycin BNM		Two Cal HN	
Tandem Cartridge		Tobramycin Mylan		Two Cal HN RTH	
Tandem t:slim X2 with Basal-IQ		Tobrex		Tykerb	
Tap water		Tocilizumab		Tysabri	120
Tarceva		Tofranil		- U -	100
Tasigna		Tolcapone		Ultibro Breezhaler	22/
Tasıyıra		Topamax		Ultraproct	
Taurine		Topical Products for Joint and	120	Umeclidinium	20/
Tecfidera		Muscular Pain	110	Umeclidinium with vilanterol	
Tegretol		Topiderm		Univent22	
Tegretol CR		Topiramate		Ural22	
regietoi Ort	140	1 opii ai ii ai c	140	υιαι	/ (

Urea64	Viramune Suspension104
Urex Forte53	ViruPOS229
Urinary Agents75	Vit.D333
Urinary Tract Infections108	Vita-B1233
Urinorm115	VitA-POS234
Uromitexan 152	Vitabdeck34
Ursodeoxycholic acid24	Vital246
Ursosan24	Vitamin B complex33
Utrogestan81	Vitamin B6 25
-V-	Vitamins33–34
Vaccinations	Vivonex TEN246
Vaclovir	Voltaren
Valaciclovir	Voltaren D
Valganciclovir	Voltaren Ophtha230
Valganciclovir Mylan100	Volumatic228 Voriconazole
Vancomycin95	
Vannair	Vosol
Varenicline Pfizer142	Votrient
Varenicline tartrate142	Vttack97
Varicella vaccine [Chickenpox	- W -
vaccine]	Warfarin sodium
Varicella zoster virus (Oka strain) live	Wart Preparations
attenuated vaccine [shingles	Wasp venom allergy treatment220 Water
vaccine]	Blood45
Vasodilators	Extemporaneous
Vasorex	- X -
Vasorex	Xarelto44
Veletri	Xifaxan
Venclexta	XMET Maxamum
Venetoclax	Xolair
Venlafaxine 125	XP Maxamum254
Venomil	Xylocaine119
VENOX	Xylocaine 2% Jelly118
Ventavis	Xyntha40
Ventolin	- Z -
Vepesid	Zapril47
Verapamil hydrochloride52	Zarontin
Vergo 16129	Zaroxolyn53
Vermox88	Zavedos151
Versacloz131	Zeffix100
Vesanoid155	Zetlam100
Vexazone12	Ziagen104
Vfend97	Zidovudine [AZT]105
Viaderm KC63	Zidovudine [AZT] with
Vidaza146	lamivudine 105
Vigabatrin128	Zimybe55
Vildagliptin12	Zinc and castor oil64
Vildagliptin with metformin	Zinc sulphate36
hydrochloride12	Zincaps36
Vimpat126	Zinnat88
Vinblastine sulphate156	Ziprasidone132
Vincristine sulphate156	Zista221
Vinorelbine156	Zithromax89
Vinorelbine Ebewe	

Zoledronic acid	
Hormone	<mark>77</mark>
Musculoskeletal	113
Zoledronic acid Mylan	77
Zopiclone	136
Zopiclone Actavis	136
Zostavax	271
Zostrix	110
Zostrix HP	119
Zuclopenthixol decanoate	133
Zuclopenthixol hydrochloride	132
Zusdone	132
Zyban	141
Zypine	
Zypine ODT	131
Zyprexa Relprevv	132
Zytiga	