# September 2021 Volume 28 Number 2

#### **Editors:**

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

#### Circulation

Accessible in an electronic format at no cost from the Pharmac website <a href="https://www.pharmac.govt.nz/schedule">www.pharmac.govt.nz/schedule</a>.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

#### Production

Typeset automatically from XML and T<sub>E</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

### **Programmers**

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

2	Introducing Pharmac	
5	General Rules	Section A
6	Alimentary Tract & Metabolism	Section B
38	Blood & Blood Forming Organs	
48	Cardiovascular System	
62	Dermatologicals	
72	Genito Urinary System	
79	Hormone Preparations – Systemic	
90	Infections – Agents For Systemic Use	
112	Musculoskeletal System	
120	Nervous System	
149	Oncology Agents & Immunosuppressants	
225	Respiratory System & Allergies	
234	Sensory Organs	
240	Various	
242	Extemporaneous Compounds (ECPs)	Section C
245	Special Foods	Section D
265	National Immunisation Schedule	Section I
277	Index	

Introducing Pharmac

# **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

#### Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://www.pharmac.govt.nz/about">https://www.pharmac.govt.nz/about</a>.

# **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

# **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

#### Example



# Glossary

# **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

# SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 n sachet	0.	30	<b>✓</b> Ga	viscon Infant
GODIUM ALGINATE  ★ Tab 500 mg with sodium bicarbonate 267 mg and calciur carbonate 160 mg - peppermint flavour		60		viscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and cal carbonate 160 mg per 10 ml		500 ml	Ac	idex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg CALCIUM CARBONATE		100	✓ Alı	u-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) Subsidy by endorsement Only when prescribed for patients unable to swallow inappropriate and the prescription is endorsed accor	39.00	500 ml ts or whe	✓ Ro re calcium	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available  Tab 2 mg Cap 2 mg	10.75	400 400	✓ No ✓ <u>Dia</u>	dia amide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap 3 mg - Special Authority see SA1886 below - Reta pharmacy	166.50	90 alid for 6		tocort CIR r applications meeting

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
  - 2.1 Diabetes; or
  - 2.2 Cushingoid habitus; or
  - 2.3 Osteoporosis where there is significant risk of fracture; or

(N	Subsidy fanufacturer's Price)	Subsi	Fully dised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes; or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture; or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
  - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with \* is an unapproved indication.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

#### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	<ul><li>✓ Cortifoam S29</li><li>✓ Colifoam</li></ul>
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab EC 500 mg49.50	100	✓ Asamax
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	✓ Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	/	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	•	Nalcrom
SULFASALAZINE  * Tab 500 mg  Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

# Local preparations for Anal and Rectal Disorders

# Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00 Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	30 g OP 12	✓ Proctosedyl ✓ Proctosedyl

# **Management of Anal Fissures**

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

# **⇒SA1329** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

# **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE  * Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE  * Tab 135 mg	0.20	90	✓ Colofae
* Tab 100 IIIg	5.20	90	• Cololac

# **Antiulcerants**

# **Antisecretory and Cytoprotective**

MIS	SOPROSTOL STORE ST		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	Cytotec

		ALIMENTARY	'TRAC	T AND	METABOLISM
		Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
Н	elicobacter Pylori Eradication				
CL	ARITHROMYCIN Tab 500 mg – Subsidy by endorsement	10.40 14.58	14		po-Clarithromycin lacid
(Ap	<ul> <li>a) Maximum of 28 tab per prescription</li> <li>b) Subsidised only if prescribed for helicobacter pylori er Note: the prescription is considered endorsed if clarit inhibitor and either amoxicillin or metronidazole.</li> <li>co-Clarithromycin Tab 500 mg to be delisted 1 February 2022)</li> </ul>				
Н	2 Antagonists				
	MOTIDINE – Only on a prescription Tab 20 mg	4.91	100	<b>✓</b> F	amotidine Hovid §29
*	Tab 40 mg	8.48	100	<b>✓</b> F	amotidine Hovid \$29
*	Inj 10 mg per ml, 4 ml — Subsidy by endorsement		10 t of pallia		ylan S29
P	roton Pump Inhibitors				
	NSOPRAZOLE  Cap 15 mg		100	<b>√</b> L	anzol Relief
*	Cap 30 mgLanzol Relief to be Principal Supply on 1 December 2021	5.26	100	<b>✓</b> L	anzol Relief
ON	1EPRAZOLE				
	For omeprazole suspension refer Standard Formulae, page 2				
*	Cap 10 mg	1.94	90	<b>√</b> <u>0</u>	meprazole actavis 10
*	Cap 20 mg	1.86	90	<b>√</b> <u>0</u>	meprazole actavis 20
*	Cap 40 mg	3.11	90	<b>√</b> <u>0</u>	meprazole actavis 40
*	Powder – Only in combination Only in extemporaneously compounded omeprazole susp		5 g	✓ M	idwest
	1 10 11 11 11 11 11 11 11 11 11 11 11 11	00.00	-		<b>B</b>

# 

(Ocicure S29 Inj 40 mg ampoule with diluent to be delisted 1 October 2021)

✓ Panzop Relief 100 100 Panzop Relief

# **Site Protective Agents**

**PANTOPRAZOLE** 

COLLOIDAL BISMUTH SUBCITRATE

Tab 120 mg ......14.51

✓ Gastrodenol S29

50

✓ Dr Reddy's

Omeprazole ✓ Ocicure S29

	Subsidy (Manufacturer's Price \$	e) Si Per	Fully ubsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g	35.50 (48.28)	120	(	Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail Tab 550 mg		56	✓)	(ifaxan
■ SA1461 Special Authority for Subsidy  initial application only from a gastroenterologist, hepatolog  nepatologist. Approvals valid for 6 months where the patier  olerated doses of lactulose.  Renewal only from a gastroenterologist, hepatologist or Pra  nepatologist. Approvals valid without further renewal unless  benefiting from treatment.	gist or Practitioner on the nt has hepatic encephalo actitioner on the recomme	pathy des endation o	pite an a	dequate trial of maximur penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retai Cap 25 mg Cap 100 mg Oral liq 50 mg per ml  >>SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approval nypoglycaemia caused by hyperinsulinism.  Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.  GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO			For the tre	
	52.00	'	• •	<u>alucagen riypokit</u>
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml OP		Actrapid
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ F	Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen NSULIN ISOPHANE	52.15	5	<b>✓</b> N	NovoMix 30 FlexPen
▲ Inj human 100 u per ml	17.68	10 ml OP	-	Humulin NPH Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	<b>✓</b>	Humulin NPH Protaphane Penfill

	Subsidy (Manufacturer's Price	no) Subc	Fully Brand or idised Generic
	(Manufacturer 3 i iii	Per Subs	✓ Manufacturer
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
,			✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30
			✓ PenMix 40
			✓ PenMix 50
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per n	nl,		
3 ml	42.66	5	Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per n	nl,		
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE	00.00	4	/ Lambura
▲ Inj 100 u per ml, 10 ml		1	✓ Lantus
▲ Inj 100 u per ml, 3 ml		5	<ul><li>✓ Lantus</li><li>✓ Lantus SoloStar</li></ul>
Inj 100 u per ml, 3 ml disposable pen	94.50	5	Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
▲ Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	✓ NovoRapid FlexPen
NSULIN GLULISINE			·
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
▲ Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
NSULIN LISPRO			•
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
Alpha Glucosidase Inhibitors			<u> </u>
ACARBOSE	0.50	00	√ Cluschev
* Tab 50 mg		90	✓ Glucobay
Accord to be Principal Supply on 1 December 2001	8.95		✓ Accarb
Accarb to be Principal Supply on 1 December 2021  Tab 100 mg	6.40	90	✓ Glucobay
- Tab Too Hig	15.29	50	✓ Accarb
Accarb to be Principal Supply on 1 December 2021	. 0.20		
Glucobay Tab 50 mg to be delisted 1 December 2021)			
Glucobay Tab 100 mg to be delisted 1 December 2021)			
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE	7.50	100	√ Doonil
* Tab 5 mg	7.50	100	✓ Daonil
GLICLAZIDE	45.40	500	<b>4</b> 011-1-1-
* Tab 80 mg	15.18	500	✓ Glizide

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
GLIPIZIDE				
* Tab 5 mg	3.27	100	✓	Minidiab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	8.63	1,000	1	Apotex
* Tab immediate-release 850 mg	7.04	500	✓	Apotex
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓	Vexazone
* Tab 30 mg	7.30	90	✓	Vexazone
* Tab 45 mg	12.25	90	✓	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	✓	Galvus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	✓	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	1	Galvumet

# **GLP-1 Agonists**

### ⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*: or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
  - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE – Special Authority see SA2065 above – Retail pharmacy
Note: Not to be given in combination with a funded SGLT-2 inhibitor.

★ Inj 1.5mg per 0.5 ml prefilled pen .......115.23 4 ✓ Trulicity

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per 🗸	<ul> <li>Manufacturer</li> </ul>

#### SGLT2 Inhibitors

### ⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Fither:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Maori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
  - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

# EMPAGLIFLOZIN – Special Authority see SA2068 above – Retail pharmacy

*	Tab 10 mg	30	<ul><li>Jardiance</li></ul>
*	Tab 25 mg 58.56	30	✓ Jardiance

# EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE — Special Authority see SA2068 above — Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

# **Diabetes Management**

# **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

# **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips \_\_\_\_\_\_20.00 1 OP 

CareSens Dual

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes; or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

20.00

✓ CareSens N Premier

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRC

#### BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

# **Insulin Syringes and Needles**

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

### INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

IIVC	oblini bin nebelo inaximam of 200 dev per prescripti	OH		
*	29 g × 12.7 mm	10.50	100	B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	00 dev per p	prescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.00	100	✓ B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	✓ B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II

# **Insulin Pumps**

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each fou	ır year period.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

### **⇒SA1603** Special Authority for Subsidy

**Initial application** — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
<b>\$</b>	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Fither:
  - 4.1 Applicant is a relevant specialist: or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

<del></del>			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)	Su.	Fully bsidised	Brand or Generic	
(Mandacaters i nee)	Per	J	Manufacturer	
Ψ	rei	•	Manuacturei	

#### continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Fither:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.

# **Insulin Pump Consumables**

#### ⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

# All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy (Manufacturer's Price)	Fu Subsidise	,	Brand or Generic
\$	Per	<b>✓</b>	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

1 OP

1 OP

✓ Sure-T MMT-863

✓ Sure-T MMT-873

	ALIMENTAR	Y TRA	CT AND	METABOLISM
	Subsidy (Manufacturer's Price \$	e) S Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
pump therapy; and 4 The patient is continuing to derive benefit from pump the 5 The patient had achieved and is maintaining a HbA1c of 6 The patient has had no increase in severe unexplained 7 The patient's HbA1c has not deteriorated more than 5 m 8 Either:  8.1 Applicant is a relevant specialist; or	f equal to or less than hypoglycaemic episoo nmol/mol from baselin	des from		
8.2 Applicant is a nurse practitioner working within the Renewal — (Previous use before 1 September 2012) only from the control of the contro		lict or nu	rco proctiti	oner Approvale valid for G
years for applications meeting the following criteria:	oni a relevant specia	iist oi Tiu	ise praciiii	orier. Approvais valid for 2
All of the following:				
1 The patient is continuing to derive benefit according to the	he treatment plan and	l has ma	intained a	HbA1c of equal to or less
than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 m	amol/mol from initial a	nnlicatio	n: and	
3 The patient has not had an increase in severe unexplair				ne; and
4 Either:	7, 3,			.,
4.1 Applicant is a relevant specialist; or				
4.2 Applicant is a nurse practitioner working within the	neir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 19 - Retail	pharmac	:y	
a) Maximum of 3 sets per prescription				
<ul><li>b) Only on a prescription</li><li>c) Maximum of 13 packs of cartridge sets will be funded p</li></ul>	or voar			
Cartridge 300 U, t:lock × 10	50.00	1 OP	<b>√</b> T	andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Specia		85 on pa	ge 19 – Re	etail pharmacy
a) Maximum of 3 sets per prescription	,	•		, ,
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.  10 mm steel needle; 60 cm tubing × 10	120.00	1 OP	./ N	IiniMed Sure-T
10 min steer needle, 60 cm tubing x 10	130.00	TOP	• IV	MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T
				MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	<b>✓</b> N	liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	<b>✓</b> N	IiniMed Sure-T
o min otoo hoodio, oo on tabing x 10		1 01	- 14	MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	liniMed Sure-T
				MMT-876A

6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

8 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
 \$	Per	•	Manufacturer	

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.1	00 1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel

(TruSteel 6 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021) (TruSteel 8 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021)

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm clear tubing × 10 ......130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
   ✓ MiniMed Silhouette
- MMT-381A

  MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
   10 P

   6 mm teflon needle, 45 cm blue tubing × 10
   130.00
   1 OP
- ✓ MiniMed Quick-Set MMT-398A
   ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
  - ✓ MiniMed Mio MMT-943A
  - ✓ MiniMed Mio MMT-923A
  - ✓ MiniMed Quick-Set MMT-399A
  - ✓ MiniMed Mio MMT-945A
  - ✓ MiniMed Mio MMT-965A
  - ✓ MiniMed Mio MMT-925A
  - ✓ MiniMed Quick-Set MMT-387A
  - ✓ MiniMed Quick-Set MMT-396A
  - ✓ MiniMed Quick-Set MMT-397A
  - ✓ MiniMed Mio MMT-975A
  - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. \*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1985 on page 19 – Retail pharmacy  a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH II	NSERTION	I DEVICE)	- Special Authority see
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	cm 140.00	1 OP	<b>√</b> Διι	toSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cl	n	1 OP		toSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II Retail pharmacy  a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with	NSERTION) - Spe			
10 needles; luer lock	IT INSERTION WIT	1 OP H INSERT		houette MMT-373 ICE) – Special Authority
110 cm line × 10 with 10 needles	m	1 OP		toSoft 90
line x 10 with 10 needles		1 OP		toSoft 90 toSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m	1 OP		toSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy  a) Maximum of 3 sets per prescription		-		
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock	130.00	1 OP	<b>√</b> Qu	ick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock		1 OP	<b>√</b> Qu	ick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 c  a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per	n page 19 – Retail <sub>I</sub>	oharmacy	-	
10 × luer lock conversion cartridges 1.8 ml for Paradigm pun Cartridge for 5 and 7 series pump; 1.8 ml × 10	ps50.00	1 OP 1 OP	✔ Mir 1	R Cartridge 1.8 niMed .8 Reservoir //MT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		niMed 8.0 Reservoir //MT-332A

Fully

Brand or

	(Manufacturer's Price	) Sub Per	osidised	Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					_
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Cr	eon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))	*	100	✓ Pa	nzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Cr	eon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph		00 00			
Eur U)		20 g OP	✓ Cr	eon Micro	
URSODEOXYCHOLIC ACID – Special Authority see SA1739 be	•	•			
Cap 250 mg	32.95	100	✓ <u>Ur</u> s	sosan	

Subsidy

# ⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- ner:
  - 1 Patient has been diagnosed with Alagille syndrome; or
  - 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

**Initial application** — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application** — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

# Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Su	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	dised	Generic
	\$ Per	•	Manufacturer

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

#### Laxatives

# **Bulk-forming Agents**

* Powder for oral soln	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		·	
* Dry	6.02	500 g OP	
·	(17.32)	•	Normacol Plus
	2.41	200 g OP	
	(8.72)		Normacol Plus

#### **Faecal Softeners**

* Tab 50 mg 2.31 * Tab 120 mg 3.13	100 100	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>
DOCUSATE SODIUM WITH SENNOSIDES  * Tab 50 mg with sennosides 8 mg	200	✓ Laxsol
POLOXAMER – Only on a prescription  Not funded for use in the ear.		
* Oral drops 10%	30 ml OP	✓ Coloxyl

# **Opioid Receptor Antagonists - Peripheral**

DOCUSATE SODIUM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below - Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
• •	246.00	7	✓ Relistor

#### ⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$	e) Sul Per	Fully osidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL  * Suppos 3.6 g – Only on a prescription  LACTULOSE – Only on a prescription	9.25	20	<b>√</b> P	SM
* Oral liq 10 g per 15 ml	3.33	500 ml	<b>√</b> L	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIG	CARBONATE AND	SODIUM	CHLORII	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7	O,	30	✓ N	<u>lolaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	<b>√</b> F	leet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	•		P1-11-
5 ml	29.98	50	<b>✓</b> <u>IV</u>	<u>licolette</u>
Stimulant Laxatives				
BISACODYL — Only on a prescription  * Tab 5 mg  * Suppos 10 mg  Lax-Suppositories to be Principal Supply on 1 December	3.69	200 10	_	ax-Tab ax-Suppositories
SENNA – Only on a prescription  * Tab, standardised	2.17 (8.21)	100	S	enokot
	0.43 (2.06)	20	Q	enokot
SODIUM PICOSULFATE - Special Authority see SA2053 below	` '		3	GIIOROL
Oral soln 7.5 mg per ml		30 ml OP	<b>✓</b> D	ulcolax SP Drop

# SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# **Metabolic Disorder Agents**

# ⇒SA1986 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells: or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE - Sr	ecial Authority see	SA2042 helow -	- Retail nharmacy
---------------	---------------------	----------------	-------------------

Tab 1,000 mg	CBS	90	<ul><li>Clinicians</li></ul>
Cap 500 mg	CBS	50	✓ Solgar
Powder		400 a	✓ Biomed

#### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Powder for oral soln......575.00

BETAINE – Special Authority see SA1987 below – Retail pharmacy

# ⇒SA1987 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

continued...

180 g OP

✓ Cystadane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	Go Healthy

# **⇒SA2039** Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

#### ⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

# ⇒SA1623 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

# ⇒SA1695 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - F	Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

# ⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

# ⇒SA2041 Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price)		Fully lised	Brand or Generic
	\$	Per	1	Manufacturer
SAPROPTERIN DIHYDROCHLORIDE - Special Authority see S	A1989 below – Reta	il pharmacy		
Tab soluble 100 mg	1,452.70	30 OP	1	Kuvan

#### ⇒SA1989 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

# All of the following:

- 1 Fither:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - R	letail pharmacy		
Soln 100 mg per ml	CBS	100 ml	✓ Amzoate S29

#### ⇒SA1599 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE − Special Authority see SA1990 below − Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP

✓ Pheburane

# ⇒SA1990 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 on the next page	<ul> <li>Retail pharmacy</li> </ul>		
Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

Subsidy (Manufacturer's Price)

Fully Subsidised Per Brand or Generic Manufacturer

# ⇒SA2043 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

# Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

✓ Elelyso

# ⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 Pharmac PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

#### **Access Criteria**

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific
  deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
  - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
  - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
  - Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
  - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

#### \*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic
 \$	Per 🗸	Manufacturer

continued...

#### All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 7) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

# Mouth and Throat

# **Agents Used in Mouth Ulceration**

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with Endorsement	9.00 (20.31)	500 ml	Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	oral mucositis a	as a result of tro	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	Ü	Orabase
	1.52	5 g OP	
	(3.60)	•	Orabase
Powder	8.48	28 g OP	
	(10.95)	-	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
7 Carlouro gor o.7 /o mar octamornam cinonao o.o 1 /o	(6.00)	10 9 01	Bonjela
TRIAMCINOLONE ACETONIDE	(0.00)		261,1010
	F 00	5 = OD	/ Kanalan in Ovehees
Paste 0.1%	5.33	5 g OP	<ul> <li>Kenalog in Orabase</li> </ul>
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
5 5		20	• Tungiiii
MICONAZOLE			
Oral gel 20 mg per g  Decozol to be Principal Supply on 1 December 2021	4.74	40 g OP	✓ Decozol
NYSTATIN			
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$		ully Brand or ed Generic  Manufacturer
Other Oral Agents			
For folinic mouthwash, pilocarpine oral liquid or saliva substitute for THYMOL GLYCERIN	ormula refer Standa	ırd Formulae, p	page 242
★ Compound, BPC PSM Compound, BPC to be delisted 1 February 2022)	9.15	500 ml	✓ PSM
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN ★ Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS0	O1.89		✓ Neo-B12 ✓ Vita-B12
	3.15		✓ Hydroxocobalamin Mercury Pharma
PYRIDOXINE HYDROCHLORIDE  a) No more than 100 mg per dose b) Only on a prescription			
* Tab 25 mg - No patient co-payment payable			Vitamin B6 25
Tab 50 mg	13.63	500	✓ Apo-Pyridoxine
HIAMINE HYDROCHLORIDE  - Only on a prescription  ★ Tab 50 mg	7.09	100	✓ Max Health
/ITAMIN B COMPLEX ★ Tab, strong, BPC			✓ Bplex
Vitamin C			
ASCORBIC ACID  a) No more than 100 mg per dose b) Only on a prescription			
* Tab 100 mg	9.90	500	✓ <u>Cvite</u>
Vitamin D			
ALFACALCIDOL	22.22	100	
* Cap 0.25 mcg * Cap 1 mcg			✓ One-Alpha ✓ One-Alpha
♦ Oral drops 2 mcg per ml			✓ One-Alpha
CALCITRIOL	7.05	400	Containing area
<ul><li>★ Cap 0.25 mcg</li><li>★ Cap 0.5 mcg</li></ul>			✓ <u>Calcitriol-AFT</u> ✓ <u>Calcitriol-AFT</u>
COLECALCIFEROL	- · <del>-</del>		
<ul> <li>★ Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription</li> <li>★ Oral liq 188 mcg per ml (7,500 iu per ml)</li> </ul>			✓ <u>Vit.D3</u> ✓ Puria
Multivitamin Preparations			
MULTIVITAMIN RENAL - Special Authority see SA1546 on the n  * Cap			✓ Clinicians Renal Vit

✓ fully subsidised

Principal Supply

	ALIMENTARY	TRACT AND	METABOLISM
(	Subsidy Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either:  1 The patient has chronic kidney disease and is receiving eith 2 The patient has chronic kidney disease grade 5, defined as	ner peritoneal dialysi	is or haemodialys	is; or
15 ml/min/1.73 m² body surface area (BŠA).  MULTIVITAMINS – Special Authority see SA1036 below – Retail p	pharmacy	·	
* Powder	without further renev	wal unless notifie	·
VITAMINS  * Tab (BPC cap strength)	11.45	1,000 ✓ N	luita
** Cap (fat soluble vitamins A, D, E, K) — Special Authority see     ** SA1720 below — Retail pharmacy			itabdeck
Initial application from any relevant practitioner. Approvals valid the following criteria:  Any of the following:  1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut sy 3 Patient has severe malabsorption syndrome.		wal unless notifie	d for applications meeting
Minerals			
Calcium			

CALCIUM CARBONATE			
* Tab 1.25 g (500 mg elemental)	6.69	250	✓ Calci-Tab 500
* Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement	52.00	20	✓ Calcium-Sandoz
			Forte S29
	54.60	76	✓ Cacit S29
Subsidy by endorsement – Only when prescribed for paediatric considered unsuitable.	c patients (< 5	years) wher	e calcium carbonate oral liquid is
CALCIUM GLUCONATE			
* Inj 10%, 10 ml ampoule	32.00	10	✓ Max Health -
			Hameln S29
	64.00	20	✓ Max Health S29
Fluoride			
SODIUM FLUORIDE			
* Tab 1.1 mg (0.5 mg elemental)	5.75	100	✓ PSM
	5.75	100	✓ PSM

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	<u>euroTabs</u>
Iron				
FERROUS FUMARATE  * Tab 200 mg (65 mg elemental)  FERROUS FUMARATE WITH FOLIC ACID	3.09	100	<b>√</b> F	erro-tab
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	<b>√</b> F	erro-F-Tabs
FERROUS SULFATE  * Tab long-acting 325 mg (105 mg elemental)  * Oral liq 30 mg (6 mg elemental) per 1 ml	2.06 12.08 5	30 600 m		errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority se Inj 50 mg per ml, 10 ml vial		etail ¡ 1		erinject

#### ⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

#### IRON POLYMALTOSE

✓ Ferrosia

# ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
Magnesium				
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml		nillips Milk of Magnesia 829
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ <u>M</u>	artindale
Zinc				
ZINC SULPHATE  * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ <u>Zi</u>	ncaps

Subsidy
(Manufacturer's Price)
\$ Per

Fully Subsidised

Brand or Generic Manufacturer

### **Antianaemics**

### Hypoplastic and Haemolytic

### ⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus: and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus; and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with \* is an unapproved indication

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
	<u> </u>	Per		Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	s page – Retail pharm	nacv		
Wastage claimable	, page 1121am p.1am.	,		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	1	Binocrit
		6		Binocrit
Inj 2,000 iu in 1 ml, syringe		-		
Inj 3,000 iu in 0.3 ml, syringe		6		Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6		Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	/	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓	Binocrit
Inj 10,000 iu in 1 ml, syringe		6		Binocrit
Inj 40,000 iu in 1 ml, syringe		1	_	Binocrit
11] 40,000 td 111 1 1111, 0y1111go	200.00	•	•	<u>Diniodrit</u>
Magalahlastia				
Megaloblastic				
FOLIC ACID				
	21.04	1,000	./	Apo-Folic Acid
Tab 0.8 mg		,		•
* Tab 5 mg		100		Folic Acid Mylan
	12.12	500	•	Apo-Folic Acid
Folic Acid Mylan to be Sole Supply on 1 December 2021				
Oral liq 50 mcg per ml	26.00 25	5 ml Ol	•	Biomed
(Apo-Folic Acid Tab 5 mg to be delisted 1 December 2021)				

### Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia

 Inj 250 iu vial.
 612.50
 1
 ✓ Alprolix

 Inj 500 iu vial.
 1,225.00
 1
 ✓ Alprolix

 Ini 1.000 iu vial.
 2.450.00
 1
 ✓ Alprolix

✓ Revolade

28

 Inj 1,000 iu vial.
 2,450.00
 1
 ✓ Alprolix

 Inj 2,000 iu vial.
 4,900.00
 1
 ✓ Alprolix

 Inj 3,000 iu vial.
 7,350.00
 1
 ✓ Alprolix

ELTROMBOPAG - Special Authority see SA1743 below - Retail pharmacy

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist.

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter: or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

**Initial application — (severe aplastic anaemia)** only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

4 T....

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

#### EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	1	✓ Hemlibra
Inj 150 mg in 1 ml vial	1	<ul><li>Hemlibra</li></ul>

⇒SA1969 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

(1)	Subsidy Manufacturer's Price)	Fully Subsidised	
	\$	Per 🗸	Manufacturer

#### continued...

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months;
- 2 Either:
  - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
  - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
  - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
  - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

**Renewal** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1	✓ NovoSeven RT
Inj 2 mg syringe2,356.60		✓ NovoSeven RT
Inj 5 mg syringe		✓ NovoSeven RT
Inj 8 mg syringe9,426.40	1	✓ NovoSeven RT

#### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 Ŭ	1,315.00	1	✓ FĚIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Ini 2.500 U	6.575.00	1	✓ FEIBA NF

#### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Subject to officia.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	✓ Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe		1	✓ Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

	Subsidy	F	ully Brand or
	(Manufacturer's Price)	Subsidis	
	\$	Per	✓ Manufacturer
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharm]			
For patients with haemophilia. Access to funded treatment is	managed by the Hae	emophilia Tr	eaters Group in conjunction
with the National Haemophilia Management Group. Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		-	✓ RIXUBIS
Inj 2,000 iu vial		=	✓ RIXUBIS
Inj 3,000 iu vial		=	✓ RIXUBIS
• •	•	•	· III/ODIO
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – [ For patients with haemophilia. Preferred Brand of short half-		vr \/III \\ A 0000	s to funded treatment is
managed by the Haemophilia Treaters Group in conjunction v			
Inj 250 iu vial			✓ Advate
Inj 500 iu vial			✓ Advate
Inj 1,000 iu vial		-	✓ Advate
Inj 1,500 iu vial		-	✓ Advate
Inj 2,000 iu vial		•	✓ Advate
Inj 3,000 iu vial			✓ Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	•	•	7147410
For patients with haemophilia. Rare Clinical Circumstances I		, rocombinor	ot factor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in	conjunction with the r	vational Hae	mophilia Management Group,
subject to criteria.	227 50	1	√ Kaganata ES
Inj 250 iu vial		-	✓ Kogenate FS
Inj 500 iu vial		-	✓ Kogenate FS
Inj 1,000 iu vial Inj 2,000 iu vial			✓ Kogenate FS ✓ Kogenate FS
Inj 3,000 iu vial	,		✓ Kogenate FS
		1	Nogeliale F3
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]			
For patients with haemophilia A receiving prophylaxis treatme		d treatment is	s managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			
Inj 250 iu vial			✓ Adynovate
Inj 500 iu vial			✓ Adynovate
Inj 1,000 iu vial			✓ Adynovate
Inj 2,000 iu vial	2,400.00	1	✓ Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml	28.50	5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	9.45	60	✓ Mercury Pharma
Vitamin K			
DI IVTOMENIA DIONIE			
PHYTOMENADIONE	0.00	_	/ Kamaldan MM
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO		_	✓ Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO	9.21	5	✓ Konakion MM
Antithrombotic Agents			
Antitinombotic Agents			
Antiplotolot Agento			
Antiplatelet Agents			
ASPIRIN			
* Tab 100 mg	10.80	990	✓ Ethics Aspirin EC
		<del>-</del>	

	Subsidy (Manufacturer's Price) \$	Per	
CLOPIDOGREL			
* Tab 75 mg	4.60	84	✓ <u>Clopidogrel</u> <u>Multichem</u>
DIPYRIDAMOLE			
* Tab long-acting 150 mg	10.90	60	✓ Pytazen SR
TICAGRELOR - Special Authority see SA1955 below - Retail pha	armacy		
* Tab 90 mg	90.00	56	✓ Brilinta
SA1055 Special Authority for Subsidy			

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
  - 1.2 Patient is about to have a neurological stenting procedure performed\*; and
- 2 Fither:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

# **Heparin and Antagonist Preparations**

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	31.28	10	<ul><li>Clexane</li></ul>
Inj 40 mg in 0.4 ml syringe	42.49	10	<ul><li>Clexane</li></ul>
Inj 60 mg in 0.6 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 80 mg in 0.8 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 100 mg in 1 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 120 mg in 0.8 ml syringe	125.87	10	<ul> <li>Clexane Forte</li> </ul>
Inj 150 mg in 1 ml syringe		10	<ul> <li>Clexane Forte</li> </ul>

#### ⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

**Initial application** — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	
HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ DBL Heparin
, -, ,			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
, =0,000 14 por, 01=	42.40	ŭ	✓ Heparin DBL S29
LIEDADINICED CALINE	72.70		Tiopaini bbe
HEPARINISED SALINE	GE 10	50	✓ Pfizer
Inj 10 iu per ml, 5 ml	05.48	ου	♥ Pilzer
Oral Anticoagulants			
Oral Anticoagalants			
DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ Pradaxa
Cap 110 mg	76.36	60	✓ Pradaxa
Cap 150 mg	76.36	60	✓ Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ Xarelto
Tab 20 mg		28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	✓ Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	✓ Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg		100	✓ Marevan
* Tab 5 mg		50	✓ Coumadin
ŭ	11.48	100	✓ Marevan
Blood Colony-stimulating Factors			
FILGRASTIM - Special Authority see SA1259 below - Retail p	,	10	Nivestim
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Nivestim to be Principal Supply on 1 December 2021			_

✓ Nivestim 10 Nivestim to be Principal Supply on 1 December 2021

### ⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10<sup>9</sup>/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10<sup>9</sup>/L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price)	s	Fully ubsidised	Brand or Generic
	\$	Per	•	Manufacturer
PEGFILGRASTIM - Special Authority see SA1912 below - Reta	il pharmacy			
Inj 6 mg per 0.6 ml syringe	1,080.00	1	✓ No	eulastim
⇒SA1912 Special Authority for Subsidy				

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

# Fluids and Electrolytes

Intravenous Administration			
GLUCOSE [DEXTROSE]  * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO  * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO  POTASSIUM CHLORIDE		5 1	✓ Biomed ✓ Biomed
* Inj 75 mg per ml, 10 ml	55.00	50	<ul> <li>✓ AstraZeneca</li> <li>✓ Potassium Chloride</li> <li>Aguettant \$29</li> </ul>
	65.00		✓ Juno
(AstraZeneca Inj 75 mg per ml, 10 ml to be delisted 1 November 20	021)		
(Potassium Chloride Aguettant §29 Inj 75 mg per ml, 10 ml to be	delisted 1 Nove	mber 2021)	
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			_
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
<ul><li>a) Up to 5 inj available on a PSO</li><li>b) Not in combination</li></ul>			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebuliser u for nebuliser use.	ise except wher	n used in conju	unction with an antibiotic intended
Inj 0.9%, bag - Up to 2000 ml available on a PSO	1.23	500 ml	✓ Baxter
	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, mate	rnity or post-na	tal care in the l	home of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs)	00.00	_	<b>4</b> D: 1
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standard	Formulae, page	20	✓ Fresenius Kabi
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		20 50	✓ Fresenius Kabi
Inj 0.9%, 20 ml ampoule		20	✓ Fresenius Kabi
ing 0.070, 20 ini ampoulo		20	- I TOOTHING HUDI

✓	fully subsidised
Pri	ncipal Supply

TOTAL PARENTERAL NUTRITION (TPN)

1 OP

✓ TPN

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

#### WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule - Up to 5 inj available on a PSO7.19	50	✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO5.00	20	✓ Fresenius Kabi
		✓ Multichem

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE  Powder169.85  COMPOUND ELECTROLYTES	300 g OP	✓ Calcium Resonium
Powder for oral soln — Up to 5 sach available on a PSO9.77	50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)6.55	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS		
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	60	
(11.85)		Chlorvescent
* Tab long-acting 600 mg (8 mmol)8.90	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	✓ Sodibic
		✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder	454 a OP	✓ Resonium-A

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

216.67

100

✓ Dibenzyline S29

# Alpha-Adrenoceptor Blockers

### Alpha Adrenoceptor Blockers

#### **DOXAZOSIN**

*	Tab 2 mg17.	.35	500	✓ Apo-Doxazosin
	Tab 4 mg20.		500	✓ Apo-Doxazosin
PH	IENOXYBENZAMINE HYDROCHLORIDE			
*	Cap 10 mg	.00	30	✓ BNM S29

#### PRAZOSIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking prazosin prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of prazosin.

*	Tab 1 mg5.53	100	✓ Apo-Prazosin
*	Tab 2 mg7.00	100	✓ Apo-Prazosin
*	Tab 5 mg11.70	100	✓ Apo-Prazosin

(Apo-Prazosin Tab 1 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 2 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 5 mg to be delisted 1 May 2022)

# Agents Affecting the Renin-Angiotensin System

### **ACE Inhibitors**

**CAPTOPRIL** 

T- - 0 F -

*	Oral liq 5 mg per ml	94.99	95 ml OP	✓ Capoten
		135.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

(Captopril-Mylan S29 Oral liq 5 mg per ml to be delisted 1 January 2022)

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

* Tab 0.5 mg	2.09	90	✓ ∠aprii
* Tab 2.5 mg		90	✓ Zapril
Tab 5 mg	8.35	90	✓ Zapril
ENALAPRIL MALEATE			
* Tab 5 mg	1.82	100	✓ Acetec
* Tab 10 mg	2.02	100	✓ Acetec
* Tab 20 mg	2.42	100	✓ Acetec
LISINOPRIL			
* Tab 5 mg	2.07	90	✓ Ethics Lisinopril
* Tab 10 mg	2.36	90	✓ Ethics Lisinopril
* Tab 20 mg	3.17	90	✓ Ethics Lisinopril

	Subsidy (Manufacturer's Price)	_	Fully	I Generic
PERMIT APPLI	\$	Per		Manufacturer
PERINDOPRIL Tob 2 mg	2.75	20	./	Ana Darindanril
Tab 2 mg		30		Apo-Perindopril Coversyl
Tab 4 mg		30		Apo-Perindopril
Tab 4 Hig	6.30	30		Coversyl
(Apo-Perindopril Tab 2 mg to be delisted 1 October 2021) (Apo-Perindopril Tab 4 mg to be delisted 1 October 2021) QUINAPRIL	0.30		·	Coversyi
* Tab 5 mg	5.97	90	✓	Arrow-Quinapril 5
* Tab 10 mg		90		Arrow-Quinapril 10
* Tab 20 mg	7.95	90	•	Arrow-Quinapril 20
ACE Inhibitors with Diuretics				
QUINAPRIL WITH HYDROCHLOROTHIAZIDE				
Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	•	Accuretic
<i>,</i>	3.83	30	✓	Accuretic 10
* Tab 20 mg with hydrochlorothiazide 12.5 mg	4.92	30	•	Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg	2.00	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021				
* Tab 8 mg	2.28	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021				
* Tab 16 mg	3.31	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021	5.00	00	,	0
* Tab 32 mg	5.26	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021				
LOSARTAN POTASSIUM				
* Tab 12.5 mg		84		Losartan Actavis
* Tab 25 mg		84		Losartan Actavis
* Tab 50 mg		84		Losartan Actavis
* Tab 100 mg	3.50	84	•	Losartan Actavis
Angiotensin II Antagonists with Diuretics				
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	•	Arrow-Losartan &
				Hydrochlorothiazide
Angiotensin II Antagonists with Neprilysin Inhii	bitors			
		Dot	nil pharma	01/
SACUBITRIL WITH VALSARTAN – Special Authority see SA19 Note: Due to the angiotensin II receptor blocking activity of ACE inhibitor or another ARB.				
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	/	Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg		56		Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg		56		Entresto 97/103
g				

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### **⇒SA1905** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III; or
  - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 120

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

### **Antiarrhythmics**

AMIODARONE HYDROCHLORIDE	,,	9	
▲ Tab 100 mg	3.80	30	✓ Aratac
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a	16.37	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	12.07	10	✓ Hameln S29
	15.09		Martindale
(Hameln S29 Inj 600 mcg per ml, 1 ml ampoule to be delisted 1 Ja	anuary 2022)		
DIGOXIN			
* Tab 62.5 mcg – Up to 30 tab available on a PSO		240	✓ Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO		240	Lanoxin
* Oral liq 50 mcg per ml	16.60	60 ml	<ul><li>✓ Lanoxin</li><li>✓ Lanoxin Paediatric</li></ul>
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	✓ Rythmodan
FLECAINIDE ACETATE			
▲ Tab 50 mg		60	Flecainide BNM
▲ Cap long-acting 100 mg	39.51	90	✓ <u>Flecainide</u> <u>Controlled</u> <u>Release Teva</u>
▲ Cap long-acting 200 mg	61.06	90	✓ <u>Flecainide</u> <u>Controlled</u> Release Teva
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
MENU ETIME UMPROCULORIDE	Ψ	rei		Manuacturer
MEXILETINE HYDROCHLORIDE	100.00	400	,	ANIL
▲ Cap 150 mg	162.00	100		ANI \$29 Mexiletine
			•	Hydrochloride
				USP S29
			1	Teva S29
▲ Cap 250 mg	202.00	100	1	Mexiletine
				Hydrochloride
			_	USP S29
			/	Teva S29
(ANI S29 Cap 150 mg to be delisted 1 January 2022)				
(Mexiletine Hydrochloride USP \$29 Cap 150 mg to be delisted	, ,			
(Mexiletine Hydrochloride USP S29 Cap 250 mg to be delisted	1 January 2022)			
PROPAFENONE HYDROCHLORIDE			_	
▲ Tab 150 mg	40.90	50	•	Rytmonorm
Antihypotensives				
MIDODRINE - Special Authority see SA1474 below - Retail pha	armacy			
Tab 2.5 mg		100	1	Gutron
Tab 5 mg	79.00	100	1	Gutron
<b>⇒SA1474</b> Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	d for 2 years where p	atien	has disab	ling orthostatic hypotension
not due to drugs.		n, ۱۱	martanaia	n abauld ba ayaidad and
Note: Treatment should be started with small doses and titrated the usual target is a standing systolic blood pressure of 90 mm H		ıy. n	ypertensio	n should be avoided, and
<b>Renewal</b> from any relevant practitioner. Approvals valid for 2 years		ent re	mains app	ropriate and the patient is
benefiting from treatment.				
Data Advanagantas Disakasa				
Beta-Adrenoceptor Blockers				
Beta Adrenoceptor Blockers				
ATENOLOL				
* Tab 50 mg		500		Mylan Atenolol
* Tab 100 mg		500		Mylan Atenolol Atenolol AFT
* Oral liq 25 mg per 5 ml	21.25 30	0 ml	-	Atenolol AFT
			•	S29 S29
	38.20		1	Essential
	00.20		•	=000.74441

Restricted to children under 12 years of age.

**BISOPROLOL FUMARATE** 

**CARVEDILOL** 

Generics \$29

✓ Bisoprolol Mylan

✓ Bisoprolol Mylan

✓ Bisoprolol Mylan

✓ Carvedilol Sandoz

✓ Carvedilol Sandoz✓ Carvedilol Sandoz

90

90

60

60

60

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

## Tab 200 mg			Subsidy		Fully	
ABETALOL			,	Per		
## Tab 100 mg	ΙΔ	RETALOI	· · · · · · · · · · · · · · · · · · ·			
# Tab 200 mg	-∧ *		14 50	100	1	Trandate
Find Sing per ml, 20 ml ampoule   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   5   1.59.06   1.59.06   5   1.59.06   1.59	*	•				
(88.60)   Trandate	*	•				
According   Acco	•	, og po, =0 apoao		ŭ		Trandate
METOPROLOL SUCCINATE	*	ini 5 mg per ml. 20 ml vial	\ /	1		
## Tab long-acting 23.75 mg		, - 3r - ,				Alvogen S29
## Tab long-acting 47.5 mg	ME	TOPROLOL SUCCINATE	, ,			· ·
Tab long-acting 47.5 mg	*	Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
## Tab long-acting 95 mg	*			30	1	Betaloc CR
Tab 50 mg	*			30	✓	Betaloc CR
Tab 50 mg	*	Tab long-acting 190 mg	4.27	30	✓	Betaloc CR
Tab 50 mg	MF					
Tab 100 mg			5.66	100	/	Apo-Metoprolol
## Tab long-acting 200 mg		· ·				
INDOLOL Tab 40 mg	*	•		28		
Tab 40 mg	*			5		•
Tab 80 mg	NA	DOLOL				
Tab 80 mg		Tab 40 mg	16.69	100	✓	Apo-Nadolol
Subsidy by endorsement – Subsidised for patients who were taking pindolol prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of pindolol.  **Tab 5 mg				100		•
Subsidy by endorsement – Subsidised for patients who were taking pindolol prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of pindolol.  **Tab 5 mg	PII	NDOLOL - Subsidy by endorsement				•
dispensing of pindolol.  # Tab 5 mg	-	Subsidy by endorsement – Subsidised for patients who we	01		0	' '
# Tab 10 mg		, ,				
# Tab 15 mg	*	Tab 5 mg	13.22	100	✓	Apo-Pindolol
Apo-Pindolol Tab 5 mg to be delisted 1 May 2022)  Apo-Pindolol Tab 10 mg to be delisted 1 May 2022)  Apo-Pindolol Tab 15 mg to be delisted 1 May 2022)  ROPRANOLOL  Tab 10 mg	*	Tab 10 mg	23.12	100	✓	Apo-Pindolol
Apo-Pindolol Tab 10 mg to be delisted 1 May 2022)  Apo-Pindolol Tab 15 mg to be delisted 1 May 2022)  ROPRANOLOL  Tab 10 mg	*	Tab 15 mg	33.31	100	✓	Apo-Pindolol
Apo-Pindolol Tab 15 mg to be delisted 1 May 2022)  ROPRANOLOL  Tab 10 mg	(A <sub>l</sub>	po-Pindolol Tab 5 mg to be delisted 1 May 2022)				
ROPRANOLOL	(A)	po-Pindolol Tab 10 mg to be delisted 1 May 2022)				
Tab 10 mg       4.64       100       ✓ Apo-Propranolol         Tab 40 mg       5.72       100       ✓ Apo-Propranolol         € Cap long-acting 160 mg       18.17       100       ✓ Cardinol LA         © Oral liq 4 mg per ml       Special Authority see SA1327 below –       CBS       500 ml       ✓ Roxane-	(A)	po-Pindolol Tab 15 mg to be delisted 1 May 2022)				
Tab 40 mg	P	OPRANOLOL				
Tab 40 mg			4.64	100	1	Apo-Propranolol
<ul> <li>Cap long-acting 160 mg</li></ul>		9		100		
Foral liq 4 mg per ml − Special Authority see SA1327 below − Retail pharmacyCBS 500 ml Foxane-	*	· ·		100		
Retail pharmacyCBS 500 ml ✓ Roxane-	*					
·				500 m	nl 🗸	Roxane-
		,				Propranolol \$29

### ⇒SA1327 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

	Subsidy lanufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OTALOL				
Tab 80 mg	32.58	500	✓	<u>Mylan</u>
Tab 160 mg	10.98	100	1	<u>Mylan</u>
Calcium Channel Blockers				
Dihydropyridine Calcium Channel Blockers				
MLODIPINE			_	
Tab 2.5 mg		90		<u>Vasorex</u>
Tab 5 mg		90		Vasorex
Tab 10 mg	1.19	90	•	<u>Vasorex</u>
ELODIPINE				
Tab long-acting 2.5 mg		30		Plendil ER
Tab long-acting 5 mg		90		Felo 5 ER
Tab long-acting 10 mg	4.32	90	•	Felo 10 ER
FEDIPINE				
Tab long-acting 10 mg	18.80	56	✓	Tensipine MR10 S29
Tab long-acting 20 mg	9.12	50	1	Mylan (12 hr
<u>Jgg</u>				release) S29
	17.72	100	1	Nyefax Retard
Tab long-acting 30 mg		100		Mylan (24 hr
rab long dolling oo mg	07.10	100	•	
Tab long acting 60 mg	50.01	100	./	release) \$29 Mylan (24 br
Tab long-acting 60 mg	32.81	100	•	Mylan (24 hr
				release) \$29
Other Calcium Channel Blockers				
LTIAZEM HYDROCHLORIDE			_	
Tab 60 mg		100		Dilzem
Tab 60 mg Cap long-acting 120 mg	33.42	500	1	Apo-Diltiazem CD
Tab 60 mg Cap long-acting 120 mg Cap long-acting 180 mg	33.42 50.05	500 500	<b>√</b>	Apo-Diltiazem CD Apo-Diltiazem CD
Tab 60 mg	33.42 50.05	500	<b>√</b>	Apo-Diltiazem CD
Tab 60 mg	33.42 50.05	500 500	<b>√</b>	Apo-Diltiazem CD Apo-Diltiazem CD
Tab 60 mg	33.42 50.05 66.76	500 500 500	<i>y y</i>	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD
Tab 60 mg	33.42 50.05 66.76	500 500	<i>y y</i>	Apo-Diltiazem CD Apo-Diltiazem CD
Tab 60 mg	33.42 50.05 66.76	500 500 500	<i>y y y</i>	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD Pexsig
Tab 60 mg	33.42 50.05 66.76	500 500 500 100	<i>y y y</i>	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD Pexsig
Tab 60 mg	33.42 50.05 66.76 62.90	500 500 500	· · · · · · · · · · · · · · · · · · ·	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD Pexsig Isoptin Isoptin
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74	500 500 500 100	· · · · · · · · · · · · · · · · · · ·	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD Pexsig
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100	· · · · · · · · · · · · · · · · · · ·	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100	· · · · · · · · · · · · · · · · · · ·	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD Pexsig Isoptin Isoptin Isoptin Retard 529
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100 100	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100 100	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100 100 30	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02	500 500 500 100 100 100 30	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02 15.12	500 500 500 100 100 100 30 5	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard 229 Isoptin SR Isoptin SR
Tab 60 mg	33.42 50.05 66.76 62.90 7.01 11.74 36.02 15.12 25.00	500 500 500 100 100 100 30	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Apo-Diltiazem CD Apo-Diltiazem CD Apo-Diltiazem CD  Pexsig  Isoptin Isoptin Isoptin Retard \$29 Isoptin SR Isoptin SR

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	
CLONIDINE HYDROCHLORIDE				
<b>★</b> Tab 25 mcg	8.75	112	1	Clonidine BNM
₭ Tab 150 mcg		100	1	Catapres
K Inj 150 mcg per ml, 1 ml ampoule	29.68	10	/	Medsurge
METHYLDOPA				•
k Tab 250 mg	15 10	100	1	Methyldopa Mylan
140 200 ng	52.85	500		Methyldopa Mylan S29 S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
<b>₭</b> Tab 1 mg	4.91	30	1	Burinex S29 S29
	16.36	100	1	Burinex
₭ Inj 500 mcg per ml, 4 ml vial	7.95	5	✓	Burinex
UROSEMIDE [FRUSEMIDE]				
Tab 40 mg – Up to 30 tab available on a PSO	7 24	1,000	) 🗸	Apo-Furosemide
* Tab 500 mg		50		Urex Forte
	89.48			Furosemid-
	100.00	400	,	Ratiopharm S29
	169.96	100	V	Furosemid- Ratiopharm S29
★ Oral lig 10 mg per ml	11.20	30 ml C	OP 🗸	Lasix
k Inj 10 mg per ml, 25 ml ampoule		6		Lasix
Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	PSO 1.15	5	•	Furosemide-Baxter
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE			_	
Oral liq 1 mg per ml	30.00	25 ml (	OP 🗸	Biomed
EPLERENONE - Special Authority see SA1728 below - Retail p	harmacy			
Tab 50 mg	17.00	30	✓	Inspra
Tab 25 mg	11.87	30	✓	Inspra
⇒SA1728 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals valid ne following criteria:	d without further ren	ewal u	nless notif	ied for applications meet
Soth: 1 Patient has heart failure with ejection fraction less than 40	%; and			
<ul><li>2 Either:</li><li>2.1 Patient is intolerant to optimal dosing of spironolac</li></ul>	tone; or			
2.2 Patient has experienced a clinically significant adve		optima	al dosing o	f spironolactone.
METOLAZONE				

1 50 ✓ Metolazone S29

✓ Zaroxolyn S29

	Subsidy		Fully Brand or
	(Manufacturer's Pric	e) Subsid	dised Generic
DIDONOL ACTONE	\$	Per	✓ Manufacturer
PIRONOLACTONE ← Tab 25 mg	4 38	100	✓ Spiractin
€ Tab 100 mg		100	✓ Spiractin
Oral liq 5 mg per ml		25 ml OP	✓ Biomed
Potassium Sparing Combination Diuretics			
• •			
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE	0.00	00	/ F!!
Tab 5 mg with furosemide 40 mg		28	✓ Frumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIA			<b></b>
Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓ Moduretic
Thiazide and Related Diuretics			
ENDROFLUMETHIAZIDE [BENDROFLUAZIDE]			
Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ <u>Arrow-</u>
			<u>Bendrofluazide</u>
May be supplied on a PSO for reasons other than eme	raencv.		
F Tab 5 mg	• ,	500	✓ Arrow-
•			Bendrofluazide
HLOROTHIAZIDE	06.00	05 ml OD	✓ Biomed
Oral liq 50 mg per ml	20.00	25 ml OP	▼ bioinea
HLORTALIDONE [CHLORTHALIDONE]	2.22	00	<b>41</b>
Tab 25 mg		30	✓ Igroton S29
	6.50	50	✓ <u>Hygroton</u>
NDAPAMIDE ₹ Tab 2.5 mg	10.45	90	✓ Dapa-Tabs
1 ab 2.5 mg	11.61	100	✓ <u>Dapa-Tabs</u> ✓ Mylan
	11.01	100	Indapamide \$29
Lipid-Modifying Agents			
Fibrates			
EZAFIBRATE			
EZAFIDNATE Tab 200 mg	19.46	90	✓ Bezalip
Tab long-acting 400 mg		30	✓ Bezalip Retard
			•
Other Lipid-Modifying Agents			
CIPIMOX	0:		
Cap 250 mg	21.56	30	✓ Olbetam
			✓ Olbetam S29 S29
Resins			
OLESTIPOL HYDROCHLORIDE			
Grans for oral liq 5 g	32.89	30	✓ Colestid

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg	6.16	500	<b>√</b> L	orstat
Lorstat to be Principal Supply on 1 December 2021				
* Tab 20 mg	9.24	500	✓L	orstat
Lorstat to be Principal Supply on 1 December 2021	14.00	F00		
* Tab 40 mg Lorstat to be Principal Supply on 1 December 2021	14.92	500	<b>V</b> L	orstat
* Tab 80 mg	26 54	500	<b>√</b> I	orstat
Lorstat to be Principal Supply on 1 December 2021	20.04	000		orotat
PRAVASTATIN				
* Tab 20 mg	2.11	28	<b>√</b> P	ravastatin Mylan
* Tab 40 mg		28	_	ravastatin Mylan
SIMVASTATIN			_	<del></del> _
* Tab 10 mg	1.23	90	<b>√</b> S	Simvastatin Mylan
* Tab 20 mg		90	<b>√</b> <u>S</u>	imvastatin Mylan
* Tab 40 mg	3.58	90	_	imvastatin Mylan
* Tab 80 mg	7.12	90	<b>√</b> <u>s</u>	Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors				

# ⇒SA1045 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 x normal) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

✓ Zimybe	30	vastatin 10 mg5.15	Tab 10 mg with
✓ Zimybe	30	vastatin 20 mg6.15	Tab 10 mg with
✓ Zimybe	30	vastatin 40 mg7.15	Tab 10 mg with
✓ Zimybe	30	vastatin 80 mg8.15	Tab 10 mg with

30

✓ Ezetimibe Sandoz

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	•	Manufacturer

### ⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

#### **Nitrates**

GLYCERYL TRINITRATE			
* Oral pump spray, 400 mcg per dose - Up to 250 dose	Э		
available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
ISOSORBIDE MONONITRATE			
* Tab 20 mg	19.55	100	✓ Ismo 20
* Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
* Tab long-acting 60 mg	9.25	90	✓ Duride

Sym	path	omim	netics

ADRENALINE		
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	✓ Aspen Adrenaline

### **Vasodilators**

#### HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 on the next page -		
	Retail pharmacyCBS	1	<ul> <li>Hydralazine</li> </ul>
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink S29
*	Inj 20 mg ampoule25.90	5	✓ Apresoline

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

### ⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL			
▲ Tab 10 mg70	.00	100	Loniten
NICORANDIL			
▲ Tab 10 mg25	.57	60	✓ Ikorel
▲ Tab 20 mg32	.28	60	/ Ikorel
PAPAVERINE HYDROCHLORIDE			_
* Inj 12 mg per ml, 10 ml ampoule217	.90	5	<ul><li>Hospira</li></ul>
PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg42	.26	50	✓ Trental 400

### **Endothelin Receptor Antagonists**

AMBRISENTAN - Special Authority see SA1/02 below - R	letail pharmacy		
Tab 5 mg	1,550.00	30	<ul> <li>Ambrisentan Mylan</li> </ul>
Tab 10 mg	1,550.00	30	✓ Ambrisentan Mylan

### ⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1991 below - Retail pharmacy

Tab 62.5 mg	119.85	60	✓ Bosentan Dr Reddy's
Bosentan Dr Reddy's to be Principal Supply on 1 Tab 125 mg		60	✓ Bosentan Dr Reddy's

Bosentan Dr Reddy's to be Principal Supply on 1 December 2021

### ⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 Bosentan is to be used as PAH monotherapy; and
    - 4.1.2 Either:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	•	

continued...

- 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
- 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
- 4.2 Both:
  - 4.2.1 Bosentan is to be used as PAH dual therapy; and
  - 4.2.2 Either:
    - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
    - 4.2.2.2 Patient deteriorated while on a PAH monotherapy: or
- 4.3 Both:
  - 4.3.1 Bosentan is to be used as PAH triple therapy; and
  - 4.3.2 Any of the following:
    - 4.3.2.1 Patient is on the lung transplant list; or
    - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
    - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
    - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

**Renewal** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Bosentan is to be used as PAH monotherapy; and
  - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
  - 2.1 Bosentan is to be used as PAH dual therapy; and
  - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
  - 3.1 Bosentan is to be used as PAH triple therapy; and
  - 3.2 Any of the following:
    - 3.2.1 Patient is on the lung transplant list; or
    - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
    - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
    - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

# Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail pharmacy		
Tab 25 mg	4	✓ Vedafil
Tab 50 mg1.70	4	✓ Vedafil
Tab 100 mg10.20	12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Subsidy			Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 Any of the following:
  - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
  - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
  - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
  - 3.1 PAH is in NYHA/WHO functional class II; or
  - 3.2 PAH is in NYHA/WHO functional class III; or
  - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.2 Either:
      - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
      - 4.1.2.2 Patient is peri Fontan repair; and
    - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
  - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with \* are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
  - Patient has a decommend history of tradmatte of non-traditional spiral cord injury, and
     Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Prostacyclin Analogues**



### ⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

✓ Ventavis

30

	Subsidy (Manufacturer's Price) \$	- · · · · · <b>/</b>		Brand or Generic Manufacturer
ILOPROST - Special Authority see SA1705 below - Retail phar	macy			

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Nebuliser soln 10 mcg per ml, 2 ml .......740.10

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- b) Only on a prescription

Crm 0.1%	22.89	30 g OP	<ul><li>Differin</li></ul>
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below -	Retail pharmacy	· ·	
Cap 5 mg	8.14	60	<ul><li>Oratane</li></ul>
Cap 10 mg	13.34	120	<ul><li>Oratane</li></ul>
Cap 20 mg	20.49	120	<ul><li>Oratane</li></ul>

#### ⇒SA2023 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

#### **TRFTINOIN**

Crm 0.5 mg per g − Maximum of 50 g per prescription ......15.57 50 g OP ✓ ReTrieve

# **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

HYDROGEN PEROXIDE

111	DITOGEN I ETIONIDE		
*	Crm 1%8.56	10 g OP	<ul><li>Crystaderm</li></ul>
		15 g OP	<ul> <li>Crystaderm</li> </ul>

	Subsidy (Manufacturer's P \$	rice) Subs	Fully Brand or sidised Generic  Manufacturer
MUPIROCIN			
Oint 2%	6.60 (10.50)	15 g OP	Bactroban
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
SODIUM FUSIDATE [FUSIDIC ACID]  Crm 2%	1.59	5 g OP	✓ Foban
<ul><li>a) Maximum of 5 g per prescription</li><li>b) Only on a prescription</li><li>c) Not in combination</li></ul>			
d) Foban to be Principal Supply on 1 December 2021 Oint 2%		5 g OP	✓ Foban
<ul> <li>a) Maximum of 5 g per prescription</li> <li>b) Only on a prescription</li> <li>c) Not in combination</li> <li>d) Foban to be Principal Supply on 1 December 2021</li> </ul>	ı		
SULFADIAZINE SILVER Crm 1%	10.80	50 g OP	✓ Flamazine
a) Up to 250 g available on a PSO     b) Not in combination		50 g 51	- Fundanc
Antifungals Topical			
For systemic antifungals, refer to INFECTIONS, Antifungals, p	age 97		
AMOROLFINE	_		
a) Only on a prescription     b) Not in combination Nail soln 5%	14.93	5 ml OP	✓ MycoNail
CICLOPIROX OLAMINE			
a) Only on a prescription     b) Not in combination			
Nail-soln 8%	5.72	7 ml OP	✓ Apo-Ciclopirox
CLOTRIMAZOLE  * Crm 1%	0.77	20 g OP	✓ Clomazol
a) Only on a prescription		- <b>3</b> -	
b) Not in combination  * Soln 1%	4.36	20 ml OP	
	(7.55)		Canesten
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
ECONAZOLE NITRATE	4.00	00 00	
Crm 1%	1.00 (7.48)	20 g OP	Pevaryl
a) Only on a prescription	( - /		· · · <b>,</b>
b) Not in combination Foaming soln 1%, 10 ml sachets	9.89	3	
1 Saming Som 170, 10 mil Saurioto	(17.23)	3	Pevaryl
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g OP	✓ N	<u>Multichem</u>
a) Only on a prescription				
b) Not in combination				
* Lotn 2%		30 ml OP	_	·
	(10.03)		L	Daktarin
a) Only on a prescription     Net in combination				
b) Not in combination  * Tinct 2%	4.26	30 ml OP		
T ΠΟΙ Δ /0	(12.10)	JU IIII UP	Г	Daktarin
a) Only on a prescription	(12.10)			
b) Not in combination				
,				
Antipruritic Preparations				
CALAMINE				
a) Only on a prescription				
b) Not in combination	4.00	100		
Crm, aqueous, BP	1.26	100 g	✓ h	nealthE Calamine
				Aqueous Cream BP
CROTAMITON				DF
a) Only on a prescription				
b) Not in combination				
Crm 10%	3.29	20 g OP	✓ II	tch-Soothe
Itch-Soothe to be Principal Supply on 1 December 2021		_0 g Oi	- 11	•••••
MENTHOL – Only in combination				
Only in combination with a dermatological base or prop     With or without other dermatological galenicals.	orietary Topical Co	orticosteriod –	Plain	
Crystals	6.92	25 g	<b>✓</b> N	/lidWest
-,	29.60	100 g		/lidWest
0				
Corticosteroids Topical				
For systemic corticosteroids, refer to CORTICOSTEROIDS AND	RELATED AGEN	NTS, page 80		
Corticosteroids - Plain				
BETAMETHASONE DIPROPIONATE				
Crm 0.05%	2.96	15 g OP	_	Diprosone
	36.00	50 g OP	_	<u>Diprosone</u>
Oint 0.05%		15 g OP	_	<u>Diprosone</u>
0:10.050/:	36.00	50 g OP	_	Diprosone
Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ [	Diprosone OV
BETAMETHASONE VALERATE			_	
* Crm 0.1%	4.53	50 g OP	<b>✓</b> E	Beta Cream

CLOBETASOL PROPIONATE

Crm 0.05%......2.18

Oint 0.05%......2.12

✓ Beta Ointment

✓ Betnovate

✓ <u>Dermol</u>
✓ Dermol

50 g OP

50 ml OP

30 g OP

30 g OP

Subsidity   Subsidity   Subsidity   Subsidity   Subsidition   Subsidit					
S   Per			)	•	
CLOBETASONE BUTYRATE					er
Crm 0.05%	CLOBETASONE BUTYRATE	·			
HYDROCORTISONE   # Crm 1% - Only on a prescription.   3.70   100 g OP		5.38	30 a OP		
# Crm 1% - Only on a prescription			55 g 5.	Eumovate	
# Crm 1% - Only on a prescription	HYDROCORTISONE	(13133)			
Powder - Only in combination		3.70	100 a OP	✓ Hydrocortise	one
Powder - Only in combination	The Strip of a procomption		100 g O1		<u> </u>
# Powder – Only in combination		17.15	500 a		one
Up to 5% in a dermatological base (not proprietary Topical Corticosteriod − Plain) with or without other dermatological galenicals  HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN  Loto 1% with paraffin liquid 15.9% and lanolin 0.6% − Only on a prescription			3	_	
galenicals HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% — Only on a prescription	* Powder – Only in combination	49.95	25 g	✓ ABM	
Selection   Control   Co				r without other deri	matological
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% − Only on a prescription			,		· ·
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% − Only on a prescription	HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
a prescription		n			
HYDROCORTISONE BUTYRATE			250 ml	✓ DP Lotn HC	
Lipocream 0.1%	• •				
Dint 0.1%		4.85	100 a OP	✓ Locoid Lipo	cream
Locoid to be Principal Supply on 1 December 2021  Milky emul 0.1%			•		oroun.
Milky emul 0.1%					
Locoid Crelo to be Principal Supply on 1 December 2021	,	12.33	100 ml OP	✓ Locoid Crelo	<b>o</b>
METHYLPREDNISOLONE ACEPONATE  Crm 0.1%	•				
Crm 0.1%       4.46       15 g OP       ✓ Advantan         Oint 0.1%       4.46       15 g OP       ✓ Advantan         MOMETASONE FUROATE	METHYL PREDNISOLONE ACEPONATE				
Oint 0.1%		4.46	15 a OP	✓ Advantan	
Crm 0.1%	Oint 0.1%	4.46	•		
Crm 0.1%	MOMETASONE FUROATE		ŭ	<del></del>	
3.10   50 g OP		1.95	15 a OP	✓ Elocon Alco	hol Free
Oint 0.1%					
Lotn 0.1%	Oint 0.1%	1.95	•	✓ Elocon	
TRIAMCINOLONE ACETONIDE  Crm 0.02%			50 g OP	✓ Elocon	
Crm 0.02%	Lotn 0.1%	4.50	30 ml OP	✓ Elocon	
Oint 0.02%	FRIAMCINOLONE ACETONIDE				
Corticosteroids - Combination  BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID] Crm 0.1% with sodium fusidate (fusidic acid) 2%	Crm 0.02%	6.30	100 g OP	✓ Aristocort	
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]  Crm 0.1% with sodium fusidate (fusidic acid) 2%	Oint 0.02%	6.35	100 g OP	✓ Aristocort	
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]  Crm 0.1% with sodium fusidate (fusidic acid) 2%					
Crm 0.1% with sodium fusidate (fusidic acid) 2%	Corticosteroids - Combination				
Crm 0.1% with sodium fusidate (fusidic acid) 2%	 	SIDIC VCIDI			
(10.45) Fucicort  a) Maximum of 15 g per prescription b) Only on a prescription  HYDROCORTISONE WITH MICONAZOLE – Only on a prescription  ** Crm 1% with miconazole nitrate 2%	•	•	15 a OP		
a) Maximum of 15 g per prescription b) Only on a prescription HYDROCORTISONE WITH MICONAZOLE – Only on a prescription  ★ Crm 1% with miconazole nitrate 2%	Offit 0.176 with Social Hasiatic (lastate acid) 276		10 g O1	Fucicort	
b) Only on a prescription  HYDROCORTISONE WITH MICONAZOLE – Only on a prescription  ** Crm 1% with miconazole nitrate 2%	a) Maximum of 15 g per prescription	(10.10)		radioon	
HYDROCORTISONE WITH MICONAZOLE — Only on a prescription  ★ Crm 1% with miconazole nitrate 2%	, , , ,				
* Crm 1% with miconazole nitrate 2%	, , ,	tion			
Micreme H to be Principal Supply on 1 December 2021  HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN − Only on a prescription  Crm 1% with natamycin 1% and neomycin sulphate 0.5%			15 a OP	✓ Micrama H	
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Only on a prescription  Crm 1% with natamycin 1% and neomycin sulphate 0.5%		1.03	13 y OF	A MINOLETTIC U	
Crm 1% with natamycin 1% and neomycin sulphate 0.5%		nly on a procesis	ation		
Oint 1% with natamycin 1% and neomycin sulphate 0.5%				✓ Dimofuoort	
			•		

DERMATOLOGICALS			
	Subsidy (Manufacturer's   \$		Fully Brand or dised Generic  Manufacturer
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMY Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 and gramicidin 250 mcg per g - Only on a prescriptio	mg n3.49	TIN 15 g OP	
	(9.28)		Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE  * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE  Dimethicone 10%
ZINC AND CASTOR OIL  * Oint	4.25	500 g	✓ Boucher
Emollients			
* Crm	1.92	500 g	<ul><li>✓ Basic AquaCream</li><li>✓ Boucher</li><li>✓ Medco</li></ul>
CETOMACROGOL  * Crm BP  CETOMACROGOL WITH GLYCEROL	2.48	500 g	✓ healthE
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ Boucher ✓ Kenkay Sorbolene ✓ Pharmacy Health Sorbolene with Glycerin
	3.10	1,000 ml OP	✓ ADE ✓ Boucher
* Oint BP	3.40	500 g	✓ <u>Emulsifying</u> <u>Ointment ADE</u>
OIL IN WATER EMULSION  * Crm	2.19	500 g	✓ O/W Fatty Emulsion Cream

Oint liquid paraffin 50% with white soft paraffin 50%......5.35

500 ml OP

100 g OP

✓ healthE

✓ healthE Urea Cream

**PARAFFIN** 

UREA

	Subsidy		Fully Brand or
	(Manufacturer's F		idised Generic
	\$	Per	✓ Manufacturer
NOOL FAT WITH MINERAL OIL - Only on a prescription			
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
·	(11.95)		DP Lotion
	1.40	250 ml OP	
	(4.53)		DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91)		BK Lotion
	1.40	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft - Only in combination	4.99	450 g	✓ healthE
•	19.99	2,500 g	✓ healthE

### **Minor Skin Infections**

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	2.55	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
·	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

# **Parasiticidal Preparations**

DIMETHICONE

* Lotn 4%	4.98	200 ml OP	✓ <u>healthE</u> <u>Dimethicone 4%</u> <u>Lotion</u>
IVERMECTIN - Special Authority see SA1225 below - F	letail pharmacy		
Tab 3 mg - Up to 100 tab available on a PSO	17.20	4	✓ Stromectol

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

### **⇒SA1225** Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:



-					
	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Su	bsidised	Generic	
	\$	Per	/	Manufacturer	

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
    - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

	Subsidy	F	ully	Brand or
(Man	ufacturer's Price)	Subsidi	sed	Generic
	\$	Per	•	Manufacturer

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

**Renewal** — **(Other parasitic infections)** only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

#### **PERMETHRIN**

Crm 5%	•	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN		
Shampoo 0.5%11.36	200 ml OP	Parasidose
(Parasidose Shampoo 0.5% to be delisted 1 January 2022)		

## **Psoriasis and Eczema Preparations**

		ACITRETIN - Special Authority see SA2024 below - Retail pharmacy	Α
✓ Novatretin	60	Cap 10 mg17.86	
✓ Novatretin	60	Cap 25 mg41.36	

#### ⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

#### BETAMETHASONE DIPROPIONATE WITH CAI CIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per gGel 500 mcg with calcipotriol 50 mcg per g		60 g OP 60 g OP	<ul><li>✓ Enstilar</li><li>✓ Daivobet</li></ul>
Daivobet to be Principal Supply on 1 December 2021 Oint 500 mcg with calcipotriol 50 mcg per g Daivobet to be Principal Supply on 1 December 2021		30 g OP	✓ Daivobet
CALCIPOTRIOL Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex

	Subsidy (Manufacturer's F	Price) Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
COAL TAR				
Soln BP – Only in combination		200 ml	_	<u>lidwest</u>
<ol> <li>Up to 10% only in combination with a dermatolo</li> <li>With or without other dermatological galenicals.</li> </ol>	gical base or propri	etary Topical (	Corticos	teriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SU	JLPHUR			
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5%				
allantoin crm 2.5%		75 g OP	_	
	(8.00) 3.43	20 a OB	E	Egopsoryl TA
	(4.35)	30 g OP	F	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR	(4.00)		_	gopooryi 174
Soln 12% with salicylic acid 2% and sulphur 4% oint	<i>1</i> 97	25 g OP	<b>/</b> (	Coco-Scalp
Con 12/0 with Sancy ite acid 2/0 and Sulphur 4/0 cint	7.95	40 g OP		Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Re		.0 9 0.		, 000 00a.p
a) Maximum of 15 g per prescription	tali priarriacy			
b) Note: a maximum of 15 g per prescription and no mor	e than one prescrip	tion per 12 we	eks.	
Cream 1%		15 g OP		ilidel
⇒SA1970 Special Authority for Subsidy		Ü	_	
of a dermatologist, paediatrician or ophthalmologist. Approval meeting the following criteria:  Both:	3 valid without furth	ici iciicwai diii	033 1101	med for applications
<ol> <li>Patient has atopic dermatitis on the eyelid; and</li> </ol>				
<ul> <li>Patient has atopic dermatitis on the eyelid; and</li> <li>Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.</li> </ul>				
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to	topical corticostero	ids, cataracts,	glaucor	
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.	topical corticostero	ids, cataracts,	glaucoi n	
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure. PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOR	topical corticostero	ids, cataracts,	glaucoi n	ma, or raised intraocular
<ul> <li>2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.</li> <li>PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOR</li> <li>Soln 2.3% with trolamine laurilsulfate and fluorescein sodi</li> </ul>	topical corticostero	ids, cataracts,	glaucor n ✓ <u>F</u>	ma, or raised intraocular  Pinetarsol  Aidwest
Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID	topical corticostero RESCEIN - Only o um444	n a prescriptio 500 ml	glaucon	ma, or raised intraocular  Pinetarsol  Aidwest  SM
Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF  ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID  Powder – Only in combination	topical corticostero RESCEIN - Only o um444	n a prescriptio 500 ml	glaucon	ma, or raised intraocular  Pinetarsol  Aidwest  SM
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder – Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic	n a prescriptio 500 ml	glaucon n	ma, or raised intraocular  Pinetarsol  Aidwest  SM
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder – Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostere	glaucon  P  N  P  poid – Pla	ma, or raised intraocular  Pinetarsol  Midwest  PSM  ain or collodion flexible
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder — Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostere	glaucon  P  N  P  poid – Pla	ma, or raised intraocular  Pinetarsol  Midwest  PSM  ain or collodion flexible
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder — Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostere	glaucon  P  N  P  poid – Pla	ma, or raised intraocular  Pinetarsol  Midwest  PSM  ain or collodion flexible
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder — Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic6.35 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostere	glaucon  P  N  P  poid – Pla	ma, or raised intraocular  Pinetarsol  Midwest  PSM  ain or collodion flexible
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder — Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic6.35 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostere	glaucon  Pi  N  Pi  Did – Pla	ma, or raised intraocular  Pinetarsol  Midwest  PSM  ain or collodion flexible
2 Patient has at least one of the following contraindication documented epidermal atrophy, documented allergy to pressure.  PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUOF ** Soln 2.3% with trolamine laurilsulfate and fluorescein sodi SALICYLIC ACID Powder — Only in combination	topical corticostero RESCEIN - Only o um4.4418.88 or proprietary Topic6.35 or proprietary Topic	ids, cataracts, n a prescriptio 500 ml 250 g cal Corticostero 100 g cal Corticostero	glaucon  Pi  N  Pi  Did – Pla	ma, or raised intraocular  Pinetarsol  Aidwest  PSM  ain or collodion flexible  Aidwest  ain

### **DERMATOLOGICALS**

	Subsidy (Manufacturer's \$	Price) Sub	Fully sidised	Brand or Generic Manufacturer
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	6.57	100 ml OP	<b>✓</b> L	ocoid
Locoid to be Principal Supply on 1 December 2021 KETOCONAZOLE				
Shampoo 2%	3.23	100 ml OP	<b>√</b> <u>S</u>	<u>sebizole</u>

### **Sunscreens**

# **Wart Preparations**

### **Other Skin Preparations**

# Antineoplastics

# **GENITO-URINARY SYSTEM**

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

Brand or

Generic

Manufacturer

Subsidy Fully
(Manufacturer's Price) Subsidised

\$ Per 
✓

# **Contraceptives - Non-hormonal**

### **Condoms**

-	IDOMS			
	49 mm - Up to 144 dev available on a PSO		144	✓ <u>Moments</u>
	53 mm		10	✓ <u>Moments</u>
	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	11.64	144	✓ <u>Moments</u>
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO		, .	
	53 mm, 0.05 mm thickness		10	✓ <u>Moments</u>
	\	11.42	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	2.05	40	<b>/ 11</b>
	53 mm, chocolate, brown		10	✓ Moments
	-\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	0.05	40	/ Name : :- te
	53 mm, strawberry, red		10	✓ Moments
	.) 11-1-00 1	11.64	144	✓ <u>Moments</u>
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription	2.27	40	<b>/ 11</b>
	56 mm		10	✓ Moments
		11.64	144	✓ <u>Moments</u>
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
	56 mm, 0.05 mm thickness		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	<ul> <li>a) Maximum of 60 dev per prescription</li> </ul>			
	b) Up to 60 dev available on a PSO			_
	56 mm, 0.08 mm thickness		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, 0.08 mm thickness, red		10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, chocolate		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	56 mm, strawberry	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
	60 mm	1.42	12	✓ Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

b) Up to 60 dev available on a PSO

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	<b>√</b> <u>(</u>	Gold Knight XL
	a) Maximum of 60 dev per prescription				

### **Contraceptive Devices**

#### INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IÚD 29.1 mm length × 23.2 mm width	18.45	1	✓	Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓	Choice
					TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	1	Choice Load 375

### **Contraceptives - Hormonal**

### **Combined Oral Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

**Renewal** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to		
	84 tab available on a PSO10.00	84	✓ Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab6.62	84	
	(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

(Marvelon 28 Tab 30 mcg with desogestrel 150 mcg and 7 inert tab to be delisted 1 November 2021)

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer
-			
2.18	84	✓ M	licrogynon 20 ED
6.45	112	✓ F	emme-Tab ED
Jp			
·9.45	84	✓ M	licrogynon 50 ED
6.62	63		
(16.50)		M	licrogynon 30
thority see SA0500 on	the pr	evious pag	е
•	•		
_			
	84	<b>√</b> L	evlen ED
6.45	112	✓ F	emme-Tab ED
•			
	0.4	<b>/</b> D	
	84	ν <u>Β</u>	revinor 1/28
•			
21.99	84	✓ N	orimin
	(Manufacturer's Price) 2.18 6.45  Jp9.456.62 (16.50) thority see SA0500 on 1.77	(Manufacturer's Price) \$ Per \$	(Manufacturer's Price)     Subsidised Per       -

### **Progestogen-only Contraceptives**

### ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither
  - 1.1 Patient is on a Social Welfare benefit; or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ᄕ	VONORGESTREL			
*	Tab 30 mcg – Up to 84 tab available on a PSO	.50	84	✓ Microlut
	22.0	.00 1	12	✓ Microlut
*	Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
	on a PSO106.9	.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO7.9	.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	6.25	84	<b>√</b> 1	Noriday 28	
<b>Emergency Contraceptives</b>					
LEVONORGESTREL  * Tab 1.5 mg	4.95	1	<b>√</b> F	Postinor-1	

### **Antiandrogen Oral Contraceptives**

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

### **Gynaecological Anti-infectives**

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.00)		Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	✓ Micreme
NYSTATIN	,	
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

# **Myometrial and Vaginal Hormone Preparations**

RGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj ava		5	✓ DBL Ergometrine
DESTRIOL			
Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
Pessaries 500 mcg	6.86	15	✓ Ovestin
OXYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	<ul> <li>Oxytocin BNM</li> </ul>
Inj 10 iu per ml, 1 ml ampoule		5	<ul> <li>Oxytocin BNM</li> </ul>
DXYTOCIN WITH ERGOMETRINE MALEATE - Up to	5 ini available on a PSO		-
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1	•	5	✓ Syntometrine

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

# **Pregnancy Tests - hCG Urine**

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Pregnancy Test

✓ Smith BioMed Rapid

**Pregnancy Test** 

### **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 108

### 5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

★ Tab 5 mg .......4.81 100 ✓ Ricit

#### ⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

# Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

### **⇒SA1032** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

### **Other Urinary Agents**

OXYBUTYNIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking oxybutynin prior to 1 June 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of oxybutynin.

(Apo-Oxybutynin Oral liq 5 mg per 5 ml to be delisted 1 May 2022)

POTASSIUM CITRATE

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

### ⇒SA1083 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE	2.22	20	<b>4</b> 11 1
* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	<ul> <li>Solifenacin Mylan</li> </ul>
Solifenacin Mylan to be Principal Supply on 1 Decemb	er 2021		
Tab 10 mg	3.72	30	✓ Solifenacin Mylan
Solifenacin Mylan to be Principal Supply on 1 Decemb	er 2021		•

### **Detection of Substances in Urine**

ORT	THO-	TOL	IDINI	=

* Compound diagnostic sticks	7.50 50	0 test OP
,	(8.25)	Hemastix
TETRABROMOPHENOL		

# **Obstetric Preparations**

# Antiprogesterones

		NE	MIFEPRISTONE
✓ Mifegyne	1	ng60.00	Tab 200 mg.
✓ Mifegyne	3	180.00	_

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

			_
Subsidy	Fully	/ Brand or	_
(Manufacturer's Price)	Subsidised	d Generic	
\$	Por 🗸	Manufacturer	

### **Calcium Homeostasis**

CALCITONIN

CINACALCET - Special Authority see SA1618 below - Retail pharmacy

Tab 30 mg − Wastage claimable ......210.30 28 ✓ Sensipar

### ⇒SA1618 Special Authority for Subsidy

**Initial application** only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
  - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
  - 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

**Renewal** only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

#### ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA2031 below −
Retail pharmacy.......18.00 1 ✓ Zoledronic acid
Mylan

Zoledronic acid Mylan to be Principal Supply on 1 December 2021

### ⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- Patient has hypercalcaemia of malignancy: or
- 2 Both:
  - 2.1 Patient has bone metastases or involvement; and
  - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
  - 3.1 Patient has bone metastases or involvement: and
  - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy	Fully	Brand or
(Manufacturer's	Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

# Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

W. In: 0.0 man with histograph acceptance of the control of the co		_	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml		5	
(3)	36.96)		Celestone
			Chronodose
DEXAMETHASONE			
* Tab 0.5 mg - Up to 60 tab available on a PSO	1 50	30	✓ Dexmethsone
* Tab 4 mg - Up to 30 tab available on a PSO			✓ Dexmethsone
			✓ Biomed
Oral liq 1 mg per ml	15.00 2	o IIII OP	bioinea
DEXAMETHASONE PHOSPHATE			
Dexamethasone phosphate injection will not be funded for oral use.			
* Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO	.9.25	10	✓ Dexamethasone
			Phosphate
			Panpharma
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO1	16 27	10	✓ Dexamethasone
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO1	10.37	10	
			<u>Phosphate</u>
			<u>Panpharma</u>
FLUDROCORTISONE ACETATE			
* Tab 100 mcg	14.32	100	✓ Florinef
HYDROCORTISONE			
	0.40	100	/ Davidas
* Tab 5 mg			✓ Douglas
* Tab 20 mg			✓ Douglas
* Inj 100 mg vial	.4.38	1 '	✓ Solu-Cortef
<ul> <li>a) Up to 5 inj available on a PSO</li> </ul>			
b) Only on a PSO			
c) Solu-Cortef to be Principal Supply on 1 November 2021			
METHYLPREDNISOLONE			
* Tab 4 mg11	12.00	100	✓ Medrol
* Tab 100 mg19	94.00	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial1	18.90	1 ,	✓ Solu-Medrol-Act-
, •			O-Vial
Inj 125 mg vial2	28.90	1 ,	✓ Solu-Medrol-Act-
. •			O-Vial
Inj 500 mg vial	22.78	1 ,	✓ Solu-Medrol-Act-
,			O-Vial
			• • • • • • • • • • • • • • • • • • • •
Inj 1 g vial2	27.83	1 ,	✓ Solu-Medrol
METHYLPREDNISOLONE ACETATE			
	14.40	-	/ Dama Madual
Inj 40 mg per ml, 1 ml vial4	14.40	5	✓ Depo-Medrol

	Subsidy		Fully	Brand or
(	Manufacturer's Pric	ce) Subs Per	idised •	Generic Manufacturer
	<u></u>	rei		Manuacturei
PREDNISOLONE				
* Oral liq 5 mg per ml - Up to 30 ml available on a PSO	6.00	30 ml OP	<b>✓</b> [	Redipred
<ul> <li>a) Restricted to children under 12 years of age.</li> </ul>				
b) Redipred to be Principal Supply on 1 December 2021				
PREDNISONE				
* Tab 1 mg	18.58	500	1	Apo-Prednisone
* Tab 2.5 mg		500	1	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	1	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	1	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen S29
, , , , , , , , , , , , , , , , , , , ,				AU Synacthen
			1	Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Depot
, .,			<b>√</b> 9	Synacthene
				Retard \$29
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	<b>✓</b>	Kenacort-A 10
, , , , , , , , , , , , , , , , , , , ,	26.62		1	Adcortyl \$29
Inj 40 mg per ml, 1 ml ampoule		1		Triaver S29
ing 40 mg por mi, 1 mi ampodio	51.10	5		Kenacort-A 40
	70.62	Ū	-	Kenalog S29
(Adcortyl \$29 Inj 10 mg per ml, 1 ml ampoule to be delisted 1 Oct	. 0.02		•	tolialog
, , , , , , , , , , , , , , , , , , , ,	,			
(Triaver S29 Inj 40 mg per ml, 1 ml ampoule to be delisted 1 Octo	,			
(Kenalog S29 Inj 40 mg per ml, 1 ml ampoule to be delisted 1 Oct	oper 2021)			

# **Sex Hormones Non Contraceptive**

# **Androgen Agonists and Antagonists**

CYPROTERONE ACETATE		
Tab 50 mg14.	37 50	✓ Rex S29
•		✓ Siterone
Tab 100 mg28.	03 50	✓ Siterone
TESTOSTERONE		
Patch 5 mg per day90.	00 30	✓ Androderm
TESTOSTERONE CIPIONATE		
Inj 100 mg per ml, 10 ml vial85.	00 1	✓ Depo-Testosterone
TESTOSTERONE ESTERS		·
Inj 250 mg per ml, 1 ml12.	98 1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE		- Cuctanon / impoured
	00 00	Andrial Tastasana
Cap 40 mg21.		Andriol Testocaps
Inj 250 mg per ml, 4 ml vial86.	00 1	Reandron 1000

# Hormone Replacement Therapy - Systemic

### **Prescribing Guideline**

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

		Subsidy (Manufacturer's Price	e) Sub	Fully sidised	Brand or Generic
		\$	Per	<b>√</b>	Manufacturer
Oe	estrogens				
DES	TRADIOL - See prescribing guideline on the previous page				
*	Tab 1 mg	4.12	28 OP		
		(11.10)		Е	strofem
*	Tab 2 mg		28 OP	_	
		(11.10)			strofem
*	Patch 100 mcg per 24 hours	7.91	4	✓ (	Climara
	No more than 1 patch per week				
	b) Only on a prescription				
*	Patch 50 mcg per 24 hours	7.04	4	✓ (	Climara
	a) No more than 1 patch per week				
	b) Only on a prescription				
	Patch 25 mcg per day		8		stradot
		7.85		<b>✓</b> E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 50 mcg per day	7.04	8	<b>√</b> E	stradot 50 mcg
		9.22		<b>✓</b> E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 75 mcg per day	7.91	8	<b>√</b> E	stradot
		10.60		<b>✓</b> E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				•
	b) Only on a prescription				
	Patch 100 mcg per day	7.91	8	<b>√</b> E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
Clin	nara Patch 100 mcg per 24 hours to be delisted 1 January 20	122)			
	nara Patch 50 mcg per 24 hours to be delisted 1 January 202				
	TRADIOL VALERATE - See prescribing guideline on the pro-				
	Tab 1 mg		84	<b>√</b> □	rogynova
	Tab 1 mg		84		rogynova Progynova
	· ·		04	• -	Togyttova
	TROGENS – See prescribing guideline on the previous page		00		
*	Conjugated, equine tab 300 mcg		28	_	
	Operior and a service state OOF areas	(17.50)	00	۲	remarin
*	Conjugated, equine tab 625 mcg		28	_	
		(17.50)		۲	Premarin
Pr	ogestogens				
ΛΕΓ	PROXYPROGESTERONE ACETATE - See prescribing guid	eline on the previo	ous page		
	Tab 2.5 mg		30	<b>√</b> P	rovera
	Tab 5 mg		100	-	rovera
	Tab 10 mg		30		rovera
••	144 14 mg		00	• 1	

		Subsidy (Manufacturer's Price \$	) Su Per	Fully bsidised	Brand or Generic Manufacturer			
P	Progestogen and Oestrogen Combined Preparations							
OE	STRADIOL WITH NORETHISTERONE - See prescribing gui	ideline on page 81						
*	Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP					
		(18.10)		Kl	iovance			
*	Tab 2 mg with 1 mg norethisterone acetate		28 OP					
		(18.10)		KI	iogest			
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg							
	oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	т.,				
		(18.10)		ır	isequens			
0	ther Oestrogen Preparations							
Ŭ	ther ocotrogen reparations							
ETI	HINYLOESTRADIOL							
*	Tab 10 mcg	17.60	100		Z Medical and			
					Scientific			
	STRIOL			_				
*	Tab 2 mg	7.00	30	<b>✓</b> <u>0</u>	vestin			
	they Dyegestegen Dyeneyetiens							
U	ther Progestogen Preparations							
LE	VONORGESTREL							
*	Intra-uterine device 52 mg	269.50	1	✓ Mi	<u>irena</u>			
*	Intra-uterine device 13.5 mg	215.60	1	✓ <u>Ja</u>	<u>ıydess</u>			
ME	DROXYPROGESTERONE ACETATE							
	Tab 100 mg	116.15	100	✓ Pr	overa HD			
NO	RETHISTERONE							
*	Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓ Pr	imolut N			
	OGESTERONE							
	Cap 100 mg - Special Authority see SA1609 below - Retail							
	pharmacy	16.50	30	✓ Ut	rogestan			

#### **⇒SA1609** Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 For the prevention of pre-term labour\*; and
- 2 Either:
  - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

**Renewal** only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour\*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
  - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are unapproved indications.

Cubaidy

Eully.

Drand or

	Subsidy (Manufacturer's Price) \$	Per	Subsidised	
Thyroid and Antithyroid Agents	·			
CARBIMAZOLE				
* Tab 5 mg	10.80	100	_	Neo-Mercazole Neo-Mercazole S29 S29
LEVOTHYROXINE				
* Tab 25 mcg	5.55	90	✓	Synthroid
* Tab 50 mcg	1.71	28	✓	Mercury Pharma
•	5.79	90	✓	Synthroid
	64.28	1,000	<b>/</b>	Eltroxin
* Tab 100 mcg	1.78	28	✓	Mercury Pharma
•	6.01	90	✓	Synthroid
	66.78	1,000	<b>/</b>	Eltroxin
PROPYLTHIOURACIL – Special Authority see SA1199 below Propylthiouracil is not recommended for patients under the treatments are contraindicated.		the	patient is p	regnant and other
Tab 50 mg	35.00	100	/	PTU S29

### ⇒SA1199 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

### **Trophic Hormones**

### **Growth Hormones**

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 bel	low – Retail pha	armacy	
*	Inj 5 mg cartridge	69.75	1	<ul><li>Omnitrope</li></ul>
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope

#### ⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
  - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
  - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
  - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In

Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer	
	. 0.		mananananan	

continued...

children who are 5 years or older, GH testing with sex steroid priming is required; and

- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate: and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and

(Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer		Subsidy Fully		Fully	Brand or
\$ Per ✓ Manufacturer	(M	anufacturer's Price)			
		\$	Per		Manufacturer

continued...

- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

**Initial application** — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and</p>
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Fither
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m<sup>2</sup> in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months...

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

**Initial application — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
  - 5.1 Both:

continued...

- 5.1.1 The patient is aged two years or older; and
- 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
- 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

**Initial application** — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and
  - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
  - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and

Subsidy	Subsidy Fully		Brand or	
(Manufacturer's		ubsidised	Generic	
<u> </u>	Per		Manufacturer	_
				_

continued...

- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

Implant 3.6 mg, syringe .......65.68

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues	
----------------	--

**GOSFREI IN** 

	Implant 10.8 mg, syringe	122.37	1	✓ <u>Teva</u>
LEU	JPRORELIN			
	Additional subsidy by endorsement where the patient is a child or goserelin and the prescription is endorsed accordingly.	adolescent and is	s unable to	tolerate administration of
	Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of			
	\$221.60 per 1 inj with Endorsement	66.48	1	
		(221.60)		Lucrin Depot 1-month
	Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
	of \$591.68 per 1 inj with Endorsement	177.50	1	
		(591.68)		Lucrin Depot 3-month

Wago	nrace	in Ac	ionie	
v asu	pressi	III AV	UIIIO	
		-		

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

1

Teva

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic
	\$	Per	1	Manufacturer
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	Minirin
Tab 200 mcg	54.45	30	✓ N	Minirin
▲ Nasal drops 100 mcg per ml	39.03	2.5 ml OP	✓ N	Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	✓ [	<u>PH&amp;T</u>
Inj 4 mcg per ml, 1 ml(Minirin Nasal drops 100 mcg per ml to be delisted 1 January 2022		10	<b>✓</b> N	Minirin

### **Other Endocrine Agents**

#### **CABERGOLINE**

#### ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

00 04

#### **CLOMIFENE CITRATE**

rab 50 mg	29.84	10	Clomiphen S29
METYRAPONE Cap 250 mg	558 00	50	✓ Metopirone
Cab 250 IIIu	556.00	50	▼ INICLODITOTIC

INFECTIONS - AGENTS FOR SYSTEMIC US	E			
	Subsidy (Manufacturer's Price) \$	Sı Per	Fully bsidised	Brand or Generic Manufacturer
Anthelmintics				
ALBENDAZOLE - Special Authority see SA1318 below - Retail	pharmacy			
Tab 400 mg	469.20	60	<b>√</b> E	skazole S29
▶ SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or clipatient has hydatids.	inical microbiologist.	Approv	als valid f	or 6 months where the
Renewal only from an infectious disease specialist or clinical mic remains appropriate and the patient is benefitting from the treatment of the control of t		ıls valid f	or 6 mont	hs where the treatment
MEBENDAZOLE – Only on a prescription Tab 100 mg Oral liq 100 mg per 5 ml		6 15 ml	-	ermox
PRAZIQUANTEL Tab 600 mg	68.00	8	<b>✓</b> B	liltricide
Antibacterials				
a) For topical antibacterials, refer to DERMATOLOGICALS, pag     b) For anti-infective eye preparations, refer to SENSORY ORGA				
Cephalosporins and Cephamycins				
CEFACLOR MONOHYDRATE Cap 250 mg	24.70	100	_	lanbaxy-Cefaclor lanbaxy-Cefaclor S29 S29
Grans for oral liq 125 mg per 5 ml  – Wastage claimable	3.53	100 ml		tanbaxy-Cefaclor tanbaxy-Cefaclor S29 S29
CEFALEXIN				
Cap 250 mg		20		ephalexin ABM
Cap 500 mgGrans for oral liq 25 mg per ml – Wastage claimable		20 100 ml		ephalexin ABM efalexin Sandoz
Grans for oral liq 50 mg per ml – Wastage claimable		100 ml		efalexin Sandoz
CEFAZOLIN – Subsidy by endorsement  Only if prescribed for dialysis or cellulitis in accordance with a accordingly.	a DHB approved prot	ocol and	I the pres	cription is endorsed
Inj 500 mg vial	3.39	5	✓ A	<u>FT</u>

### CEFTRIAXONE - Subsidy by endorsement

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

11.j 1 g 1101	
CEFUROXIME AXETIL	<ul> <li>Subsidy by endorsement</li> </ul>

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.

✓ Zinnat 50

✓ Ceftriaxone-AFT

✓ Ceftriaxone-AFT

Ini 1 a vial

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

### **Macrolides**

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8 19	30	✓ Apo-Azithromycin
Tab 500 mg – Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
3 op 11 mm m m m m	2.57		✓ Zithromax
Zithromax to be Sole Supply on 1 December 2021			
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	14.38	15 ml	✓ Zithromax

(Apo-Azithromycin Tab 250 mg to be delisted 1 May 2022) (Apo-Azithromycin Tab 500 mg to be delisted 1 December 2021)

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under: and
- 3 Either:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

	Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
CLARITHROMYCIN				

Tab 250 mg - Maximum of 56 tab per prescription; can be 14 ✓ Apo-Clarithromycin ✓ Klacid 8.53 50 ml ✓ Klacid

- a) Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below
- b) Wastage claimable

(Apo-Clarithromycin Tab 250 mg to be delisted 1 February 2022)

### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Atypical mycobacterial infection: or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omegrazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

#### **ERYTHROMYCIN (AS LACTOBIONATE)**

Ervthrocin IV

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
DVTUDOMVOM ETUVI QUOCIMATE	Ψ			Wandactarci
RYTHROMYCIN ETHYL SUCCINATE	16.05	100	./	E Myoin
Tab 400 mg	10.93	100	•	E-Mycin
a) Up to 20 tab available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 200 mg per 5 ml	5.00	100 m		E-Mycin
		100 111	. •	E-IMYCIII
a) Up to 300 ml available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Grans for oral lig 400 mg per 5 ml	6.77	100 m	. <i>.</i>	E-Mycin
a) Up to 200 ml available on a PSO		100 111		L myom
b) Wastage claimable				
,				
RYTHROMYCIN STEARATE	14.05	100		
Tab 250 mg - Up to 30 tab available on a PSO		100		ERA
Tab 500 mg	(22.29)	100		ENA
Tab 500 Hig	(44.58)	100		ERA
ERA Tab 250 mg to be delisted 1 April 2022)	(44.30)			LIIA
ERA Tab 250 mg to be delisted 1 April 2022) ERA Tab 500 mg to be delisted 1 September 2022)				
, ,				
ROXITHROMYCIN	0.00	40	,	Desired D
Tab disp 50 mg	8.29	10	•	Rulide D
Restricted to children under 12 years of age.	0.00	50	./	Arrow-
Tab 150 mg	0.20	50	•	Roxithromycin
				<u>noxidirolliyelli</u>
Tab 300 mg	16.33	50	1	Arrow-
•				Roxithromycin
Penicillins				
AMOXICILLIN	22.50	500	J	Alphamov
AMOXICILLIN Cap 250 mg	22.50	500	•	Alphamox
MOXICILLIN Cap 250 mga) Up to 30 cap available on a PSO	22.50	500	•	Alphamox
MMOXICILLIN  Cap 250 mg  a) Up to 30 cap available on a PSO  b) Up to 10 x the maximum PSO quantity for RFPP				
AMOXICILLIN  Cap 250 mg  a) Up to 30 cap available on a PSO  b) Up to 10 x the maximum PSO quantity for RFPP  Cap 500 mg		500		Alphamox Alphamox
AMOXICILLIN  Cap 250 mg  a) Up to 30 cap available on a PSO  b) Up to 10 x the maximum PSO quantity for RFPP  Cap 500 mg				
AMOXICILLIN  Cap 250 mg	36.98	500	•	Alphamox
AMOXICILLIN  Cap 250 mg	36.98		•	
AMOXICILLIN  Cap 250 mg	36.98	500	•	Alphamox
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg	36.98	500	<b>/</b>	Alphamox Alphamox 125
AMOXICILLIN  Cap 250 mg	36.98	500 100 ml	<b>/</b>	Alphamox
AMOXICILLIN  Cap 250 mg	36.98	500 100 ml	<b>/</b>	Alphamox Alphamox 125
AMOXICILLIN  Cap 250 mg	36.98	500 100 ml	<b>/</b>	Alphamox Alphamox 125
AMOXICILLIN  Cap 250 mg	36.98	500 100 ml	/   /	Alphamox Alphamox 125
AMOXICILLIN  Cap 250 mg	36.981.401.73	500 100 ml		Alphamox 125 Alphamox 250

	Subsidy		Fully	Brand or
	(Manufacturer's Pr		idised	Generic
	\$	Per	1	Manufacturer
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab available on a PSO	0.89	10	/	Curam Duo 500/125
Grans for oral lig amoxicillin 25 mg with clavulanic acid 6.25 m	ng			
per ml	5.00	100 ml	1	Augmentin
a) Up to 200 ml available on a PSO     b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 m per ml – Up to 200 ml available on a PSO	•	100 ml OP	/	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	344.93	10	/	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial - Up to 5 inj available on a PS	O 11.09	10	1	Sandoz
LUCLOXACILLIN	40.00	050	,	<b>0</b>
Cap 250 mg — Up to 30 cap available on a PSO		250		Staphlex
Cap 500 mg – Up to 30 cap available on a PSO		500 100 ml		Staphlex AFT
Grans for oral liq 25 mg per ml	3.29	100 1111	•	AFI
Grans for oral liq 50 mg per ml	3.68	100 ml	1	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable	17.50	40		Fluelessie
Inj 250 mg vial		10 10		Flucioxin Flucioxin
Inj 500 mg vialInj 1 g vial – Up to 5 inj available on a PSO		5		Flucil
		3	٠	<u>i iucii</u>
HENOXYMETHYLPENICILLIN (PENICILLIN V)	0.04	50		Ollianian VV
Cap 250 mg – Up to 30 cap available on a PSO		50 50		Cilicaine VK Cilicaine VK
Cap 500 mg	0.00	50	•	Cilicaine VK
a) Up to 20 cap available on a PSO     b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2 99	100 ml	1	AFT
a) Up to 200 ml available on a PSO     b) Wastage claimable	2.00	100 1111	·	<u>Al I</u>
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	1	AFT
a) Up to 300 ml available on a PSO				<u> </u>
b) Up to 2 x the maximum PSO quantity for RFPP c) Wastage claimable				
ROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	1	Cilicaine
Tetracyclines				
DOXYCYCLINE	04.40	F00	,	Davina
Fab 100 mg - Up to 30 tab available on a PSO	64.43	500	•	Doxine

	INFECTIONS - AC	žΕΝ	ITS FOR SYSTEMIC USE
	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic  Manufacturer
MINOCYCLINE HYDROCHLORIDE	·		
* Tab 50 mg - Additional subsidy by Special Authority see			
SA1355 below – Retail pharmacy	5.79	60	
dt 0 400	(12.05)	400	Mino-tabs
* Cap 100 mg	19.32 (52.04)	100	Minomycin
<b>⇒SA1355</b> Special Authority for Manufacturers Price	(32.04)		MillottiyCill
Initial application from any relevant practitioner. Approvals val	id without further renev	wal u	nless notified where the patient has
TETRACYCLINE - Special Authority see SA1332 below - Reta	il pharmacy		
Tab 250 mg	21.42	28	✓ Accord S29
<b>⇒SA1332</b> Special Authority for Subsidy			
<b>Initial application</b> from any relevant practitioner. Approvals val Both:	id for 3 months for app	licati	ions meeting the following criteria:
<ul><li>1 For the eradication of helicobacter pylori following unsuc</li><li>2 For use only in combination with bismuth as part of a quantum</li></ul>			opriate first-line therapy; and
Other Antibiotics			
For topical antibiotics, refer to DERMATOLOGICALS, page 62			
CIPROFLOXACIN			
Recommended for patients with any of the following:			
i) microbiologically confirmed and clinically significant ps	seudomonas infection;	or	
ii) prostatitis; or	,		
iii) pyelonephritis; or			
iv) gonorrhoea.			
Tab 250 mg - Up to 5 tab available on a PSO	2 42	28	✓ Cipflox
Tab 500 mg – Up to 5 tab available on a PSO		28	✓ Cipflox
Tab 750 mg		28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	✓ <u>Dalacin C</u>
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓ Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -			
Only if prescribed for dialysis or cystic fibrosis patient and th			
Inj 150 mg	65.00	1	✓ Colistin-Link
GENTAMICIN SULPHATE	05.00	_	/ DDI Contonicio
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement. Only if prescribed for a dialysis or cystic fibrosis patient		5	✓ DBL Gentamicin
endorsed accordingly.	or complicated unitary	liaci	timection and the prescription is
Inj 10 mg per ml, 2 ml ampoule – Subsidy by endorsement.	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient	or complicated urinary		
endorsed accordingly.			
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement.		10	✓ Pfizer
Only if preservined for a distrain or quetic fibracia nations	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	, ,		i intection and the prescription is
MOXIFLOXACIN - Special Authority see SA1740 on the next p	age – Retail pharmacy	,	
No patient co-payment payable	40.00	_	Augler:
Tab 400 mg	42.00	5	✓ <u>Avelox</u>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

### ⇒SA1740 Special Authority for Subsidy

**Initial application** — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*; and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications: or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
    - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
    - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
    - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Either:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with \* are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

#### ⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Fither:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

**Renewal** only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 on the next page - Retail pharmacy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ ⇒SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg .......34.50 12 ✓ Fucidin SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy Tab 500 mg .......543.20 56 Wockhardt \$29 ⇒SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. **TOBRAMYCIN** ✓ Tobramycin Mylan Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly. Solution for inhalation 60 mg per ml. 5 ml - Subsidy by ✓ Tobramycin BNM endorsement......395.00 56 dose a) Wastage claimable b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly. **TRIMFTHOPRIM** 50 ✓ TMP TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE] Tab trimethoprim 80 mg and sulphamethoxazole 400 mg - Up to 30 tab available on a PSO......64.80 500 ✓ Trisul Oral lig 8 mg sulphamethoxazole 40 mg per ml - Up to 200 ml 100 ml Deprim VANCOMYCIN - Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly. ✓ Mylan **Antifungals** a) For topical antifungals refer to DERMATOLOGICALS, page 63 b) For topical antifungals refer to GENITO URINARY, page 76 FLUCONAZOLE 28 1 Mylan 28 Mylan Powder for oral suspension 10 mg per ml - Special Authority 35 ml ✓ Diflucan Wastage claimable

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	1	Manufacturer

### ⇒SA1359 Special Authority for Subsidy

**Initial application — (Systemic candidiasis)** from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

**Renewal** — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Т	R	Δ	C.	n	N	Δ	7	n	П	F

Cap 100 mg	4.27	15	✓ Itrazole
Oral lig 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy	141.80	150 ml OP	✓ Sporanox

#### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

CDC

### KETOCONAZOLE

Tab 200 mg DCT

Tab 200 mg - PC1		30	✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 on t	he next page – Retail ph	armacy	
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral lig 40 mg per ml		105 ml OP	✓ Noxafil

/ Link Hoalthoare con

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

### **⇒SA1285** Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

**Renewal** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

#### **TERBINAFINE**

<b>*</b> Tab 250 mg8.15 84 <b>✓ Dec</b>	olate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy	
Tab 50 mg91.00 56 ✓ Vtta	ack
Tab 200 mg350.00 56 ✓ Vtta	ack
Powder for oral suspension 40 mg per ml - Wastage	
claimable1,437.00 70 ml <b>✓ Vfe</b>	nd

### ⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis: or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### **Antimalarials**

PRIMAQUINE - Special Authority see SA1684 below - Retail pharmacy		
Tab 15 mg400.00	100	✓ Sanofi
		Primaguine \$29

### ⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

#### Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

### Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

# **Antitrichomonal Agents**

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	Arrow-Ornidazole
Arrow-Ornidazole to be Principal Supply on 1 December 2	021		

# **Antituberculotics and Antileprotics**

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

### CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

*	Cap 50 mg	442.00	100	✓ Lamprene S29
---	-----------	--------	-----	----------------

#### CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician. 29

Cap 250 mg	344 00	60	✓ Cyclorin S29
Cab 230 IIIu		00	▼ CVCIOIIII ©2

#### DAPSONE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Tab 25 mg	268.50	100	Dapsone
Tab 100 mg	329.50	100	Dapsone

		a Litt		O I O I E IIII O O O E
	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Special	ist			
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommenda respiratory physician</li> </ul>		lisease	physicia	n, clinical microbiologist or
Tab 100 mg	85.73	100	✓	EMB Fatol S29
Tab 400 mg	49.34	56	•	Myambutol S29
ISONIAZID - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician</li> </ul>	tion of, an internal me	dicine p		
* Tab 100 mg	23.00	100	/	PSM
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist				
<ul> <li>a) No patient co-payment payable</li> <li>b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician</li> </ul>	tion of, an internal me	dicine p	ohysician	, paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg		100		Rifinah
* Tab 150 mg with rifampicin 300 mg	179.13	100	/	Rifinah
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommenda respiratory physician  Crops for each line Are peoplet.		lisease 30	•	et, clinical microbiologist or
Grans for oral liq 4 g sachet	200.00	30	•	PdSel 329
PROTIONAMIDE – Retail pharmacy-Specialist  a) No patient co-payment payable  b) Prescriptions must be written by, or on the recommenda respiratory physician  Tab 250 mg		lisease 100	•	t, clinical microbiologist or
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable     b) Prescriptions must be written by, or on the recommenda respiratory physician	tion of, an infectious d	lisease	physicia	n, clinical microbiologist or
* Tab 500 mg	59.00	100	/	AFT-Pyrazinamide
RIFABUTIN – Retail pharmacy-Specialist  a) No patient co-payment payable  b) Prescriptions must be written by, or on the recommenda gastroenterologist	tion of, an infectious d	lisease	physicia	n, respiratory physician or
* Cap 150 mg	299.75	30	1	Mycobutin
RIFAMPICIN – Subsidy by endorsement				
<ul> <li>a) No patient co-payment payable</li> <li>b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescripti Retail pharmacy - Specialist. Specialist must be an inte paediatrician, or public health physician.</li> </ul>	on is endorsed accord	lingly; c	an be wa	aived by endorsement -
* Cap 150 mg	58.54	100	✓	Rifadin
* Cap 300 mg		100		Rifadin
* Oral liq 100 mg per 5 ml	12.60	60 ml	1	Rifadin

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

### **Antivirals**

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 234

### **Hepatitis B Treatment**

ENTECAVIR			
* Tab 0.5 mg	52.00	30	<ul> <li>Entecavir Sandoz</li> </ul>
LAMIVUDINE - Special Authority see SA1685 below - Retail p	harmacy		
Tab 100 mg	6.95	28	✓ Zetlam
Oral lig 5 mg per ml	270.00	240 ml OP	✓ Zeffix

### ⇒SA1685 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 105

# **Herpesvirus Treatments**

ACICLOVIR			
* Tab dispersible 200 mg	1.60	25	✓ <u>Lovir</u>
* Tab dispersible 400 mg	5.38	56	✓ <u>Lovir</u>
* Tab dispersible 800 mg	5.98	35	✓ <u>Lovir</u>
VALACICLOVIR			
Tab 500 mg	6.50	30	✓ Vaclovir
Tab 1,000 mg		30	✓ Vaclovir
VALGANCICLOVIR - Special Authority see SA1993 b	elow – Retail pharmacy		
Tab 450 mg		60	✓ Valganciclovir
·			Mylan

Valganciclovir Mylan to be Principal Supply on 1 December 2021

#### ⇒SA1993 Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

#### 1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or

#### 2 Both:

2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and

continued...

2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone. Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

# **Hepatitis C Treatment**

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <a href="https://pharmac.govt.nz/maviret">https://pharmac.govt.nz/maviret</a>

Tab 100 mg with pibrentasvir 40 mg ......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Authority see SA1605 on the next page
No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni** 

Subsidy (Manufacturer's Price)

Fully Subsidised Per Brand or Generic Manufacturer

### **⇒SA1605** Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

### **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 105 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

.

✓ Teva

### ⇒SA1994 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks;
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

### **Antiretrovirals**

### **⇒SA1651** Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

**Initial application — (Percutaneous exposure)** only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

### Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous p	oage – Retail pharr	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	page – Retail phar	macy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	page - Retail pha	rmacy	
Tab 200 mg	84.00	60	<ul><li>Nevirapine Alphapharm</li></ul>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune  Suspension

	Cubaidu		T. Ilsa	Drandar
	Subsidy (Manufacturer's I		Fully dised	Brand or Generic
	\$	Per	1	Manufacturer
Nucleosides Reverse Transcriptase Inhibitors				
·				
ABACAVIR SULPHATE – Special Authority see SA1651 on page				
Tab 300 mg Oral lig 20 mg per ml		60 240 ml OP		<u>agen</u> agen
				•
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts				
anti-retroviral Special Authority.	as two anti-retion	oviiai illeulcalion	5 101 11	ie purposes or trie
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ K	ivexa
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPR		I Authority see S	— A1651	on page 105 – Retail
pharmacy				on page 100 Tiolan
Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority	unts as three a	nti-retroviral med	lication	s for the purposes of the
Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	il			
245 mg (300 mg as a maleate)		30	<b>✓</b> M	vlan
EMTRICITABINE – Special Authority see SA1651 on page 105 –		ev.		
Cap 200 mg		30	✓ E	mtriva
LAMIVUDINE - Special Authority see SA1651 on page 105 - Re	tail pharmacy			
Tab 150 mg		60	<b>✓</b> La	amivudine
·				Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	<b>√</b> 3	TC
ZIDOVUDINE [AZT] - Special Authority see SA1651 on page 10	5 – Retail pharr	nacy		
Cap 100 mg		100	✓ R	etrovir
Oral liq 10 mg per ml	30.45	200 ml OP	✓ R	etrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE - Special Authority see				
Note: zidovudine [AZT] with lamivudine (combination tablets)	counts as two	anti-retroviral me	edication	ons for the purposes of
the anti-retroviral Special Authority.  Tab 300 mg with lamivudine 150 mg	33.00	60	./ A	lphapharm
Tab 500 mg with amivudine 150 mg			• ^	ірпарпапп
Protease Inhibitors				
ATAZANAVIR SULPHATE - Special Authority see SA1651 on pa	age 105 – Retai	l pharmacy		
Cap 150 mg		60	✓ Tell	<u>eva</u>
Cap 200 mg	188.91	60	✓ <u>Te</u>	<u>eva</u>
DARUNAVIR - Special Authority see SA1651 on page 105 - Ref			_	
Tab 400 mg		60		arunavir Mylan
Tab 600 mg		60	<b>✓</b> <u>D</u>	arunavir Mylan
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651				
Tab 100 mg with ritonavir 25 mg	150.00	60		opinavir/Ritonavir
	183.75			Mylan aletra
Tab 200 mg with ritonavir 50 mg		120		opinavir/Ritonavir
		0		Mylan
	463.00			aletra
Oral liq 80 mg with ritonavir 20 mg per ml		300 ml OP	✓ K	aletra
(Kaletra Tab 100 mg with ritonavir 25 mg to be delisted 1 Februar	,			
(Kaletra Tab 200 mg with ritonavir 50 mg to be delisted 1 Februar	y 2022)			
RITONAVIR - Special Authority see SA1651 on page 105 - Reta				
Tab 100 mg	43.31	30	✓ N	<u>orvir</u>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

#### Strand Transfer Inhibitors

DOLUTEGRAVIR - Special Authority see SA1651 on page	e 105 - Retail pharmacy		
Tab 50 mg	1,090.00	30	Tivicay
RALTEGRAVIR POTASSIUM - Special Authority see SA1	651 on page 105 - Reta	il pharmacy	
Tab 400 mg	1,090.00	60	✓ Isentress
Tab 600 mg	1.090.00	60	✓ Isentress HD

# Immune Modulators

#### Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

#### Criteria for Treatment

- 1) Diagnosis
  - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
  - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
  - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

### **Exclusion Criteria**

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10<sup>9</sup>) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

#### Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

#### **Exit Criteria**

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

### ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and

## **INFECTIONS - AGENTS FOR SYSTEMIC USE**

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

2 Maximum of 48 weeks therapy.

### Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400.000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

## All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; or
  - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and

## INFECTIONS - AGENTS FOR SYSTEMIC USE

	ubsidy cturer's Price) Subs	Fully	Brand or Generic
·	\$ Per	•	Manufacturer

continued...

- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

**Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
  - 3.1 Patient has a cutaneous T cell lymphoma\*; or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Either:
      - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
      - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with \* are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet quidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

**Initial application** — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

# **Urinary Tract Infections**

	THENAMINE (HEXAMINE) HIPPURATE Tab 1 g	40.01	100	✓ Hiprex	
NITI	ROFURANTOIN				
*	Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran	
*	Tab 100 mg	37.50	100	✓ Nifuran	
*	Cap modified-release 100 mg - Wastage claimable	86.40	100	✓ Macrobid	

## **INFECTIONS - AGENTS FOR SYSTEMIC USE**

Subsidy (Manufacturer's Price)	Ful Subsidise	,
 \$	Per •	Manufacturer

## NORFLOXACIN

	Subsidy (Manufacturerla Di	rian) Cuda	Fully	Brand or
	(Manufacturer's Pr \$	Per	sidised •	Generic Manufacturer
nticholinesterases				
EOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule		10		Ino S29 ax Health
	29.40 98.00	50		ax neam straZeneca
VIDIDOCTIONAINE DECOMIDE	90.00	30	V A3	sti azerieca
YRIDOSTIGMINE BROMIDE Tab 60 mg	45.70	100	✓ M	estinon
rab oo mg		100	• <u>IVI</u>	<u>country</u>
Non-Steroidal Anti-Inflammatory Drugs				
CLOFENAC SODIUM				
Tab EC 25 mg	1.99	50	✓ Di	clofenac Sandoz
Tab 50 mg dispersible		20		oltaren D
Tab EC 50 mg		50	_	clofenac Sandoz
Tab long-acting 75 mg		100		oltaren SR
	22.80	500		oo-Diclo SR
Tab long-acting 100 mg		500		oo-Diclo SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj availal		5		oltaren
Suppos 12.5 mg		10		oltaren
Suppos 25 mg		10		oltaren
Suppos 50 mg — Up to 10 supp available on a PSC Suppos 100 mg		10 10		oltaren oltaren
po-Diclo SR Tab long-acting 75 mg to be delisted 1 N po-Diclo SR Tab long-acting 100 mg to be delisted 1 UPROFEN				
Tab 200 mg	21.40	1,000	✓ Re	elieve
Tab long-acting 800 mg	3.05	30	✓ Br	ufen SR
	5.99		✓ Ib	uprofen SR BNM
	5.99			hics
1 - 31	1.88	200 ml	✓ Et	11100
ouprofen SR BNM Tab long-acting 800 mg to be delist	1.88	200 ml	✓ Et	
ouprofen SR BNM Tab long-acting 800 mg to be delist ETOPROFEN	1.88 ted 1 January 2022)			
ouprofen SR BNM Tab long-acting 800 mg to be delist ETOPROFEN Cap long-acting 200 mg	1.88 ted 1 January 2022)	200 ml 28		ruvail SR
ouprofen SR BNM Tab long-acting 800 mg to be delist  TOPROFEN  Cap long-acting 200 mg  EFENAMIC ACID	1.88 ted 1 January 2022)	28		
uprofen SR BNM Tab long-acting 800 mg to be delist TOPROFEN Cap long-acting 200 mg EFENAMIC ACID			<b>√</b> Oı	ruvail SR
uprofen SR BNM Tab long-acting 800 mg to be delist TOPROFEN Cap long-acting 200 mg EFENAMIC ACID		28 50	<b>√</b> Oı	
uprofen SR BNM Tab long-acting 800 mg to be delist TOPROFEN Cap long-acting 200 mg EFENAMIC ACID		28	<b>✓ Oi</b>	ruvail SR onstan
ouprofen SR BNM Tab long-acting 800 mg to be delist ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg		28 50	<b>✓ Oi</b>	ruvail SR
puprofen SR BNM Tab long-acting 800 mg to be delist ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg	1.88  ted 1 January 2022)	28 50 20	<b>✓ Oi</b> Po	onstan
puprofen SR BNM Tab long-acting 800 mg to be delisted  ETOPROFEN Cap long-acting 200 mg  EFENAMIC ACID Cap 250 mg		28 50 20	✓ Oil Po	ruvail SR onstan onstan oflam 250
APROXEN Tab 500 mg Tab 750 mg Tab 500 mg Tab 500 mg Tab 500 mg		28 50 20 500 250	✓ OI  PC  PC  ✓ NG	ruvail SR onstan onstan oflam 250 oflam 500
APROXEN Tab 500 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg	1.88  ted 1 January 2022)  12.07  12.07  1.25 (9.16) 0.50 (5.60)  32.69 28.71 6.47	28 50 20 500 250 28	Po Po	ruvail SR onstan onstan offlam 250 offlam 500 aprosyn SR 750
APROXEN Tab 250 mg Tab 500 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 7 g	1.88  ted 1 January 2022)  12.07  12.07  1.25 (9.16) 0.50 (5.60)  32.69 28.71 6.47	28 50 20 500 250	Po Po	ruvail SR onstan onstan oflam 250 oflam 500
APROXEN Tab 250 mg Tab 500 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 750 mg Tab 10ng-acting 1 g	1.88  ted 1 January 2022)  12.07  12.07  1.25 (9.16) 0.50 (5.60)  32.69 28.71 6.47 8.62	28 50 20 500 250 28 28	PC PC V No V No V No	ruvail SR onstan onstan oflam 250 oflam 500 aprosyn SR 750 aprosyn SR 1000
APROXEN Tab 250 mg Tab 500 mg Tab 100 mg	1.88  ted 1 January 2022)  12.07  12.07  1.25 (9.16) 0.50 (5.60)  28.71	28 50 20 500 250 28 28 28	Po Po No	ruvail SR onstan onstan oflam 250 oflam 500 aprosyn SR 750 aprosyn SR 1000
buprofen SR BNM Tab long-acting 800 mg to be delist ETOPROFEN Cap long-acting 200 mg EFENAMIC ACID Cap 250 mg APROXEN Tab 250 mg Tab 500 mg Tab long-acting 750 mg ULINDAC	1.88  ted 1 January 2022)  12.07  12.07  1.25 (9.16) 0.50 (5.60)  28.71	28 50 20 500 250 28 28	Po Po Po No	ruvail SR onstan onstan oflam 250 oflam 500 aprosyn SR 750 aprosyn SR 1000

MUSCULUSKELETAL SYSTEM							
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer			
TENOXICAM * Tab 20 mg * Inj 20 mg vial		100 1		<u>Tilcotil</u> AFT			
NSAIDs Other							
CELECOXIB Cap 100 mg Cap 200 mg		60 30	✓	Celecoxib Pfizer Celebrex Celecoxib Pfizer			
Topical Products for Joint and Muscular Pain							
CAPSAICIN  Crm 0.025% - Special Authority see SA1289 below - Ret pharmacy	9.75 4 alid without further rene		nless notifi				
Antirheumatoid Agents							
Subsidised only if prescribed for rheumatoid arthritis, systessuppression, relevant dermatological conditions (cutaneoumucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary and non-pulmonary and system and sendorsed with the prescription as endorsed with a * is a * Tab 200 mg	is forms of lupus and lic monary)*, and the presc here there exists a reco an unapproved indication	hen p riptio ord of	planus, cuta n is endors prior dispe	aneous vasculitides and ed accordingly.			
LEFLUNOMIDE				<b></b>			
Tab 10 mg		30		<u>Arava</u>			
Tab 20 mg	6.00	30	/	<u>Arava</u>			
PENICILLAMINE  Tab 125 mg	67 23	100	1	D-Penamine			
Tab 250 mg		100		D-Penamine			
Drugs Affecting Bone Metabolism							
Alendronate for Osteoporosis							
ALENDRONATE SODIUM							
* Tab 70 mg	2.44	4	✓	Fosamax			
ALENDRONATE SODIUM WITH COLECALCIFEROL  * Tab 70 mg with colecalciferol 5,600 iu	1.51	4	/	Fosamax Plus			
Other Treatments							
DENOSUMAB – Special Authority see SA1777 on the next pa	na – Retail nharmany						
Inj 60 mg prefilled syringe	•	1	1	Prolia			

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

## ⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

## All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
  - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

## PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	79.95	1	Pamisol

RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779 on the next page - Retail pharmacy

Si	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsi	dised	Generic
	\$ Per	1	Manufacturer

## ⇒SA1779 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

### RISEDRONATE SODIUM

Tab 35 mg3.10	0 4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy		
Inj 250 mcg per ml, 2.4 ml490.00	0 1	✓ Forteo

## ⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops

(Mar	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
	\$	Per	<b>✓</b>	Manufacturer

continued...

during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

### ZOLEDRONIC ACID

## ⇒SA1780 Special Authority for Subsidy

**Initial application** — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

**Initial application — (Underlying cause - Osteoporosis)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

**Initial application — (Underlying cause - glucocorticosteroid therapy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
  - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

Subsidy	ce)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
  - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
  - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

## Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

- -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

# **Hyperuricaemia and Antigout**

ALLOPURINOL			
* Tab 100 mg	11.47	500	<ul><li>DP-Allopurinol</li></ul>
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see S	SA1963 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite S29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	✓ Benzbromaron AL
			<b>100</b> \$29

## ⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

## 

FEBUXOSTAT - Special Authority see SA2054 below	w – Retail pharmacy		
Tab 80 mg	20.00	28	✓ Febuxostat multichem
	39.50		✓ Adenuric
Tab 120 mg	20.00	28	✓ Febuxostat multichem
	39.50		✓ Adenuric

(Adenuric Tab 80 mg to be delisted 1 January 2022)

(Adenuric Tab 120 mg to be delisted 1 January 2022)

## **⇒SA2054** Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

## Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or

continued...

100

Colgout

Subsidy (Manufacturer's Price)	Full <sub>y</sub> Subsidise	
 \$	Per 🗸	Manufacturer

continued...

- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

**Renewal** — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

### **PROBENECID**

DACI OFFN

### Muscle Relaxants

DA	GLOFEN		
*	Tab 10 mg4.20	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement11.55	1	<ul> <li>Lioresal Intrathecal</li> </ul>

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.

Inj 2 mg per ml, 5 ml ampoule − Subsidy by endorsement...........306.82 5 **Medsurge** 

- a) Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.
- b) Medsurge to be Principal Supply on 1 December 2021

## DANTROLENE

Cap 25 mg97.50	100	✓ Dantrium ✓ Dantrium S29 S29
Cap 50 mg77.00	100	✓ Dantrium
ORPHENADRINE CITRATE		
Tab 100 mg 20.76	100	✓ Norflex

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	
Agents for Parkinsonism and Related Disorde	rs			
Dopamine Agonists and Related Agents				
AMANTADINE HYDROCHLORIDE  ▲ Cap 100 mg	38.24	60	✓	Symmetrel
APOMORPHINE HYDROCHLORIDE  ▲ Inj 10 mg per ml, 2 ml ampoule		5 5		Movapo Movapo
BROMOCRIPTINE MESYLATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of bromocriptine mesylate.				
* Tab 2.5 mg		30		Parlodel \$29
	32.08	100	•	Apo-Bromocriptine
(Parlodel S29 Tab 2.5 mg to be delisted 1 March 2022) (Apo-Bromocriptine Tab 2.5 mg to be delisted 1 March 2022)				
ENTACAPONE	00.00	400	,	
▲ Tab 200 mg	22.00	100	•	Entapone
LEVODOPA WITH BENSERAZIDE				
* Tab dispersible 50 mg with benserazide 12.5 mg		100		Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100 100		Madopar 62.5 Madopar 125
Cap 100 mg with benserazide 25 mg  Cap long-acting 100 mg with benserazide 25 mg		100		Madopar HBS
Cap 101g-acting 100 mg with benserazide 20 mg      Cap 200 mg with benserazide 50 mg		100		Madopar 250
	20.20	100	•	madopai 200
LEVODOPA WITH CARBIDOPA	01.11	100	./	Sinemet
<ul> <li>Tab 100 mg with carbidopa 25 mg</li> <li>Tab long-acting 200 mg with carbidopa 50 mg</li> </ul>		100		Sinemet CR
* Tab 250 mg with carbidopa 25 mg		100		Sinemet
PRAMIPEXOLE HYDROCHLORIDE		100	-	<u>unioniot</u>
▲ Tab 0.25 mg	6 12	100	1	Ramipex
▲ Tab 1 mg		100		Ramipex
RASAGILINE	20.70	100	•	паппрех
	E0 E0	20	./	A-ilest con
* Tab 1 mg	33.30	30	٧	Azilect S29
ROPINIROLE HYDROCHLORIDE				
▲ Tab 0.25 mg		84		Ropin
A Tab Amar	3.39	100		Mylan S29
▲ Tab 1 mg		84		Ropin
A. Tala O area	4.70	100		Mylan S29
▲ Tab 2 mg		84		Ropin
▲ Tab 5 mg	12.50	84	•	Ropin

## SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of

	prior dispensing of selegiline hydrochloride.	ic the prescription	i as criadisca	Where there exists a record o
*	Tab 5 mg	22.00	100	✓ Apo-Selegiline

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TOLCAPONE  A Tab 100 mg	152.38	100	✓:	<b>Tasmar</b>
Anticholinergics				
BENZATROPINE MESYLATE  Tab 2 mg Inj 1 mg per ml, 2 ml  a) Up to 10 inj available on a PSO b) Only on a PSO PROCYCLIDINE HYDROCHLORIDE		60 5		Benztrop Phebra
Tab 5 mg	7.40	100	<b>✓</b> I	Kemadrin
Agents for Essential Tremor, Chorea and Relate	ed Disorders			
RILUZOLE - Special Authority see SA1403 below - Retail pharr Wastage claimable Tab 50 mg Rilutek to be Principal Supply on 1 December 2021  SA1403 Special Authority for Subsidy	•	56	<b>√</b>	Rilutek
Initial application only from a neurologist or respiratory specialis	st. Approvals valid fo	r 6 m	onths for an	polications meeting the

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
  - 5.1 The patient is ambulatory: or
  - 5.2 The patient is able to use upper limbs; or
  - 5.3 The patient is able to swallow.

**Renewal** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
  - 3.1 The patient is ambulatory: or
  - 3.2 The patient is able to use upper limbs; or
  - 3.3 The patient is able to swallow.

## TETRABENAZINE

Tab 25 mg	91.10	112	✓ Motetis
-----------	-------	-----	-----------

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Price) \$	Per	Subsidised ✓	Manufacturer
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE]				
Gel 2%, tube – Subsidy by endorsement		30 ml		Xylocaine 2% Jelly
<ul> <li>b) Subsidised only if prescribed for urethral or cervical a</li> <li>Gel 2%, 11 ml urethral syringe – Subsidy by endorsement</li> <li>a) Up to 5 each available on a PSO</li> </ul>		e pres 10		endorsed accordingly.  Instillagel Lido
<ul> <li>b) Subsidised only if prescribed for urethral, cervical or r accordingly.</li> </ul>	rectal administration	and th	ne prescrip	otion is endorsed
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%		200 m		Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25		Lidocaine-Baxter Lidocaine-Claris
	17.50	50		
	(35.00)		_	Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO	8.25	25		<u>Lidocaine-Baxter</u> <u>Lidocaine-Claris</u>
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO	12.00 (20.00)	5		Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	( /	5	1	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO		5	✓	Lidocaine-Baxter Lidocaine-Claris
(Lidocaine-Claris Inj 1%, 5 ml ampoule to be delisted 1 January 2 (Lidocaine-Claris Inj 2%, 5 ml ampoule to be delisted 1 January 2				
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE	,			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –			_	
Subsidy by endorsement	103.32	10	•	Pfizer
<ul><li>a) Up to 5 each available on a PSO</li><li>b) Subsidised only if prescribed for urethral or cervical a</li></ul>	dministration and the	e pres	cription is	endorsed accordingly.
Topical Local Anaesthetics				
■ SA0906 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid condition requiring frequent injections or venepuncture.  Renewal from any relevant practitioner. Approvals valid for 2 years.				
benefiting from treatment.				
LIDOCAINE ILICNOCAINEL Special Authority and SA0006 abo	Dotoil phormas	.,		

LIDOCAINE [LIGNOCAINE] — Special Authority see SA0906 above -	- Retall phar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Retai	I pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

## **NERVOUS SYSTEM**

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

90

✓ Acupan

Brand or Generic Manufacturer

# **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 112

Tab 30 mg ......23.40

,,	- 3		
Non-opioid Analgesics			
ASPIRIN  * Tab dispersible 300 mg - Up to 30 tab available on a PSO.	4.50	100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or of accordingly.	diabetic periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream   \$29
NEFOPAM HYDROCHLORIDE			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PARACETAMOL	· · ·			
Tab 500 mg - blister pack	0.50	20	1	Medco
			✓	Pharmacy Health
	1.12		✓	Ethics Paracetamol
				Classic
	2.48	100	✓	Pharmacy Health
	5.01	50	✓	Panadol
	11.75	96	✓	Panadol Mini Caps
	19.75	1,000	✓	Pacimol
	24.82		✓	Paracetamol
				Pharmacare
			✓	Pharmacare

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

Tab 500 mg - bottle pack - Maximum of 300 tab per		
prescription; can be waived by endorsement17.92	1,000	✓ Noumed
		Paracetamol
24.82		<ul> <li>Paracetamol</li> </ul>
		Pharmacare

a)

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed nations), then dispense in repeat dispensings not exceeding 100 tab her dispensing

(for non-endorsed patients), then dispense in repe b) Noumed Paracetamol to be Sole Supply on 1 December	, ,	s not exceedin	g 100 tab per dispensing.
* Oral lig 120 mg per 5 ml	5.45	1,000 ml	✓ Paracare
a) Up to 200 ml available on a PSO			
b) Not in combination			
* Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double
			<u>Strength</u>
a) Up to 100 ml available on a PSO			
b) Not in combination			
	3.29	10	✓ Gacet
	3.79	10	✓ Gacet
	12.40	50	✓ Gacet
Medco Tab 500 mg - blister pack to be delisted 1 February 2022)			
Pharmacy Health Tab 500 mg - blister pack to be delisted 1 Februa	ary 2022)		
Ethics Paracetamol Classic Tab 500 mg - blister pack to be deliste	d 1 February 2	2022)	
Pharmany Haalth Tah 500 mg blister neak to be delicted 1 Eabrus	ani 2022)		

(Pharmacy Health Tab 500 mg - blister pack to be delisted 1 February 2022)

(Panadol Tab 500 mg - blister pack to be delisted 1 February 2022)

(Panadol Mini Caps Tab 500 mg - blister pack to be delisted 1 February 2022)

(Paracetamol Pharmacare Tab 500 mg - blister pack to be delisted 1 February 2022)

(Pharmacare Tab 500 mg - blister pack to be delisted 1 February 2022)

(Paracetamol Pharmacare Tab 500 mg - bottle pack to be delisted 1 December 2021)

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
Opioid Analgesics				
CODEINE PHOSPHATE - Safety medicine; prescriber may det	ermine dispensing fre	quency		
Tab 15 mg	6.25	100	<b>√</b> <u>F</u>	<u>PSM</u>
Tab 30 mg		100	<b>√</b> <u>F</u>	
Tab 60 mg	14.25	100	<b>✓</b> <u>F</u>	<u>'SM</u>
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg	8.60	60	✓ [	HC Continus
FENTANYL				
<ul> <li>a) Only on a controlled drug form</li> </ul>				
b) No patient co-payment payable				
Safety medicine; prescriber may determine dispensing fr Inj 50 mcg per ml, 2 ml ampoule		10	./ 5	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10		Boucher and Muir
Patch 12.5 mcg per hour		5		entanyl Sandoz
Patch 25 mcg per hour		5		entanyi Sandoz
Patch 50 mcg per hour		5		entanyl Sandoz
Patch 75 mcg per hour	17.99	5	<b>√</b> F	entanyl Sandoz
Patch 100 mcg per hour	18.59	5	<b>√</b> F	entanyl Sandoz
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr				
d) Extemporaneously compounded methadone will only be	reimbursed at the rate	e of the c	neapest	form available
<ul><li>(methadone powder, not methadone tablets).</li><li>e) For methadone hydrochloride oral liquid refer Standard F</li></ul>	Formulae page 040			
Tab 5 mg		10	<b>✓</b> 1	/lethatabs
Oral lig 2 mg per ml		200 ml		Biodone
Oral liq 5 mg per ml		200 ml	_	Biodone Forte
Oral liq 10 mg per ml		200 ml	<b>√</b> E	Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	<b>√</b> µ	\FT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
<ul> <li>c) Safety medicine; prescriber may determine dispensing fr</li> </ul>				
Oral liq 1 mg per ml		200 ml		RA-Morph
Oral liq 2 mg per ml		200 ml		RA-Morph
Oral liq 5 mg per ml	19.44	200 ml	-	Ordine S29
Oral lia 10 ma nor mi	07.74	200 ml		RA-Morph
Oral liq 10 mg per ml	27.74	200 ml	-	Ordine S29

✓ RA-Morph

## **NERVOUS SYSTEM**

	Subsidy		Fully Subsidised	
<b>'</b>	Manufacturer's Price) \$	Per		Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free	uency			
Tab immediate-release 10 mg	2.80	10	✓	Sevredol
Tab immediate-release 20 mg	5.52	10	✓	Sevredol
Cap long-acting 10 mg	2.05	10	✓	m-Eslon
Cap long-acting 30 mg	3.00	10	✓	m-Eslon
Cap long-acting 60 mg		10	✓	m-Eslon
Cap long-acting 100 mg	7.13	10	✓	m-Eslon
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC	D6.99	5	•	DBL Morphine Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	SO5.61	5	•	DBL Morphine Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO7.08	5	•	DBL Morphine Sulphate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO7.28	5	✓	DBL Morphine

Sulphate

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacturer
OXYCODONE HYDROCHLORIDE	*			
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	equency			
Tab controlled-release 5 mg		20	1	Oxycodone Sandoz
	3.01	28		Oxycodone Sandoz
				S29 S29
Tab controlled-release 10 mg	2 15	20	/	Oxycodone Sandoz
Tab controlled foldage to frig	3.23	30		Oxycodone Sandoz
				S29 S29
	5.38	50	/	Oxycodone Sandoz
	0.00	00	•	S29 S29
	10.75	100	1	Oxycodone Sandoz
	10.75	100	•	S29 S29
	11.50	28	1	OxyContin
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg	5.38	50		Oxycodone Sandoz
	3.00	50	•	S29 S29
	10.75	100		0_0
	10.75	100	•	Oxycodone Sandoz
	40.05	00	,	S29 S29
Tab controlled valence 40 mm	13.25	28		OxyContin
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20 20		Oxycodone Sandoz OxyNorm
OxyNorm to be Principal Supply on 1 December 2021	1.00	20	•	Oxynoriii
Cap immediate-release 10 mg	3 32	20	1	OxyNorm
OxyNorm to be Principal Supply on 1 December 2021		20	•	Oxymoniii
Cap immediate-release 20 mg	5.23	20	1	OxyNorm
OxyNorm to be Principal Supply on 1 December 2021				,
Oral lig 5 mg per 5 ml	11.20	250 m	· •	OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5	✓	OxyNorm
Inj 50 mg per ml, 1 ml ampoule	30.60	5	1	OxyNorm
PARACETAMOL WITH CODEINE - Safety medicine; prescribe	r mav determine dispe	ensino	ı freauenc	V
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
		,		Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fr	equency			
Tab 50 mg		10	1	PSM
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a l	PSO29.88	5	✓	DBL Pethidine
				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a l	PSO30.72	5	✓	DBL Pethidine
				Hydrochloride
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	1	Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
. •				

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 10 mg		100		Arrow-Amitriptyline
Tab 25 mg Tab 50 mg		100		Arrow-Amitriptyline Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; pres				
Tab 10 mg	•	30		Clomipramine
· • • • • • • • • • • • • • • • • • • •				Teva S29
	13.99	100	1	Apo-Clomipramine
Tab 25 mg	9.46	100		Apo-Clomipramine
	11.99	30	•	Clomipramine
(A O) : T   (O)   (A   (A   (A   (A   (A   (A   (A   (				Teva S29
(Apo-Clomipramine Tab 10 mg to be delisted 1 February 2022) (Apo-Clomipramine Tab 25 mg to be delisted 1 February 2022)				
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by a Safety medicine; prescriber may determine dispensing				
<ul><li>b) Subsidy by endorsement – Subsidised for patients who</li></ul>		[dothi	eninl hvdr	ochloride prior to 1 June
2019 and the prescription is endorsed accordingly. Pha				
exists a record of prior dispensing of dosulepin [dothiep				
Tab 75 mg		30	_	Dosulepin Mylan
Cap 25 mg	7.83	50	/	Dosulepin
				Mylan S29
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescribe	,	•		
Tab 10 mg	5.48 10.96	50 100		Tofranil Tofranil
Tab 25 mg		50		Tofranil
NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; pres				
Tab 10 mg		100		Norpress
Tab 25 mg		180	_	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non	Solootivo			
	Selective			
TRANYLCYPROMINE SULPHATE			_	
Tab 10 mg		28	_	Parnate S29 S29
	22.94	50		Parnate
	45.88	100	_	Parnate S29 S29
	96.00			Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE				
* Tab 150 mg		60		Aurorix
* Tab 300 mg	19.25	60	<b>✓</b>	Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE				
* Tab 20 mg	1.91	84	1	PSM Citalopram

<del></del>			
	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
SCITALOPRAM			
€ Tab 10 mg	1.07	28	✓ Escitalopram (Ethics)
	1.40		✓ Escitalopram- Apotex
Escitalopram (Ethics) to be Sole Supply on 1 October 2	2021		
: Tab 20 mg	1.92	28	✓ Escitalopram (Ethics)
	2.49		<ul><li>✓ Escitalopram- Apotex</li></ul>
Escitalopram (Ethics) to be Sole Supply on 1 October 2 Socitalopram-Apotex Tab 10 mg to be delisted 1 October 2021 Socitalopram-Apotex Tab 20 mg to be delisted 1 October 2021 LUOXETINE HYDROCHLORIDE	')		
Tab dispersible 20 mg, scored — Subsidy by endorsement     Subsidised by endorsement	1.98	30	✓ Fluox
When prescribed in a daily dose that is not a mul endorsed. Note: Tablets should be combined w  Cap 20 mg	ith capsules to facilitate		
1 0			
AROXETINE			TIUOX
	1.20	30	✓ Paxtine
F Tab 20 mg	1.20 3.61	30 90	
Tab 20 mg			✓ Paxtine
· Tab 20 mg Paxtine Tab 20 mg to be delisted 1 January 2022) ERTRALINE	3.61		✓ Paxtine
Tab 20 mg	3.61 0.92 3.05	90	✓ Paxtine ✓ Loxamine ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline
Fab 20 mg	3.61 0.92 3.05 1.61	90 30 90 30	✓ Paxtine ✓ Loxamine ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona
Faxtine Tab 20 mg to be delisted 1 January 2022)  ERTRALINE  Tab 50 mg  Tab 100 mg  Setrona AU Tab 50 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021)  Setrona AU Tab 100 mg to be delisted 1 October 2021)	3.61 0.92 3.05	90 30 90	✓ Paxtine ✓ Loxamine ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona
E Tab 20 mg	3.61 0.92 3.05 1.61	90 30 90 30	✓ Paxtine ✓ Loxamine ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona
Paxtine Tab 20 mg to be delisted 1 January 2022)  ERTRALINE Tab 50 mg Tab 100 mg  Ertrona AU Tab 50 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021)  Setrona AU Tab 100 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)  Other Antidepressants	3.61 0.92 3.05 1.61	90 30 90 30	✓ Paxtine ✓ Loxamine ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona
Paxtine Tab 20 mg to be delisted 1 January 2022)  ERTRALINE Tab 50 mg Tab 100 mg  Ertrona AU Tab 50 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021)  Setrona AU Tab 100 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)  Other Antidepressants	3.61 0.92 3.05 1.61 5.25	90 30 90 30	✓ Paxtine ✓ Loxamine  ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline
Paxtine Tab 20 mg to be delisted 1 January 2022)  ERTRALINE Tab 50 mg  Tab 100 mg  Setrona AU Tab 50 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021)  Setrona AU Tab 100 mg to be delisted 1 October 2021)  Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021)  Other Antidepressants  IIRTAZAPINE Tab 30 mg	3.61 0.92 3.05 1.61 5.25	90 30 90 30 90 28 30	✓ Paxtine ✓ Loxamine  ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline  ✓ Apo-Mirtazapine
Setrona AU Tab 50 mg to be delisted 1 October 2021) Arrow-Sertraline Tab 50 mg to be delisted 1 October 2021) Setrona AU Tab 100 mg to be delisted 1 October 2021) Arrow-Sertraline Tab 100 mg to be delisted 1 October 2021) Other Antidepressants IIRTAZAPINE	3.61 0.92 3.05 1.61 5.25	90 30 90 30 90	✓ Paxtine ✓ Loxamine  ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline ✓ Setrona ✓ Setrona AU ✓ Arrow-Sertraline

<sup>(</sup>Apo-Mirtazapine Tab 30 mg to be delisted 1 January 2022) (Apo-Mirtazapine Tab 45 mg to be delisted 1 January 2022)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VENLAFAXINE				
* Cap 37.5 mg		84		Enlafax XR
* Cap 75 mg		84		Enlafax XR
* Cap 150 mg	11.16	84	•	Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM - Safety medicine; prescriber may determin	ne dispensing frequency			
Inj 1 mg per ml, 1 ml	21.00	5	1	Rivotril
(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)				
DIAZEPAM - Safety medicine; prescriber may determine d				
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorseme	ent23.66	5	1	Hospira
<ul> <li>a) Up to 5 inj available on a PSO</li> </ul>				
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic pro				
Rectal tubes 5 mg - Up to 5 tube available on a PSO	43.50	5	✓	Stesolid
PARALDEHYDE				
* Inj 5 ml	1.500.00	5	1	AFT S29
AFT S29 Inj 5 ml to be delisted 1 October 2021)		Ū		
•				
PHENYTOIN SODIUM	- DOO 00.00	_	,	
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available o		5	•	Hospira
★ Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available of		_		
PSO	133.92	5	•	Hospira
Control of Epilepsy				
CARBAMAZEPINE				_
* Tab 200 mg		100		Tegretol
* Tab long-acting 200 mg		100		Tegretol CR
* Tab 400 mg		100		Tegretol
* Tab long-acting 400 mg		100		Tegretol CR
* Oral liq 20 mg per ml		250 m	ı 🗸	Tegretol
CLOBAZAM – Safety medicine; prescriber may determine of	dispensing frequency			
Tab 10 mg	9.12	50	1	Frisium
CLONAZEPAM - Safety medicine; prescriber may determine	ne dispensing frequency			
Oral drops 2.5 mg per ml		0 ml C	)P 🗸	Rivotril
ETHOSUXIMIDE				
Cap 250 mg	140.88	100	1	Zarontin
Oral lig 250 mg per 5 ml		200 m		Zarontin

Oral liq 250 mg per 5 ml ......56.35

200 ml

✓ Zarontin

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
GABAPENTIN				
Note: Not subsidised in combination with subsidised pregaba	alin			
* Cap 100 mg	2.65	100	✓	Apo-Gabapentin
	6.45		1	Nupentin .
* Cap 300 mg	4.07	100	1	Apo-Gabapentin
	8.45			Nupentin
* Cap 400 mg	5.64	100		Apo-Gabapentin
	10.26			Nupentin
(Apo-Gabapentin Cap 100 mg to be delisted 1 February 2022) (Apo-Gabapentin Cap 300 mg to be delisted 1 February 2022) (Apo-Gabapentin Cap 400 mg to be delisted 1 February 2022)				
LACOSAMIDE - Special Authority see SA1125 below - Retail ph	narmacv			
▲ Tab 50 mg	•	14	1	Vimpat
▲ Tab 100 mg		14		Vimpat
<b>3</b>	200.24	56		Vimpat
▲ Tab 150 mg	75.10	14		Vimpat
•	300.40	56	_	Vimpat
▲ Tab 200 mg	400.55	56		Vimpat

## ⇒SA1125 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

### LAMOTRIGINE

$\blacktriangle$	Tab dispersible 2 mg	55.00	30	✓ Lamictal
$\blacktriangle$	Tab dispersible 5 mg	50.00	30	✓ Lamictal
*	Tab dispersible 25 mg		56	✓ Logem
*	Tab dispersible 50 mg		56	✓ Logem
*	Tab dispersible 100 mg		56	✓ Logem
LE	VETIRACETAM			
	Tab 250 mg	4.99	60	✓ Everet
	Tab 500 mg	8.79	60	✓ Everet
	Tab 750 mg		60	✓ Everet
	Tab 1,000 mg	18.59	60	✓ Everet
	Oral liq 100 mg per ml		300 ml OP	✓ Levetiracetam-AFT
РΗ	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae, p	age 242		
*	Tab 15 mg		500	✓ PSM
*	Tab 30 mg	40.00	500	✓ PSM

	Subsidy		Fully	Brand or
(	(Manufacturer's F		Subsidised	
	\$	Per		Manufacturer
PHENYTOIN SODIUM				
* Tab 50 mg	75.00	200	1	Dilantin Infatab
Cap 30 mg	74.00	200	✓	Dilantin
Cap 100 mg	37.00	200	✓	Dilantin
Oral liq 30 mg per 5 ml	22.03	500 m	✓	Dilantin
REGABALIN				
Note: Not subsidised in combination with subsidised gabapen	ıtin			
€ Cap 25 mg		56	✓	Pregabalin Pfizer
€ Cap 75 mg		56	✓	Pregabalin Pfizer
Cap 150 mg		56		Lyrica
•			1	Pregabalin Pfizer
Cap 300 mg	7.38	56	✓	Pregabalin Pfizer
RIMIDONE				
F Tab 250 mg	17.25	100	1	Apo-Primidone
ODIUM VALPROATE				•
Tab 100 mg	13.65	100	1	Epilim Crushable
Tab 200 mg EC		100		Epilim
Tab 500 mg EC		100		Epilim
Oral liq 200 mg per 5 ml		300 m		Epilim S/F Liquid
				Epilim Syrup
€ Inj 100 mg per ml, 4 ml	41.50	1		Epilim IV
TIRIPENTOL - Special Authority see SA1330 below - Retail pha				•
Cap 250 mg	•	60	1	Diacomit S29
Powder for oral lig 250 mg sachet		60		Diacomit \$29
Fowder for oral lig 200 mg Sachet	509.29	00	•	Diacollill

## **⇒SA1330** Special Authority for Subsidy

**Initial application** only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TO	PIRA	MΑ	ΤE
	T - 1-	0-	

$\blacktriangle$	Tab 25 mg	11.07	60	Arrow-Topiramate
	-			✓ Topiramate Actavis
		26.04		✓ Topamax
$\blacktriangle$	Tab 50 mg	18.81	60	✓ Arrow-Topiramate
	-			✓ Topiramate Actavis
		44.26		✓ Topamax
$\blacktriangle$	Tab 100 mg	31.99	60	✓ Arrow-Topiramate
	-			✓ Topiramate Actavis
		75.25		✓ Topamax
$\blacktriangle$	Tab 200 mg	55.19	60	✓ Arrow-Topiramate
				✓ Topiramate Actavis
		129.85		✓ Topamax
$\blacktriangle$	Sprinkle cap 15 mg	20.84	60	✓ Topamax
$\blacktriangle$	Sprinkle cap 25 mg	26.04	60	✓ Topamax
VIG	ABATRIN - Special Authority see SA1997 on the next page -	- Retail pharmacy	,	
	Tab 500 mg		100	✓ Sabril

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

## ⇒SA1997 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and
    - 1.2.2 Fither:
      - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
      - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Fither:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

. -

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either
  - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

# **Antimigraine Preparations**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 112

# **Acute Migraine Treatment**

RIZATRIPTAN			
Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	14.41	90	✓ Sumagran
	24.44	100	✓ Apo-Sumatriptan
Tab 100 mg	22.68	90	✓ Sumagran
	46.23	100	✓ Apo-Sumatriptan
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	34.00	2 OP	✓ Imigran
(Apo-Sumatriptan Tab 50 mg to be delisted 1 February 2022)			<del></del> -

(Apo-Sumatriptan Tab 100 mg to be delisted 1 February 2022)



Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
 \$	Per	1	Manufacturer

## **Prophylaxis of Migraine**

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 51

**PIZOTIFEN** 

# **Antinausea and Vertigo Agents**

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

Emend Tri-Pack to be Principal Supply on 1 December 2021

## **⇒SA0987** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

## BETAHISTINE DIHYDROCHLORIDE

* Tab 16 mg	3.88	84	✓ Vergo 16
CYCLIZINE HYDROCHLORIDE			
Tab 50 mg	0.49	10	✓ Nausicalm
Nausicalm to be Principal Supply on 1 December 2021			
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml	21.53	10	✓ <u>Hameln</u>
DOMPERIDONE			
* Tab 10 mg	2.85	100	Pharmacy Health
HYOSCINE HYDROBROMIDE			
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1998 below - Retail			
pharmacy	14.11	2	Scopoderm TTS

## ⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

## METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO	1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO	9.50	10	✓ Pfizer

_		Subsidy		Fully	
		(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
<u></u>	IDANSETRON				
*	Tab 4 mg	2.68	50	1	Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO		10		Ondansetron
	3 -1				ODT-DRLA
*	Tab 8 mg	4.57	50	1	Onrex
*	Tab disp 8 mg – Up to 10 tab available on a PSO		10		Ondansetron
•			. •		ODT-DRLA
P۲	OCHLORPERAZINE				
*	Tab 3 mg buccal	5.97	50		
	<b>3</b>	(30.00)			Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO		250	1	Nausafix
	Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO		10		Stemetil
A	ntipsychotics				
C	ieneral				
_					
٩N	IISULPRIDE – Safety medicine; prescriber may determine d			_	
	Tab 100 mg		30	/	Sulprix
		17.16	100	•	Amisulpride
					Mylan S29
	Tab 200 mg	14.96	60	1	Sulprix
	Tab 400 mg	29.78	60	1	Sulprix
۸R	IPIPRAZOLE – Safety medicine; prescriber may determine	dispensing frequency			— <del>.</del>
	Tab 5 mg		30	1	Aripiprazole Sandoz
	Tab 10 mg		30		Aripiprazole Sandoz
	Tab 15 mg		30		Aripiprazole Sandoz
	Tab 20 mg		30		Aripiprazole Sandoz
	Tab 30 mg		30		Aripiprazole Sandoz
-ш	ILORPROMAZINE HYDROCHLORIDE – Safety medicine; p				
חכ	Tab 10 mg – Up to 30 tab available on a PSO		100		Largactil
	Tab 25 mg - Up to 30 tab available on a PSO		100		Largactil
	Tab 100 mg - Up to 30 tab available on a PSO		100		Largactil
	Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO		100		Largactil
٥.			10	•	Largactii
JL	OZAPINE – Hospital pharmacy [HP4]				
	Safety medicine; prescriber may determine dispensing freq			,	01
	Tab 25 mg		50		Clozaril
		6.69	400		Clopine
		11.36	100		Clozaril
	T   50	13.37			Clopine
	Tab 50 mg		50		Clopine
	Tab 400	17.33	100		Clopine
	Tab 100 mg		50		Clozaril
		17.33	400		Clopine
		29.45	100		Clozaril
	<b>-</b>	34.65			Clopine
	Tab 200 mg		50		Clopine
		69.30	100		Clopine
	Suspension 50 mg per ml	17.33	100 n		Clopine
		67.62			Vorcanioz

67.62

✓ Versacloz

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
HALOPERIDOL - Safety medicine; prescriber may determine of	lispensina frequency			
Tab 500 mcg - Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
	29.72	100		Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO	23.84	100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a F		10	✓	Serenace
LEVOMEPROMAZINE – Safety medicine; prescriber may dete		ulonov		
Tab 25 mg (33.8 mg as a maleate)	, ,	100		Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan (Swiss)
· · · · · · · · · · · · · · · · · · ·		100		Nozinan (Swiss)
Tab 100 mg (135 mg as a maleate)		100	_	Nozinan (Swiss)
Tab 100 mg as a maleate				
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;				, ,
Inj 25 mg per ml, 1 ml ampoule	33.50	10	•	<u>Nozinan</u>
LITHIUM CARBONATE - Safety medicine; prescriber may dete	ermine dispensing free	quency		
Tab long-acting 400 mg	72.00	100	✓	Priadel
Cap 250 mg	9.42	100	✓	Douglas
OLANZAPINE - Safety medicine; prescriber may determine dis				
Tab 2.5 mg		28	1	Zypine
Tab 5 mg		28	_	Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28		Zypine
Tab orodispersible 10 mg		28		Zypine ODT
		20	•	Lypine OD I
PERICYAZINE – Safety medicine; prescriber may determine di		0.4		Naulastil
Tab 2.5 mg		84		Neulactil
T.I. 40	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
	44.45	100	•	Neulactil
QUETIAPINE – Safety medicine; prescriber may determine disp				
Tab 25 mg		90		Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg	12.86	90	/	Quetapel
RISPERIDONE - Safety medicine; prescriber may determine d	ispensing frequency			
Tab 0.5 mg	1.86	60	✓	Risperidone (Teva)
Tab 1 mg	2.06	60	✓	Risperidone (Teva)
Tab 2 mg	2.29	60	✓	Risperidone (Teva)
Tab 3 mg	2.50	60	✓	Risperidone (Teva)
Tab 4 mg	3.42	60	✓	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 ml	✓	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine di	snensing frequency			·
Cap 20 mg		60	1	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60		Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; pro	•		-	
Tab 10 mg	31.45	100	•	Clopixol

Subsidy		Fully	Brand or	
(Manufacturer's Price)	) Su	ıbsidised	Generic	
\$	Per	<b>✓</b>	Manufacturer	

## **Depot Injections**

FLUPENTHIXOL DECANOATE - Safety medicine; preso	criber may determine dispen	sing frequ	iency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PS	iO13.14	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PS	O20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a P	SO40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescr	riber may determine dispens	ing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PS	O28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a P	SO55.90	5	✓ Haldol Concentrate
			✓ Haldol
			Decanoas \$29
OLANZAPINE - Special Authority see SA1428 below - F	Retail pharmacy		
Safety medicine; prescriber may determine dispensin	g frequency		
Inj 210 mg vial	252.00	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓ Zyprexa Relprevv

## ⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

## PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensir	ng frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

## ⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the



	Subsidy	Full	/ Brand or
(Manu	ufacturer's Price)	Subsidise	d Generic
	\$ F	Per 🗸	Manufacturer

continued...

initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing f	requency		
Inj 25 mg vial	135.98	1	<ul> <li>Risperdal Consta</li> </ul>
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

## ⇒SA1427 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:

Anxiolytics

2.1 The patient has schizophrenia or other psychotic disorder; and

Tab 15 mg ......8.53

- 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
- 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO ......19.80 5 **Clopixol** 

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine dispe	ensing frequenc	у	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispensir	ng frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine dispen	sing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Ativan to be Principal Supply on 1 December 2021			
Tab 2.5 mg Ativan to be Principal Supply on 1 December 2021	12.50	100	✓ Ativan
OXAZEPAM – Safety medicine; prescriber may determine dispens	0 ,	100	✓ Ox-Pam

Ox-Pam

100

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

## **Multiple Sclerosis Treatments**

⇒SA2051 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
  - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
  - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
  - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
  - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
  - 5.5 Either:
    - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
    - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
  - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
  - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
  - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
  - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
  - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. **Renewal — (Multiple sclerosis)** only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2051 above - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg	520.00	14	Tecfidera
Cap 240 mg	2,000.00	56	✓ Tecfidera

FINGOLIMOD - Special Authority see SA2051 above - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 0.5 mg......2,200.00 28 **✓ Gilenya** 



	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully Brand or idised Generic ✓ Manufacturer
GLATIRAMER ACETATE - Special Authority see SA2051 on th			•
Note: Treatment on two or more funded multiple sclerosis tr Inj 40 mg prefilled syringe		ısly is not p 12	permitted. <b>✓ Copaxone</b>
INTERFERON BETA-1-ALPHA – Special Authority see SA2051 Note: Treatment on two or more funded multiple sclerosis tr Inj 6 million iu prefilled syringe	eatments simultaneou 1,170.00		•
INTERFERON BETA-1-BETA – Special Authority see SA2051 of Note: Treatment on two or more funded multiple sclerosis truling 8 million iu per 1 ml	eatments simultaneou		
NATALIZUMAB – Special Authority see SA2051 on the previous  Note: Treatment on two or more funded multiple sclerosis tr Inj 20 mg per ml, 15 ml vial	eatments simultaneou	•	permitted. <b>✓ Tysabri</b>
OCRELIZUMAB – Special Authority see SA2051 on the previous Note: Treatment on two or more funded multiple sclerosis truin 30 mg per ml, 10 ml vial	eatments simultaneou	•	permitted.  ✓ Ocrevus
TERIFLUNOMIDE - Special Authority see SA2051 on the previous	ous page - Retail pha	rmacy	
a) Wastage claimable     b) Note: Treatment on two or more funded multiple scleros     Tab 14 mg		eously is r 28	not permitted.  ✓ Aubagio

## Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

30

✓ Circadin

## ⇒SA1666 Special Authority for Subsidy

**Initial application** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

**Renewal** only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

## All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

Both:  1 For the treatment of terminal agitation that is unresponsive to other agents; and 2 The applicant is part of a multidisciplinary team working in palliative care.  TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg				NE	RVOUS SYSTEM
Inj 1 mg per ml, 5 ml ampoule			Per	Subsidised	Generic
1,1 mg per ml, 5 ml plastic ampoule — Up to 10 inj available on a PSO	MIDAZOLAM – Safety medicine; prescriber may determine dispe	ensing frequency			
On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. Inj 5 mg per ml, 3 ml ampoule	Inj 1 mg per ml, 5 ml ampoule		10		
Inj 5 mg per ml, 3 ml ampoule	on a PSO	14.90			
a PSO	Inj 5 mg per ml, 3 ml ampoule	4.50			
PHENOBARBITONE SODIUM — Special Authority see SA1386 below — Retail pharmacy Inj 200 mg per ml, 1 ml ampoule	a PSO	11.90			
Inj 200 mg per ml, 1 ml ampoule	On a PSO for status epilepticus use only. PSO must be	endorsed for status e	epilep	ticus use	only.
Inj 200 mg per ml, 1 ml ampoule	PHENOBARBITONE SODIUM - Special Authority see SA1386 b	oelow – Retail pharma	acy		
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:  Both:  1 For the treatment of terminal agitation that is unresponsive to other agents; and 2 The applicant is part of a multidisciplinary team working in palliative care.  TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency Tab 10 mg			-	1	Max Health S29
Tab 10 mg	Initial application from any relevant practitioner. Approvals valid the following criteria:  Both:  1 For the treatment of terminal agitation that is unresponsive	e to other agents; and		nless notif	ied for applications meeting
Tab 125 mcg			25	•	Normison
Tab 250 mcg		5.10	100		I home on
(11.20) Hypam  ZOPICLONE — Safety medicine; prescriber may determine dispensing frequency Tab 7.5 mg	Tah 250 mcg	` '	100		нурат
ZOPICLONE - Safety medicine; prescriber may determine dispensing frequency Tab 7.5 mg	Tab 250 Hicy		100		Hynam
Tab 7.5 mg       10.80       ✓ Zopiclone Actavis         Stimulants/ADHD Treatments         ATOMOXETINE       Cap 10 mg       18.41       28       ✓ Generic Partners         Cap 18 mg       27.06       28       ✓ Generic Partners         Cap 25 mg       29.22       28       ✓ Generic Partners         Cap 40 mg       29.22       28       ✓ Generic Partners         Cap 60 mg       46.51       28       ✓ Generic Partners         Cap 80 mg       56.45       28       ✓ Generic Partners	ZODICI ONE Safaty modicina: proceribor may determine dispe	( -/			Турат
ATOMOXETINE         Cap 10 mg       18.41       28	· · · · · · · · · · · · · · · · · · ·	•	500	•	Zopiclone Actavis
Cap 10 mg.       18.41       28       Generic Partners         107.03       Strattera         Cap 18 mg.       27.06       28       Generic Partners         107.03       Strattera         Cap 25 mg.       29.22       28       Generic Partners         Cap 40 mg.       29.22       28       Generic Partners         107.03       Strattera         Cap 60 mg.       46.51       28       Generic Partners         Cap 80 mg.       56.45       28       Generic Partners	Stimulants/ADHD Treatments				
107.03   Strattera	ATOMOXETINE				
Cap 18 mg.       27.06       28       Generic Partners         107.03       Strattera         Cap 25 mg.       29.22       28       Generic Partners         Cap 40 mg.       29.22       28       Generic Partners         107.03       Strattera         Cap 60 mg.       46.51       28       Generic Partners         Cap 80 mg.       56.45       28       Generic Partners	Cap 10 mg	18.41	28		
107.03       Strattera         Cap 25 mg       29.22       28       Generic Partners         Cap 40 mg       29.22       28       Generic Partners         107.03       Strattera         Cap 60 mg       46.51       28       Generic Partners         Cap 80 mg       56.45       28       Generic Partners					
Cap 25 mg.       29.22       28       ✓ Generic Partners         Cap 40 mg.       29.22       28       ✓ Generic Partners         107.03       ✓ Strattera         Cap 60 mg.       46.51       28       ✓ Generic Partners         Cap 80 mg.       56.45       28       ✓ Generic Partners	Cap 18 mg		28		
Cap 40 mg.       29.22       28       ✓ Generic Partners         107.03       ✓ Strattera         Cap 60 mg.       46.51       28       ✓ Generic Partners         Cap 80 mg.       56.45       28       ✓ Generic Partners	0 05		00		
107.03       ✓ Strattera         Cap 60 mg       46.51       28       ✓ Generic Partners         Cap 80 mg       56.45       28       ✓ Generic Partners	_ 1				
Cap 60 mg	Cap 40 mg		20		
Cap 80 mg	Cap 60 mg		28		•
· · ·	. •			_	

DEXAMFETAMINE SULFATE - Special Authority see SA1149 on the next page - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg ......21.00 100 ✓ PSM



	Subsidy	Fi	ıllv	Brand or
(Mar	nufacturer's Price)	Subsidis	,	Generic
<b>(</b>		Per	/	Manufacturer

## **⇒SA1149** Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 on the next page - Retail pharmacy

a	) (	)ni	٧	on	а	con	trol	led	С	Iruq	torm	

<ul> <li>b) Safety medicine; prescriber may determine disp</li> </ul>	ensing frequency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg	7.75	30	<ul><li>Methylphenidate ER</li></ul>
			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg		30	<ul><li>Methylphenidate ER</li></ul>
			- Teva
Tab extended-release 36 mg	15.50	30	<ul><li>Methylphenidate ER</li></ul>
•			- Teva
Tab extended-release 54 mg	22.25	30	<ul><li>Methylphenidate ER</li></ul>
Ç			- Teva

Subsidy	Fu	ılly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	✓ Manufacturer	

## ⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine: prescriber may determine dispensing frequency

by carety meaning, processes may actermine alopeneing i			
Tab extended-release 18 mg	58.96	30	<ul><li>Concerta</li></ul>
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg	71.93	30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg		30	✓ Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	✓ Ritalin LA
Cap modified-release 30 mg		30	✓ Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

## ⇒SA1965 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid



(Ma	Subsidy nufacturer's Price)	Sub	Fully	Brand or Generic
<u> </u>	\$	Per	✓	Manufacturer

continued...

for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 Fither:
    - 2.1 Applicant is a paediatrician or psychiatrist; or
    - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy Tab 100 mg ......64.00 60 ✓ Modaviqil

## ⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Fither:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatme	nts for	Dementia
DONEDEZII	LIVDDO	

טט	NEFEZIL HTDNOCHLONIDE		
*	Tab 5 mg4.34	90	✓ Donepezil-Rex
*	Tab 10 mg	90	✓ Donepezil-Rex

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
RIVASTIGMINE - Special Authority see SA1488 below - Retail	pharmacy			
Patch 4.6 mg per 24 hour	38.00	30	✓ [	Rivastigmine Patch BNM 5
	48.75		1	Generic Partners
Patch 9.5 mg per 24 hour	38.00	30	✓ [	Rivastigmine Patch BNM 10
	48.75		✓ (	Generic Partners
(Generic Partners Patch 4.6 mg per 24 hour to be delisted 1 Feb (Generic Partners Patch 9.5 mg per 24 hour to be delisted 1 Feb				

# ⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

# Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

✓ Buprenorphine Naloxone BNM

Tab sublingual 8 mg with naloxone 2 mg ......53.12

✓ Buprenorphine Naloxone BNM

28

⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent: and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health;
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use



	Subsidy	Fully	Brand or
(Manut	facturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

and another attempt is planned; and

- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 Patient received but failed detoxification with buprenorphine with naloxone; and
  - 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
  - 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
  - 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE			
Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM			
Tab 200 mg	236.40	100	Antabuse
Antabuse to be Principal Supply on 1 November 2021			
NALTREXONE HYDROCHLORIDE - Special Authority see SA1408 b	elow – Retail ph	armacy	
Tab 50 mg	133.33	30	✓ <u>Naltraccord</u>
' '		,	✓ <u>Naltraccord</u>

#### ⇒SA1408 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

✓ Habitrol

✓ Habitrol

384

96

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO ......18.14 ✓ Habitrol 28 Patch 7 mg for direct distribution only - [Xpharm]......3.94 7 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO ......19.95 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......4.52 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO ......22.86 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......5.18 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......19.18 216 ✓ Habitrol 36 ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......21.02 216 ✓ Habitrol 36 ✓ Habitrol ✓ Habitrol Gum 2 mg (Fruit) - Up to 384 piece available on a PSO ......38.21 384 96 ✓ Habitrol ✓ Habitrol Gum 2 mg (Mint) - Up to 384 piece available on a PSO......38.21 384 Gum 2 mg (Mint) for direct distribution only - [Xpharm]......8.64 96 ✓ Habitrol ✓ Habitrol Gum 4 mg (Fruit) - Up to 384 piece available on a PSO ......44.17 384 Gum 4 mg (Fruit) for direct distribution only - [Xpharm].................10.01 96 ✓ Habitrol

Gum 4 mg (Mint) for direct distribution only – [Xpharm]..................10.01

VARENICLINE TARTRATE – Special Authority see SA1845 below – Retail pharmacy

Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 4216.67	53 OP	✓ Varenicline Pfizer
25.64		Champix
Tab 1 mg17.62	56	✓ Varenicline Pfizer
27.10		Champix

(Champix Tab 0.5 mg  $\times$  11 and 1 mg  $\times$  42 to be delisted 1 January 2022) (Champix Tab 1 mg to be delisted 1 January 2022)

### **⇒SA1845** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and

# **NERVOUS SYSTEM**

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

# **Chemotherapeutic Agents**

### Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2046 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

### ⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 All of the following:
    - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
    - 3.2.2 The patient has not received prior bendamustine therapy; and
    - 3.2.3 Fither:
      - 3.2.3.1 Both:
        - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
        - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
      - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Fither:
  - 2.1 Both:

Subsidy (Manufacturer's Price)	Subi	Fully sidised	Brand or Generic
(Manufacturer's Price)	Per	siuiseu •	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1,387.00	1	✓ BiCNU
, ,	•		✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		ŭ	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ DBL Cisplatin
, 9 po,	21.00	·	✓ Cisplatin Ebewe
Inj 1 mg for ECP	0.25	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE		· ·	
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
	145.00		✓ Cyclonex
	158.00	100	✓ Procytox S29
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
, <b>3</b>	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
(Endoxan S29 Tab 50 mg to be delisted 1 January 2022)			
(Procytox \$29) Tab 50 mg to be delisted 1 January 2022)			
IFOSFAMIDE – PCT only – Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter
		-	

	Subsidy		Fully	Brand or
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	1	CeeNU
Cap 40 mg		20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1	1	Alkeran
, , ,			1	Alkeran S29 S29
	420.00		1	Tillomed S29
(Tillomed S29 Inj 50 mg to be delisted 1 December 2021)				
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
iiij 100 iiig viai	25.01	'	•	100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP		1 mg		Baxter
, ,		9		
THIOTEPA – PCT only – Specialist	ODO	1		Dedford 200
Inj 15 mg vial		ı		Bedford S29
				THIO-TEPA S29
				Tepadina S29
Inj 100 mg vial	CBS	1	/	Tepadina S29
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see	SA1467 helow			
Inj 100 mg vial		1	1	Azacitidine Dr

1  ✓ Azacitidine Dr	1	Inj 100 mg vial75.06
Reddy's		, 0
✓ Vidaza		605.00
		Azacitidine Dr Reddy's to be Principal Supply on 1 December 2021
1 mg Saxter	1 mg	Inj 1 mg for ECP0.83

#### ⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome: or
  - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
  - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

**Renewal** only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy	>	Fully	
	(Manufacturer's Pric	ce) Per	Subsidised •	I Generic Manufacturer
ALCIUM FOLINATE	·			
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	•	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specia		1		Calcium Folinate Sandoz
			✓	Calcium Folinate Sandoz S29 S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	•	Calcium Folinate Ebewe
Inj 300 mg - PCT only - Specialist	22.51	1	✓	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓	Calcium Folinate Sandoz
			•	Calcium Folinate Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	✓	Calcium Folinate Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓	Calcium Folinate Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	/	Baxter
APECITABINE - Retail pharmacy-Specialist				
Tab 150 mg		60	✓	Capercit
Tab 500 mg	49.00	120	/	Capercit
LADRIBINE - PCT only - Specialist				
Inj 1 mg per ml, 10 ml	749.96	1	✓	Leustatin
Inj 10 mg for ECP	749.96	10 mg C	)P 🗸	Baxter
/TARABINE				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia Inj 100 mg per ml, 20 ml vial – PCT – Retail		5		Pfizer
pharmacy-Specialist		1		Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg		Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specia	alist80.00	100 mg (	OP 🗸	Baxter
UDARABINE PHOSPHATE				
Tab 10 mg - PCT - Retail pharmacy-Specialist		20		Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg C	)P 🗸	Baxter
UOROURACIL		_	_	
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist		1		Fluorouracil Accord
Initia managari 400 milatah BOT da O da Mar	12.00			Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist		1		Fluorouracil Accord
	30.00		•	Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist		100 mg		Baxter

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e)	Subsidised	I Generic
	\$	Per	•	Manufacturer
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	✓	DBL Gemcitabine
lnj 1 g		1	✓	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg	•	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	71.44	1	✓	Irinotecan
				Accord S29
			✓	Irinotecan Actavis
				100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.75	1 mg	•	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist	37.00	25	✓	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist	_			
Special Authority see SA1725 below	428.00	100 ml	OP 🗸	Allmercap

# **⇒SA1725** Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

**Renewal** only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

ME	THOTREXATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	<ul><li>Methotrexate Sandoz</li></ul>
*	Inj 10 mg prefilled syringe14.66	1	<ul><li>Methotrexate Sandoz</li></ul>
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate  Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate  Sandoz
*	Inj 25 mg prefilled syringe	1	<ul><li>Methotrexate Sandoz</li></ul>
*	Inj 30 mg prefilled syringe	1	<ul><li>Methotrexate Sandoz</li></ul>
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter
PE	METREXED - PCT only - Specialist - Special Authority see SA1679 on the	next page	
	Inj 100 mg vial	1	✓ Juno Pemetrexed
	Inj 500 mg vial217.77	1	✓ Juno Pemetrexed
	Inj 1 mg for ECP	1 mg	✓ Baxter

Subsidy	Fu	ly Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

### ⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient has chemotherapy-naïve disease; and
    - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
  - 2.2 All of the following:
    - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
    - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
    - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2. The treatment remains appropriate and the patient is benefitting from treatment; and

<ul> <li>I he treatment remains appropriate and the patient is benefittin</li> <li>Pemetrexed is to be administered at a dose of 500mg/m² every</li> </ul>		; and	
THIOGUANINE - PCT - Retail pharmacy-Specialist Tab 40 mg	.126.31	25	✓ Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule1	,500.00	6	✓ Amsidine S29
4	,736.00		✓ Amsidine S29
Inj 75 mg1	,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Special			•
Cap 0.5 mg1		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial4	,817.00	10	✓ Phenasen
Inj 10 mg for ECP		mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	.161.01	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45 1	,000 iu	✓ Baxter

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	I Generic
BORTEZOMIB – PCT only – Specialist – Special Authority see S Inj 3.5 mg vial		1		Bortezomib
Inj 1 mg for ECP		1 mg		Dr-Reddy's Baxter

### **⇒SA1889** Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis \*.

Note: Indications marked with \* are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	✓ DBL Dacarbazine
, •	580.60	10	✓ Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		3 -	
Inj 2 mg per ml, 10 ml	140.50	1	✓ Pfizer
Inj 20 mg for ECP		20 mg OP	✓ Baxter
	143.50	20 mg Oi	Daxiei
DOCETAXEL – PCT only – Specialist	40.75	4	( December of Occupies
Inj 20 mg		1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
			Accord S29
Inj 80 mg		1	Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	Doxorubicin Ebewe
	17.00		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	Arrow-Doxorubicin
	69.99		<ul> <li>Doxorubicin Ebewe</li> </ul>
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial	30.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial	99.99	1	✓ Epirubicin Ebewe
Inj 1 mg for ECP	0.50	1 mg	✓ Baxter
ETOPOSIDE			
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	✓ Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓ Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialis		1	✓ Rex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
, , , , , , , , , , , , , , , , , , , ,		3	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
ETOPOSIDE PHOSPHATE — PCT only — Specialist Inj 100 mg (of etoposide base)		1 1 mg	_	Etopophos Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phat Cap 500 mgIDARUBICIN HYDROCHLORIDE	, ,	100	•	<u>Devatis</u>
Inj 5 mg vial – PCT only – Specialist Inj 10 mg vial – PCT only – Specialist Inj 1 mg for ECP – PCT only – Specialist	198.00	1 1 1 mg	1	Zavedos Zavedos Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authorit Wastage claimable	•	28		Revlimid
Cap 5 mg Cap 10 mg	4,655.25 6,207.00	21 28	•	Revlimid Revlimid
Cap 15 mg  Cap 25 mg	7,239.18	21 28 21	•	Revlimid Revlimid Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

**Initial application — (Maintenance following first-line autologous stem cell transplant (SCT))** only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

### MESNA

0		
Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 20 mg vial3,275.00	1	✓ Omegapharm S29
		✓ Teva
Inj 1 mg for ECP470.75	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1883 below		
Tab 100 mg3,701.00	56	✓ Lynparza
Tab 150 mg3,701.00	56	✓ Lynparza

### **⇒SA1883** Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib: and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

	Subsidy		Fully	Brand or
	Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
PACLITAXEL - PCT only - Specialist				
Inj 30 mg	47.30	5	1	Paclitaxel Ebewe
Inj 100 mg		1	1	Paclitaxel Ebewe
	91.67		1	Paclitaxel Actavis
Inj 150 mg	26.69	1	1	Paclitaxel Ebewe
	137.50		1	Anzatax
			1	Paclitaxel Actavis
Inj 300 mg	44.00	1	1	Paclitaxel Ebewe
	275.00		1	Anzatax
			1	Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	<b>/</b>	Baxter
EGASPARGASE - PCT only - Special Authority see SA1979 be	low	_		
Inj 750 iu per ml, 5 ml vial		1	1	Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

**Initial application** — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - S	pecialist		
Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pha	armacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 on the	e next page – Retail phar	macy	
Cap 5 mg	9.13	5	✓ <u>Temaccord</u>
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ <u>Temaccord</u>
	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ <u>Temaccord</u>
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ <u>Temaccord</u>
	688.00		✓ Amneal S29

Subsidy Fully (Manufacturer's Price) Per

Subsidised

Brand or Generic Manufacturer

### ⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m<sup>2</sup> per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m2 per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has glioblastoma multiforme; and
  - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
  - 2.1 Patient has anaplastic astrocytoma\*; and
  - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
  - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Authority see SA1124 on the next page

✓ Thalomid	28	378.00	 · 	Cap 50 mg
✓ Thalomid	28	756.00	 	Cap 100 mg

Subsidy		Fully	Brand or
(Manufacturer's Pri	ice)	Subsidised	Generic
<u> </u>	Per	1	Manufacturer

### ⇒SA1124 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an unapproved indication.

#### **TRETINOIN**

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	y see SA1868 belo	W	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		14 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

### ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

	Subsidy		Fully	Brand or
	(Manufacturer's Price	) Si	ubsidised	
	\$	Per	✓	
/INBLASTINE SULPHATE				
Inj 1 mg per ml, 10 ml vial - PCT - Retail pharmacy-Speciali	st270.37	5	1	DBL Vinblastine S29
			1	Hospira
Inj 1 mg for ECP - PCT only - Specialist	6.00	1 mg	1	Baxter
DBL Vinblastine S29 Inj 1 mg per ml, 10 ml vial to be delisted 1	November 2021)	•		
/INCRISTINE SULPHATE				
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialis	t74.52	5	•	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialis	t 102.73	5	•	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist	12.60	1 mg	1	Baxter
/INORELBINE - PCT only - Specialist		ŭ		
Inj 10 mg per ml, 1 ml vial	12.00	1	1	Navelbine
.,	42.00	•	/	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	/	Navelbine
, 01	210.00		1	Vinorelbine Ebewe
	328.65		1	Sagent S29
Inj 1 mg for ECP	1.25	1 mg		Baxter
Inj 50 mg for ECP		i0 mg OF	•	Baxter (Sagent)

# Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below		
Wastage claimable		
Cap 150 mg	224	Alecensa

#### ⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable	•		
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	.7,692.58	60	✓ Sprycel

### ⇒SA1805 Special Authority for Subsidy

**Initial application** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 <b>`</b> \$	Per	✓	Manufacturer

continued...

Any of the following:

- 1 Both:
  - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
  - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
  - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
  - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Maximum dose of 100 mg/day; and
  - 3.3 Any of the following:
    - 3.3.1 Patient has documented treatment failure\* with imatinib; or
    - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
    - 3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

**Renewal** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority see \$A2000 below

Tab 100 mg	764.00	30	Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

### ⇒SA2000 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither
  - 3.1 Patient is treatment naive; or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB - Retail pharmacy-Specialist - Special Authority see SA2001 below

Tab 250 mg .......1,700.00 30 ✓ Iressa

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
  - 2.1 Patient is treatment naive: or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

#### **IMATINIB MESILATE**

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460			
	below2,	400.00	60	✓ Glivec
*	Cap 100 mg	58.23	60	✓ Imatinib-Rex
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex

### ⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

#### Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST)
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

### ⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
NILOTINIB - Special Authority see SA1489 below - Retail pharm	acy				
Wastage claimable					
Cap 150 mg	4,680.00	120	<b>√</b> T	asigna	
Cap 200 mg	6,532.00	120	<b>✓</b> T	asigna	

### **⇒SA1489** Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
  - 2.1 Patient has documented CML treatment failure\* with imatinib: or
  - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

# PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable			
Cap 75 mg	4,000.00	21	Ibrance
Cap 100 mg	4,000.00	21	Ibrance
Cap 125 mg	4,000.00	21	✓ Ibrance

### ⇒SA1894 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
  - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
  - 4.2.2.2 All of the following:
    - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
    - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
    - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

#### continued...

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

#### PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

### ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive: or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

# RUXOLITINIB – Special Authority see SA1890 on the next page – Retail pharmacy

wasiaye dalmable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	·	56	✓ Jakavi
<u> </u>			

Sub	osidy Fully	Brand or
(Manufactu	. '	Generic
	\$ Per ✓	Manufacturer

### **⇒SA1890** Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:
    - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
    - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2002 below - Retail pharmacy

Cap 12.5 mg2,315.38	28	✓ Sutent
Cap 25 mg4,630.77	28	✓ Sutent
Cap 50 mg9,261.54	28	✓ Sutent

### ⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
    - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		sidised	Generic
 \$	Per	/	Manufacturer

continued...

- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Fither:
  - 2.1 The patient's disease has progressed following treatment with imatinib: or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

# **Endocrine Therapy**

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has prostate cancer; and

Subs (Manufactur		
\$	Per	Manufacturer

continued...

- 2 Patient has metastases: and
- 3 Patient's disease is castration resistant; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
    - 4.1.3 Patient has ECOG performance score of 0-1; and
    - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
  - 4.2 All of the following:
    - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
    - 4.2.2 Patient has ECOG performance score of 0-2; and
    - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

#### **BICALUTAMIDE**

Tab 50 mg	1.36	10	✓ Calutide-50 S29
· ·	4.21	28	✓ Binarex
(Calutide-50 S29	Tab 50 mg to be delisted 1 January 2022)		
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT -	Retail pharmacy-Specialist - Special Authority see SA1895 below	٧	
	ml, 5 ml prefilled syringe1,068.00	2	✓ Faslodex

#### ⇒SA1895 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy		Fully	
(	Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
MEGESTROL ACETATE - Subsidy by endorsement	*			
Subsidy by endorsement – Subsidised for patients who were to prescription is endorsed accordingly. Pharmacists may annote prior dispensing of megestrol acetate.				
Tab 160 mg	63.53	30	/	Apo-Megestrol
OCTREOTIDE		00		Tipo mogocilo:
Inj 100 mcg per ml, 1 ml ampoule	18 69	5	/	Octreotide GH \$29
Inj 50 mcg per ml, 1 ml ampoule		5		Octreotide GH \$29
Inj 50 mcg per ml, 1 ml vial		5		Octreotide
, ooog po,		ŭ		MaxRx S29
	56.87		/	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	1	DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule		5	1	Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	1	DBL Octreotide
	222.00		✓	Octreotide
				(Sun) \$29
OCTREOTIDE LONG-ACTING – Special Authority see SA2072 be	elow – Retail pharm	асу		
Inj depot 10 mg prefilled syringe	439.97	1	/	Octreotide Depot Teva
	1,772.50		1	Sandostatin LAR
Inj depot 20 mg prefilled syringe	647.03	1	/	Octreotide Depot Teva
	2,358.75		✓	Sandostatin LAR
Inj depot 30 mg prefilled syringe	718.55	1	•	Octreotide Depot Teva
	2,951.25		1	Sandostatin LAR
Sandostatin LAR Inj depot 10 mg prefilled syringe to be delisted 1 Sandostatin LAR Inj depot 20 mg prefilled syringe to be delisted 1 Sandostatin LAR Inj depot 30 mg prefilled syringe to be delisted 1	March 2022)			

⇒SA2072 Special Authority for Subsidy

**Initial application — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist

(Ma	Subsidy nufacturer's Price)	Sub	Fully	Brand or Generic
<u> </u>	\$	Per	✓	Manufacturer

continued...

has failed: or

2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma: and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas: and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

#### TAMOXIFEN CITRATE

* Tab 20 mg	60	✓ Tamoxifen Sandoz
Aromatase Inhibitors		
ANASTROZOLE		

*	Tab 1 mg	4.55	30	✓ Anatrole
ΕX	EMESTANE			
*	Tab 25 mg	.14.50	30	✓ Pfizer Exemestane

60

✓ Tamoxifen Sandoz

165 ml OP

	Subsidy (Manufacturer's Price)	Subsi Per	Fully dised	Brand or Generic Manufacturer
LETROZOLE  * Tab 2.5 mg	<u> </u>	30		etrole
Immunosuppressants				
Cytotoxic Immunosuppressants				
AZATHIOPRINE  * Tab 25 mg  * Tab 50 mg  * Inj 50 mg vial	7.60	60 100 1	✓ A	zamun zamun nuran
MYCOPHENOLATE MOFETIL  Tab 500 mg  Cap 250 mg	35.90 35.90	50 100		ellcept ellcept

Powder for oral liq 1 g per 5 ml - Subsidy by endorsement........... 187.25 ✓ Cellcept Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

### **Fusion Proteins**

ETANERCEPT - Special Authority see SA2048 below -	- Retail pharmacy		
Inj 25 mg	690.00	4	<ul><li>Enbrel</li></ul>
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Fither:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg. non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	<b>√</b>	Manufacturer

continued...

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application** — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less;

Subsidy	Full	/ Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

and

- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:
    - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Fither:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application** — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsid	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Fither:
      - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 2.2 Both:
    - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 2.2.2 Fither:
      - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's F	rice)	Subsidised	Generic	
\$	Pe	er 🗸	Manufacturer	

continued...

- 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

### **Immune Modulators**

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specia	alist		
Inj 50 mg per ml, 5 ml	2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only	- Specialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial	176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG \$29 Ini 40 mg per ml. vial to be delisted 1 Apri	1 2022)		

#### Monoclonal Antibodies

		below – Retail pharmacy	ADALIMUMAB – Special Authority see SA2049 b
<ul><li>Humira</li></ul>	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
<ul><li>Humira</li></ul>	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

### ⇒SA2049 Special Authority for Subsidy

**Initial application — (adult-onset Still's disease)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both
  - 1.1 Either:
    - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD): or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	<b>✓</b>	Manufacturer	

continued...

- 1.2 Either:
  - 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992:19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Fither
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs. CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
    - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
  - 2 Both:
    - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
    - 2.2 Any of the following:
      - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
      - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
      - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
    - 2.1.2 CDAI score is 150 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- of the following 1 Either:
  - 1.1 Applicant is a gastroenterologist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab: or
    - 2.1.2 PCDAI score is 15 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (fistulising Crohn's disease)** only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

**Initial application — (hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage III or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA: or
- 2 All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
  - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
  - 2.5 Any of the following:
    - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

(A)	Subsidy fanufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
	\$	Per	<b>✓</b>	Manufacturer

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Fither:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
  - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

**Initial application** — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
  - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a

 $dermatologist. \ \ Approvals\ valid\ for\ 6\ months\ for\ applications\ meeting\ the\ following\ criteria:$ 

All of the following:

- 1 Either:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Either:
      - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

- 2.2 Both:
  - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2.2 Either:
    - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
    - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
  - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

## ⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy: or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	<ul><li>Erbitux</li></ul>
Inj 1 mg for ECP	3.82	1 mg	<ul><li>Baxter</li></ul>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ⇒SA1697 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAR	- PCT only -	- Special Authority	see SA2050 below
	I O I OIIIY	Opedial Additionity	JOE ONE DOLON

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

### ⇒SA2050 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

**Initial application — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
  - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
  - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
  - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
  - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
  - 2.2 There has been a marked reduction in prednisone dose; and
  - 2.3 Fither:
    - 2.3.1 There has been an improvement in MRI appearances: or
    - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Both:
    - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

Subsidy	Fully	Brand or
(Manufacturer's		Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
  - 1.2.2 Either:
    - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

### Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis: or
  - 2.2 Ankylosing spondylitis: or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation: or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults); or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease: or
  - 2.9 Severe fulminant ulcerative colitis: or
  - 2.10 Severe ulcerative colitis: or
  - 2.11 Plague psoriasis; or
  - 2.12 Neurosarcoidosis: or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
  - 2 Fither:
    - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
    - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application — (severe Behcet's disease)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
  - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

**Renewal — (ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
  - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - Re	etail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Ini 100 mg vial	1.638.00	1	✓ Nucala

### ⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10<sup>9</sup> cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
  - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	Baxter

## ⇒SA1627 Special Authority for Subsidy

**Initial application — (chronic lymphocytic leukaemia)** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

	-p			
Inj 150 mg p	refilled syringe	450.00	1	Xolair
lni 150 ma v	rial	450.00	1	✓ Xolair

### ⇒SA1744 Special Authority for Subsidy

**Initial application** — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	•	Manufacturer	

continued...

or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months. unless contraindicated or not tolerated: and

- 6 Either:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
  - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks: or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
  - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
  - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal** — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

### Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Roth:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	<b>✓</b>	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	<b>✓</b>	Baxter

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

## **⇒SA1606** Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
  - 2 Either:
    - 2.1 Patient is chemotherapy treatment naïve; or
    - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
  - 3 The patient has good performance status (ECOG grade 0-1); and
  - 4 Pertuzumab to be administered in combination with trastuzumab; and
  - 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
  - 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

## RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

Inj 100 mg per 10 ml vial	1,075.50	2	<ul><li>Mabthera</li></ul>
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

### **⇒SA1976** Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 7 Either:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2061 below

inj 100 mg per 10 ml vial	2/5.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Ini 1 mg for ECP	1.38	1 ma	✓ Baxter (Riximvo)

## ⇒SA2061 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

**Initial application — (Antibody-mediated organ transplant rejection)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

**Initial application — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Either:
    - 2.2.1 The patient is chemotherapy treatment naive: or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Either:
  - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
  - 1.2 All of the following:
    - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
    - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
    - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
    - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
=	<u> </u>				

continued...

375 mg/m2 administered weekly for four weeks; and

- 2 Either
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD: and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

**Initial application** — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
  - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
  - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal** — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

**Renewal — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

**Initial application — (immune thrombocytopenic purpura (ITP))** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
  - 2.1 Treatment with steroids and splenectomy have been ineffective; or
  - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
  - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Subsidy	Fı	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy: and
  - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy:
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms: and

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m<sup>2</sup> weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of  $2 \times 1,000$ mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m<sup>2</sup> of body surface area per week for a total of

(Man	Subsidy	Fully		Brand or
	ufacturer's Price)	Subsidised		Generic
(wich	. ,	Per	<b>✓</b>	Manufacturer

continued...

4 weeks

**Initial application** — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

Subsidy (Manufacturer's Price) Subsi \$ Per	Fully idised	Brand or Generic Manufacturer	
---	-----------------	-------------------------------------	--

continued...

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*: and
- 2 Either:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

#### Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB – Special Authority see SA2044 below – Retail pharmacy

### **⇒SA2044** Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

**Initial application** — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
<b>\$</b>	Per	1	Manufacturer	

continued...

- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal** — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy	Fully	Brand or
acturer's Price)	Subsidised	Generic
 \$ Per	•	

continued...

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

# SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

## ⇒SA1596 Special Authority for Subsidy

**Initial application** only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

### TOCILIZUMAB - PCT only - Special Authority see SA1977 below

Inj 20 mg per ml, 4 ml vial	220.00	1	Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓ Actemra
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

## ⇒SA1977 Special Authority for Subsidy

**Initial application — (cytokine release syndrome)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
  - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
  - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
  - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
  - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
  - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis: or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 systemic juvenile idiopathic arthritis; or
- 2.3 adult-onset Still's disease; or
- 2.4 polyarticular juvenile idiopathic arthritis; or
- 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fithor
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
  - 3.1 Treatment with methotrexate is contraindicated; or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

#### 1 Both:

- 1.1 Either:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and

#### 1.2 Fither:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

### 2 All of the following:

- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

### 1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

### 2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.4 Any of the following:
  - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
  - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	ce)	Fully	Brand or
(Manufacturer's Pric		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	 	1,350.00	1	<ul> <li>Herceptin</li> </ul>
Inj 440 mg vial	 	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	 	9.36	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 2.2 Both:
    - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned: or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
  - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 3.2 Both:
    - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 3.2.2 The cancer did not progress whilst on lapatinib: or
  - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
  - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 4.2 All of the following:
    - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and

Subsidy	Full		
 (Manufacturer's Price) \$	Subsidise Per 🗸	d Generic Manufacturer	

continued...

- 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
- 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial2.320.00	1	✓ Kadcyla
Inj 160 mg vial	1	✓ Kadcyla
Inj 1 mg for ECP23.20	1 mg	✓ Baxter

### ⇒SA1871 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine;
- 2 Treatment to be discontinued at disease progression.

Note: \*Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

# Programmed Cell Death-1 (PD-1) Inhibitors

		NIVOLUMAB – PCT only – Specialist – Special Authority see SA2006 below
Opdivo	1	Inj 10 mg per ml, 4 ml vial1,051.98
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96
✓ Baxter	1 mg	Inj 1 mg for ECP27.62

### ⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
  - 5 Baseline measurement of overall tumour burden is documented (see Note); and
  - 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Fither:
    - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan). following the most recent treatment period: or
    - 1.2.2 Both:
      - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
      - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA. et al. Eur J Cancer 2009:45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
 \$	Per	1	Manufacturer

continued...

Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive

PEMBROLIZUMAB - PCT only - Specialist - Special Authority see SA2007 below ✓ Keytruda 1 mg Baxter

### ⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Fither:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and

1.2 Either:

- 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
- 1.2.2 Both:
  - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
  - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression: and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm.</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

## Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	<ul><li>Neoral</li></ul>
Cap 50 mg	88.91	50	<ul><li>Neoral</li></ul>
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - F	Retail pharmacy		
Wastage claimable			
Tab 10 mg	6,512.29	30	<ul><li>Afinitor</li></ul>
Tab 5 mg	4,555.76	30	Afinitor

### ⇒SA2008 Special Authority for Subsidy

**Initial application** only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg	100	Rapamune
Tab 2 mg	100	✓ Rapamune
Oral lig 1 mg per ml	60 ml OP	✓ Rapamune

### ⇒SA2005 Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or</li>
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis: or
- . HUS or TTP; or
- Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease: and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

 Subsidy Manufacturer's Price)	٤	Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:
    - 2.2.1 Vigabatrin is contraindicated; and
    - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

### ⇒SA1745 Special Authority for Subsidy

**Initial application — (organ transplant)** only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
  - 1 Patient requires long-term systemic immunosuppression; and
  - 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with \* are unapproved indications

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Antiallergy Preparations**

## Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

### ⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

## **Allergy Desensitisation**

### ⇒SA1367 Special Authority for Subsidy

**Initial application** only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1	367 above – Ret	ail pharmacy	
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml		1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent .	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see S	A1367 above – R	etail pharma	су
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			•
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29

	Subsidy		Fully	Brand or
	(Manufacturer's Price		idised	Generic
	\$	Per		Manufacturer
Authistonius				
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1 12	100	1	Zista
* Oral liq 1 mg per ml		200 ml		Histaclear
CHLORPHENIRAMINE MALEATE				
* Oral liq 2 mg per 5 ml	0.27	500 ml	./	Histafen
	9.37	500 1111	•	пізіаівіі
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg	4	40		
	(8.40)			Polaramine
	1.01	20		<b>-</b>
4. 0 . 11. 0	(5.99)			Polaramine
* Oral liq 2 mg per 5 ml		100 ml		<b>D</b>
	(10.29)			Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
	(8.23)			Telfast
* Tab 120 mg	4.74	10		
	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
LORATADINE				
* Tab 10 mg	1.69	100	1	Lorafix
* Oral liq 1 mg per ml	1.43	100 ml	1	Haylor syrup
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg	1.68	50	1	Allersoothe
* Tab 25 mg		50	1	Allersoothe
* Oral lig 1 mg per 1 ml		100 ml	1	Allersoothe
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a		5		Hospira
, 01 , 1 , 1				•
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE			_	_
Aerosol inhaler, 50 mcg per dose		200 dose OP		Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP		Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	200 dose OP	•	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00 2	200 dose OP	1	Pulmicort
				Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	200 dose OP	1	Pulmicort
				Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	200 dose OP	1	Pulmicort
<del>-</del> ·				Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's	Price) Subs	idised Generic
	\$	Per	✓ Manufacturer
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	sts		
EFORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose dev	rice20.64	60 dose	
· · · · · · · · · · · · · · · · · · ·	(35.80)		Foradil
EFORMOTEROL FUMARATE DIHYDRATE	(55.55)		
Powder for inhalation 4.5 mcg per dose, breath activated	-) 10.00	CO dana OD	
(equivalent to eformoterol fumarate 6 mcg metered dos	,	60 dose OP	Orio Trutubalas
	(16.90)		Oxis Turbuhaler
INDACATEROL			
Powder for inhalation 150 mcg		30 dose OP	<ul><li>Onbrez Breezhaler</li></ul>
Powder for inhalation 300 mcg	61.00	30 dose OP	Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	✓ Serevent Accuhaler
habitated Analysis at any life with Lorent Anthon But-	A .l		
Inhaled Corticosteroids with Long-Acting Beta-	-Aarenocept	or Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide v	with		
6 mcg eformoterol fumarate metered dose)	41.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma	rate		
per dose (equivalent to 400 mcg budesonide with 12 mc			
eformoterol fumarate metered dose) - No more than 2	J		
dose per day	82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	18.23	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6	mcg33.74	120 dose OP	✓ Symbicort
ř	-		Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40	120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6		120 dose OP	✓ Symbicort
ř	-		Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	44.08	60 dose OP	✓ Symbicort
, ,			Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	<i>44</i> 08	30 dose OP	✓ Breo Ellipta
1 5 macri for initialiation 100 mag with vitalitorol 20 mag		50 dosc OI	- Dico Empla

	Subsidy		Fully Brand or
	(Manufacturer's		
	\$	Per	✓ Manufacturer
FLUTICASONE WITH SALMETEROL			
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP	✓ <u>Seretide</u>
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No	)		
more than 2 dose per day	33.74	60 dose OP	<ul> <li>Seretide Accuhaler</li> </ul>
Powder for inhalation 250 mcg with salmeterol 50 mcg - No			
more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler
Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Oral lig 400 mcg per ml	20.00	150 ml	✓ Ventolin
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO	53.00	5	✓ Ventolin
injour mag per mi, 1 mi op to e injuvanable en a 1 ee			- 7011.01111
Inhaled Beta-Adrenoceptor Agonists			
SALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000			
dose available on a PSO	3.80	200 dose OP	✓ Respigen
		200 0000 0.	✓ SalAir
	(6.00)		Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule - Up to 30 neb			VOITOIIII
available on a PSO		20	✓ Asthalin
		20	▼ ASuldilli
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb		00	/ A adh alin
available on a PSO	9.43	20	✓ Asthalin
FERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Bricanyl Turbuhaler
-			•
Anticholinergic Agents			
PRATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	✓ Atrovent
a) Up to 400 dose available on a PSO			
b) No patient co-payment payable			
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne	eh		
available on a PSO		20	✓ Univent
Inhaled Beta-Adrenoceptor Agonists with Antic	holineraic /	\aente	<del></del>
minute beta Autonocopior Agomata with Antic	indinici gic F	igenio	
SALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg			
dose CFC-free	12.19	200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per			
vial, 2.5 ml ampoule – Up to 20 neb available on a PSC	11 04	20	✓ Duolin
viai, 2.5 mi ampoulo op to 20 nob available on a 1 00		20	- Duoini

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per Manufacturer

## **Long-Acting Muscarinic Antagonists**

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

## Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

**Renewal** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL — Special Authority see SA1584 above — Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg.......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg ......81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

## **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

### **⇒SA2012** Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	<ul><li>Esbriet</li></ul>
Tab 267 mg	1,215.00	90	<ul><li>Esbriet</li></ul>
Cap 267 mg - Wastage claimable		270	✓ Esbriet

(Esbriet Cap 267 mg to be delisted 1 January 2022)

#### ⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Manuacturers Frice)	Per	Jubsiuiseu ✓	Manufacturer	

# Leukotriene Receptor Antagonists

МО	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
*	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg3.95	28	✓ Montelukast Mylan

## **Mast Cell Stabilisers**

SODIUM CROMOGLICATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking sodium cromoglicate prior to 1 July 2020 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of sodium cromoglicate.

✓ Intal Forte CFC Free

# Methylxanthines

#### **AMINOPHYLLINE**

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	180.00	5	✓ DBL Aminophylline
THI	EOPHYLLINE			
*	Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
*	Oral liq 80 mg per 15 ml	16.60	500 ml	✓ <u>Nuelin</u>

# **Mucolytics**

DORNASE ALFA - Special Authority see SA1978 below - F	Retail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

### ⇒SA1978 Special Authority for Subsidy

**Initial application — (cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 on the next page

Tab 150 mg29,386	5.00 56	✓ Kalydeco
Oral granules 50 mg, sachet		✓ Kalydeco
Oral granules 75 mg, sachet	5.00 56	✓ Kalydeco

Subsidy (Manufacturer's Pr	Subsidy Fully (Manufacturer's Price) Subsidised		Brand or Generic	
\$	Per	✓	Manufacturer	

## ⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system: and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

#### SODIUM CHLORIDE

Not funded for use as a nasal drop.

✓ Biomed 90 ml OP

# Nasal Preparations

## Allergy Prophylactics

BUDESONIDE		
Metered aqueous nasal spray, 50 mcg per dose2.54	200 dose OP	✓ SteroClear
Metered aqueous nasal spray, 100 mcg per dose2.84	200 dose OP	✓ SteroClear
FLUTICASONE PROPIONATE		
Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ Flixonase Hayfever
		& Allergy
Flixonase Hayfever & Allergy to be Principal Supply on 1 December 20	21	
100 170 000 110 000 110 0		

**IPRATROPIUM BROMIDE** 

Aqueous nasal spray, 0.03%......5.23 15 ml OP ✓ Univent

# **Respiratory Devices**

### MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

Small 2.20 ✓ e-chamber Mask

## PEAK FLOW METER

- a) Up to 25 dev available on a PSO
- b) Only on a PSO

✓ Mini-Wright AFS Low Range

Normal range 9.54 ✓ Mini-Wright Standard

25 ml OP

✓ Biomed

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSO     b) Only on a PSO				
220 ml (single patient)	2.95	1	✓	e-chamber Turbo
510 ml (single patient)	5.12	1	•	e-chamber La Grande
800 ml	6.50	1	•	Volumatic
Respiratory Stimulants				
CAFFEINE CITRATE				

Oral liq 20 mg per ml (10 mg base per ml)......15.10

	(Manufacturer's F	Price) Subs	idised Generic
	\$	Per	✓ Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND B	ENZETHONIUM		
For Vosol ear drops with hydrocortisone powder refer Stand	lard Formulae, pa	age 242	
Ear drops 2% with 1, 2-Propanediol diacetate 3% and benzethonium chloride 0.02%	6 97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE	0.07	00 1111 01	70301
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYO	OIN AND NYSTAT	ΓIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5 16	7.5 ml OP	✓ Kenacomb
		7.01111 01	· Kenadomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml		8 ml OP	0.4
EDAM/OFTIN OUR BUATE	(9.27)		Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4 13	8 ml OP	
Eui/Eyo diopo 0.0/0	(8.65)	01111 01	Soframycin
			•
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expl	icitly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL			4
Eye oint 1% Eye drops 0.5%		5 g OP 10 ml OP	✓ <u>Devatis</u> ✓ Chlorafast
Funded for use in the ear*. Indications marked with * a			Cilioralast
CIPROFLOXACIN	• • •		
Eye drops 0.3% - Subsidy by endorsement		5 ml OP	<ul> <li>Ciprofloxacin Teva</li> </ul>
a) When prescribed for the treatment of bacterial kerat		•	
or for the second line treatment of chronic suppurati accordingly. Note: Indication marked with a * is an			e prescription is endorsed
b) Ciprofloxacin Teva to be Principal Supply on 1 Nove		J. J	
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE	2.27	40 1 0.0	
* Eye drops 0.1%	2.97 (14.55)	10 ml OP	Brolene
SODIUM FUSIDATE (FUSIDIC ACID)	(14.00)		DIOIOIIC
Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
		-	

Subsidy

Fully

Brand or

					_
	Subsidy		Fully	Brand or	
	(Manufacturer's F	Price) Sub	sidised	Generic	
	\$	Per	•	Manufacturer	
TOBRAMYCIN					
Eye oint 0.3%	10.45	3.5 g OP	✓ T	obrex	
Eye drops 0.3%		5 ml OP	<b>✓</b> T	obrex	
Corticosteroids and Other Anti-Inflammatory Pr	reparations				
DEXAMETHASONE					
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	laxidex	
* Eye drops 0.1%	4.50	5 ml OP	✓ N	laxidex	
Ocular implant 700 mcg - Special Authority see SA1680 bel	low				
- Retail pharmacy		1	<b>✓</b> 0	zurdex	

### ⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not vet completed a family: and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin			
	b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
	Voltaren Ophtha to be Principal Supply on 1 November 2021			•

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's P	rice) Sub:	Fully	Brand or Generic
	\$	Per	1	Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	<b>√</b> F	ML
	5.20		<b>√</b> F	lucon
KETOROLAC TROMETAMOL - Special Authority see SA1981	below – Retail ph	armacy		
Eye drops 0.5%	9.50	5 ml OP	<b>√</b> A	Acular
OA4004 On a dal Anathanita fan Ontada				

### ⇒SA1981 Special Authority for Subsidy

**Initial application — (macular oedema)** only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
  - 2.1 The patient is at risk of postoperative macular oedema; and
  - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

#### LEVOCABASTINE

Eye drops 0.5 mg per ml	8.71 (10.34)	4 ml OP	Livostin
LODOXAMIDE			•
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC	10.00	0 1 0 0	<b>/</b> II
Eye drops 0.3%	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE			
Eye drops 1%	5.93	10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	– Retail pharm	nacy
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	<ul><li>Minims</li><li>Prednisolone</li></ul>

## ⇒SA1715 Special Authority for Subsidy

**Initial application** only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

## SODIUM CROMOGLICATE

Eye drops 2%	✓ Rexacrom
--------------	------------

# Glaucoma Preparations - Beta Blockers

BETAXOLOL			
* Eye drops 0.25%	11.80	5 ml OP	✓ Betoptic S
* Eye drops 0.5%	7.50	5 ml OP	✓ Betoptic
TIMOLOL			
* Eye drops 0.25%	1.81	5 ml OP	Arrow-Timolol
* Eye drops 0.5%	2.04	5 ml OP	✓ Arrow-Timolol
* Eve drops 0.5% gel forming	3.78	2.5 ml OP	✓ Timontol XF

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or idised Generic  Manufacturer
Glaucoma Preparations - Carbonic Anhydrase	Inhibitors		
ACETAZOLAMIDE  * Tab 250 mg	17.03	100	✓ Diamox
K Eye drops 1%	7.30	5 ml OP	✓ Azopt
Eye drops 2%	9.77 (17.44)	5 ml OP	Trusopt
ORZOLAMIDE WITH TIMOLOL  Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ Dortimopt
Glaucoma Preparations - Prostaglandin Analog	jues		
BIMATOPROST * Eye drops 0.03%	3.30	3 ml OP	✓ Bimatoprost Multichem
ATANOPROST  * Eye drops 0.005%	1.82	2.5 ml OP	✓ Teva
FRAVOPROST  * Eye drops 0.004%	7.30 9.75	5 ml OP 2.5 ml OP	<ul><li>✓ Travopt</li><li>✓ Travatan</li></ul>
Travatan to be Principal Supply on 1 December 2021 Travopt Eye drops 0.004% to be delisted 1 December 2021) Mylan S29 Eye drops 0.004% to be delisted 1 December 2021	10.50	5 ml OP	✓ Mylan S29
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE  * Eye drops 0.2%	4.29	5 ml OP	✓ Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE  Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
ATANOPROST WITH TIMOLOL  Eye drops 0.005% with timolol 0.5%  PILOCARPINE HYDROCHLORIDE	2.49	2.5 ml OP	✓ Arrow - Lattim
Eye drops 1%  Eye drops 2%		15 ml OP 15 ml OP	<ul><li>✓ Isopto Carpine</li><li>✓ Isopto Carpine</li></ul>
Subsidised for oral use pursuant to the Standard Formu		15 ml OP	✓ Isopto Carpine
Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine

1 Patient has to use an unpreserved solution due to an allergy to the preservative; or

Either:



	Subsidy	Fully	Brand or
(Manut	facturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

# **Mydriatics and Cycloplegics**

ATROPINE SULPHATE  * Eye drops 1%	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE  * Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a prescription	20 dose	✓ Minims Cyclopentolate
TROPICAMIDE  * Eye drops 0.5%	15 ml OP 15 ml OP	✓ Mydriacyl ✓ Mydriacyl

## **Preparations for Tear Deficiency**

For acetylcysteine eye drops refer Standard Formulae, page 242

HYPROMELLOSE

* Eye drops 0.5%19.50	15 ml OP	✓ Methopt
HYPROMELLOSE WITH DEXTRAN		
* Eve drops 0.3% with dextran 0.1%	15 ml OP	✓ Poly-Tears

## **Preservative Free Ocular Lubricants**

### ⇒SA1388 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eve drops more than four times daily on a regular basis: or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

aropo and nao ponemou nom noamona			
CARBOMER - Special Authority see SA1388 above - Retail pha	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Author	ity see SA1388 at	ove – Reta	il pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	✓ Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth	ority see SA1388	above – Re	tail pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pha	armacy Procedure	s Manual re	striction allowing one bottle per
month is not relevant and therefore only the prescribed d	losage to the near	est OP may	be claimed.

# **SENSORY ORGANS**

	Subsidy (Manufacturer's Prio \$	ce) Subsi Per	Fully idised	Brand or Generic Manufacturer
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%	4.15	15 ml OP	<b>✓</b> N	laphcon Forte
OLOPATADINE Eye drops 0.1%	2.20	5 ml OP	<b>√</b> <u>C</u>	Diopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT  * Eye oint 3% with wool fat 3%	3.63	3.5 g OP	<b>✓</b> P	Poly-Visc
RETINOL PALMITATE  Eye oint 138 mcg per g	3.80	5 g OP	✓ V	'itA-POS



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# Agents Used in the Treatment of Poisonings

#### Antidotes

ACFTYL	CVCTE	

Pharma \$29

#### NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

### Removal and Elimination

#### CHARCOAL

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

## DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

 Tab 125 mg dispersible
 276.00
 28
 ✓ Exjade

 Tab 250 mg dispersible
 552.00
 28
 ✓ Exjade

 Tab 500 mg dispersible
 1,105.00
 28
 ✓ Exjade

## **⇒SA1492** Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or
  - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
  - 3.3 Treatment with deferiprone has resulted in arthritis; or
  - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

#### Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE – Special Authority see SA1480 on the nex	<mark>rt page –</mark> Retail pharm	acy	
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox



Subsidy (Manufacturer's Price)	Subsid	Fully lised	Brand or Generic
 \$	Per	•	Manufacturer

## ⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

* Inj 500 mg vial	84.53	10	DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE		_	
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate



# Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Acetylcysteine inj 200 mg per ml, 10 ml	qs	mg per ml)	
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
		Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
00DEINE   INOTHIO (45		Water	to 500 ml
CODEINE LINCTUS (15 mg per 5 ml)	000	(Preservative should be used if quantity supplied is	for more
Codeine phosphate	300 mg	than 5 days.)	
Glycerol	40 ml	CALIVA CURCTITUTE FORMULA	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	<b>5</b>
Water	to 100 ml	Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative	qs
Calcium folinate 15 mg tab	1 tab	Water	to 500 ml
Preservative		(Preservative should be used if quantity supplied is	or more
Water	qs to 500 ml	than 5 days. Maximum 500 ml per prescription.)	
(Preservative should be used if quantity supplied is		SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)	ioi iliole	Sodium chloride inj 23.4%, 20 ml	qs
than 5 days. Maximum 500 mi per prescription.)		Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatra	
Methadone powder	qs	(Only failed in procession for a caution of hypothane	ισιτια
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection	10 vials
		Glycerol BP	40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridiu	m difficile
Propylene glycol	to 100 ml	following metronidazole failure)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	id mixture)		
OMEPRAZOLE SUSPENSION		VOSOL EAR DROPS	
		WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder	qs	Hydrocortisone powder	1%
Sodium bicarbonate powder BP	8.4 g	Vosol Ear Drops	to 35 ml
Water	to 100 ml		
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Water	to 100 ml		
Water	100 1111		

## EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or Generic (Manufacturer's Price) Subsidised Per

Manufacturer

Extemporaneously	<b>Compounded Pr</b>	reparations and Galenica	Is
------------------	----------------------	--------------------------	----

CODEINE PHOSPHATE - Safety medicine; prescriber may de	etermine dispensing t	requency	
Powder - Only in combination	63.09	25 g	
·	(90.09)	•	Douglas
Only in automorphism and a solution in the state of the s			

Only in extemporaneously compounded codeine linctus.

#### COLLODION FLEXIBLE

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

✓ PSM 100 ml

## COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

100 ml ✓ Midwest

## GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus.

473 ml **Ora-Sweet SF** 

## GLYCERIN WITH SUCROSE - Only in combination

Only in combination with Ora-Plus.

Suspension......30.95 473 ml ✓ Ora-Sweet

#### GI YCFROI

Powder

500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations.

### METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

7 9/

✓ AET

/ .84	1 g	♥ AFI
0.00	0.5	
8.98	25 g	✓ <u>Midwest</u>
36.95	100 a	✓ MidWest
	473 ml	✓ Ora-Plus
- Only in cor	nbination	
30.95	473 ml	✓ Ora-Blend SF
ombination		
30.95	473 ml	✓ Ora-Blend
52.50	10 g	✓ MidWest
325.00	100 g	✓ MidWest
10% solution.		
11.25	500 ml	✓ Midwest
10.05	500 g	✓ Midwest
	. 3	
	30.95 ombination 30.95 52.50 325.00 10% solution. 11.25	8.98 25 g 36.95 100 g 30.95 473 ml  - Only in combination 30.95 473 ml  ombination 30.95 473 ml 52.50 10 g  325.00 100 g

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

# **EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS**

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	✓ <u>M</u>	idwest
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

# **Nutrient Modules**

## Carbohydrate

### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

# Carbohydrate And Fat

## **⇒SA1376** Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

✓ fully subsidised 245



Subsidy		Fully	Brand or
(Manufacturer's Price)	Su	ıbsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Fat

### ⇒SA1523 Special Authority for Subsidy

**Initial application — (Inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy	5	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

## **Protein**

### ⇒SA1524 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	rmacy [HP3]	PROTEIN SUPPLEMENT – Special Authority see SA1524 above – Hospital pha
✓ Protifar	225 g OP	Powder7.90
✓ Resource	227 g OP	8.95
Beneprotein	•	

✓ fully subsidised 247

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic 
\$ Per 
Manufacturer

## **Oral and Enteral Feeds**

#### Diabetic Products

### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1 Liquid		500 ml OP	acy [HP3]  ✓ Glucerna Select ✓ Diason RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 Liquid (strawberry) Liquid (vanilla)	1.50	spital pharmacy [ 200 ml OP 200 ml OP 237 ml OP	[HP3] ✓ Diasip ✓ Diasip
(Suctagen Diabatic Liquid (vanilla) to be delicted 1 February 2022)	(2.10) (2.10)	200 ml OP	Sustagen Diabetic Nutren Diabetes

(Sustagen Diabetic Liquid (vanilla) to be delisted 1 February 2022)

## **Fat Modified Products**

## ⇒SA1525 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above -	– Hospitai pnarma	Cy [HP3]	
Powder	60.48	400 g OP	<ul><li>Monogen</li></ul>

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

# **Paediatric Products For Children Awaiting Liver Transplant**

## ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

## Paediatric Products For Children With Chronic Renal Failure

# ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

## **Paediatric Products**

## ⇒SA1379 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

✓ fully subsidised 249

Subsidy	Fully	Brand or
(Manufacturer's	Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

•			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see Liquid		the previous page 500 ml OP	ge – Hospital pharmacy [HP3]  ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see S/Liquid		e previous page 500 ml OP	<ul> <li>Hospital pharmacy [HP3]</li> <li>Nutrini RTH</li> <li>Pediasure RTH</li> </ul>
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special pharmacy [HP3]	Authority se	e SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1: Liquid (strawberry) Liquid (vanilla)	1.60	orevious page – 200 ml OP 200 ml OP	Hospital pharmacy [HP3]  Fortini Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA137 Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.07 1.07	evious page – H 200 ml OP 200 ml OP 200 ml OP 250 ml OP	ospital pharmacy [HP3]  Pediasure Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Auth pharmacy [HP3]	ority see SA	A1379 on the pre	evious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.60 1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	<ul> <li>✓ Fortini Multi Fibre</li> <li>✓ Fortini Multi Fibre</li> <li>✓ Fortini Multi Fibre</li> <li>✓ Fortini Multi Fibre</li> </ul>
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		page – Hospital 400 g OP	pharmacy [HP3]  Peptamen Junior

### **Renal Products**

## ⇒SA1101 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML - Special Authority :	see SA1101 above –	Hospital pharma	cy [HP3]
Liquid	6.08	500 ml OP	✓ Nepro HP RTH

	Subsidy (Manufacturer's Pric		Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA11 Liquid		page – Hosp 220 ml OP	✓ N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110 Liquid	2.88	o <mark>age</mark> – Hospit 237 ml OP	·	
Liquid (apricot) 125 ml		4 OP 4 OP	<b>✓</b> R	lovaSource Renal Renilon 7.5 Renilon 7.5

## **Specialised And Elemental Products**

### ⇒SA1377 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

  ENTERAL (ORAL SEMI ELEMENTAL EEED 1.5KCAL(ML). Special Authority see SA1277 above. Hospital pharmacy (HR2)

LiquidLiquid		1,000 ml OP	, , , ,			
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]						
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra			
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]						
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN			
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Authority see SA1377 above - Hospital pharmacy [HP3]						
Liquid	12.04	1,000 ml OP	✓ Peptisorb			

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

## Paediatric Products For Children With Low Energy Requirements

## ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years: and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML  –	<ul> <li>Special Authority</li> </ul>	see SA1196 ab	ove -	<ul> <li>Hospital pharmacy [HP3]</li> </ul>
Liquid	4.00	500 ml OP	1	Nutrini Low Energy
				Multi Fibre

## Standard Supplements

### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- - 1 The patient is under 18 years of age; and
  - 2 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	S	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

**Renewal — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

**Initial application** — **(Long-term medical condition)** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm<sup>3</sup>); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 2	252 – Hos	pital pharmacy	[HP3]
Liquid	'.00 1	,000 ml OP	<ul><li>Nutrison Energy</li></ul>

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic  ✓ Manufacturer
ENTERN FEED 1/CAL/MI Createl Authority and CA10F0 or	Ψ OFO		
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on Liquid		spitai pharmacy 250 ml OP	✓ Isosource Standard
Liquid	5.29	1,000 ml OP	✓ Nutrison Standard
	3.23	1,000 1111 01	RTH
			✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authorit	v see SA1859 o	n nage 252 – H	
Liquid	•	1.000 ml OP	✓ Nutrison
1.		,	800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s	ee SA1859 on p	age 252 – Hosp	oital pharmacy [HP3]
Liquid	5.29	1,000 ml OP	✓ Jevity RTH
			Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority		page 252 - Hos	
Liquid		250 ml OP	✓ Ensure Plus HN
	7.00	1,000 ml OP	✓ Ensure Plus RTH
			✓ Jevity HiCal RTH
			✓ Nutrison Energy Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on page	no 252 Hospita	al pharmacy [HE	
Powder (chocolate)	,	840 g OP	✓ Sustagen Hospital
. 5.55. (6.5554.5)		0.0 g 0.	Formula Active
	26.00	850 g OP	✓ Ensure
Powder (vanilla)	14.00	840 g OP	<ul> <li>Sustagen Hospital</li> </ul>
			Formula Active
	26.00	850 g OP	✓ Ensure
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa			
Additional subsidy by endorsement is available for patients b			
epidermolysis bullosa, or as exclusive enteral nutrition in chil			
disease, or for patients with COPD and hypercapnia, defined endorsed accordingly.	as CO2 value e	exceeding Somm	nng. The prescription must be
Liquid (banana) — Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	5 Di
	(1.26)		Ensure Plus
Liquid (fruit of the forcet) . Higher subside of \$1.00 per 200	(1.26)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 with Endorsement		200 ml OP	
Har Endorsomont	(1.26)	200 1111 01	Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml w			
Endorsement		237 ml OP	5 Di
	(1.33)	000 ml OD	Ensure Plus
	0.72 (1.26)	200 ml OP	Ensure Plus
	(1.26)		Fortisip
	(1.20)		i ordorp

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 252 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Liquid (Chocolate) – Higher Subsidy of \$1.26 per 200 fill with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

## **High Calorie Products**

#### ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 of		•		,
Liquid	5.50	500 ml OP	<b>✓</b> N	Nutrison Concentrated
	11.00	1,000 ml OP	<b>✓</b> E	Ensure Two Cal HN RTH
			<b>√</b> 1	Two Cal HN RTH
(Two Cal HN RTH Liquid to be delisted 1 February 2022)				
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients lepidermolysis bullosa. The prescription must be endorsed a	being bolus fed t		, .	
Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with	0.00	000   OD		
Endorsement	0.96	200 ml OP	7	Two Cal HN

## **Food Thickeners**

#### ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER - Special Authority see SA	1106 above - Hospital pharmacy	[HP3]	
Powder	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	✓ Feed Thickener
		-	Karicare Aptamil

### **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

## ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

	Subsidy		Fully Brand or
	(Manufacturer's Pric \$	e) Subsid	dised Generic  ✓ Manufacturer
GLUTEN FREE BAKING MIX - Special Authority see SA17.	<del>-</del>		
Powder		,000 g OP	namacy [m oj
1 OWGOT	(5.15)	,000 g Oi	Healtheries Simple
	(0.10)		Baking Mix
NUTEN EDEE DDEAD MIV - Or stiel Authority - 0 0476	00 11 1	11 9.1 . 1	ŭ
GLUTEN FREE BREAD MIX – Special Authority see SA172			narmacy [HP3]
Powder		,000 g OP	NZD L Olutara
	(7.32)		NZB Low Gluten Bread Mix
	3.51		Dreau IVIIX
	(10.87)		Horleys Bread Mix
	( /		•
GLUTEN FREE FLOUR - Special Authority see SA1729 on			acy [HP3]
Powder		2,000 g OP	
	(18.10)		Horleys Flour
SLUTEN FREE PASTA - Special Authority see SA1729 on	the previous page - Ho	spital pharma	acy [HP3]
Buckwheat Spirals	2.00	250 g OP	,
•	(3.11)	· ·	Orgran
Corn and Vegetable Shells	2.00 <sup>′</sup>	250 g OP	v
•	(2.92)	•	Orgran
Corn and Vegetable Spirals	2.00	250 g OP	•
	(2.92)		Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP	
	(3.82)		Orgran
Rice and Corn Macaroni	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Penne	2.00	250 g OP	
	(2.92)		Orgran
Rice and Maize Pasta Spirals		250 g OP	
	(2.92)		Orgran
Rice and Millet Spirals	2.00	250 g OP	
	(3.11)		Orgran
Rice and corn spaghetti noodles		375 g OP	
	(2.92)		Orgran
Vegetable and Rice Spirals		250 g OP	_
	(2.92)		Orgran
Italian long style spaghetti		220 g OP	_
	(3.11)		Orgran

# Foods And Supplements For Inborn Errors Of Metabolism

### ⇒SA1108 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

# **Supplements For Homocystinuria**

AMINOACID FORMULA WITHOUT METHIONINE – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

## **Supplements For MSUD**

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

## **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet		30	✓ PKU Ánamix Junior Orange
Powder (chocolate) 36 g sachet	393.00	30	<ul> <li>PKU Anamix Junior Chocolate</li> </ul>
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	PKU Anamix Junior
Powder (vanilla) 36 g sachet		30	<ul><li>PKU Anamix Junior Vanilla</li></ul>
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	<ul> <li>Easiphen Liquid</li> </ul>
Liquid (juicy tropical) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml		30 OP	✓ PKU Lophlex LQ 20

## Foods

LOW PROTEIN BAKING MIX - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Powder ......8.22 500 g OP ✓ Loprofin Mix

## **SPECIAL FOODS**

	Subsidy		Fully	Brand or
	(Manufacturer's Pri	ice) Subsi	dised	Generic
	\$	Per		Manufacturer
LOW PROTEIN PASTA - Special Authority see SA1108 on page	e 258 – Hospital p	harmacy [HP3	3]	
Animal shapes	11.91	500 g OP	<b>✓</b> L	.oprofin
Lasagne	5.95	250 g OP	<b>✓</b> L	.oprofin
Low protein rice pasta	11.91	500 g OP	<b>✓</b> L	.oprofin
Macaroni	5.95	250 g OP	<b>√</b> L	.oprofin
Penne	11.91	500 g OP	<b>√</b> L	.oprofin
Spaghetti	11.91	500 g OP	<b>√</b> L	.oprofin
Spirals	11.91	500 g OP	<b>√</b> L	.oprofin
		-		•

#### Infant Formulae

## For Williams Syndrome

## ⇒SA1110 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

# Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA - Special Authority see SA1940 below - Hospital phar	macy [HP3]	
Powder43.60	400 g OP	<ul><li>✓ Alfamino</li><li>✓ Alfamino Junior</li></ul>
Powder (unflavoured)53.00	400 g OP	✓ Elecare ✓ Elecare LCP ✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	<ul><li>✓ Neocate SYNEO</li><li>✓ Elecare</li><li>✓ Neocate Junior</li><li>✓ Vanilla</li></ul>

## **⇒SA1940** Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or

continued...

Subsidy	Fı	illy Brand or	
(Manufacturer's Pric	e) Subsidis	ed Generic	
\$	Per	✓ Manufacturer	

continued...

- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
  - 6.2 Fither:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
  - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
  - 2.2 All of the following:
    - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
    - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and

continued...

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

S Per 
Manufacturer

continued...

- 2.2.3 Amino acid formula is required for a nutritional deficit; and
- 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA -	<ul> <li>Special Authority see SA1953 below</li> </ul>	- Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

## ⇒SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea: or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption; or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and
    - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

continued...

Subsidy	Fully	Brand or
acturer's Price) Subs	sidised	Generic
\$ Per	1	

continued...

- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

#### ⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Fither:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Sov milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula; and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...



Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	∡DSIGISCG ✓	Manufacturer

continued...

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

#### Fluid Restricted

## ⇒SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

**Renewal** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

## **Ketogenic Diet**

## ⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)35.50	300 g OP	✓ KetoCal 4:1

### **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

# **Vaccinations**

#### BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > 0 requal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00 10 

✓ BCG Vaccine

#### DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
  Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

## DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE			
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	ND HAEMOPHILUS I	NFLUENZAE TY	PE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of			and a considerable and a second
2) An additional four doses (as appropriate) are funded for			
10 who are patients post haematopoietic stem cell trans post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up p		'	
to complete full primary immunisation. Please refer to the Imi			
programmes.			
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10 <b>✓</b> <u>In</u>	<u>ifanrix-hexa</u>
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)im			
transplantation, or chemotherapy; functional asplenic; pr			olid organ transplant, pre-
or post cochlear implants, renal dialysis and other sever	, , , , , , , , , , , , , , , , , , , ,	•	val madiaina physisian ar
<ol> <li>For use in testing for primary immunodeficiency disease paediatrician.</li> </ol>	s, on the recommend	allon of an intern	iai medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg	:		
prefilled syringe plus vial 0.5 ml		1 🗸 H	iberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver di	sease; or		
3) One dose of vaccine for close contacts of known hepatit	is A cases.		

✓ Havrix

✓ Havrix Junior

1

	NATIONAL	IIVIIVI	UNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE - [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the following criteria		1	<b>✓</b> E	ngerix-B
<ol> <li>for household or sexual contacts of known acute legal for children born to mothers who are hepatitis B second for children up to and under the age of 18 years in serology and require additional vaccination or require additional vaccination or require for HIV positive patients; or</li> <li>for hepatitis C positive patients; or</li> <li>for patients following non-consensual sexual inter</li> <li>for patients following immunosuppression; or</li> <li>for solid organ transplant patients; or</li> <li>for post-haematopoietic stem cell transplant (HSC)</li> <li>following needle stick injury.</li> </ol>	hepatitis B patients or h urface antigen (HBsAg nclusive who are consicuire a primary course of rcourse; or	, posi dered	tive; or not to have	
Inj 20 mcg per 1 ml prefilled syringe  Funded for patients meeting any of the following criteria  1) for household or sexual contacts of known acutel  2) for children born to mothers who are hepatitis B s  3) for children up to and under the age of 18 years in serology and require additional vaccination or req  4) for HIV positive patients; or  5) for hepatitis C positive patients; or  6) for patients following non-consensual sexual inter  7) for patients following immunosuppression; or  8) for solid organ transplant patients; or  9) for post-haematopoietic stem cell transplant (HSC)  10) following needle stick injury; or  11) for dialysis patients; or  2) for liver or kidney transplant patients.	a: hepatitis B patients or h urface antigen (HBsAg nclusive who are consicuire a primary course of	, posi dered	tis B carriers tive; or not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND Any of the following:  1) Maximum of two doses for children aged 14 years and 2) Maximum of three doses for patients meeting any of the second	d under; or he following criteria: or			
Inj 270 mcg in 0.5 ml syringe	0.00	10	<b>√</b> <u>G</u>	ardasil 9

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

#### INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

✓ Afluria Quad Junior (2021 Formulation)

#### A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
  - a) ischaemic heart disease, or
  - b) congestive heart failure, or
  - c) rheumatic heart disease, or
  - d) congenital heart disease, or
- e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
  - a) asthma, if on a regular preventative therapy, or
  - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
  - a) autoimmune disease, or
  - b) immune suppression or immune deficiency, or
  - c) HIV, or
  - d) transplant recipients, or
  - e) neuromuscular and CNS diseases/disorders, or
  - f) haemoglobinopathies, or
  - g) on long term aspirin, or
  - h) have a cochlear implant, or
  - i) errors of metabolism at risk of major metabolic decompensation, or
  - i) pre and post splenectomy, or
  - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine ini 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable
- c)

#### A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes: or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV. or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - j) pre and post splenectomy, or
    - k) down syndrome, or
  - vii) are pregnant:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine).......90.00 10 ✓ Fluad Quad (2021 Formulation)

Subsi	idy Fu	ly Brand or
(Manufacture	er's Price) Subsidise	ed Generic
\$	Per	<ul> <li>Manufacturer</li> </ul>

- a) Only on a prescription
- b) No patient co-payment payable

С

- INFLUENZA VACCINE people 65 years and over
  is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

inj 60 mcg in 0.5 mi syringe (paediatric quadrivalent vaccir	ie) –		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

#### A) INFLUENZA VACCINE – people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
  - a) ischaemic heart disease, or
  - b) congestive heart failure, or
  - c) rheumatic heart disease, or
  - d) congenital heart disease, or
  - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
  - a) asthma, if on a regular preventative therapy, or
  - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
  - a) autoimmune disease, or
  - b) immune suppression or immune deficiency, or
  - c) HIV, or
  - d) transplant recipients, or
  - e) neuromuscular and CNS diseases/disorders, or
  - f) haemoglobinopathies, or
  - g) are children on long term aspirin, or
  - h) have a cochlear implant, or
  - i) errors of metabolism at risk of major metabolic decompensation, or
  - i) pre and post splenectomy, or
  - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

## A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

# MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
  - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
  - 2) One dose for close contacts of meningococcal cases of any group; or
  - 3) One dose for person who has previously had meningococcal disease of any group; or
  - 4) A maximum of two doses for bone marrow transplant patients; or
  - 5) A maximum of two doses for person pre- and post-immunosuppression\*; or
- B) Both:
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either
    - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
    - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

NATIONAL IMMUNISATION SCHEDULE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression\*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients: or v) up to two doses for person pre- and post-immunosuppression\*. \*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression\*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. \*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B. 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10

Synflorix

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

#### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) with primary immune deficiencies; or
  - c) with HIV infection; or
  - d) with renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) with cochlear implants or intracranial shunts; or
  - g) with cerebrospinal fluid leaks; or
  - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) with cardiac disease, with cyanosis or failure; or
  - I) with diabetes; or
  - m) with Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Ini 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]		
<ol> <li>Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochl</li> <li>All of the following:</li> </ol>	tional asplenia, pre- or p	post-solid organ t	ransplant, renal dialysis,
<ul> <li>a) Patient is a child under 18 years for (re-)immur</li> <li>b) Treatment is for a maximum of two doses; and</li> <li>c) Any of the following:</li> </ul>	isation; and		
i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or	or	·	
or vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more prednisone of 2 mg/kg per day or greater	than two weeks, and wh		
20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or fail xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	estation; or ure; or	gh-dose corticost	eroid therapy); or
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ng: ndividuals; or	_	neumovax 23
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe ROTAVIRUS ORAL VACCINE – [Xpharm]		tch-up programm 1 ✓ <u>II</u>	
Maximum of two doses for patients meeting the following:  1) first dose to be administered in infants aged under 1- 2) no vaccination being administered to children aged 2			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

	Subsidy (Manufacturer's Price \$	e) S Per	Fully Subsidised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xph Either:	narm]			
1) Maximum of one dose for primary vaccination fo	r either:			
<ul> <li>a) Any infant born on or after 1 April 2016; or</li> <li>b) For previously unvaccinated children turnir varicella infection (chickenpox), or</li> </ul>		1 July 20	017, who h	ave not previously had a
Maximum of two doses for any of the following:				
a) Any of the following for non-immune patier	its:			
i) with chronic liver disease who may in ii) with deteriorating renal function before iii) prior to solid organ transplant; or iv) prior to any elective immunosuppress v) for post exposure prophylaxis who are considered by for patients at least 2 years after bone mand c) For patients at least 6 months after comple d) For HIV positive non immune to varicella we) For patients with inborn errors of metabolis varicella, or f) For household contacts of paediatric patient immune compromise where the household g) For household contacts of adult patients we immunocompromised, or undergoing a prohas no clinical history of varicella.  * immunosuppression due to steroid or other immunos	re transplantation; or sion*, or e immune competent inpa rrow transplantation, on action of chemotherapy, on with mild or moderate immuner at risk of major metabonts who are immunocompricontact has no clinical his ho have no clinical history cedure leading to immune	tients.; of livice of the device of the deconomised, tory of vof varice compro	or their special their spec ression on appensation or undergo aricella, or ella and wh mise where	cialist, or advice of HIV specialist, or , with no clinical history of oing a procedure leading to o are severely e the household contact
28 days Inj 1350 PFU prefilled syringe	0.00	1		<u>'arivax</u>
		10	_	<u>'arivax</u>
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTE		GLES V	ACCINE] -	- [Xpharm]
Funded for patients meeting either of the following crit  1) One dose for all people aged 65 years; or	ena.			
2) One dose for all people aged between 66 and 80	0 years inclusive from 1 Ar	ril 2018	and 31 De	cember 2021.
_, p.s.p.s ag.s	, ,			
Inj 19,400 PFU prefilled syringe plus vial	0.00	1 10		ostavax ostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	<b>√</b> <u>T</u>	ubersol

- Symbols -	Renin-Angiotensin System 48	AmsaLyo154
UK Synacthen81	Agents for Parkinsonism and Related	Amsidine154
3TC107	Disorders 120	Amzoate3
- A -	Agents Used in the Treatment of	Anaesthetics122
A-Scabies69	Poisonings240	Anagrelide hydrochloride154
Abacavir sulphate107	Agrylin154	Analgesics120
Abacavir sulphate with	Albendazole90	Anastrozole170
lamivudine 107	Albey225	Anatrole170
Abiraterone acetate167	Albustix78	Andriol Testocaps8
Acarbose11	Aldurazyme30	Androderm8
Accarb11	Alecensa161	ANI5
Accuretic49	Alectinib161	Anoro Ellipta229
Accuretic 1049	Alendronate sodium113	Antabuse146
Accuretic 2049	Alendronate sodium with	Antacids and Antiflatulents
Acetazolamide237	colecalciferol113	Anthelmintics90
Acetec48	Alfacalcidol34	Antiacne Preparations62
Acetic acid with 1, 2- propanediol	Alfamino260	Antiallergy Preparations225
diacetate and	Alfamino Junior260	Antianaemics38
benzethonium234	Alginic acid6	Antiandrogen Oral
Acetic acid with hydroxyquinoline and	Alglucosidase alfa27	Contraceptives76
ricinoleic acid76	Alkeran151	Antiarrhythmics50
Acetylcysteine240	Alkeran S29151	Antibacterials90
Aci-Jel76	Allersoothe226	Antibacterials Topical62
Aciclovir	Allmercap	Anticholinergic Agents228
Infection 102	Allopurinol118	Anticholinesterases112
Sensory234	Alpha-Adrenoceptor Blockers48	Antidepressants128
Acidex6	Alpha-Keri Lotion67	Antidiarrhoeals
Acipimox55	Alphamox93	Antiepilepsy Drugs130
Acitretin69	Alphamox 12593	Antifibrinolytics, Haemostatics and
Aclasta116	Alphamox 25093	Local Sclerosants39
Aclin	Alprolix39	Antifibrotics229
Actemra214	Alu-Tab6	Antifungals9
Actinomycin D155	Aluminium hydroxide6	Antifungals Topical63
Actrapid10	Alvogen52	Antihistamines226
Actrapid Penfill10	Amantadine hydrochloride120	Antihypotensives5
Acular236	Ambrisentan58	Antimalarials100
Acupan123	Ambrisentan Mylan58	Antimigraine Preparations133
Adalimumab178	Amiloride hydrochloride54	Antinausea and Vertigo Agents 134
Adapalene62	Amiloride hydrochloride with	Antipruritic Preparations64
Adcortyl81	furosemide55	Antipsychotics135
Adenuric118	Amiloride hydrochloride with	Antiretrovirals105
ADR Cartridge 1.824	hydrochlorothiazide55	Antirheumatoid Agents113
Adrenaline57	Aminophylline231	Antispasmodics and Other Agents
Advantan65	Amiodarone hydrochloride50	Altering Gut Motility
Advate42	Amisulpride135	Antithrombotic Agents42
Adynovate42	Amisulpride Mylan135	Antithymocyte globulin
Afinitor222	Amitriptyline	(equine) 178
Aflibercept	Amlodipine53	Antitrichomonal Agents100
Afluria Quad	Amneal158	Antituberculotics and
(2021 Formulation)	Amorolfine63	Antileprotics100
Afluria Quad Junior	Amoxicillin93	Antiulcerants
(2021 Formulation)	Amoxicillin with clavulanic acid94	Antivirals102
AFT-Pyrazinamide101	Amphotericin B33	Anxiolytics138
Agents Affecting the	Amsacrine154	Anzatax158

Apidra	11	Arrow-Roxithromycin	03	Basic AquaCream	6
Apidra SoloStar		Arrow-Sertraline		BCG Vaccine	
Apo-Azithromycin		Arrow-Timolol		Beclazone 100	
		Arrow-Timolor		Beclazone 250	
Apo-Bromocriptine		·			
Apo-Ciclopirox	03	Arrow-Tramadol		Beclazone 50	
Apo-Clarithromycin	0	Arsenic trioxide		Beclomethasone dipropionate	
Alimentary		Asacol		Bee venom allergy treatment	
Infection		Asamax		Bendamustine hydrochloride	
Apo-Clomipramine		Ascorbic acid		Bendrofluazide	5
Apo-Diclo SR		Aspen Adrenaline	5/	Bendroflumethiazide	_
Apo-Diltiazem CD		Aspirin		[Bendrofluazide]	
Apo-Doxazosin		Blood		Benzathine benzylpenicillin	
Apo-Folic Acid		Nervous		Benzatropine mesylate	
Apo-Furosemide		Asthalin		Benzbromaron AL 100	
Apo-Gabapentin		Atazanavir sulphate		Benzbromarone	
Apo-Megestrol		Atenolol	51	Benztrop	
Apo-Metoprolol	52	Atenolol AFT		Benzydamine hydrochloride	3
Apo-Mirtazapine	129	Atenolol AFT S29	51	Benzylpenicillin sodium [Penicillin	
Apo-Nadolol	52	ATGAM	178	G]	9
Apo-Oxybutynin	77	Ativan	138	Beta Cream	6
Apo-Perindopril	49	Atomoxetine	141	Beta Ointment	6
Apo-Pindolol	52	Atorvastatin	56	Beta Scalp	7
Apo-Prazosin		Atropine sulphate		Beta-Adrenoceptor Agonists	22
Apo-Prednisone		Cardiovascular	50	Beta-Adrenoceptor Blockers	
Apo-Primidone		Sensory	238	Betadine	
Apo-Propranolol	52	Atropt		Betadine Skin Prep	
Apo-Pyridoxine		Atrovent		Betaferon	14
Apo-Selegiline S29		AU Synacthen		Betahistine dihydrochloride	
Apo-Sumatriptan		Aubagio		Betaine	
Apo-Temozolomide	158	Augmentin		Betaloc CR	
Apomorphine hydrochloride		Aurorix		Betamethasone dipropionate	
Aprepitant		AutoSoft 30		Betamethasone dipropionate with	
Apresoline		AutoSoft 90		calcipotriol	6
Aptamil AllerPro SYNEO 1		Avelox		Betamethasone sodium phosphate	
Aptamil AllerPro SYNEO 2		Avonex		with betamethasone acetate	
Aptamil Gold+ Pepti Junior		Avonex Pen		Betamethasone valerate	
Aqueous cream		Azacitidine		Betamethasone valerate with sodiu	,
Aratac		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Arava		Azamun		Betaxolol	
Arginine		Azathioprine		Betnovate	
		Azilect		Betoptic	
Aripiprazole Sandoz		Azithromycin		Betoptic S	20
Aripiprazole Sandoz					
Aristocort		Azopt		Bexsero	
Arrow - Lattim		AZT	107	Bezafibrate	
Arrow-Amitriptyline		<del>-</del>	40	Bezalip	
Arrow-Bendrofluazide		B-D Micro-Fine		Bezalip Retard	
Arrow-Brimonidine		B-D Ultra Fine		Bicalutamide	
Arrow-Diazepam		B-D Ultra Fine II	16	Bicillin LA	
Arrow-Doxorubicin	155	Bacillus Calmette-Guerin (BCG)		BiCNU	
Arrow-Losartan &		vaccine	178	Bicnu Heritage	
Hydrochlorothiazide		Bacillus Calmette-Guerin		Bile and Liver Therapy	
Arrow-Norfloxacin		vaccine		Biltricide	
Arrow-Ornidazole		Baclofen		Bimatoprost	
Arrow-Quinapril 10		Bactroban		Bimatoprost Multichem	
Arrow-Quinapril 20		Balance		Binarex	
Arrow-Quinapril 5	49	Barrier Creams and Emollients	66	Binocrit	3

Biodone	Calcipotriol69	Cefuroxime axetil9
Biodone Extra Forte125	Calcitonin79	Celebrex11
Biodone Forte125	Calcitriol34	Celecoxib11
Bisacodyl27	Calcitriol-AFT34	Celecoxib Pfizer11
Bisoprolol fumarate51	Calcium carbonate6, 35	Celestone Chronodose8
Bisoprolol Mylan51	Calcium Channel Blockers53	Cellcept17
BK Lotion67	Calcium Disodium Versenate241	Centrally-Acting Agents5
Bleomycin sulphate154	Calcium folinate152	Cephalexin ABM9
Blood Colony-stimulating	Calcium Folinate Ebewe152	Cetirizine hydrochloride22
Factors45	Calcium Folinate Sandoz152	Cetomacrogol6
Blood glucose diagnostic test	Calcium Folinate Sandoz S29 152	Cetomacrogol with glycerol6
meter 15	Calcium gluconate35	Cetuximab18
Blood glucose diagnostic test	Calcium Homeostasis79	Champix14
strip15	Calcium polystyrene sulphonate47	Charcoal24
Blood glucose test strips (visually	Calcium Resonium47	Chemotherapeutic Agents14
impaired)15	Calcium-Sandoz Forte35	Chickenpox vaccine27
Blood Ketone Diagnostic Test	Calogen247	Chlorafast23
Strip 14	Calutide-50168	Chlorambucil15
Bonjela33	Candesartan cilexetil49	Chloramphenicol23
Boostrix265	Candestar49	Chlorothiazide5
Bortezomib	Canesten63	Chlorpheniramine maleate22
Bortezomib Dr-Reddy's	Capecitabine152	Chlorpromazine hydrochloride 13
Bosentan	Capercit	Chlortalidone [Chlorthalidone]5
Bosentan Dr Reddy's58	Capoten48	Chlorthalidone5
Bplex34	Capsaicin	Chlorvescent4
Breo Ellipta227	Musculoskeletal113	Choice Load 3757
Brevinor 1/28	Nervous123	Choice TT380 Short74
Bricanyl Turbuhaler228	Captopril48	Choice TT380 Standard74
Brilinta	Captopril-Mylan48	Choline salicylate with cetalkonium
Brimonidine tartrate237	Carafate10	chloride3
Brimonidine tartrate with timolol	Carbaccord150	Ciclopirox olamine6
maleate237	Carbamazepine130	Ciclosporin22
Brinzolamide237	Carbimazole84	Cilazapril4
Brolene	Carbomer238	Cilicaine9
Bromocriptine mesylate120	Carboplatin150	Cilicaine VK9
Brufen SR112	Carboplatin Ebewe150	Cinacalcet7
Buccastem135	Carbosorb-X240	Cipflox9
Budesonide	Cardinol LA52	Ciprofloxacin
Alimentary6	CareSens Dual14	Infection9
Respiratory226, 232	CareSens N	Sensory23
Budesonide with eformoterol227	CareSens N POP	Ciprofloxacin Teva23
Bumetanide54	CareSens N Premier	Circadin14
Buprenorphine Naloxone BNM 145	CareSens PRO	Cisplatin15
Buprenorphine with naloxone145	Carmellose sodium with gelatin and	Cisplatin Ebewe15
Bupropion hydrochloride146	pectin	Citalopram hydrobromide12
Burinex54	Carmustine150	Cladribine
Burinex S29	Carvedilol51	Clarithromycin
Buscopan8	Carvedilol Sandoz51	Alimentary
Buspirone hydrochloride138	Catapres 54	Infection9
Busulfan150	CeeNU151	Clexane4
- C -	Cefaclor monohydrate90	Clexane Forte4
Cabergoline89	Cefalexin90	Climara8
Cacit	Cefalexin Sandoz90	Climara
Caffeine citrate	Cefazolin90	Clindamycin9
Calamine	Ceftriaxone 90	Clinicians2
Calci-Tab 500	Ceftriaxone-AFT90	Clinicians Renal Vit3
Jaio. 140 000	JOHN 10 / 11 1	Omnolario i loriar vil

Clobazam1	30	Copaxone140	DBL Acetylcysteine24
Clobetasol propionate64,	70	Corticosteroids and Related Agents	DBL Adrenaline5
Clobetasone butyrate	65	for Systemic Use80	DBL Aminophylline23
Clofazimine1	00	Corticosteroids Topical64	DBL Bleomycin Sulfate15
Clomazol		Cortifoam7	DBL Carboplatin15
Dermatological	63	Cosentyx212	DBL Cisplatin15
Genito-Urinary		Cosmegen 155	DBL Dacarbazine15
Clomifene citrate		Coumadin45	DBL Desferrioxamine Mesylate for Inj
Clomipramine hydrochloride1	28	Country Life30	BP24
Clomipramine Teva1		Coversyl49	DBL Docetaxel15
Clonazepam 130, 1		Creon 1000025	DBL Ergometrine7
Clonidine		Creon 2500025	DBL Gemcitabine15
Clonidine BNM	54	Creon Micro25	DBL Gentamicin9
Clonidine hydrochloride	54	Crotamiton64	DBL Heparin Sodium4
Clopidogrel		Crystaderm62	DBL Leucovorin Calcium15
Clopidogrel Multichem		Curam94	DBL Methotrexate Onco-Vial15
Clopine1		Curam Duo 500/12594	DBL Morphine Sulphate12
Clopixol136, 1		Cvite34	DBL Naloxone Hydrochloride24
Clotrimazole		Cyclizine hydrochloride134	DBL Octreotide16
Dermatological	63	Cyclizine lactate134	DBL Pethidine Hydrochloride12
Genito-Urinary		Cyclogyl238	DBL Vinblastine16
Clozapine1	35	Cyclonex	DBL Vincristine Sulfate16
Clozaril1		Cyclopentolate hydrochloride238	Decozol3
Co-trimoxazole		Cyclophosphamide	Deferasirox24
Coal tar		Cyclorin	Deferiprone24
Coal tar with allantoin, menthol,		Cycloserine100	Denosumab11
phenol and sulphur	70	Cyproterone acetate81	Deolate9
Coal tar with salicylic acid and	, ,	Cyproterone acetate with	Deoxycoformycin15
sulphur	70	ethinyloestradiol	Depo-Medrol8
Coco-Scalp		Cystadane28	Depo-Provera7
Codeine phosphate	, ,	Cytarabine	Depo-Testosterone8
Extemporaneous2	43	Cytotec8	Deprim9
Nervous1		Cytoxan	Dermol
Coenzyme Q10		- D -	Desferrioxamine mesilate24
Colchicine		D-Penamine113	Desmopressin8
Colecalciferol		Dabigatran45	Desmopressin acetate8
Colestid		Dacarbazine	Desmopressin-PH&T8
Colestipol hydrochloride		Dacarbazine APP	Desuric11
Colgout1		Dactinomycin [Actinomycin D]155	Detection of Substances in
Colifoam		Daivobet69	Urine
Colistin sulphomethate		Daivonex69	Dexamethasone
Colistin-Link		Daktarin	Hormone8
Collodion flexible		Dalacin C	Sensory23
Colloidal bismuth subcitrate		Dantrium	Dexamethasone phosphate8
Colofac		Dantrium S29	Dexamethasone Phosphate
Coloxyl		Dantrolene	Panpharma8
Combigan2		Daonil	Dexamethasone with framycetin and
Compound electrolytes	٥ <i>١</i> 47		
	47	Dapa-Tabs	gramicidin
Compound electrolytes with glucose	47	Dapsone	Dexamethasone with neomycin
[Dextrose]		Daraprim	sulphate and polymyxin B
Compound hydroxybenzoate2		Darunavir	sulphate
Concerta1		Darunavir Mylan	Dexamfetamine sulfate
Condoms		Dasatinib	Dexmethsone8
Condyline		Daunorubicin	Dextrochlorpheniramine
Contraceptives - Hormonal		David One Step Cassette Pregnancy	maleate22
Contraceptives - Non-hormonal	/2	Test77	Dextrose46–4

DHC Continus125	Dostinex89	Emend Tri-Pack	.13
Diabetes10	Dosulepin [Dothiepin]	Emicizumab	
Diabetes Management14	hydrochloride128	EMLA	. 12
Diacomit132	Dosulepin Mylan128	Empagliflozin	13
Diagnostic Agents276	Dothiepin	Empagliflozin with metformin	
Diamide Relief6	Doxazosin48	hydrochloride	13
Diamox237	Doxine94	Emtricitabine	. 10
Diasip248	Doxorubicin Ebewe155	Emtricitabine with tenofovir	
Diason RTH248	Doxorubicin hydrochloride155	disoproxil	. 104
Diazepam130, 138	Doxycycline94	Emtriva	.10
Diazoxide10	DP Lotion67	Emulsifying ointment	
Dibenzyline48	DP Lotn HC65	Emulsifying Ointment ADE	6
Diclofenac Sandoz112	DP-Allopurinol118	Enalapril maleate	4
Diclofenac sodium	Dr Reddy's Omeprazole9	Enbrel	
Musculoskeletal112	Drugs Affecting Bone	Endocrine Therapy	
Sensory235	Metabolism 113	Endoxan	
Differin	Dual blood glucose and blood ketone	Engerix-B	
Difflam33	diagnostic test meter14	Enlafax XR	
Diflucan97	Dulaglutide12	Enoxaparin sodium	4
Digestives Including Enzymes25	Dulcolax SP Drop27	Enstilar	
Digoxin50	Duocal Super Soluble Powder246	Ensure	
Dihydrocodeine tartrate125	Duolin228	Ensure Plus	. 25
Dilantin	Duolin HFA228	Ensure Plus HN	. 25
Dilantin Infatab132	DuoResp Spiromax227	Ensure Plus RTH	. 25
Diltiazem hydrochloride53	Duride57	Ensure Two Cal HN RTH	
Dilzem53	- E -	Entacapone	. 12
Dimethicone66–67	e-chamber La Grande233	Entapone	
Dimethyl fumarate139	e-chamber Mask232	Entecavir	
Dipentum7	e-chamber Turbo233	Entecavir Sandoz	
Diphtheria, tetanus and pertussis	E-Mycin93	Entocort CIR	
vaccine 265	Ear Preparations234	Entresto 24/26	
Diphtheria, tetanus, pertussis and	Ear/Eye Preparations234	Entresto 49/51	49
polio vaccine265	Easiphen Liquid259	Entresto 97/103	
Diphtheria, tetanus, pertussis, polio,	Econazole nitrate63	Epilim	
hepatitis B and haemophilus	Efavirenz106	Epilim Crushable	. 13
influenzae type B vaccine 266	Efavirenz with emtricitabine and	Epilim IV	. 13
Diprosone64	tenofovir disoproxil107	Epilim S/F Liquid	
Diprosone OV64	Eformoterol fumarate227	Epilim Syrup	
Dipyridamole43	Eformoterol fumarate dihydrate227	Epirubicin Ebewe	
Disopyramide phosphate50	Eftrenonacog alfa [Recombinant	Epirubicin hydrochloride	. 15
Disulfiram146	factor IX]39	Eplerenone	5
Diuretics54	Efudix71	Epoetin alfa	3
Docetaxel	Egopsoryl TA70	Epoprostenol	6
Docetaxel Accord155	Elaprase29	Eptacog alfa [Recombinant factor	
Docetaxel Sandoz155	Eldepryl120	VIIa]	
Docusate sodium26	Elecare260	ERA	
Docusate sodium with	Elecare LCP260	Erbitux	
sennosides26	Electral47	Ergometrine maleate	7
Dolutegravir108	Elelyso32	Erlotinib	. 16
Domperidone 134	Elemental 028 Extra251	Erythrocin IV	
Donepezil hydrochloride 144	Elidel70	Erythromycin (as lactobionate)	
Donepezil-Rex144	Elocon	Erythromycin ethyl succinate	9
Dornase alfa231	Elocon Alcohol Free65	Erythromycin stearate	9
Dortimopt237	Eltrombopag39	Esbriet	
Dorzolamide hydrochloride237	Eltroxin84	Escitalopram	
Dorzolamide with timolol237	EMB Fatol101	Escitalopram (Ethics)	. 12

Escitalopram-Apotex	129	Fentanyl Sandoz	125	Fluticasone	22
Eskazole	90	Ferinject		Fluticasone furoate with	
Essential Generics	51	Ferodan	36	vilanterol	22
Essential Prednisolone	8	Ferriprox	240	Fluticasone propionate	
Estradiol TDP Mylan	82	Ferro-F-Tabs	36	Fluticasone with salmeterol	
Estradot		Ferro-tab		FML	23
Estradot 50 mcg	82	Ferrograd	36	Foban	6
Estrofem		Ferrosig	36	Folic acid	3
Etanercept	171	Ferrous fumarate		Folic Acid Mylan	3
Ethambutol hydrochloride	101	Ferrous fumarate with folic acid	36	Food Thickeners	
Ethics Aspirin		Ferrous sulfate	36	Foods And Supplements For Int	
Ethics Aspirin EC		Fexofenadine hydrochloride	226	Errors Of Metabolism	
Ethics Lisinopril		Fibro-vein	42	Foradil	22
Ethics Paracetamol Classic		Filgrastim	45	Forteo	11
Ethinyloestradiol	83	Finasteride		Fortini	25
Ethinyloestradiol with		Fingolimod	139	Fortini Multi Fibre	
desogestrel	74	Firazyr		Fortisip	
Ethinyloestradiol with		Flagyl		Fortisip Multi Fibre	
levonorgestrel	75	Flagyl-S		Fosamax	
Ethinyloestradiol with		Flamazine	63	Fosamax Plus	
norethisterone	75	Flecainide acetate		Framycetin sulphate	
Ethosuximide		Flecainide BNM		Frisium	
Etopophos		Flecainide Controlled Release		Frumil	
Etoposide		Teva	50	Frusemide	
Etoposide phosphate		Fleet Phosphate Enema		Fucicort	
Etravirine		Flixonase Hayfever & Allergy		Fucidin	
Eumovate		Flixotide		Fucithalmic	
Everet		Flixotide Accuhaler		Fulvestrant	
Everolimus		Florinef		Fungilin	
Evista		Fluad Quad		Furosemid-Ratiopharm	
Exemestane		(2021 Formulation)	269	Furosemide [Frusemide]	
Exjade		Fluanxol		Furosemide-Baxter	
Extemporaneously Compound		Flucil		fusidic acid	
Preparations and	aca	Flucloxacillin		Dermatological	63 6
Galenicals	243	Flucloxin		Infection	
Eye Preparations		Flucon		Sensory	
Eylea		Fluconazole		- G -	
Ezetimibe		Fludara Oral		Gabapentin	13
Ezetimibe Sandoz		Fludarabine Ebewe		Gacet	
Ezetimibe with simvastatin		Fludarabine phosphate		Galsulfase	
-F-		Fludrocortisone acetate		Galvumet	
Factor eight inhibitor bypassir	na	Fluids and Electrolytes		Galvus	
fraction		Flumetasone pivalate		Gardasil 9	
Famotidine		Fluocortolone caproate with	204	Gastrodenol	
Famotidine Hovid		fluocortolone pivalate and		Gaviscon Double Strength	
Faslodex		cinchocaine	8	Gaviscon Infant	
Febuxostat		Fluorometholone		Gazyva	
Febuxostat multichem		Fluorouracil		Gefitinib	
Feed Thickener Karicare	110	Fluorouracil Accord		Gemcitabine Ebewe	
Aptamil	257	Fluorouracil Ebewe		Gemcitabine hydrochloride	
FEIBA NF		Fluorouracil sodium		Genoptic	
Felo 10 ER		Fluox		Gentamicin sulphate	20
Felo 5 ER		Fluoxetine hydrochloride		Infection	۵
Felodipine		Flupenthixol decanoate		Sensory	
Femme-Tab ED		Flutamide		Gilenya	
Fentanyl		Flutamin		Ginet	
1 Omany	120	: :u:a:::::::::::::::::::::::::::::::::		On 101	

Glatiramer acetate		Heparon Junior		Hyoscine hydrobromide	
Glecaprevir with pibrentasvir		Hepatitis A vaccine	266	Hypam	
Glibenclamide		Hepatitis B recombinant		Hyperuricaemia and Antigout	
Gliclazide		vaccine		Hypromellose	
Glipizide		Herceptin		Hypromellose with dextran	23
Glivec		Hiberix		-1-	_
Glizide		Hiprex		Ibiamox	
Glucagen Hypokit		Histaclear		Ibrance	
Glucagon hydrochloride	10	Histafen		Ibuprofen	
Glucerna Select		Holoxan	.150	Ibuprofen SR BNM	
Glucobay		Horleys Bread Mix		Icatibant	
Glucose [Dextrose]	46	Horleys Flour	258	Idarubicin hydrochloride	
Gluten Free Foods	257	Hormone Replacement Therapy -		Idursulfase	2
Glycerin with sodium saccharin	243	Systemic	81	Ifosfamide	150
Glycerin with sucrose	243	HPV	267	Igroton	5
Glycerol		Humalog	11	Ikorel	5
Alimentary	27	Humalog Mix 25	11	llevro	23
Extemporaneous	243	Humalog Mix 50		lloprost	6
Glyceryl trinitrate		Human papillomavirus (6, 11, 16, 18		Imatinib mesilate	
Alimentary	8	31, 33, 45, 52 and 58) vaccine	,	Imatinib-Rex	16
Cardiovascular		[HPV]	267	Imigran	
Glycopyrronium		Humatin		Imipramine hydrochloride	
Glycopyrronium bromide		Humira		Imiquimod	
Glycopyrronium with		HumiraPen		Immune Modulators	
indacaterol	229	Humulin 30/70		Immunosuppressants	
Go Healthy		Humulin NPH		Imuran	
Gold Knight		Humulin R		Incruse Ellipta	
Gold Knight XL		Hyaluronic acid		Indacaterol	
Goserelin		Hydralazine		Indapamide	
Gutron		Hydralazine hydrochloride		Infanrix IPV	
Gynaecological Anti-infectives		Hydrocortisone	01	Infanrix-hexa	
- H -	70	Dermatological	65	Infant Formulae	
Habitrol	1/17	Hormone		Infatrini	
Haemophilus influenzae type B	147	Hydrocortisone (PSM)		Infliximab	
vaccine	266	Hydrocortisone acetate		Influenza vaccine	
Haldol		Hydrocortisone acetate with	/	Influvac Tetra	200
		pramoxine hydrochloride	7		07
Haldol Concentrate			/	(2021 Formulation) Inhaled Corticosteroids	
Haldol Decanoas		Hydrocortisone and paraffin liquid	e E		221
Haloperidol		and lanolin		Inhaled Long-acting	00.
Haloperidol decanoate		Hydrocortisone butyrate65		Beta-adrenoceptor Agonists	
Harvoni		Hydrocortisone with cinchocaine		Inspra	
Havrix		Hydrocortisone with miconazole	65	Instillagel Lido	
Havrix Junior		Hydrocortisone with natamycin and	0.5	Insulin aspart	1
Haylor syrup		neomycin		Insulin aspart with insulin aspart	4.
healthE Calamine Aqueous Crea		Hydrogen peroxide		protamine	
BP		Hydroxocobalamin	34	Insulin glargine	
healthE Dimethicone 10%		Hydroxocobalamin Mercury		Insulin glulisine	
healthE Dimethicone 4% Lotion		Pharma		Insulin isophane	10
healthE Dimethicone 5%		hydroxycarbamide		Insulin isophane with insulin	
healthE Glycerol BP		Hydroxychloroquine	113	neutral	1
healthE Urea Cream		Hydroxyurea		Insulin lispro	1
Healtheries Simple Baking Mix		[hydroxycarbamide]		Insulin lispro with insulin lispro	
Hemastix		Hygroton		protamine	
Hemlibra		Hylo-Fresh	.238	Insulin neutral	
Heparin sodium		Hymenoptera	.225	Insulin pen needles	
Heparinised saline	45	Hyoscine butylbromide	8	Insulin pump	10

Insulin pump cartridge21	Jakavi	165	Lanvis	15
Insulin pump infusion set (steel	Jardiamet		Lanzol Relief	
cannula)21	Jardiance	13	Lapatinib ditosylate	
Insulin pump infusion set (steel	Jaydess		Largactil	
cannula, straight insertion) 22	Jevity HiCal RTH		Laronidase	3
Insulin pump infusion set (teflon	Jevity RTH		Lasix	
cannula)23	Juno Pemetrexed		Latanoprost	
Insulin pump infusion set (teflon	- K -		Latanoprost with timolol	
cannula, angle insertion with	Kadcyla	219	Lax-Suppositories	
insertion device)24	Kaletra		Lax-Tab	
Insulin pump infusion set (teflon	Kalydeco		Laxatives	
cannula, angle insertion) 24	Kemadrin		Laxsol	
Insulin pump infusion set (teflon	Kenacomb		Ledipasvir with sofosbuvir	
cannula, straight insertion with	Kenacort-A 10		Leflunomide	
insertion device)24	Kenacort-A 40		Lenalidomide	
Insulin pump infusion set (teflon	Kenalog		Letrole	
cannula, straight insertion) 24	Kenalog in Orabase		Letrozole	
• ,			Leukeran FC	
Insulin pump reservoir24	Kenkay Sorbolene			15
Insulin syringes, disposable with	Ketocal 3:1		Leukotriene Receptor	00
attached needle	KetoCal 4:1	204	Antagonists	
Intal Forte CFC Free231	Ketoconazole	74	Leuprorelin	
Intelence	Dermatological		Leustatin	
Interferon beta-1-alpha140	Infection		Levetiracetam	
Interferon beta-1-beta140	Ketogenic Diet		Levetiracetam-AFT	
Intra-uterine device74	Ketoprofen		Levlen ED	
Invega Sustenna137	Ketorolac trometamol		Levocabastine	
IPOL275	KetoSens		Levocarnitine	
Ipratropium bromide228, 232	Keytruda		Levodopa with benserazide	
Iressa162	Kindergen	249	Levodopa with carbidopa	12
Irinotecan Accord153	Kivexa	107	Levomepromazine	13
Irinotecan Actavis 100153	Klacid		Levomepromazine	
Irinotecan hydrochloride153	Alimentary	9	hydrochloride	13
Irinotecan-Rex153	Infection		Levonorgestrel	
Iron (as ferric carboxymaltose)36	Kliogest	83	Genito-Urinary	75-7
Iron polymaltose36	Kliovance		Hormone	8
Isentress108	Kogenate FS	42	Levothyroxine	
Isentress HD108	Konakion MM	42	Lidocaine [Lignocaine]	
Ismo 2057	Konsyl-D	<mark>26</mark>	Lidocaine [Lignocaine]	
Ismo 40 Retard57	Kuvan		hydrochloride	12
Isoniazid101	-L-		Lidocaine [Lignocaine] with	
Isoniazid with rifampicin101	Labetalol	52	chlorhexidine	12
Isoptin53	Lacosamide	131	Lidocaine [Lignocaine] with	
Isoptin Retard53	Lactulose		prilocaine	12
Isoptin SR53	Laevolac		Lidocaine-Baxter	
Isopto Carpine237	Lamictal	131	Lidocaine-Claris	
Isosorbide mononitrate57	Lamivudine		Life Extension	
Isosource Standard255	Lamivudine Alphapharm		Lignocaine	
Isotretinoin	Lamotrigine		Lioresal Intrathecal	
Ispaghula (psyllium) husk26	Lamprene		Lipid-Modifying Agents	
Itch-Soothe64	Lanoxin		Liquigen	24
Itraconazole 98	Lanoxin Paediatric Elixir		Lisinopril	24 //
Itrazole 98	Lanoxin PG		Lithium carbonate	
lvacaftor231	Lanoxin S29		Livostin	
Ivermectin67	Lansoprazole		LMX4	
I-			Livix4Locacorten-Viaform ED's	
<b>J</b>	Lantus Lantus SoloStar			23
Jadelle75	Lantus Joiostar	11	Local preparations for Anal and	

Rectal Disorders	8	Maxitrol235	Methylphenidate hydrochloride
Locasol	260	MCT oil (Nutricia)247	extended-release143
Locoid	65, 71	Measles, mumps and rubella	Methylprednisolone80
Locoid Crelo	65	vaccine 272	Methylprednisolone (as sodium
Locoid Lipocream	65	Mebendazole90	succinate) 80
Locorten-Vioform	234	Mebeverine hydrochloride8	Methylprednisolone aceponate65
Lodoxamide	236	Medco	Methylprednisolone acetate80
Logem	131	Dermatological66	Methylxanthines231
Lomide	236	Nervous124	Metoclopramide Actavis 10134
Lomustine	151	Medrol80	Metoclopramide hydrochloride 134
Loniten	<u>58</u>	Medroxyprogesterone acetate	Metolazone54
Loperamide hydrochloride	<mark>6</mark>	Genito-Urinary75	Metopirone89
Lopinavir with ritonavir	107	Hormone82–83	Metoprolol IV Mylan52
Lopinavir/Ritonavir Mylan	107	Mefenamic acid112	Metoprolol succinate52
Loprofin	260	Megestrol acetate169	Metoprolol tartrate52
Loprofin Mix	259	Melatonin140	Metrogyl100
Lorafix		Melphalan151	Metronidazole100
Loratadine		Menactra272	Metyrapone89
Lorazepam	138	Meningococcal (groups A, C, Y and	Mexiletine hydrochloride51
Lorstat	<mark>56</mark>	W-135) conjugate vaccine 272	Mexiletine Hydrochloride USP51
Losartan Actavis	49	Meningococcal B multicomponent	Miacalcic79
Losartan potassium	49	vaccine 273	Micolette27
Losartan potassium with		Meningococcal C conjugate	Miconazole33
hydrochlorothiazide	49	vaccine273	Miconazole nitrate
Lovir	102	Menthol64	Dermatological64
Loxamine	129	Mepolizumab197	Genito-Urinary76
Lucrin Depot 1-month	88	Mercaptopurine153	Micreme76
Lucrin Depot 3-month		Mercilon 2874	Micreme H65
Lyderm	69	Mesalazine7	Microgynon 20 ED
Lynparza	157	Mesna157	Microgynon 3075
Lyrica		Mestinon112	Microgynon 50 ED
- M -		Metabolic Disorder Agents27	Microlut
m-Eslon	126	Metformin hydrochloride12	Midazolam141
Mabthera	200	Methadone hydrochloride	Midazolam-Baxter141
Macrobid	110	Extemporaneous243	Midodrine51
Macrogol 3350 with potassium	n	Nervous125	Mifegyne78
chloride, sodium bicarbona	ite and	Methatabs125	Mifepristone78
sodium chloride	<u>27</u>	Methenamine (hexamine)	Minerals35
Macrogol 400 and propylene		hippurate110	Mini-Wright AFS Low Range232
glycol	238	Methopt238	Mini-Wright Standard232
Madopar 125	120	Methotrexate153	Minidiab12
Madopar 250	120	Methotrexate DBL Onco-Vial153	MiniMed 1.8 Reservoir
Madopar 62.5	120	Methotrexate Ebewe153	MMT-326A24
Madopar HBS	120	Methotrexate Sandoz153	MiniMed 3.0 Reservoir
Madopar Rapid	120	Methyl hydroxybenzoate243	MMT-332A24
Magnesium hydroxide	37	Methylcellulose243	MiniMed 640G16
Magnesium sulphate	37	Methylcellulose with glycerin and	MiniMed Mio MMT-921A23
Mantoux	276	sodium saccharin243	MiniMed Mio MMT-923A23
Marevan	45	Methylcellulose with glycerin and	MiniMed Mio MMT-925A23
Marine Blue Lotion SPF 50+ .	71	sucrose243	MiniMed Mio MMT-941A23
Martindale Pharma	240	Methyldopa54	MiniMed Mio MMT-943A23
Marvelon 28		Methyldopa Mylan54	MiniMed Mio MMT-945A23
Mask for spacer device	232	Methyldopa Mylan S2954	MiniMed Mio MMT-965A23
Mast Cell Stabilisers	231	Methylnaltrexone bromide26	MiniMed Mio MMT-975A23
Maviret	103	Methylphenidate ER - Teva142	MiniMed Quick-Set MMT-386A23
Maxidex	235	Methylphenidate hydrochloride 142	MiniMed Quick-Set MMT-387A23

MiniMed Quick-Set MMT-396A					
	23	Mucolytics	231	Nepro HP RTH	25
MiniMed Quick-Set MMT-397A	23	Mucosoothe	122	Neulactil	13
MiniMed Quick-Set MMT-398A	23	Multiple Sclerosis Treatments	139	Neulastim	4
MiniMed Quick-Set MMT-399A	23	Multivitamin renal	34	NeuroTabs	3
MiniMed Silhouette MMT-368A	23	Multivitamins		Nevirapine	10
MiniMed Silhouette MMT-377A	23	Mupirocin	63	Nevirapine Alphapharm	
MiniMed Silhouette MMT-378A	23	Muscle Relaxants	119	Nicorandil	
MiniMed Silhouette MMT-381A	23	Mvite	35	Nicotine	14
MiniMed Silhouette MMT-382A	23	Myambutol	101	Nifedipine	5
MiniMed Silhouette MMT-383A	23	Mycobutin	101	Nifuran	11
MiniMed Silhouette MMT-384A	23	MycoNail	63	Nilotinib	
MiniMed Sure-T MMT-864A	21	Mycophenolate mofetil		Nilstat	
MiniMed Sure-T MMT-866A	21	Mydriacyl	238	Alimentary	3
MiniMed Sure-T MMT-874A	21	Mylan (12 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-876A	21	Mylan (24 hr release)		Infection	9
MiniMed Sure-T MMT-884A	21	Mylan Atenolol		Nintedanib	
MiniMed Sure-T MMT-886A	21	Mylan Clomiphen		Nipent	15
Minims Cyclopentolate	238	Mylan Indapamide		Nitrates	
Minims Pilocarpine		Mylan Midazolam		Nitroderm TTS	
Minims Prednisolone		Myleran		Nitrofurantoin	
Minirin		Myometrial and Vaginal Hormone		Nitrolingual Pump Spray	
Minirin Melt		Preparations		Nivestim	
Mino-tabs		Myozyme		Nivolumab	
Minocycline hydrochloride		- N -		Nizoral	
Minomycin		Nadolol	52	Nodia	
Minor Skin Infections		Naglazyme		Noflam 250	
Minoxidil		Nalcrom		Noflam 500	
WIII IOAIUII				Nonam 500	! !!
Mirena	83	Naloxone hydrochloride	240		
Mirena		Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	111
Mirtazapine	129	Naltraccord	146	Non-Steroidal Anti-Inflammatory Drugs	11
Mirtazapine	129 8	NaltraccordNaltrexone hydrochloride	146 146	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant	
Misoprostol	129 8 157	Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride	146 146 239	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX]	
Mirtazapine	129 8 157 157	Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte	146 146 239 239	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe	129 8 157 157	Naltraccord	146 146 239 239	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone Genito-Urinary	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30	129 8 157 157 157	Naltraccord	146 239 239 112	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone Genito-Urinary Hormone	4: 7: 8:
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II	129 8 157 157 157 11	Naltraccord	146 239 239 112 112	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone Genito-Urinary Hormone Norflex	4: 7: 8: 11:
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide	129 8 157 157 157 11 272	Naltraccord	146 146 239 112 112 112	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone Genito-Urinary Hormone Norflex Norfloxacin	4: 7: 8: 11: 11
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil	129 157 157 157 11 272 128	Naltraccord	146 146 239 239 112 112 112 118 232	Non-Steroidal Anti-Inflammatory Drugs  Nonacog gamma, [Recombinant Factor IX]  Norethisterone Genito-Urinary  Hormone  Norflex  Norfloxacin  Noriday 28	4/ 7/ 11/ 11
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil	129815715715711272128144	Naltraccord	146 146 239 239 112 112 112 118 232 140	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX]. Norethisterone Genito-Urinary Hormone Norflex Norfloxacin Noriday 28 Norimin	4/ 8/ 11/ 11 7/
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic		Naltraccord	146 146 239 112 112 112 118 232 140 158	Non-Steroidal Anti-Inflammatory Drugs Nonacog gamma, [Recombinant Factor IX] Norethisterone Genito-Urinary Hormone Norflex Norfloxacin Noriday 28 Norimin Normacol Plus	4: 7: 11: 11: 7: 7: 7:
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole		Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix	146239239112112118232140158	Non-Steroidal Anti-Inflammatory Drugs	4: 7: 11: 11: 7: 7: 2: 14
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm	146146239239112112118232140158135	Non-Steroidal Anti-Inflammatory Drugs	4 7 11 11 7 2 14 12
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Nacaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine	146146239239112112118232140158135134	Non-Steroidal Anti-Inflammatory Drugs	
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Nacaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Navelbine Nefopam hydrochloride	146146239239112112112118232140158134161123	Non-Steroidal Anti-Inflammatory Drugs	
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C	146146239239112112112118232158134161123273	Non-Steroidal Anti-Inflammatory Drugs	4 7 11 11 7 7 12 12 12 12
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Montelukast Montelukast		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12	14614623923911211211823214015813513416112327334	Non-Steroidal Anti-Inflammatory Drugs	4 7 11 11 7 7 12 10 12 12
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Montelukast Mylan Moroctocog alfa [Recombinant f		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole	1461462392391121121182321401581351341611232733484	Non-Steroidal Anti-Inflammatory Drugs	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Montelukast Montelukast Montelukast Mylan Moroctocog alfa [Recombinant f	129815715715711272128144	Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole S29	1461462392391121121182321401581351341611232733484	Non-Steroidal Anti-Inflammatory Drugs	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Mometasone furoate Monogen Montelukast Montelukast Mylan Moroctocog alfa [Recombinant f VIII] Morphine hydrochloride		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole S29 Neocate Gold	146146239239112112118232140158135134161123273348484	Non-Steroidal Anti-Inflammatory Drugs	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Monotelukast Moroctocog alfa [Recombinant t VIII] Morphine hydrochloride Morphine sulphate		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations. Natalizumab Natulan Nausicalm Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured	146146239239112112118232140158135134161123273348484260260	Non-Steroidal Anti-Inflammatory Drugs	4 7/ 11 11 12 12 12 12 12 12 12 12 12
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Mylan Moroctocog alfa [Recombinant f VIII] Morphine hydrochloride Morphine sulphate Motetis		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla		Non-Steroidal Anti-Inflammatory Drugs	4 7,
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Mylan Moroctocog alfa [Recombinant f VIII] Morphine hydrochloride Morphine sulphate Motetis Motetis Motetis Motetis Motetis Mothand Throat		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla Neocate SYNEO		Non-Steroidal Anti-Inflammatory Drugs	4 7 8 11 11 7 2 14 12 10 12 12 10 12 11 11 11 11 11 11 11 11 11 11
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Montelukast Mylan Moroctocog alfa [Recombinant I VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla Neocate SYNEO Neoral		Non-Steroidal Anti-Inflammatory Drugs	4 7 8 11 11 7 2 14 12 10 12 10 12 11 11 1 1 1 1 1
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Mylan Moroctocog alfa [Recombinant f		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla Neocate SYNEO Neoral Neostigmine metilsulfate		Non-Steroidal Anti-Inflammatory Drugs	4
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Montelukast Montelukast Mylan Moroctocog alfa [Recombinant to VIII] Morphine hydrochloride Morphine sulphate Motetis Mouth and Throat Movapo Moxifloxacin MSUD Maxamum		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulian Nausafix Nausicalm Navelbine Nefopam hydrochloride Nefopam hydrochloride Neo-B12 Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla Neocate SYNEO Neoral Neostigmine metilsulfate Neostigmine metilsulfate		Non-Steroidal Anti-Inflammatory Drugs	
Mirtazapine Misoprostol Mitomycin C Mitozantrone Mitozantrone Ebewe Mixtard 30 MMR II Moclobemide Modafinil Modavigil Moduretic Molaxole Moments Mometasone furoate Monogen Montelukast Mylan Moroctocog alfa [Recombinant f		Naltraccord Naltraccord Naltrexone hydrochloride Naphazoline hydrochloride Naphcon Forte Naprosyn SR 1000 Naprosyn SR 750 Naproxen Narcaricin mite Nasal Preparations Natalizumab Natulan Nausafix Nausicalm Navelbine Nefopam hydrochloride Neisvac-C Neo-B12 Neo-Mercazole Neo-Mercazole Neo-Mercazole S29 Neocate Gold Neocate Junior Unflavoured Neocate Junior Vanilla Neocate SYNEO Neoral Neostigmine metilsulfate		Non-Steroidal Anti-Inflammatory Drugs	

Nuelin	231	Olopatadine Teva	239	maleate	76
Nuelin-SR	231	Olsalazine	7	Ozurdex	235
Nupentin	131	Omalizumab	198	- P -	
Nutilis	257	Omeprazole	9	Pacifen	119
Nutren Diabetes		Omeprazole actavis 10	9	Pacimol	
Nutrient Modules	245	Omeprazole actavis 20	9	Paclitaxel	158
Nutrini Energy Multi Fibre	250	Omeprazole actavis 40	9	Paclitaxel Actavis	158
Nutrini Energy RTH	250	Omnitrope	84	Paclitaxel Ebewe	158
Nutrini Low Energy Multi Fibre		Onbrez Breezhaler	227	Paediatric Seravit	35
Nutrini Peptisorb	262	Oncaspar LYO	158	Palbociclib	164
Nutrini Peptisorb Energy		OncoTICE	178	Paliperidone	
Nutrini RTH	250	Ondansetron		Pamidronate disodium	114
Nutrison 800 Complete Multi		Ondansetron ODT-DRLA	135	Pamisol	
Fibre	255	One-Alpha	34	Panadol	124
Nutrison Concentrated	257	Onrex	135	Panadol Mini Caps	124
Nutrison Energy	254	Opdivo		Pancreatic enzyme	25
Nutrison Energy Multi Fibre		Ora-Blend	243	Pantoprazole	9
Nutrison Multi Fibre		Ora-Blend SF		Panzop Relief	
Nutrison Standard RTH		Ora-Plus		Panzytrat	
Nyefax Retard		Ora-Sweet		Papaverine hydrochloride	
Nystatin		Ora-Sweet SF		Para-amino salicylic acid	
Alimentary	. 33	Orabase	33	Paracare	
Genito-Urinary		Oral and Enteral Feeds	248	Paracare Double Strength	
Infection	. 98	Oratane		Paracetamol	
NZB Low Gluten Bread Mix	258	Ordine	125	Paracetamol + Codeine	
- 0 -		Orgran	258	(Relieve)	127
O/W Fatty Emulsion Cream	. 66	Ornidazole		Paracetamol Pharmacare	124
Obinutuzumab		Orphenadrine citrate		Paracetamol with codeine	
Obstetric Preparations	78	Ortho-tolidine		Paraffin	
Ocicure		Oruvail SR	112	Paraffin liquid with wool fat	239
Ocrelizumab	140	Osmolite RTH	255	Paraldehyde	
Ocrevus		Other Endocrine Agents	89	Parasidose	
Octocog alfa [Recombinant factor		Other Oestrogen Preparation		Parasiticidal Preparations	
VIII] (Advate)	. 42	Other Progestogen		Parlodel	
Octocog alfa [Recombinant factor		Preparations	83	Parnate	
VIII] (Kogenate FS)	. 42	Other Skin Preparations		Parnate S29	128
Octreotide		Ovestin		Paromomycin	96
Octreotide (Sun)		Genito-Urinary	76	Paroxetine	
Octreotide Depot Teva		Hormone		Paser	101
Octreotide GH		Ox-Pam	138	Paxam	138
Octreotide long-acting		Oxaliplatin	151	Paxtine	129
Octreotide MaxRx		Oxaliplatin Accord		Pazopanib	165
Oestradiol	.82	Oxaliplatin Actavis 100		Peak flow meter	
Oestradiol valerate	.82	Oxaliplatin Ebewe		Pedialyte - Bubblegum	47
Oestradiol with norethisterone	.83	Oxazepam		Pediasure	250
Oestriol		Oxis Turbuhaler	227	Pediasure RTH	250
Genito-Urinary	.76	Oxpentifylline		Pegaspargase	
Hormone		Oxybutynin		Pegasys	
Oestrogens	.82	Oxycodone hydrochloride		Pegfilgrastim	
Ofev		Oxycodone Sandoz		Pegylated interferon alfa-2a	
Oil in water emulsion		Oxycodone Sandoz S29		Pembrolizumab	
Olanzapine136-		OxyContin		Pemetrexed	
Olaparib	157	OxyNorm		Penicillamine	
Olbetam		Oxytocin		Penicillin G	
Olbetam S29	.55	Oxytocin BNM		PenMix 30	
Olopatadine		Oxytocin with ergometrine		PenMix 40	11
		· ·			

PenMix 50	11	PKU Lophlex Sensation 20	259	Proctosedyl	
Pentasa		Plaquenil		Procyclidine hydrochloride	
Pentostatin [Deoxycoformycin]	158	Plendil ER		Procytox	
Pentoxifylline [Oxpentifylline]		Pneumococcal (PCV10) conjugate	е	Progesterone	
Peptamen Junior		vaccine		Proglicem	
Peptisorb		Pneumococcal (PCV13) conjugat		Proglycem	
Perhexiline maleate		vaccine		Progynova	
Pericyazine		Pneumococcal (PPV23)		Prolia	
Perindopril		polysaccharide vaccine	275	Promethazine hydrochloride	
Perjeta		Pneumovax 23		Propafenone hydrochloride	
Permethrin		Podophyllotoxin	71	Propamidine isethionate	23
Perrigo	71	Polaramine	226	Propranolol	5
Pertuzumab		Poliomyelitis vaccine		Propylene glycol	
Peteha	101	Poloxamer		Propylthiouracil	
Pethidine hydrochloride	127	Poly-Gel	238	Prostacur	16
Pevaryl		Poly-Tears		Protaphane	
Pexsig		Poly-Visc		Protaphane Penfill	
Pfizer Exemestane	170	Polycal		Protifar	
Pharmacy Health Sorbolene with		Ponstan		Protionamide	
Glycerin		Posaconazole	98	Provera	8
Pheburane		Postinor-1	76	Provera HD	8
Phenasen	154	Potassium chloride	46-47	PSM Citalopram	
Phenobarbitone	131	Potassium Chloride Aguettant		Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate		Preparations	6
Extemporaneous	243	Potassium iodate		PTU	
Nervous		Povidone iodine	67	Pulmicort Turbuhaler	22
Phenothrin	69	Pradaxa	45	Pulmozyme	23
Phenoxybenzamine		Pramipexole hydrochloride		Puri-nethol	
hydrochloride	48	Pravastatin		Puria	3
Phenoxymethylpenicillin (Penicill		Pravastatin Mylan	56	Pyrazinamide	
V)		Praziquantel		Pyridostigmine bromide	11
Phenytoin sodium1		Prazosin		Pyridoxine hydrochloride	
Phillips Milk of Magnesia		Pred Forte	236	Pyrimethamine	
Phlexy 10		Prednisolone	81	Pytazen SR	
Phosphate Phebra		Prednisolone acetate	236	. Q -	
Phosphorus		Prednisolone sodium	8	Quetapel	13
Phytomenadione	42	Prednisolone sodium		Quetiapine	13
Pilocarpine hydrochloride		phosphate	236	Quick-Set MMT-392	
Pimafucort		Prednisolone-AFT		Quick-Set MMT-393	
Pimecrolimus	70	Prednisone	81	Quinapril	
Pindolol	52	Pregabalin	132	Quinapril with	
Pine tar with trolamine laurilsulfat	te	Pregabalin Pfizer	132	hydrochlorothiazide	4
and fluorescein		Pregnancy Tests - hCG Urine	77	Qvar	
Pinetarsol	70	Premarin		- R -	
Pioglitazone	12	Prevenar 13		RA-Morph	12
Pirfenidone	230	Priadel	136	Raloxifene hydrochloride	
Pizotifen	134	Primaquine		Raltegravir potassium	
PKU Anamix Infant	259	Primidone	132	Ramipex	
PKU Anamix Junior	259	Primolut N	83	Ranbaxy-Cefaclor	9
PKU Anamix Junior Chocolate	259	Priorix	272	Ranbaxy-Cefaclor S29	9
PKU Anamix Junior LQ	259	Probenecid		Rapamune	22
PKU Anamix Junior Orange	259	Probenecid-AFT	119	Rasagiline	12
PKU Anamix Junior Vanilla		Procaine penicillin	94	Reandron 1000	8
PKU Lophlex LQ 10	259	Procarbazine hydrochloride	158	Recombinant factor IX	
PKU Lophlex LQ 20		Prochlorperazine		Recombinant factor VIIa	
PKU Lophlex Powder		Proctofoam	<mark>7</mark>	Recombinant factor VIII	

Rectogesic	8	Rubifen SR	142	Sinemet CR	120
Redipred		Rugby Capsaicin Topical		Sirolimus	222
Relieve		Cream	123	Siterone	81
Relistor		Rulide D		Slow-Lopresor	
Remicade		Rurioctocog alfa pegol [Reco		Smith BioMed Rapid Pregnancy	
Renilon 7.5		factor VIII]		Test	77
Resonium-A		Ruxolitinib		Sodibic	
Resource Beneprotein		Rythmodan		Sodium acid phosphate	
Respigen	228	Rytmonorm		Sodium alginate	6
Respiratory Devices		- S -		Sodium benzoate	31
Respiratory Stimulants		Sabril	132	Sodium bicarbonate	
Retinol palmitate		Sacubitril with valsartan		Blood	46_47
ReTrieve		Sagent		Extemporaneous	
Retrovir		SalAir		Sodium calcium edetate	
Revlimid		Salazopyrin		Sodium chloride	241
Revolade		. ,		Blood	10
		Salazopyrin EN			
Rexacrom		Salbutamol	220	Respiratory	202
Riboflavin		Salbutamol with ipratropium	000	Sodium citrate with sodium lauryl	07
Ribomustin		bromide		sulphoacetate	
Ricit		Salicylic acid		Sodium citro-tartrate	/8
Rifabutin		Salmeterol		Sodium cromoglicate	
Rifadin		Sandomigran		Alimentary	
Rifampicin		Sandostatin LAR		Respiratory	
Rifaximin		Sanofi Primaquine		Sensory	
Rifinah		Sapropterin dihydrochloride		Sodium fluoride	35
Rilutek		Scalp Preparations		Sodium Fusidate [fusidic acid]	
Riluzole		Scopoderm TTS		Dermatological	
Riodine		Sebizole		Infection	
Risedronate Sandoz		Secukinumab		Sensory	234
Risedronate sodium		Sedatives and Hypnotics	140	Sodium hyaluronate [Hyaluronic	
Risperdal Consta		Seebri Breezhaler		acid]	
Risperidone	136, 138	Selegiline hydrochloride	120	Sodium phenylbutyrate	
Risperidone (Teva)		Senna	27	Sodium picosulfate	27
Risperon	136	Senokot		Sodium polystyrene sulphonate	47
Ritalin	142	Sensipar	79	Sodium tetradecyl sulphate	42
Ritalin LA	143	SensoCard	15	Sodium valproate	132
Ritonavir	107	Serenace	136	Sofradex	234
Rituximab (Mabthera)	200	Seretide	228	Soframycin	234
Rituximab (Riximyo)	202	Seretide Accuhaler	228	Solgar	28-31
Rivaroxaban	45	Serevent	227	Solifenacin Mylan	
Rivastigmine	145	Serevent Accuhaler	227	Solifenacin succinate	
Rivastigmine Patch BNM 10.	145	Sertraline	129	Solu-Cortef	80
Rivastigmine Patch BNM 5		Setrona	129	Solu-Medrol	
Rivotril	130	Setrona AU	129	Solu-Medrol-Act-O-Vial	80
Riximyo	202	Sevredol	126	Somatropin (Omnitrope)	84
RIXUBIS	42	Sex Hormones Non		Sotalol	
Rizamelt	133	Contraceptive	81	Spacer device	233
Rizatriptan	133	Shield XL		Span-K	
Ropin		shingles vaccine		Spiolto Respimat	229
Ropinirole hydrochloride		SII-Onco-BCG		Spiractin	55
Rotarix		Sildenafil		Spiriva	229
Rotavirus oral vaccine		Silhouette MMT-373		Spiriva Respimat	229
Roxane		Siltuximab		Spironolactone	
Roxane-Propranolol		Simvastatin		Sporanox	
Roxithromycin		Simvastatin Mylan		Sprycel	
Rubifen		Sinemet		Staphlex	
		OII 1011 101	120	Otapi IIOA	34

## **INDEX: Generic Chemicals and Brands**

Stemetil	135	Tarceva	162	Tolcapone	121
SteroClear	232	Tasigna	164	Topamax	
Stesolid	130	Tasmar		Topical Products for Joint and	
Stimulants/ADHD Treatments	141	Taurine		Muscular Pain	113
Stiripentol	132	Tecfidera	139	Topiramate	
Stocrin		Tegretol		Topiramate Actavis	
Stomahesive		Tegretol CR		Total parenteral nutrition (TPN)	
Strattera		Telfast		TPN	
Strides Shasun		Teligent		Tramadol hydrochloride	
Stromectol		Temaccord		Tramal SR 100	
Sucralfate		Temazepam		Tramal SR 150	
Sulfadiazine Silver		Temozolomide		Tramal SR 200	
Sulfadiazine sodium		Tenofovir disoproxil		Trandate	
Sulfasalazine		Tenofovir Disoproxil Teva		Tranexamic acid	
Sulindac		Tenoxicam		Tranylcypromine sulphate	
Sulindac Mylan		Tensipine MR10		Trastuzumab	
Sulphur		Tepadina		Trastuzumab emtansine	
Sulprix		Terbinafine		Travatan	
Sumagran		Terbutaline sulphate		Travoprost	
Sumatriptan		Teriflunomide		Travopt	
Sunitinib		Teriparatide		Treatments for Dementia	144
Sunscreens		Testosterone		Treatments for Substance	
Sunscreens, proprietary		Testosterone cipionate		Dependence	
Sure-T MMT-863		Testosterone esters		Trental 400	58
Sure-T MMT-873		Testosterone undecanoate		Tretinoin	
Sustagen Diabetic	248	Tetrabenazine		Dermatological	
Sustagen Hospital Formula		Tetrabromophenol		Oncology	
Active		Tetracosactrin		Trexate	153
Sustanon Ampoules	81	Tetracycline	95	Triamcinolone acetonide	
Sutent		Thalidomide	159	Alimentary	33
Sylvant	214	Thalomid	159	Dermatological	65
Symbicort Turbuhaler 100/6	227	Theophylline	231	Hormone	81
Symbicort Turbuhaler 200/6	227	Thiamine hydrochloride	34	Triamcinolone acetonide with	
Symbicort Turbuhaler 400/12	227	THIO-TEPA	151	gramicidin, neomycin and nysta	atin
Symmetrel	120	Thioguanine	154	Dermatological	66
Sympathomimetics		Thiotepa		Sensory	
Synacthen		Thymol glycerin		Triaver	
Synacthen Depot		Thyroid and Antithyroid Agents		Triazolam	141
Synacthene Retard		Ticagrelor		Trimethoprim	
Synflorix		Tilcotil		Trimethoprim with	
Synthroid		Tillomed		sulphamethoxazole	
Syntometrine	76	Timolol		[Co-trimoxazole]	97
Syrup (pharmaceutical grade)		Timoptol XE		Trisequens	
Systane Unit Dose		Tiotropium bromide		Trisul	
- T -	200	Tiotropium bromide with	220	Trophic Hormones	
Tacrolimus	224	olodaterol	220	Tropicamide	
Tacrolimus Sandoz		Tivicay		Trulicity	12
Taliglucerase alfa		TMP	31	Trusopt TruSteel	
		Tobramycin	07		
Tamoxifen citrate		Infection		Tuberculin PPD [Mantoux] test	
Tamoxifen Sandoz		Sensory		Tubersol	
Tamsulosin hydrochloride		Tobramycin BNM	97	Two Cal HN	
Tamsulosin-Rex		Tobramycin Mylan		Two Cal HN RTH	
Tandem Cartridge		Tobrex		Tykerb	
Tandem t:slim X2 with Basal-IQ		Tocilizumab		Tysabri	140
Tap water	244	Tofranil	128	- U -	

## **INDEX: Generic Chemicals and Brands**

Ultibro Breezhaler229	hydrochloride12	Zinc sulphate
Ultraproct8	Vimpat131	Zincaps
Umeclidinium229	Vinblastine sulphate161	Zinnat
Umeclidinium with vilanterol229	Vincristine sulphate161	Ziprasidone
Univent228, 232	Vinorelbine161	Zista
Ural78	Vinorelbine Ebewe161	Zithromax
Urea66	Viramune Suspension106	Zoledronic acid
Urex Forte54	ViruPOS234	Hormone
Urinary Agents77	Vit.D334	Musculoskeletal
Urinary Tract Infections110	Vita-B1234	Zoledronic acid Mylar
Urinorm118	VitA-POS239	Zopiclone
Uromitexan 157	Vitabdeck35	Zopiclone Actavis
Ursodeoxycholic acid25	Vital251	Zostavax
Ursosan	Vitamin B complex34	Zostrix
Utrogestan83	Vitamin B6 2534	Zostrix HP
- V -	Vitamins34–35	Zuclopenthixol decan
Vaccinations265	Vivonex TEN251	Zuclopenthixol hydro
Vaclovir102	Voltaren112	Zusdone
Valaciclovir102	Voltaren D112	Zyban
Valganciclovir102	Voltaren Ophtha235	Zypine
Valganciclovir Mylan102	Voltaren SR112	Zypine ODT
Vancomycin97	Volumatic233	Zyprexa Relprevv
Vannair227	Voriconazole99	Zytiga
Varenicline Pfizer147	Vosol234	, 0
Varenicline tartrate147	Votrient165	
Varicella vaccine [Chickenpox	Vttack99	
vaccine] 276	- W -	
Varicella zoster virus (Oka strain) live	Warfarin sodium45	
attenuated vaccine shingles	Wart Preparations71	
vaccine] 276	Wasp venom allergy treatment225	
Varivax276	Water	
Vasodilators57	Blood47	
Vasopressin Agonists88	Extemporaneous244	
Vasorex53	Wool fat with mineral oil67	
Vedafil59	- X -	
Veletri60	Xarelto45	
Venclexta160	Xifaxan10	
Venetoclax160	XMET Maxamum259	
Venlafaxine130	Xolair198	
Venomil225	XP Maxamum259	
VENOX225	Xylocaine122	
Ventavis61	Xylocaine 2% Jelly122	
Ventolin228	Xyntha41	
Vepesid155	- Z -	
Verapamil hydrochloride53	Zapril48	
Vergo 16134	Zarontin	
Vermox90	Zaroxolyn54	
Versacloz135	Zavedos	
Vesanoid	Zeffix102	
Vexazone12	Zetlam102	
Vfend99	Ziagen107	
Viaderm KC66	Zidovudine [AZT]107	
Vidaza151	Zidovudine [AZT] with	
Vigabatrin132	lamivudine	
Vildagliptin	Zimybe	
Vildagliptin with metformin	Zinc and castor oil66	

Zinc sulphate	37
Zincaps	37
Zinnat	90
Ziprasidone	136
Zista	226
Zithromax	91
Zoledronic acid	
Hormone	79
Musculoskeletal	116
Zoledronic acid Mylan	79
Zopiclone	
Zopiclone Actavis	
Zostavax	
Zostrix	113
Zostrix HP	
Zuclopenthixol decanoate	138
Zuclopenthixol hydrochloride	
Zusdone	
Zyban	
Zypine	
Zypine ODT	
Zyprexa Relprevy	
Zvtina	