October 2021 Volume 28 Number 2

Editors:

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at no cost from the Pharmac website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	Introducing Pharmac I	2
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	63
	Genito Urinary System	73
	Hormone Preparations – Systemic	80
	Infections – Agents For Systemic Use	91
	Musculoskeletal System	113
	Nervous System	122
	Oncology Agents & Immunosuppressants	150
	Respiratory System & Allergies	228
	Sensory Organs	237
	Various	243
Section C	Extemporaneous Compounds (ECPs)	245
Section D	Special Foods	248
Section I	National Immunisation Schedule	268
	Index	280

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 n sachet	0.	30	✓ Ga	viscon Infant
GODIUM ALGINATE ★ Tab 500 mg with sodium bicarbonate 267 mg and calciur carbonate 160 mg - peppermint flavour		60		viscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and cal carbonate 160 mg per 10 ml		500 ml	Ac	idex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE		100	✓ Alı	u-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) Subsidy by endorsement Only when prescribed for patients unable to swallow inappropriate and the prescription is endorsed accor	39.00	500 ml ts or whe	✓ Ro re calcium	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available Tab 2 mg Cap 2 mg	10.75	400 400	✓ No ✓ <u>Dia</u>	dia amide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Reta pharmacy	166.50	90 alid for 6		tocort CIR r applications meeting

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy		Fully	Brand or
(Manufacturer's Price)) :	Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROC	HLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg		100	✓ Asamax
Tab long-acting 500 mg	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g	118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml	41.30	7	✓ Pentasa
Suppos 500 mg		20	✓ Asacol
Suppos 1 g	50.96	28	✓ Pentasa
(Asamax Tab EC 500 mg to be delisted 1 March 2022)			
OLSALAZINE			
Tab 500 mg	93.37	100	✓ Dipentum
Cap 250 mg	53.00	100	✓ Dipentum

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	/	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	•	Nalcrom
SULFASALAZINE * Tab 500 mg Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CI	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00 Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	30 g OP 12	✓ Proctosedyl ✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE * Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE * Tab 135 mg	0.20	90	✓ Colofae
* Tab 100 IIIg	5.20	90	• Cololac

Antiulcerants

Antisecretory and Cytoprotective

MIS	SOPROSTOL STORE ST		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	Cytotec

Subsidy		Fully	Brand or	
(Manufacturer's Pr	rice)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Helicobacter Pylori Eradication

CLARITHROMYCIN

- a) Maximum of 28 tab per prescription
- Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.
 Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.

(Apo-Clarithromycin Tab 500 mg to be delisted 1 February 2022)

H2 Antagonists

	MOTIDINE - Only on a prescription	4.04	400	
*	Tab 20 mg	4.91	100	✓ Famotidine Hovid \$29
*	Tab 40 mg	8.48	100	✓ Famotidine Hovid S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	57.02	10	✓ Mylan S29
	Subsidy by endorsement – Subsidised for patients receiving to	reatment as par	t of palliative	e care.

Proton	Pump	Inhibitors
--------	------	------------

LANSOPRAZOLE		
* Cap 15 mg4.20 Lanzol Relief to be Principal Supply on 1 December 2021	100	✓ Lanzol Relief
* Cap 30 mg	100	✓ Lanzol Relief
OMEPRAZOLE		
For omeprazole suspension refer Standard Formulae, page 245		
* Cap 10 mg1.94	90	✓ Omeprazole actavis 10
* Cap 20 mg	90	✓ Omeprazole actavis 20
* Cap 40 mg3.11	90	✓ Omeprazole actavis 40
* Powder – Only in combination42.50 Only in extemporaneously compounded omeprazole suspension.	5 g	✓ Midwest
* Inj 40 mg ampoule with diluent	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
PANTOPRAZOLE		
* Tab EC 20 mg	100 100	 ✓ Panzop Relief ✓ Panzop Relief
Site Protective Agents		

COLLOIDAL BISMUTH SUBCITRATE			
Tab 120 mg	14.51	50	✓ Gastrodenol S29
SUCRALFATE			
Tab 1 g	35.50	120	
	(48.28)		Carafate

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Bile and Liver Therapy

RIFAXIMIN - Special Authority see SA1461 below - Retail pharmacy 56

Xifaxan

⇒SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Retail pharmacy

Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral lig 50 mg per ml	620.00	30 ml OP	✓ Proglycem S29

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit - Up to 5 kit available on a PSO......32.00 ✓ Glucagen Hypokit

✓ Protaphane Penfill

Insulin - Short-acting Preparations

INSULIN NEUTRAL

•	Inj human 100 u per ml25.26	10 ml OP	✓ Actrapid✓ Humulin R
•	Inj human 100 u per ml, 3 ml42.66	5	✓ Actrapid Penfill ✓ Humulin R

Insulin - Intermediate-acting Preparations

INSULIN ASPART WITH INSULIN ASPART PROTAMINE		
▲ Inj 100 iu per ml, 3 ml prefilled pen52.15	5	✓ NovoMix 30 FlexPen
INCLUDIT CODUANT		

IIV	JULIN ISOFTANE			
\blacktriangle	Inj human 100 u per ml	17.68	10 ml OP	Humulin NPH

					Fiolaphane
lack	Inj human 100 u per	ml, 3 ml	29.86	5	Humulin NPH

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic Manufacturer
SULIN ISOPHANE WITH INSULIN NEUTRAL	Ψ		That land to the l
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
nij naman war neatar maaiir 100 a per mi	25.20	10 1111 01	✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30
			✓ PenMix 40
			✓ PenMix 50
SULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			· ·
3 ml		5	✓ Humalog Mix 50
noulin Long acting Proporations			
nsulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml		1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓ Lantus SoloStar
nsulin - Rapid Acting Preparations			
SULIN ASPART			
Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	✓ NovoRapid FlexPen
SULIN GLULISINE			
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
SULIN LISPRO			•
Inj 100 u per ml, 10 ml	34 92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
· ·			-
Alpha Glucosidase Inhibitors			
CARBOSE			
Tab 50 mg		90	✓ Glucobay
	8.95		✓ Accarb
Accarb to be Principal Supply on 1 December 2021	2.42	00	(Observe)
Tab 100 mg		90	✓ Glucobay
Accarb to be Principal Supply on 1 December 2021	15.29		✓ Accarb
Alucobay Tab 50 mg to be delisted 1 December 2021) Alucobay Tab 100 mg to be delisted 1 December 2021)			
,			
Oral Hypoglycaemic Agents			
LIBENCLAMIDE			
Tab 5 mg	7.50	100	✓ Daonil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
GLICLAZIDE				
* Tab 80 mg	15.18	500	1	Glizide
GLIPIZIDE				
* Tab 5 mg	4.58	100	/	Minidiab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	8.63	1,000	/	Apotex
·	14.74		✓	Metformin Mylan
* Tab immediate-release 850 mg	7.04	500		Apotex
	11.28		✓	Metformin Mylan
(Apotex Tab immediate-release 850 mg to be delisted 1 March 2 PIOGLITAZONE	,	00	./	Veverene
* Tab 15 mg Vexazone to be Principal Supply on 1 January 2022	6.80	90	•	Vexazone
* Tab 30 mg	7.30	90	1	Vexazone
Vexazone to be Principal Supply on 1 January 2022				
* Tab 45 mg Vexazone to be Principal Supply on 1 January 2022	12.25	90	✓	Vexazone
,				
VILDAGLIPTIN Tab 50 mg	35.00	60	1	Galvus
		00	•	Gaivus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE	05.00	00	,	Calminat
Tab 50 mg with 1,000 mg metformin hydrochloride		60		Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	•	Galvumet

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*: or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

 a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN – Special Authority see SA2068 above – Retail pharmacy

	110to. 110t to be given in combination that a landed GET 1 agente.		
*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see SA2068 above – Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
	Tab 5 mg with 500 mg metformin hydrochloride	Tab 5 mg with 1,000 mg metformin hydrochloride	Tab 5 mg with 500 mg metformin hydrochloride 58.56 60 Tab 12.5 mg with 1,000 mg metformin hydrochloride 58.56 60

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips ______20.00 1 OP

CareSens Dual

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

biood glucose lest strips	Blood glucose test strips	26.20	50 test OP	✓ SensoCar
---------------------------	---------------------------	-------	------------	------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

IIVC	oblivi biv ivabbas iviaximum or 200 dev per presemptio	11		
*	29 g × 12.7 mm	10.50	100	✓ B-D Micro-Fine
*	31 g × 5 mm	11.75	100	✓ B-D Micro-Fine
*	31 g × 6 mm		100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDL	E - Maximum of 2	00 dev per p	rescription
*	Syringe 0.3 ml with 29 g × 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	B-D Ultra Fine
		1.30	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g x 8 mm needle	13.00	100	B-D Ultra Fine II
		1.30	10	
		(1.99)		B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year pe	riod.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
			✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

(MiniMed 640G Min basal rate 0.025 U/h to be delisted 1 January 2022)

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment: and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy: and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Fither
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump

Subsidy (Manufacturer's Price) Subsidised Subsidised Generic Manufacturer continued treatment; and 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 5 The patient is continuing to derive benefit from pump therapy; and 5 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 3 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription c) Maximum of 3 sets per p		ALIMENTAR	Y TR	ACT AND	METABOLISM
treatment; and 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient has adhered to and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 33 sets per prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		(Manufacturer's Price)		Subsidised	Generic
3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.3 Applicant is a relevant specialist; or 4.4 Applicant is a relevant specialist; or 4.5 Applicant is a relevant specialist; or 4.6 Applicant is a relevant specialist; or 4.7 Applicant is a relevant specialist; or 4.8 Applicant is a relevant specialist; or 4.9 Maximum of 3 sets per prescription 4 Donly on a prescription 5 Only on a prescription 6 Only on a prescription 7 Only on a prescription 8 Only on a prescription 9 Only on a prescription 10 Only on a prescription 11 Only on a prescription 12 Maximum of 13 infusion sets will be funded per year. 13 Only on a prescription 14 Only on a prescription 15 Only on a prescription 16 Maximum of 13 infusion sets will be funded per year. 17 Only on a prescription 18 MiniMed Sure-T 19 MiniMed Sure-T					
pump therapy; and 4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10	· · · · · · · · · · · · · · · · · · ·				
4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		ising analogue insulins	for at	least six mo	nths prior to initiating
5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tilock × 10	1 1 127	aranu and			
6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tilock × 10			80 mm	nal/mal an ni	ımn therany: and
7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10					
8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE – Special Authority see SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10					
8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	•		,		
Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	8.1 Applicant is a relevant specialist; or				
years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		•			
All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		from a relevant speciali	st or n	urse practiti	oner. Approvals valid for 2
1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10					
than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	3		l	ata ta ta a at a	III. Ada at a sucal ta an lasa
2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		tne treatment plan and	nas m	iaintained a	HDA1c of equal to or less
3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	,	mmol/mol from initial ar	nlicati	on and	
4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	•		•		ne: and
4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		, , , , , , , , , , , , , , , , , , ,			,
INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	4.1 Applicant is a relevant specialist; or				
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	4.2 Applicant is a nurse practitioner working within t	heir vocational scope.			
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	INSULIN PUMP CARTRIDGE - Special Authority see SA198	5 on page 19 – Retail p	harma	acv	
c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10				,	
Cartridge 300 U, t:lock × 10	b) Only on a prescription				
INSULIN PUMP INFUSION SET (STEEL CANNULA) — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10					
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	Cartridge 300 U, t:lock × 10	50.00	1 OP	✓ T	andem Cartridge
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	, , ,	al Authority see SA198	5 on p	age 19 – Re	etail pharmacy
c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	, , , , ,				
10 mm steel needle; 60 cm tubing × 10	, , , ,				
MMT-884A 10 mm steel needle; 80 cm tubing × 10130.00 1 OP ✓ MiniMed Sure-T		120.00	1 OD		liniMad Sura-T
10 mm steel needle; 80 cm tubing × 10130.00 1 OP ✓ MiniMed Sure-T	To min steer needle, oo chi tubing x 10	130.00	I OF	▼ IV	
·	10 mm steel needle: 80 cm tubing × 10	130.00	1 OP	✓ N	
mini ooon	3 · · · · · · · · · · · · · · · · · · ·			-	MMT-886A
6 mm steel needle; 60 cm tubing × 10	6 mm steel needle; 60 cm tubing x 10	130.00	1 OP	✓ N	liniMed Sure-T

MMT-864A

MMT-866A

MMT-874A

✓ MiniMed Sure-T

✓ MiniMed Sure-T

✓ MiniMed Sure-T

MMT-876A

✓ Sure-T MMT-863

✓ Sure-T MMT-873

1 OP

1 OP

1 OP

1 OP

1 OP

8 mm steel needle; 60 cm tubing × 10130.00

6 mm steel needle: 29 G: manual insertion: 60 cm tubing x

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
 \$	Per	•	Manufacturer	

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.1	00 1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 81 cm line x 10 with 10 needles130.	00 1 OP	✓ TruSteel

(TruSteel 6 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021) (TruSteel 8 mm steel cannula; straight insertion; 81 cm line × 10 with 10 needles to be delisted 1 November 2021)

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm blue tubing.......130.00

6 mm teflon needle, 80 cm clear tubing × 10130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
 ✓ MiniMed Silhouette
- MMT-381A

 MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
 10 P

 6 mm teflon needle, 45 cm blue tubing × 10
 130.00
 1 OP
- ✓ MiniMed Quick-Set MMT-398A
 ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
 - ✓ MiniMed Mio MMT-943A
 - ✓ MiniMed Mio MMT-923A
 - ✓ MiniMed Quick-Set MMT-399A
 - ✓ MiniMed Mio MMT-945A
 - ✓ MiniMed Mio MMT-965A
 - ✓ MiniMed Mio MMT-925A
 - ✓ MiniMed Quick-Set MMT-387A
 - ✓ MiniMed Quick-Set MMT-396A
 - ✓ MiniMed Quick-Set MMT-397A
 - ✓ MiniMed Mio MMT-975A
 - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH II	NSERTION	I DEVICE)	- Special Authority see
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	cm 140.00	1 OP	√ Διι	toSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cl	n	1 OP		toSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with	NSERTION) - Spe			
10 needles; luer lock	IT INSERTION WIT	1 OP H INSERT		houette MMT-373 ICE) – Special Authority
110 cm line × 10 with 10 needles	m	1 OP		toSoft 90
line x 10 with 10 needles		1 OP		toSoft 90 toSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m	1 OP		toSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy a) Maximum of 3 sets per prescription		-		
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock	130.00	1 OP	√ Qu	ick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock		1 OP	√ Qu	ick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 c a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per	n page 19 – Retail _I	oharmacy	-	
10 × luer lock conversion cartridges 1.8 ml for Paradigm pun Cartridge for 5 and 7 series pump; 1.8 ml × 10	ps50.00	1 OP 1 OP	✔ Mir 1	R Cartridge 1.8 niMed .8 Reservoir //MT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		niMed 8.0 Reservoir //MT-332A

Fully

Brand or

	(Manufacturer's Price) Sub Per	osidised	Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					_
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Cr	eon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))	*	100	✓ Pa	nzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Cr	eon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph		00 00			
Eur U)		20 g OP	✓ Cr	eon Micro	
URSODEOXYCHOLIC ACID – Special Authority see SA1739 be	•	•			
Cap 250 mg	32.95	100	✓ <u>Ur</u> s	sosan	

Subsidy

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- ner:
 - 1 Patient has been diagnosed with Alagille syndrome; or
 - 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Su	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	dised	Generic
	\$ Per	•	Manufacturer

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soln	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
•	(17.32)	-	Normacol Plus
	2.41	200 g OP	
	(8.72)	•	Normacol Plus

Faecal Softeners

*	Tab 50 mg	.2.31	100	✓ Coloxyl
	Tab 120 mg		100	✓ Coloxyl
	CUSATE SODIUM WITH SENNOSIDES			
*	Tab 50 mg with sennosides 8 mg	.3.10	200	✓ Laxsol
	OXAMER - Only on a prescription			
	Not funded for use in the ear.			
*	Oral drops 10%	.3.98	30 ml OP	Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below – Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
,	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$	e) Sul Per	Fully osidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL * Suppos 3.6 g – Only on a prescription LACTULOSE – Only on a prescription	9.25	20	√ P	SM
* Oral liq 10 g per 15 ml	3.33	500 ml	√ L	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIG	CARBONATE AND	SODIUM	CHLORII	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7	O,	30	✓ N	<u>lolaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	√ F	leet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	•		Parlana
5 ml	29.98	50	✓ <u>IV</u>	<u>licolette</u>
Stimulant Laxatives				
BISACODYL — Only on a prescription * Tab 5 mg * Suppos 10 mg Lax-Suppositories to be Principal Supply on 1 December	3.69	200 10	_	ax-Tab ax-Suppositories
SENNA – Only on a prescription * Tab, standardised	2.17 (8.21)	100	S	enokot
	0.43 (2.06)	20	Q	enokot
SODIUM PICOSULFATE - Special Authority see SA2053 below	` '		3	GIIOROL
Oral soln 7.5 mg per ml		30 ml OP	✓ D	ulcolax SP Drop

SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE	 Special Authority 	see SA2042 held	ow – Retail pharmacy

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 below - Retail pharmacy

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
----------------------------------------------------------------------------------------	--

continued...

- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - F	Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price)		Fully lised	Brand or Generic
	\$	Per	1	Manufacturer
SAPROPTERIN DIHYDROCHLORIDE - Special Authority see S	A1989 below – Reta	il pharmacy		
Tab soluble 100 mg	1,452.70	30 OP	1	Kuvan

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see S	A1599 below – Retail pharmacy		
Soln 100 ma per ml	CBS	100 ml	✓ Amzoate S29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE − Special Authority see SA1990 below − Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP

✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 on the next page -	 Retail pharmacy 		
Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

✓ Elelvso

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 Pharmac PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, 6) thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g., hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
` \$	Per	•	

continued...

All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 7) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with

BENZYDAMINE HYDROCHLORIDE

Odil 0.1070 Trigilar subsidy of \$20.01 per 500 fill with			
Endorsement	9.00	500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	s oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	J	Orabase
	1.52	5 g OP	
	(3.60)	- 9	Orabase
Powder	٠,	28 g OP	
	(10.95)	_0 g 0.	Stomahesive
CHOLINE CALICYLATE WITH OFTALKONILINA CHI OPIDE	(13133)		
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE	0.00	45 - 00	
* Adhesive gel 8.7% with cetalkonium chloride 0.01%		15 g OP	Davida
	(6.00)		Bonjela
TRIAMCINOLONE ACETONIDE			
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			3
Oral gel 20 mg per g	4.74	40 a OB	✓ Decozol
Decozol to be Principal Supply on 1 December 2021	4.74	40 g OP	♥ Decozoi
,			
NYSTATIN			
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pric \$		ully Brand or sed Generic Manufacturer		
Other Oral Agents					
For folinic mouthwash, pilocarpine oral liquid or saliva substitute	formula refer Stand	ard Formulae,	page 245		
THYMOL GLYCERIN			4 		
* Compound, BPC(PSM Compound, BPC to be delisted 1 February 2022)	9.15	500 ml	✓ PSM		
Vitamins					
Vitamin B					
HYDROXOCOBALAMIN			_		
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PS	SO1.89		✓ Neo-B12 ✓ Vita-B12		
	3.15		✓ Hydroxocobalamin		
			Mercury Pharma		
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose					
b) Only on a prescription					
* Tab 25 mg – No patient co-payment payable			✓ <u>Vitamin B6 25</u>		
Tab 50 mg THIAMINE HYDROCHLORIDE – Only on a prescription	13.03	500	✓ Apo-Pyridoxine		
* Tab 50 mg	7.09	100	✓ Max Health		
VITAMIN B COMPLEX			4 - .		
* Tab, strong, BPC	/.15	500	✓ Bplex		
Vitamin C					
ASCORBIC ACID					
a) No more than 100 mg per dose b) Only on a prescription					
* Tab 100 mg	9.90	500	✓ <u>Cvite</u>		
Vitamin D					
ALFACALCIDOL					
* Cap 0.25 mcg			✓ One-Alpha		
* Cap 1 mcg * Oral drops 2 mcg per ml			✓ One-Alpha ✓ One-Alpha		
CALCITRIOL					
* Cap 0.25 mcg			✓ Calcitriol-AFT		
* Cap 0.5 mcg COLECALCIFEROL	13./5	100	✓ <u>Calcitriol-AFT</u>		
* Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per prescript		12	✓ <u>Vit.D3</u>		
* Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	4.8 ml OP	✓ Puria		
Multivitamin Preparations					
MULTIVITAMIN RENAL - Special Authority see SA1546 on the			(a) 11 - 11		
* Cap	6.49	30	✓ Clinicians Renal Vit		

ALIMENTARY TRACT AND METABOLISM						
	Subsidy (Manufacturer's Price \$) Subsi Per	Fully dised	Brand or Generic Manufacturer		
■ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either: 1 The patient has chronic kidney disease and is receiving ei 2 The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m² body surface area (BSA).	ther peritoneal dialy	sis or haem	odialys	sis; or		
MULTIVITAMINS – Special Authority see SA1036 below – Retail * Powder	72.00 2		notifie			
* Tab (BPC cap strength) * Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy		1,000	✓ <u>N</u>	Ivite itabdeck		
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut s 3 Patient has severe malabsorption syndrome.	d without further ren					
Minerals Calcium						
CALCIUM CARBONATE	0.00	050		ala: Tab 500		

CALCIUM CARBONATE		
* Tab 1.25 g (500 mg elemental)	250	✓ Calci-Tab 500
* Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement52.00	20	✓ Calcium-Sandoz
		Forte S29
54.60	76	✓ Cacit S29
Subsidy by endorsement – Only when prescribed for paediatric patients (considered unsuitable.	< 5 years) wh	ere calcium carbonate oral liquid is
CALCIUM GLUCONATE		
* Inj 10%, 10 ml ampoule32.00	10	✓ Max Health -
, , , , , , , , , , , , , , , , , , , ,		Hameln S29
64.00	20	✓ Max Health S29
Fluoride		
SODIUM FLUORIDE		
* Tab 1.1 mg (0.5 mg elemental)	100	✓ PSM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	euroTabs
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.09	100	✓ Fe	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	✓ Fe	erro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) — Special Authority so Inj 50 mg per ml, 10 ml vial		etail p 1		erinject

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

✓ Ferrosia

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
Magnesium				
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml		nillips Milk of Magnesia 829
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ <u>M</u>	artindale
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ <u>Zi</u>	ncaps

Subsidy
(Manufacturer's Price)
\$ Per

Fully Subsidised

Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price)		Fully Subsidised	Generic
	<u> </u>	Per		Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previous	s page – Retail pharm	nacv		
Wastage claimable	, page 1121am p.1am.	,		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	1	Binocrit
		6		Binocrit
Inj 2,000 iu in 1 ml, syringe		-		
Inj 3,000 iu in 0.3 ml, syringe		6		Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6		Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	/	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	✓	Binocrit
Inj 10,000 iu in 1 ml, syringe		6		Binocrit
Inj 40,000 iu in 1 ml, syringe		1	_	Binocrit
11] 40,000 td 111 1 1111, 0y1111go	200.00	•	•	<u>Diniodrit</u>
Magalahlastia				
Megaloblastic				
FOLIC ACID				
	21.04	1,000	./	Apo-Folic Acid
Tab 0.8 mg		,		•
* Tab 5 mg		100		Folic Acid Mylan
	12.12	500	•	Apo-Folic Acid
Folic Acid Mylan to be Sole Supply on 1 December 2021				
Oral liq 50 mcg per ml	26.00 25	5 ml Ol	•	Biomed
(Apo-Folic Acid Tab 5 mg to be delisted 1 December 2021)				

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia

 Inj 250 iu vial.
 612.50
 1
 ✓ Alprolix

 Inj 500 iu vial.
 1,225.00
 1
 ✓ Alprolix

 Ini 1.000 iu vial.
 2.450.00
 1
 ✓ Alprolix

✓ Revolade

28

 Inj 1,000 iu vial.
 2,450.00
 1
 ✓ Alprolix

 Inj 2,000 iu vial.
 4,900.00
 1
 ✓ Alprolix

 Inj 3,000 iu vial.
 7,350.00
 1
 ✓ Alprolix

ELTROMBOPAG - Special Authority see SA1743 below - Retail pharmacy

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist.

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

4 T....

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial	1	✓ Hemlibra
Inj 150 mg in 1 ml vial	1	Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

continued...

(1)	Subsidy Manufacturer's Price)	Fully Subsidised	
	\$	Per 🗸	Manufacturer

continued...

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months;
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1	✓ NovoSeven RT
Inj 2 mg syringe2,356.60		✓ NovoSeven RT
Inj 5 mg syringe		✓ NovoSeven RT
Inj 8 mg syringe9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 Ŭ	1,315.00	1	✓ FĚIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Ini 2.500 U	6.575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Subject to officia.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe		1	✓ Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

	Subsidy	F	ully Brand or
	(Manufacturer's Price)	Subsidis	
	\$	Per	✓ Manufacturer
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharm]			
For patients with haemophilia. Access to funded treatment is	managed by the Hae	emophilia Tr	eaters Group in conjunction
with the National Haemophilia Management Group. Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		-	✓ RIXUBIS
Inj 2,000 iu vial		=	✓ RIXUBIS
Inj 3,000 iu vial		=	✓ RIXUBIS
• •	•	•	· III/ODIO
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – [For patients with haemophilia. Preferred Brand of short half-		vr \/III \\ A 0000	s to funded treatment is
managed by the Haemophilia Treaters Group in conjunction v			
Inj 250 iu vial			✓ Advate
Inj 500 iu vial			✓ Advate
Inj 1,000 iu vial		-	✓ Advate
Inj 1,500 iu vial		-	✓ Advate
Inj 2,000 iu vial		•	✓ Advate
Inj 3,000 iu vial			✓ Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	•	•	7147410
For patients with haemophilia. Rare Clinical Circumstances I		, rocombinor	ot factor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in	conjunction with the r	vational Hae	mopnilia Management Group,
subject to criteria.	227 50	1	√ Kaganata ES
Inj 250 iu vial		-	✓ Kogenate FS
Inj 500 iu vial		-	✓ Kogenate FS
Inj 1,000 iu vial Inj 2,000 iu vial			✓ Kogenate FS ✓ Kogenate FS
Inj 3,000 iu vial	,		✓ Kogenate FS
		1	Nogeliale F3
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]			
For patients with haemophilia A receiving prophylaxis treatme		d treatment is	s managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			
Inj 250 iu vial			✓ Adynovate
Inj 500 iu vial			✓ Adynovate
Inj 1,000 iu vial			✓ Adynovate
Inj 2,000 iu vial	2,400.00	1	✓ Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml	28.50	5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	9.45	60	✓ Mercury Pharma
Vitamin K			
DI IVTOMENIA DIONE			
PHYTOMENADIONE	0.00	_	/ Kamaldan MM
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO		_	✓ Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO	9.21	5	✓ Konakion MM
Antithrombotic Agents			
Antitinombotic Agents			
Antiplotolot Agento			
Antiplatelet Agents			
ASPIRIN			
* Tab 100 mg	10.80	990	✓ Ethics Aspirin EC
		-	

	Subsidy (Manufacturer's Price) \$	Per	
CLOPIDOGREL			
* Tab 75 mg	4.60	84	✓ <u>Clopidogrel</u> <u>Multichem</u>
DIPYRIDAMOLE			
* Tab long-acting 150 mg	10.90	60	✓ Pytazen SR
TICAGRELOR - Special Authority see SA1955 below - Retail pha	armacy		
* Tab 90 mg	90.00	56	✓ Brilinta
SA1055 Special Authority for Subsidy			

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Fither:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe		10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe	125.87	10	 Clexane Forte
Inj 150 mg in 1 ml syringe		10	 Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	
HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ DBL Heparin
, -, ,			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
, =0,000 14 por, 01=	42.40	ŭ	✓ Heparin DBL S29
LIEDADINICED CALINE	72.70		Tiopaini bbe
HEPARINISED SALINE	GE 10	50	✓ Pfizer
Inj 10 iu per ml, 5 ml	05.48	ου	♥ Pilzer
Oral Anticoagulants			
Oral Anticoagalants			
DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ Pradaxa
Cap 110 mg	76.36	60	✓ Pradaxa
Cap 150 mg	76.36	60	✓ Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ Xarelto
Tab 20 mg		28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	✓ Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	✓ Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg		100	✓ Marevan
* Tab 5 mg		50	✓ Coumadin
ŭ	11.48	100	✓ Marevan
Blood Colony-stimulating Factors			
FILGRASTIM - Special Authority see SA1259 below - Retail p	,	10	Nivestim
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Nivestim to be Principal Supply on 1 December 2021			_

✓ Nivestim 10 Nivestim to be Principal Supply on 1 December 2021

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price)	s	Fully ubsidised	Brand or Generic
	\$	Per	•	Manufacturer
PEGFILGRASTIM – Special Authority see SA1912 below – Retail pharmacy				
Inj 6 mg per 0.6 ml syringe	1,080.00	1	✓ No	eulastim
⇒SA1912 Special Authority for Subsidy				

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration			
GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO POTASSIUM CHLORIDE		5 1	✓ Biomed ✓ Biomed
* Inj 75 mg per ml, 10 ml	55.00	50	 ✓ AstraZeneca ✓ Potassium Chloride Aguettant \$29
	65.00		✓ Juno
(AstraZeneca Inj 75 mg per ml, 10 ml to be delisted 1 November 20	121)		
(Potassium Chloride Aguettant \$29 Inj 75 mg per ml, 10 ml to be of	delisted 1 Nove	mber 2021)	
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
a) Up to 5 inj available on a PSOb) Not in combination			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebuliser u	se except wher	n used in conji	unction with an antibiotic intended
for nebuliser use. Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.00	500 ml	✓ Baxter
III] 0.9 %, day = 0p to 2000 fill available off a F30	1.26	1.000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, mate		,	
for emergency use. (500 ml and 1,000 ml packs)	, p		рашен, ет ет ет
Inj 23.4% (4 mmol/ml), 20 ml ampoule	33.00	5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standard	Formulae, page	e 245	
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	✓ Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO		50	Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓ Fresenius Kabi

TOTAL PARENTERAL NUTRITION (TPN)

Infusion......CBS

1 OP

✓ TPN

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule - Up to 5 inj available on a PSO7.19	50	✓ Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO5.00	20	✓ Fresenius Kabi
		✓ Multichem

Oral Administration		
CALCIUM POLYSTYRENE SULPHONATE Powder169.85 COMPOUND ELECTROLYTES	300 g OP	✓ Calcium Resonium
Powder for oral soln — Up to 5 sach available on a PSO9.77	50	✓ <u>Electral</u>
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)6.55	1,000 ml OP	✓ Pedialyte - Bubblegum
PHOSPHORUS		
Tab eff 500 mg (16 mmol)82.50	100	Phosphate Phebra
POTASSIUM CHLORIDE		
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)5.26	60	
(11.85)		Chlorvescent
* Tab long-acting 600 mg (8 mmol)8.90	200	✓ Span-K
SODIUM BICARBONATE		
Cap 840 mg8.52	100	✓ Sodibic
		✓ Sodibic
SODIUM POLYSTYRENE SULPHONATE		
Powder	454 a OP	✓ Resonium-A

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
΄ ΄ ΄	Dox /	Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DC		

* Tab 2 mg	17.35	500	✓ Apo-Doxazosin
* Tab 4 mg	20.94	500	✓ Apo-Doxazosin
PHENOXYBENZAMINE HYDROCHLORIDE			
₩ Can 10 mg	65.00	30	✓ RNM 929

PRAZOSIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking prazosin prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of prazosin.

216.67

100

Dibenzyline S29

	1 0 1		
*	Tab 1 mg5.53	100	✓ Apo-Prazosin
*	Tab 2 mg7.00	100	✓ Apo-Prazosin
	Tab 5 mg	100	✓ Apo-Prazosin

(Apo-Prazosin Tab 1 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 2 mg to be delisted 1 May 2022) (Apo-Prazosin Tab 5 mg to be delisted 1 May 2022)

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

*	Oral liq 5 mg per ml94.99	95 ml OP	✓ Capoten
	135.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

(Captopril-Mylan S29 Oral liq 5 mg per ml to be delisted 1 January 2022)

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

*	Tab 0.5 mg2.09	9	90 •	✓ Zapril
*	Tab 2.5 mg4.8	0	90 •	Zapril
	Tab 5 mg		90 •	Zapril
ENA	LAPRIL MALEATE			
*	Tab 5 mg1.8	2 .	100 •	/ Acetec
*	Tab 10 mg2.0	2 .	100	Acetec
*	Tab 20 mg2.4	2 .	100	Acetec
LISII	NOPRIL			
*	Tab 5 mg2.0	7	90 •	Ethics Lisinopril
*	Tab 10 mg2.3	6	90 •	Ethics Lisinopril
*	Tab 20 mg	7	90 •	Ethics Lisinopril
PER	INDOPRIL			
	Tab 2 mg4.9	5	30	✓ Coversyl
	Tab 4 mg6.3		30	Coversyl

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	Generic Manufacturer
QUINAPRIL				
★ Tab 5 mg Tab 10 mg		90 90		Arrow-Quinapril 5 Arrow-Quinapril 10
≰ Tab 20 mg		90		Arrow-Quinapril 20
ACE Inhibitors with Diuretics				
QUINAPRIL WITH HYDROCHLOROTHIAZIDE				
Tab 10 mg with hydrochlorothiazide 12.5 mg		28		Accuretic
★ Tab 20 mg with hydrochlorothiazide 12.5 mg	4.10	30 30		Accuretic 10 Accuretic 20
Accuretic Tab 10 mg with hydrochlorothiazide 12.5 mg to be del		30	•	Accuretic 20
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
₭ Tab 4 mg	2.00	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021	0.00	00	,	Odd
Tab 8 mg Candestar to be Principal Supply on 1 December 2021	2.28	90	•	Candestar
* Tab 16 mg	3.31	90	1	Candestar
Candestar to be Principal Supply on 1 December 2021				
₹ Tab 32 mg	5.26	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021				
OSARTAN POTASSIUM	4.50			
₹ Tab 12.5 mg ₹ Tab 25 mg		84 84		Losartan Actavis Losartan Actavis
k Tab 25 mg		84		Losartan Actavis
₭ Tab 100 mg		84		Losartan Actavis
Angiotensin II Antagonists with Diuretics				
OSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE				
Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	•	Arrow-Losartan & Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:

continued...

Subsidy (Manufacturer's Price)	Fu Subsidise	,	nd or neric
\$	Per •	/ Mai	nufacturer

continued...

- 2.1 Patient is in NYHA/WHO functional class II; or
- 2.2 Patient is in NYHA/WHO functional class III; or
- 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaestr AMIODARONE HYDROCHLORIDE	netics, Local, pa	ge 122	
▲ Tab 100 mg	3.80	30	✓ Aratac
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a		00	- <u>Alutuo</u>
PSO	16 37	10	✓ Max Health
	10.57	10	WIAX HEAILH
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a			
PSO		10	✓ Hameln S29
	15.09		Martindale
Martindale to be Principal Supply on 1 January 2022			
(Hameln S29 Inj 600 mcg per ml, 1 ml ampoule to be delisted 1 Ja	anuary 2022)		
DIGOXIN			
* Tab 62.5 mcg - Up to 30 tab available on a PSO	7.00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO	15.20	240	✓ Lanoxin
* Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			 Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	✓ Rythmodan
FLECAINIDE ACETATE	20.07		,
▲ Tab 50 mg	10.05	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg		90	✓ Flecainide Bivim
ap long-acting rooming	39.31	90	Controlled
			Release Teva
▲ Cap long-acting 200 mg	61.06	90	✓ Flecainide
Cap long-acting 200 mg	01.00	90	Controlled
			Release Teva
Ini 10 ma par ml. 15 ml ampaula	100.00	_	
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor

Subsidy (Manufacturer's Price) \$162.00	Per 100	✓	I Generic
		✓	Mexiletine Hydrochloride USP \$29
		✓	Mexiletine Hydrochloride USP \$29
202.00	100		Toyo 620
		•	Mexiletine Hydrochloride USP \$29
		•	Teva S29
January 2022) January 2022)			
40.90	50	•	Rytmonorm
macy			
53.00 79.00	100 100		Gutron Gutron
for 2 years where pay pwards as necessar	y. H	ypertensio	on should be avoided, and
r .	macy53.0079.00 for 2 years where prowards as necessar	macy 53.00 100 100 for 2 years where patient owards as necessary. Hy	January 2022) January 2022)40.90 50 macy53.00 100 for 2 years where patient has disab

Beta Adrenoce	ptor Blo	ockers
----------------------	----------	--------

ATENOLOL		
* Tab 50 mg9.33 Mylan Atenolol to be Principal Supply on 1 January 2022	500	✓ Mylan Atenolol
* Tab 100 mg	500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml	300 ml OP	 ✓ Atenolol AFT ✓ Atenolol AFT S29 S29
38.20		✓ Essential Generics S29
Restricted to children under 12 years of age.		
BISOPROLOL FUMARATE		
* Tab 2.5 mg	90	✓ Bisoprolol Mylan
* Tab 5 mg	90	✓ Bisoprolol Mylan
* Tab 10 mg 3.62	90	✓ Bisoprolol Mylan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	1	Manufacturer
CARVEDILOL				
* Tab 6.25 mg	2.24	60	✓	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	✓	Carvedilol Sandoz
* Tab 25 mg	2.95	60	✓	Carvedilol Sandoz
ABETALOL				
* Tab 100 mg	14.50	100	✓	Trandate
★ Tab 200 mg	27.00	100	✓	Trandate
k Inj 5 mg per ml, 20 ml ampoule	59.06	5		
, ,	(88.60)			Trandate
k inj 5 mg per ml, 20 ml vial	42.29	1		
•	(48.20)			Alvogen S29
METOPROLOL SUCCINATE				
★ Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
★ Tab long-acting 47.5 mg		30	1	Betaloc CR
★ Tab long-acting 95 mg		30	1	Betaloc CR
★ Tab long-acting 190 mg		30	1	Betaloc CR
METOPROLOL TARTRATE				
Tab 50 mg	5.66	100	1	Apo-Metoprolol
745 00 mg		.00		IPCA-Metoprolol
Tab 100 mg	7 55	60		Apo-Metoprolol
7 ab 700 mg		00		IPCA-Metoprolol
★ Tab long-acting 200 mg	23.40	28		Slow-Lopresor
k Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
Apo-Metoprolol Tab 50 mg to be delisted 1 March 2022)	20.00	ŭ		
Apo-Metoprolol Tab 100 mg to be delisted 1 March 2022)				
IADOLOL				
Tab 40 mg	16.69	100	/	Apo-Nadolol
	19.19			Nadolol BNM S29
Tab 80 mg		100		Apo-Nadolol
140 00 119	30.39	100		Nadolol BNM S29
Apo-Nadolol Tab 40 mg to be delisted 1 March 2022) Apo-Nadolol Tab 80 mg to be delisted 1 March 2022)	00.09		•	HAUGIOI DINIVI 1023
PINDOLOL – Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who we endorsed accordingly. Pharmacists may annotate the pre dispensing of pindolol.				
taspensing or pindoloi. ★ Tab 5 mg	13 22	100	1	Apo-Pindolol
* Tab 10 mg		100		Ano-Pindolol

	dioponioning of pindolon		
*	Tab 5 mg13.22	100	✓ Apo-Pindolol
*	Tab 10 mg23.12	100	✓ Apo-Pindolol
	Tab 15 mg33.31	100	✓ Apo-Pindolol

(Apo-Pindolol Tab 5 mg to be delisted 1 May 2022)

(Apo-Pindolol Tab 10 mg to be delisted 1 May 2022)

(Apo-Pindolol Tab 15 mg to be delisted 1 May 2022)

52

	Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
PROPRANOLOL				
Tab 10 mg	4.64	100	✓ A	po-Propranolol
Ÿ	7.04		✓ D	rofate
Tab 40 mg	5.72	100	✓ A	po-Propranolol
· ·	8.75		✓ II	PCA-Propranolol
★ Cap long-acting 160 mg	18.17	100	✓ 0	ardinol LA
FOral lig 4 mg per ml - Special Authority see SA				
Retail pharmacy		500 ml	✓ R	Roxane- Propranolol S29

(Apo-Propranolol Tab 10 mg to be delisted 1 March 2022) (Apo-Propranolol Tab 40 mg to be delisted 1 March 2022)

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTAL OL

AND ODIDING

*	Tab 80 mg	.32.58	500	Mylan
	Tab 160 mg			Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑI	ILODIPINE			
*	Tab 2.5 mg	1.08	90	✓ Vasorex
*	Tab 5 mg	0.96	90	✓ Vasorex
*	Tab 10 mg	1.19	90	✓ Vasorex
FE	ELODIPINE			
*	Tab long-acting 2.5 mg	1.45	30	Plendil ER
	Tab long-acting 5 mg		90	✓ Felo 5 ER
	Felo 5 ER to be Principal Supply on 1 January 2022			
*	Tab long-acting 10 mg	4.32	90	✓ Felo 10 ER
	Felo 10 ER to be Principal Supply on 1 January 2022			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
IFEDIPINE				
Fab long-acting 10 mg	18.80	56	✓	Tensipine MR10 S29
Tab long-acting 20 mg	9.12	50	✓	Mylan (12 hr
				release) S29
	17.72	100	1	Nyefax Retard
Fab long-acting 30 mg	4.78	14	1	Mylan Italy (24 hr
				release) S29
	34.10	100	1	Mylan (24 hr
				release) S29
Tab long-acting 60 mg	52.81	100	1	Mylan (24 hr
				release) \$29
Other Calcium Channel Blockers				
ILTIAZEM HYDROCHLORIDE				
← Tab 60 mg	8.50	100	1	Dilzem
Cap long-acting 120 mg	33.42	500		Apo-Diltiazem CD
Cap long-acting 180 mg	7.00	30	1	Cardizem CD
	50.05	500		Apo-Diltiazem CD
Cap long-acting 240 mg	9.30 66.76	30 500		Cardizem CD Apo-Diltiazem CD
Dilzem Tab 60 mg to be delisted 1 January 2022) Apo-Diltiazem CD Cap long-acting 180 mg to be del Apo-Diltiazem CD Cap long-acting 240 mg to be del ERHEXILINE MALEATE				
€ Tab 100 mg	62 90	100	1	Pexsig
ERAPAMIL HYDROCHLORIDE				
Tab 40 mg	7.01	100	1	Isoptin
€ Tab 80 mg		100		Isoptin
€ Tab long-acting 120 mg		100		Isoptin Retard \$29
rab long-acting 120 mg	30.02	100		Isoptin SR
Fab long-acting 240 mg	15 12	30		Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj av		00	•	isopiiii ori
PSO		5	1	Isoptin
Centrally-Acting Agents				
LONIDINE				
RECITIONNE Fatch 2.5 mg, 100 mcg per day − Only on a pre	scription 10.34	4	1	Mylan
Patch 5 mg, 200 mcg per day – Only on a presc		4		Mylan
Patch 7.5 mg, 300 mcg per day — Only on a pre	•	4		Mylan
0. 01 , ,				<u>y</u>
LONIDINE HYDROCHLORIDE	0 75	112	.1	Clonidine BNM
4 Tab 0E mag				
	27 07	1()()		
F Tab 150 mcg		100	•	Catapres
Tab 25 mcg Tab 150 mcg Catapres to be Principal Supply on 1 Janual Inj 150 mcg per ml, 1 ml ampoule	ry 2022	100		Medsurge

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	I Generic
METHYLDOPA	Ψ	1 01		Managaror
★ Tab 250 mg	15.10 52.85	100 500		Methyldopa Mylan Methyldopa Mylan S29 S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
₭ Tab 1 mg	4.91 16.36	30 100		Burinex S29 S29 Burinex
k Inj 500 mcg per ml, 4 ml vial	7.95	5	•	Burinex
Tab 40 mg - Up to 30 tab available on a PSO	8.00	1,000	1	Apo-Furosemide IPCA-Frusemide
★ Tab 500 mg	25.00 89.48	50		Urex Forte Furosemid- Ratiopharm S29
	169.96	100	•	Furosemid- Ratiopharm ©29
Oral liq 10 mg per ml		30 ml C		Lasix
k Inj 10 mg per ml, 25 ml ampoule k Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a l Apo-Furosemide Tab 40 mg to be delisted 1 March 2022)		6 5		<u>Lasix</u> <u>Furosemide-Baxter</u>
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml		25 ml C)P ✓	Biomed
Tab 50 mg	•	30	1	Inspra
Tab 25 mg Special Authority for Subsidy		30		Inspra
nitial application from any relevant practitioner. Approvals val ne following criteria: oth:	d without further ren	ewal u	nless notif	ied for applications meeti
Patient has heart failure with ejection fraction less than 40 Either:	0%; and			
2.1 Patient is intolerant to optimal dosing of spironolac2.2 Patient has experienced a clinically significant adv		optima	ıl dosing o	f spironolactone.
METOLAZONE				
Tab 5 mg	CBS	1 50		Metolazone S29 Zaroxolyn S29
SPIRONOLACTONE				•

* Tab 100 mg11.80

✓ Spiractin

✓ Spiractin

✓ Biomed

100

100

25 ml OP

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Out state		F. II.	December 1
1	Subsidy (Manufacturer's Price \$) ;	Fully Subsidised	
Potassium Sparing Combination Diuretics				
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE			_	
★ Tab 5 mg with furosemide 40 mg		28	•	Frumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID * Tab 5 mg with hydrochlorothiazide 50 mg		50	✓	Moduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
★ Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	•	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge	•			
₭ Tab 5 mg	34.55	500	•	Arrow- Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	26.00 2	25 ml O	P 🗸	Biomed
HLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	3 00	30	,	Igroton \$29
740 23 Hg	6.50	50		Hygroton
NDAPAMIDE				
F Tab 2.5 mg	10.45 11.61	90 100		<u>Dapa-Tabs</u> Mylan
	11.01	100	·	Indapamide \$29
Lipid-Modifying Agents				
Fibrates				
EZAFIBRATE				
 ₹ Tab 200 mg ₹ Tab long-acting 400 mg 		90 30		Bezalip Bezalip Retard
	21.21	30	•	bezanp netaru
Other Lipid-Modifying Agents				
CIPIMOX Cap 250 mg	21 56	30	,	Olbetam
	21.50	50		Olbetam S29 S29
Resins				
OLESTIPOL HYDROCHLORIDE	00.00	00		0-1
Grans for oral liq 5 g	32.89	30	•	Colestid

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg Lorstat to be Principal Supply on 1 December 2021	6.16	500	✓	Lorstat
* Tab 20 mg	9.24	500	1	Lorstat
* Tab 40 mg	14.92	500	1	Lorstat
* Tab 80 mg Lorstat to be Principal Supply on 1 December 2021	26.54	500	•	Lorstat
PRAVASTATIN				
* Tab 20 mg	2.11	28	✓	Pravastatin Mylan
* Tab 40 mgSIMVASTATIN	3.61	28	•	Pravastatin Mylan
* Tab 10 mg	1.23	90	✓	Simvastatin Mylan
* Tab 20 mg		90	✓	Simvastatin Mylan
* Tab 40 mg		90	✓	Simvastatin Mylan
* Tab 80 mg		90	✓	Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors				
EZETIMIBE - Special Authority see SA1045 below - Retail ph	armacy			

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and

- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

30

✓ Ezetimibe Sandoz

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

30 ✓ Zimybe	30	5.15	Tab 10 mg with simvastatin 10 mg.
30 ✓ Zimybe	30	6.15	Tab 10 mg with simvastatin 20 mg.
30 ✓ Zimybe	30	7.15	Tab 10 mg with simvastatin 40 mg.
30 ✓ Zimybe	30	8.15	Tab 10 mg with simvastatin 80 mg.

Subsidy		Fully	Brand or
(Manufacturer's \$	Price)	Subsidised <	Generic Manufacturer
⇒SA1046 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid for 2 years fo All of the following:	r application	is meeting th	ne following criteria:
Patient has a calculated absolute risk of cardiovascular disease of at lea	st 15% over	5 year; and	
2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and			
3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/ atoryastatin.	litre with the	use of the r	naximal tolerated dose of
Notes: A patient who has failed to reduce their LDL cholesterol to less than or e	egual to 2.0	mmol/litre w	ith the use of a less poten
statin should use a more potent statin prior to consideration being given to the u	ise of non-st	atin therapie	es.
Other treatment options including fibrates, resins and nicotinic acid should be co			
f a patient's LDL cholesterol cannot be calculated because the triglyceride level performed and if the LDL cholesterol again cannot be calculated then it can be or			
2.0 mmol/litre.	301101001001	nat the LBL	onologici or lo groater trial
Renewal from any relevant practitioner. Approvals valid for 2 years where the t	reatment re	mains appro	priate and the patient is
			p
penefiting from treatment.			F
Nitrates			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Nitrates GLYCERYL TRINITRATE			
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose			
Nitrates GLYCERYL TRINITRATE	250 dose		itrolingual Pump Spray
Nitrates SLYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30	OP ✔ N	itrolingual Pump Spray itroderm TTS
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO		OP ✔ N	itrolingual Pump Spray
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30	OP	itrolingual Pump Spray itroderm TTS
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose	30 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS
Nitrates GLYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose − Up to 250 dose available on a PSO	30 30 100	OP	itrolingual Pump Spray itroderm TTS itroderm TTS
Nitrates SILYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose − Up to 250 dose available on a PSO	30 30 100 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30 90	OP VNVNVN	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard uride
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30	OP VNVNVNVN	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30 90	OP V N V N V N V N N V N N N N N N N N N	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard uride

ΗY	DR	Al	LAZII	NE F	IYD	R
	_				_	

Vasodilators

HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 on the next page -		
	Retail pharmacyCBS	1	Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink S29
*	lnj 20 mg ampoule25.90	5	✓ Apresoline

Subsidy (Manufacturer's Price)	F Subsidis	ully	Brand or Generic
(Manufacturer's Price)	Per	√	Manufacturer

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

MINIOVIDII

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIE ▲ Tab 10 mg70.00	100	✓ Loniten
NICORANDIL ▲ Tab 10 mg 25.57 ▲ Tab 20 mg 32.28	60 60	✓ <u>Ikorel</u> ✓ <u>Ikorel</u>
PAPAVERINE HYDROCHLORIDE * Inj 12 mg per ml, 10 ml ampoule	5	✓ Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg42.26	50	✓ Trental 400

Endothelin Receptor Antagonists

AMBRISENTAN - Special Authority see SA1/02 below	– Retail pharmacy		
Tab 5 mg	1,550.00	30	 Ambrisentan Mylan
Tab 10 mg	1,550.00	30	✓ Ambrisentan Mylan

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or: The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

BOSENTAN - Special Authority see SA1991 below - Retail pharmacy

✓ Bosentan Dr. Reddv's Bosentan Dr Reddy's to be Principal Supply on 1 December 2021 60 ✓ Bosentan Dr

Bosentan Dr Reddy's to be Principal Supply on 1 December 2021

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:

continued...

Reddy's

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
- 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
- 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy: or
- 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA1992 on the next page - Retail pharmacy		
Tab 25 mg0.85	4	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022		
Tab 50 mg1.70	4	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022		
Tab 100 mg10.20	12	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022		

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications: or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dvn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 on the r	next page – Retail pharr	nacy	
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 91

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	✓ Differin
ISOTRETINOIN - Special Authority see SA2023 below - Ret	ail pharmacy		
Cap 5 mg		60	Oratane
Cap 10 mg	18.75	120	Oratane
Cap 20 mg	26.73	120	Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription15.57 50 g OP ✓ ReTrieve
ReTrieve to be Principal Supply on 1 January 2022

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 91

HYDROGEN PEROXIDE

	Subsidy (Manufacturer's Pri \$	ice) Subs	Fully Brand or sidised Generic Manufacturer
MUPIROCIN			
Oint 2%	6.60 (10.50)	15 g OP	Bactroban
a) Only on a prescriptionb) Not in combination			
SODIUM FUSIDATE [FUSIDIC ACID] Crm 2%	1.59	5 g OP	✓ Foban
a) Maximum of 5 g per prescriptionb) Only on a prescriptionc) Not in combination			
d) Foban to be Principal Supply on 1 December 2021 Oint 2%	1.59	5 g OP	✓ Foban
a) Maximum of 5 g per prescriptionb) Only on a prescriptionc) Not in combination			
d) Foban to be Principal Supply on 1 December 2021 SULFADIAZINE SILVER			
Crm 1% a) Up to 250 g available on a PSO b) Not in combination	10.80	50 g OP	✓ Flamazine
AMOROLFINE a) Only on a prescription b) Not in combination		- 100	
Nail soln 5%	14.93	5 ml OP	✓ <u>MycoNail</u>
a) Only on a prescription b) Not in combination			
Nail-soln 8%	5.72	7 ml OP	✓ Apo-Ciclopirox
CLOTRIMAZOLE * Crm 1%	0.77	20 g OP	✓ Clomazol
a) Only on a prescription b) Not in combination			
* Soln 1%	4.36 (7.55)	20 ml OP	Canesten
a) Only on a prescriptionb) Not in combination			
ECONAZOLE NITRATE Crm 1%	1.00	20 g OP	
	(7.48)	5	Pevaryl
a) Only on a prescription			
D) NOU IN COMBINATION		•	
b) Not in combination Foaming soln 1%, 10 ml sachets	9.89 (17.23)	3	Pevaryl

	Subsidy (Manufacturer's P \$	rice) Subs	Fully sidised	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g OP	✓ <u>N</u>	<u>lultichem</u>
a) Only on a prescriptionb) Not in combination				
* Lotn 2%	4.36	30 ml OP		
	(10.03))aktarin
a) Only on a prescriptionb) Not in combination				
* Tinct 2%		30 ml OP	_	And the site
	(12.10)		L	aktarin
a) Only on a prescriptionb) Not in combination				

Antipruritic Preparations

CALAMINE

a) Only on a prescription

b) Not in combination

Crm, aqueous, BP......1.26

100 g

 healthE Calamine Aqueous Cream BP

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%

9 20 g OP

✓ Itch-Soothe

Itch-Soothe to be Principal Supply on 1 December 2021

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Crystals	6.92	25 g	✓ MidWest
	29.60	100 g	✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 81

Corticosteroids - Plain

BE	FAMETHASONE DIPROPIONATE			
	Crm 0.05%	2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
BE	FAMETHASONE VALERATE			
*	Crm 0.1%	4.53	50 g OP	✓ Beta Cream
	Beta Cream to be Principal Supply on 1 January 2022		_	
*	Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
	Beta Ointment to be Principal Supply on 1 January 2022			
*	Lotn 0.1%	25.00	50 ml OP	✓ Betnovate

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's Price		dised	Generic
	\$	Per	_	Manufacturer
CLOBETASOL PROPIONATE				
* Crm 0.05%		30 g OP		Dermol
* Oint 0.05%	2.12	30 g OP	/	Dermol
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(10.00)			Eumovate
HYDROCORTISONE				
* Crm 1% - Only on a prescription	3.70	100 g OP	1	Hydrocortisone
				<u>(PSM)</u>
	17.15	500 g	•	<u>Hydrocortisone</u>
				(PSM)
* Powder – Only in combination		25 g		ABM
Up to 5% in a dermatological base (not proprietary Topic	cal Corticosteriod –	Plain) with or	with	out other dermatological
galenicals				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only of				
a prescription	10.57	250 ml	/	DP Lotn HC
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%		100 g OP		Locoid Lipocream
Oint 0.1%	10.28	100 g OP	•	Locoid
Locoid to be Principal Supply on 1 December 2021	10.00		,	
Milky emul 0.1%		100 ml OP	•	Locoid Crelo
Locoid Crelo to be Principal Supply on 1 December 202	I			
METHYLPREDNISOLONE ACEPONATE	4.40	45 00	,	
Crm 0.1% Oint 0.1%		15 g OP		Advantan
	4.40	15 g OP	٧	<u>Advantan</u>
MOMETASONE FUROATE	4.05	45 00	,	
Crm 0.1%		15 g OP		Elocon Alcohol Free Elocon Alcohol Free
Elocon Alcohol Free to be Principal Supply on 1 January	3.10	50 g OP	٧	Elocon Alconol Free
Oint 0.1%		15 g OP	1	Elocon
Olite 0.170	2.90	50 g OP		Elocon
Lotn 0.1%		30 ml OP		Elocon
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	1	Aristocort
Oint 0.02%		100 g OP		Aristocort
		J -		
Corticosteroids - Combination				
DETAMETUA COME VALEDATE MUTU CODUNA ELICIDATE (ELI	OIDIO AOIDI			
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU		15 ~ OD		
Crm 0.1% with sodium fusidate (fusidic acid) 2%	(10.45)	15 g OP		Fucicort
a) Maximum of 15 a per proceription	(10.43)			i ucicult
a) Maximum of 15 g per prescriptionb) Only on a prescription				
	ution.			
HYDROCORTISONE WITH MICONAZOLE – Only on a prescrip * Crm 1% with miconazole nitrate 2%		15 g OP	_	Micreme H
Micreme H to be Principal Supply on 1 December 2021	1.03	13 y OF	•	MICIGINE II
mioreme i i to be i illidipal dappiy dii i Decellibel 2021				

	Subsidy Manufacturer's l \$		Fully Brand or dised Generic Manufacturer
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Onl Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% (Pimafucort Crm 1% with natamycin 1% and neomycin sulphate 0.3 TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg	3.35 3.35 5% to be delis	15 g OP 15 g OP ted 1 May 2022)	✓ Pimafucort ✓ Pimafucort
and gramicidin 250 mcg per g - Only on a prescription	3.49 (9.28)	15 g OP	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE Dimethicone 10%
ZINC AND CASTOR OIL * Oint	4.25	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM * Crm	1.92	500 g	✓ Basic AquaCream ✓ Boucher ✓ Medco
CETOMACROGOL * Crm BP	2.48	500 g	✓ healthE
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ <u>Boucher</u> ✓ Kenkay Sorbolene ✓ Pharmacy Health Sorbolene with Glycerin
	3.10	1,000 ml OP	✓ ADE ✓ <u>Boucher</u>
(ADE Crm 90% with glycerol 10% to be delisted 1 January 2022) (Kenkay Sorbolene Crm 90% with glycerol 10% to be delisted 1 January 2022) (ADE Crm 90% with glycerol 10% to be delisted 1 January 2022)	nuary 2022)		
EMULSIFYING OINTMENT * Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION * Crm	2.19	500 g	✓ O/W Fatty Emulsion Cream
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ healthE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's I	Price) Subs Per	sidised Generic Manufacturer
	\$	Per	Wanufacturer
JREA			•
* Crm 10%	1.37	100 g OP	healthE Urea Cream
WOOL FAT WITH MINERAL OIL - Only on a prescription			
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
•	(11.95)		DP Lotion
	1.40	250 ml OP	
	(4.53)		DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91)		BK Lotion
	` 1.40 [′]	250 ml OP	
	(7.73)		BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft — Only in combination	A QQ	450 g	✓ healthE
Willie 30tt Only in combination	19.99	2,500 g	✓ healthE
Only in combination with a dermatological galenical of			
Only in combination with a connactorogreal galerilear c	a di	propriotary rop	noar cornocotorola Thain.
Minor Skin Infections			
POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
 a) Maximum of 130 g per prescription 			
b) Only on a prescription			
	4.15	100 ml	✓ Riodine
Antiseptic Solution 10%			/ B' ''
Antiseptic Solution 10%Antiseptic soln 10%		15 ml	✓ Riodine
		15 ml 500 ml	✓ Riodine ✓ Riodine
	3.83 5.40		
Antiseptic soln 10%	3.83 5.40	500 ml	
Antiseptic soln 10%	3.83 5.40 1.63 (3.48)	500 ml	✓ Riodine

DIMETHICONE

*	Lotn 4%4.98	200 MI OP	•	Dimethicone 4% Lotion
IVF	RMECTIN - Special Authority see SA1225 on the next page - Retail pharm:	acv		

- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Fer ✓ Manufacturer

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or

dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

- Any of the following:
 - 1 Filaricides; or
 - 2 Cutaneous larva migrans (creeping eruption); or
 - 3 Strongyloidiasis.

PERMETHRIN

Crm 5% 5.75 Lotn 5% 3.99	0 -	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN		
Shampoo 0.5%11.36	200 ml OP	Parasidose
(Parasidose Shampoo 0.5% to be delisted 1 January 2022)		

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail p	harmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Fither:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

PETAMETA MOCKE BIT NOT TOTAL WITH CALCUITOTHEE			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP	Daivobet
Daivobet to be Principal Supply on 1 December 2021		_	
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	Daivobet
Daiyobet to be Principal Supply on 1 December 2021		•	

		_	
	Subsidy		Fully Brand or
	(Manufacturer's F	rice) Sub: Per	sidised Generic Manufacturer
CALCIPOTRIOL	•		
Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex
COAL TAR		3 -	
Soln BP – Only in combination	36.25	200 ml	✓ Midwest
 Up to 10% only in combination with a dermatological With or without other dermatological galenicals. 		etary Topical (Corticosteriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUI	_PHUR		
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% at			
allantoin crm 2.5%		75 g OP	
	(8.00)	_	Egopsoryl TA
	3.43	30 g OP	
	(4.35)		Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint		25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp
PIMECROLIMUS - Special Authority see SA1970 below - Reta	ail pharmacy		
a) Maximum of 15 g per prescription			
b) Note: a maximum of 15 g per prescription and no more			_
Cream 1% SA1970 Special Authority for Subsidy	28.50	15 g OP	✓ <u>Elidel</u>
Initial application only from a dermatologist, paediatrician, oph of a dermatologist, paediatrician or ophthalmologist. Approvals meeting the following criteria: Both:			
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. 			
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORI * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium		n a prescriptio 500 ml	n ✓ <u>Pinetarsol</u>
SALICYLIC ACID Rounday Only in combination	10.00	050 ~	./ Midwood
Powder - Only in combination	18.88	250 g	✓ Midwest✓ PSM
 Only in combination with a dermatological base o With or without other dermatological galenicals. 	r proprietary Topid	cal Corticostero	
SULPHUR			
Precipitated – Only in combination	6.35	100 g	✓ Midwest
Only in combination with a dermatological base o		•	
2) With or without other dermatological galenicals.	r proprietary ropic	ai Corticostero	oiu – Fiairi
TACDOLIMIE			
TACROLIMUS Oint 0.19/ Special Authority see \$42074 on the payt page.	0		
Oint 0.1% – Special Authority see SA2074 on the next pag Retail pharmacy		30 g OP	✓ Zematop
a) Maximum of 30 g per prescription		30 g Oi	- Zematop
b) Note: a maximum of 30 g per prescription and no n	nore than one pre	scription per 12	2 weeks.
,			



Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE * Scalp app 0.1% Beta Scalp to be Principal Supply on 1 January 2022	9.84	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%	5.69	30 ml OP	✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%	6.57	100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%	3.23	100 ml OP	✓ <u>Sebizole</u>

Sunscreens

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 70 $\,$

IMIQUIMOD

Crm 5%, 250 mg sachet......21.72 24 **✔ Perrigo**

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM		
Crm 5%6.95	20 g OP	✓ Efudix
Efudix to be Principal Supply on 1 December 2021	ŭ	

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

Contraceptives - Non-hormonal

Condoms

	a) Maximum of 60 dev per prescription	S20 Hnann		
		17.02		✓ Gold Knight XL
		14.87	144	✓ Shield XL
⊬ 6	60 mm		12	✓ Gold Knight XL
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		15.57	144	✓ Gold Knight
÷ (56 mm, strawberry		12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		15.57	144	✓ Gold Knight
: !	56 mm, chocolate		12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	 a) Up to 60 dev available on a PSO 			
		11.64	144	✓ Moments
. [56 mm, 0.08 mm thickness, red		10	✓ <u>Moments</u>
	b) Maximum of 60 dev per prescription			•
	a) Up to 60 dev available on a PSO			
		11.64	144	✓ Moments
. [56 mm, 0.08 mm thickness		10	✓ <u>Moments</u>
	b) Up to 60 dev available on a PSO			
	a) Maximum of 60 dev per prescription			
į	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
	\	15.57	144	✓ Gold Knight
	56 mm, 0.05 mm thickness		12	✓ Gold Knight
,	b) Up to 60 dev available on a PSO	1.00	10	Cold Kalak
	a) Maximum of 60 dev per prescription			
	Alexander of OO decreases which	11.64	144	✓ <u>Moments</u>
;	ווווו סכ			
. į	b) Maximum of 60 dev per prescription 56 mm	0.07	10	✓ Moments
	a) Up to 60 dev available on a PSO			
	a) Un to 60 day available on a BSO	11.04	144	• WOUNGHES
•	oo mm, shawbeny, leu	11.64	144	✓ Moments
. [53 mm, strawberry, red	0.95	10	✓ Moments
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	11.07	177	· momonto
	oo min, onoooiato, brown	11.64	144	✓ Moments
	53 mm, chocolate, brown	0.95	10	✓ Moments
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			<u></u>
•	,	11.42	144	✓ Moments
	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
	b) Up to 60 dev available on a PSO			
	a) Maximum of 60 dev per prescription	11.04	177	Monicita
•		11.64	144	✓ Moments
	49 mm – Up to 144 dev available on a PSO53 mm.		144 10	✓ Moments✓ Moments

a) Maximumosidisedev per prescription b) Hindopassdayiqvailable on a PSO

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	√ <u>G</u>	old Knight XL
	a) Maximum of 60 dev per prescription				

Contraceptive Devices

INTRA-UTERINE DEVICE

a) Up to 40 dev available on a PSO

b) Up to 60 dev available on a PSO

b) Only on a PSO

	b) Only on a PSO			
*	IUD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓ Choice
	-			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up	to		
	84 tab available on a PSO	10.00	84	Mercilon 28
*	Tab 30 mcg with desogestrel 150 mcg and 7 inert tab	6.62	84	
		(19.80)		Marvelon 28

- a) Higher subsidy of \$13.80 per 84 tab with Special Authority see \$A0500 above
- b) Up to 84 tab available on a PSO

(Marvelon 28 Tab 30 mcg with desogestrel 150 mcg and 7 inert tab to be delisted 1 November 2021)

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ETH	HINYLOESTRADIOL WITH LEVONORGESTREL				
*	Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
	Up to 112 tab available on a PSO	2.18	84	✓	Microgynon 20 ED
		6.45	112	✓	Femme-Tab ED
*	Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - U	р			
	to 84 tab available on a PSO	9.45	84	✓	Microgynon 50 ED
*	Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
		(16.50)			Microgynon 30
	a) Higher subsidy of \$15.00 per 63 tab with Special Autl	hority see SA0500 on	the	previous p	age
	b) Up to 63 tab available on a PSO				
*	Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
	Up to 112 tab available on a PSO	1.77	84	✓	Levien ED
		6.45	112	✓	Femme-Tab ED
ETH	HINYLOESTRADIOL WITH NORETHISTERONE				
	Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to	1			
	84 tab available on a PSO		84	1	Brevinor 1/28
	Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – U		•	-	
	to 84 tab available on a PSO		84	1	Norimin
	10 0 1 100 01010 011 0 1 00		07	•	

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO1	6.50	84	✓ <u>Microlut</u>
	2	22.00	112	✓ <u>Microlut</u>
*	Subdermal implant (2×75 mg rods) – Up to 3 pack available			
	on a PSO10	06.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO	7.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised		
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	/	Noriday 28	
Emergency Contraceptives					
LEVONORGESTREL * Tab 1.5 mg		1 Part I o		Postinor-1	

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

• \$5.00 prescription charge (patient co-payment) will apply.

ACETIC ACID WITH HADDOAAOHINOLINE AND DICINOLEIC ACID

• prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator6.89	40 g OP	✓ <u>Micreme</u>
NYSTATIN		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat

Myometrial and Vaginal Hormone Preparations

,			
ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a		5	✓ DBL Ergometrine
OESTRIOL			
* Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
* Pessaries 500 mcg	6.86	15	✓ Ovestin
OXYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	 Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	 Oxytocin BNM
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj avail	able on a PSO		
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml Syntometrine to be Principal Supply on 1 January 2022		5	✓ Syntometrine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Su	ubsidy F	ully	Brand or
(Manufact	cturer's Price) Subsidi	sed	Generic
	\$ Per	1	Manufacturer

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

✓ Smith BioMed Rapid Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 109

5-Alpha Reductase Inhibitors

FINASTERIDE - Special Authority see SA0928 below - Retail pharmacy

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking oxybutynin prior to 1 June 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of oxybutynin.

* Tab 5 mg	11.70	500	✓ Apo-Oxybutynin
* Oral liq 5 mg per 5 ml	60.40	473 ml	✓ Apo-Oxybutynin
(Apo-Oxybutynin Oral liq 5 mg per 5 ml to be delisted	May 2022)		

POTASSIUM CITRATE

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	9	Fully Subsidised	Brand or Generic
 \$	Per	Jubsidised ✓	Manufacturer

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22 28	✓	<u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05 30	1	Solifenacin Mylan
Solifenacin Mylan to be Principal Supply on 1 Decer	mber 2021		
Tab 10 mg	3.72 30	1	Solifenacin Mylan
Solifenacin Mylan to be Principal Supply on 1 Decer	mber 2021		-

Detection of Substances in Urine

ORTHO-TOLIDINE			
* Compound diagnostic sticks	7.50	50 test OP	
, ,	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
	(13.92)		Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE			
Tab 200 mg	60.00	1	✓ Mifegyne
-	180.00	3	✓ Mifegyne

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Calcium Homeostasis

CALCITONIN		
* Inj 100 iu per ml, 1 ml ampoule121.00	5	Miacalcic
CINACALCET – Special Authority see SA1618 below – Retail pharmacy		
Tab 30 mg - Wastage claimable210.30	28	Sensipar

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID



Zoledronic acid Mylan to be Principal Supply on 1 December 2021

⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement: and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — **(early breast cancer)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Cubaidu		Fully	Drand or	
Subsidy		. ,	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETAT	Έ	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	5	
(36.96)		Celestone
, ,		Chronodose
DEXAMETHASONE		
* Tab 0.5 mg – Up to 60 tab available on a PSO	30	✓ Dexmethsone
Dexmethsone to be Principal Supply on 1 January 2022		20/1111011100110
* Tab 4 mg – Up to 30 tab available on a PSO2.65	30	✓ Dexmethsone
Dexmethsone to be Principal Supply on 1 January 2022		
Oral liq 1 mg per ml45.00	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ Dexamethasone
ing 4 mg per mi, 1 mi ampoule op to o mg available on a 1 00	10	Phosphate
		Panpharma
* Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO 16.37	10	✓ Dexamethasone
ing 4 mg per mi, 2 mi ampoule op to o mg available on a 1 50 10.07	10	Phosphate
		Panpharma
FLUDROCORTISONE ACETATE		<u>r unpriumu</u>
	100	✓ Florinef
* Tab 100 mcg	100	▼ Florillei
HYDROCORTISONE		
* Tab 5 mg	100	✓ Douglas
* Tab 20 mg	100	✓ Douglas
* Inj 100 mg vial	1	✓ Solu-Cortef
a) Up to 5 inj available on a PSO		
b) Only on a PSO		
c) Solu-Cortef to be Principal Supply on 1 November 2021		
METHYLPREDNISOLONE		_
* Tab 4 mg112.00	100	✓ Medrol
* Tab 100 mg194.00	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)		
Inj 40 mg vial18.90	1	✓ Solu-Medrol-Act-
		O-Vial
Inj 125 mg vial28.90	1	✓ Solu-Medrol-Act-
		O-Vial
Inj 500 mg vial22.78	1	✓ Solu-Medrol-Act-
11 JUU 11 19 VIGI	1	O-Vial
		O-Viai
Inj 1 g vial27.83	1	✓ Solu-Medrol
, ,		

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
METHYLPREDNISOLONE ACETATE Inj 40 mg per ml, 1 ml vial PREDNISOLONE	44.40	5	1	Depo-Medrol
** Oral liq 5 mg per ml – Up to 30 ml available on a PSO a) Restricted to children under 12 years of age. b) Redipred to be Principal Supply on 1 December 2021		30 ml OP	•	Redipred
PREDNISONE * Tab 1 mg * Tab 2.5 mg * Tab 5 mg - Up to 30 tab available on a PSO * Tab 20 mg - Up to 30 tab available on a PSO TETRACOSACTRIN	21.04 19.30	500 500 500 500	1	Apo-Prednisone Apo-Prednisone Apo-Prednisone Apo-Prednisone
 Inj 250 mcg per ml, 1 ml ampoule Inj 1 mg per ml, 1 ml ampoule 		1	1	UK Synacthen S29 AU Synacthen Synacthen Synacthen Depot Synacthene Retard S29
TRIAMCINOLONE ACETONIDE Inj 10 mg per ml, 1 ml ampoule Inj 40 mg per ml, 1 ml ampoule		5 5		Kenacort-A 10 Kenacort-A 40

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CYPROTERONE ACETATE			
Tab 50 mg	14.37	50	✓ Rex S29
-			✓ Siterone
Siterone to be Principal Supply on 1 January 2022			
Tab 100 mg	28.03	50	✓ Siterone
Siterone to be Principal Supply on 1 January 2022			
(Rex S29 Tab 50 mg to be delisted 1 January 2022)			
TESTOSTERONE			
Patch 5 mg per day	90.00	30	✓ Androderm
TESTOSTERONE CIPIONATE			
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ Depo-Testosterone
	05.00	'	• Depo-Testosterone
TESTOSTERONE ESTERS	40.00		
Inj 250 mg per ml, 1 ml	12.98	1	✓ Sustanon Ampoules
TESTOSTERONE UNDECANOATE			
Cap 40 mg	21.00	60	Andriol Testocaps
Inj 250 mg per ml, 4 ml vial	86.00	1	✓ Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

_		Subsidy		Fully	Brand or
		(Manufacturer's Pri	ce) Sub	sidised	Generic
_		\$	Per		Manufacturer
0	estrogens				
ΟE	STRADIOL - See prescribing guideline on the previous page				
	Tab 1 mg	4.12	28 OP		
	· ·	(11.10)		Е	strofem
*	Tab 2 mg	4.12	28 OP		
		(11.10)		_	strofem
*	Patch 100 mcg per 24 hours	7.91	4	✓ C	limara
	a) No more than 1 patch per week b) Only on a prescription				
*	Patch 50 mcg per 24 hours	7.04	4	✓ C	limara
	a) No more than 1 patch per week				
	b) Only on a prescription				
	Patch 25 mcg per day	6.12	8	√ E	stradot
	,	7.85		√ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				•
	b) Only on a prescription				
	Patch 50 mcg per day	7.04	8	✓ E	stradot 50 mcg
		9.22		✓ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 75 mcg per day	7.91	8		stradot
		10.60		✓ E	stradiol TDP
					Mylan S29
	a) No more than 2 patch per week				
	b) Only on a prescription				
	Patch 100 mcg per day	7.91	8	✓ E	stradot
	a) No more than 2 patch per week				
	b) Only on a prescription				
•	imara Patch 100 mcg per 24 hours to be delisted 1 January 20	,			
•	imara Patch 50 mcg per 24 hours to be delisted 1 January 202	,			
	STRADIOL VALERATE – See prescribing guideline on the pro-				
	Tab 1 mg		84		rogynova
	Tab 2 mg		84	✓ P	rogynova
	STROGENS - See prescribing guideline on the previous page				
*	Conjugated, equine tab 300 mcg		28	_	
*	Conjugated equips tob COE mag	(17.50)	00	Р	remarin
*	Conjugated, equine tab 625 mcg		28	D	remarin
		(17.50)		P	ı Gınıdılı I
Р	rogestogens				
•	. 0 9 0 1 0 9 0 1 0				
	DROXYPROGESTERONE ACETATE - See prescribing guid			_	
	Tab 2.5 mg		30	-	rovera
	Tab 5 mg		100	-	rovera
*	Tab 10 mg	8.94	30	✓ P	rovera

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

		Subsidy (Manufacturer's Price)) Per	Fully Subsidised	Brand or Generic Manufacturer			
P	Progestogen and Oestrogen Combined Preparations							
OE	STRADIOL WITH NORETHISTERONE - See prescribing gui	ideline on page 82						
*	Tab 1 mg with 0.5 mg norethisterone acetate	, ,	28 OF	•				
		(18.10)			Kliovance			
*	Tab 2 mg with 1 mg norethisterone acetate		28 OF					
		(18.10)			Kliogest			
*	Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	5 40	00.05					
	oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OF		T.:			
		(18.10)			Trisequens			
C	ther Oestrogen Preparations							
ET	HINYLOESTRADIOL							
*	Tab 10 mcg	17.60	100	✓	NZ Medical and Scientific			
OE	STRIOL							
*	Tab 2 mg	7.00	30	✓	Ovestin			
C	ther Progestogen Preparations							
LE	VONORGESTREL							
*	Intra-uterine device 52 mg	269.50	1	1	Mirena			
*	Intra-uterine device 13.5 mg	215.60	1	✓	<u>Jaydess</u>			
ME	DROXYPROGESTERONE ACETATE							
	Tab 100 mg	116.15	100	✓	Provera HD			
NC	RETHISTERONE							
*	Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	✓	Primolut N			
PF	OGESTERONE							
	Cap 100 mg - Special Authority see SA1609 below - Retail							
	, , , , , , , , , , , , , , , , , , , ,							

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

pharmacy......16.50

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

30

✓ Utrogestan

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully Brand or sidised Generic Manufacturer
Thyroid and Antithyroid Agents			
CARBIMAZOLE * Tab 5 mg	10.80	100	✓ Neo-Mercazole ✓ Neo-Mercazole S29 S29
(Neo-Mercazole S29 S29 Tab 5 mg to be delisted 1 January 20	022)		
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg		28	Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1.78	28	Mercury Pharma
	6.01	90	Synthroid
	66.78	1,000	✓ Eltroxin
PROPYLTHIOURACIL — Special Authority see SA1199 below— Propylthiouracil is not recommended for patients under the a treatments are contraindicated.		s the patie	nt is pregnant and other
Tab 50 mg	35.00	100	✓ PTU S29
■ Special Authority for Subsidy Initial application from any relevant practitioner. Approvals val Both:		cations m	eeting the following criteria:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 belo	ow – Retail pha	armacy	
*	Inj 5 mg cartridge	69.75	ĺ	✓ Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			-
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			-
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			•

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:

Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and</p>
- 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — **(short stature without growth hormone deficiency)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 \$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician

Subsidy (Manufacturaria Price)	Ç.	Fully bsidised	Brand or
 (Manufacturer's Price) \$	Per	⊅Sidised ✓	Generic Manufacturer

continued...

and/or ENT surgeon; and

- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
 - 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
 - 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
 - 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
 - 5 No malignancy has developed after growth hormone therapy was commenced; and
 - 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
- 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN			
Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva
LEUPRORELIN			
Additional subsidy by endorsement where the patient is a goserelin and the prescription is endorsed accordingly.	child or adolescent a	ind is unable	to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsi	dy of		
\$221.60 per 1 inj with Endorsement	66.48	1	
	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subs	sidy		
of \$591.68 per 1 inj with Endorsement		1	
	(591.68)		Lucrin Depot 3-month

Vasopressin Agonists

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

	Subsidy (Manufacturer's I	Price) Subs	Fully	Brand or Generic
	\$	Per	1	Manufacturer
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	Minirin
Tab 200 mcg	54.45	30	✓ N	Minirin
▲ Nasal drops 100 mcg per ml	39.03	2.5 ml OP	✓ N	Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	√ [<u>PH&T</u>
Inj 4 mcg per ml, 1 ml(Minirin Nasal drops 100 mcg per ml to be delisted 1 January 202		10	✓ N	Minirin

Other Endocrine Agents

CABERGOLINE

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE			
Can 250 mg	558 00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 ✓ Eskazole S29 **⇒SA1318** Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription			
Tab 100 mg	7.97	6	✓ Vermox
Vermox to be Principal Supply on 1 January 2022			
Oral liq 100 mg per 5 ml	2.18	15 ml	
	(7.53)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 63
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 237

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE		
Cap 250 mg24.70	100	✓ Ranbaxy-Cefaclor
		Ranbaxy-Cefactor
		S29 S29
Grans for oral liq 125 mg per 5 ml - Wastage claimable3.53	100 ml	✓ Ranbaxy-Cefactor
		✓ Ranbaxy-Cefactor
		S29 S29
CEFALEXIN		
Cap 250 mg3.33	20	✓ Cephalexin ABM
Cap 500 mg	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml – Wastage claimable8.75	100 ml	✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable11.75	100 ml	✓ Cefalexin Sandoz
CEFAZOLIN - Subsidy by endorsement		
Only if prescribed for dialysis or cellulitis in accordance with a DHB approved	protocol and	the prescription is endorsed
accordingly.		
Inj 500 mg vial	5	✓ AFT
lnj 1 g vial	5	✓ AFT
CEFTRIAXONE – Subsidy by endorsement		
a) Up to 10 inj available on a PSO		
h) Cubaidiand only if preparited for a dialysis or ayotic fibragic nations or the	trootmont of a	anarrhaga ar tha traatmant a

b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.89	1	✓ Ceftriaxone-AFT
lnj 1 g vial	3.99	5	✓ Ceftriaxone-AFT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Subsic Per	Fully dised	Brand or Generic Manufacturer
CEFUROXIME AXETIL – Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the pre Tab 250 mg		accordingly 50		Zinnat

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
ů i	2.57		✓ Zithromax
Zithromax to be Sole Supply on 1 December 2021			
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage			

(Apo-Azithromycin Tab 250 mg to be delisted 1 May 2022) (Apo-Azithromycin Tab 500 mg to be delisted 1 December 2021)

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and

claimable14.38

- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

✓ Zithromax

15 ml

	Subsidy (Manufacturer's Price)	Fully Subsidised		Brand or Generic	
	\$	Per	✓	Manufacturer	
CLARITHROMYCIN					
Tab 250 mg - Maximum of 56 tab per prescription; can be					

14 ✓ Apo-Clarithromycin ✓ Klacid 8.53 50 ml ✓ Klacid

- a) Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below
 - b) Wastage claimable

(Apo-Clarithromycin Tab 250 mg to be delisted 1 February 2022)

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omegrazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets: and
- 2 For use only in combination with ome prazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ERYTHROMYCIN (AS LACTOBIONATE)

✓ Ervthrocin IV

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	
RYTHROMYCIN ETHYL SUCCINATE			_	
Tab 400 mg	16.95	100	/	E-Mycin
a) Up to 20 tab available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	•	E-Mycin
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable	0.77	100 1		□ M
Grans for oral liq 400 mg per 5 ml		100 ml	•	E-Mycin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
RYTHROMYCIN STEARATE				
Tab 250 mg - Up to 30 tab available on a PSO		100		50.
T 500	(22.29)	400		ERA
Tab 500 mg		100		50.
	(44.58)			ERA
ERA Tab 250 mg to be delisted 1 April 2022)				
ERA Tab 500 mg to be delisted 1 September 2022)				
ROXITHROMYCIN				
Tab disp 50 mg	8.29	10	1	Rulide D
Restricted to children under 12 years of age.				_
Tab 150 mg	8.28	50	/	Arrow-
				Roxithromycin
Tab 300 mg	16 33	50	1	Arrow-
140 500 mg				Roxithromycin
Penicillins				
MOXICILLIN				
Cap 250 mg	22.50	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500	1	Alphamox
a) Up to 30 cap available on a PSO				-
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	1	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial		10		Ibiamox
Inj 500 mg vial	17.43	10		Ibiamox Ibiamox
Inj 1 g vial - Up to 5 inj available on a PSO		10		

urer's Price)	sidy F		
\$	Subsidis Per		eneric anufacturer
89	10	✓ Cura	m Duo 500/125
03	10	Cuia	III Duo 300/123
00 1	100 ml		
00 1	100 ml	✓ Augr	nenun
00 10			
20 100	0 ml OP	Cura	m
93	10	✓ Bicill	in LA
09	10	✓ Sand	loz
00	10	<u> </u>	. <u></u>
00	050	. Ctan	hla
83		✓ Stap	
61		✓ Stap	niex
29 1	100 ml	✓ AFT	
68 1	100 ml	✓ AFT	
56		✓ Fluci	
87		Fluci	
70	5	✓ Fluci	<u>l</u>
84	50	✓ Cilica	aine VK
86	50	✓ Cilica	aine VK
99 1	00 ml	✓ AFT	
99 1	100 ml	✓ AFT	
50	5	✓ Cilia	nino
50	ວ 	- CIIIC	anne
43	500	✓ Doxi	ne
	.43		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

95

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	· •	Manufacturer
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
	(12.05)		M	lino-tabs
* Cap 100 mg	19.32	100		
	(52.04)		M	1inomycin
⇒SA1355 Special Authority for Manufacturers Price				

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy

Tab 250 mg21.42 ✓ Accord \$29

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 63

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	✓ Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓ Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist -	Subsidy by endorse	ment	
Only if prescribed for dialysis or cystic fibrosis patient and t	he prescription is en	dorsed acc	ordingly.
Inj 150 mg		1	✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	25.00	5	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urina	ary tract inf	ection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urina	ary tract inf	ection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓ Pfizer
	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient	t or complicated uring	ary tract inf	ection and the prescription is

endorsed accordingly.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MOXIFLOXACIN - Special Authority see SA1740 below - Retail	pharmacy			
No patient co-payment payable				
Tab 400 mg	42.00	5	✓ A	velox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case. Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

⇒SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 on the next page - Retail pharmacy

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approval the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with F 2 For pregnant patients for the term of the pregnancy;	HV for a period of 3 month		nless notific	ed for applications meetin
3 For infants with congenital toxoplasmosis until 12 mg	onths of age.			
SODIUM FUSIDATE [FUSIDIC ACID]	07.05	00		Provide a
Tab 250 mg		36		Fucidin
SULFADIAZINE SODIUM - Special Authority see SA1331	•	EC		Weekheudt 000
Tab 500 mg ■ SA1331 Special Authority for Subsidy	543.20	56		Wockhardt S29
the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with F 2 For pregnant patients for the term of the pregnancy; 3 For infants with congenital toxoplasmosis until 12 mg	or	ıs; or		
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement a) Only if prescribed for dialysis or cystic fibrosis p b) Tobramycin Mylan to be Principal Supply on 1 c	patient and the prescription January 2022	5 n is en		Tobramycin Mylan ordingly.
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement		56 dos	se 🗸	Tobramycin BNM
b) Only if prescribed for a cystic fibrosis patient an	d the prescription is endo	rsed a	ccordingly.	
TRIMETHOPRIM * Tab 300 mg - Up to 30 tab available on a PSO TMP to be Principal Supply on 1 January 2022	18.55	50	•	ТМР
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRI	IMOXAZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 m to 30 tab available on a PSO Trisul to be Principal Supply on 1 January 2022	ng – Up 64.80	500	✓.	Trisul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to available on a PSO		100 m	nl 🗸	Deprim
VANCOMYCIN - Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient			is or for trea	atment of Clostridium
difficile following metronidazole failure and the prescrip	tion is endorsed according	jly.		Madan

Inj 500 mg vial2.35

✓ Mylan

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 64
- b) For topical antifungals refer to GENITO URINARY, page 77

FLUCONAZOLE

OCONAZOLL		
Cap 50 mg2.75	28	✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy109.34	35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOI F

Cap 100 mg4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -		
Retail pharmacy141.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KFTOCONAZOI F

Tab 200 mg - PCTCBS	30	✓ Link Healthcare S29
		✓ Nizoral S29
	100	✓ Strides Shasun S29

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pric	-,	Fully obsidised	Brand or Generic
	\$	Per		Manufacturer
NYSTATIN				
Tab 500,000 u	14.16	50		
	(17.09)		N	lilstat
Cap 500,000 u	12.81	50		
·	(15.47)		N	lilstat
POSACONAZOLE - Special Authority see SA1285 below - Re	tail pharmacy			
Tab modified-release 100 mg	869.86	24	✓ N	loxafil
Oral liq 40 mg per ml	761.13	105 ml OP	✓ N	loxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE ★ Tab 250 mg 8.15 84 ✓ Deolate VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy 91.00 56 ✓ Vttack Tab 50 mg 91.00 56 ✓ Vttack Tab 200 mg 350.00 56 ✓ Vttack Powder for oral suspension 40 mg per ml - Wastage 1,437.00 70 ml ✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist.

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued...

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - Retail ph	armacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaquine S29

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl	
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl	
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S	
Suppos 500 mg	24.48	10	✓ Flagyl	
ORNIDAZOLE				
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole	
Arrow-Ornidazole to be Principal Supply on 1 December	er 2021			

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Generic
YCLOSERINE - Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendar respiratory physician.	ation of, an infectious d	iseas	e physicia	n, clinical microbiologist
Cap 250 mg	344.00	60	/	Cyclorin S29
APSONE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendadermatologist		iseas	e physicia	n, clinical microbiologist
Tab 25 mg		100		Dapsone
Tab 100 mg	329.50	100	/	Dapsone
THAMBUTOL HYDROCHLORIDE – Retail pharmacy-Special a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation		iseas	e physicia	n, clinical microbiologist
respiratory physician Tab 100 mg	95.72	100	1	EMB Fatol \$29
Tab 400 mg		56	_	Myambutol S29
ONIAZID - Retail pharmacy-Specialist	43.04	50	•	wyambutor
b) Prescriptions must be written by, or on the recommendar microbiologist, dermatologist or public health physician Tab 100 mg		100		PSM
 SONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician 	ation of, an internal me	dicine	physician	ı, paediatrician, clinical
Tab 100 mg with rifampicin 150 mgRifinah to be Principal Supply on 1 January 2022		100		Rifinah
 Tab 150 mg with rifampicin 300 mg	179.13	100	•	Rifinah
ARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda	ation of, an infectious d	iseas	e specialis	st, clinical microbiologist
respiratory physician			_	
Grans for oral liq 4 g sachet	280.00	30	•	Paser S29
ROTIONAMIDE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendar respiratory physician				-
Tab 250 mg	305.00	100	•	Peteha S29
/RAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendation	ation of, an infectious d	iseas	e physicia	n, clinical microbiologist
respiratory physician				

✓ Entecavir Sandoz

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	•	Manufacturer

RIFABUTIN - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or gastroenterologist
- **★** Cap 150 mg......299.75 30 **✓ Mycobutin**

RIFAMPICIN - Subsidy by endorsement

- a) No patient co-payment payable
- b) For confirmed recurrent Staphylococcus aureus infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement -Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.

*	Cap 150 mg58.54	100	Rifadin
	Cap 300 mg122.06		✓ Rifadin
	Oral lig 100 mg per 5 ml		✓ Rifadin

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 237

Hepatitis B Treatment

ENTECAVIR		
* Tab 0.5 mg	52.00	30

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 106

Herpesvirus Treatments

ACICLOVIR		
* Tab dispersible 200 mg1.60	25	✓ Lovir
* Tab dispersible 400 mg5.38	56	✓ Lovir
* Tab dispersible 800 mg5.98	35	✓ Lovir
VALACICLOVIR		
Tab 500 mg6.50	30	✓ Vaclovir
Vaclovir to be Principal Supply on 1 January 2022		
Tab 1,000 mg13.76	30	✓ Vaclovir
Vaclovir to be Principal Supply on 1 January 2022		
VALGANCICLOVIR - Special Authority see SA1993 on the next page - Retail	pharmacy	
Tab 450 mg	60	✓ Valganciclovir
-		Mylan

Valganciclovir Mylan to be Principal Supply on 1 December 2021

_			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Subsidy (Manufacturer's Price) \$ Pe

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 28 **✓ Harvoni**

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac. PO Box 10-254. WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

below
Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is
co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated
accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 106 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

⇒SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and

Subsidy (Manufacturer's Pric	:e)	Fully Subsidised	Brand or Generic	
(Mandactary 8 1 1 1 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2 2	Per	✓	Manufacturer	

continued...

- 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
- 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (**Confirmed HIV**) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous	page - Retail pharm	acy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	page – Retail pharn	nacy	
Tab 200 mg	770.00	60	✓ Intelence

	Subsidy		Fully	Brand or
	(Manufacturer's Pi \$	rice) Subs Per	idised •	Generic Manufacturer
NEVIRAPINE – Special Authority see SA1651 on page	106 – Retail pharmacy			
Tab 200 mg	84.00	60	√ 1	Nevirapine Alphapharm
Nevirapine Alphapharm to be Principal Supply o Oral suspension 10 mg per ml	•	240 ml	•	/iramune Suspension
Nucleosides Reverse Transcriptase Inhib	itors			
ABACAVIR SULPHATE - Special Authority see SA1651	l on page 106 – Retail ph	armacy		
Tab 300 mg	180.00	60		<u>Ziagen</u>
Oral liq 20 mg per ml		240 ml OP		Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special A Note: abacavir with lamivudine (combination tablets anti-retroviral Special Authority.				
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ <u>I</u>	<u> Kivexa</u>
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR charmacy Note: Efavirenz with emtricitabine and tenofovir disc anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir 245 mg (300 mg as a maleate)	oproxil counts as three an	·	dicatio	. •
EMTRICITABINE - Special Authority see SA1651 on pa	ige 106 – Retail pharmac	30	_	-
Oap 200 mg		30	• .	
AMIVI IDINE Chariel Authority and CA16E1 on page	106 Poteil phermony		_	<u>Emtriva</u>
AMIVUDINE – Special Authority see SA1651 on page Tab 150 mg		60	✓ <u>I</u>	<u>-mtriva</u> <u>-amivudine</u> Alphapharm
	84.50	60 240 ml OP	✓ <u>!</u>	_amivudine Alphapharm
Tab 150 mg Oral liq 10 mg per ml ZIDOVUDINE [AZT] – Special Authority see SA1651 on		240 ml OP		_amivudine Alphapharm
Tab 150 mg Oral liq 10 mg per ml IDOVUDINE [AZT] – Special Authority see SA1651 on Cap 100 mg		240 ml OP acy 100	√ 3	<u>Lamivudine</u> <u>Alphapharm</u> BTC Retrovir
Tab 150 mg Oral liq 10 mg per ml ZIDOVUDINE [AZT] – Special Authority see SA1651 on		240 ml OP	√ 3	<u>_amivudine</u> <u>Alphapharm</u> BTC
Oral liq 10 mg per ml ZIDOVUDINE [AZT] – Special Authority see SA1651 on Cap 100 mg		240 ml OP acy 100 200 ml OP e 106 – Retail	✓ 3 ✓ I ✓ I	Lamivudine Alphapharm BTC Retrovir Retrovir acy
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail	✓ 3 ✓ I ✓ Inpharmaledicat	Lamivudine Alphapharm BTC Retrovir Retrovir acy
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m	✓ 3 ✓ I ✓ Inpharmaledicat	_amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m	✓ 3 ✓ I ✓ Inpharmaledicat	_amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m 60	y i pharmnedicat	amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of Alphapharm
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m 60	y i pharmnedicat	Amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of
Oral liq 10 mg per ml		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m 60	y i y i pharmhedicat	Amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of Alphapharm
Tab 150 mg		240 ml OP acy 100 200 ml OP e 106 – Retail inti-retroviral m 60	y i y i pharm nedicat	amivudine Alphapharm BTC Retrovir Retrovir acy ions for the purposes of Alphapharm

	Subsidy (Manufacturer's I	Price) Subsi		Brand or Generic
	\$	Per	✓	Manufacturer
LODINAVID WITH DITONAVID Special Authority see SA1661	on page 106	Dotail pharmaou		
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651				
Tab 100 mg with ritonavir 25 mg	150.00	60	• [∟opinavir/Ritonavir Mylan
	183.75		✓ }	Kaletra
Tab 200 mg with ritonavir 50 mg	295.00	120	√ L	opinavir/Ritonavir
				Mylan
	463.00		✓ H	Caletra
Oral lig 80 mg with ritonavir 20 mg per ml	735.00	300 ml OP	✓	Caletra
(Kaletra Tab 100 mg with ritonavir 25 mg to be delisted 1 Februa				
(Kaletra Tab 200 mg with ritonavir 50 mg to be delisted 1 Februa	, ,			
,	,			
RITONAVIR – Special Authority see SA1651 on page 106 – Ret	ail pharmacy			
Tab 100 mg	43.31	30	✓ <u>N</u>	<u>Norvir</u>
•				
Strand Transfer Inhibitors				
DOLUTEGRAVIR - Special Authority see SA1651 on page 106	– Retail nharma	CV		
Tab 50 mg	•	30	/ 1	Tivicay
-			•	iviouy
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 of		etail pharmacy		
Tab 400 mg	1,090.00	60	✓	sentress
Tab 600 mg	1,090.00	60	✓	sentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
 \$	Per	✓	Manufacturer

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.
- Inj 180 mcg prefilled syringe......500.00 ✓ Pegasys

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed: or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive: and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml: and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2.000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use: and
- 8 Not co-infected with HCV, HIV or HDV: and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate: or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet auidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	•	Manufacturer

continued...

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	40.01	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg		100	✓ Nifuran
* Cap modified-release 100 mg - Wastage claimable	86.40	100	✓ <u>Macrobid</u>
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	135.00	100	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

				_
	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per		Manufacturer
Anticholinesterases				
IEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	19.60	10	1	Juno S29
,g r,	33.81			Max Health
	98.00	50		AstraZeneca
Juno ⁸²⁹ Inj 2.5 mg per ml, 1 ml ampoule to be delisted 1 Mai AstraZeneca Inj 2.5 mg per ml, 1 ml ampoule to be delisted 1 M YRIDOSTIGMINE BROMIDE	March 2022)			
▲ Tab 60 mg	45.79	100	/	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
ICLOFENAC SODIUM				
← Tab EC 25 mg	1 00	50	1	Diclofenac Sandoz
Diclofenac Sandoz to be Principal Supply on 1 January		50	•	Diciolellac Salluoz
Tab 50 mg dispersible		20	1	Voltaren D
₹ Tab 50 mg dispersible		50		Diclofenac Sandoz
Diclofenac Sandoz to be Principal Supply on 1 January		50	•	Diciolellac Salluoz
Tab long-acting 75 mg		100	1	Voltaren SR
Tab long acting 75 mg	22.80	500		Apo-Diclo SR
Tab long-acting 100 mg		500		Apo-Diclo SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5		Voltaren
Suppos 12.5 mg		10		Voltaren
Suppos 25 mg		10		Voltaren
		10		Voltaren
- Triber		10		Voltaren
Suppos 100 mgpo-Diclo SR Tab long-acting 75 mg to be delisted 1 May 2022 Apo-Diclo SR Tab long-acting 100 mg to be delisted 1 May 2023 BUPROFEN	?)	10	·	voltaren
₹ Tab 200 mg	21.40	1,000	<i>•</i>	Relieve
Tab long-acting 800 mg		30		Brufen SR
1 45 15 19 45 and 9 50 5 119	5.99	00		Ibuprofen SR BNM
Brufen SR to be Principal Supply on 1 January 2022	0.00			
Oral liq 20 mg per mlbuprofen SR BNM Tab long-acting 800 mg to be delisted 1 Jar		200 n	nl 🗸	Ethics
ETOPROFEN	idary LoLL)			
ETOPROFEIN Cap long-acting 200 mg	12.07	28	.1	Oruvail SR
	12.07	20	•	Oruvali Sh
EFENAMIC ACID				
F Cap 250 mg		50		
	(9.16)			Ponstan
	0.50	20		

(5.60)

Ponstan

		Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
NAPRO	OXEN				
* Ta	b 250 mg Noflam 250 to be Principal Supply on 1 January 2022	32.69	500	•	Noflam 250
∗ Ta	b 500 mg Noflam 500 to be Principal Supply on 1 January 2022	28.71	250	✓	Noflam 500
* Ta	b long-acting 750 mgNaprosyn SR 750 to be Principal Supply on 1 January 20		28	•	Naprosyn SR 750
* Ta	b long-acting 1 g Naprosyn SR 1000 to be Principal Supply on 1 January 2	8.62	28	•	Naprosyn SR 1000
SULINI	DAC				
* Ta	b 100 mg	9.57	56	1	Mylan S29
* Ta	b 200 mg	15.10	50	1	Aclin
(Aclin 7	Tab 200 mg to be delisted 1 January 2022)	16.91	56	✓	Sulindac Mylan S29
•	lac Mylan 👀 Tab 200 mg to be delisted 1 January 2022,)			
TENO	KICAM				
	b 20 mg		100		Tilcotil
* Inj	20 mg vial	9.95	1	•	AFT
NSA	IDs Other				
CELEC	· · · · · -			_	
	up 100 mg		60 30		Celecoxib Pfizer Celebrex
Ga	p 200 mg	3.30	30		Celecoxib Pfizer
Topi	ical Products for Joint and Muscular Pain				
CAPSA	AICIN				
Cri	m 0.025% - Special Authority see SA1289 below - Retail	0.75	45 a C	ND ./	Zootriy
₩ \$Λ1	pharmacy 289 Special Authority for Subsidy	9.75	45 g C	,r •	<u>Zostrix</u>
Initial a	application from any relevant practitioner. Approvals valid rithritis that is not responsive to paracetamol and oral non-s	d without further renesteroidal anti-inflam	ewal u natorie	inless notif es are conf	ied where the patient has traindicated.
Anti	rheumatoid Agents				
	OXYCHLOROQUINE - Subsidy by endorsement				
Su suj mi Ph	bisidised only if prescribed for rheumatoid arthritis, systemi ppression, relevant dermatological conditions (cutaneous f ucosal ulceration)*, sarcoidosis (pulmonary and non-pulmo larmacists may annotate the prescription as endorsed whe droxychloroquine. Note: Indication marked with a * is an i	orms of lupus and li nary)*, and the pres re there exists a rec	chen p criptio ord of	olanus, cut n is endors	aneous vasculitides and sed accordingly.
* Ta	b 200 mg		100	•	Plaquenil
	NOMIDE			_	_
	b 10 mg		30		Arava
ıa	b 20 mg	6.00	30	•	<u>Arava</u>

Tab 125 mg67.23

Tab 250 mg110.12

100

100

✓ D-Penamine

✓ D-Penamine

PENICILLAMINE

Principal Supply

Fosamax Plus

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer	
Drugs Affecting Bone Metabolism				

Alendronate for Osteoporosis

ALI	ENDRONATE SODIUM			
*	Tab 70 mg	2 44	4	✓ Fosamax

ALENDRONATE SODIUM WITH COLECALCIFEROL

Other Treatments

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither
 - 2.1 The patient is female and postmenopausal; or

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically: or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a

Subs (Manufactur	.,	ully Brand or sed Generic	
	Per	✓ Manufacture	er

continued...

fall from a standing height or less

- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	27.53	1	Pamisol
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see	SA1779 below – Retail p	harmacy	

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or

* Tab 60 mg53.76

- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Tab 35 mg3.10	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 on the next page - Retail pharm	nacy	
Inj 250 mcg per ml, 2.4 ml490.00	1	✓ Forteo

✓ Evista

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

⇒SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease: and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or

Subsidy		Fully	Brand or
(Manufacturer's Pric	·0)	Subsidised	Generic
(Wallactarer 3 File		Oubbialoca	
\$	Per	/	Manufacturer

continued...

- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid: or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg		500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA	1963 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite S29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	 Benzbromaron AL
			100 S29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

COLCHICINE

*	Tab 500 mcg	9.58	100	Colgout
---	-------------	------	-----	---------------------------

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer	
FEBUXOSTAT - Special Authority see SA2054 below - Retail ph	armacy				
Tab 80 mg	20.00	28	√ F	Febuxostat multichem	
	39.50		✓ A	Adenuric	
Febuxostat multichem to be Sole Supply on 1 January 20	22				
Tab 120 mg	20.00	28	√ F	Febuxostat multichem	
Echuyactat multicham ta ba Cala Supply on 1 January 20	39.50		✓ A	Adenuric	

Febuxostat multichem to be Sole Supply on 1 January 2022

(Adenuric Tab 80 mg to be delisted 1 January 2022)

(Adenuric Tab 120 mg to be delisted 1 January 2022)

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- Both:
 - 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
 - 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

חח	D			\sim 1	\Box
PR	ᄱ	LΙΝ	ᅜ	U	υ

	SPENEOID			
*	Tab 500 mg	55.00	100	✓ Probenecid-AFT

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Muscle Relaxants

BACLOFEN			
* Tab 10 mg	4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo			ents have been ineffective or have
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement	306.82	5	✓ Medsurge
a) Subsidised only for use in a programmable pump in pat have caused intolerable side effects and the prescription b) Medsurge to be Principal Supply on 1 December 2021			agents have been ineffective or
DANTROLENE			
Cap 25 mg	97.50	100	✓ Dantrium
			✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			
Tab 100 mg Norflex to be Principal Supply on 1 January 2022	20.76	100	✓ Norflex

NERVOUS SYSTEM				
	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
Agents for Parkinsonism and Related Disorders	;			
Dopamine Agonists and Related Agents				
AMANTADINE HYDROCHLORIDE				
▲ Cap 100 mg	38.24	60	•	Symmetrel
APOMORPHINE HYDROCHLORIDE			_	
▲ Inj 10 mg per ml, 2 ml ampoule		5	_	Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	•	<u>Movapo</u>
BROMOCRIPTINE MESYLATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were prescription is endorsed accordingly. Pharmacists may anno prior dispensing of bromocriptine mesylate. * Tab 2.5 mg	tate the prescription		idorsed w	
	32.08	100	•	Apo-Bromocriptine
(Parlodel S29 Tab 2.5 mg to be delisted 1 March 2022) (Apo-Bromocriptine Tab 2.5 mg to be delisted 1 March 2022) ENTACAPONE Tab 200 mg	22.00	100		Entapone
LEVODOPA WITH BENSERAZIDE	22.00	100	•	Ептаропе
* Tab dispersible 50 mg with benserazide 12.5 mg	13 25	100	/	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100		Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	1	Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100		Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	•	Madopar 250
LEVODOPA WITH CARBIDOPA				
* Tab 100 mg with carbidopa 25 mg		100		Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg	43.65	100		Sinemet CR
* Tab 250 mg with carbidopa 25 mg	38.39	100	•	Sinemet
PRAMIPEXOLE HYDROCHLORIDE			_	
▲ Tab 0.25 mg		100	_	Ramipex
▲ Tab 1 mg	20.73	100	•	Ramipex
RASAGILINE				
* Tab 1 mg	53.50	30	•	Azilect S29
ROPINIROLE HYDROCHLORIDE				
▲ Tab 0.25 mg		84		Ropin
A. Teli dama	3.39	100	_	Mylan S29
▲ Tab 1 mg		84		Ropin
A Tob 0 mg	4.70	100		Mylan S29
▲ Tab 2 mg		84 84		Ropin Ropin
· ·	12.00	04	•	ПОРШ
SELEGILINE HYDROCHLORIDE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were	taking salegiline bud	rochl	orida prior	to 1 August 2021 and the
prescription is endorsed accordingly. Pharmacists may anno prior dispensing of selegiline hydrochloride.				
* Tab 5 mg	22.00	100	•	Apo-Selegiline

1	fully subsidised
Pri	ncipal Supply

S29 S29 ✓ Eldepryl S29

48.00

				VOOD OTOTEM
	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
TOLCAPONE A Tab 100 mg	152.38	100	√ T	'asmar
Anticholinergics				
BENZATROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 10 inj available on a PSO b) Only on a PSO		60 5		Benztrop Phebra
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	√ K	Cemadrin
Agents for Essential Tremor, Chorea and Relati	ed Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail phar Wastage claimable Tab 50 mg Rilutek to be Principal Supply on 1 December 2021	•	56	✓ F	Rilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialifolium oritoriogist or respiratory specialifolium oritoriogist.	st. Approvals valid fo	r 6 month	ns for app	plications meeting the

following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory: or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory: or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE

Tab 25 mg	91.10	112	✓ Motetis
-----------	-------	-----	-----------

NERVOUS SYSTEM					
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
Anaesthetics					
Local					
LIDOCAINE [LIGNOCAINE] Gel 2%, tube — Subsidy by endorsement		30 ml		Kylocaine 2% Jelly	
Gel 2%, 11 ml urethral syringe – Subsidy by endorsement a) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral, cervical or r		10 and the		ion is endorsed	
accordingly. LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE					
Oral (gel) soln 2%	38.00	200 ml	1	Mucosoothe	
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO		25	√ i	Lidocaine-Baxter Lidocaine-Claris	
	17.50 (35.00)	50)	Kylocaine	
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO		25	-	Lidocaine-Baxter Lidocaine-Claris	
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	(20.00)	5		Kylocaine	
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO Inj 2%, 20 ml vial – Up to 5 inj available on a PSO		5 5	√ <u>[</u>	<u>Lidocaine-Claris</u> <u>Lidocaine-Baxter</u> Lidocaine-Claris	
(Lidocaine-Claris Inj 1%, 5 ml ampoule to be delisted 1 January 2 (Lidocaine-Claris Inj 2%, 5 ml ampoule to be delisted 1 January 2					
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –					
Subsidy by endorsementa) Up to 5 each available on a PSO b) Subsidised only if prescribed for urethral or cervical a		10 e preso		Pfizer endorsed accordingly.	
Topical Local Anaesthetics					
➤ SA0906 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid condition requiring frequent injections or venepuncture. Renewal from any relevant practitioner. Approvals valid for 2 year.	•	•			
benefiting from treatment.					

LIDOCAINE [LIGNOCAINE] — Special Authority see SA0906 above -	- Retall phar	macy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special Authority	see SA0906	above – Retai	I pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

NERVOUS SYSTEM

Subsidy (Manufacturer's Price)

Fully Subsidised Per

90

✓ Acupan

Brand or Generic Manufacturer

Analgesics

NEFOPAM HYDROCHLORIDE

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 113

Tab 30 mg23.40

Non-opioid Analgesics		
ASPIRIN * Tab dispersible 300 mg - Up to 30 tab available on a PSO4.50	100	✓ Ethics Aspirin
CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or diabetic periphera accordingly.	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	45 g OP 57 g OP	✓ <u>Zostrix HP</u> ✓ Rugby Capsaicin Topical Cream \$29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PARACETAMOL				
Tab 500 mg - blister pack	0.50	20	✓	Medco
•			✓	Pharmacy Health
	1.12		1	Ethics Paracetamol
				Classic
	2.48	100	✓	Pharmacy Health
	5.01	50	✓	Panadol
	11.75	96	✓	Panadol Mini Caps
	19.75	1,000	/	Pacimol
	24.82		1	Paracetamol
				Pharmacare
			✓	Pharmacare

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.

Tab 500 mg - bottle pack - Maximum of 300 tab per			
prescription; can be waived by endorsement	17.92	1,000	✓ Noumed
			Paracetamol
	24.82		Paracetamol
			Pharmacare

a)

- 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.
- 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing. b) Noumed Paracetamol to be Sole Supply on 1 February 2022

*	Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓ Paracare
	a) Up to 200 ml available on a PSO			
	b) Not in combination			
*	Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓ Paracare Double
				<u>Strength</u>
	a) Up to 100 ml available on a PSO			
	b) Not in combination			
*	Suppos 125 mg	3.29	10	✓ Gacet
*	Suppos 250 mg	3.79	10	✓ Gacet
*	Suppos 500 mg	12.40	50	✓ Gacet
(M	edco Tab 500 mg - blister pack to be delisted 1 February 2022)			

(Pharmacy Health Tab 500 mg - blister pack to be delisted 1 February 2022)

(Ethics Paracetamol Classic Tab 500 mg - blister pack to be delisted 1 February 2022)

(Pharmacy Health Tab 500 mg - blister pack to be delisted 1 February 2022)

(Panadol Tab 500 mg - blister pack to be delisted 1 February 2022)

(Panadol Mini Caps Tab 500 mg - blister pack to be delisted 1 February 2022)

(Paracetamol Pharmacare Tab 500 mg - blister pack to be delisted 1 February 2022)

(Pharmacare Tab 500 mg - blister pack to be delisted 1 February 2022)

(Paracetamol Pharmacare Tab 500 mg - bottle pack to be delisted 1 February 2022)

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully Brand or idised Generic Manufacturer
Opioid Analgesics			
CODEINE PHOSPHATE - Safety medicine; prescriber may dete	ermine dispensing fre	quency	
Tab 15 mg		100	✓ PSM
Tab 30 mg		100	✓ PSM
Tab 60 mg		100	✓ PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8 60	60	✓ DHC Continus
FENTANYL			
a) Only on a controlled drug form b) No potient as payment payable.			
b) No patient co-payment payablec) Safety medicine; prescriber may determine dispensing from	earrency		
Inj 50 mcg per ml, 2 ml ampoule		10	✓ Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	✓ Boucher and Muir
Patch 12.5 mcg per hour		5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 20			,,
Patch 25 mcg per hour		5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 20)22		•
Patch 50 mcg per hour	9.49	5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 20			_
Patch 75 mcg per hour		5	Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 20		_	4
Patch 100 mcg per hour		5	Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 20)22		
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	'		
d) Extemporaneously compounded methadone will only be	reimbursed at the rate	e of the che	eapest form available
(methadone powder, not methadone tablets).e) For methadone hydrochloride oral liquid refer Standard F	formulae page 045		
Tab 5 mg		10	✓ Methatabs
Oral lig 2 mg per ml		200 ml	✓ Biodone
Biodone to be Principal Supply on 1 January 2022			Diodolio
Oral lig 5 mg per ml	6.40	200 ml	✓ Biodone Forte
Biodone Forte to be Principal Supply on 1 January 2022	!		
Oral liq 10 mg per ml	7.50	200 ml	✓ Biodone Extra Forte
Biodone Extra Forte to be Principal Supply on 1 January	/ 2022		
Inj 10 mg per ml, 1 ml	61.00	10	✓ AFT
MORPHINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	'		
Oral liq 1 mg per ml		200 ml	✓ RA-Morph
Oral liq 2 mg per ml		200 ml	✓ RA-Morph
Oral liq 5 mg per ml	19.44 2	200 ml	✓ Ordine \$29
			✓ RA-Morph
Oral liq 10 mg per ml	27.74 2	200 ml	✓ Ordine S29
			✓ RA-Morph

NERVOUS SYSTEM

	Subsidy Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing free	uency			
Tab immediate-release 10 mg	2.80	10	✓ Se	evredol
Tab immediate-release 20 mg	5.52	10	✓ Se	evredol
Cap long-acting 10 mg	2.05	10	✓ m	-Eslon
Cap long-acting 30 mg	3.00	10	✓ m	-Eslon
Cap long-acting 60 mg	6.12	10	✓ m	-Eslon
Cap long-acting 100 mg	7.13	10	✓ <u>m</u>	-Eslon
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC)6.99	5		BL Morphine Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	6O5.61	5	✓ DI	BL Morphine Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	O7.08	5		BL Morphine Sulphate
Ini 30 mg per ml. 1 ml ampoule – Up to 5 ini available on a PS	O7.28	5	✓ DI	BL Morphine

Sulphate

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	
	\$	Per	•	Manufacturer
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f	roguonov			
Tab controlled-release 5 mg		20	./	Oxycodone Sandoz
Tab controlled-release 5 mg	3.01	28		Oxycodone Sandoz
	3.01	20	•	•
			_	S29 S29
Tab controlled-release 10 mg		20		Oxycodone Sandoz
	3.23	30	•	Oxycodone Sandoz
				S29 S29
	5.38	50	✓	Oxycodone Sandoz
				S29 S29
	10.75	100	1	Oxycodone Sandoz
	10.75	100	•	•
			_	S29 S29
	11.50	28		OxyContin
Tab controlled-release 20 mg	2.15	20		Oxycodone Sandoz
	5.38	50	✓	Oxycodone Sandoz
				S29 S29
	10.75	100	1	Oxycodone Sandoz
	10.70			S29 S29
	10.05	00		
Tab acostrollad valance 40 mm	13.25	28		OxyContin
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg	1.88	20	•	OxyNorm
OxyNorm to be Principal Supply on 1 December 2021			_	
Cap immediate-release 10 mg	3.32	20	•	OxyNorm
OxyNorm to be Principal Supply on 1 December 2021				
Cap immediate-release 20 mg	5.23	20	✓	OxyNorm
OxyNorm to be Principal Supply on 1 December 2021				
Oral liq 5 mg per 5 ml	11.20	250 m		OxyNorm
Inj 10 mg per ml, 1 ml ampoule	7.28	5	✓	OxyNorm
Inj 10 mg per ml, 2 ml ampoule	14.36	5	✓	OxyNorm
Inj 50 mg per ml, 1 ml ampoule	30.60	5	✓	OxyNorm
PARACETAMOL WITH CODEINE - Safety medicine; prescribe		ancin	r fraguanc	
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
* Tab paracetation 500 mg with codeline phosphate 6 mg	20.51	1,000	•	
				Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f	requency			
Tab 50 mg	4.70	10	✓	PSM
PSM to be Principal Supply on 1 January 2022				
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a	PSO 29.88	5	/	DBL Pethidine
, , , , , , , , , , , , , , , , , , , ,				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 30.72	5	1	DBL Pethidine
ing 55 mg por mi, 2 mi ampould to p to 5 mg available on a	. 5555.72	J	•	Hydrochloride
				riyarocinonae

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
FRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	✓	Tramal SR 100
Tab sustained-release 150 mg	2.10	20	✓	Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg		100		Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determ				
Tab 10 mg	2.49	100	✓	Arrow-Amitriptyline
Tab 25 mg	1.51	100		Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; pr	escriber may determine d	isper	sing freau	ency
Tab 10 mg	•	30		Clomipramine Teva
	13.99	100		Apo-Clomipramine
Tab 25 mg		100		Apo-Clomipramine
	11.99	30		Clomipramine Teva
Apo-Clomipramine Tab 10 mg to be delisted 1 February 202 Apo-Clomipramine Tab 25 mg to be delisted 1 February 202	22)	00		olollipianinio rova
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy b				
a) Safety medicine; prescriber may determine dispensir	•			
, , , , , , , , , , , , , , , , , , , ,	0 1 7			
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth]	ho were taking dosulepin Pharmacists may annotate			
 b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. 	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride.		prescriptio	n as endorsed where th
 Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth 	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride. 3.85	the	prescriptio	n as endorsed where th
 Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg 	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83	30 50	prescriptio	n as endorsed where th Dosulepin Mylan Dosulepin Mylan S29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83	30 50	prescriptio	n as endorsed where th Dosulepin Mylan Dosulepin Mylan S29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83 riber may determine dispe5.48 10.96	30 50 nsing 50 100	prescriptio	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83 riber may determine dispe5.48 10.96	30 50 nsing	prescriptio	n as endorsed where th Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83 riber may determine dispe5.48 10.968.80	30 50 nsing 50 100 50	prescriptio	n as endorsed where the Dosulepin Mylan Mylan S29 y Tofranil Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50	prescription frequency nsing frequency	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Lency
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride3.857.83 riber may determine dispe5.48 10.968.80 rescriber may determine d2.44	30 50 nsing 50 100 50	rescription frequency	n as endorsed where the Dosulepin Mylan Mylan S29 y Tofranil Tofranil Tofranil
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 disper	rescription frequency	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Lency Norpress
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 disper	rescription frequency	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Lency Norpress
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 disper	orescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Lency Norpress
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 disper 100 180	prescription	n as endorsed where the Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Jency Norpress Norpress
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 nsing 50 100 50 100 180 28 50	prescription	Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate \$29 \$29 Parnate
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 nsing 50 100 50 100 180 28	orescription	Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate \$29 \$29 Parnate Parnate \$29 \$29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 nsing 50 100 50 100 180 28 50	orescription	Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate \$29 \$29 Parnate
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 nsing 50 100 50 100 180 28 50	orescription	Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate \$29 \$29 Parnate Parnate \$29 \$29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. If exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 sdisper 100 180	orescription	Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate S29 S29 Parnate Parnate S29 S29 Parnate S29 S29 Parnate S29 S29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 nsing 50 100 50 100 180 28 50	prescription	Dosulepin Mylan Dosulepin Mylan \$29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate \$29 \$29 Parnate Parnate \$29 \$29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	the 30 50 solution of 50 100 50 solution of 50 100 100 180 100 100 100 100 100 100 10	prescription	Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate S29 S29 Parnate Parnate S29 S29 Parnate S29 S29 Parnate S29 S29 Parnate S29 S29
b) Subsidy by endorsement – Subsidised for patients w 2019 and the prescription is endorsed accordingly. I exists a record of prior dispensing of dosulepin [doth Tab 75 mg	ho were taking dosulepin Pharmacists may annotate iepin] hydrochloride	30 50 nsing 50 100 50 sdisper 100 180	prescription	Dosulepin Mylan Dosulepin Mylan S29 y Tofranil Tofranil Tofranil Jency Norpress Norpress Parnate S29 S29 Parnate Parnate S29 S29 Parnate S29 S29 Parnate S29 S29

	Subsidy (Manufacturer's Price) \$	F Subsidi Per	ully Brand or sed Generic Manufacturer
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			4
* Tab 20 mg	1.91	84	✓ PSM Citalopram
ESCITALOPRAM No. Tab. 10 and	4.07	00	/ Faciliation
* Tab 10 mg	1.07	28	✓ Escitalopram (Ethics)
* Tab 20 mg	1.92	28	✓ Escitalopram (Ethics)
FLUOXETINE HYDROCHLORIDE			 ,
* Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	1.98	30	✓ Fluox
 When prescribed for a patient who cannot swallow accordingly; or 	whole tablets or caps	ules and the	e prescription is endorsed
When prescribed in a daily dose that is not a multip endorsed. Note: Tablets should be combined with			
Cap 20 mgPAROXETINE	2.91	84	✓ Fluox
* Tab 20 mg	1.20	30	✓ Paxtine
· · · · · · · · · · · · · · · · · · ·	3.61	90	✓ Loxamine
(Paxtine Tab 20 mg to be delisted 1 January 2022)			
SERTRALINE			
* Tab 50 mg		30	✓ <u>Setrona</u>
* Tab 100 mg	1.61	30	✓ <u>Setrona</u>
Other Antidepressants			
MIRTAZAPINE			
Tab 30 mg	2.60	28	✓ Noumed
N	2.63	30	✓ Apo-Mirtazapine
Noumed to be Sole Supply on 1 January 2022	2.45	28	✓ Noumed
Tab 45 mg	3.45 3.48	30	✓ Apo-Mirtazapine
Noumed to be Sole Supply on 1 January 2022	0.40	00	Apo inintazapino
(Apo-Mirtazapine Tab 30 mg to be delisted 1 January 2022) (Apo-Mirtazapine Tab 45 mg to be delisted 1 January 2022)			
VENLAFAXINE			4
* Cap 37.5 mg		84	✓ Enlafax XR
* Cap 75 mg		84 84	✓ Enlafax XR ✓ Enlafax XR
* Cap 150 mg	11.10	U4	▼ LilididX Aft
Antiepilepsy Drugs			
Agents for Control of Status Epilepticus			
CLONAZEPAM – Safety medicine; prescriber may determine dis Inj 1 mg per ml, 1 ml(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)		5	✓ Rivotril

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy	-) 0 !	Fully Brand or
	(Manufacturer's Price \$	e) Sui Per	bsidised Generic Manufacturer
AZEPAM – Safety medicine; prescriber may determine disp	ensing frequency		
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement.		5	✓ Hospira
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
c) PSO must be endorsed "not for anaesthetic proced	dures".		
Rectal tubes 5 mg - Up to 5 tube available on a PSO	43.50	5	✓ Stesolid
ENYTOIN SODIUM			
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	a PSO88.63	5	✓ Hospira
Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a			•
PSO		5	✓ Hospira
ontrol of Epilepsy			
RBAMAZEPINE			_
Tab 200 mg		100	✓ Tegretol
Tab long-acting 200 mg		100	✓ Tegretol CR
Tab 400 mg		100	✓ Tegretol
Tab long-acting 400 mg		100	✓ Tegretol CR
Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
OBAZAM - Safety medicine; prescriber may determine disp	pensing frequency		
Tab 10 mg	9.12	50	✓ Frisium
ONAZEPAM - Safety medicine; prescriber may determine	dispensing frequency		
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
THOSUXIMIDE			
Cap 250 mg	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml		200 ml	✓ Zarontin
BAPENTIN			
Note: Not subsidised in combination with subsidised prega	ahalin		
Cap 100 mg		100	✓ Apo-Gabapentin
oup 100 mg	6.45	100	✓ Apo-Gabaperiiiii ✓ Nupentin
Cap 300 mg		100	✓ Apo-Gabapentin
- · T - · · · · •	8.45		✓ Nupentin
Cap 400 mg		100	✓ Apo-Gabapentin
, ,	10.26		✓ Nupentin
po-Gabapentin Cap 100 mg to be delisted 1 February 2022)			•
po-Gabapentin Cap 300 mg to be delisted 1 February 2022)			
po-Gabapentin Cap 400 mg to be delisted 1 February 2022)			
COSAMIDE - Special Authority see SA1125 below - Retail	pharmacy		
Tab 50 mg		14	✓ Vimpat
Tab 100 mg		14	✓ Vimpat
····	200.24	56	✓ Vimpat
Tab 150 mg		14	✓ Vimpat
•	300.40	56	✓ Vimpat
Tab 200 mg		56	✓ Vimpat

■ SA1125 Special Authority for Subsidy
Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

I AMOTRIGINE

- 1 Patient has partial-onset epilepsy; and
 - 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LA	WOTHIGHT			_	
	Tab dispersible 2 mg	55.00	30	/	Lamictal
	Tab dispersible 5 mg	50.00	30	1	Lamictal
*	Tab dispersible 25 mg	2.76	56	1	Logem
*	Tab dispersible 50 mg	3.31	56	•	Logem
*	Tab dispersible 100 mg	4.40	56	1	Logem
۱F	VETIRACETAM				
	Tab 250 mg	4.99	60	/	Everet
	Tab 500 mg		60		Everet
	Tab 750 mg		60		Everet
	Tab 1,000 mg		60		Everet
	Oral lig 100 mg per ml		300 ml OP	1	Levetiracetam-AFT
ווח	ENOBARBITONE				
ГΠ	For phenobarbitone oral liquid refer Standard Formulae, pa	000 04E			
×	Tab 15 mg	•	500	_	PSM
*	· ·		500		PSM
*	Tab 30 mg	40.00	500	v	PSIVI
	ENYTOIN SODIUM				
*	Tab 50 mg		200		Dilantin Infatab
	Cap 30 mg		200		Dilantin
	Cap 100 mg		200		Dilantin
*	Oral liq 30 mg per 5 ml	22.03	500 ml	•	Dilantin
PR	EGABALIN				
	Note: Not subsidised in combination with subsidised gaba	pentin			
*		•	56	1	Pregabalin Pfizer
*	Cap 75 mg		56		Pregabalin Pfizer
*			56		Lyrica
					Pregabalin Pfizer
*	Cap 300 mg	7.38	56		Pregabalin Pfizer
DD	IMIDONE				Ū
	Tab 250 mg	17.05	100	_	Apo-Primidone
	· ·	17.25	100	•	Apo-Fillilluolle
SO	DIUM VALPROATE				
	Tab 100 mg		100		Epilim Crushable
	Tab 200 mg EC		100		Epilim
	Tab 500 mg EC		100		Epilim
*	Oral liq 200 mg per 5 ml	20.48	300 ml		Epilim S/F Liquid
					Epilim Syrup
*	Inj 100 mg per ml, 4 ml	41.50	1	•	Epilim IV

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
STIRIPENTOL - Special Authority see SA1330 below - Retail p	harmacy				
Cap 250 mg	509.29	60	1	Diacomit \$29	
Powder for oral liq 250 mg sachet	509.29	60	•	Diacomit S29	

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

▲ Tab 500 mg		100	✓ Sabril
VIGABATRIN - Special Authority see SA1997 below - Reta			•
▲ Sprinkle cap 25 mg	26.04	60	✓ Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓ Topamax
	129.85		✓ Topamax
ŭ			✓ Topiramate Actavis
▲ Tab 200 mg	55.19	60	✓ Arrow-Topiramate
	75.25		✓ Topamax
			Topiramate Actavis
▲ Tab 100 mg	31.99	60	Arrow-Topiramate
	44.26		✓ Topamax
			Topiramate Actavis
▲ Tab 50 mg	18.81	60	Arrow-Topiramate
	26.04		Topamax
			Topiramate Actavis
▲ Tab 25 mg	11.07	60	Arrow-Topiramate

⇒SA1997 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy: and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

continued Vigabatrin is associated with a risk of irreversible visual field defects, v Renewal from any relevant practitioner. Approvals valid without furthe following criteria: Both: 1 The patient has demonstrated a significant and sustained imprecation of treatment with vigabatrin; or 2.1 Patient is receiving regular automated visual field testing of treatment with vigabatrin; or 2.2 It is impractical or impossible (due to comorbid condition Notes: As a guideline, clinical trials have referred to a notional 50% reference.	er renewal unless ovement in seizu g (ideally every 6 ns) to monitor the eduction in seizu	ymptomati s notified f ure rate or 6 months) e patient's re frequen	or appl severity on an o	ications meeting the y and or quality of life; ar engoing basis for duration fields
Vigabatrin is associated with a risk of irreversible visual field defects, versible from any relevant practitioner. Approvals valid without further following criteria: Both: 1 The patient has demonstrated a significant and sustained improversible: 2.1 Patient is receiving regular automated visual field testing of treatment with vigabatrin; or 2.2 It is impractical or impossible (due to comorbid condition)	er renewal unless ovement in seizu g (ideally every 6 ns) to monitor the eduction in seizu	s notified fure rate or months) e patient's re frequen	or appl severity on an o	ications meeting the y and or quality of life; ar engoing basis for duration fields
The patient has demonstrated a significant and sustained improvements. Either: 2.1 Patient is receiving regular automated visual field testing of treatment with vigabatrin; or 2.2 It is impractical or impossible (due to comorbid condition).	g (ideally every 6 ns) to monitor the eduction in seizu	6 months) e patient's re frequen	on an c	ongoing basis for duration
	eduction in seizu	re frequen		
anticonvulsant therapy and have assessed quality of life from the patie /igabatrin is associated with a risk of irreversible visual field defects, v			c in the	
Antimigraine Preparations				
For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 1	13			
Acute Migraine Treatment				
RIZATRIPTAN Tab orodispersible 10 mg	3.65	30	✓ <u>R</u>	tizamelt
SUMATRIPTAN Tab 50 mg	14.41	90 100		umagran po-Sumatriptan
Tab 100 mg	22.68	90	✓ S	umagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per	46.23	100	V A	po-Sumatriptan
prescription	34.00	2 OP	√ <u>Ir</u>	<u>migran</u>
Prophylaxis of Migraine				
For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTE PIZOTIFEN				
* Tab 500 mcg	23.21	100	√ S	andomigran
Antinausea and Vertigo Agents				
For Antispasmodics refer to ALIMENTARY TRACT, page 8				
APREPITANT – Special Authority see SA0987 below – Retail pharma Cap 2 × 80 mg and 1 × 125 mg Emend Tri-Pack to be Principal Supply on 1 December 2021 SA0987 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid for	30.00	3 OP		mend Tri-Pack

84 ✓ Vergo 16

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic

emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE Tab 50 mg Nausicalm to be Principal Supply on 1 December 2021	0.49	10	✓ N	Nausicalm
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml	21.53	10	✓ <u>F</u>	lameln_
DOMPERIDONE * Tab 10 mg	2.85	100	√ F	Pharmacy Health
HYOSCINE HYDROBROMIDE * Inj 400 mcg per ml, 1 ml ampoule		10	✓ N	Martindale \$29
Patch 1.5 mg – Special Authority see SA1998 below – Retail pharmacy		2	√ 9	Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

ME	TOCLOPRAMIDE HYDROCHLORIDE		
*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50 IDANSETRON	10	✓ <u>Pfizer</u>
*	Tab 4 mg2.68	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ <u>Ondansetron</u> <u>ODT-DRLA</u>
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ Ondansetron ODT-DRLA
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal	50	Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
- 1			✓ Stemetil
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	Stemetii

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may deterr	nine dispensing frequency	1	
Tab 100 mg	5.15	30	✓ Sulprix
·	17.16	100	✓ Amisulpride Mylan \$29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per		Manufacturer
ARIPIPRAZOLE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 5 mg	17.50	30	1	Aripiprazole Sandoz
Tab 10 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 15 mg	17.50	30	✓	Aripiprazole Sandoz
Tab 20 mg	17.50	30	1	Aripiprazole Sandoz
Tab 30 mg	17.50	30	✓	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may determi	ne dis	pensina fr	eauencv
Tab 10 mg - Up to 30 tab available on a PSO		100		Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100		Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100		Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10		Largactil
CLOZAPINE - Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing freq	III Ancv			
Tab 25 mg	•	50	1	Clozaril
1 ab 25 mg	6.69	50		Clopine
	11.36	100		Clozaril
	13.37	100		Clopine
Tab 50 mg		50		Clopine
1 ab 30 mg	17.33	100	_	Clopine
Tab 100 mg		50	_	Clozaril
Tab Too mg	17.33	50		Clopine
	29.45	100		Clozaril
	34.65	100		Clopine
Tab 200 mg		50		Clopine
Tab 200 Mg	69.30	100		Clopine
Suspension 50 mg per ml		100 m		Clopine
Ouspension of mg per mi	67.62	10011		Versacloz
HALODERIDOL Octobrondicion and disconnection			•	VCIGUOIOZ
HALOPERIDOL – Safety medicine; prescriber may determine		400	,	0
Tab 500 mcg – Up to 30 tab available on a PSO		100		<u>Serenace</u>
Tab 1.5 mg — Up to 30 tab available on a PSO		100		Serenace Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace Serenace
Oral lin O man man and	29.72	100		Serenace Serenace
Oral liq 2 mg per ml — Up to 200 ml available on a PSO		100 m		Serenace Serenace
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a l		10		<u>Serenace</u>
LEVOMEPROMAZINE - Safety medicine; prescriber may dete		uency		
Tab 25 mg (33.8 mg as a maleate)	16.10	100	1	Nozinan (Swiss)
Tab 25 mg as a maleate		100		<u>Nozinan</u>
Tab 100 mg (135 mg as a maleate)		100	_	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	•	<u>Nozinan</u>
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	prescriber may deterr	nine d	ispensing	frequency
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
LITHIUM CARBONATE - Safety medicine; prescriber may det	armina disnansina frac	III LANCI	,	
Tab long-acting 400 mg		100		Priadel
Cap 250 mg		100	_	Douglas
		100	•	- Jugius
OLANZAPINE – Safety medicine; prescriber may determine di				7 !
Tab 2.5 mg		28		Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28		Zypine ODT
Tab orodispersible 10 mg	2.38	28	•	Zypine ODT

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PERICYAZINE - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 2.5 mg	10.49	84	✓	Neulactil
•	12.49	100	✓	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
•	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensina frequency			
Tab 25 mg	•	90	/	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	1	Quetapel
RISPERIDONE – Safety medicine; prescriber may determine dis	spensing frequency			
Tab 0.5 mg		60	/	Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60	/	Risperidone (Teva)
Tab 3 mg		60	1	Risperidone (Teva)
Tab 4 mg	3.42	60	1	Risperidone (Teva)
Oral liq 1 mg per ml		30 m	✓	Risperon
ZIPRASIDONE – Safety medicine; prescriber may determine dis	pensing frequency			
Cap 20 mg		60	1	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60	_	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre		a die	nancina fra	adilenov
Tab 10 mg	•	100	•	Clopixol
1 ab 10 mg	51.45	100	•	Ciopixoi

Depot Injections

• •		
FLUPENTHIXOL DECANOATE - Safety medicine; prescriber may determine disp	ensing freq	uency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispe	ensing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO55.90	5	✓ Haldol Concentrate
, ,		✓ Haldol
		Decanoas \$29
OLANZAPINE - Special Authority see SA1428 below - Retail pharmacy		
Safety medicine; prescriber may determine dispensing frequency		
Inj 210 mg vial252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial504.00	1	✓ Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
- 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe		1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing for	requency		
Inj 25 mg vial	135.98	1	Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.



	Gubsidy		uny	Diana oi
	(Manufacturer's Price)	Subsi	dised	Generic
	\$	Per	<u> </u>	Manufacturer
ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescribe	r may determine disn	ensing fred	nuency	
Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO	,	5		lopixol
ing 200 mg per mi, 1 mi - Op to 3 mg available on a 1 00	10.00	3		ТОРІХОІ
Anxielyties				
Anxiolytics				
BUSPIRONE HYDROCHLORIDE				
	20.22	100	√ 0	rion
* Tab 10 mg		100	✓ 0	rion
CLONAZEPAM – Safety medicine; prescriber may determine dis	pensing frequency			
Tab 500 mcg	5.64	100	✓ P	axam
Tab 2 mg	10.78	100	✓ P	axam
DIAZEPAM - Safety medicine; prescriber may determine dispen	sing fraguancy			
		500	✓ A	rrow-Diazepam
Tab 2 mg			_	
Tab 5 mg	73.00	500	V A	rrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency			
Tab 1 mg	9.72	250	✓ A	tivan
Ativan to be Principal Supply on 1 December 2021				
Tab 2.5 mg	12.50	100	✓ A	tivan
Ativan to be Principal Supply on 1 December 2021				
,	! f			
OXAZEPAM – Safety medicine; prescriber may determine dispe	. ,	400		D
Tab 10 mg		100		x-Pam
Tab 15 mg	8.53	100	√ 0	x-Pam

Subsidy

Fully

Brand or

Multiple Sclerosis Treatments

SA2051 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or

			NEI	RVOUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
continued				
5.5.2 Each significant relapse is a recurrent pare trigeminal neuralgia, Lhermitte's symptom		ultipl	e sclerosis	(tonic seizures/spasms,
6 Evidence of new inflammatory activity on an MR scan wit7 Any of the following:	thin the past 24 months	s; and	d	
 7.1 A sign of that new inflammatory activity is a gadol 7.2 A sign of that new inflammatory activity is a lesior 7.3 A sign of that new inflammatory is a T2 lesion with 7.4 A sign of that new inflammatory activity is a proming a recent relapse that occurred within the last 2 ye 7.5 A sign of that new inflammatory activity is new T2 	n showing diffusion resi n associated local swel inent T2 lesion that cle ars; or	trictio lling; arly is	or s responsil	
Note: Natalizumab can only be dispensed from a pharmacy regoperated by the supplier. Treatment on two or more funded mul Renewal — (Multiple sclerosis) only from a neurologist or ger had an EDSS score of 0 to 6.0 (inclusive) with or without the use (i.e. the patient has walked 100 metres or more with or without Note: Natalizumab can only be dispensed from a pharmacy regoperated by the supplier. Treatment on two or more funded multiple of the supplier is the supplier of the supplier is the supplier of the supplier is the supplier in the supplier in the supplier is the	Itiple sclerosis treatmenteral physician. Approse of unilateral or bilateral distribution the last six more istered in the Tysabri Attiple sclerosis treatments.	nts si vals v al aid nths). Austra nts si	imultaneou valid for 12 ds at any tii alasian Pre imultaneou	sly is not permitted. months where patient has me in the last six months escribing Programme
DIMETHYL FUMARATE – Special Authority see SA2051 on the	e previous page – Reta	ail pha	armacy	
a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Cap 120 mg Cap 240 mg	520.00	eous 14 56	1	rmitted. Tecfidera Tecfidera
FINGOLIMOD – Special Authority see SA2051 on the previous	page – Retail pharmad	су		
a) Wastage claimable b) Note: Treatment on two or more funded multiple scleros Cap 0.5 mg		eous 28		rmitted. Gilenya
GLATIRAMER ACETATE – Special Authority see SA2051 on the Note: Treatment on two or more funded multiple sclerosis to Inj 40 mg prefilled syringe	reatments simultaneou	tail pl usly is 12	s not permi	tted. Copaxone
INTERFERON BETA-1-ALPHA – Special Authority see SA205 Note: Treatment on two or more funded multiple sclerosis t Inj 6 million iu prefilled syringe	on the previous page reatments simultaneou1,170.00		s not permi	
INTERFERON BETA-1-BETA – Special Authority see SA2051 Note: Treatment on two or more funded multiple sclerosis t Inj 8 million iu per 1 ml	reatments simultaneou		s not permi	
NATALIZUMAB – Special Authority see SA2051 on the previou Note: Treatment on two or more funded multiple sclerosis t Inj 20 mg per ml, 15 ml vial	reatments simultaneou			tted. Tysabri
OCRELIZUMAB – Special Authority see SA2051 on the previous Note: Treatment on two or more funded multiple sclerosis to Inj 30 mg per ml, 10 ml vial	reatments simultaneou			tted. Ocrevus

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

TERIFLUNOMIDE - Special Authority see SA2051 on the previous page - Retail pharmacy

✓ Aubagio

28

a) Wastage claimable



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy

Tab modified-release 2 mg - No more than 5 tab per day28.22

30

Circadin

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine disper	nsing frequency		
Inj 1 mg per ml, 5 ml ampoule	3.95	10	Mylan Midazolam
	5.50		✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be e	endorsed for stat	us epilepticu	ıs use only.
Inj 5 mg per ml, 3 ml ampoule	4.50	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available o	on		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be e	endorsed for state	us epilepticu	is use only.
(Mylan Midazolam Inj 1 mg per ml, 5 ml ampoule to be delisted 1	January 2022)		·
PHENOBARBITONE SODIUM - Special Authority see SA1386 be	<mark>elow</mark> – Retail pha	armacy	
Inj 200 mg per ml, 1 ml ampoule	78.20	10	✓ Max Health S29
⇒SA1386 Special Authority for Subsidy			

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determ	ine dispensing frequency		
Tab 10 mg	1.33	25	✓ Normison
TRIAZOLAM - Safety medicine; prescriber may determine	ne dispensing frequency		
Tab 125 mcg	5.10	100	
	(9.85)		Hypam
Tab 250 mcg	4.10	100	
·	(11.20)		Hypam

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ZOPICLONE – Safety medicine; prescriber may determine dis Tab 7.5 mg		500	/	Zopiclone Actavis
Stimulants/ADHD Treatments				
ATOMOXETINE				
Cap 10 mg	18.41	28	✓	Generic Partners
	107.03		✓	Strattera
Cap 18 mg	27.06	28	✓	Generic Partners
	107.03		✓	Strattera
Cap 25 mg	29.22	28	✓	Generic Partners
Cap 40 mg		28	✓	Generic Partners
	107.03		✓	Strattera
Cap 60 mg	46.51	28	✓	Generic Partners
Cap 80 mg	56.45	28	✓	Generic Partners
Cap 100 mg	58.48	28	✓	Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see SA114	9 below – Retail pharma	acv		
a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing Tab 5 mg	requency	100	/	PSM
PSM to be Principal Supply on 1 January 2022				

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.



	\$	Per	✓ Manufacturer
METHYLPHENIDATE HYDROCHLORIDE - Special Authority s	ee SA1964 below	- Retail pha	ırmacy
a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing fre	equency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg	3.00	30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg	7.75	30	Methylphenidate ER
			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg		30	Methylphenidate ER
			- Teva
Tab extended-release 36 mg	15.50	30	Methylphenidate ER
•			- Teva
Tab extended-release 54 mg	22.25	30	Methylphenidate ER
G			- Teva

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

b) Salety medicine, prescriber may determine dispensing i	requericy		
Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	Concerta
Tab extended-release 54 mg		30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see SA1999 below – Retail pharmacy
Tab 100 mg29.13 60 ✓ Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or

NERVOUS SYSTEM

Subsidy (Manufacturer's	Price) Sub	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below - Retail	pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	Rivastigmine Patch BNM 5
	48.75		Generic Partners
Patch 9.5 mg per 24 hour	38.00	30	Rivastigmine Patch BNM 10
	48.75		✓ Generic Partners
(Generic Partners Patch 4.6 mg per 24 hour to be delisted 1 Feb	ruary 2022)		
(Generic Partners Patch 9.5 mg per 24 hour to be delisted 1 Feb	ruary 2022)		

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine: prescriber may determine dispensing frequency

-,					
Tab su	ıblingual 2	2 mg with	naloxone 0.5 mg.	1	8.37 28

Tab sublingual 8 mg with naloxone 2 mg53.12

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

1 Patient is opioid dependent; and

continued...

28

Subsidy (Manufacturer's Price)	,		Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient received but failed detoxification with buprenorphine with naloxone; and
 - 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
 - 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
 - 4 Applicant works in an opioid treatment service approved by the Ministry of Health.


Tab modified-release 150 mg11.00	✓ Zyban
DISULFIRAM Tab 200 mg236.40 100 Antabuse to be Principal Supply on 1 November 2021	0 ✓ Antabuse
NALTREXONE HYDROCHLORIDE – Special Authority see SA1408 below – Retail pharr Tab 50 mg133.33 30	

⇒SA1408 Special Authority for Subsidy

BLIDDODIONI HADDOCHI ODIDE

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

Patch 7 mg - Up to 28 patch available on a PSO	b) Note: Direct Provision by a pharmacist permitted under the provisions	in Part I of Section A.	
Patch 14 mg — Up to 28 patch available on a PSO	Patch 7 mg - Up to 28 patch available on a PSO18.14	28	Habitrol
Patch 14 mg for direct distribution only – [Xpharm]	Patch 7 mg for direct distribution only – [Xpharm]3.94	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO		28	Habitrol
Patch 21 mg for direct distribution only – [Xpharm]	Patch 14 mg for direct distribution only – [Xpharm]4.52	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO		28	Habitrol
Lozenge 1 mg for direct distribution only – [Xpharm]	Patch 21 mg for direct distribution only - [Xpharm]5.18	7	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	Lozenge 1 mg - Up to 216 loz available on a PSO19.18	216	Habitrol
Lozenge 2 mg for direct distribution only – [Xpharm]	Lozenge 1 mg for direct distribution only - [Xpharm]3.20	36	Habitrol
Gum 2 mg (Fruit) — Up to 384 piece available on a PSO	Lozenge 2 mg - Up to 216 loz available on a PSO21.02	216	Habitrol
Gum 2 mg (Fruit) for direct distribution only – [Xpharm]	Lozenge 2 mg for direct distribution only - [Xpharm]3.24	36	Habitrol
Gum 2 mg (Mint) — Up to 384 piece available on a PSO	Gum 2 mg (Fruit) - Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only – [Xpharm]8.64 96	Gum 2 mg (Fruit) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 4 mg (Fruit) – Up to 384 piece available on a PSO	Gum 2 mg (Mint) - Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 4 mg (Fruit) for direct distribution only – [Xpharm]	Gum 2 mg (Mint) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 4 mg (Mint) − Up to 384 piece available on a PSO44.17 384 ✔ Habitrol	Gum 4 mg (Fruit) – Up to 384 piece available on a PSO44.17	384	Habitrol
	Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.01	96	Habitrol
Gum 4 mg (Mint) for direct distribution only − [Xpharm]10.01 96 ✓ Habitrol	Gum 4 mg (Mint) – Up to 384 piece available on a PSO44.17	384	Habitrol
, , ,	Gum 4 mg (Mint) for direct distribution only - [Xpharm]10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg x 11 and 1 mg x 42	16.67 25.64	53 OP	✓ Varenicline Pfizer✓ Champix
Varenicline Pfizer to be Principal Supply on 1 January 2022			
Tab 1 mg	17.62	56	✓ Varenicline Pfizer
·	27.10		✓ Champix

Varenicline Pfizer to be Principal Supply on 1 January 2022

(Champix Tab 0.5 mg x 11 and 1 mg x 42 to be delisted 1 January 2022) (Champix Tab 1 mg to be delisted 1 January 2022)

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and

NERVOUS SYSTEM

	Subsidy	0.1	Fully	Brand or
(1)	Manufacturer's Price) \$	Per	sidised •	Generic Manufacturer

continued...

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2046 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy (Manufacturer's Price)	Full Subsidise	
\$	Per 🗸	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+): and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2: and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

bosoli AN - FCT - netali pilatiliacy-specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1,387.00	1	✓ BiCNU
, ,	•		✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		-	
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Ebewe
ing i ing por ini, roo ini viai	29.66	•	✓ DBL Cisplatin
Inj 1 mg for ECP		1 mg	✓ Baxter
CYCLOPHOSPHAMIDE		9	
	70.00	50	✓ Endoxan S29
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
		100	•
Cyclonex to be Principal Supply on 1 January 2022	158.00	100	✓ Procytox S29
Inj 1 g vial – PCT – Retail pharmacy-Specialist	35.65	1	✓ Endoxan
IIIJ I g viai – I O I – Hetali phaimacy-opecialist	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist		1	✓ Endoxan
Inj 1 mg for ECP – PCT only – Specialist		1 mg	✓ Baxter
(Endoxan S29 Tab 50 mg to be delisted 1 January 2022)		٠٠	
(Procytox \$29 Tab 50 mg to be delisted 1 January 2022)			
(1 100ylox see Tab 30 mg to be delisted 1 January 2022)			

	Subsidy (Manufacturer's Price)		Fully Subsidised	I Generic
	\$	Per		Manufacturer
FOSFAMIDE - PCT only - Specialist				
Inj 1 g	96.00	1	1	Holoxan
Inj 2 g		1	1	Holoxan
Inj 1 mg for ECP	0.10	1 mg	/	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	1	CeeNU
Cap 40 mg	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1	1	Alkeran
			1	Alkeran S29 S29
	420.00		1	Tillomed S29
Tillomed S29 Inj 50 mg to be delisted 1 December 2021)				
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis
.,		•		100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1		Oxaliplatin Accord
Inj 1 mg for ECP		1 mg	_	Baxter
HIOTEPA - PCT only - Specialist		·		
Inj 15 mg vial	CBS	1	1	Bedford S29
ing to mg via				Max Health S29
				THIO-TEPA \$29
				Tepadina S29
lai 100 ma vial	CDC	4		•
Inj 100 mg vial	B3	1		Max Health S29
			•	Tepadina S29

Antimetabolites

AZACITIDINE – PCT only – Specialist – Special Authority see SA1467 below		
Inj 100 mg vial75.06	1	 Azacitidine Dr Reddy's
605.00		✓ Vidaza
Azacitidine Dr Reddy's to be Principal Supply on 1 December 2021		
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

CALCIUM FOLINATE		
Tab 15 mg - PCT - Retail pharmacy-Specialist114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist7.28	1	✓ Calcium Folinate
		<u>Sandoz</u>
		 Calcium Folinate
		Sandoz S29 S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49	1	 Calcium Folinate
		Sandoz
Inj 100 mg - PCT only - Specialist7.33	1	 Calcium Folinate
		Ebewe
Inj 300 mg – PCT only – Specialist22.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14	1	✓ Calcium Folinate
ing to mg per mi, so mi viai i or only opecialist25.14		Sandoz
		✓ Calcium Folinate
		Sandoz S29 S29
Inj 1 g - PCT only - Specialist67.51	1	✓ Calcium Folinate
ing r g r or only opposition		Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00	1	✓ Calcium Folinate
,g por,	·	Sandoz
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
CAPECITABINE – Retail pharmacy-Specialist	· ·	
Tab 150 mg	60	✓ Capercit
Tab 500 mg	120	✓ Capercit
CLADRIBINE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml749.96	1	✓ Leustatin
Inj 10 mg for ECP	10 mg OP	✓ Baxter
CYTARABINE	ŭ	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist400.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail	-	
pharmacy-Specialist	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist	10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist80.00	100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE		
Tab 10 mg - PCT - Retail pharmacy-Specialist412.00	20	✓ Fludara Oral
Inj 50 mg vial - PCT only - Specialist576.45	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist115.29	50 mg OP	✓ Baxter

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		Subsidised	
	\$	Per		Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	✓	Fluorouracil Accord
	12.00		✓	Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
	30.00		✓	Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg		Baxter
(Fluorouracil Ebewe Inj 50 mg per ml, 20 ml vial to be delisted 1 l	February 2022)			
(Fluorouracil Ebewe Inj 50 mg per ml, 100 ml vial to be delisted 1	February 2022)			
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g		1	/	Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg		Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist		J		
Inj 20 mg per ml, 5 ml vial	52 57	1	1	Accord
11) 20 119 por 111, 0 111 1141	71.44	•		Irinotecan
				Accord S29
			1	Irinotecan Actavis
			•	100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP		1 mg		Baxter
, ,		i ilig	•	Daxiei
(Irinotecan Accord S29 Inj 20 mg per ml, 5 ml vial to be delisted	1 Watch 2022)			
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist		25	/	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist			_	
Special Authority see SA1725 below	428.00	100 ml C)P 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

		Subsidy		Fully	Brand or
		(Manufacturer's Price		Subsidised	Generic
_		\$	Per		Manufacturer
ME	THOTREXATE				
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	9.98	90	1	Trexate
-	Trexate to be Principal Supply on 1 January 2022				
*	Tab 10 mg - PCT - Retail pharmacy-Specialist	33.71	90	1	Trexate
•	Trexate to be Principal Supply on 1 January 2022				
*	Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist	47 50	5	1	Methotrexate DBL
*	Inj 7.5 mg prefilled syringe		1		Methotrexate
*	ing 7.5 mg promied symige	17.01	'	•	Sandoz
*	Ini 10 ma profilled arrings	14.00	1	./	Methotrexate
*	Inj 10 mg prefilled syringe	14.00	ı	V	Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	•	Methotrexate
					Sandoz
*	Inj 20 mg prefilled syringe	14.88	1	✓	Methotrexate
					Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	✓	Methotrexate
	, , , ,				Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	/	Methotrexate
-	, g p , g		-		Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Speciali	et 30.00	5	1	Methotrexate DBL
*	mij 25 mg per mi, 2 mi viai - 1 O1 - Hetali phamacy opecial	31	3	•	Onco-Vial
	Ini OF man and OO militial DOT Datail abannasi Casain	.II1 4E 00	4		DBL Methotrexate
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specia	IIIST45.00	1	V	
				_	Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist	t25.00	1	•	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial - PCT - Retail				
	pharmacy-Specialist		1	✓	Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓	Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist.	4.73	5 mg Ol	P 🗸	Baxter
PF	METREXED - PCT only - Specialist - Special Authority see	SA1679 below			
-	Inj 100 mg vial		1	1	Juno Pemetrexed
	Inj 500 mg vial		1		Juno Pemetrexed
	Inj 1 mg for ECP		1 mg		Baxter
_			1 1119	•	-unioi

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

Subsidy (Manufacturer's Price)	s	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Permetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE	- PCT - Ref	tail pharmac	y-Specialist
T 1 40			

Tab 40 mg126.31 25 ✓ Lanvis

Other Cytotoxic Agents

AMSACRINE – PCT only – Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmac	/-Specialist		
Cap 0.5 mg	1,175.87	100	Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP		10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist			
Inj 15,000 iu, vial	161.01	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	12.45	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority s	ee SA1889 below		
Inj 3.5 mg vial		1	✓ Bortezomib Dr-Reddy's
			✓ Bortezomib Juno \$29
Inj 1 mg for ECP	31.20	1 mg	✓ Baxter

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

	Subsidy		Fully	Brand or
	(Manufacturer's Pri	ce) Subs	idised	
	\$	Per	_	Manufacturer
DACARBAZINE – PCT only – Specialist				
Inj 200 mg vial	62.70	1	1	DBL Dacarbazine
, 3	580.60	10	1	Dacarbazine
				APP S29
Inj 200 mg for ECP	62.70	200 mg OP	1	Baxter
	02.70	200 mg Oi	•	Daxiei
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist				<u> </u>
Inj 0.5 mg vial		1		Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	•	Baxter
DAUNORUBICIN - PCT only - Specialist				
Inj 2 mg per ml, 10 ml	149.50	1	1	Pfizer
Inj 20 mg for ECP	149.50	20 mg OP	1	Baxter
DOCETAXEL - PCT only - Specialist				
Inj 20 mg	48 75	1	1	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1		DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1		Docetaxel
inj 20 mg per mi, 4 mi vidi	20.00	•	•	Accord \$29
la: 00	405.00		,	
Inj 80 mg		1		Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	•	Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	10.00	1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	1	Doxorubicin Ebewe
	17.00		1	Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	1	Arrow-Doxorubicin
	69.99		1	Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	1	Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	1	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1		Epirubicin Ebewe
Inj 1 mg for ECP		1 mg		Baxter
ETOPOSIDE		9		
	240.72	00	./	Venesid
Cap 50 mg - PCT - Retail pharmacy-Specialist		20		Vepesid Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		Vepesid Day Madical
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia		1		Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	v	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist				
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	1	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha	rmacy-Specialist			
Cap 500 mg		100	1	Devatis
		•		
IDARUBICIN HYDROCHLORIDE	00.00	4		Zavadas
Inj 5 mg vial – PCT only – Specialist		1		Zavedos
Inj 10 mg vial – PCT only – Specialist		1		Zavedos
Inj 1 mg for ECP - PCT only - Specialist	21.84	1 mg	•	Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LENALIDOMIDE - Retail pharmacy-Specialist - Special Authorit	y see SA2047 below			
Wastage claimable				
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg	4,655.25	21	✓	Revlimid
•	6,207.00	28	✓	Revlimid
Cap 15 mg	5,429.39	21	✓	Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg	7,627.00	21	1	Revlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 0.4.1.5
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MFSNA

Tab 400 mg - PCT - Retail pharmacy-Specialist3	14.00 5	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist4	48.50 5	50	Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist1	77.45 1	5	Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	07.40 1	5	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	.2.96 100	mg 🗸	Baxter

	Subsidy Manufacturer's Price	١	Fully	
(\$	Per	Jubsiuiseu ✓	
/ITOMYCIN C - PCT only - Specialist				
Inj 20 mg vial	3,275.00	1		Omegapharm \$29
Inj 1 mg for ECP	470.75	1 ma	_	Teva Baxter
MITOZANTRONE - PCT only - Specialist	470.73	i ilig	•	Daxiei
Inj 2 mg per ml, 10 ml vial	97.50	1	/	Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	•	Baxter
DLAPARIB - Retail pharmacy-Specialist - Special Authority see S	A1883 below			
Tab 100 mg	3,701.00	56	•	Lynparza
Tab 150 mg	3,701.00	56	✓	Lynparza

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	Paclitaxel Ebewe
•	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
•	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
•	275.00		✓ Anzatax
			Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see \$	SA1979 on the next page		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

	Subsidy	F	ully	Brand or
(N	Manufacturer's Price)	Subsid	ised	Generic
	\$	Per	•	Manufacturer

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

DENITOSTATINI	[DEOXYCOFORMYCIN]	DCT only	, Specialist
PENTOSTATIN		- PC I OIII	/ – Specialist

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail	pharmacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 be	low – Retail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ Temaccord
, ,	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ <u>Temaccord</u>
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ <u>Temaccord</u>
	688.00		✓ Amneal S29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m2 per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special A	authority see SA1124 below		
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

100 ✓ Vesanoid

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 below			
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OF	•	Venclexta
Tab 10 mg	95.78	14 OF	•	Venclexta
Tab 50 mg	239.44	7 OP	✓	Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓	Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SUI PHATE

inj i mg per mi, 10 mi viai – PCT – Retail pharmacy-Specialist270.37	5	♥ DBL vindiastine 529
Inj 1 mg for ECP – PCT only – Specialist	1 mg	✓ Hospira✓ Baxter
(DBL Vinblastine 229 Inj 1 mg per ml, 10 ml vial to be delisted 1 November 2021)		
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	✓ DBL Vincristine Sulfate
Ini 1 mg for ECP - PCT only - Specialist	1 ma	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
VINORELBINE - PCT only - Specialist	· · · · · · · · · · · · · · · · · · ·			
Inj 10 mg per ml, 1 ml vial	12.00	1	✓	Navelbine
	42.00		1	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	✓	Navelbine
,	210.00		1	Vinorelbine Ebewe
	328.65		1	Sagent S29
Inj 1 mg for ECP		1 mg		Baxter
Inj 50 mg for ECP) mg OP	✓	Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

wasiage ciaimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel
3	**		

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

Mostogo eleimoble

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	Manufacturer

continued...

- 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
- 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
- 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority s	see SA2000 below		
Tab 100 mg	764.00	30	✓ Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

⇒SA2000 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive: or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malionant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg – [Xpharm] – Special Authority see SA1460		
	below2,400.00	60	✓ Glivec
÷	Cap 100 mg58.23	60	✓ Imatinib-Rex
÷	Cap 400 mg84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

* *

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy Note – no new patients to be initiated on lapatinib ditosylate.

70 ✓ Tykerb

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable ✓ Tasigna 120 120 ✓ Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- - 2.1 Patient has documented CML treatment failure* with imatinib; or

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines: and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable	•		
Tab 75 mg	4,000.00	21	✓ Ibrance
Tab 100 mg	4,000.00	21	Ibrance
Tab 125 mg	4,000.00	21	Ibrance
Cap 75 mg	4,000.00	21	Ibrance
Cap 100 mg	4,000.00	21	Ibrance
Cap 125 mg	4,000.00	21	Ibrance

(Ibrance Cap 75 mg to be delisted 1 March 2022)

(Ibrance Cap 100 mg to be delisted 1 March 2022)

(Ibrance Cap 125 mg to be delisted 1 March 2022)

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	1	Manufacturer
PAZOPANIB - Special Authority see SA1190 below - Retail ph	armacy			
Tab 200 mg	1,334.70	30	✓	Votrient
Tab 400 mg	2,669.40	30	1	Votrient
- CA1100 Chariel Authority for Cubaidy				

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive: or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5,000.00	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or

2.2 Both:

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	Manufacturer	

continued...

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2002 below - Retail pharmacy

Cap 12.5 mg2,315.38	28	✓ Sutent
Cap 25 mg	28	✓ Sutent
Cap 50 mg	28	✓ Sutent

⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and 4 The disease is of predominant clear cell histology; and
- The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 85

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and

Subsidy		Fully	Brand or
(Manufacturer's		sidised	Generic
\$	Per		Manufacturer

continued...

- 4.2.2 Patient has ECOG performance score of 0-2; and
- 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg		10 28	✓ Calutide-50 S29 ✓ Binarex
(Calutide-50 \$29	Tab 50 mg to be delisted 1 January 2022)	20	▼ <u>Billarex</u>
FLUTAMIDE			
Tab 250 mg.	107.55	90	✓ Prostacur S29
-	119.50	100	✓ Flutamin
FULVESTRANT -	- Retail pharmacy-Specialist - Special Authority see SA1895 below	V	
	ml, 5 ml prefilled syringe1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised <	
OCTREOTIDE				
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	1	Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	1	Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial		5	1	Octreotide
				MaxRx S29
	56.87		1	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓	DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule	72.50	5	1	Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓	DBL Octreotide
	222.00		1	Octreotide
				(Sun) S29
OCTREOTIDE LONG-ACTING - Special Authority see SA20	072 below – Retail pharm	acy		
Inj depot 10 mg prefilled syringe		ĺ	1	Octreotide Depot
			_	Teva
	1,772.50			Sandostatin LAR
Inj depot 20 mg prefilled syringe	647.03	1	•	Octreotide Depot Teva
	2,358.75		1	Sandostatin LAR
Inj depot 30 mg prefilled syringe	718.55	1	•	Octreotide Depot Teva
	2,951.25		/	Sandostatin LAR
Sandostatin LAR Inj depot 10 mg prefilled syringe to be delis	sted 1 March 2022)			

(Sandostatin LAR Inj depot 10 mg prefilled syringe to be delisted 1 March 2022) (Sandostatin LAR Inj depot 20 mg prefilled syringe to be delisted 1 March 2022) (Sandostatin LAR Inj depot 30 mg prefilled syringe to be delisted 1 March 2022)

⇒SA2072 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed: and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma: and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

	Tab 20 mg	60	✓ Tamoxifen Sandoz
A	romatase Inhibitors		

ANAS	TROZOLE	

* Tab 1 mg4.55	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE	30	✓ Letrole

Letrole to be Principal Supply on 1 January 2022

60

✓ Tamoxifen Sandoz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Immunosuppressants

Cytotoxic Immunosuppressants

ΑZ	ATH	H	OΡ	RI	ΝE
	_				

*	1 ab 25 mg	/.35	60	Azamun
*	Tab 50 mg	7.60	100	✓ Azamun
	Inj 50 mg vial		1	✓ Imuran
MY	COPHENOLATE MOFETIL			
	T 1 500	05.00		

TCOPHENOLATE MOPETIL		
Tab 500 mg35.90	50	Cellcept
Cap 250 mg35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml — Subsidy by endorsement	165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT - Special Authority see SA2048 below - Reta	iil pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector		4	✓ Enbrel
Ini 50 ma prefilled syringe		4	✓ Enbrel

⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application. Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

2 Fither:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plague psoriasis; and
 - 1.2 Either.
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy	Fu	lly Brand or
(Manufacturer's Price)	Subsidise	ed Generic
\$	Per	 Manufacturer

continued...

- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

ANTITHYMOCYTE GLOBILLIN (FOLLINE) - PCT only - Specialist

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITITIMOOTTE GEODOLIN (EQUINE) - 1 OT OHIJ - Specialist		
Inj 50 mg per ml, 5 ml2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only - Specialist		
Subsidised only for bladder cancer.		
Inj 2-8 × 100 million CFU149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG 529 Inj 40 mg per ml, vial to be delisted 1 April 2022)		

Monoclonal Antibodies

		below – Retail pharmacy	ADALIMUMAB – Special Authority see SA2049 be
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
Humira	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2049 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

continued...

55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
 - 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss: and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- of the following 1 Either:
 - 1.1 Applicant is a gastroenterologist; or

Subsidy	Fully	Brand or
(Manufacturer's		Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab: or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage III or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:

Subsidy	Fully	Brand or	_
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a

dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
S Per ✔ Manufacturer

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy: or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	Baxter

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAR	- PCT only -	- Special Authority	see SA2050 below
	I O I OIII	Opedial Additionity	JOE UNEUUU DUIUW

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA2050 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

Subsi	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:

S	Subsidy	Fully	Brand or
(Manufa	cturer's Price)	Subsidised	Generic
	\$ Per	r √	Manufacturer

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sul	bsidised	Generic
\$	Per	1	Manufacturer

continued...

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Fither:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for severe chronic plaque psoriasis; and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis: or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
 - 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy	-	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

(1)	Subsidy Manufacturer's Price)	Subs	Fully idised	Brand or Generic
	\$	Per	✓	Manufacturer

continued...

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema): or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subs	sidy Full	/ Brand or
(Manufactur	rer's Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - Re	etail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Ini 100 mg vial	1.638.00	1	✓ Nucala

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg j	orefilled syringe	450.00	1	Xolair
Ini 150 mg	vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	l Generic	
\$	Per 🗸	Manufacturer	

continued...

or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months. unless contraindicated or not tolerated: and

- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
 - 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

	nj 100 mg per 10 mi viai)	2	•	viaptnera
- 1	nj 500 mg per 50 ml vial	2,688.3)	1 •	/ I	Mabthera
- 1	nj 1 mg for ECP	5.6	1 1	mg •	✓ E	Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2061 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Ini 1 ma for ECP	1.38	1 ma	✓ Baxter (Riximvo)

⇒SA2061 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive: or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

continued...

375 mg/m2 administered weekly for four weeks; and

- - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks: and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective: or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective;
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- - 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
 - 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy: and
 - 1.2 To be used for a maximum of 6 treatment cycles: or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy:
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms: and

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and

Subsidy (Manufacturer's Price)	Fully Subsidised		
(ivialitulacture) 5 Filoe)	Per 🗸	Manufacturer	
Φ	rei •	Manuacturei	

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of $2 \times 1,000$ mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function: and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria: Fither:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*: and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB – Special Authority see SA2044 below – Retail pharmacy

⇒SA2044 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or

Subsid (Manufacturer	•	Fully	Brand or Generic	
(Manuacture)	Per	J3iui3eu ✓	Manufacturer	

continued...

- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or etanercept for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or etanercept to meet the renewal criteria for adalimumab or etanercept for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	

continued...

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
 - 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

✓ Sylvant 1 1 ✓ Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA2078 below

Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
Inj 20 mg per ml, 10 ml vial		1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP	,	1 mg	✓ Baxter

⇒SA2078 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
 - 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 rheumatoid arthritis; or
- 2.2 systemic juvenile idiopathic arthritis; or
- 2.3 adult-onset Still's disease; or
- 2.4 polyarticular juvenile idiopathic arthritis; or
- 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Subsidy (Manufacturer's Price)	Su	Fully ibsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and

1.2 Fither:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient has significantly increased laboratory markers of systemic inflammation (eg CRP, PCT or ferritin); and
- 4 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 5 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose.

Note: Indications marked with * are unapproved indications.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Au	thority see SA1632 below		
Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	·	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
- 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

 Renewal (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

specialist. Approvals valid for 12 months for applications meeting the following criteria:

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	 Subsidy		Fully	Brand or
\$ Per ✓ Manutacturer	(Manufacturer's Price)		dised	
ų 13. maiatata	\$	Per		Manutacturer

continued...

- 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
- 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial	2,320.00	1	✓ Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	23.20	1 mg	✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB - PCT only - Specialist - Special Authori	ity see SA2006 on the next	page	
Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Fither:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period: or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.

Subsidy (Manufacturer's Price)	Full Subsidise	
\$	Per 🗸	Manufacturer

continued...

- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special A	uthority see SA2007 below	
Inj 25 mg per ml, 4 ml vial	4,680.00	✓ Keytruda
Inj 1 mg for ECP	49.14 1 m	ng 🗸 Baxter

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Either
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and

Subsidy (Manufacturer's Price)	Suk	Fully	Brand or Generic
 \$	Per	√	Manufacturer

continued...

2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN		
Cap 25 mg44	.63 5	0 ✓ Neoral
Cap 50 mg88	.91 5	0 ✓ Neoral
Cap 100 mg177	.81 5	0 ✓ Neoral
Oral liq 100 mg per ml198	.13 50 m	nl OP 🗸 Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pharmacy		
Wastage claimable		
Tab 10 mg6,512	.29 3	30 ✓ Afinitor
Tab 5 mg4,555	.76 3	30 ✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS – Special Authorit	y see SA2005 on the next	page – Retail pharmacy
------------------------------	--------------------------	------------------------

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

 $\textbf{Renewal--(renal angiomyolipoma(s) associated with tuberous sclerosis complex*)} \ \ \text{from any relevant practitioner}.$

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound;
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg		100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
S Per ✔ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1	367 above -	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent .	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA	A1367 above	– Retail pharn	nacy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			4
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	005.00	4 OD	✓ Allean
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze	205.00	1 OP	✓ Venomil S29
dried venom, with diluent	303.00	1 01	Verionili 529

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		idised	Generic
	\$	Per	✓	Manufacturer
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1.12	100	✓ Z	ista
* Oral liq 1 mg per ml	2.84	200 ml	✓ H	listaclear
Histaclear to be Principal Supply on 1 January 2022				
CHLORPHENIRAMINE MALEATE				
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ H	listafen
		300 1111	• 11	iistaicii
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg		40	_	
	(8.40)		Р	olaramine
	1.01	20	_	
	(5.99)		Р	olaramine
* Oral liq 2 mg per 5 ml		100 ml	_	
	(10.29)		Р	olaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
ů	(8.23)		Т	elfast
* Tab 120 mg	4.74 [′]	10		
, and a g	(8.23)		Т	elfast
	14.22	30		
	(26.44)		Т	elfast
LORATADINE	(- /			
* Tab 10 mg	1.60	100	./ 1	orafix
· · · · · · · · · · · · · · · · · · ·		100 ml		laylor syrup
	1.43	100 1111	V <u>1</u>	iayioi syrup
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg		50		Illersoothe
* Tab 25 mg		50		Illersoothe
* Oral liq 1 mg per 1 ml		100 ml		Illersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	a PSO 17.87	5	✓ H	lospira
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose		200 dose OP	√ 0	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP		Seclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	√ 0	
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP		Seclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ B	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ P	ulmicort
.				Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	√ P	ulmicort
			•	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	√ D	ulmicort
Toward for initial ation, 400 filey per 4036		_00 003E OF	· F	Tambook alam

Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's P \$	rice) Subs Per	idised Generic Manufacturer
	Ψ	rei	Wanuacturer
FLUTICASONE			4
Aerosol inhaler, 50 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose	7.50	60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ <u>Flixotide</u>
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide ✓ Flixotide Accuhaler
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	Flixotide Accunaier
Inhaled Long-acting Beta-adrenoceptor Agonists			
initialed Long-acting Deta-adrenoceptor Agonists	,		
EFORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose device	e20.64	60 dose	
	(35.80)		Foradil
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose).	10.32	60 dose OP	
(equivalent to clothioteror familiate of mog motored dood).	(16.90)	00 0000 01	Oxis Turbuhaler
INDACATEROL	(10.00)		CAIC Turburiaior
INDACATEROL Powder for inhelation 150 mag	61.00	30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 150 mcgPowder for inhalation 300 mcg		30 dose OP	✓ Onbrez Breezhaler ✓ Onbrez Breezhaler
ŭ	61.00	30 dose OF	• Officez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose		120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-A	drenocepto	or Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide with	h		
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumara:		120 0000 01	5 Buortesp opirolliax
per dose (equivalent to 400 mcg budesonide with 12 mcg	ic		
eformoterol fumarate metered dose) – No more than 2			
dose per day	82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mc		120 dose OP	✓ Symbicort
To the state of th	·9 ····· ·	0 0000 0.	Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40	120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 mc		120 dose OP	✓ Symbicort
1 0 1 0 0 1 0 1 1 1 1 1 1 1 1 1 1 1 1 1	·9 ····· · · · · · ·	0 0000 0.	Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg – No more than 2 dose per day	44.08	60 dose OP	✓ Symbicort
			Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.00	30 dose OP	✓ Breo Ellipta
Toward for initial autor 100 micy with vitalite of 25 micy	44.00	JU UUSE UP	- DIEO EIIIPIA

	Subsidy (Manufacturer's \$	Price) Subs	idised	Brand or Generic Manufacturer
FLUTICASONE WITH SALMETEROL				
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP	✓ Ser	
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Ser	<u>etide</u>
Powder for inhalation 100 mcg with salmeterol 50 mcg – No		00 1 00		
more than 2 dose per day		60 dose OP	✓ Ser	etide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg - No more than 2 dose per day		60 dose OP	✓ Ser	etide Accuhaler
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral liq 400 mcg per ml	40.00	150 ml	✓ Ver	itolin
Infusion 1 mg per ml, 5 ml	118.38	10	✓ Ver	
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	53.00	5	✓ Ver	itolin
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP	✓ Res	pigen
	4		✓ Sal	
Nobelian ale 4 managed 0.5 ml amagela. He to 00 ml	(6.00)		Ver	tolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ Ast	halin
Asthalin to be Principal Supply on 1 January 2022	0.90	20	▼ ASI	IIdiiii
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb				
available on a PSO		20	✓ Ast	halin
Asthalin to be Principal Supply on 1 January 2022				
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to				
250 mcg metered dose), breath activated	22.20	120 dose OP	✓ Brid	anyl Turbuhaler
Anticholinergic Agents				
IPRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free	16.20	200 dose OP	✓ Atr	ovent
a) Up to 400 dose available on a PSO				
b) No patient co-payment payable				
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne	eb			
available on a PSO	11.73	20	✓ Uni	vent
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	gents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p	per			
dose CFC-free		200 dose OP	✓ Due	olin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per				
vial, 2.5 ml ampoule — Up to 20 neb available on a PSO Duolin to be Principal Supply on 1 January 2022	11.04	20	✓ Due	olin

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

✓ fully subsidised S29 Unapproved m Principal Supply Sole Subsidised Su

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note. I inclident is not subsidiscu in combination wit	ii subsidisca iiii ilcadiiib.		
Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	·	90	✓ Esbriet
Cap 267 mg - Wastage claimable		270	✓ Esbriet

(Esbriet Cap 267 mg to be delisted 1 January 2022)

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

Leukotriene Receptor Antagonists

MOM	NTELUKAST		
*	Tab 4 mg4.25	28	✓ Montelukast Mylan
	Tab 5 mg4.25	28	✓ Montelukast Mylan
*	Tab 10 mg	28	✓ Montelukast Mylan

Mast Cell Stabilisers

SODIUM CROMOGLICATE - Subsidy by endorsement

Methylxanthines

AMINOPHYLLINE

*	Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	180.00	5	✓ DBL Aminophylline
TH	EOPHYLLINE			
*	Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
*	Oral lig 80 mg per 15 ml	16.60	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA – Special Authority see SA1978 below –	· Retail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 on the next page

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	·	56	✓ Kalydeco

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Fither:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N and S549R) in the CFTR gene on at least 1 allele; and
- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

RUDESONIDE

Not funded for use as a nasal drop.

Nasal Preparations

Allergy Prophylactics

Metered aqueous nasal spray, 50 mcg per dose		✓ <u>SteroClear</u> ✓ <u>SteroClear</u>
FLUTICASONE PROPIONATE		
Metered aqueous nasal spray, 50 mcg per dose1.98	120 dose OP	✓ Flixonase Hayfever

Flixonase Hayfever & Allergy to be Principal Supply on 1 December 2021

IPRATROPIUM BROMIDE

Respiratory Devices

MASK FOR SPACER DEVICE

- a) Up to 50 dev available on a PSO
- b) Only on a PSO
- c) Only for children aged six years and under

PEAK FLOW METER

- a) Up to 25 dev available on a PSO
- b) Only on a PSO

✓ Mini-Wright AFS Low Range

& Alleray

Normal range......9.54

Mini-Wright Standard

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
220 ml (single patient)	2.95	1		e-chamber Turbo
510 ml (single patient)	5.12	1		e-chamber La Grande
800 ml	6.50	1	1	Volumatic

CAFFEINE CITRATE			
Oral lig 20 mg per ml (10 mg base per ml)	15.10	25 ml OP	Biomed

	Subsidy (Manufacturer's I \$	Price) Subs Per	Fully Brand or idised Generic Manufacturer
Ear Preparations			
ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BE For Vosol ear drops with hydrocortisone powder refer Stand Ear drops 2% with 1, 2-Propanediol diacetate 3% and			
benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol
FLUMETASONE PIVALATE Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTA	TIN	✓ Locorten-Vioform
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate		1111	
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50	8 ml OP	
	(9.27)	· · · · · · ·	Sofradex
FRAMYCETIN SULPHATE Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expli	citly stated other	wise.	
Anti-Infective Preparations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%Funded for use in the ear*. Indications marked with * au		10 ml OP	✓ Chlorafast
CIPROFLOXACIN	о апарриотоа п		
Eye drops 0.3% – Subsidy by endorsement	tis or severe bac ve otitis media (C unapproved indic	CSOM)*; and the	

* Eye drops 0.1%......2.97

GENTAMICIN SULPHATE

PROPAMIDINE ISETHIONATE

SODIUM FUSIDATE [FUSIDIC ACID]

✓ Genoptic

Brolene

✓ Fucithalmic

5 ml OP

10 ml OP

5 g OP

(14.55)

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F	Price) Sub Per	Fully sidised	Brand or Generic Manufacturer	
TOBRAMYCIN Eye oint 0.3% Eye drops 0.3%		3.5 g OP 5 ml OP	-	obrex obrex	
Corticosteroids and Other Anti-Inflammatory Pr	eparations				
BEXAMETHASONE Eye oint 0.1% Eye drops 0.1% Ocular implant 700 mcg – Special Authority see SA1680 belance.	4.50	3.5 g OP 5 ml OP	••	Maxidex Maxidex	

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Either:
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	LOFENAC SODIUM			
	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
	Voltaren Onbtha to be Principal Supply on 1 November 2021			

Ozurdex

	Subsidy (Manufacturer's Pr	ice) Sub	Fully	Brand or Generic	
	\$	Per	1	Manufacturer	
FLUOROMETHOLONE					
* Eye drops 0.1%	3.09	5 ml OP	√ F	ML	
	5.20		✓ F	lucon	
KETOROLAC TROMETAMOL - Special Authority see SA1981	below – Retail pha	armacy			
Eye drops 0.5%	9.50 ·	5 ml OP	✓ A	Acular	
(Acular Eye drops 0.5% to be delisted 1 November 2021)					

⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

Evo drang 0 E ma nor ml	0.71	4 ml OP	
Eye drops 0.5 mg per ml	(10.34)	4 IIII OP	Livostin
LODOXAMIDE Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC Eye drops 0.3%(Ilevro Eye drops 0.3% to be delisted 1 November 2021)	13.80	3 ml OP	✓ Ilevro
PREDNISOLONE ACETATE Eye drops 1%		10 ml OP	✓ Prednisolone-AFT
	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below	– Retail pharr	nacy

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Eye drops 0.5%, single dose (preservative free)......38.50

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

20 dose

5 ml OP

✓ Minims

✓ Rexacrom

Prednisolone

SODIUM CROMOGLICATE

, ,				
Glaucoma Preparations - Beta Blockers				
BETAXOLOL				
* Eye drops 0.25%	11.80	5 ml OP	✓ Betoptic S	
* Eye drops 0.5%		5 ml OP	✓ Betoptic	
TIMOLOL			-	
* Eye drops 0.25%	1.81	5 ml OP	✓ Arrow-Timolol	
* Eye drops 0.5%		5 ml OP	✓ Arrow-Timolol	
* Eye drops 0.5%, gel forming		2.5 ml OP	✓ Timoptol XE	

	Subsidy (Manufacturer's F \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
Glaucoma Preparations - Carbonic Anhydrase	Inhibitors			
ACETAZOLAMIDE				
米 Tab 250 mg	17.03	100	✓ D	iamox
BRINZOLAMIDE				
* Eye drops 1%	7.30	5 ml OP	✓ A	zopt
OORZOLAMIDE HYDROCHLORIDE				
* Eye drops 2%	9.77	5 ml OP		
	(17.44)		Т	rusopt
OORZOLAMIDE WITH TIMOLOL				
* Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ D	ortimopt
Glaucoma Preparations - Prostaglandin Analog	gues			
BIMATOPROST				
* Eye drops 0.03%	3 30	3 ml OP	√ R	imatoprost
- 2,0 0.000 0.00 /0		0 1111 01	ں ۔	Multichem
ATANOPROST				
* Eye drops 0.005%	1 82	2.5 ml OP	✓ T	eva
	1.02	2.5 1111 01	• .	ova
FRAVOPROST * Eye drops 0.004%	7 20	5 ml OP	√ T	ravopt
κ Eye αιορs 0.004 /o	9.75	2.5 ml OP		ravopi ravatan
	10.50	5 ml OP		vlan S29
Travatan to be Principal Supply on 1 December 2021	10.50	3 IIII OF	♥ IV	ylair 023
Travopt Eye drops 0.004% to be delisted 1 December 2021)				
Mylan S29 Eve drops 0.004% to be delisted 1 December 2021	1)			
liviyian be delisted it December 202	')			
Glaucoma Preparations - Other				
BRIMONIDINE TARTRATE				
* Eye drops 0.2%		5 ml OP	✓ A	rrow-Brimonidine
Arrow-Brimonidine to be Principal Supply on 1 January	2022			
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ C	ombigan
ATANOPROST WITH TIMOLOL				
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ A	rrow - Lattim
PILOCARPINE HYDROCHLORIDE				
* Eye drops 1%	4.26	15 ml OP	✓ Is	opto Carpine
* Eye drops 2%	5.35	15 ml OP		opto Carpine
* Eye drops 4%		15 ml OP	✓ Is	opto Carpine
Subsidised for oral use pursuant to the Standard Formu				
★ Eye drops 2% single dose – Special Authority see SA0895			_	
below - Retail pharmacy		20 dose		inims Pilocarpine

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics a	nd Cyc	loplegics
--------------	--------	-----------

ATROPINE SULPHATE * Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) – Only on a prescription	20 dose	✓ Minims Cyclopentolate
(Minims Cyclopentolate Eye drops 1%, single dose (preservative free) to be det	listed 1 April 2022	, ,
* Eye drops 0.5%	15 ml OP 15 ml OP	MydriacylMydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 245

HYPROMELLOSE

	Eye drops 0.5%	19.50	15 ml OP	✓ Methopt
111/	DDOMELL OCE WITH DEVEDAN			

HYPROMELLOSE WITH DEXTRAN 15 ml OP ✓ Poly-Tears

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye: and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eve drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail pharmacy ✓ Polv-Gel MACROGOL 400 AND PROPYLENE GLYCOL - Special Authority see SA1388 above - Retail pharmacy Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml4.30 ✓ Systane Unit Dose SODIUM HYALURONATE [HYALURONIC ACID] - Special Authority see SA1388 above - Retail pharmacy 10 ml OP ✓ Hylo-Fresh

a) Hylo-Fresh has a 6 month expiry after opening. The Pharmacy Procedures Manual restriction allowing one bottle per

- month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.
- b) Hylo-Fresh to be Principal Supply on 1 January 2022

SENSORY ORGANS

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully dised	Brand or Generic Manufacturer
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	4.15	15 ml OP	✓ N	laphcon Forte
OLOPATADINE Eye drops 0.1%	2.20	5 ml OP	√ <u>0</u>	Nopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.63	3.5 g OP	✓ P	oly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	3.80	5 g OP	✓ V	itA-POS

Subsidy Fully (Manufacturer's Price) Per

Subsidised

Brand or Generic Manufacturer

Agents Used in the Treatment of Poisonings

Antidotes

ACETYL CYSTEINE

Inj 200 mg per ml, 10 ml ampoule58.76 10 ✓ DBL Acetylcysteine ✓ Martindale Pharma \$29

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO
- ✓ DBL Naloxone 5 Hydrochloride

Removal and Elimination

CHARCOAL

250 ml OP ✓ Carbosorb-X

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels. liver or cardiac MRI T2*: or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis: or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 on the nex	xt page - Retail pharn	nacy	
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE * Inj 500 mg vial	84.53	10	✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	52 21	6	
* IIIJ 200 IIIg pei IIII, 3 IIII		U	
	(156.71)		Calcium Disodium
			Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml	qs	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml)	LIQUID (10
Suitable eye drop base	qs	Phenobarbitone Sodium Glycerol BP	400 mg 4 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate	60 mg	Water	to 40 ml
Glycerol Preservative	40 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg per 5 ml)		Water (Preservative should be used if quantity supplied is	to 500 ml for more
Codeine phosphate Glycerol	300 mg 40 ml	than 5 days.)	
Preservative Water	qs to 100 ml	SALIVA SUBSTITUTE FORMULA Methylcellulose	5 g
FOLINIC MOUTHWASH		Preservative Water	qs to 500 ml
Calcium folinate 15 mg tab Preservative	1 tab qs	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	for more
Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.)	to 500 ml for more	SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml	qs
METHADONE MIXTURE		Water (Only funded if prescribed for treatment of hyponatra	qs aemia)
Methadone powder Glycerol	qs qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
Water	to 100 ml	Vancomycin 500 mg injection Glycerol BP	10 vials 40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate	10 g	Water (Only funded if prescribed for treatment of Clostridiu	to 100 ml
Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqu	to 100 ml	following metronidazole failure)	m annone
OMEPRAZOLE SUSPENSION	ila Illixtule)	VOSOL EAR DROPS	
Omeprazole Sostelivsion Omeprazole capules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	WITH HYDROCORTISONE POWDER 1% Hydrocortisone powder Vosol Ear Drops	1% to 35 ml
	10 100 1111		
PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully Brand or
	(Manufacturer's Pi	rice) Sub	sidised Generic Manufacturer
Extemporaneously Compounded Preparations	and Galenica	ils	
CODEINE PHOSPHATE - Safety medicine; prescriber may de		g frequency	
Powder – Only in combination		25 g	December
Only in extemporaneously compounded codeine linctus	(90.09)		Douglas
COLLODION FLEXIBLE	·•		
Note: This product is no longer being manufactured by the determined.	supplier and will b	e delisted fror	m the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination			
Only in extemporaneously compounded oral mixtures.			
Soln		100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination	า		
Only in combination with Ora-Plus. Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination			<u> </u>
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prep	arations.		
METHADONE HYDROCHLORIDE a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing f			
d) Extemporaneously compounded methadone will only be	reimbursed at the	rate of the ch	neapest form available
(methadone powder, not methadone tablets). Powder	7 84	1 g	✓ AFT
METHYL HYDROXYBENZOATE		' 9	· All
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE		-	
Powder		100 g	✓ <u>MidWest</u>
Suspension – Only in combination		473 ml	✓ <u>Ora-Plus</u>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH Suspension		combination 473 ml	√ Ove Bland CE
•			✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Or Suspension	,	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM		1701111	<u>ora Bioria</u>
Powder – Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years			
PROPYLENE GLYCOL		_	
Only in extemporaneously compounded methyl hydroxyben Liq		n. 500 ml	✓ Midwest
SODIUM BICARBONATE		555 IIII	
Powder BP - Only in combination		500 g	✓ <u>Midwest</u>
Only in aytomographously compounded amongszala ar	id lanconrazola cu	enoncion	

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>M</u>	lidwest	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see SA1930 above – Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

0.1.1		- "	D -	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia: or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 249

Subsidy		Fully	Brand or	
(Manufacturer's Price))	Subsidised	Generic	
	Per	✓	Manufacturer	

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)12.30	200 ml OP	✓ Calogen
30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)12.30	200 ml OP	✓ Calogen
Oil	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	 Special Authority see SA1524 above – Hospital p 	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy (Manufacturer's Price) Subsidised Per

Brand or Generic Manufacturer

Fully

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Autho	•	500 ml OP	nacy [HP3] Glucerna Select Diason RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority s Liquid (strawberry) Liquid (vanilla)	1.50	spital pharmacy 200 ml OP 200 ml OP 237 ml OP	[HP3] ✓ Diasip ✓ Diasip
(Containing Dishatial implied (conflict to a delicated of Fahrum	(2.10) (2.10)	200 ml OP	Sustagen Diabetic Nutren Diabetes

(Sustagen Diabetic Liquid (vanilla) to be delisted 1 February 2022)

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above - Hospital pharmacy [HP3] 400 g OP Monogen

251 ✓ fully subsidised

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy (HP3)

400 g OP ✓ Heparon Junior

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 vears where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3] Powder54.00 400 a OP ✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see S. Liquid		he previous pag 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SALiquid		previous page 500 ml OP	Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Ai pharmacy [HP3]	uthority see	SA1379 on the	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA13 Liquid (strawberry) Liquid (vanilla)	1.60	revious page – 200 ml OP 200 ml OP	Hospital pharmacy [HP3] Fortini Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML — Special Authority see SA1379 Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.07 1.07	evious page – He 200 ml OP 200 ml OP 200 ml OP 250 ml OP	ospital pharmacy [HP3] ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Autho pharmacy [HP3]	rity see SA	1379 on the pre	vious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.60 1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		oage – Hospital 400 g OP	pharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML − Special Authority see SA1101 above − Hospital pharmacy [HP3]

Liquid.......6.08 500 ml OP ✓ Nepro HP RTH

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully idised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML — Special Authority see SA1 Liquid		o <mark>us page</mark> – Hos 220 ml OP	✓ N	harmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110		s page – Hospi 237 ml OP	tal pha	rmacy [HP3]
Liquid (apricot) 125 ml Liquid (caramel) 125 ml	(3.31) 11.52	4 OP 4 OP	✓ F	NovaSource Renal Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

 ENTERAL (ORAL SEMI ELEMENTAL EEED 1.5KCAL(ML). Special Authority see SA1277 above. Hespital pharmacy (HR2)

Liquid	•	1,000 ml OP	
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority se	ee SA1377 above	– Hospital phar	macy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above -	Hospital pharm	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Au	thority see SA137	7 above – Hosp	oital pharmacy [HP3]
Liquid	12.04	1,000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 Child aged one to eight years: and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML - Special Authority see SA1196 above - Hospital pharmacy [HP3] 500 ml OP ✓ Nutrini Low Energy Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

255



(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (**Long-term medical condition**) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 255 - Hospital pharmacy [HP3]
Liquid.......7.00 1,000 ml OP ✓ Nutrison Energy

	Subsidy		Fully Brand or
	(Manufacturer's		,
	\$	Per	✓ Manufacturer
	Ψ	1 61	Wallulaciulei
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	page 255 - Ho	spital pharmacy [HP3]
Liquid		250 ml OP	✓ Isosource Standard
Liquid			
	5.29	1,000 ml OP	✓ Nutrison Standard
			RTH
			✓ Osmolite RTH
ENTERN FEED WITH FIRRE COOKON AND COOK ON THE	044050	055 11	
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority			
Liquid	5.29	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority se	e SA1859 on	page 255 – Hospi	tal pharmacy [HP3]
Liquid	5 29	1,000 ml OP	✓ Jevity RTH
<u> </u>		1,000 1111 01	✓ Nutrison Multi Fibre
			• Nutrison waiti Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority	see SA1859 on	page 255 - Hosi	oital pharmacy [HP3]
Liquid		250 ml OP	✓ Ensure Plus HN
Liquid	7.00	1,000 ml OP	✓ Ensure Plus RTH
	7.00	1,000 1111 0P	
			Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
ODAL EEED (DOM/DED)			
ORAL FEED (POWDER) – Special Authority see SA1859 on page			-
Powder (chocolate)	14.00	840 g OP	Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
December (com/He)			
Powder (vanilla)	14.00	840 g OP	 Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
		ŭ	
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa	ge 255 – Hosp	oital pharmacy [HF	23]
Additional subsidy by endorsement is available for patients be	eina bolus fed t	through a feeding	tube, who have severe
epidermolysis bullosa, or as exclusive enteral nutrition in child			
disease, or for patients with COPD and hypercapnia, defined	as CO2 value	exceeding bonnin	ng. The prescription must be
endorsed accordingly.			
Liquid (banana) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
Literation	(1.26)	200 1111 01	Ensure Plus
	' '		
	(1.26)		Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 r	nl		
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Limit (street and) High and their total OO and OOO and with			Ellouic Fluo
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml wi	, ,		•
		007 00	
Endorsement		237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
	(1.20)		i Ortioip

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 255 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Elquid (chocolate) Trighter subsidy of \$1.20 per 200 fill with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy (Manufacturer's \$	Price) Subs Per	Fully sidised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML – Special Authority see SA1195 c		age – Hospital 500 ml OP		cy [HP3] Nutrison Concentrated
	11.00	1,000 ml OP		Ensure Two Cal HN RTH Two Cal HN RTH
(Two Cal HN RTH Liquid to be delisted 1 February 2022)			٠.	WO GUITHVITTI
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients epidermolysis bullosa. The prescription must be endorsed at liquid (contile). History subsides of \$1.00 per 2000 polywith.	being bolus fed t			
Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with Endorsement	0.96 (1.90)	200 ml OP	Т	wo Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER -	Special Authority see SA1106 above - Hospital pharma	cy [HP3]	
Powder	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	Feed Thickener
			Karicare Antamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

	Cubaide	-	Fully Propd or
	Subsidy (Manufacturer's Price		Fully Brand or ised Generic
	\$	Per	✓ Manufacturer
GLUTEN FREE BAKING MIX - Special Authority see SA17	'29 on the previous page	- Hospital ph	narmacy [HP3]
Powder		.000 g OP	
	(5.15)	,	Healtheries Simple
	(00)		Baking Mix
CLUTEN EDEE DDEAD MIX Chaolial Authority and CA17	20 on the provious page	Hoonital ph	ŭ
GLUTEN FREE BREAD MIX – Special Authority see SA172 Powder		– поѕрцагрп ,000 g OP	amacy [nroj
rowdei		,000 g OF	NZB Low Gluten
	(7.32)		
	3.51		Bread Mix
			Harlana Dragal Min
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 or	the previous page – Ho	spital pharma	cy [HP3]
Powder	5.62 2,	,000 g OP	
	(18.10)		Horleys Flour
GLUTEN FREE PASTA – Special Authority see SA1729 on	the previous page - Hos	spital pharma	cv [HP3]
Buckwheat Spirals		250 g OP	-) []
Busininous opirale	(3.11)	200 g O.	Orgran
Corn and Vegetable Shells	\ /	250 g OP	Orgium
Containa Vogotable Chollo	(2.92)	-00 g Oi	Orgran
Corn and Vegetable Spirals		250 g OP	Orgini
Com and vegetable opirals	(2.92)	200 g Oi	Orgran
Rice and Corn Lasagne Sheets	\ /	200 g OP	Olgian
Tilde and Com Lasagne Oneels	(3.82)	200 g Oi	Orgran
Rice and Corn Macaroni	\ /	250 g OP	Olgian
Tilde and Com Macaroni	(2.92)	200 g Oi	Orgran
Rice and Corn Penne		250 g OP	Olgian
nice and Com Femile	(2.92)	250 g OF	Orgran
Rice and Maize Pasta Spirals	\ /	250 g OP	Olylali
nice and maize rasia spirals		250 g OF	Orgran
Rice and Millet Spirals	(2.92)	050 a OB	Olylan
nice and willer opirals		250 g OP	0
Dies and same anarhalti asselles	(3.11)	77 OD	Orgran
Rice and corn spaghetti noodles		375 g OP	0
Manadahla and Bias Osinala	(2.92)	250 - OD	Orgran
Vegetable and Rice Spirals		250 g OP	0
to P. T. T. T. T. W.	(2.92)		Orgran
Italian long style spaghetti		220 g OP	
	(3.11)		Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer Supplements For Homocystinuria AMINOACID FORMULA WITHOUT METHIONINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 500 a OP ✓ XMET Maxamum Powder 461.94 Supplements For MSUD AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] ✓ MSUD Maxamum 500 q OP Supplements For PKU AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 75 OP ✓ Phlexv 10 30 PKU Anamix Junior Orange Powder (chocolate) 36 g sachet......393.00 ✓ PKU Anamix Junior 30 Chocolate Powder (unflavoured) 28 g sachets......936.00 30 ✓ PKU Lophlex Powder ✓ PKU Anamix Junior 30 ✓ PKU Anamix Junior Powder (vanilla) 36 g sachet393.00 30 Vanilla ✓ PKU Anamix Infant 400 g OP ✓ XP Maxamum 500 a OP 500 g OP ✓ XP Maxamum 125 ml OP ✓ PKU Anamix Junior LQ 125 ml OP ✓ PKU Anamix Junior ✓ PKU Anamix Junior 125 ml OP LQ Liquid (forest berries), 250 ml carton......540.00 18 OP ✓ Easiphen Liquid Liquid (juicy tropical) 125 ml......936.00 ✓ PKU Lophlex LQ 20 30 OP 36 OP ✓ PKU Lophlex

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page – Hospital pharmacy [HP3]

Powder8.22 500 g OP ✓ Loprofin Mix

Liquid (juicy berries) 62.5 ml......939.00

Liquid (juicy orange) 125 ml936.00

Sensation 20

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 20

✓ PKU Lophlex LQ 20

60 OP

60 OP

60 OP

30 OP

30 OP

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
LOW PROTEIN PASTA - Special Authority see SA1108 on page	e 261 – Hospital	pharmacy [HP:	3]	
Animal shapes	11.91	500 g OP	✓ L	.oprofin
Lasagne	5.95	250 g OP	√ L	.oprofin
Low protein rice pasta	11.91	500 g OP	√ L	.oprofin
Macaroni	5.95	250 g OP	√ L	.oprofin
Penne	11.91	500 g OP	√ L	.oprofin
Spaghetti	11.91	500 g OP	√ L	.oprofin
Spirals	11.91	500 g OP	√ L	oprofin

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see SA1940 below – Hospital pha Powder43.60	400 g OP	✓ Alfamino✓ Alfamino Junior
Powder (unflavoured)53.00	400 g OP	✓ Elecare ✓ Elecare LCP ✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	 ✓ Neocate SYNEO ✓ Elecare ✓ Neocate Junior ✓ Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or

continued...

Subsi	sidy Fu	ly Brand or
(Manufacture	rer's Price) Subsidise	d Generic
\$	Per	Manufacturer

- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and

continued...

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

- 2.2.3 Amino acid formula is required for a nutritional deficit; and
- 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis: or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA	 Special Authority see SA1953 below 	- Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Fither:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Sov milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Vaccinations** BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm] For infants at increased risk of tuberculosis. Increased risk is defined as: 1) living in a house or family with a person with current or past history of TB; or 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm] Funded for any of the following criteria: 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or 5) A single dose for vaccination of patients aged from 65 years old; or 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or 7) For vaccination of previously unimmunised or partially immunised patients: or 8) For revaccination following immunosuppression; or 9) For boosting of patients with tetanus-prone wounds. Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous 10 **Boostrix Boostrix** DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm] Funded for any of the following: 1) A single dose for children up to the age of 7 who have completed primary immunisation; or 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive 4) Five doses will be funded for children requiring solid organ transplantation. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

Infanrix IPV

10

Subsidised

Fully

Brand or

Generic

✓ <u>Havrix</u>✓ Havrix Junior

Subsidy

(Manufacturer's Price)

·	\$	Per	✓ Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HA [Xpharm]	EMOPHILUS	INFLUENZ	ZAE TYPE B VACCINE -
Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of 10 for	nrimary immi	inisation: o	nr
2) An additional four doses (as appropriate) are funded for (re-)in			
10 who are patients post haematopoietic stem cell transplantat			
post solid organ transplant, renal dialysis and other severely in			
3) Up to five doses for children up to and under the age of 10 rec			
Note: A course of up-to four vaccines is funded for catch up program	•		
to complete full primary immunisation. Please refer to the Immunisa			
programmes.	lion Handboo	ik ioi iiie a	ppropriate scriedule for catch up
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	✓ Infanrix-hexa
	.0.00	10	IIIIaiiiix-iiexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
An additional dose (as appropriate) is funded for (re-)immunisa			
transplantation, or chemotherapy; functional asplenic; pre or po			
or post cochlear implants, renal dialysis and other severely imr		•	· ·
For use in testing for primary immunodeficiency diseases, on the	ne recommen	dation of a	ın internal medicine physician oı
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg;			
prefilled syringe plus vial 0.5 ml	.0.00	1	✓ Hiberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
1) Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver disease;	or		
3) One dose of vaccine for close contacts of known hepatitis A ca			
,			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the following crite		1	√ E	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual int for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HS) following needle stick injury. 	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of ercourse; or) pos derec	itive; or I not to have	
Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following crite		1	✓ <u>E</u>	ngerix-B
 for household or sexual contacts of known acut for children born to mothers who are hepatitis B for children up to and under the age of 18 years serology and require additional vaccination or reference of the patients or for hepatitis C positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual int for patients following immunosuppression; or for solid organ transplant patients; or following needle stick injury; or for dialysis patients; or for liver or kidney transplant patients. 	s surface antigen (HBsAgs s inclusive who are consic equire a primary course of tercourse; or) pos derec	itive; or I not to have	
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AN Any of the following:	,	- [Xpl	narm]	
1) Maximum of two doses for children aged 14 years a 2) Maximum of three doses for patients meeting any o 1) People aged 15 to 26 years inclusive; or 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: 3) Maximum of four doses for people aged 9 to 26 years.	f the following criteria: or	nerap	у	
Inj 270 mcg in 0.5 ml syringe	0.00	10	√ <u>G</u>	ardasil 9

Subsidy (Manufacturer's Price)	Sı	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

- [Xpharm]......9.00 1 ✓ Afluria Quad Junior (2021 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Sı	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable

C)

A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes: or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine).......90.00 10 ✓ Fluad Quad (2021 Formulation)

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

C)

- INFLUENZA VACCINE people 65 years and over
 is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine)	_		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

	Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

A) INFLUENZA VACCINE - people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

((Subsidy Fully Manufacturer's Price) Subsidised		Brand or Generic
	\$	Per 🗸	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant: or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C 1 PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00

✓ Synflorix

10

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Immunisation Handbook for the appropriate appropriate appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for the appropriate please refer to the Immunisation Handbook for	priate schedule for	catch up programmes
Inj 30.	8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,	•	

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml		
syringe	10	Prevenar 13
	1	✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]			
Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochl All of the following:	ctional asplenia, pre- or p	oost-solid	organ t	ransplant, renal dialysis,
a) Patient is a child under 18 years for (re-)immurb) Treatment is for a maximum of two doses; andc) Any of the following:	isation; and			
 i) on immunosuppressive therapy or radiati immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or or vi) with cochlear implants or intracranial shu vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more prednisone of 2 mg/kg per day or greater 20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks greater 	or gan transplantation (incl nts; or than two weeks, and wh , or children who weigh or g asthma treated with high	o are on a	ematopo an equiv 1 10 kg o	vietic stem cell transplant); ralent daily dosage of on a total daily dosage of
xi) with cardiac disease, with cyanosis or failxii) with diabetes; orxiii) with Down syndrome; orxiv) who are pre-or post-splenectomy, or with				
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ng:	1	√ <u>P</u>	neumovax 23
2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe	ropriate schedule for cal	tch-up pro 1	ogramm ✓ <u>IF</u>	
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator		10	√ <u>R</u>	otarix

	NATIONAL II	MMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm] Either:			
1) Maximum of one dose for primary vaccination for eith a) Any infant born on or after 1 April 2016; or b) For previously unvaccinated children turning 11 varicella infection (chickenpox), or 2) Maximum of two doses for any of the following: a) Any of the following for non-immune patients: i) with chronic liver disease who may in futur ii) with deteriorating renal function before tra iii) prior to solid organ transplant; or iv) prior to any elective immunosuppression*, v) for post exposure prophylaxis who are imressions.	years old on or after 1 Ju re be candidates for trans nsplantation; or		ave not previously had a

- b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
- c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
- d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
- e) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
- f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
- g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* immu	unosuppression due to	steroid or other imr	nunosuppressive th	herapy must be fo	or a treatment perio	od of greater than
28 day	'S					

Inj 1350 PFU prefilled syringe	0.00	1	✓ Varivax
		10	✓ Varivax

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting either of the following criteria:

1) One dose for all people aged 65 years; or

2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 December 2021.

Inj 19,400 PFU prefilled syringe plus vial	0.00	1	✓ Zostavax
		10	✓ Zostavax

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]			
Ini 5 TU per 0.1 ml. 1 ml vial	0.00	1	✓ Tuberso

- Symbols -	Agents for Parkinsonism and Related	Amsidine156
UK Synacthen82	Disorders 122	Amzoate31
3TC108	Agents Used in the Treatment of	Anaesthetics124
- A -	Poisonings243	Anagrelide hydrochloride156
A-Scabies70	Agrylin156	Analgesics
Abacavir sulphate108	Albendazole91	Anastrozole172
Abacavir sulphate with	Albey228	Anatrole
lamivudine 108	Albustix79	Andriol Testocaps82
Abiraterone acetate169	Aldurazyme30	Androderm82
Acarbose11	Alecensa163	ANI51
Accarb11	Alectinib163	Anoro Ellipta232
Accuretic49	Alendronate sodium115	Antabuse147
Accuretic 1049	Alendronate sodium with	Antacids and Antiflatulents6
Accuretic 2049	colecalciferol115	Anthelmintics91
Acetazolamide240	Alfacalcidol34	Antiacne Preparations63
Acetec48	Alfamino263	Antiallergy Preparations228
Acetic acid with 1, 2- propanediol	Alfamino Junior263	Antianaemics38
diacetate and	Alginic acid6	Antiandrogen Oral
benzethonium237	Alglucosidase alfa27	Contraceptives77
Acetic acid with hydroxyquinoline and	Alkeran	Antiarrhythmics50
ricinoleic acid77	Alkeran S29152	Antibacterials91
Acetylcysteine243	Allersoothe229	Antibacterials Topical63
Aci-Jel77	Allmercap154	Anticholinergic Agents231
Aciclovir	Allopurinol119	Anticholinesterases113
Infection103	Alpha-Adrenoceptor Blockers48	Antidepressants130
Sensory237	Alpha-Keri Lotion68	Antidiarrhoeals6
Acidex6	Alphamox94	Antiepilepsy Drugs131
Acipimox56	Alphamox 12594	Antifibrinolytics, Haemostatics and
Acitretin70	Alphamox 25094	Local Sclerosants39
Aclasta117	Alprolix39	Antifibrotics232
Aclin114	Alu-Tab6	Antifungals99
Actemra216	Aluminium hydroxide6	Antifungals Topical64
Actinomycin D157	Alvogen52	Antihistamines229
Actrapid10	Amantadine hydrochloride122	Antihypotensives51
Actrapid Penfill10	Ambrisentan59	Antimalarials101
Acular239	Ambrisentan Mylan59	Antimigraine Preparations 135
Acupan125	Amiloride hydrochloride55	Antinausea and Vertigo Agents 135
Adalimumab180	Amiloride hydrochloride with	Antipruritic Preparations65
Adapalene63	furosemide56	Antipsychotics136
Adenuric120	Amiloride hydrochloride with	Antiretrovirals106
ADR Cartridge 1.824	hydrochlorothiazide56	Antirheumatoid Agents114
Adrenaline58	Aminophylline234	Antispasmodics and Other Agents
Advantan66	Amiodarone hydrochloride50	Altering Gut Motility8
Advate42	Amisulpride136	Antithrombotic Agents42
Adynovate42	Amisulpride Mylan136	Antithymocyte globulin
Afinitor224	Amitriptyline130	(equine) 180
Aflibercept190	Amlodipine53	Antitrichomonal Agents101
Afluria Quad	Amneal160	Antituberculotics and
(2021 Formulation) 272	Amorolfine64	Antileprotics101
Afluria Quad Junior	Amoxicillin94	Antiulcerants8
(2021 Formulation) 271	Amoxicillin with clavulanic acid95	Antivirals103
AFT-Pyrazinamide102	Amphotericin B33	Anxiolytics140
Agents Affecting the	Amsacrine156	Anzatax159
Renin-Angiotensin System 48	AmsaLyo156	Apidra11

Apidra SoloStar	11	Arrow-Topiramate		Beclazone 250	22
Apo-Azithromycin	92	Arrow-Tramadol	130	Beclazone 50	22
Apo-Bromocriptine	122	Arsenic trioxide	156	Beclomethasone dipropionate	
Apo-Ciclopirox	64	Asacol	<mark>7</mark>	Bee venom allergy treatment	22
Apo-Clarithromycin		Asamax	7	Bendamustine hydrochloride	
Alimentary	9	Ascorbic acid	34	Bendrofluazide	5
Infection	93	Aspen Adrenaline	58	Bendroflumethiazide	
Apo-Clomipramine	130	Aspirin		[Bendrofluazide]	50
Apo-Diclo SR	113	Blood	42	Benzathine benzylpenicillin	9
Apo-Diltiazem CD	54	Nervous	125	Benzatropine mesylate	12
Apo-Doxazosin	48	Asthalin	231	Benzbromaron AL 100	
Apo-Folic Acid		Atazanavir sulphate	108	Benzbromarone	119
Apo-Furosemide	<u>55</u>	Atenolol	51	Benztrop	12
Apo-Gabapentin		Atenolol AFT	51	Benzydamine hydrochloride	
Apo-Megestrol		Atenolol AFT S29	51	Benzylpenicillin sodium [Penicillin	
Apo-Metoprolol		ATGAM	180	Gj	9!
Apo-Mirtazapine		Ativan		Beta Cream	
Apo-Nadolol		Atomoxetine	143	Beta Ointment	6
Apo-Oxybutynin		Atorvastatin		Beta Scalp	
Apo-Pindolol		Atropine sulphate		Beta-Adrenoceptor Agonists	
Apo-Prazosin		Cardiovascular	50	Beta-Adrenoceptor Blockers	
Apo-Prednisone		Sensory		Betadine	
Apo-Primidone		Atropt		Betadine Skin Prep	
Apo-Propranolol		Atrovent		Betaferon	
Apo-Pyridoxine		AU Synacthen		Betahistine dihydrochloride	
Apo-Selegiline S29		Aubagio		Betaine	
Apo-Sumatriptan		Augmentin		Betaloc CR	
Apo-Temozolomide		Aurorix		Betamethasone dipropionate	
Apomorphine hydrochloride		AutoSoft 30		Betamethasone dipropionate with	0
Aprepitant		AutoSoft 90		calcipotriol	71
Apresoline		Avelox		Betamethasone sodium phosphate	/ \
Aptamil AllerPro SYNEO 1		Avonex		with betamethasone acetate	0
•		Avonex Pen		Betamethasone valerate6	
Aptamil AllerPro SYNEO 2					
Aptamil Gold+ Pepti Junior		Azacitidine		Betamethasone valerate with sodiu	
Aqueous cream		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aratac		Azamun		Betaxolol	
Arava		Azathioprine		Betnovate	
Arginine		Azilect		Betoptic	
Aripiprazole		Azithromycin		Betoptic S	
Aripiprazole Sandoz		Azopt		Bexsero	
Aristocort		AZT	108	Bezafibrate	
Arrow - Lattim		- B -		Bezalip	
Arrow-Amitriptyline		B-D Micro-Fine		Bezalip Retard	
Arrow-Bendrofluazide		B-D Ultra Fine		Bicalutamide	
Arrow-Brimonidine		B-D Ultra Fine II	16	Bicillin LA	9
Arrow-Diazepam		Bacillus Calmette-Guerin (BCG)		BiCNU	
Arrow-Doxorubicin	157	vaccine	180	Bicnu Heritage	
Arrow-Losartan &		Bacillus Calmette-Guerin		Bile and Liver Therapy	10
Hydrochlorothiazide	49	vaccine	268	Biltricide	9
Arrow-Norfloxacin	112	Baclofen	121	Bimatoprost	24
Arrow-Ornidazole	101	Bactroban	64	Bimatoprost Multichem	
Arrow-Quinapril 10	49	Balance		Binarex	170
Arrow-Quinapril 20		Barrier Creams and Emollients	67	Binocrit	3
Arrow-Quinapril 5		Basic AquaCream		Biodone	12
Arrow-Roxithromycin		BCG Vaccine		Biodone Extra Forte	12
Arrow-Timolol		Beclazone 100		Biodone Forte	
***************************************			-		-

			_
Bisacodyl27	Calcitriol34	Celebrex1	1
Bisoprolol fumarate51	Calcitriol-AFT34	Celecoxib1	1
Bisoprolol Mylan51	Calcium carbonate6, 35	Celecoxib Pfizer1	1
BK Lotion68	Calcium Channel Blockers53	Celestone Chronodose	3
Bleomycin sulphate156	Calcium Disodium Versenate244	Cellcept17	7
Blood Colony-stimulating	Calcium folinate153	Centrally-Acting Agents	5
Factors45	Calcium Folinate Ebewe153	Cephalexin ABM	
Blood glucose diagnostic test	Calcium Folinate Sandoz153	Cetirizine hydrochloride22	
meter 15	Calcium Folinate Sandoz S29 153	Cetomacrogol	3
Blood glucose diagnostic test	Calcium gluconate35	Cetomacrogol with glycerol	
strip	Calcium Homeostasis80	Cetuximab19	
Blood glucose test strips (visually	Calcium polystyrene sulphonate47	Champix14	
impaired)15	Calcium Resonium47	Charcoal24	4
Blood Ketone Diagnostic Test	Calcium-Sandoz Forte35	Chemotherapeutic Agents15	5
Strip 14	Calogen250	Chickenpox vaccine27	7
Bonjela33	Calutide-50 170	Chlorafast23	3
Boostrix268	Candesartan cilexetil49	Chlorambucil15	5
Bortezomib	Candestar49	Chloramphenicol23	3
Bortezomib Dr-Reddy's156	Canesten64	Chlorothiazide	
Bortezomib Juno156	Capecitabine153	Chlorpheniramine maleate22	
Bosentan59	Capercit	Chlorpromazine hydrochloride 13	
Bosentan Dr Reddy's59	Capoten48	Chlortalidone [Chlorthalidone]	
Bplex34	Capsaicin	Chlorthalidone	
Breo Ellipta230	Musculoskeletal114	Chlorvescent	4
Brevinor 1/2876	Nervous125	Choice Load 375	7
Bricanyl Turbuhaler231	Captopril48	Choice TT380 Short	
Brilinta43	Captopril-Mylan48	Choice TT380 Standard	
Brimonidine tartrate240	Carafate9	Choline salicylate with cetalkonium	
Brimonidine tartrate with timolol	Carbaccord151	chloride	3
maleate240	Carbamazepine132	Ciclopirox olamine	ô
Brinzolamide240	Carbimazole85	Ciclosporin22	
Brolene237	Carbomer241	Cilazapril	4
Bromocriptine mesylate122	Carboplatin 151	Cilicaine	Э.
Brufen SR113	Carboplatin Ebewe151	Cilicaine VK	Э.
Buccastem136	Carbosorb-X243	Cinacalcet	
Budesonide	Cardinol LA53	Cipflox	Э
Alimentary6	Cardizem CD54	Ciprofloxacin	
Respiratory229, 235	CareSens Dual14	Infection	Э
Budesonide with eformoterol230	CareSens N15	Sensory23	3
Bumetanide55	CareSens N POP15	Ciprofloxacin Teva23	
Buprenorphine Naloxone BNM146	CareSens N Premier15	Circadin14	4
Buprenorphine with naloxone146	CareSens PRO15	Cisplatin15	
Bupropion hydrochloride147	Carmellose sodium with gelatin and	Cisplatin Ebewe15	
Burinex55	pectin33	Citalopram hydrobromide 13	
Burinex S2955	Carmustine 151	Cladribine15	5
Buscopan8	Carvedilol52	Clarithromycin	
Buspirone hydrochloride140	Carvedilol Sandoz52	Alimentary	
Busulfan151	Catapres54	Infection	9
- C -	CeeNU152	Clexane	4
Cabergoline90	Cefaclor monohydrate91	Clexane Forte	
Cacit	Cefalexin91	Climara	
Caffeine citrate236	Cefalexin Sandoz91	Climara	
Calamine65	Cefazolin91	Clindamycin	
Calci-Tab 50035	Ceftriaxone91	Clinicians	
Calcipotriol71	Ceftriaxone-AFT91	Clinicians Renal Vit	
Calcitonin80	Cefuroxime axetil92	Clobazam13	

Clobetasol propionate66, 72	Corticosteroids and Related Agents	DBL Adrenaline50
Clobetasone butyrate66	for Systemic Use81	DBL Aminophylline234
Clofazimine101	Corticosteroids Topical65	DBL Bleomycin Sulfate150
Clomazol	Cortifoam7	DBL Carboplatin15
Dermatological64	Cosentyx214	DBL Cisplatin15
Genito-Urinary77	Cosmegen	DBL Dacarbazine15
Clomifene citrate90	Coumadin45	DBL Desferrioxamine Mesylate for Inj
Clomipramine hydrochloride130	Country Life30	BP24
Clomipramine Teva130	Coversyl48	DBL Docetaxel15
Clonazepam 131–132, 140	Creon 1000025	DBL Ergometrine7
Clonidine54	Creon 2500025	DBL Gemcitabine154
Clonidine BNM54	Creon Micro25	DBL Gentamicin9
Clonidine hydrochloride54	Crotamiton65	DBL Heparin Sodium4
Clopidogrel43	Crystaderm63	DBL Leucovorin Calcium15
Clopidogrel Multichem43	Curam95	DBL Methotrexate Onco-Vial15
Clopine137	Curam Duo 500/12595	DBL Morphine Sulphate128
Clopixol138, 140	Cvite34	DBL Naloxone Hydrochloride24
Clotrimazole	Cyclizine hydrochloride136	DBL Octreotide17
Dermatological64	Cyclizine lactate136	DBL Pethidine Hydrochloride 129
Genito-Urinary77	Cyclogyl241	DBL Vinblastine162
Clozapine137	Cyclonex151	DBL Vincristine Sulfate162
Clozaril137	Cyclopentolate hydrochloride241	Decozol3
Co-trimoxazole98	Cyclophosphamide151	Deferasirox24
Coal tar71	Cyclorin102	Deferiprone24
Coal tar with allantoin, menthol,	Cycloserine102	Denosumab11
phenol and sulphur71	Cyproterone acetate82	Deolate100
Coal tar with salicylic acid and	Cyproterone acetate with	Deoxycoformycin160
sulphur71	ethinyloestradiol77	Depo-Medrol82
Coco-Scalp71	Cystadane28	Depo-Provera70
Codeine phosphate	Cytarabine	Depo-Testosterone82
Extemporaneous246	Cytotec8	Deprim98
Nervous127	Cytoxan151	Dermol66, 72
Coenzyme Q1029	- D -	Desferrioxamine mesilate24
Colchicine	D-Penamine114	Desmopressin8
Colecalciferol34	Dabigatran45	Desmopressin acetate9
Colestid56	Dacarbazine157	Desmopressin-PH&T9
Colestipol hydrochloride56	Dacarbazine APP157	Desuric119
Colgout119	Dactinomycin [Actinomycin D]157	Detection of Substances in
Colifoam7	Daivobet70	Urine79
Colistin sulphomethate96	Daivonex71	Dexamethasone
Colistin-Link96	Daktarin65	Hormone8
Collodion flexible246	Dalacin C96	Sensory23
Colloidal bismuth subcitrate9	Dantrium 121	Dexamethasone phosphate8
Colofac8	Dantrium S29121	Dexamethasone Phosphate
Coloxyl26	Dantrolene121	Panpharma8
Combigan240	Daonil11	Dexamethasone with framycetin and
Compound electrolytes47	Dapa-Tabs56	gramicidin 23
Compound electrolytes with glucose	Dapsone 102	Dexamethasone with neomycin
[Dextrose]47	Daraprim97	sulphate and polymyxin B
Compound hydroxybenzoate246	Darunavir108	sulphate23
Concerta145	Darunavir Mylan108	Dexamfetamine sulfate14
Condoms74	Dasatinib163	Dexmethsone8
Condyline72	Daunorubicin157	Dextrochlorpheniramine
Contraceptives - Hormonal75	David One Step Cassette Pregnancy	maleate22
Contraceptives - Non-hormonal73	Test78	Dextrose 46–4
Copaxone141	DBL Acetylcysteine243	DHC Continus12

Diabetes	Dosulepin [Dothiepin]		Emend Tri-Pack	13
Diabetes Management		130	Emicizumab	
Diacomit13			EMLA	124
Diagnostic Agents27			Empagliflozin	1
Diamide Relief			Empagliflozin with metformin	
Diamox24		95	hydrochloride	13
Diasip25		157	Emtricitabine	
Diason RTH25			Emtricitabine with tenofovir	
Diazepam132, 14			disoproxil	10
Diazoxide			Emtriva	
Dibenzyline			Emulsifying ointment	
Diclofenac Sandoz11			Emulsifying Ointment ADE	
Diclofenac sodium	Dr Reddy's Omeprazole		Enalapril maleate	
Musculoskeletal11			Enbrel	
Sensory23			Endocrine Therapy	
Differin		115	Endoxan	
Difflam			Engerix-B	
Diflucan	•		Enlafax XR	
Digestives Including Enzymes			Enoxaparin sodium	
Digoxin			Enstilar	
Dihydrocodeine tartrate			Ensure	
			Ensure Plus	
Dilantin			Ensure Plus HN	
			Ensure Plus RTH	
Diltiazem hydrochloride				
Dilzem		56	Ensure Two Cal HN RTH	
Dimethicone		000	Entacapone	124
Dimethyl fumarate14			Entapone	
Dipentum			Entecavir	
Diphtheria, tetanus and pertussis	e-chamber Turbo		Entecavir Sandoz	
vaccine			Entocort CIR	
Diphtheria, tetanus, pertussis and	Ear Preparations		Entresto 24/26	
polio vaccine			Entresto 49/51	
Diphtheria, tetanus, pertussis, polio,	Easiphen Liquid		Entresto 97/103	
hepatitis B and haemophilus	Econazole nitrate		Epilim	
influenzae type B vaccine 26		107	Epilim Crushable	
Diprosone			Epilim IV	
Diprosone OV			Epilim S/F Liquid	
Dipyridamole			Epilim Syrup	
Disopyramide phosphate		230	Epirubicin Ebewe	
Disulfiram14	ŭ .		Epirubicin hydrochloride	
Diuretics	•		Eplerenone	
Docetaxel15			Epoetin alfa	
Docetaxel Accord15		71	Epoprostenol	6
Docetaxel Sandoz15	57 Elaprase	29	Eptacog alfa [Recombinant factor	
Docusate sodium			VIIa]	
Docusate sodium with	Elecare	263	ERA	94
sennosides	Elecare LCP	263	Erbitux	19
Dolutegravir10	9 Electral	47	Ergometrine maleate	
Domperidone13	B6 Elelyso	32	Erlotinib	164
Donepezil hydrochloride14	Elemental 028 Extra	254	Erythrocin IV	
Donepezil-Rex14	16 Elidel	71	Erythromycin (as lactobionate)	
Dornase alfa23	34 Elocon		Erythromycin ethyl succinate	94
Dortimopt24		66	Erythromycin stearate	94
Dorzolamide hydrochloride24			Esbriet	233
Dorzolamide with timolol24			Escitalopram	
Dostinex			Escitalopram (Ethics)	
			the state of the s	

Eskazole	91	Ferinject	36	Fluticasone furoate with	
Essential Generics		Ferodan		vilanterol	230
Essential Prednisolone		Ferriprox	243	Fluticasone propionate	235
Estradiol TDP Mylan	83	Ferro-F-Tabs	36	Fluticasone with salmeterol	231
Estradot	83	Ferro-tab		FML	
Estradot 50 mcg	83	Ferrograd	36	Foban	64
Estrofem	83	Ferrosig	36	Folic acid	
Etanercept	173	Ferrous fumarate	36	Folic Acid Mylan	39
Ethambutol hydrochloride	102	Ferrous fumarate with folic acid	36	Food Thickeners	260
Ethics Aspirin	125	Ferrous sulfate	36	Foods And Supplements For In	born
Ethics Aspirin EC		Fexofenadine hydrochloride	229	Errors Of Metabolism	261
Ethics Lisinopril	48	Fibro-vein	42	Foradil	230
Ethics Paracetamol Classic	126	Filgrastim	45	Forteo	116
Ethinyloestradiol	84	Finasteride	78	Fortini	253
Ethinyloestradiol with		Fingolimod	141	Fortini Multi Fibre	253
desogestrel	75	Firazyr	228	Fortisip	258
Ethinyloestradiol with		Flagyl		Fortisip Multi Fibre	259
levonorgestrel	76	Flagyl-S	101	Fosamax	
Ethinyloestradiol with		Flamazine		Fosamax Plus	
norethisterone	76	Flecainide acetate	50	Framycetin sulphate	237
Ethosuximide	132	Flecainide BNM	50	Frisium	
Etopophos	157	Flecainide Controlled Release		Frumil	
Etoposide	157	Teva	<u>50</u>	Frusemide	55
Etoposide phosphate		Fleet Phosphate Enema	27	Fucicort	66
Etravirine		Flixonase Hayfever & Allergy	235	Fucidin	98
Eumovate	66	Flixotide		Fucithalmic	237
Everet	133	Flixotide Accuhaler		Fulvestrant	170
Everolimus	224	Florinef	81	Fungilin	
Evista	116	Fluad Quad		Furosemid-Ratiopharm	
Exemestane	172	(2021 Formulation)	272	Furosemide [Frusemide]	
Exjade		Fluanxol		Furosemide-Baxter	
Extemporaneously Compoun		Flucil		fusidic acid	
Preparations and		Flucloxacillin	95	Dermatological	64. 66
Galenicals	246	Flucloxin		Infection	
Eye Preparations		Flucon		Sensory	
Eylea		Fluconazole		´ - G -	
Ezetimibe		Fludara Oral		Gabapentin	132
Ezetimibe Sandoz		Fludarabine Ebewe		Gacet	
Ezetimibe with simvastatin		Fludarabine phosphate		Galsulfase	
-F-		Fludrocortisone acetate		Galvumet	
Factor eight inhibitor bypassi	na	Fluids and Electrolytes		Galvus	
fraction	-	Flumetasone pivalate		Gardasil 9	
Famotidine		Fluocortolone caproate with		Gastrodenol	
Famotidine Hovid		fluocortolone pivalate and		Gaviscon Double Strength	
Faslodex		cinchocaine	8	Gaviscon Infant	
Febuxostat		Fluorometholone		Gazyva	
Febuxostat multichem		Fluorouracil		Gefitinib	
Feed Thickener Karicare		Fluorouracil Accord		Gemcitabine Ebewe	
Aptamil		Fluorouracil Ebewe		Gemcitabine hydrochloride	
FEIBA NF		Fluorouracil sodium		Genoptic	
Felo 10 ER		Fluox		Gentamicin sulphate	
Felo 5 ER		Fluoxetine hydrochloride		Infection	96
Felodipine		Flupenthixol decanoate		Sensory	237
Femme-Tab ED	76	Flutamide		Gilenya	141
Fentanyl		Flutamin		Ginet	
Fentanyl Sandoz		Fluticasone		Glatiramer acetate	
. Jinanyi Janabeninini				GIGHT ATTION GOOD TO THE TOTAL THE TOTAL TO THE TOTAL TOT	

Glecaprevir with pibrentasvir	5 Hepatitis A vaccine269	Hypam	14
Glibenclamide1	·	Hyperuricaemia and Antigout	11
Gliclazide1	2 vaccine270	Hypromellose	
Glipizide1		Hypromellose with dextran	
Glivec16		-1-	
Glizide1	2 Hiprex112	Ibiamox	9
Glucagen Hypokit1	0 Histaclear229	Ibrance	16
Glucagon hydrochloride1		Ibuprofen	11
Glucerna Select25		Ibuprofen SR BNM	11
Glucobay1	1 Horleys Bread Mix261	Icatibant	
Glucose [Dextrose]	6 Horleys Flour261	Idarubicin hydrochloride	15
Gluten Free Foods26		Idursulfase	2
Glycerin with sodium saccharin 24		Ifosfamide	15
Glycerin with sucrose24		Igroton	5
Glycerol	Humalog11	lkorel	
Alimentary2		llevro	23
Extemporaneous24		lloprost	6
Glyceryl trinitrate	Human papillomavirus (6, 11, 16, 18,	Imatinib mesilate	16
Alimentary		Imatinib-Rex	
Cardiovascular		Imigran	
Glycopyrronium23		Imipramine hydrochloride	
Glycopyrronium bromide		Imiquimod	
Glycopyrronium with	HumiraPen180	Immune Modulators	10
indacaterol23	2 Humulin 30/7011	Immunosuppressants	
Go Healthy2	9 Humulin NPH10	Imuran	
Gold Knight7		Incruse Ellipta	
Gold Knight XL74–7		Indacaterol	23
Goserelin	•	Indapamide	
Gutron		Infanrix IPV	
Gynaecological Anti-infectives		Infanrix-hexa	26
́ -Н-	Dermatological66	Infant Formulae	26
Habitrol14	8 Hormone81	Infatrini	26
Haemophilus influenzae type B	Hydrocortisone (PSM)66	Infliximab	19
vaccine26		Influenza vaccine	
Haldol13	•	Influvac Tetra	
Haldol Concentrate13	pramoxine hydrochloride	(2021 Formulation)	27
Haldol Decanoas13		Inhaled Corticosteroids	
Haloperidol13		Inhaled Long-acting	
Haloperidol decanoate13		Beta-adrenoceptor Agonists	23
Harvoni10	•	Inspra	
Havrix26	9 Hydrocortisone with miconazole 66	Instillagel Lido	
Havrix Junior26	9 Hydrocortisone with natamycin and	Insulin aspart	
Haylor syrup22		Insulin aspart with insulin aspart	
healthE Calamine Aqueous Cream	Hydrogen peroxide63	protamine	1
BP6		Insulin glargine	1
healthE Dimethicone 10%		Insulin glulisine	
healthE Dimethicone 4% Lotion6		Insulin isophane	1
healthE Dimethicone 5%	7 hydroxycarbamide157	Insulin isophane with insulin	
healthE Glycerol BP24		neutral	1
healthE Urea Cream		Insulin lispro	
Healtheries Simple Baking Mix26		Insulin lispro with insulin lispro	
Hemastix7		protamine	1
Hemlibra		Insulin neutral	
Heparin sodium		Insulin pen needles	
Heparinised saline		Insulin pump	
Heparon Junior25		Insulin pump cartridge	

			1 1 0 1 0:	-
Insulin pump infusion set (steel	- J -		Lantus SoloStar	
cannula)	Jadelle		Lanvis	
Insulin pump infusion set (steel	Jakavi		Lanzol Relief	
cannula, straight insertion) 22	Jardiamet		Lapatinib ditosylate	
Insulin pump infusion set (teflon	Jardiance		Largactil	
cannula)23	Jaydess		Laronidase	
Insulin pump infusion set (teflon	Jevity HiCal RTH		Lasix	
cannula, angle insertion with	Jevity RTH		Latanoprost	
insertion device)24	Juno Pemetrexed	155	Latanoprost with timolol	
Insulin pump infusion set (teflon	- K -		Lax-Suppositories	
cannula, angle insertion) 24	Kadcyla		Lax-Tab	
Insulin pump infusion set (teflon	Kaletra		Laxatives	
cannula, straight insertion with	Kalydeco	234	Laxsol	
insertion device)24	Kemadrin	123	Ledipasvir with sofosbuvir	
Insulin pump infusion set (teflon	Kenacomb	2 <mark>37</mark>	Leflunomide	114
cannula, straight insertion) 24	Kenacort-A 10	82	Lenalidomide	158
Insulin pump reservoir24	Kenacort-A 40	82	Letrole	
Insulin syringes, disposable with	Kenalog in Orabase	33	Letrozole	172
attached needle16	Kenkay Sorbolene	67	Leukeran FC	151
Intal Forte CFC Free234	Ketocal 3:1	267	Leukotriene Receptor	
Intelence107	KetoCal 4:1	267	Antagonists	234
Interferon beta-1-alpha141	Ketoconazole		Leuprorelin	
Interferon beta-1-beta141	Dermatological	72	Leustatin	
Intra-uterine device75	Infection		Levetiracetam	133
Invega Sustenna139	Ketogenic Diet	267	Levetiracetam-AFT	
IPCA-Frusemide55	Ketoprofen		Levlen ED	
IPCA-Metoprolol52	Ketorolac trometamol		Levocabastine	
IPCA-Propranolol53	KetoSens		Levocarnitine	
IPOL278	Keytruda		Levodopa with benserazide	
Ipratropium bromide231, 235	Kindergen		Levodopa with carbidopa	
Iressa	Kivexa		Levomepromazine	
Irinotecan Accord154	Klacid		Levomepromazine	
Irinotecan Actavis 100 154	Alimentary	9	hydrochloride	137
Irinotecan hydrochloride	Infection		Levonorgestrel	
Irinotecan-Rex	Kliogest		Genito-Urinary	76–77
Iron (as ferric carboxymaltose)36	Kliovance		Hormone	
Iron polymaltose36	Kogenate FS		Levothyroxine	
Isentress	Konakion MM		Lidocaine [Lignocaine]	
Isentress HD	Konsyl-D		Lidocaine [Lignocaine]	
Ismo 20	Kuvan		hydrochloride	124
Ismo 40 Retard58	-L-		Lidocaine [Lignocaine] with	12-
Isoniazid102	Labetalol	52	chlorhexidine	124
Isoniazid with rifampicin	Lacosamide		Lidocaine [Lignocaine] with	12-
Isoptin54	Lactulose		prilocaine	12/
Isoptin Retard54	Laevolac		Lidocaine-Baxter	
Isoptin SR54	Lamictal		Lidocaine-Claris	
Isopto Carpine240	Lamivudine		Life Extension	124
Isosorbide mononitrate	Lamivudine Alphapharm		LignocaineLioresal Intrathecal	124
	Lamotrigine			
Isotretinoin	Lamprene		Lipid-Modifying Agents	
Ispaghula (psyllium) husk	Lanoxin Lanoxin Paediatric Elixir		Liquigen	25(
Itch-Soothe 65			Lisinopril	48
Itraconazole	Lanoxin PG		Lithium carbonate	
Itrazole	Lanoxin S29		Livostin	
lvacaftor234	Lansoprazole		LMX4 FDI:	
Ivermectin68	Lantus	11	Locacorten-Viaform ED's	237

Local preparations for Anal and	Maxidex	238	Methylphenidate ER - Teva	144
Rectal Disorders	Maxitrol	238	Methylphenidate hydrochloride	144
Locasol263		250	Methylphenidate hydrochloride	
Locoid66, 72	Measles, mumps and rubella		extended-release	145
Locoid Crelo66		275	Methylprednisolone	81
Locoid Lipocream66	Mebendazole	91	Methylprednisolone (as sodium	
Locorten-Vioform237	Mebeverine hydrochloride	8	succinate)	81
Lodoxamide239			Methylprednisolone aceponate	66
Logem133	B Dermatological	67	Methylprednisolone acetate	82
Lomide239			Methylxanthines	234
Lomustine152	Medrol	81	Metoclopramide Actavis 10	136
Loniten59			Metoclopramide hydrochloride	136
Loperamide hydrochloride	Genito-Urinary	76	Metolazone	55
Lopinavir with ritonavir109		83–84	Metopirone	
Lopinavir/Ritonavir Mylan 109	Mefenamic acid	113	Metoprolol IV Mylan	52
Loprofin263	Megestrol acetate	170	Metoprolol succinate	52
Loprofin Mix262		142	Metoprolol tartrate	52
Lorafix229	Melphalan	152	Metrogyl	
Loratadine229	Menactra	275	Metronidazole	
Lorazepam140	Meningococcal (groups A, C, Y and	d	Metyrapone	90
Lorstat57	W-135) conjugate vaccine	275	Mexiletine hydrochloride	51
Losartan Actavis49	Meningococcal B multicomponent		Mexiletine Hydrochloride USP	51
Losartan potassium49	vaccine	276	Miacalcic	80
Losartan potassium with	Meningococcal C conjugate		Micolette	27
hydrochlorothiazide49			Miconazole	33
Lovir103	Menthol	65	Miconazole nitrate	
Loxamine13	Mepolizumab	199	Dermatological	65
Lucrin Depot 1-month89	Mercaptopurine	154	Genito-Urinary	77
Lucrin Depot 3-month89	Mercilon 28	75	Micreme	77
Lyderm70	Mesalazine	7	Micreme H	66
Lynparza159	Mesna	158	Microgynon 20 ED	76
Lyrica133	Mestinon	113	Microgynon 30	76
- M -	Metabolic Disorder Agents	27	Microgynon 50 ED	76
m-Eslon 128	Metformin hydrochloride	12	Microlut	76
Mabthera202	Metformin Mylan	12	Midazolam	142
Macrobid112	Methadone hydrochloride		Midazolam-Baxter	142
Macrogol 3350 with potassium	Extemporaneous	246	Midodrine	51
chloride, sodium bicarbonate and	Nervous	127	Mifegyne	79
sodium chloride27	Methatabs	127	Mifepristone	79
Macrogol 400 and propylene	Methenamine (hexamine)		Minerals	35
glycol241	hippurate		Mini-Wright AFS Low Range	235
Madopar 125122	Methopt	241	Mini-Wright Standard	235
Madopar 250122	Methotrexate	155	Minidiab	12
Madopar 62.5122	Methotrexate DBL Onco-Vial	155	MiniMed 1.8 Reservoir	
Madopar HBS122	Methotrexate Ebewe	155	MMT-326A	<mark>2</mark> 4
Madopar Rapid122	Methotrexate Sandoz	155	MiniMed 3.0 Reservoir	
Magnesium hydroxide37	Methyl hydroxybenzoate	246	MMT-332A	24
Magnesium sulphate37	Methylcellulose	246	MiniMed 640G	16
Mantoux279	Methylcellulose with glycerin and		MiniMed 770G	
Marevan45	sodium saccharin	246	MiniMed Mio MMT-921A	23
Marine Blue Lotion SPF 50+72	Methylcellulose with glycerin and		MiniMed Mio MMT-923A	23
Martindale Pharma243			MiniMed Mio MMT-925A	23
Marvelon 2875	Methyldopa	<u>55</u>	MiniMed Mio MMT-941A	23
Mask for spacer device235	Methyldopa Mylan	55	MiniMed Mio MMT-943A	23
Mast Cell Stabilisers234	Methyldopa Mylan S29	55	MiniMed Mio MMT-945A	
Maviret105			MiniMed Mio MMT-965A	23

MiniMed Mio MMT-975A	23	MSUD Maxamum	262	Neoral	22
MiniMed Quick-Set MMT-386A	23	Mucilaginous laxatives with		Neostigmine metilsulfate	113
MiniMed Quick-Set MMT-387A	23	stimulants	26	Nepafenac	23
MiniMed Quick-Set MMT-396A	23	Mucolytics	234	Nepro HP (strawberry)	25
MiniMed Quick-Set MMT-397A	23	Mucosoothe	124	Nepro HP (vanilla)	
MiniMed Quick-Set MMT-398A	23	Multiple Sclerosis Treatments	140	Nepro HP RTH	25
MiniMed Quick-Set MMT-399A	23	Multivitamin renal	34	Neulactil	
MiniMed Silhouette MMT-368A	23	Multivitamins	35	Neulastim	
MiniMed Silhouette MMT-377A	23	Mupirocin	64	NeuroTabs	3
MiniMed Silhouette MMT-378A	23	Muscle Relaxants		Nevirapine	
MiniMed Silhouette MMT-381A	23	Mvite	35	Nevirapine Alphapharm	
MiniMed Silhouette MMT-382A	23	Myambutol		Nicorandil	
MiniMed Silhouette MMT-383A	23	Mycobutin		Nicotine	14
MiniMed Silhouette MMT-384A	23	MycoNail		Nifedipine	
MiniMed Sure-T MMT-864A	21	Mycophenolate mofetil		Nifuran	
MiniMed Sure-T MMT-866A		Mydriacyl		Nilotinib	
MiniMed Sure-T MMT-874A		Mylan (12 hr release)		Nilstat	
MiniMed Sure-T MMT-876A		Mylan (24 hr release)		Alimentary	3
MiniMed Sure-T MMT-884A		Mylan Atenolol		Genito-Urinary	
MiniMed Sure-T MMT-886A		Mylan Clomiphen		Infection	
Minims Cyclopentolate		Mylan Indapamide		Nintedanib	
Minims Pilocarpine		Mylan Italy (24 hr release)		Nipent	
Minims Prednisolone		Mylan Midazolam		Nitrates	
Minirin		Myleran		Nitroderm TTS	
Minirin Melt		Myometrial and Vaginal Hormone		Nitrofurantoin	
Mino-tabs		Preparations		Nitrolingual Pump Spray	
				Nivestim	
Minocycline hydrochloride		Myozyme N -	21	Nivolumab	
Minor Skin Infactions		Nadolol	E0	Nizoral	
Minor Skin Infections Minoxidil				Nodia	
Mirena		Nadolol BNM		Noflam 250	
		Naglazyme Nalcrom			
Mirtazapine				Noflam 500	114
Misoprostol	450	Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	441
Mitomycin C		Naltraccord		Drugs	11
Mitozantrone		Naltrexone hydrochloride		Nonacog gamma, [Recombinant	4
Mitozantrone Ebewe		Naphazoline hydrochloride		Factor IX]	47
Mixtard 30		Naphcon Forte		Norethisterone	_
MMR II		Naprosyn SR 1000		Genito-Urinary	
Moclobemide		Naprosyn SR 750		Hormone	
Modafinil		Naproxen		Norflex	
Modavigil		Narcaricin mite		Norfloxacin	
Moduretic		Nasal Preparations		Noriday 28	
Molaxole		Natalizumab		Norimin	
Moments		Natulan		Normacol Plus	
Mometasone furoate		Nausafix		Normison	
Monogen		Nausicalm	136	Norpress	
Montelukast		Navelbine		Nortriptyline hydrochloride	
Montelukast Mylan	234	Nefopam hydrochloride	125	Norvir	
Moroctocog alfa [Recombinant fact		Neisvac-C		Noumed	
VIII]		Neo-B12		Noumed Paracetamol	
Morphine hydrochloride		Neo-Mercazole		NovaSource Renal	
Morphine sulphate		Neo-Mercazole S29	85	Novatretin	
Motetis	123	Neocate Gold		NovoMix 30 FlexPen	
Mouth and Throat	33	Neocate Junior Unflavoured	263	NovoRapid	1
Movapo		Neocate Junior Vanilla	263	NovoRapid FlexPen	
Moxifloxacin		Neocate SYNEO	263	NovoRapid Penfill	

INDEX: Generic Chemicals and Brands

NovoSeven RT	41	Olaparib	159	OxyNorm	12
Noxafil		Olbetam	56	Oxytocin	
Nozinan	137	Olbetam S29	56	Oxytocin BNM	
Nozinan (Swiss)	137	Olopatadine	242	Oxytocin with ergometrine	
Nucala		Olopatadine Teva		maleate	<mark>7</mark>
Nuelin		Olsalazine		Ozurdex	23
Nuelin-SR	234	Omalizumab	200	- P -	
Nupentin	132	Omeprazole	9	Pacifen	12
Nutilis	260	Omeprazole actavis 10	9	Pacimol	12
Nutren Diabetes	251	Omeprazole actavis 20		Paclitaxel	15
Nutrient Modules	248	Omeprazole actavis 40	9	Paclitaxel Actavis	15
Nutrini Energy Multi Fibre	253	Omnitrope		Paclitaxel Ebewe	
Nutrini Energy RTH	253	Onbrez Breezhaler	230	Paediatric Seravit	3
Nutrini Low Energy Multi Fibre		Oncaspar LYO	159	Palbociclib	16
Nutrini Peptisorb	265	OncoTICE		Paliperidone	13
Nutrini Peptisorb Energy	265	Ondansetron	136	Pamidronate disodium	
Nutrini RTH		Ondansetron ODT-DRLA	136	Pamisol	11
Nutrison 800 Complete Multi		One-Alpha	34	Panadol	12
Fibre	258	Onrex	136	Panadol Mini Caps	12
Nutrison Concentrated	260	Opdivo	221	Pancreatic enzyme	
Nutrison Energy	257	Ora-Blend	246	Pantoprazole	
Nutrison Energy Multi Fibre		Ora-Blend SF	246	Panzop Relief	
Nutrison Multi Fibre	258	Ora-Plus	246	Panzytrat	
Nutrison Standard RTH	258	Ora-Sweet	246	Papaverine hydrochloride	5
Nyefax Retard	54	Ora-Sweet SF	246	Para-amino salicylic acid	10
Nystatin		Orabase	33	Paracare	
Alimentary	33	Oral and Enteral Feeds	251	Paracare Double Strength	12
Genito-Urinary	77	Oratane	63	Paracetamol	
Infection	100	Ordine	127	Paracetamol + Codeine	
NZB Low Gluten Bread Mix	261	Orgran	261	(Relieve)	12
- 0 -		Ornidazole	101	Paracetamol Pharmacare	12
O/W Fatty Emulsion Cream	67	Orphenadrine citrate	121	Paracetamol with codeine	
Obinutuzumab	200	Ortho-tolidine	79	Paraffin	67–6
Obstetric Preparations	79	Oruvail SR	113	Paraffin liquid with wool fat	24
Ocrelizumab		Osmolite RTH		Parasidose	
Ocrevus	141	Other Endocrine Agents	90	Parasiticidal Preparations	6
Octocog alfa [Recombinant factor		Other Oestrogen Preparations	84	Parlodel	12
VIII] (Advate)	42	Other Progestogen		Parnate	13
Octocog alfa [Recombinant factor		Preparations	84	Parnate S29	13
VIII] (Kogenate FS)	42	Other Skin Preparations	72	Paromomycin	9
Octreotide	171	Ovestin		Paroxetine	13
Octreotide (Sun)	171	Genito-Urinary	77	Paser	10
Octreotide Depot Teva		Hormone		Paxam	14
Octreotide GH	171	Ox-Pam	140	Paxtine	13
Octreotide long-acting	171	Oxaliplatin	152	Pazopanib	16
Octreotide MaxRx	171	Oxaliplatin Accord	152	Peak flow meter	23
Oestradiol	83	Oxaliplatin Actavis 100	152	Pedialyte - Bubblegum	4
Oestradiol valerate	83	Oxaliplatin Ebewe	152	Pediasure	25
Oestradiol with norethisterone	84	Oxazepam		Pediasure RTH	25
Oestriol		Oxis Turbuhaler	230	Pegaspargase	15
Genito-Urinary	77	Oxpentifylline		Pegasys	
Hormone		Oxybutynin		Pegfilgrastim	4
Oestrogens	83	Oxycodone hydrochloride		Pegylated interferon alfa-2a	11
Ofev	232	Oxycodone Sandoz	129	Pembrolizumab	22
Oil in water emulsion	67	Oxycodone Sandoz S29		Pemetrexed	
Olanzapine13		OxyContin		Penicillamine	11

Penicillin G	95	PKU Lophlex LQ 10	262	Procarbazine hydrochloride	160
PenMix 30	11	PKU Lophlex LQ 20	262	Prochlorperazine	130
PenMix 40	11	PKU Lophlex Powder		Proctofoam	
PenMix 50	11	PKU Lophlex Sensation 20		Proctosedyl	
Pentasa	7	Plaquenil		Procyclidine hydrochloride	12
Pentostatin [Deoxycoformycin].	160	Plendil ER		Procytox	
Pentoxifylline [Oxpentifylline]	59	Pneumococcal (PCV10) conjug	ate	Progesterone	
Peptamen Junior	253	vaccine		Proglicem	10
Peptisorb		Pneumococcal (PCV13) conjug	ate	Proglycem	
Perhexiline maleate		vaccine		Progynova	
Pericyazine	138	Pneumococcal (PPV23)		Prolia	
Perindopril		polysaccharide vaccine	278	Promethazine hydrochloride	22
Perjeta		Pneumovax 23	278	Propafenone hydrochloride	5
Permethrin		Podophyllotoxin	72	Propamidine isethionate	
Perrigo	72	Polaramine	229	Propranolol	5
Pertuzumab	201	Poliomyelitis vaccine	278	Propylene glycol	
Peteha	102	Poloxamer	26	Propylthiouracil	
Pethidine hydrochloride	129	Poly-Gel		Prostacur	
Pevaryl		Poly-Tears		Protaphane	10
Pexsig		Poly-Visc		Protaphane Penfill	
Pfizer Exemestane		Polycal		Protifar	
Pharmacy Health Sorbolene wit		Ponstan		Protionamide	
Glycerin		Posaconazole		Provera	
Pheburane		Postinor-1		Provera HD	
Phenasen		Potassium chloride		PSM Citalopram	
Phenobarbitone		Potassium Chloride Aguettant		Psoriasis and Eczema	
Phenobarbitone sodium		Potassium citrate		Preparations	70
Extemporaneous	246	Potassium iodate		PTU	
Nervous		Povidone iodine		Pulmicort Turbuhaler	
Phenothrin		Pradaxa		Pulmozyme	
Phenoxybenzamine		Pramipexole hydrochloride		Puri-nethol	
hydrochloride	48	Pravastatin		Puria	
Phenoxymethylpenicillin (Penici		Pravastatin Mylan		Pyrazinamide	
V)		Praziquantel		Pyridostigmine bromide	
Phenytoin sodium		Prazosin		Pyridoxine hydrochloride	
Phillips Milk of Magnesia		Pred Forte		Pyrimethamine	
Phlexy 10		Prednisolone		Pytazen SR	
Phosphate Phebra		Prednisolone acetate		- Q -	
Phosphorus		Prednisolone sodium		Quetapel	138
Phytomenadione		Prednisolone sodium	•	Quetiapine	
Pilocarpine hydrochloride		phosphate	239	Quick-Set MMT-392	2
Pimafucort		Prednisolone-AFT		Quick-Set MMT-393	
Pimecrolimus		Prednisone		Quinapril	
Pindolol		Pregabalin		Quinapril with	
Pine tar with trolamine laurilsulfa		Pregabalin Pfizer		hydrochlorothiazide	49
and fluorescein		Pregnancy Tests - hCG Urine		Qvar	
Pinetarsol		Premarin		- R -	
Pioglitazone		Prevenar 13		RA-Morph	12
Pirfenidone		Priadel		Raloxifene hydrochloride	
Pizotifen		Primaguine		Raltegravir potassium	109
PKU Anamix Infant		Primidone		Ramipex	12
PKU Anamix Junior		Primolut N		Ranbaxy-Cefaclor	
PKU Anamix Junior Chocolate.		Priorix		Ranbaxy-Cefaclor S29	
PKU Anamix Junior LQ		Probenecid		Rapamune	
PKU Anamix Junior Orange		Probenecid-AFT		Rasagiline	
PKU Anamix Junior Vanilla		Procaine penicillin		Reandron 1000	

INDEX: Generic Chemicals and Brands

Recombinant factor IX	39, 42	Roxane	6	Simvastatin	5
Recombinant factor VIIa	41	Roxane-Propranolol	<u>53</u>	Simvastatin Mylan	
Recombinant factor VIII	41–42	Roxithromycin	94	Sinemet	122
Rectogesic	8	Rubifen	144	Sinemet CR	122
Redipred		Rubifen SR	144	Sirolimus	22
Relieve		Rugby Capsaicin Topical		Siterone	82
Relistor	26	Cream	125	Slow-Lopresor	52
Remicade	192	Rulide D		Smith BioMed Rapid Pregnancy	
Renilon 7.5	254	Rurioctocog alfa pegol [Reco	mbinant	Test	78
Resonium-A	47	factor VIII]		Sodibic	4
Resource Beneprotein	250	Ruxolitinib		Sodium acid phosphate	2
Respigen		Rythmodan		Sodium alginate	
Respiratory Devices		Rytmonorm		Sodium benzoate	
Respiratory Stimulants		· -S-		Sodium bicarbonate	
Retinol palmitate		Sabril	134	Blood	46–47
ReTrieve	63	Sacubitril with valsartan		Extemporaneous	
Retrovir		Sagent		Sodium calcium edetate	
Revlimid		SalAir		Sodium chloride	
Revolade		Salazopyrin		Blood	46
Rexacrom		Salazopyrin EN		Respiratory	
Riboflavin		Salbutamol		Sodium citrate with sodium lauryl	200
Ribomustin		Salbutamol with ipratropium	201	sulphoacetate	2
Ricit		bromide	221	Sodium citro-tartrate	
Rifabutin		Salicylic acid		Sodium cromoglicate	/ 3
		Salmeterol			
Rifadin		Sandomigran		Alimentary	
		· ·		Respiratory	
Rifaximin		Sandostatin LAR		Sensory	
Rifinah		Sanofi Primaquine		Sodium fluoride	J:
Rilutek		Sapropterin dihydrochloride		Sodium Fusidate [fusidic acid]	•
Riluzole		Scalp Preparations		Dermatological	
RINVOQ		Scopoderm TTS		Infection	
Riodine		Sebizole		Sensory	23
Risedronate Sandoz		Secukinumab		Sodium hyaluronate [Hyaluronic	
Risedronate sodium		Sedatives and Hypnotics		acid]	
Risperdal Consta		Seebri Breezhaler		Sodium phenylbutyrate	
Risperidone		Selegiline hydrochloride		Sodium picosulfate	
Risperidone (Teva)		Senna		Sodium polystyrene sulphonate	
Risperon		Senokot		Sodium tetradecyl sulphate	
Ritalin		Sensipar		Sodium valproate	
Ritalin LA		SensoCard		Sofradex	
Ritonavir		Serenace		Soframycin	
Rituximab (Mabthera)		Seretide		Solgar	
Rituximab (Riximyo)	204	Seretide Accuhaler	231	Solifenacin Mylan	
Rivaroxaban	45	Serevent	<mark>230</mark>	Solifenacin succinate	79
Rivastigmine	146	Serevent Accuhaler		Solu-Cortef	8
Rivastigmine Patch BNM 10	146	Sertraline	131	Solu-Medrol	
Rivastigmine Patch BNM 5	146	Setrona	131	Solu-Medrol-Act-O-Vial	
Rivotril	131–132	Sevredol	128	Somatropin (Omnitrope)	8
Riximyo	204	Sex Hormones Non		Sotalol	5
RIXUBIS		Contraceptive		Spacer device	236
Rizamelt	135	Shield XL	74	Span-K	4
Rizatriptan	135	shingles vaccine		Spiolto Respimat	232
Ropin		SII-Onco-BCG		Spiractin	5
Ropinirole hydrochloride		Sildenafil		Spiriva	
Rotarix	278	Silhouette MMT-373		Spiriva Respimat	
Rotavirus oral vaccine		Siltuximab		Spironolactone	
	-	***************************************	-	,	

Sporanox	90	Tamsulosin hydrochloride	78	Tobramycin BNM	QS
Sprycel		Tamsulosin-Rex		Tobramycin Mylan	
Staphlex		Tandem Cartridge		Tobrex	
Stemetil		Tandem t:slim X2 with Basal-IQ.		Tocilizumab	
SteroClear				Tofranil	
		Tap water			
Stesolid		Tarceva		Tolcapone	
Stimulants/ADHD Treatments		Tasigna		Topamax	134
Stiripentol		Tasmar		Topical Products for Joint and	
Stocrin		Taurine		Muscular Pain	
Stomahesive		Tecfidera		Topiramate	
Strattera		Tegretol		Topiramate Actavis	
Strides Shasun		Tegretol CR		Total parenteral nutrition (TPN)	
Stromectol		Telfast		TPN	
Sucralfate		Teligent		Tramadol hydrochloride	
Sulfadiazine Silver		Temaccord		Tramal SR 100	
Sulfadiazine sodium		Temazepam		Tramal SR 150	
Sulfasalazine		Temozolomide		Tramal SR 200	
Sulindac	114	Tenofovir disoproxil	103	Trandate	52
Sulindac Mylan	114	Tenofovir Disoproxil Teva	103	Tranexamic acid	
Sulphur	71	Tenoxicam	114	Tranylcypromine sulphate	130
Sulprix	136	Tensipine MR10	54	Trastuzumab	219
Sumagran	135	Tepadina	152	Trastuzumab emtansine	221
Sumatriptan		Terbinafine	100	Travatan	240
Sunitinib		Terbutaline sulphate	231	Travoprost	240
Sunscreens	72	Teriflunomide		Travopt	
Sunscreens, proprietary		Teriparatide		Treatments for Dementia	
Sure-T MMT-863		Testosterone		Treatments for Substance	
Sure-T MMT-873		Testosterone cipionate		Dependence	146
Sustagen Diabetic		Testosterone esters		Trental 400	
Sustagen Hospital Formula		Testosterone undecanoate		Tretinoin	
Active	258	Tetrabenazine		Dermatological	63
Sustanon Ampoules		Tetrabromophenol		Oncology	
Sutent		Tetracosactrin		Trexate	
Sylvant		Tetracycline		Triamcinolone acetonide	100
Symbicort Turbuhaler 100/6		Thalidomide		Alimentary	31
Symbicort Turbuhaler 200/6		Thalomid		Dermatological	66
Symbicort Turbuhaler 400/12		Theophylline		Hormone	
Symmetrel		Thiamine hydrochloride		Triamcinolone acetonide with	02
Sympathomimetics		THIO-TEPA		gramicidin, neomycin and nysta	tin
Synacthen		Thioguanine		Dermatological	
		Thiotepa		Sensory	
Synacthen Depot				Triazolam	
Synacthene Retard		Thymol glycerin			
Synflorix		Thyroid and Antithyroid Agents		Trimethoprim	90
Synthroid		Ticagrelor		Trimethoprim with	
Syntometrine		Tilcotil		sulphamethoxazole	00
Syrup (pharmaceutical grade)		Tillomed		[Co-trimoxazole]	
Systane Unit Dose	241	Timolol		Trisequens	
-Т-		Timoptol XE	239	Trisul	
Tacrolimus		Tiotropium bromide	232	Trophic Hormones	85
Dermatological		Tiotropium bromide with		Tropicamide	
Oncology		olodaterol		Trulicity	
Tacrolimus Sandoz		Tivicay		Trusopt	
Taliglucerase alfa		TMP	98	TruSteel	
Tambocor		Tobramycin		Tuberculin PPD [Mantoux] test	
Tamoxifen citrate		Infection		Tubersol	
Tamoxifen Sandoz	172	Sensory	238	Two Cal HN	260

INDEX: Generic Chemicals and Brands

Two Cal HN RTH	260	Viaderm KC	67	Ziagen	108
Tykerb		Vidaza		Zidovudine [AZT]	
Tysabri	141	Vigabatrin	134	Zidovudine [AZT] with	
- U -		Vildagliptin	12	lamivudine	108
Ultibro Breezhaler	232	Vildagliptin with metformin		Zimybe	57
Ultraproct	8	hydrochloride	12	Zinc and castor oil	67
Umeclidinium		Vimpat		Zinc sulphate	37
Umeclidinium with vilanterol	232	Vinblastine sulphate	162	Zincaps	
Univent	231, 235	Vincristine sulphate	162	Zinnat	92
Upadacitinib	226	Vinorelbine		Ziprasidone	138
Ural		Vinorelbine Ebewe	163	Zista	
Urea		Viramune Suspension	108	Zithromax	
Urex Forte	55	ViruPOS		Zoledronic acid	
Urinary Agents	78	Vit.D3	34	Hormone	80
Urinary Tract Infections		Vita-B12	34	Musculoskeletal	117
Urinorm		VitA-POS	242	Zoledronic acid Mylan	80
Uromitexan		Vitabdeck	35	Zopiclone	
Ursodeoxycholic acid		Vital	254	Zopiclone Actavis	
Ursosan		Vitamin B complex		Zostavax	
Utrogestan		Vitamin B6 25		Zostrix	
- V -		Vitamins		Zostrix HP	
Vaccinations	268	Vivonex TEN		Zuclopenthixol decanoate	
Vaclovir		Voltaren		Zuclopenthixol hydrochloride	
Valaciclovir		Voltaren D		Zusdone	
Valganciclovir		Voltaren Ophtha		Zyban	
Valganciclovir Mylan		Voltaren SR		Zypine	
Vancomycin		Volumatic		Zypine ODT	
Vannair		Voriconazole		Zyprexa Relprevv	
Varenicline Pfizer		Vosol		Zytiga	
Varenicline tartrate		Votrient			
Varicella vaccine [Chickenpox		Vttack			
vaccine]		- W -			
Varicella zoster virus (Oka stra		Warfarin sodium	45		
attenuated vaccine [shingle		Wart Preparations			
vaccine]		Wasp venom allergy treatment			
Varivax		Water			
Vasodilators		Blood	47		
Vasopressin Agonists		Extemporaneous			
Vasorex		Wool fat with mineral oil			
Vedafil		- X -			
Veletri		Xarelto	45		
Venclexta		Xifaxan			
Venetoclax		XMET Maxamum			
Venlafaxine		Xolair			
Venomil		XP Maxamum			
VENOX		Xylocaine			
Ventavis		Xylocaine 2% Jelly			
Ventolin		Xyntha			
Venesid		- Z -	41		
		-	//0		
Verapamil hydrochloride		Zapril			
Vergo 16		Zarontin			
Vermox Versacloz		Zaroxolyn			
Vesanoid		ZavedosZeffix			
Vexazone		Zematop			
V EXALUITE	12	Zematop	/1		