November 2021 Volume 28 Number 2

Editors:

Kaye Wilson, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at no cost from the Pharmac website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
@Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

		_
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	63
	Genito Urinary System	74
	Hormone Preparations – Systemic	81
	Infections – Agents For Systemic Use	92
	Musculoskeletal System	114
	Nervous System	123
	Oncology Agents & Immunosuppressants	152
	Respiratory System & Allergies	230
	Sensory Organs	239
	Various	245
Section C	Extemporaneous Compounds (ECPs)	247
Section D	Special Foods	250
Section I	National Immunisation Schedule	270
	ladau	000

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 n sachet	0.	30	✓ Ga	viscon Infant
GODIUM ALGINATE ★ Tab 500 mg with sodium bicarbonate 267 mg and calciur carbonate 160 mg - peppermint flavour		60		viscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and cal carbonate 160 mg per 10 ml		500 ml	Ac	idex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE		100	✓ Alı	u-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) Subsidy by endorsement Only when prescribed for patients unable to swallow inappropriate and the prescription is endorsed accor	39.00	500 ml ts or whe	✓ Ro re calcium	
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available Tab 2 mg Cap 2 mg	10.75	400 400	✓ No ✓ <u>Dia</u>	dia amide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Reta pharmacy	166.50	90 alid for 6		tocort CIR r applications meeting

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy		Fully	Brand or
(Manufacturer's Price)) :	Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)	26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROC	HLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%	26.55	10 g OP	✓ Proctofoam S29
MESALAZINE			
Tab 400 mg	49.50	100	✓ Asacol
Tab EC 500 mg		100	✓ Asamax
Tab long-acting 500 mg	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g	118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml	41.30	7	✓ Pentasa
Suppos 500 mg		20	✓ Asacol
Suppos 1 g	50.96	28	✓ Pentasa
(Asamax Tab EC 500 mg to be delisted 1 March 2022)			
OLSALAZINE			
Tab 500 mg	93.37	100	✓ Dipentum
Cap 250 mg	53.00	100	✓ Dipentum

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	/	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	•	Nalcrom
SULFASALAZINE * Tab 500 mg Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE			
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g	30 g OP	✓ Ultraproct	
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg2.66	12	✓ Ultraproct	
HYDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00 Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	30 g OP 12	✓ Proctosedyl ✓ Proctosedyl	

Management of Anal Fissures

GLYCERYL TRINITRATE − Special Authority see SA1329 below − Retail pharmacy

★ Oint 0.2%......22.00 30 g OP

✓ Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE * Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE * Tab 135 mg	0.20	90	✓ Colofae
* Tab 100 IIIg	5.20	90	• Cololac

Antiulcerants

Antisecretory and Cytoprotective

MIS	SOPROSTOL STORE ST		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	Cytotec

	ALIMENTALLI	THAVE AND	DIMETABOLISM
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per 🗸	Brand or Generic Manufacturer
Helicobacter Pylori Eradication			

CLARITHROMYCIN

14 ✓ Apo-Clarithromycin ✓ Klacid

- a) Maximum of 28 tab per prescription
- b) Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly. Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.
- c) Klacid to be Sole Supply on 1 February 2022

(Apo-Clarithromycin Tab 500 mg to be delisted 1 February 2022)

H2 Antagonists

FA	MOTIDINE - Only on a prescription			
	Tab 20 mg	4.91	100	✓ Famotidine
	•			Hovid S29
*	Tab 40 mg	8.48	100	✓ Famotidine
	-			Hovid S29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement	57.02	10	✓ Mylan S29
	Subsidy by endorsement - Subsidised for patients receiving to	reatment as par	t of palliative	e care.

Proton Pump Inhi	bitors
------------------	--------

LANSOPRAZOLE			
* Cap 15 mg	4.20	100	Lanzol Relief
Lanzol Relief to be Principal Supply on 1 December 2021			
* Cap 30 mg	5.26	100	Lanzol Relief
Lanzol Relief to be Principal Supply on 1 December 2021			
OMEPRAZOLE			
For omeprazole suspension refer Standard Formulae, page 24	7		
* Cap 10 mg	1.94	90	✓ Omeprazole actavis 10
* Cap 20 mg	1.86	90	✓ Omeprazole actavis 20
* Cap 40 mg	3.11	90	✓ Omeprazole actavis 40
* Powder – Only in combination	42.50	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspe	ension.	-	
* Inj 40 mg ampoule with diluent	33.98	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
PANTOPRAZOLE			
* Tab EC 20 mg	2.02	100	✓ Panzop Relief
* Tab EC 40 mg		100	✓ Panzop Relief

Site Protective Agents

COLLOIDAL BISMUTH SUBCITRATE

✓ Gastrodenol S29 Tab 120 mg14.51 50

	Subsidy (Manufacturer's Price \$	e) Si Per	Fully ubsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g	35.50 (48.28)	120	(Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail Tab 550 mg		56	✓)	(ifaxan
■ SA1461 Special Authority for Subsidy initial application only from a gastroenterologist, hepatolog nepatologist. Approvals valid for 6 months where the patier olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Pra nepatologist. Approvals valid without further renewal unless benefiting from treatment.	gist or Practitioner on the nt has hepatic encephalo actitioner on the recomme	pathy des endation o	pite an a	dequate trial of maximur penterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE – Special Authority see SA1320 below – Retai Cap 25 mg Cap 100 mg Oral liq 50 mg per ml >>SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approval nypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO			For the tre	
	52.00	'	• •	<u>alucagen riypokit</u>
Insulin - Short-acting Preparations				
NSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml OP		Actrapid
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓ F	Humulin R Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
NSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen NSULIN ISOPHANE	52.15	5	✓ N	NovoMix 30 FlexPen
▲ Inj human 100 u per ml	17.68	10 ml OP	-	Humulin NPH Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓	Humulin NPH Protaphane Penfill

	Subsidy		Fully Brand or
	(Manufacturer's F	Price) Subs Per	idised Generic Manufacturer
SULIN ISOPHANE WITH INSULIN NEUTRAL	Ψ		That land to the l
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
nij naman war neatar maaiin 100 a per mi	25.20	10 1111 01	✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30
			✓ PenMix 40
			✓ PenMix 50
SULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,			
3 ml		5	✓ Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,			· ·
3 ml		5	✓ Humalog Mix 50
noulin Long acting Proporations			
nsulin - Long-acting Preparations			
SULIN GLARGINE			
Inj 100 u per ml, 10 ml		1	✓ Lantus
Inj 100 u per ml, 3 ml		5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓ Lantus SoloStar
nsulin - Rapid Acting Preparations			
SULIN ASPART			
Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	✓ NovoRapid FlexPen
SULIN GLULISINE			
Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen		5	✓ Apidra SoloStar
SULIN LISPRO			•
Inj 100 u per ml, 10 ml	34 92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml		5	✓ Humalog
· ·			-
Alpha Glucosidase Inhibitors			
CARBOSE			
Tab 50 mg		90	✓ Glucobay
	8.95		✓ Accarb
Accarb to be Principal Supply on 1 December 2021	2.42	00	(Observe)
Tab 100 mg		90	✓ Glucobay
Accarb to be Principal Supply on 1 December 2021	15.29		✓ Accarb
Alucobay Tab 50 mg to be delisted 1 December 2021) Alucobay Tab 100 mg to be delisted 1 December 2021)			
,			
Oral Hypoglycaemic Agents			
LIBENCLAMIDE			
Tab 5 mg	7.50	100	✓ Daonil

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
GLICLAZIDE				
* Tab 80 mg	15.18	500	1	Glizide
GLIPIZIDE				
* Tab 5 mg	4.58	100	•	Minidiab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg	8.63	1,000	/	Apotex
·	14.74		✓	Metformin Mylan
* Tab immediate-release 850 mg	7.04	500		Apotex
	11.28		✓	Metformin Mylan
(Apotex Tab immediate-release 850 mg to be delisted 1 March 2 PIOGLITAZONE	,	00	./	Veverene
* Tab 15 mg Vexazone to be Principal Supply on 1 January 2022	6.80	90	•	Vexazone
* Tab 30 mg	7.30	90	1	Vexazone
Vexazone to be Principal Supply on 1 January 2022				
* Tab 45 mg Vexazone to be Principal Supply on 1 January 2022	12.25	90	✓	Vexazone
,				
VILDAGLIPTIN Tab 50 mg	35.00	60	1	Galvus
		00	•	Gaivus
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE	05.00	00	,	Calminat
Tab 50 mg with 1,000 mg metformin hydrochloride		60		Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	•	Galvumet

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*: or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

 a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack,

Subsidy		Fully	Brand or
(Manufacturer's Price	,	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.

b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*: or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 above - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58.56	30	Jardiance
*	Tab 25 mg58.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see SA2068 above – Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

	Trotor Trot to be given in combination that a famaca dis. I age inch			
*	Tab 5 mg with 1,000 mg metformin hydrochloride	.58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	.58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	.58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	.58.56	60	Jardiamet

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips ______20.00 1 OP

CareSens Dual

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

20.00

✓ CareSens N Premier

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRC

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood g	lucose test stri	os26.20	50 test OP	✓ SensoCard
---------	------------------	---------	------------	-------------

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES – Maximum	of 200 deviper prescription
-------------------------------	-----------------------------

* * *	31 g × 6 mm	11.75 9.50 10.50 10.50	100 100 100 100 100	111	B-D Micro-Fine B-D Micro-Fine Berpu B-D Micro-Fine B-D Micro-Fine
INS	ULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE				
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.00	100	/	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.00	100	1	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.00	100	1	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.00	100	1	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.00	100	1	B-D Ultra Fine
		1.30	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.00	100	1	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

 c) Maximum of 1 insulin pump per patient each four year 	period.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 640G
			✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

(MiniMed 640G Min basal rate 0.025 U/h to be delisted 1 January 2022)

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	F	ılly	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment: and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy: and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Fither
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Fither:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump

Subsidy (Manufacturer's Price) Subsidised Subsidised Generic Manufacturer continued treatment; and 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 5 The patient is continuing to derive benefit from pump therapy; and 5 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 3 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 3 sets per prescription c) Maximum of 3 sets per p		ALIMENTAR	Y TR	ACT AND	METABOLISM
treatment; and 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient has adhered to and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 33 sets per prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		(Manufacturer's Price)		Subsidised	Generic
3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and 4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.3 Applicant is a relevant specialist; or 4.4 Applicant is a relevant specialist; or 4.5 Applicant is a relevant specialist; or 4.6 Applicant is a relevant specialist; or 4.7 Applicant is a relevant specialist; or 4.8 Applicant is a relevant specialist; or 4.9 Maximum of 3 sets per prescription 4 Donly on a prescription 5 Only on a prescription 6 Only on a prescription 7 Only on a prescription 8 Only on a prescription 9 Only on a prescription 10 Only on a prescription 11 Only on a prescription 12 Maximum of 13 infusion sets will be funded per year. 13 Only on a prescription 14 Only on a prescription 15 Only on a prescription 16 Maximum of 13 infusion sets will be funded per year. 17 Only on a prescription 18 MiniMed Sure-T 19 MiniMed Sure-T					
pump therapy; and 4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10	•				
4 The patient is continuing to derive benefit from pump therapy; and 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient as not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		ising analogue insulins	for at	least six mo	nths prior to initiating
5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tilock × 10	1 1 127	aranu and			
6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tilock × 10			80 mm	nal/mal an ni	ımn therany: and
7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and 8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10					
8 Either: 8.1 Applicant is a relevant specialist; or 8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE – Special Authority see SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10					
8.2 Applicant is a nurse practitioner working within their vocational scope. Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	•		,		
Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	8.1 Applicant is a relevant specialist; or				
years for applications meeting the following criteria: All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, tlock × 10		•			
All of the following: 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		from a relevant speciali	st or n	urse practiti	oner. Approvals valid for 2
1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10					
than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	3		l	ata ta ta a at a	III. Ada at annial ta an lasa
2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		tne treatment plan and	nas m	iaintained a	HDA1c of equal to or less
3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and 4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	,	mmol/mol from initial ar	nlicati	on and	
4 Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	•		•		ne: and
4.2 Applicant is a nurse practitioner working within their vocational scope. INSULIN PUMP CARTRIDGE − Special Authority see SA1985 on page 19 − Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10		, , , , , , , , , , , , , , , , , , ,			,
INSULIN PUMP CARTRIDGE — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	4.1 Applicant is a relevant specialist; or				
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	4.2 Applicant is a nurse practitioner working within t	heir vocational scope.			
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10	INSULIN PUMP CARTRIDGE - Special Authority see SA198	5 on page 19 – Retail p	harma	acv	
c) Maximum of 13 packs of cartridge sets will be funded per year. Cartridge 300 U, t:lock × 10				,	
Cartridge 300 U, t:lock × 10	b) Only on a prescription				
INSULIN PUMP INFUSION SET (STEEL CANNULA) — Special Authority see SA1985 on page 19 — Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10					
a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	Cartridge 300 U, t:lock × 10	50.00	1 OP	✓ T	andem Cartridge
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	, , ,	al Authority see SA198	5 on p	age 19 – Re	etail pharmacy
c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing × 10	, , , , ,				
10 mm steel needle; 60 cm tubing × 10	, , , ,				
MMT-884A 10 mm steel needle; 80 cm tubing × 10130.00 1 OP ✓ MiniMed Sure-T		120.00	1 OD		liniMad Sura-T
10 mm steel needle; 80 cm tubing × 10130.00 1 OP ✓ MiniMed Sure-T	To min steer needle, oo chi tubing x 10	130.00	I OF	▼ IV	
·	10 mm steel needle: 80 cm tubing × 10	130.00	1 OP	✓ N	
mini ooon	3 · · · · · · · · · · · · · · · · · · ·			-	MMT-886A
6 mm steel needle; 60 cm tubing × 10	6 mm steel needle; 60 cm tubing x 10	130.00	1 OP	✓ N	liniMed Sure-T

MMT-864A

MMT-866A

MMT-874A

✓ MiniMed Sure-T

✓ MiniMed Sure-T

✓ MiniMed Sure-T

MMT-876A

✓ Sure-T MMT-863

✓ Sure-T MMT-873

1 OP

1 OP

1 OP

1 OP

1 OP

8 mm steel needle; 60 cm tubing × 10130.00

6 mm steel needle: 29 G: manual insertion: 60 cm tubing x

8 mm steel needle; 29 G; manual insertion; 60 cm tubing x

Subsidy	H	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	✓	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles		1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line × 10 with	130 00	1 OP	✓ TruSteel

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing × 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm clear tubing × 10130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
 ✓ MiniMed Silhouette
- MMT-381A

 MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
 10 P

 6 mm teflon needle, 45 cm blue tubing × 10
 130.00
 1 OP
- ✓ MiniMed Quick-Set MMT-398A
 ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
 - ✓ MiniMed Mio MMT-943A
 - ✓ MiniMed Mio MMT-923A
 - ✓ MiniMed Quick-Set MMT-399A
 - ✓ MiniMed Mio MMT-945A
 - ✓ MiniMed Mio MMT-965A
 - ✓ MiniMed Mio MMT-925A
 - ✓ MiniMed Quick-Set MMT-387A
 - ✓ MiniMed Quick-Set MMT-396A
 - ✓ MiniMed Quick-Set MMT-397A
 - ✓ MiniMed Mio MMT-975A
 - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH II	NSERTION	I DEVICE)	- Special Authority see
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	cm 140.00	1 OP	√ Διι	toSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cl	n	1 OP		toSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with	NSERTION) - Spe			
10 needles; luer lock	IT INSERTION WIT	1 OP H INSERT		houette MMT-373 ICE) – Special Authority
110 cm line × 10 with 10 needles	m	1 OP		toSoft 90
line x 10 with 10 needles		1 OP		toSoft 90 toSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m	1 OP		toSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy a) Maximum of 3 sets per prescription		-		
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock	130.00	1 OP	√ Qu	ick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock		1 OP	√ Qu	ick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 c a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per	n page 19 – Retail _I	oharmacy	-	
10 × luer lock conversion cartridges 1.8 ml for Paradigm pun Cartridge for 5 and 7 series pump; 1.8 ml × 10	ps50.00	1 OP 1 OP	✔ Mir 1	R Cartridge 1.8 niMed .8 Reservoir //MT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		niMed 8.0 Reservoir //MT-332A

Fully

Brand or

	(Manufacturer's Price) Sub Per	osidised	Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					_
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Cr	eon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))	*	100	✓ Pa	nzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Cr	eon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph		00 00			
Eur U)		20 g OP	✓ Cr	eon Micro	
URSODEOXYCHOLIC ACID – Special Authority see SA1739 be	•	•			
Cap 250 mg	32.95	100	✓ <u>Ur</u> s	sosan	

Subsidy

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- ner:
 - 1 Patient has been diagnosed with Alagille syndrome; or
 - 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6

Su	Subsidy	Fully	Brand or
(Manufac	cturer's Price) Subsid	dised	Generic
	\$ Per	•	Manufacturer

continued...

months where the patient continues to benefit from treatment.

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

* Powder for oral soln	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
•	(17.32)	-	Normacol Plus
	2.41	200 g OP	
	(8.72)	-	Normacol Plus

Faecal Softeners

*	Tab 50 mg	.2.31	100	✓ Coloxyl
	Tab 120 mg		100	✓ Coloxyl
	CUSATE SODIUM WITH SENNOSIDES			
*	Tab 50 mg with sennosides 8 mg	.3.10	200	✓ Laxsol
	OXAMER – Only on a prescription			
	Not funded for use in the ear.			
*	Oral drops 10%	.3.98	30 ml OP	Coloxyl

Opioid Receptor Antagonists - Peripheral

DOCUSATE SODIUM - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	see SA1691 below – Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
,	246.00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Fither:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

	Subsidy (Manufacturer's Price \$	e) Sul Per	Fully osidised	Brand or Generic Manufacturer
Osmotic Laxatives				
GLYCEROL * Suppos 3.6 g – Only on a prescription LACTULOSE – Only on a prescription	9.25	20	√ P	SM
* Oral liq 10 g per 15 ml	3.33	500 ml	√ L	aevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIG	CARBONATE AND	SODIUM	CHLORII	DE
Powder for oral soln 13.125 g with potassium chloride 46.6 m sodium bicarbonate 178.5 mg and sodium chloride 350.7	O,	30	✓ <u>N</u>	<u>lolaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	√ F	leet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,	, ,	•		Parlana
5 ml	29.98	50	✓ <u>IV</u>	<u>licolette</u>
Stimulant Laxatives				
BISACODYL — Only on a prescription * Tab 5 mg * Suppos 10 mg Lax-Suppositories to be Principal Supply on 1 December	3.69	200 10	_	ax-Tab ax-Suppositories
SENNA – Only on a prescription * Tab, standardised	2.17 (8.21)	100	S	enokot
	0.43 (2.06)	20	Q	enokot
SODIUM PICOSULFATE - Special Authority see SA2053 below	` '		3	GIIOROL
Oral soln 7.5 mg per ml		30 ml OP	✓ D	ulcolax SP Drop

SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE	 Special Authority 	see SA2042 held	ow – Retail pharmacy

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 below - Retail pharmacy

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

- 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
- 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

Cap 120 mg	CBS	30	Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts; or

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - F	Retail pharmacy		
Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral lig 500 mg per 10 ml		300 ml	✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

	Subsidy (Manufacturer's Price)		-ully ised	Brand or Generic	
	\$	Per	1	Manufacturer	
SAPROPTERIN DIHYDROCHLORIDE - Special Authority see S	A1989 below – Reta	il pharmacy			
Tab soluble 100 mg	1,452.70	30 OP	1	Kuvan	

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - R	letail pharmacy		
Soln 100 mg per ml	CBS	100 ml	✓ Amzoate S29

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE − Special Authority see SA1990 below − Retail pharmacy
Grans 483 mg per g......2,016.00 174 g OP

✓ Pheburane

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 on the next page	 Retail pharmacy 		
Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy

✓ Elelvso

⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 Pharmac PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, 6) thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or massive symptomatic splenomegaly; or
 - 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g., hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
 - 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher
 - 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated: and
- Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 7) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with

BENZYDAMINE HYDROCHLORIDE

Endorsement		500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly.	s oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)	_	Orabase
	1.52	5 g OP	
	(3.60)	•	Orabase
Powder	8.48	28 g OP	
	(10.95)	_	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
7. Allouvo goro /o mar obtainorilain cinoriae c.e. //o	(6.00)	10 g 01	Bonjela
TRIAMCINOLONE ACETONIDE	(0.00)		201,014
Paste 0.1%	E 22	F a OB	✓ Kanalag in Orahaga
Fasie 0.176	3.33	5 g OP	 Kenalog in Orabase
Oropharyngeal Anti-infectives			
Oropharyngear Anti-Infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ Decozol
Decozol to be Principal Supply on 1 December 2021		. o g o .	20020.
NYSTATIN			
****	1 76	24 ml OP	✓ Nilstat
Oral liq 100,000 u per ml	1.70	24 IIII UP	▼ INIISIAL

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's l \$	Price) Sub Per	Fully Brand or sidised Generic ✓ Manufacturer
Other Oral Agents		
For folinic mouthwash, pilocarpine oral liquid or saliva substitute formula refer Sta FHYMOL GLYCERIN	andard Formul	ae, page 247
Compound, BPC9.15 PSM Compound, BPC to be delisted 1 February 2022)	500 ml	✓ PSM
Vitamins		
Vitamin B		
HYDROXOCOBALAMIN		€ No Bd0
* Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO1.89	3	✓ Neo-B12 ✓ Vita-B12
3.15	5	✓ Hydroxocobalamin Mercury Pharma
PYRIDOXINE HYDROCHLORIDE		
a) No more than 100 mg per dose b) Only on a prescription		
* Tab 25 mg - No patient co-payment payable2.70	90	✓ Vitamin B6 25
Tab 50 mg	500	✓ Apo-Pyridoxine
23.45		✓ Pyridoxine multichem
(Apo-Pyridoxine Tab 50 mg to be delisted 1 May 2022)		
THIAMINE HYDROCHLORIDE - Only on a prescription * Tab 50 mg7.09	100	✓ Max Health
VITAMIN B COMPLEX	100	▼ Wax Health
* Tab, strong, BPC7.15	500	✓ Bplex
Vitamin C		
ASCORBIC ACID		
a) No more than 100 mg per dose		
b) Only on a prescription * Tab 100 mg9.90	500	✓ Cvite
Vitamin D	000	- <u> </u>
vitamin D		
ALFACALCIDOL	400	Comp Alerte
* Cap 0.25 mcg	100 100	✓ One-Alpha✓ One-Alpha
* Oral drops 2 mcg per ml 60.68	20 ml OP	✓ One-Alpha
CALCITRIOL		•
* Cap 0.25 mcg7.95	100	✓ Calcitriol-AFT
* Cap 0.5 mcg	100	✓ Calcitriol-AFT
COLECALCIFEROL	40	/ VIII DO

* Cap 1.25 mg (50,000 iu) - Maximum of 12 cap per prescription......2.95

✓ Vit.D3

✓ Puria

12

4.8 ml OP

ALIMENTARY TRACT AND METABOLISM Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Multivitamin Preparations** MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pharmacy 30 ✓ Clinicians Renal Vit ⇒SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither: 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA). MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy 200 g OP ✓ Paediatric Seravit ⇒SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins. **VITAMINS** 1.000 Mvite * Cap (fat soluble vitamins A, D, E, K) - Special Authority see 60 Vitabdeck ⇒SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut syndrome; or 3 Patient has severe malabsorption syndrome. **Minerals** Calcium **CALCIUM CARBONATE** ✓ Calci-Tab 500 250 * Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement.......52.00 20 ✓ Calcium-Sandoz Forte S29 ✓ Cacit \$29 54.60 76 Subsidy by endorsement - Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable. CALCIUM GLUCONATE 10 ✓ Max Health -Hameln \$29 ✓ Max Health S29 64.00 20

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Fluoride

SODIUM FLUORIDE

✓ PSM

100

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	euroTabs
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental) FERROUS FUMARATE WITH FOLIC ACID	3.09	100	✓ F	erro-tab
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	4.68	60	✓ F	erro-F-Tabs
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 m		errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) — Special Authority se Inj 50 mg per ml, 10 ml vial		etail 1		erinject

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$		Fully dised	Brand or Generic Manufacturer
Magnesium				
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml		nillips Milk of Magnesia 829
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ <u>M</u>	artindale
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ <u>Zi</u>	ncaps

Subsidy
(Manufacturer's Price)
\$ Per

Fully Subsidised

Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Pric	20)	Fully Subsidised	Brand or Generic
	(Manufacturer's Fric	Per	Jubsiuiseu	Manufacturer
EPOETIN ALFA - Special Authority see SA1775 on the previo	us page – Retail pha	ırmacy		
Wastage claimable		•		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓ E	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	✓ E	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	√ E	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	√ E	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	√ E	Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	✓ E	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	√ <u>E</u>	<u> Binocrit</u>
Inj 10,000 iu in 1 ml, syringe	197.50	6	✓ <u>E</u>	<u> Binocrit</u>
Inj 40,000 iu in 1 ml, syringe		1	✓ <u>E</u>	<u> Binocrit</u>
Megaloblastic				
FOLIC ACID				
Tab 0.8 mg	21.84	1,000	✓	Apo-Folic Acid
•	26.60		√ F	olic Acid multichem
₭ Tab 5 mg	5.82	100	√ F	Folic Acid Mylan
,	12.12	500		Apo-Folic Acid
Folic Acid Mylan to be Sole Supply on 1 December 202	21			
Oral lig 50 mcg per ml		25 ml O	P 🗸 E	Biomed
Apo-Folic Acid Tab 0.8 mg to be delisted 1 May 2022)				
Apo-Folic Acid Tab 5 mg to be delisted 1 December 2021)				

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	 Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
TROMBOPAG - Special Authority see SA1743 below	/ – Retail pharmacy		

ELTROMBOPAG – Special Authority see SA1743 below – Retail pharma Wastage claimable

Revolade	28	Tab 25 mg1,550.00
✓ Revolade	28	Tab 50 mg

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
- 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (**idiopathic thrombocytopenic purpura - preparation for splenectomy**) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1	969 on the next page		
Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial		1	✓ Hemlibra
Inj 150 mg in 1 ml vial		1	✓ Hemlibra

	Subsidy	Fully	Brand or
(Manuf	acturer's Price) Su	ubsidised	Generic
	\$ Per	✓	Manufacturer

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Fither:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe		1	✓ NovoSeven RT
Inj 5 mg syringe	,	1	✓ NovoSeven RT
Ini 8 mg syringe		1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

io managou by and maciniopini	na rreatere ereap in eerijaneiter tiiti tiit rtat		
Inj 500 U	1,315.00	1	FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Ini 2.500 U	6.575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

subject to criteria.			
Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe	575.00	1	Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

	Subsidy	F	ully Brand or
	(Manufacturer's Price)	Subsidis	
	\$	Per	✓ Manufacturer
NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharm]			
For patients with haemophilia. Access to funded treatment is	managed by the Hae	emophilia Tr	eaters Group in conjunction
with the National Haemophilia Management Group. Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial		-	✓ RIXUBIS
Inj 2,000 iu vial		=	✓ RIXUBIS
Inj 3,000 iu vial		=	✓ RIXUBIS
• •	•	•	· III/ODIO
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – [For patients with haemophilia. Preferred Brand of short half-		vr \/III \\ A 0000	s to funded treatment is
managed by the Haemophilia Treaters Group in conjunction v			
Inj 250 iu vial			✓ Advate
Inj 500 iu vial			✓ Advate
Inj 1,000 iu vial		-	✓ Advate
Inj 1,500 iu vial		-	✓ Advate
Inj 2,000 iu vial		•	✓ Advate
Inj 3,000 iu vial			✓ Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	•	•	7147410
For patients with haemophilia. Rare Clinical Circumstances I		, rocombinor	ot factor VIII. Access to funded
treatment is managed by the Haemophilia Treaters Group in	conjunction with the r	vational Hae	mopnilia Management Group,
subject to criteria.	227 50	1	√ Kaganata ES
Inj 250 iu vial		-	✓ Kogenate FS
Inj 500 iu vial		-	✓ Kogenate FS
Inj 1,000 iu vial Inj 2,000 iu vial			✓ Kogenate FS ✓ Kogenate FS
Inj 3,000 iu vial	,		✓ Kogenate FS
		1	Nogeliale F3
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]			
For patients with haemophilia A receiving prophylaxis treatme		d treatment is	s managed by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			
Inj 250 iu vial			✓ Adynovate
Inj 500 iu vial			✓ Adynovate
Inj 1,000 iu vial			✓ Adynovate
Inj 2,000 iu vial	2,400.00	1	✓ Adynovate
SODIUM TETRADECYL SULPHATE			
* Inj 3% 2 ml	28.50	5	
	(73.00)		Fibro-vein
TRANEXAMIC ACID			
Tab 500 mg	9.45	60	✓ Mercury Pharma
Vitamin K			
DI IVTOMENIA DIONIE			
PHYTOMENADIONE	0.00	_	/ Kamaldan MM
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO		_	✓ Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO	9.21	5	✓ Konakion MM
Antithrombotic Agents			
Antitinombotic Agents			
Antiplotolot Agento			
Antiplatelet Agents			
ASPIRIN			
* Tab 100 mg	10.80	990	✓ Ethics Aspirin EC
		-	

	Subsidy (Manufacturer's Price) \$	Per	
CLOPIDOGREL			
* Tab 75 mg	4.60	84	✓ <u>Clopidogrel</u> <u>Multichem</u>
DIPYRIDAMOLE			
* Tab long-acting 150 mg	10.90	60	✓ Pytazen SR
TICAGRELOR - Special Authority see SA1955 below - Retail pha	armacy		
* Tab 90 mg	90.00	56	✓ Brilinta
SA1055 Special Authority for Subsidy			

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Fither:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA1646 below - Retail pharmacy

Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe		10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe	125.87	10	 Clexane Forte
Inj 150 mg in 1 ml syringe		10	 Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	
HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	58.57	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml		5	✓ DBL Heparin
, -, ,			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule		50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
, =0,000 14 por, 01=	42.40	ŭ	✓ Heparin DBL S29
LIEDADINICED CALINE	72.70		Tiopaini bbe
HEPARINISED SALINE	GE 10	50	✓ Pfizer
Inj 10 iu per ml, 5 ml	05.48	ου	♥ Pilzer
Oral Anticoagulants			
Oral Anticoagalants			
DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ Pradaxa
Cap 110 mg	76.36	60	✓ Pradaxa
Cap 150 mg	76.36	60	✓ Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ Xarelto
Tab 20 mg		28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	✓ Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	✓ Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg		100	✓ Marevan
* Tab 5 mg		50	✓ Coumadin
ŭ	11.48	100	✓ Marevan
Blood Colony-stimulating Factors			
FILGRASTIM - Special Authority see SA1259 below - Retail p	,	10	Nivestim
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓ Nivestim
Nivestim to be Principal Supply on 1 December 2021			_

✓ Nivestim 10 Nivestim to be Principal Supply on 1 December 2021

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price)	Ş	Fully Subsidised	Brand or Generic
	\$	Per	✓	Manufacturer
PEGFILGRASTIM - Special Authority see SA1912 below - Reta	il pharmacy			
Inj 6 mg per 0.6 ml syringe	1,080.00	1	✓ N	eulastim

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

⇒SA1912 Special Authority for Subsidy

GLUCOSE [DEXTROSE]			
* Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO	30.65	5	✓ Biomed
* Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO	15.00	1	✓ Biomed
POTASSIUM CHLORIDE			
* Inj 75 mg per ml, 10 ml	65.00	50	✓ Juno
SODIUM BICARBONATE			
Inj 8.4%, 50 ml	19.95	1	✓ Biomed
a) Up to 5 inj available on a PSO			
b) Not in combination			
Inj 8.4%, 100 ml	20.50	1	✓ Biomed
 a) Up to 5 inj available on a PSO 			
b) Not in combination			
SODIUM CHLORIDE			
Not funded for use as a nasal drop. Not funded for nebuliser	use except whe	n used in conj	unction with an antibiotic intended
for nebuliser use.			_
Inj 0.9%, bag – Up to 2000 ml available on a PSO			✓ Baxter
	1.26	1,000 ml	✓ Baxter
Only if prescribed on a prescription for renal dialysis, mater	ernity or post-na	atal care in the	home of the patient, or on a PSO
for emergency use. (500 ml and 1,000 ml packs)		_	4.51
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5	✓ Biomed
For Sodium chloride oral liquid formulation refer Standard			/ For earlier Male!
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO	5.40	50	✓ Fresenius Kabi

WATER

- On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order; or

TOTAL PARENTERAL NUTRITION (TPN)

3) When used in the extemporaneous compounding of eye drops; or

Infusion......CBS

4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule – Up to 5 inj available on a PSO7.19	50	✓ Pfizer
Inj 20 ml ampoule - Up to 5 inj available on a PSO5.00	20	Fresenius Kabi
		✓ Multichem

20

1 OP

Fresenius Kabi

✓ TPN

	Subsidy (Manufacturer's Pr \$	ice) Subsi Per	Fully dised	Brand or Generic Manufacturer
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	✓ (Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	9.77	50	✓ <u>E</u>	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)		1,000 ml OP	✓ F	Pedialyte - Bubblegum
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓ F	Phosphate Phebra
POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (11.85)	60	(Chlorvescent
* Tab long-acting 600 mg (8 mmol)	, ,	200	✓ 9	Span-K
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	-	Sodibic Sodibic
SODIUM POLYSTYRENE SULPHONATE			_	
Powder	84.65	454 a OP	√ F	Resonium-A

-			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN

* Tab 2 mg	17.35	500	✓ Apo-Doxazosin
* Tab 4 mg	20.94	500	✓ Apo-Doxazosin
PHENOXYBENZAMINE HYDROC	CHLORIDE		
* Cap 10 mg	65.00	30	✓ BNM S29
	216.67	100	✓ Dibenzyline S29
PRAZOSIN			
* Tab 1 mg	5.53	100	✓ Apo-Prazosin
			✓ Apo-Prazosin
			S29 S29
* Tab 2 mg	7.00	100	✓ Apo-Prazosin
			✓ Apo-Prazosin
			S29 S29

(Apo-Prazosin Tab 1 mg to be delisted 1 May 2022) (Apo-Prazosin Tab 2 mg to be delisted 1 May 2022)

(Apo-Prazosin Tab 5 mg to be delisted 1 May 2022)

Agents Affecting the Renin-Angiotensin System

Tab 5 mg11.70

ACE Inhibitors

CA			

Oral liq 5 mg per ml94	1.99	95 ml OP	✓ Capoten
135	5.00	100 ml OP	✓ Captopril-Mylan S29

Oral liquid restricted to children under 12 years of age.

(Captopril-Mylan S29 Oral lig 5 mg per ml to be delisted 1 January 2022)

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

* Tab 0.5 mg	2.09	90	✓ Zapril
* Tab 2.5 mg		90	✓ Zapril
	8.35	90	✓ Zapril
ENALAPRIL MALEATE			
* Tab 5 mg	1.82	100	✓ Acetec
* Tab 10 mg	2.02	100	✓ Acetec
* Tab 20 mg		100	✓ Acetec
LISINOPRIL			
* Tab 5 mg	2.07	90	✓ Ethics Lisinopril
* Tab 10 mg	2.36	90	✓ Ethics Lisinopril
* Tab 20 mg	3.17	90	 Ethics Lisinopril

✓ Apo-Prazosin

✓ Apo-Prazosin
S29 S29

100

	Subsidy (Manufacturer's Price)		Fully	
	(Manufacturers Frice)	Per		Manufacturer
PERINDOPRIL				
Tab 2 mg	4.95	30	✓	Coversyl
Tab 4 mg	6.30	30	•	Coversyl
QUINAPRIL				
* Tab 5 mg		90	✓	Arrow-Quinapril 5
Arrow-Quinapril 5 to be Principal Supply on 1 February				
* Tab 10 mg		90	•	Arrow-Quinapril 10
Arrow-Quinapril 10 to be Principal Supply on 1 February * Tab 20 mg		90	/	Arrow-Quinapril 20
Arrow-Quinapril 20 to be Principal Supply on 1 February		00	·	Arrow Gumapin 20
ACE Inhibitors with Diuretics				
QUINAPRIL WITH HYDROCHLOROTHIAZIDE				
Tab 10 mg with hydrochlorothiazide 12.5 mg	3.57	28	✓	Accuretic
	4.10	30		Accuretic 10
* Tab 20 mg with hydrochlorothiazide 12.5 mg		30	/	Accuretic 20
(Accuretic Tab 10 mg with hydrochlorothiazide 12.5 mg to be de-	listed 1 March 2022)			
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL				
* Tab 4 mg	2.00	90	•	Candestar
Candestar to be Principal Supply on 1 December 2021	0.00	00	,	O
* Tab 8 mg Candestar to be Principal Supply on 1 December 2021	2.28	90	•	Candestar
* Tab 16 mg	3.31	90	/	Candestar
Candestar to be Principal Supply on 1 December 2021		00		- Curia Cotar
* Tab 32 mg	5.26	90	✓	Candestar
Candestar to be Principal Supply on 1 December 2021				
LOSARTAN POTASSIUM				
* Tab 12.5 mg	1.56	84	✓	Losartan Actavis
* Tab 25 mg		84	_	Losartan Actavis
* Tab 50 mg		84	_	Losartan Actavis
* Tab 100 mg	3.50	84	-	Losartan Actavis
Angiotensin II Antagonists with Diuretics				
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE		_	=	
Tab 50 mg with hydrochlorothiazide 12.5 mg	1.88	30	/	Arrow-Losartan &
				Hydrochlorothiazide
Angiotensin II Antagonists with Neprilysin Inhil	bitors			
SACUBITRIL WITH VALSARTAN - Special Authority see SA19	05 on the next nage –	Ret	ail nharma	^v
Note: Due to the angiotensin II receptor blocking activity of				
ACE inhibitor or another ARB.			23.331	
Tab 24.3 mg with valsartan 25.7 mg	190.00	56	1	Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg		56	_	Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg		56	_	Entresto 97/103

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure: and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, page 123

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

To light calle hydrochionae relet to NETTV 003 3131EW, Anaestile	ilos, Local, po	age 120	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg	3.80	30	✓ Aratac
▲ Tab 200 mg		30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a			
PSO	16.37	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a			
PSO	12 07	10	✓ Hameln S29
1 00	15.09	10	✓ Martindale
Martindale to be Principal Supply on 1 January 2022	10.00		· martinadic
(Hameln \$29 Inj 600 mcg per ml, 1 ml ampoule to be delisted 1 Jan	nuary 2022)		
DIGOXIN	iddi'y 2022)		
	7.00	0.40	/ Lawrente BO
* Tab 62.5 mcg – Up to 30 tab available on a PSO		240	✓ <u>Lanoxin PG</u>
* Tab 250 mcg – Up to 30 tab available on a PSO		240	Lanoxin
* Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	✓ Rythmodan
FLECAINIDE ACETATE			•
▲ Tab 50 mg	19 95	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg		90	✓ Flecainide
Supporting doubling footing	00.01	00	Controlled
			Release Teva
▲ Cap long-acting 200 mg	61.06	90	✓ Flecainide
■ Oap long adding 200 mg	01.00	30	Controlled
			Release Teva
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor
ing to my per mi, to mi ampoule	100.00	J	· I allibucui

Subsidy (Manufacturer's Price) \$162.00	Per 100	✓	I Generic
		✓	Mexiletine Hydrochloride USP \$29
		✓	Mexiletine Hydrochloride USP \$29
202.00	100		Toyo 620
		•	Mexiletine Hydrochloride USP \$29
		•	Teva S29
January 2022) January 2022)			
40.90	50	•	Rytmonorm
macy			
53.00 79.00	100 100		Gutron Gutron
for 2 years where payards as necessar	y. H	ypertensio	on should be avoided, and
r .	macy53.0079.00 for 2 years where prowards as necessar	macy 53.00 100 100 for 2 years where patient owards as necessary. Hy	January 2022) January 2022)40.90 50 macy53.00 100 for 2 years where patient has disab

Beta Adrenoce	ptor Blo	ockers
----------------------	----------	--------

ATENOLOL		
* Tab 50 mg9.33 Mylan Atenolol to be Principal Supply on 1 January 2022	500	✓ Mylan Atenolol
* Tab 100 mg	500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml	300 ml OP	 ✓ Atenolol AFT ✓ Atenolol AFT S29 S29
38.20		✓ Essential Generics S29
Restricted to children under 12 years of age.		
BISOPROLOL FUMARATE		
* Tab 2.5 mg	90	✓ Bisoprolol Mylan
* Tab 5 mg	90	✓ Bisoprolol Mylan
* Tab 10 mg 3.62	90	✓ Bisoprolol Mylan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	1	Manufacturer
CARVEDILOL				
* Tab 6.25 mg	2.24	60	✓	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	✓	Carvedilol Sandoz
* Tab 25 mg	2.95	60	✓	Carvedilol Sandoz
ABETALOL				
* Tab 100 mg	14.50	100	✓	Trandate
★ Tab 200 mg	27.00	100	✓	Trandate
k Inj 5 mg per ml, 20 ml ampoule	59.06	5		
, ,	(88.60)			Trandate
k inj 5 mg per ml, 20 ml vial	42.29	1		
•	(48.20)			Alvogen S29
METOPROLOL SUCCINATE				
★ Tab long-acting 23.75 mg	1.45	30	1	Betaloc CR
★ Tab long-acting 47.5 mg		30	1	Betaloc CR
★ Tab long-acting 95 mg		30	1	Betaloc CR
★ Tab long-acting 190 mg		30	1	Betaloc CR
METOPROLOL TARTRATE				
Tab 50 mg	5.66	100	1	Apo-Metoprolol
745 00 mg				IPCA-Metoprolol
Tab 100 mg	7 55	60		Apo-Metoprolol
7 ab 700 mg		00		IPCA-Metoprolol
★ Tab long-acting 200 mg	23.40	28		Slow-Lopresor
k Inj 1 mg per ml, 5 ml vial		5		Metoprolol IV Mylan
Apo-Metoprolol Tab 50 mg to be delisted 1 March 2022)	20.00	ŭ		
Apo-Metoprolol Tab 100 mg to be delisted 1 March 2022)				
IADOLOL				
Tab 40 mg	16.69	100	/	Apo-Nadolol
	19.19			Nadolol BNM S29
Tab 80 mg		100		Apo-Nadolol
140 00 119	30.39	100		Nadolol BNM S29
Apo-Nadolol Tab 40 mg to be delisted 1 March 2022) Apo-Nadolol Tab 80 mg to be delisted 1 March 2022)	00.09		•	HAUGIOI DINIVI 1023
PINDOLOL – Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who we endorsed accordingly. Pharmacists may annotate the pre dispensing of pindolol.				
taspensing or pindoloi. ★ Tab 5 mg	13 22	100	1	Apo-Pindolol
* Tab 10 mg		100		Ano-Pindolol

	dioponioning of pindolon		
*	Tab 5 mg13.22	100	✓ Apo-Pindolol
*	Tab 10 mg23.12	100	✓ Apo-Pindolol
	Tab 15 mg33.31	100	✓ Apo-Pindolol

(Apo-Pindolol Tab 5 mg to be delisted 1 May 2022)

(Apo-Pindolol Tab 10 mg to be delisted 1 May 2022)

(Apo-Pindolol Tab 15 mg to be delisted 1 May 2022)

52

	Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
PROPRANOLOL				
Tab 10 mg	4.64	100	✓ A	po-Propranolol
Ÿ	7.04		✓ D	rofate
Tab 40 mg	5.72	100	✓ A	po-Propranolol
· ·	8.75		✓ II	PCA-Propranolol
★ Cap long-acting 160 mg	18.17	100	✓ 0	ardinol LA
FOral lig 4 mg per ml - Special Authority see SA				
Retail pharmacy		500 ml	✓ R	Roxane- Propranolol S29

(Apo-Propranolol Tab 10 mg to be delisted 1 March 2022) (Apo-Propranolol Tab 40 mg to be delisted 1 March 2022)

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTAL OL

AND ODIDING

*	Tab 80 mg	.32.58	500	Mylan
	Tab 160 mg			Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑI	ILODIPINE			
*	Tab 2.5 mg	1.08	90	✓ Vasorex
*	Tab 5 mg	0.96	90	✓ Vasorex
*	Tab 10 mg	1.19	90	✓ Vasorex
FE	ELODIPINE			
*	Tab long-acting 2.5 mg	1.45	30	Plendil ER
	Tab long-acting 5 mg		90	✓ Felo 5 ER
	Felo 5 ER to be Principal Supply on 1 January 2022			
*	Tab long-acting 10 mg	4.32	90	✓ Felo 10 ER
	Felo 10 ER to be Principal Supply on 1 January 2022			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
IFEDIPINE				
Fab long-acting 10 mg	18.80	56	✓	Tensipine MR10 S29
Tab long-acting 20 mg	9.12	50	✓	Mylan (12 hr
				release) S29
	17.72	100	1	Nyefax Retard
Fab long-acting 30 mg	4.78	14	1	Mylan Italy (24 hr
				release) S29
	34.10	100	1	Mylan (24 hr
				release) S29
Tab long-acting 60 mg	52.81	100	1	Mylan (24 hr
				release) \$29
Other Calcium Channel Blockers				
ILTIAZEM HYDROCHLORIDE				
← Tab 60 mg	8.50	100	1	Dilzem
Cap long-acting 120 mg	33.42	500		Apo-Diltiazem CD
Cap long-acting 180 mg	7.00	30	1	Cardizem CD
	50.05	500		Apo-Diltiazem CD
Cap long-acting 240 mg	9.30 66.76	30 500		Cardizem CD Apo-Diltiazem CD
Dilzem Tab 60 mg to be delisted 1 January 2022) Apo-Diltiazem CD Cap long-acting 180 mg to be del Apo-Diltiazem CD Cap long-acting 240 mg to be del ERHEXILINE MALEATE				
€ Tab 100 mg	62 90	100	1	Pexsig
ERAPAMIL HYDROCHLORIDE				
Tab 40 mg	7.01	100	1	Isoptin
€ Tab 80 mg		100		Isoptin
€ Tab long-acting 120 mg		100		Isoptin Retard \$29
rab long-acting 120 mg	30.02	100		Isoptin SR
Fab long-acting 240 mg	15 12	30		Isoptin SR
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj av		00	•	isopiiii ori
PSO		5	1	Isoptin
Centrally-Acting Agents				
LONIDINE				
RECITIONNE Fatch 2.5 mg, 100 mcg per day − Only on a pre	scription 10.34	4	1	Mylan
Patch 5 mg, 200 mcg per day – Only on a presc		4		Mylan
Patch 7.5 mg, 300 mcg per day — Only on a pre	•	4		Mylan
0. 01 , ,				<u>y</u>
LONIDINE HYDROCHLORIDE	0 75	112	.1	Clonidine BNM
4 Tab 0E mag				
	27 07	1()()		
F Tab 150 mcg		100	•	Catapres
Tab 25 mcg Tab 150 mcg Catapres to be Principal Supply on 1 Janual Inj 150 mcg per ml, 1 ml ampoule	ry 2022	100		Medsurge

(M	Subsidy anufacturer's Price \$) Per	Fully Subsidised	
METHYLDOPA	т.			
* Tab 250 mg	15.10 52.85	100 500		Methyldopa Mylan Methyldopa Mylan S29 S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
k Tab 1 mg		30		Burinex S29 S29
(Ini FOO mag nor m) 4 m) vial	16.36	100		Burinex Burinex
Inj 500 mcg per ml, 4 ml vial	7.95	5	•	Durinex
UROSEMIDE [FRUSEMIDE] Tab 40 mg - Up to 30 tab available on a PSO	7 24	1,000	•	Apo-Furosemide
Tab 40 mg Op to 00 tab available on a 1 00	8.00	1,000		IPCA-Frusemide
← Tab 500 mg	25.00	50	✓	Urex Forte
	89.48		•	Furosemid- Ratiopharm S29
	169.96	100	•	Furosemid- Ratiopharm S29
♦ Oral liq 10 mg per ml	11.20	80 ml (OP 🗸	<u>Lasix</u>
l≮ Inj 10 mg per ml, 25 ml ampoule l√ Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSC Apo-Furosemide Tab 40 mg to be delisted 1 March 2022)		6 5		<u>Lasix</u> <u>Furosemide-Baxter</u>
Potassium Sparing Diuretics				
MILORIDE HYDROCHLORIDE	00.00)	ND	Biomed
Oral liq 1 mg per ml		25 ml (JP •	Biomea
EPLERENONE – Special Authority see SA1728 below – Retail pha Tab 50 mg		30	J	Inspra
Tab 55 mg		30		Inspra
➤ SA1728 Special Authority for Subsidy nitial application from any relevant practitioner. Approvals valid w ne following criteria: noth: 1 Patient has heart failure with ejection fraction less than 40%; 2 Either: 2.1 Patient is intolerant to optimal dosing of spironolacton 2.2 Patient has experienced a clinically significant advers	ithout further ren and e; or	ewal u	nless notif	· ied for applications mea
METOLAZONE				
Tab 5 mg	CBS	1	1	Metolazone S29
•		50		Zaroxolyn S29
PIRONOLACTONE				
★ Tab 25 mg	4.38	100	✓.	Spiractin

* Tab 100 mg11.80

✓ Spiractin

✓ Biomed

100

25 ml OP

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Out state		F. II.	December 1
1	Subsidy (Manufacturer's Price \$) ;	Fully Subsidised	
Potassium Sparing Combination Diuretics				
MILORIDE HYDROCHLORIDE WITH FUROSEMIDE			_	
★ Tab 5 mg with furosemide 40 mg		28	•	Frumil
MILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZID * Tab 5 mg with hydrochlorothiazide 50 mg		50	✓	Moduretic
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
★ Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	•	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerge	•			
₭ Tab 5 mg	34.55	500	•	Arrow- Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	26.00 2	25 ml O	P 🗸	Biomed
HLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	3 00	30	,	Igroton \$29
740 23 Hg	6.50	50		Hygroton
NDAPAMIDE				
F Tab 2.5 mg	10.45 11.61	90 100		<u>Dapa-Tabs</u> Mylan
	11.01	100	·	Indapamide \$29
Lipid-Modifying Agents				
Fibrates				
EZAFIBRATE				
₹ Tab 200 mg ₹ Tab long-acting 400 mg		90 30		Bezalip Bezalip Retard
		30	•	bezanp netaru
Other Lipid-Modifying Agents				
CIPIMOX Cap 250 mg	21 56	30	,	Olbetam
	21.00	50		Olbetam S29 S29
Resins				
OLESTIPOL HYDROCHLORIDE	00.00	00		0-1
Grans for oral liq 5 g	32.89	30	•	Colestid

✓ Ezetimibe Sandoz

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN				
* Tab 10 mg	6.16	500	•	Lorstat
Lorstat to be Principal Supply on 1 December 2021			_	
* Tab 20 mg	9.24	500	•	Lorstat
Lorstat to be Principal Supply on 1 December 2021 * Tab 40 mg	14.00	500	1	Lorstat
Lorstat to be Principal Supply on 1 December 2021	14.32	500	•	LUISIAI
* Tab 80 mg	26.54	500	1	Lorstat
Lorstat to be Principal Supply on 1 December 2021				
PRAVASTATIN				
* Tab 20 mg	2.11	28	✓	Pravastatin Mylan
* Tab 40 mg	3.61	28	•	Pravastatin Mylan
SIMVASTATIN				
* Tab 10 mg	1.23	90		Simvastatin Mylan
* Tab 20 mg		90		Simvastatin Mylan
* Tab 40 mg		90		Simvastatin Mylan
* Tab 80 mg	7.12	90	•	Simvastatin Mylan
Selective Cholesterol Absorption Inhibitors				
EZETIMIBE - Special Authority see SA1045 below - Retail phar	macv			

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 on the next page - Retail pharmacy

5.15 30 ✓ Zimybe	Tab 10 mg with simvastatin 10 mg5.15
6.15 30 ✓ Zimybe	Tab 10 mg with simvastatin 20 mg
7.15 30 ✓ Zimybe	Tab 10 mg with simvastatin 40 mg7.15
8.15 30 Zimybe	Tab 10 mg with simvastatin 80 mg8.15

Subsidy		Fully	Brand or
(Manufacturer's \$	Price)	Subsidised <	Generic Manufacturer
⇒SA1046 Special Authority for Subsidy			
Initial application from any relevant practitioner. Approvals valid for 2 years fo All of the following:	r application	is meeting th	ne following criteria:
Patient has a calculated absolute risk of cardiovascular disease of at lea	st 15% over	5 year; and	
2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and			
3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/ atoryastatin.	litre with the	use of the r	naximal tolerated dose of
Notes: A patient who has failed to reduce their LDL cholesterol to less than or e	egual to 2.0	mmol/litre w	ith the use of a less poten
statin should use a more potent statin prior to consideration being given to the u	ise of non-st	atin therapie	es.
Other treatment options including fibrates, resins and nicotinic acid should be co			
f a patient's LDL cholesterol cannot be calculated because the triglyceride level performed and if the LDL cholesterol again cannot be calculated then it can be or			
2.0 mmol/litre.	301101001001	nat the LBL	onologici or lo groater trial
Renewal from any relevant practitioner. Approvals valid for 2 years where the t	reatment re	mains appro	priate and the patient is
			p
penefiting from treatment.			F
Nitrates			,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,
Nitrates GLYCERYL TRINITRATE			
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose			
Nitrates GLYCERYL TRINITRATE	250 dose		itrolingual Pump Spray
Nitrates SLYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30	OP ✔ N	itrolingual Pump Spray itroderm TTS
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO		OP ✔ N	itrolingual Pump Spray
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30	OP	itrolingual Pump Spray itroderm TTS
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose	30 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS
Nitrates GLYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose − Up to 250 dose available on a PSO	30 30 100	OP	itrolingual Pump Spray itroderm TTS itroderm TTS
Nitrates SILYCERYL TRINITRATE ★ Oral pump spray, 400 mcg per dose − Up to 250 dose available on a PSO	30 30 100 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30	OP	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30 90	OP VNVNVN	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard uride
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30	OP VNVNVNVN	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard
Nitrates GLYCERYL TRINITRATE * Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	30 30 100 30 90	OP V N V N V N V N N V N N N N N N N N N	itrolingual Pump Spray itroderm TTS itroderm TTS smo 20 smo 40 Retard uride

ΗY	DR	Al	LAZII	NE F	IYD	R
	_				_	

Vasodilators

HYDRALAZINE HYDROCHLORIDE

*	Tab 25 mg - Special Authority see SA1321 on the next page -		
	Retail pharmacyCBS	1	Hydralazine
		56	✓ Onelink S29
		84	✓ AMDIPHARM \$29
		100	✓ Onelink S29
*	lnj 20 mg ampoule25.90	5	✓ Apresoline

Subs		Fully	Brand or
(Manufactur		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

MINIOVIDII

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL ▲ Tab 10 mg	70.00	100	✓ Loniten
NICORANDIL ▲ Tab 10 mg	25.57	60	✓ <u>Ikorel</u> ✓ Ikorel
Tab 20 mg	32.28	60	✓ <u>Ikorel</u>
* Inj 12 mg per ml, 10 ml ampoule	217.90	5	✓ Hospira
Tab 400 mg	42.26	50	✓ Trental 400

Endothelin Receptor Antagonists

AMBRISENTAN - Special Authority see SA1702 below - Retai	II pharmacy		
Tab 5 mg	1,550.00	30	✓ Ambrisentan Mylan
Tab 10 mg	1,550.00	30	✓ Ambrisentan Mylan

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u> or: The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz
BOSENTAN – Special Authority see SA1991 below – Retail pharmacy

Tab 62.5 mg	119.85	60	✓ Bosentan Dr Reddy's
Bosentan Dr Reddy's to be Principal Supply on 1 Decer	mber 2021		
Tab 125 mg	119.85	60	Bosentan Dr

Bosentan Dr Reddy's to be Principal Supply on 1 December 2021

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Fither:

continued...

Reddy's

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
- 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
- 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy: or
- 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA1992 on the next page -	- Retail pharmacy		
Tab 25 mg	0.85	4	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022			
Tab 50 mg	1.70	4	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022			
Tab 100 mg	10.20	12	✓ Vedafil
Vedafil to be Principal Supply on 1 January 2022			

S

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 on the r	next page – Retail pharr	nacy	
Inj 500 mcg vial	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 92

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	✓ Differin
ISOTRETINOIN - Special Authority see SA2023 below - Ret	ail pharmacy		
Cap 5 mg		60	Oratane
Cap 10 mg	18.75	120	Oratane
Cap 20 mg	26.73	120	Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per g − Maximum of 50 g per prescription15.57 50 g OP ✓ ReTrieve
ReTrieve to be Principal Supply on 1 January 2022

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 92

HYDROGEN PEROXIDE

				Fully Brand or		
	(Manufacturer's F		sidised Generic			
	\$	Per	✓ Manufacturer			
MUPIROCIN						
Oint 2%	6.60 (10.50)	15 g OP	Bactroban			
a) Only on a prescriptionb) Not in combination						
SODIUM FUSIDATE [FUSIDIC ACID]						
Crm 2%	1.59	5 g OP	✓ Foban			
 a) Maximum of 5 g per prescription 						
b) Only on a prescription						
c) Not in combination						
d) Foban to be Principal Supply on 1 December 2021 Oint 2%	1.50	E a OB	✓ Foban			
a) Maximum of 5 g per prescription	1.59	5 g OP	▼ FUDAII			
b) Only on a prescription						
c) Not in combination						
d) Foban to be Principal Supply on 1 December 2021						
SULFADIAZINE SILVER						
Crm 1%	10.80	50 g OP	✓ Flamazine			
a) Up to 250 g available on a PSO		5				
b) Not in combination						
,						
Antifungals Topical						
Formula with modern to the INFECTION O. Antiferrorle	- 00					
For systemic antifungals, refer to INFECTIONS, Antifungals, pag	je 99					
AMOROLFINE						
a) Only on a prescription						
b) Not in combination Nail soln 5%	14.02	5 ml OP	✓ MycoNail			
	14.33	3 IIII OF	♥ <u>IVIYCOIVAII</u>			
CICLOPIROX OLAMINE						
a) Only on a prescription b) Not in combination						
Nail-soln 8%	5 72	7 ml OP	✓ Apo-Ciclopirox			
'Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022)		7 1111 01	• Apo-ololopilox			
CLOTRIMAZOLE						
* Crm 1%	0.77	20 g OP	✓ Clomazol			
a) Only on a prescription		20 9 01	- Olomazor			
b) Not in combination						
* Soln 1%	4.36	20 ml OP				
	(7.55)		Canesten			
a) Only on a prescription	• •					
b) Not in combination						
ECONAZOLE NITRATE						
Crm 1%	1.00	20 g OP				
	(7.48)	•	Pevaryl			
a) Only on a prescription						
b) Not in combination						
Foaming soln 1%, 10 ml sachets		3				
	(17.23)		Pevaryl			
A Contract of the contract of						
a) Only on a prescriptionb) Not in combination						

	Subsidy (Manufacturer's F	Orion) Cub	Fully sidised	Brand or Generic
	(Manufacturer's F	Per	siuiseu 🗸	Manufacturer
MICONAZOLE NITRATE				
* Crm 2%	0.81	15 g OP	✓ N	<u>lultichem</u>
a) Only on a prescription				
b) Not in combination				
Lotn 2%	4.36	30 ml OP		
	(10.03))aktarin
a) Only on a prescription				
b) Not in combination				
米 Tinct 2%	4.36	30 ml OP		
	(12.10)			aktarin
a) Only on a prescription				
b) Not in combination				

Antipruritic Preparations

CALAMINE

a) Only on a prescription

b) Not in combination

100 g

✓ healthE Calamine Aqueous Cream BP

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%

9 20 g OP

✓ Itch-Soothe

Itch-Soothe to be Principal Supply on 1 December 2021

MENTHOL - Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteriod Plain
- 2) With or without other dermatological galenicals.

Crystals	6.92	25 g	✓ MidWest
	29.60	100 g	✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 82

Corticosteroids - Plain

BE	TAMETHASONE DIPROPIONATE			
	Crm 0.05%	2.96	15 g OP	✓ <u>Diprosone</u>
		36.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	Diprosone OV
BE	TAMETHASONE VALERATE			
*	Crm 0.1%	4.53	50 g OP	✓ Beta Cream
	Beta Cream to be Principal Supply on 1 January 2022		_	
*	Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
	Beta Ointment to be Principal Supply on 1 January 2022			
*	Lotn 0.1%	25.00	50 ml OP	✓ Betnovate

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Eully	Prand or
	(Manufacturer's Pr	rice) Sub	Fully sidised	Brand or Generic
	\$	Per	√	Manufacturer
CLOBETASOL PROPIONATE				
* Crm 0.05%	2.18	30 g OP	✓	Dermol
* Oint 0.05%	2.12	30 g OP	√ i	Dermol
CLOBETASONE BUTYRATE		•		
Crm 0.05%	5.38	30 g OP		
	(10.00)	22 9 21		Eumovate
HYDROCORTISONE	,			
* Crm 1% – Only on a prescription	3 70	100 g OP	✓ I	Hydrocortisone
on the one processiple.			-	(PSM)
	17.15	500 g	✓ I	Hydrocortisone
		500 g	-	(PSM)
* Powder – Only in combination	49.95	25 g	1	ABM
Up to 5% in a dermatological base (not proprietary Topi			-	
galenicals		,		g
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only	on			
a prescription		250 ml	✓ I	DP Lotn HC
		200 1111		<u> </u>
HYDROCORTISONE BUTYRATE Lipocream 0.1%	A 95	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OP	_	Locoid
Locoid to be Principal Supply on 1 December 2021	10.20	100 g O1	•	Locolu
Milky emul 0.1%	12 33	100 ml OP	✓ I	Locoid Crelo
Locoid Crelo to be Principal Supply on 1 December 202		100 1111 01		200014 01010
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	1	Advantan
Oint 0.1%		15 g OP		Advantan Advantan
MOMETASONE FUROATE		10 9 01		- Tavarran
Crm 0.1%	1 05	15 g OP	1	Elocon Alcohol Free
OIIII 0.176	3.10	50 g OP		Elocon Alcohol Free
Elocon Alcohol Free to be Principal Supply on 1 Februa		30 g Oi	•	LIOCOII AICOIIOI I 166
Oint 0.1%	,	15 g OP	✓ I	Elocon
	2.90	50 g OP		Elocon
Elocon to be Principal Supply on 1 February 2022		oug o.		
Lotn 0.1%	4.50	30 ml OP	✓	Elocon
Elocon to be Principal Supply on 1 February 2022				
TRIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	1	Aristocort
Oint 0.02%	6.35	100 g OP	-	Aristocort
Corticosteroids - Combination				
DETAMETI IA CONE MAI EDATE MUTU CODUINA ELICIDATE (EL	IOIDIO AOIDI			
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU		15 ~ OD		
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		Fucicort
a) Maximum of 1E a new properties	(10.45)		ı	UCICUIT
a) Maximum of 15 g per prescriptionb) Only on a prescription				
, , ,				
HYDROCORTISONE WITH MICONAZOLE – Only on a prescri		15 - 00	, .	Mianama II
* Crm 1% with miconazole nitrate 2%	1.89	15 g OP	✓	Micreme H
Micreme H to be Principal Supply on 1 December 2021				

	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully Brand or idised Generic ✓ Manufacturer
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN — Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% (Pimafucort Crm 1% with natamycin 1% and neomycin sulphate TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	3.35 35 <i>0.5% to be delis</i> CIN AND NYSTA	15 g OP 15 g OP sted 1 May 2022)	✓ Pimafucort ✓ Pimafucort
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m and gramicidin 250 mcg per g - Only on a prescription		15 g OP	Viaderm KC
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE * Crm 5% pump bottle	4.48	500 ml OP	✓ <u>healthE</u> Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE Dimethicone 10%
ZINC AND CASTOR OIL * Oint	4.65	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM			
* Crm	1.73	500 g	✓ GEM Aqueous Cream
	1.92		✓ Basic AquaCream✓ Boucher✓ Medco
(Basic AquaCream Crm to be delisted 1 April 2022) (Boucher Crm to be delisted 1 April 2022) (Medco Crm to be delisted 1 April 2022) CETOMACROGOL			
CETOMACROGOL ★ Crm BP CETOMACROGOL WITH GLYCEROL	2.48	500 g	✓ healthE
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ ADE ✓ <u>Boucher</u> ✓ Kenkay Sorbolene ✓ Pharmacy Health Sorbolene with Glycerin
(ADE Crm 90% with glycerol 10% to be delisted 1 January 2022 (Kenkay Sorbolene Crm 90% with glycerol 10% to be delisted 1	,	1,000 ml OP	✓ ADE ✓ <u>Boucher</u>
(ADE Crm 90% with glycerol 10% to be delisted 1 January 2022			
EMULSIFYING OINTMENT * Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's I \$	Price) Subs	Fully idised	Brand or Generic Manufacturer
OIL IN WATER EMULSION				
€ Crm	2.19	500 g	✓ (D/W Fatty Emulsion Cream
ARAFFIN				
Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ ł	nealthE
REA			٠.	
€ Crm 10%	1.37	100 g OP	✓ ł	ealthE Urea Cream
VOOL FAT WITH MINERAL OIL - Only on a prescription				
Lotn hydrous 3% with mineral oil	5.60	1,000 ml		
	(11.95)		[OP Lotion
	1.40	250 ml OP	_	
	(4.53)		[OP Lotion
	5.60	1,000 ml		Halaa Kadi Labaa
	(20.53)			Alpha-Keri Lotion
	(23.91) 1.40	250 ml OP	E	3K Lotion
	(7.73)	250 IIII OF	F	3K Lotion
Other Dermatological Bases				
ARAFFIN				
White soft - Only in combination		450 g	_	nealthE
Only to a south to all any other decount of a final and a straight of	19.99	2,500 g	_	nealthE
Only in combination with a dermatological galenical or a	is a diluent for a	proprietary Topi	cal Co	rticosteroid – Plain.
Minor Skin Infections				
OVIDONE IODINE				
Oint 10%	7.40	65 g OP	✓ [Betadine
a) Maximum of 130 g per prescription				
b) Only on a prescription				
Antiseptic Solution 10%		100 ml	-	Riodine
Antiseptic soln 10%		15 ml		Riodine
	5.40	500 ml	✓ F	Riodine
Skin preparation, povidone iodine 10% with 30% alcohol		100 ml	_	
Olds are continued as in the 400/ with 700/	(3.48)	4001	Е	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol		100 ml		Ofizor
	(7.78)		h	Pfizer
Parasiticidal Preparations				
Parasiticidal Preparations				
Parasiticidal Preparations IMETHICONE Lotn 4%		200 ml OP		nealthE

Lotion

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO.......17.20 4 ✓ Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Roth:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5% 5.75 Lotn 5% 3.99	30 g OP 30 ml OP	✓ <u>Lyderm</u> ✓ <u>A-Scabies</u>
PHENOTHRIN		
Shampoo 0.5%11.36	200 ml OP	✓ Parasidose
(Parasidose Shamnoo 0.5% to be delisted 1. January 2022)		

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 below - Retail	pharmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of

continued...

		[DERM	ATOLOGICALS
	Subsidy (Manufacturer's Pr \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
continued				
treatment and patient has been counselled and understa and that they must not become pregnant during treatmer or				
2 Patient is not of child bearing potential.				
BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL Foam spray 500 mcg with calcipotriol 50 mcg per g Gel 500 mcg with calcipotriol 50 mcg per g Daivobet to be Principal Supply on 1 December 2021		60 g OP 60 g OP		nstilar aivobet
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	✓ D	aivobet
CALCIPOTRIOL Oint 50 mcg per g	40.00	120 g OP	✓ D	aivonex
COAL TAR	22.25	000		
Soln BP - Only in combination		200 ml etary Topical C	_	<u>lidwest</u> teriod – Plain
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% ar				
allantoin crm 2.5%	6.59 (8.00) 3.43	75 g OP 30 g OP	E	gopsoryl TA
	(4.35)	Ü	E	gopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97 7.95	25 g OP 40 g OP		coco-Scalp coco-Scalp
PIMECROLIMUS – Special Authority see SA1970 below – Reta a) Maximum of 15 g per prescription	il pharmacy	-		р
b) Note: a maximum of 15 g per prescription and no more Cream 1%		ion per 12 we 15 g OP		lidel
■ SA1970 Special Authority for Subsidy Initial application only from a dermatologist, paediatrician, ophi of a dermatologist, paediatrician or ophthalmologist. Approvals meeting the following criteria: Both:				
 Patient has atopic dermatitis on the eyelid; and Patient has at least one of the following contraindications documented epidermal atrophy, documented allergy to to pressure. 				
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE	SCEIN - Only on	a prescription	n	

- PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN Onl
 * Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.........44 500 ml ✓ Pinetarsol SALICYLIC ACID 250 g ✓ Midwest
 - ✓ PSM
 - 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
 - 2) With or without other dermatological galenicals.

DERMATOLOGICALS				
	Subsidy (Manufacturer's Pri \$	ice) Subs	Fully sidised	Brand or Generic Manufacturer
SULPHUR				
Precipitated – Only in combination		100 g		idwest
 Only in combination with a dermatological base or p With or without other dermatological galenicals. 	oroprietary Topica	al Corticostero	oid – Pla	in
TACROLIMUS				
Oint 0.1% - Special Authority see SA2074 below - Retail				
pharmacya) Maximum of 30 g per prescription	33.00	30 g OP	✓ Ze	ematop
b) Note: a maximum of 30 g per prescription and no mo	re than one preso	cription per 12	2 weeks.	
⇒SA2074 Special Authority for Subsidy				
Initial application only from a dermatologist, paediatrician or any paediatrician, . Approvals valid without further renewal unless no Both:				
 Patient has atopic dermatitis on the face; and Patient has at least one of the following contraindications t documented epidermal atrophy or documented allergy to t 			orificial d	ermatitis, rosacea,
Scalp Preparations				
BETAMETHASONE VALERATE				
* Scalp app 0.1% Beta Scalp to be Principal Supply on 1 January 2022	9.84	100 ml OP	✓ B	eta Scalp
CLOBETASOL PROPIONATE				
* Scalp app 0.05%	5.69	30 ml OP	✓ <u>D</u>	ermol
HYDROCORTISONE BUTYRATE	6.57	100 00	./ 1.	aaid
Scalp lotn 0.1% Locoid to be Principal Supply on 1 December 2021	0.57	100 ml OP	♥ L(ocoid

KETOCONAZOLE

100 ml OP

 Sebizole Sebizole

a) Maximum of 100 ml per prescription

b) Only on a prescription

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Lotn,......5.10 200 g OP

✓ Marine Blue Lotion SPF 50+

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 70

IMIQUIMOD

24 Crm 5%, 250 mg sachet......21.72 ✓ Perrigo

DERMATOLOGICALS

	Subsidy (Manufacturer's Pr \$	ice) Subs	Fully idised	Brand or Generic Manufacturer
PODOPHYLLOTOXIN Soln 0.5%	33.60	3.5 ml OP	✓ C	Condyline
 a) Maximum of 3.5 ml per prescription b) Only on a prescription 				

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

Efudix to be Principal Supply on 1 December 2021

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Contraceptives - Non-hormonal

Condoms

00	NDOMO			
	NDOMS 10 mm — Unito 144 day ayailahla on a PSO	11 40	144	✓ Moments
	49 mm – Up to 144 dev available on a PSO		10	✓ Moments
不	33 HIIII	11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription	11.04	144	• INIOILIEILE
	b) Up to 60 dev available on a PSO			
*	53 mm. 0.05 mm thickness	0.05	10	✓ Moments
~	JO HIIII, U.UJ HIIII HIIUNIESS	11.42	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.74	177	· momenta
	b) Maximum of 60 dev per prescription			
*	53 mm, chocolate, brown	0.95	10	✓ Moments
	oo min, onoonato, brown	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.07	177	- Invinorito
	b) Maximum of 60 dev per prescription			
*	53 mm, strawberry, red	0.95	10	✓ Moments
	55, 51.5017, 100	11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO	11.01		111011101110
	b) Maximum of 60 dev per prescription			
*	56 mm	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
ĸ	56 mm, 0.05 mm thickness	1.30	12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
*	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	a) Maximum of 60 dev per prescription			
	b) Up to 60 dev available on a PSO			
K	56 mm, 0.08 mm thickness	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
K	56 mm, 0.08 mm thickness, red	0.97	10	✓ Moments
		11.64	144	✓ Moments
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
K	56 mm, chocolate		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			
K	56 mm, strawberry		12	✓ Gold Knight
		15.57	144	✓ Gold Knight
	a) Up to 60 dev available on a PSO			
	b) Maximum of 60 dev per prescription			_
K	60 mm		12	✓ Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		✓ Gold Knight XL

_					
		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	I Generic
		\$	Per	•	Manufacturer
*	60 mm (bulk pack)	14.87	144	1	Gold Knight XL
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IÚD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	.18.45	1	✓ Choice
				TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

★ Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO.......10.00 84 Mercilon 28

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	Per	Subsidised	Generic Manufacturer	
ETHINN CECTRARIOL WITH FLONOROFOTOE					_
ETHINYLOESTRADIOL WITH LEVONORGESTREL					
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_				
Up to 112 tab available on a PSO	2.18	84	✓ M	licrogynon 20 ED	
	6.45	112	✓ F	emme-Tab ED	
* Tab 50 mcg with levonorgestrel 125 mcg and 7 inert tab - U	lp g				
to 84 tab available on a PSO		84	✓ M	licrogynon 50 ED	
* Tab 30 mcg with levonorgestrel 150 mcg		63		0,	
3 3	(16.50)		M	licrogynon 30	
a) Higher subsidy of \$15.00 per 63 tab with Special Aut	hority see SA0500 on	the pr		0,	
b) Up to 63 tab available on a PSO	,				
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_				
Up to 112 tab available on a PSO		84	✓ 1.	evlen ED	
op to 112 tab available off a 1 00	6.45	112	_	emme-Tab ED	
	0.43	112	• 1	Cililic-Tab Lb	
ETHINYLOESTRADIOL WITH NORETHISTERONE					
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to)				
84 tab available on a PSO	6.95	84	✓ <u>B</u>	revinor 1/28	
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - L	Jp				
to 84 tab available on a PSO		84	✓ N	orimin	

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ᄕ	VONORGESTREL			
*	Tab 30 mcg - Up to 84 tab available on a PSO	.16.50	84	✓ Microlut
		22.00	112	✓ Microlut
*	Subdermal implant (2 × 75 mg rods) – Up to 3 pack available			
	on a PSO	106.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO	7.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	1	Noriday 28
Emergency Contraceptives				
LEVONORGESTREL * Tab 1.5 mg a) Maximum of 2 tab per prescription b) Up to 5 tab available on a PSO	4.95	1	•	Postinor-1

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

ACETIC ACID WITH HADDOAAOUINOLINE VAID DICINOLEIC VOID

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme
•	.0 9 0.	<u></u>
NYSTATIN	75 · OD	Alliana
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ <u>Nilstat</u>

Myometrial and Vaginal Hormone Preparations

RGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a PSO	160.00	5	✓ DBL Ergometrine
ESTRIOL			
Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
Pessaries 500 mcg	6.86	15	✓ Ovestin
XYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	 Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	 Oxytocin BNM
XYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj availab	le on a PSO		-
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml Syntometrine to be Principal Supply on 1 January 2022		5	✓ Syntometrine

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Pregnancy Test

✓ Smith BioMed Rapid

Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 110

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

* Tab 5 mg4.81 100

Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking oxybutynin prior to 1 June 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of oxybutynin.

(Apo-Oxybutynin Tab 5 mg to be delisted 1 May 2022)

(Apo-Oxybutynin Oral liq 5 mg per 5 ml to be delisted 1 May 2022)

79

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 below -

Retail pharmacy......31.80 200 ml OP **✓ Biomed**

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

*	Grans eff 4 g sachets	22	28	✓ <u>Ural</u>
SO	LIFENACIN SUCCINATE			
	Tab 5 mg	05	30	✓ Solifenacin Mylan
	Solifenacin Mylan to be Principal Supply on 1 December 2021			
	Tab 10 mg	72	30	✓ Solifenacin Mylan
	Solifenacin Mylan to be Principal Supply on 1 December 2021			

Detection of Substances in Urine

ORTHO-TOLIDINE

*	Compound diagnostic sticks	7.50	50 test OP	
		(8.25)		Hemastix
T E	TD A DD O MODULEN OL			

TETRABROMOPHENOL

Obstetric Preparations

Antiprogesterones

				 -
MI	H۲	РΚ	IST	l⊢ .

✓ Mifegyne	1	Tab 200 mg60.00
✓ Mifegyne	3	180.00

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy

nufacturaria Drica

Fully

Cubaidiaad

Brand or

	(Manufacturer's Price) \$	Per	sidised ✓	Manufacturer
Calcium Homeostasis				
CALCITONIN * Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ M	iacalcic
CINACALCET - Special Authority see SA1618 below - Retail p	harmacy			
Tab 30 mg - Wastage claimable	42.06	28	✓ C	inacalet Devatis
	210.30		✓ Se	ensipar
Tab 60 mg - Wastage claimable	84.12	28	✓ C	inacalet Devatis

⇒SA1618 Special Authority for Subsidy

(Sensipar Tab 30 mg to be delisted 1 April 2022)

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Zoledronic acid Mylan to be Principal Supply on 1 December 2021

⇒SA2031 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	✓	Manufacturer

continued...

Initial application — (early breast cancer) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 Treatment to be used as adjuvant therapy for early breast cancer; and

PETAMETHA COME CODILIM DUOCDUATE MITH PETAMETHA COME ACETATE

- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 2 years.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASO	ONE ACETAT	E	
* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20	5	
, ,	(36.96)		Celestone
	,		Chronodose
DEXAMETHASONE			
* Tab 0.5 mg - Up to 60 tab available on a PSO	1.50	30	✓ Dexmethsone
Dexmethsone to be Principal Supply on 1 January 2022	1.50	30	Dexilienisone
* Tab 4 mg – Up to 30 tab available on a PSO	2.65	30	✓ Dexmethsone
Dexmethsone to be Principal Supply on 1 January 2022	2.05	30	Dexilienisone
Oral lig 1 mg per ml	45.00	25 ml OP	✓ Biomed
	45.00	23 IIII OF	▼ bioilieu
DEXAMETHASONE PHOSPHATE			
Dexamethasone phosphate injection will not be funded for oral			
* Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO	9.25	10	✓ <u>Dexamethasone</u>
			Phosphate
			<u>Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO	16.37	10	✓ <u>Dexamethasone</u>
			<u>Phosphate</u>
			<u>Panpharma</u>
FLUDROCORTISONE ACETATE			
* Tab 100 mcg	14.32	100	✓ Florinef
HYDROCORTISONE			
	0.40	100	/ Davidas
* Tab 5 mg		100	✓ Douglas
* Tab 20 mg		100	✓ Douglas
* Inj 100 mg vial	4.38	1	✓ Solu-Cortef
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
METHYLPREDNISOLONE			
* Tab 4 mg	112.00	100	✓ Medrol
* Tab 100 mg	194.00	20	✓ Medrol
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial	18.90	1	✓ Solu-Medrol-Act-
.,g		•	O-Vial
			•
Inj 125 mg vial	28.90	1	✓ Solu-Medrol-Act-
, ,			O-Vial
Inj 500 mg vial	22.78	1	✓ Solu-Medrol-Act-
			O-Vial
Inj 1 g vial	27.83	1	✓ Solu-Medrol

	Subsidy		Fully	/ Brand or
	(Manufacturer's Price	ر د	runy Subsidised	
	\$	Per		
METHYLPREDNISOLONE ACETATE	<u> </u>			
Inj 40 mg per ml, 1 ml vial	44.40	5	_	Depo-Medrol
		J	•	Depo-iniedi Oi
PREDNISOLONE	0.00			
 Oral liq 5 mg per ml - Up to 30 ml available on a PSO a) Restricted to children under 12 years of age. b) Redipred to be Principal Supply on 1 December 202 		30 ml OF	· •	Redipred
PREDNISONE				
* Tab 1 mg		500	•	Apo-Prednisone
* Tab 2.5 mg	21.04	500	•	Apo-Prednisone
* Tab 5 mg - Up to 30 tab available on a PSO	19.30	500	•	Apo-Prednisone
* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	•	Apo-Prednisone
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	1	UK Synacthen S29
The trip 200 mag por mi, i mi ampoulo		•		AU Synacthen
				Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1		Synacthen Depot
The trig por tris, it the ampound minimum.		•		Synacthene
				Retard S29
TRIANGINGI ONE ACETONIDE				ricial a see
TRIAMCINOLONE ACETONIDE	00.00	-	,	
Inj 10 mg per ml, 1 ml ampoule		5		Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	31.10	5	•	Kenacort-A 40
Sex Hormones Non Contraceptive				
SOX HOMEONOS HOM SOMMASSPATO				
Androgen Agonists and Antagonists				
CYPROTERONE ACETATE				
	14.07	- 0		Rex \$29
Tab 50 mg	14.37	50		
Citarana ta ha Drinainal Complesa 1 January 2000			•	Siterone
Siterone to be Principal Supply on 1 January 2022	00.00	- 0		Citamana
Tab 100 mg	28.03	50	•	Siterone
Siterone to be Principal Supply on 1 January 2022				
(Rex S29 Tab 50 mg to be delisted 1 January 2022)				
TESTOSTERONE				
Patch 5 mg per day	90.00	30	•	Androderm
TESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	/	Depo-Testosterone
TESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12 08	1		Sustanon Ampoules
	12.30	'	•	oustation Ampoules
TESTOSTERONE UNDECANOATE	04.00			
Cap 40 mg - Subsidy by endorsement		60		Andriol Testocaps
Subsidy by endorsement – subsidised for patients who v	•			, ,,
1 November 2021 and the prescription is endorsed acco				
where there exists a record of prior dispensing of testost Inj 250 mg per ml, 4 ml vial				
	an ()()	1	•	Reandron 1000

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✓ Manufacturer

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

Oestrogens

OESTRADIOL - See prescribing guideline above			
* Tab 1 mg		28 OP	
,	11.10)		Estrofem
* Tab 2 mg		28 OP	
	11.10)		Estrofem
* Patch 100 mcg per 24 hours	.7.91	4	✓ Climara
a) No more than 1 patch per week			
b) Only on a prescription			
* Patch 50 mcg per 24 hours	.7.04	4	Climara
a) No more than 1 patch per week			
b) Only on a prescription			
Patch 25 mcg per day	.6.12	8	Estradot
	7.85		Estradiol TDP
			Mylan S29
a) No more than 2 patch per week			
b) Only on a prescription			
Patch 50 mcg per day	.7.04	8	 Estradot 50 mcg
	9.22		 Estradiol TDP
			Mylan S29
a) No more than 2 patch per week			•
b) Only on a prescription			
Patch 75 mcg per day	.7.91	8	✓ Estradot
0 , ,	10.60		Estradiol TDP
			Mylan S29
a) No more than 2 patch per week			,
b) Only on a prescription			
Patch 100 mcg per day	.7.91	8	✓ Estradot
a) No more than 2 patch per week		•	
b) Only on a prescription			
(Climara Patch 100 mcg per 24 hours to be delisted 1 January 2022)			
(Climara Patch 50 mcg per 24 hours to be delisted 1 January 2022)			
(Estradiol TDP Mylan S29) Patch 25 mcg per day to be delisted 1 May 2	2022)		
(Estradiol TDP Mylan 329) Patch 50 mcg per day to be delisted 1 May 2	,		
, , , , , , , , , , , , , , , , , , , ,	,		
(Estradiol TDP Mylan S29 Patch 75 mcg per day to be delisted 1 May 2	(022)		
OESTRADIOL VALERATE – See prescribing guideline above			
* Tab 1 mg		84	Progynova
Tab 2 mg	12.36	84	Progynova
OESTROGENS - See prescribing guideline above			
* Conjugated, equine tab 300 mcg	.3.01	28	
	17.50)		Premarin
* Conjugated, equine tab 625 mcg	.4.12	28	
	17.50)		Premarin

	Subsidy (Manufacturer's Price) \$	Full Subsidise Per 🗸	d Generic
Progestogens			
MEDROXYPROGESTERONE ACETATE - See prescribing guid	deline on the previous	s page	
* Tab 2.5 mg	· ·		Provera
* Tab 5 mg			Provera
* Tab 10 mg	8.94	30	Provera
Progestogen and Oestrogen Combined Prepara	ations		
OESTRADIOL WITH NORETHISTERONE - See prescribing gu	ideline on the previou	us page	
* Tab 1 mg with 0.5 mg norethisterone acetate		28 OP	
	(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate		28 OP	
	(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg	5 40	00.00	
oestradiol tab (12) and 1 mg oestradiol tab (6)		28 OP	Trianguana
	(18.10)		Trisequens
Other Oestrogen Preparations			
ETHINYLOESTRADIOL			
* Tab 10 mcg	17.60	100	NZ Medical and Scientific
OESTRIOL			
* Tab 2 mg	7.00	30	Ovestin Ovestin
Other Progestogen Preparations			
LEVONORGESTREL			
* Intra-uterine device 52 mg	269.50	1	Mirena
* Intra-uterine device 13.5 mg		1	Jaydess
MEDROXYPROGESTERONE ACETATE			
Tab 100 mg	116.15	100	Provera HD
NORETHISTERONE			
* Tab 5 mg - Up to 30 tab available on a PSO	5.49	30	Primolut N
PROGESTERONE			
Cap 100 mg - Special Authority see SA1609 below - Retail			
pharmacy		30	' Utrogestan

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

Subsid (Manufacturer		ılly Brand or ed Generic	
	Per	✓ Manufacturer	

continued...

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

CARBIMAZOLE				
* Tab 5 mg	10	0.80	100	Neo-Mercazole Neo-Mercazole S29 S29
(Neo-Mercazole	S29 S29 Tab 5 mg to be delisted 1 January 2022)			
LEVOTHYROXI	NE			
* Tab 25 mcg	5	5.55	90	Synthroid
	1		28	Mercury Pharma
_			90	Synthroid
	64	1.28 1	,000	Eltroxin
* Tab 100 mg	rg1	.78	28	Mercury Pharma
	- 6	6.01	90	Synthroid
	66	5.78 1	,000	Eltroxin
PROPYLTHIOU	RACIL - Special Authority see SA1199 below - Retail ph	armacv		
Propylthiou	racil is not recommended for patients under the age of 18 gare contraindicated.	,	the patient is	pregnant and other
Tab 50 mg.	35	5.00	100	PTU S29
- CA4400 Cma	alal Authority for Cubaldy			

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SO	MATROPIN (OMNITROPE) - Special Authority see SA2032 of	n the next page -	Retail pha	rmacy
*	Inj 5 mg cartridge	69.75	1	Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			
*	Inj 10 mg cartridge	69.75	1	Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			
*	Inj 15 mg cartridge	139.50	1	Omnitrope
	Omnitrope to be Principal Supply on 1 January 2022			

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)		ubsidised	Generic
	\$	Per		Manufacturer

continued...

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GO	SE	RE	LIN

Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe		1	✓ Teva

	Subsidy (Manufacturer's Price		Fully	Brand or Generic
	(Manufacturer's Frice	Per	uiseu √	Manufacturer
LEUPRORELIN				
Additional subsidy by endorsement where the patient is a c goserelin and the prescription is endorsed accordingly.	hild or adolescent an	nd is unable to	toler	ate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsid	y of			
\$221.60 per 1 inj with Endorsement	66.48	1		
	(221.60)		L	ucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsi				
of \$591.68 per 1 inj with Endorsement		1		
	(591.68)		L	Lucrin Depot 3-month
Vasopressin Agonists				
Vasopiessiii Agoilisis				
DESMOPRESSIN				
Wafer 120 mcg	47.00	30	√ [Minirin Melt
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	√ [Minirin
Tab 200 mcg		30	√ I	Minirin
▲ Nasal drops 100 mcg per ml	39.03	2.5 ml OP	√ [Minirin
▲ Nasal spray 10 mcg per dose	27.95	6 ml OP	√ [<u>Desmopressin-</u> PH&T
				<u>FIIXI</u>
Ini 4 mcg per ml. 1 ml	67.18	10	✓ I	Minirin

Other Endocrine Agents

CABERGOLINE

		Tab 0.5 mg – Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below3.75
Dostinex	8	15.20

⇒SA2070 Special Authority for Waiver of Rule

(Minirin Nasal drops 100 mcg per ml to be delisted 1 January 2022)

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE			
Cap 250 mg	558.00	50	Metopirone

INFECTIONS - AGENTS FOR SYSTEMIC USE			
	Subsidy (Manufacturer's Price) \$		Fully Brand or ised Generic Manufacturer
Anthelmintics			
ALBENDAZOLE - Special Authority see SA1318 below - Retail p	harmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29
⇒SA1318 Special Authority for Subsidy			
Initial application only from an infectious disease specialist or clin patient has hydatids.	nical microbiologist.	Approvals	valid for 6 months where the
Renewal only from an infectious disease specialist or clinical microremains appropriate and the patient is benefitting from the treatment		als valid for 6	months where the treatment
MEBENDAZOLE - Only on a prescription			
Tab 100 mg	7.97	6	✓ Vermox
Vermox to be Principal Supply on 1 January 2022 Oral lig 100 mg per 5 ml	2 18	15 ml	
	(7.53)	10 1111	Vermox
PRAZIQUANTEL	, ,		
Tab 600 mg	68.00	8	✓ Biltricide
Antibacterials			
a) For topical antibacterials, refer to DERMATOLOGICALS, page b) For anti-infective eye preparations, refer to SENSORY ORGAN			
Cephalosporins and Cephamycins			
CEFACLOR MONOHYDRATE			
Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor ✓ Ranbaxy-Cefaclor S29 S29
Grans for oral liq 125 mg per 5 ml - Wastage claimable	3.53	100 ml	✓ <u>Ranbaxy-Cefaclor</u> ✓ Ranbaxy-Cefaclor S29 S29
CEFALEXIN			
Cap 250 mg		20	✓ Cephalexin ABM
Cap 500 mgGrans for oral liq 25 mg per ml – Wastage claimable		20 100 ml	✓ Cephalexin ABM✓ Cefalexin Sandoz
Grans for Graning 25 mg per mir – wastage Gallinable		100 1111	- OcialexIII Januoz

CEFTRIAXONE -	Subsidy by endorsement
---------------	------------------------

CEFAZOLIN - Subsidy by endorsement

accordingly.

Grans for oral liq 50 mg per ml – Wastage claimable......11.75

- a) Up to 10 inj available on a PSO
- b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed

Inj 500 mg vial	0.89	1	✓ <u>Ceftriaxone-AFT</u>
Inj 1 g vial	3.99	5	✓ <u>Ceftriaxone-AFT</u>

100 ml

5

✓ Cefalexin Sandoz

✓ AFT

Fully

Brand or

✓ 7ithromax

15 ml

(Manufactur \$	Pe	Subsidise r •	d Generic Manufacturer	
CEFUROXIME AXETIL — Subsidy by endorsement Only if prescribed for prophylaxis of endocarditis and the prescription is a Tab 250 mg		0,	Zinnat	

Subsidy

Macrolides

C

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Apo-Azithromycin
	2.57	_	✓ Zithromax
Zithromax to be Sole Supply on 1 December 2021			
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			

(Apo-Azithromycin Tab 250 mg to be delisted 1 May 2022) (Apo-Azithromycin Tab 500 mg to be delisted 1 December 2021)

claimable

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

	Subsidy (Manufacturer's Price) \$	Subsidi	ully ised	Brand or Generic Manufacturer
CLARITHROMYCIN – Maximum of 500 mg per prescription; can l Tab 250 mg	, ,		√	AA1857 below Apo-Clarithromycin Klacid
Klacid to be Sole Supply on 1 February 2022 Grans for oral liq 250 mg per 5 ml — Wastage claimable (Apo-Clarithromycin Tab 250 mg to be delisted 1 February 2022)	192.00	50 ml	✓ k	Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- - 1 Atypical mycobacterial infection: or

ERYTHROMYCIN (AS LACTOBIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO			•
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
 a) Up to 300 ml available on a PSO 			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable	0.77	100	✓ E Musin
Grans for oral liq 400 mg per 5 ml	6.//	100 ml	E-Mycin
a) Up to 200 ml available on a PSOb) Wastage claimable			
, •			
ERYTHROMYCIN STEARATE	44.05	400	
Tab 250 mg – Up to 30 tab available on a PSO	(22.29)	100	ERA
Tab 500 mg	(- /	100	LINA
rab 600 mg	(44.58)	100	ERA
(ERA Tab 250 mg to be delisted 1 April 2022)	(******)		
(ERA Tab 500 mg to be delisted 1 September 2022)			
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-
			<u>Roxithromycin</u>
Tab 300 mg	16.33	50	✓ Arrow-
1 ab 000 mg	10.00	50	Roxithromycin

	Subsidy		Fully	Brand or
	(Manufacturer's F	Price) Subs Per	sidised	Generic Manufacturer
	Ψ	rei	<u> </u>	Manulacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500		<u>Alphamox</u>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500		<u>Alphamox</u>
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP	1.40	100 ml	./	Alphamay 10E
Grans for oral liq 125 mg per 5 ml	1.40	100 1111		Alphamox 125
a) Up to 200 ml available on a PSOb) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	1 73	100 ml	1	Alphamox 250
a) Up to 300 ml available on a PSO	1.73	100 1111	• !	Aiphamox 230
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15.97	10	1	biamox
Inj 500 mg vial		10	✓	biamox
Inj 1 g vial - Up to 5 inj available on a PSO	21.64	10	✓	biamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab				
available on a PSO	0.89	10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25				
per ml		100 ml	1	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5				
per ml – Up to 200 ml available on a PSO	2.20	100 ml OP		Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	344.93	10	✓	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial - Up to 5 inj available on a P	SO 11.09	10	1	<u>Sandoz</u>
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO	16.83	250		Staphlex
Cap 500 mg - Up to 30 cap available on a PSO	56.61	500		Staphlex
Grans for oral liq 25 mg per ml	3.29	100 ml		AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
c) AFT to be Principal Supply on 1 January 2022	0.60	100 ml	./	AFT
Grans for oral liq 50 mg per ml	3.00	100 1111	•	AFI
b) Wastage claimable				
c) AFT to be Principal Supply on 1 January 2022				
Inj 250 mg vial	17.56	10	✓ I	Flucloxin
Inj 500 mg vial		10		Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	✓ [Flucil

 $[\]blacktriangle \textit{Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. }$

	Subsidy	_	Fully	
	(Manufacturer's Price)) Si Per	ubsidised	Generic Manufacturer
	Ψ	1 01		Marialacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)	0.04	50	,	Ollin alma VIV
Cap 250 mg – Up to 30 cap available on a PSO	3.84	50	•	Cilicaine VK
Cilicaine VK to be Principal Supply on 1 January 2022 Cap 500 mg	6 96	50	J	Cilicaine VK
a) Up to 20 cap available on a PSO	0.00	50	•	Cilicalite VK
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Cilicaine VK to be Principal Supply on 1 January 202.	2			
Grans for oral lig 125 mg per 5 ml		100 ml	1	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	✓	<u>AFT</u>
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO	123.50	5	1	Cilicaine
Tetracyclines				
DOXYCYCLINE				
* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	✓	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
	(12.05)			Mino-tabs
* Cap 100 mg	19.32	100		
	(52.04)			Minomycin
OA4055 On a lat Audio with family and a strong Bullet				

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE – Special Authority see SA1332 below – Retail pharmacy
Tab 250 mg21.42 28

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Subsidy (Manufacturer's		Fully	Brand or Generic	
\$	Per	√	Manufacturer	

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 63

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg – Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	✓ Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓ Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – St Only if prescribed for dialysis or cystic fibrosis patient and the Inj 150 mg	prescription is er		ordingly. ✓ Colistin-Link
GENTAMICIN SULPHATE			
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	95.00	5	✓ DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient o endorsed accordingly.	r complicated uri	nary tract inf	ection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient o	r complicated uri	nary tract inf	ection and the prescription is

endorsed accordingly. 10 ✓ Pfizer

87.50 50 ✓ Pfizer Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.

MOXIFLOXACIN - Special Authority see SA1740 below - Retail pharmacy

No patient co-payment payable

5 Avelox

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or

continued...

	Subsidy (Manufacturer's Price) \$	S Per	Fully subsidised	Brand or Generic Manufacturer
continued				
 Significant documented intolerance and or 	or side effects following	a reas	onable tria	I of first-line medications;
2 Mycobacterium avium-intracellulare complex not responsible.3 Patient is under five years of age and has had close or				• •
Note: Indications marked with * are unapproved indications. Renewal only from a respiratory specialist or infectious disea	ea enacialist Annrovals	valid fo	r 1 voor w	where the treatment
remains appropriate and the patient is benefiting from treatme		vallu it	n i yeai w	mere me neamem
Initial application — (Mycoplasma genitalium) only from a sexual health specialist. Approvals valid for 1 month for appli All of the following:	sexual health specialist			the recommendation of a
1 Has nucleic acid amplification test (NAAT) confirmed I2 Either:	Mycoplasma genitalium*	and is	symptoma	tic; and
2.1 Has tried and failed to clear infection using azit2.2 Has laboratory confirmed azithromycin resistar				
3 Treatment is only for 7 days.	a u latin a lus a l a sui a t		: -	
Initial application — (Penetrating eye injury) only from an requires prophylaxis following a penetrating eye injury and tre Note: Indications marked with * are unapproved indications.			ia for 1 ma	onth where the patient
PAROMOMYCIN - Special Authority see SA1689 below - R	etail pharmacy			
Cap 250 mg	126.00	16	√ F	lumatin S29
■ SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, month for applications meeting the following criteria: Either: 1 Patient has confirmed cryptosporidium infection; or	clinical microbiologist or	gastroe	enterologis	st. Approvals valid for 1
2 For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical mapplications meeting the following criteria:	icrobiologist or gastroent	erologi	st. Appro	vals valid for 1 month for
Either:				
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 				
PYRIMETHAMINE – Special Authority see SA1328 below –	Retail pharmacy			
Tab 25 mg		30	✓ [Daraprim \$29
■ SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals the following criteria: Any of the following:				•
 For the treatment of toxoplasmosis in patients with HIV For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 mon 	, '	s; or		
SODIUM FUSIDATE [FUSIDIC ACID]	67.05	0.0		···aidia

36

56

✓ Fucidin

✓ Wockhardt S29

Tab 250 mg67.85

Tab 500 mg543.20

SULFADIAZINE SODIUM - Special Authority see SA1331 on the next page - Retail pharmacy

	INFECTIONS -	- AGENTS	FOR S	SYSTEMIC USE
	Subsidy (Manufacturer's Pr \$	rice) Sub Per	Fully sidised	Brand or Generic Manufacturer
SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals the following criteria: Any of the following:	s valid without further r	renewal unles	s notifie	d for applications meeting
 1 For the treatment of toxoplasmosis in patients with H 2 For pregnant patients for the term of the pregnancy; 3 For infants with congenital toxoplasmosis until 12 mo 	or	onths; or		
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial — Subsidy by endorsement a) Only if prescribed for dialysis or cystic fibrosis pa b) Tobramycin Mylan to be Principal Supply on 1 J	atient and the prescrip	5 ition is endors		obramycin Mylan ordingly.
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by endorsement		56 dose	_	obramycin BNM
TRIMETHOPRIM * Tab 300 mg - Up to 30 tab available on a PSO TMP to be Principal Supply on 1 January 2022	18.55	50	✓ T	MP
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRII * Tab trimethoprim 80 mg and sulphamethoxazole 400 mg to 30 tab available on a PSO Trisul to be Principal Supply on 1 January 2022 * Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to available on a PSO	g – Up 64.80 200 ml	500 100 ml		risul Deprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient difficile following metronidazole failure and the prescription in 500 mg vial	ion is endorsed accord	ndocarditis or dingly. 1	for trea	•
Antifungals				
a) For topical antifungals refer to DERMATOLOGICALS, pa b) For topical antifungals refer to GENITO URINARY, page				
FLUCONAZOLE Cap 50 mg Cap 150 mg	0.65	28 1 28	✓ N	lylan Iylan Iylan

30011112022			
Cap 50 mg	2.75	28	Mylan
Cap 150 mg		1	✓ Mylan
Cap 200 mg		28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Author	ority		
see SA1359 below - Retail pharmacy		35 ml	✓ Diflucan
Wastage claimable			

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised: and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOI F

Cap 100 mg4	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 below -			
Retail pharmacy141	1.80	150 ml OP	✓ Sporanox

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare \$29
•			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81 [°]	50	
,	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Ret	tail pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml		105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation

Subsidy	Subsidy Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

chemotherapy; or

2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 below - Retail pharmacy		
Tab 50 mg91.00	56	✓ Vttack
Tab 200 mg350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage		
claimable1,437.00	70 ml	✓ Vfend

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 on the n	next page - Retail pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaguine \$29

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE

Tab 200 mg — Up to 30 tab available on a PSO	5.23 25.00	250 21 100 ml 10	✓ <u>Metrogyl</u> ✓ <u>Metrogyl</u> ✓ Flagyl-S ✓ Flagyl
	24.40	10	Flagyi
ORNIDAZOLE Tab 500 mg	26.16	10	✓ Arrow-Ornidazole
Arrow-Ornidazole to be Principal Supply on 1 December		10	Allow-Officazole
Autom Offinaazolo to be i intolpat cappily on i Bosomber	2021		
Antituberculotics and Antileprotics			
Note: There is no co-payment charge for all pharmaceuticals liste	ed in the Antitube	erculotics and	Antileprotics group regardless of
immigration status.	a iii aio 7 ii iiiao	rodiotioo dila	, introprotion group regulations of
CLOFAZIMINE - Retail pharmacy-Specialist			
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendation dermatologist.	on of, an infection	us disease ph	ysician, clinical microbiologist or
* Cap 50 mg	442 00	100	✓ Lamprene S29
CYCLOSERINE – Retail pharmacy-Specialist		100	- Lumprono
a) No patient co-payment payable			
b) Prescriptions must be written by, or on the recommendation	on of, an infection	us disease ph	ysician, clinical microbiologist or
respiratory physician. Cap 250 mg	344.00	60	✓ Cyclorin S29
	344.00	00	• Cyclorin •
DAPSONE – Retail pharmacy-Specialist			
a) No patient co-payment payableb) Prescriptions must be written by, or on the recommendation	on of an infaction	uc dicasca nh	veisian elipical microbiologist or
dermatologist	on or, an intection	us disease pi	rysician, clinical microbiologist of
Tab 25 mg	268.50	100	✓ Dapsone
Tab 100 mg	329.50	100	✓ Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist	t		
a) No patient co-payment payable			
 b) Prescriptions must be written by, or on the recommendation respiratory physician 	on of, an infection	us disease ph	ysician, clinical microbiologist or
Tab 100 mg	85.73	100	✓ EMB Fatol S29
Tab 400 mg		56	✓ Myambutol S29

INFECTIONS - AGENTS FOR SYSTEMIC USE					
	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
ISONIAZID - Retail pharmacy-Specialist					
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician 		dicine ph	nysician,	paediatrician, clinical	
* Tab 100 mg PSM to be Principal Supply on 1 January 2022	23.00	100	√ F	PSM	
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist					
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician 	tion of, an internal me	dicine ph	nysician,	paediatrician, clinical	
* Tab 100 mg with rifampicin 150 mg Rifinah to be Principal Supply on 1 January 2022		100	√ F	Rifinah	
* Tab 150 mg with rifampicin 300 mg Rifinah to be Principal Supply on 1 January 2022	179.13	100	√ F	Rifinah	
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist					
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician				v	
Grans for oral liq 4 g sachet	280.00	30	✓ F	Paser S29	
PROTIONAMIDE – Retail pharmacy-Specialist					
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician 	tion of, an infectious d	isease s	pecialist,	clinical microbiologist or	
Tab 250 mg	305.00	100	✓ F	Peteha S29	
PYRAZINAMIDE - Retail pharmacy-Specialist					
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician 	tion of, an infectious d	isease p	hysician,	clinical microbiologist or	
* Tab 500 mg	59.00	100	✓ A	AFT-Pyrazinamide	
RIFABUTIN - Retail pharmacy-Specialist					
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda gastroenterologist	tion of, an infectious d	isease p	hysician,	respiratory physician or	
* Cap 150 mg	299.75	30	✓ N	/lycobutin	
RIFAMPICIN – Subsidy by endorsement					
 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescripti Retail pharmacy - Specialist. Specialist must be an inte paediatrician, or public health physician. 	on is endorsed accord rnal medicine physicia	ingly; ca n, clinica	n be waiv al microbi	ved by endorsement - lologist, dermatologist,	
* Cap 150 mg		100	_	Rifadin	
* Cap 300 mg		100	_	Rifadin Rifadin	
* Oral liq 100 mg per 5 ml	12.60	60 ml	A F	<u>Rifadin</u>	

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Dox /	Manufacturer

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, page 239

Hepatitis B Treatment

ENTECAVIR * Tab 0.5 mg	52.00	30	✓ Entecavir Sandoz
LAMIVUDINE - Special Authority see SA1685 below - Reta	il pharmacy		
Tab 100 mg	6.95	28	✓ Zetlam
Oral liq 5 mg per ml	270.00	240 ml OP	✓ Zeffix

⇒SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B. TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., page 107

* Tab 245 mg (300.6 mg as a succinate).......38.10 30 ✓ Tenofovir Disoproxil Teva

Herpesvirus Treatments

ACICLOVIR		
* Tab dispersible 200 mg1.60	25	✓ <u>Lovir</u>
* Tab dispersible 400 mg5.38	56	✓ Lovir
* Tab dispersible 800 mg5.98	35	✓ <u>Lovir</u>
VALACICLOVIR		
Tab 500 mg6.50	30	✓ Vaclovir
Vaclovir to be Principal Supply on 1 January 2022		
Tab 1,000 mg13.76	30	✓ Vaclovir
Vaclovir to be Principal Supply on 1 January 2022		
VALGANCICLOVIR - Special Authority see SA1993 below - Retail pharmacy		
Tab 450 mg132.00	60	✓ Valganciclovir
		Mylan

Valganciclovir Mylan to be Principal Supply on 1 December 2021

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
- 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
 - 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 on the next page

No patient co-payment payable

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP). Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/mayiret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 107 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

succinate)61.15

✓ Teva

⇒SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks;
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Antiretrovirals

⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Subsidy	Fully Subsidised		Brand or
(Manufacturer's Price)			Generic
\$	Per	✓	Manufacturer

continued...

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (**Percutaneous exposure**) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previou	s page – Retail pharr	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	us page – Retail phai	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	us page – Retail phai	rmacy	
Tab 200 mg	84.00	60	NevirapineAlphapharm
Nevirapine Alphapharm to be Principal Supply on 1	January 2022		лірпарпатп
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune Suspension

	Subsidy (Manufacturer's \$	Price) Subsi	Fully Brand or dised Generic ✓ Manufacturer
Nucleosides Reverse Transcriptase Inhibitors			
ABACAVIR SULPHATE – Special Authority see SA1651 on page Tab 300 mg Oral liq 20 mg per ml	180.00	oharmacy 60 240 ml OP	✓ <u>Ziagen</u> ✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE — Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil co anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoprox	unts as three a	anti-retroviral med	dications for the purposes of the
245 mg (300 mg as a maleate)		30	✓ <u>Mylan</u>
EMTRICITABINE – Special Authority see SA1651 on page 107 – Cap 200 mg	307.20	30 30	✓ Emtriva
LAMIVUDINE – Special Authority see SA1651 on page 107 – Re Tab 150 mg	, ,	60	✓ <u>Lamivudine</u> Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 10. Cap 100 mg Oral liq 10 mg per ml	152.25	macy 100 200 ml OP	✓ Retrovir ✓ Retrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets) the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg	counts as two	•	•
Protease Inhibitors			
ATAZANAVIR SULPHATE – Special Authority see SA1651 on pa	nge 107 – Reta	il pharmacy	
Cap 150 mg	•	60	✓ <u>Teva</u>
Cap 200 mg	188.91	60	✓ <u>Teva</u>
DARUNAVIR - Special Authority see SA1651 on page 107 - Ret			
Tab 400 mg		60 60	 ✓ <u>Darunavir Mylan</u> ✓ Darunavir Mylan
Tab 600 mg			
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 c Tab 100 mg with ritonavir 25 mg		60	✓ Lopinavir/Ritonavir Mylan
Lopinavir/Ritonavir Mylan to be Principal Supply on 1 Feb Tab 200 mg with ritonavir 50 mg	•	120	✓ Kaletra✓ Lopinavir/Ritonavir
	462.00		Mylan Kolotro
Lopinavir/Ritonavir Mylan to be Principal Supply on 1 Feb Oral liq 80 mg with ritonavir 20 mg per ml	735.00 y 2022)	300 ml OP	✓ Kaletra ✓ Kaletra

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RITONAVIR – Special Authority see SA1651 on page 107 – Ret Tab 100 mg		30	✓	<u>Norvir</u>
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 107 Tab 50 mg		30	✓.	Tivicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 o Tab 400 mg Tab 600 mg	1,090.00	harm 60 60	/	Isentress Isentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

continued...

- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
 - Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

	-	,	4.	0.04	400	
*	Tab 1 g		4	0.01	100	Hiprex

METHENAMINE (HEXAMINE) HIPPURATE

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
NITROFURANTOIN				
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓	Nifuran
* Tab 100 mg	37.50	100	✓	Nifuran
* Cap modified-release 100 mg - Wastage claimable	86.40	100	1	Macrobid
NORFLOXACIN				
Tab 400 mg - Subsidy by endorsement	135.00	100	✓	Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated unwith proven resistance to first line agents and the prescribed for a patient with proven resistance to first line agents.				ve to a first line agent or

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
Anticholinesterases			
NEOSTIGMINE METILSULFATE			
	10.60	10	A lune coo
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓ Juno ©29 ✓ Max Health
		EΛ	✓ Max nealth ✓ AstraZeneca
	98.00	50	Astrazeneca
(Juno S29 Inj 2.5 mg per ml, 1 ml ampoule to be delisted 1 Marc			
(AstraZeneca Inj 2.5 mg per ml, 1 ml ampoule to be delisted 1 Ma	arch 2022)		
PYRIDOSTIGMINE BROMIDE			
▲ Tab 60 mg	45.79	100	✓ <u>Mestinon</u>
Non-Steroidal Anti-Inflammatory Drugs			
DICLOFENAC SODIUM			4.50.4
* Tab EC 25 mg		50	Diclofenac Sandoz
Diclofenac Sandoz to be Principal Supply on 1 January 2			
* Tab 50 mg dispersible		20	✓ Voltaren D
* Tab EC 50 mg		50	Diclofenac Sandoz
Diclofenac Sandoz to be Principal Supply on 1 January 2			
* Tab long-acting 75 mg		100	
	22.80	500	
* Tab long-acting 100 mg		500	
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5	✓ Voltaren
* Suppos 12.5 mg		10	✓ Voltaren
* Suppos 25 mg		10	✓ Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10	✓ Voltaren
* Suppos 100 mg	7.00	10	✓ Voltaren
(Apo-Diclo SR Tab long-acting 75 mg to be delisted 1 May 2022)			
(Apo-Diclo SR Tab long-acting 100 mg to be delisted 1 May 2022	?)		
IBUPROFEN			
* Tab 200 mg	21.40	1.000	○ ✓ Relieve
* Tab long-acting 800 mg		30	✓ Brufen SR
	5.99		✓ Ibuprofen SR BNM
Brufen SR to be Principal Supply on 1 January 2022			
* Oral lig 20 mg per ml	2.25	200 m	nl ✓ Ethics
(Ibuprofen SR BNM Tab long-acting 800 mg to be delisted 1 January			
KETOPROFEN	/		
	10.07	20	✓ Oruvail SR
* Cap long-acting 200 mg	12.07	28	Viuvaii 3n
MEFENAMIC ACID			
* Cap 250 mg		50	
	(9.16)		Ponstan
	0.50	20	
	(5.60)		Ponstan

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
NAPROXEN				
* Tab 250 mg Noflam 250 to be Principal Supply on 1 January 203		500	/	Noflam 250
* Tab 500 mg Noflam 500 to be Principal Supply on 1 January 20		250	1	Noflam 500
* Tab long-acting 750 mg Naprosyn SR 750 to be Principal Supply on 1 January	6.47	28	✓	Naprosyn SR 750
* Tab long-acting 1 g Naprosyn SR 1000 to be Principal Supply on 1 Jani	8.62	28	✓	Naprosyn SR 1000
SULINDAC	,			
* Tab 100 mg	9.57	56	1	Mylan S29
* Tab 200 mg		50		Aclin
-	16.91	56	1	Sulindac Mylan S29
, ,	2022)			
(Sulindac Mylan ©29 Tab 200 mg to be delisted 1 January TENOXICAM * Tab 20 mg	9.15	100 1		<u>Tilcotil</u> AFT
(Aclin Tab 200 mg to be delisted 1 January 2022) (Sulindac Mylan 23 Tab 200 mg to be delisted 1 January TENOXICAM * Tab 20 mg * Inj 20 mg vial NSAIDs Other	9.15			
(Sulindac Mylan 229 Tab 200 mg to be delisted 1 January TENOXICAM * Tab 20 mg * Inj 20 mg vial	9.15 9.95 5.80		<i>y</i>	
(Sulindac Mylan S29 Tab 200 mg to be delisted 1 January TENOXICAM * Tab 20 mg * Inj 20 mg vial NSAIDs Other CELECOXIB Cap 100 mg		1 60	<i>y</i>	AFT Celecoxib Pfizer Celebrex
Sulindac Mylan S29 Tab 200 mg to be delisted 1 January TENOXICAM * Tab 20 mg * Inj 20 mg vial NSAIDs Other CELECOXIB Cap 100 mg Cap 200 mg		1 60	<i>y</i>	AFT Celecoxib Pfizer Celebrex

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of

hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

	nyuroxychioroquine. Note. mulcation marked with a	is an unapproved indication	l.	
*	Tab 200 mg	7.98	100	✓ Plaquenil
LE	FLUNOMIDE			
	Tab 10 mg	6.00	30	✓ Arava
	Tab 20 mg	6.00	30	✓ Arava
PE	NICILLAMINE			
	Tab 125 mg	67.23	100	D-Penamine
	Tab 250 mg		100	✓ D-Penamine

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per
Manufacturer

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

* Tab 70 mg	2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL			

Other Treatments

AL ENDOONATE CODULA

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither:
 - 2.1 The patient is female and postmenopausal: or

- 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a

continued...

✓ Fosamax Plus

✓ fully subsidised

Principal Supply

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
\$	Per	1	Manufacturer

continued...

fall from a standing height or less

- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial	27.53	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial		1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA1779		pharmacy	
* Tab 60 mg	53.76	28	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM		
Tab 35 mg3.10	4	✓ Risedronate Sandoz
TERIPARATIDE - Special Authority see SA1139 on the next page - Retail phar	macy	
Inj 250 mcg per ml, 2.4 ml490.00	1	✓ Forteo

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
S Per ✔ Manufacturer

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

SA1780 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically: or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or

(Ma	Subsidy anufacturer's Price)	Subsidi Per	ully	Brand or Generic Manufacturer
	<u></u>	Per		Manufacturer

continued...

- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

- 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA	A1963 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite S29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	 Benzbromaron AL
			100 \$29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment: and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests

COLCHICINE

* Tab 5	500 mca	9.58	100	✓ Colgout
---------	---------	------	-----	-----------

	Subsidy (Manufacturer's Price)		Fully	Brand or Generic	
	\$	Per		Manufacturer	
FEBUXOSTAT - Special Authority see SA2054 below - Retail ph	narmacy				
Tab 80 mg	•	28	✓ F	ebuxostat	
· · · · · · · · · · · · · · · · · ·				multichem	
	39.50		✓ A	denuric	
Febuxostat multichem to be Sole Supply on 1 January 20					
Tab 120 mg		28	✓ F	ebuxostat	
145 125 Hg				multichem	
	39.50		./ ^	denuric	
			▼ A	denunc	
Febuxostat multichem to be Sole Supply on 1 January 20	122				
(Adenuric Tab 80 mg to be delisted 1 January 2022)					

⇒SA2054 Special Authority for Subsidy

Initial application — (**Gout**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Patient has been diagnosed with gout; and

(Adenuric Tab 120 mg to be delisted 1 January 2022)

- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PR	DBENECID		
*	Tab 500 mg55.00	100	✓ Probenecid-AFT

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
Muscle Relaxants					

BAG	CLOFEN			
*	Tab 10 mg	4.20	100	✓ Pacifen
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
	Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo		, ,	nts have been ineffective or have
	Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	306.82	5	✓ Medsurge
	a) Subsidised only for use in a programmable pump in pat have caused intolerable side effects and the prescriptiob) Medsurge to be Principal Supply on 1 December 2021			agents have been ineffective or
DAI	NTROLENE			
	Cap 25 mg	97.50	100	✓ Dantrium
				✓ Dantrium S29 S29
	Cap 50 mg	77.00	100	✓ Dantrium
OR	PHENADRINE CITRATE			
	Tab 100 mg	20.76	100	✓ Norflex

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

Dopamino Agomoto una Holatoa Agomo			
AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
BROMOCRIPTINE MESYLATE – Subsidy by endorsement			
Subsidy by endorsement - Subsidised for patients who were	e taking bromocrip	tine mesylate	e prior to 1 March 2021 and the
prescription is endorsed accordingly. Pharmacists may ann	notate the prescript	ion as endor	sed where there exists a record of
prior dispensing of bromocriptine mesylate.			
★ Tab 2.5 mg	11.70	30	✓ Parlodel S29
	32.08	100	Apo-Bromocriptine
Parlodel S29 Tab 2.5 mg to be delisted 1 March 2022)			
Apo-Bromocriptine Tab 2.5 mg to be delisted 1 March 2022)			
ENTACAPONE			
▲ Tab 200 mg	18.04	100	✓ Comtan
	22.00		✓ Entapone
Entapone Tab 200 mg to be delisted 1 April 2022)			
EVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	Madopar Rapid
Cap 50 mg with benserazide 12.5 mg		100	Madopar 62.5
★ Cap 100 mg with benserazide 25 mg		100	✓ Madopar 125
★ Cap long-acting 100 mg with benserazide 25 mg		100	✓ Madopar HBS
★ Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
EVODOPA WITH CARBIDOPA			
★ Tab 100 mg with carbidopa 25 mg		100	✓ Sinemet
★ Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
★ Tab 250 mg with carbidopa 25 mg	38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	6.12	100	✓ Ramipex
▲ Tab 1 mg	20.73	100	✓ Ramipex
RASAGILINE			
★ Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.85	84	✓ Ropin
	3.39	100	✓ Mylan S29
▲ Tab 1 mg		84	✓ Ropin

4.70

100

84

✓ Mylan S29✓ Ropin

✓ Ropin

▲ Tab 5 mg12.50



NEITVOOS STOTEM				
	Subsidy (Manufacturer's \$		osidised	Brand or Generic Manufacturer
	ed for patients who were taking selegili Pharmacists may annotate the presc			
* Tab 5 mg		100	•	o-Selegiline 29 S29
(Apo-Selegiline S29 S29 Tab 5 mg to be	48.00 e delisted 1 April 2022)		✓ Eld	epryl S29
TOLCAPONE Tab 100 mg	152.38	100	✓ Tas	mar
Anticholinergics				
BENZATROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 10 inj available on a PS b) Only on a PSO	95.00	60 5	✓ Ber ✓ <u>Phe</u>	•
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ Ker	nadrin
Agents for Essential Tremor,	Chorea and Related Disorder	rs		
RILUZOLE – Special Authority see SA14 Wastage claimable Tab 50 mg Rilutek to be Principal Supply on	130.00	56	✓ Rilu	utek
■ SA1403 Special Authority for Subsi Initial application only from a neurologis following criteria: All of the following:		alid for 6 month	ns for applic	cations meeting the
1 The patient has amyotrophic latera	•			tial application; and
5.1 The patient is ambulatory;5.2 The patient is able to use to 5.3 The patient is able to swall	upper limbs; or			
Renewal from any relevant practitioner. All of the following:		ications meeting	g the follow	ing criteria:
 1 The patient has not undergone a t 2 The patient has not experienced r 3 Any of the following: 3.1 The patient is ambulatory; 	espiratory failure; and			
3.2 The patient is able to use us. 3.3 The patient is able to swall	upper limbs; or			
TETRABENAZINE				

Tab 25 mg91.10

✓ Motetis

112

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	/	Manufacturer

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]			
Gel 2%, tube - Subsidy by endorsement	14.50	30 ml	Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO			
b) Subsidised only if prescribed for urethral or cervical adm	ninistration an	d the prescript	tion is endorsed accordingly.
Gel 2%, 11 ml urethral syringe - Subsidy by endorsement	42.00	10	✓ Instillagel Lido
 a) Up to 5 each available on a PSO 			
 Subsidised only if prescribed for urethral, cervical or rec accordingly. 	tal administra	tion and the p	rescription is endorsed
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE			
Oral (gel) soln 2%	38.00	200 ml	✓ Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	✓ Lidocaine-Baxter
			✓ Lidocaine-Claris
	17.50	50	
	(35.00)		Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO	8.25	25	✓ <u>Lidocaine-Baxter</u>
1:40/ 00 1 1 1 1 5:1	10.00	_	✓ Lidocaine-Claris
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO		5	W. Landina
lai 40/ 00 ant viet. Ha to 5 ini available on a DCO	(20.00)	-	Xylocaine
Inj 1%, 20 ml vial — Up to 5 inj available on a PSO		5 5	✓ <u>Lidocaine-Claris</u> ✓ Lidocaine-Baxter
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO	0.45	5	✓ Lidocaine-Baxter ✓ Lidocaine-Claris
// ideasing Clarie Ini 19/ E ml ampaule to be delicted 1 January 200	0)		▼ Liuocaine-Ciaris
(Lidocaine-Claris Inj 1%, 5 ml ampoule to be delisted 1 January 202	,		
(Lidocaine-Claris Inj 2%, 5 ml ampoule to be delisted 1 January 202	2)		
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE			
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –			•
Subsidy by endorsement	103.32	10	✓ Pfizer
 a) Up to 5 each available on a PSO 			

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority see SA09	06 above – Retail pharm	acy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - Special	al Authority see SA0906 a	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓ EMLA



Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 114

Non-o	bioid	Anal	lgesics
			9

ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabetic accordingly.	periphera	I neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE			
Tab 30 mg	23.40	90	✓ Acupan

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	5	Subsidised	Generic
	` \$	Per	•	Manufacturer
PARACETAMOL				
Tab 500 mg - blister pack	0.50	20	✓ N	ledco
• •			✓ P	harmacy Health
	1.12			thics Paracetamol
	2			Classic
	2.48	100	✓ P	harmacy Health
	5.01	50	✓ P	anadol
	11.75	96	✓ P	anadol Mini Caps
	19.75	1.000		acimol .
	24.82	,	√ P	aracetamol
	2 2			Pharmacare
			√ P	harmacare

- a) Maximum of 300 tab per prescription; can be waived by endorsement
- b) Up to 30 tab available on a PSO

c)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
 regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
 annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.
- d) Pacimol to be Sole Supply on 1 February 2022

Tab 500 mg - bottle pack - Maximum of 300 tab per			
prescription; can be waived by endorsement	17.92	1,000	✓ Noumed Paracetamol
	24.82		✓ Paracetamol Pharmacare

a)

- Subsidy by endorsement for higher quantities is available for patients with long term conditions who require
 regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may
 annotate the prescription as endorsed where dispensing history supports a long-term condition.
- Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.
 Noumed Paracetamol to be Sole Supply on 1 February 2022

*	Oral liq 120 mg per 5 ml	1,000 ml	✓ Paracare
	a) Up to 200 ml available on a PSO b) Not in combination		
*	Oral liq 250 mg per 5 ml	5 1,000 ml	✓ Paracare Double Strength
	a) Up to 100 ml available on a PSO		
	b) Not in combination		
*	Suppos 125 mg	10	✓ Gacet
*	Suppos 250 mg	10	✓ Gacet

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
* Suppos 500 mg	2022) February 2022) delisted 1 February 2022) February 2022) y 2022) 1 December 2021) lelisted 1 February 2022) ruary 2022)	50	•	Gacet

Opioid Analgesics

CODEINE PHOSPHATE - Safety medicine; prescriber may determi	ne dispensin	g frequency	
Tab 15 mg	6.25	100	✓ PSM
Tab 30 mg	7.45	100	✓ PSM
Tab 60 mg	14.25	100	✓ PSM
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	✓ DHC Continus
FENTANYL			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing frequent	encv		
Inj 50 mcg per ml, 2 ml ampoule		10	✓ Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		10	✓ Boucher and Muir
Patch 12.5 mcg per hour		5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 2022		Ü	· · · · · · · · · · · · · · · · · · ·
Patch 25 mcg per hour	7.99	5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 2022		-	
Patch 50 mcg per hour	9.49	5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 2022		-	,
Patch 75 mcg per hour	17.99	5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 2022			, ,
Patch 100 mcg per hour	18.59	5	✓ Fentanyl Sandoz
Fentanyl Sandoz to be Principal Supply on 1 January 2022			•
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing frequency	encv		
d) Extemporaneously compounded methadone will only be reim		e rate of the ch	neapest form available
(methadone powder, not methadone tablets).			
e) For methadone hydrochloride oral liquid refer Standard Form	ulae, page 24	1 7	
Tab 5 mg		10	✓ Methatabs
Oral lig 2 mg per ml		200 ml	✓ Biodone
Biodone to be Principal Supply on 1 January 2022			
Oral liq 5 mg per ml	6.40	200 ml	✓ Biodone Forte
Biodone Forte to be Principal Supply on 1 January 2022			
			_

Biodone Extra Forte to be Principal Supply on 1 January 2022 Inj 10 mg per ml, 1 ml61.00

✓ Biodone Extra Forte

✓ AFT

200 ml

10

NERVOUS SYSTEM

	Subsidy	. ,		nd or
	(Manufacturer's Pr \$	rice) Sub Per		neric nufacturer
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fro	equency			
Oral lig 1 mg per ml	9.28	200 ml	✓ RA-Mo	orph
Oral liq 2 mg per ml	16.24	200 ml	✓ RA-Mo	orph .
Oral liq 5 mg per ml	19.44	200 ml	✓ Ordine	S29
- 4 - 3 F			✓ RA-Mo	orph
Oral lig 10 mg per ml	27 74	200 ml	✓ Ordine	•
		200 1111	✓ RA-Mo	
IORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing from	equency			
Tab immediate-release 10 mg		10	✓ Sevre	dol
Tab immediate-release 20 mg	5.52	10	✓ Sevre	dol
Cap long-acting 10 mg	2.05	10	✓ m-Esle	on
Cap long-acting 30 mg	3.00	10	✓ m-Esle	on
Cap long-acting 60 mg	6.12	10	✓ m-Esle	<u>on</u>
Cap long-acting 100 mg	7.13	10	✓ m-Esle	<u>on</u>
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	SO6.99	5	✓ DBL N	orphine
			Sulp	hate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	SO5.61	5	✓ DBL N	orphine
			Sulp	hate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	SO7.08	5	✓ DBL N	orphine
, 0, , , , , , , , , , , , , , , , , ,				hate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	SO7.28	5	✓ DBL N	
,		-		hate

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
OXYCODONE HYDROCHLORIDE	•			
a) Only on a controlled drug form b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing f	requency			
Tab controlled-release 5 mg		20	✓	Oxycodone Sandoz
	3.01	28	/	Oxycodone Sandoz S29 S29
Tab controlled-release 10 mg	2.15	20	/	Oxycodone Sandoz
	3.23	30		Oxycodone Sandoz S29 S29
	5.38	50	•	Oxycodone Sandoz S29 S29
	10.75	100	•	Oxycodone Sandoz S29 S29
	11.50	28	1	OxyContin
Tab controlled-release 20 mg		20		Oxycodone Sandoz
3	5.38	50		Oxycodone Sandoz S29 S29
	10.75	100	✓	Oxycodone Sandoz S29 S29
	13.25	28	/	OxvContin
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mgOxyNorm to be Principal Supply on 1 December 2021		20		OxyNorm
Cap immediate-release 10 mg OxyNorm to be Principal Supply on 1 December 2021	3.32	20	•	OxyNorm
Cap immediate-release 20 mg OxyNorm to be Principal Supply on 1 December 2021	5.23	20	•	OxyNorm
Oral liq 5 mg per 5 ml	11.20	250 m		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		OxyNorm
Inj 10 mg per ml, 2 ml ampoule		5		OxyNorm
Inj 50 mg per ml, 1 ml ampoule	30.60	5	/	OxyNorm
PARACETAMOL WITH CODEINE - Safety medicine; prescribe	er may determine dispe	ensin	g frequenc	у
* Tab paracetamol 500 mg with codeine phosphate 8 mg	26.51	1,000		Paracetamol + Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form b) No patient co-payment payable				
Safety medicine; prescriber may determine dispensing f Tab 50 mg PSM to be Principal Supply on 1 January 2022		10	•	PSM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a	PSO29.88	5	•	DBL Pethidine Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO30.72	5	✓	DBL Pethidine Hydrochloride

			NERVOUS SYSTEM
	Subsidy (Manufacturer's Price) \$? Per	Fully Brand or Subsidised Generic ✓ Manufacturer
TRAMADOL HYDROCHLORIDE			
Tab sustained-release 100 mg		20	✓ Tramal SR 100
Tab sustained-release 150 mg		20	✓ Tramal SR 150
Tab sustained-release 200 mg		20	✓ Tramal SR 200
Cap 50 mg	2.80	100	✓ <u>Arrow-Tramadol</u>
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE - Safety medicine; prescriber may detern	nine dispensing frequency		
Tab 10 mg	2.49	100	Arrow-Amitriptyline
Tab 25 mg	1.51	100	Arrow-Amitriptyline
Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; p	rescriber may determine d	lispens	sing frequency
Tab 10 mg	10.17	30	Clomipramine Teva
	13.99	100	Apo-Clomipramine
Clomipramine Teva to be Sole Supply on 1 Februa			
Tab 25 mg		100	✓ Apo-Clomipramine
Clomipramine Teva to be Sole Supply on 1 Februa	11.99	30	Clomipramine Teva
(Apo-Clomipramine Tab 10 mg to be delisted 1 February 20 (Apo-Clomipramine Tab 25 mg to be delisted 1 February 20)22)		
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy	by endorsement		
a) Safety medicine; prescriber may determine dispens			
 b) Subsidy by endorsement – Subsidised for patients v 2019 and the prescription is endorsed accordingly. exists a record of prior dispensing of dosulepin [dott 	Pharmacists may annotate		
Tab 75 mg		30	✓ Dosulepin Mylan
Cap 25 mg		50	✓ Dosulepin
, ,			Mylan S29
		noina	frequency
IMIPRAMINE HYDROCHLORIDE - Safety medicine; preso	criber may determine dispe	Hising	
IMIPRAMINE HYDROCHLORIDE – Safety medicine; presonable 10 mg		50	✓ Tofranil
Tab 10 mg	5.48 10.96	50 100	✓ Tofranil ✓ Tofranil
Tab 10 mg	5.48 10.96 8.80	50 100 50	✓ Tofranil ✓ Tofranil ✓ Tofranil
Tab 10 mg	5.48 10.96 8.80	50 100 50	✓ Tofranil ✓ Tofranil ✓ Tofranil
Tab 10 mg	5.48 10.96 8.80 prescriber may determine of	50 100 50	✓ Tofranil ✓ Tofranil ✓ Tofranil
Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE - Safety medicine;	5.48 10.96 880 prescriber may determine of the control	50 100 50 dispens	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency
Tab 10 mg Tab 25 mg NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; Tab 10 mg		50 100 50 dispen:	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency ✓ Norpress
Tab 10 mg		50 100 50 dispen:	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency ✓ Norpress
Tab 10 mg		50 100 50 dispen:	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency ✓ Norpress
Tab 10 mg		50 100 50 dispens 100 180	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency ✓ Norpress ✓ Norpress
Tab 10 mg		50 100 50 dispens 100 180	✓ Tofranil ✓ Tofranil ✓ Tofranil sing frequency ✓ Norpress ✓ Norpress ✓ Parnate S29 S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Monoamine-Oxidase Type A Inhibitors				
MOCLOBEMIDE ★ Tab 150 mg Aurorix to be Principal Supply on 1 January 2022	11.80	60	✓.	Aurorix
* Tab 300 mgAurorix to be Principal Supply on 1 January 2022	19.25	60	✓.	Aurorix
Selective Serotonin Reuptake Inhibitors				
CITALOPRAM HYDROBROMIDE Tab 20 mgPSM Citalopram to be Principal Supply on 1 February 2		84	✓	PSM Citalopram
SCITALOPRAM Tab 10 mg	1.07	28	•	Escitalopram (Ethics)
★ Tab 20 mg	1.92	28	✓	Escitalopram (Ethics)
ELUOXETINE HYDROCHLORIDE Tab dispersible 20 mg, scored – Subsidy by endorsement. Subsidised by endorsement	1.98	30	•	Fluox
 When prescribed for a patient who cannot swallow accordingly; or When prescribed in a daily dose that is not a multiple of the control of the con	iple of 20 mg in which	case	the prescri	ption is deemed to be
accordingly; or	iple of 20 mg in which th capsules to facilitate	case	the prescri emental 10	ption is deemed to be
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	case	the prescri emental 10	otion is deemed to be mg doses.
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg PAROXETINE Tab 20 mg Paxtine Tab 20 mg to be delisted 1 January 2022)	iple of 20 mg in which th capsules to facilitate	case incre 84	the prescri emental 10	otion is deemed to be mg doses.
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg PAROXETINE Tab 20 mg Paxtine Tab 20 mg to be delisted 1 January 2022) EETRALINE Tab 50 mg	iple of 20 mg in which th capsules to facilitate2.91	case incre 84 30	the prescri emental 10	otion is deemed to be mg doses. Fluox Paxtine
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate2.91	30 90	the prescri emental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate2.91	30 90	the prescri emental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90 30	the prescri	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90 30 30 30 30	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona Noumed Apo-Mirtazapine
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 30 30 30 28	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona Noumed Apo-Mirtazapine Noumed
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90 30 30 30 30	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona Noumed Apo-Mirtazapine
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 30 30 30 28	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona Noumed Apo-Mirtazapine Noumed
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90 30 30 30 28 30	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Setrona Noumed Apo-Mirtazapine Noumed Apo-Mirtazapine
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	28 30 28 30 84	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Noumed Apo-Mirtazapine Noumed Apo-Mirtazapine Enlafax XR
accordingly; or 2) When prescribed in a daily dose that is not a mult endorsed. Note: Tablets should be combined wi Cap 20 mg	iple of 20 mg in which th capsules to facilitate	30 90 30 30 30 28 30	the prescriemental 10	otion is deemed to be mg doses. Fluox Paxtine Loxamine Setrona Noumed Apo-Mirtazapine Noumed Apo-Mirtazapine

			NERV	OUS SYSTEM
	Subsidy (Manufacturer's Price) \$	Sul Per	bsidised	Brand or Generic Manufacturer
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
CLONAZEPAM – Safety medicine; prescriber may determine di Inj 1 mg per ml, 1 ml(Rivotril Inj 1 mg per ml, 1 ml to be delisted 1 March 2022)		5	✓ Riv	rotril
DIAZEPAM – Safety medicine; prescriber may determine disper Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement a) Up to 5 inj available on a PSO b) Only on a PSO c) PSO must be endorsed "not for anaesthetic procedu	23.66	5	✓ Hos	spira
Rectal tubes 5 mg - Up to 5 tube available on a PSO		5	✓ Ste	solid
PHENYTOIN SODIUM # Inj 50 mg per ml, 2 ml ampoule — Up to 5 inj available on a l # Inj 50 mg per ml, 5 ml ampoule — Up to 5 inj available on a	PSO 88.63	5	✓ Hos	spira
PSO	133.92	5	✓ Hos	spira
Control of Epilepsy				
CARBAMAZEPINE * Tab 200 mg * Tab long-acting 200 mg * Tab 400 mg * Tab long-acting 400 mg * Oral liq 20 mg per ml CLOBAZAM — Safety medicine; prescriber may determine dispertable 10 mg	16.98 34.58 39.17 26.37	100 100 100 100 250 ml	✓ Teg	gretol CR gretol gretol CR gretol
CLONAZEPAM – Safety medicine; prescriber may determine di Oral drops 2.5 mg per ml		0 ml OP	✓ Riv	otril

100

200 ml

100

100

100

6.45

10.26

✓ Zarontin ✓ Zarontin

✓ Apo-Gabapentin ✓ Nupentin

✓ Apo-Gabapentin

✓ Apo-Gabapentin

✓ Nupentin

✓ Nupentin

ETHOSUXIMIDE

GABAPENTIN

Nupentin to be Sole Supply on 1 February 2022 (Apo-Gabapentin Cap 100 mg to be delisted 1 February 2022)

Oral liq 250 mg per 5 ml56.35

Cap 300 mg......4.07

Note: Not subsidised in combination with subsidised pregabalin

Nupentin to be Sole Supply on 1 February 2022

Nupertin to be Sole Supply on 1 February 2022

(Apo-Gabapentin Cap 300 mg to be delisted 1 February 2022)

(Apo-Gabapentin Cap 400 mg to be delisted 1 February 2022)

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
LACOSAMIDE - Special Authority see SA1125 below - Retail ph	harmacy				
▲ Tab 50 mg	25.04	14	✓ V	'impat	
▲ Tab 100 mg	50.06	14	✓ V	'impat	
•	200.24	56	✓ V	impat '	
▲ Tab 150 mg	75.10	14	✓ V	impat	
v	300.40	56	✓ V	impat	
▲ Tab 200 mg	400.55	56	✓ V	'impat	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

\blacktriangle	Tab dispersible 2 mg55.00	30	✓ Lamictal
\blacktriangle	Tab dispersible 5 mg50.00	30	✓ Lamictal
*	Tab dispersible 25 mg2.76	56	✓ Logem
*	Tab dispersible 50 mg	56	✓ Logem
*	Tab dispersible 100 mg4.40	56	Logem
1F	VETIRACETAM		
	Tab 250 mg4.99	60	✓ Everet
	Tab 500 mg8.79	60	✓ Everet
	Tab 750 mg	60	✓ Everet
	Tab 1,000 mg	60	✓ Everet
	Oral liq 100 mg per ml	300 ml OP	✓ Levetiracetam-AFT
РΗ	ENOBARBITONE		
ГП	For phenobarbitone oral liquid refer Standard Formulae, page 247		
*	Tab 15 mg40.00	500	✓ PSM
*	Tab 30 mg	500	✓ PSM
	-	300	- 1 OW
	ENYTOIN SODIUM	200	(D)
*	Tab 50 mg	200	✓ Dilantin Infatab
	Cap 30 mg	200	✓ Dilantin
	Cap 100 mg	200	✓ Dilantin
*	Oral liq 30 mg per 5 ml	500 ml	✓ Dilantin
PR	EGABALIN		
	Note: Not subsidised in combination with subsidised gabapentin		
*	Cap 25 mg2.25	56	Pregabalin Pfizer
*	Cap 75 mg2.65	56	Pregabalin Pfizer
*	Cap 150 mg4.01	56	✓ Lyrica
			Pregabalin Pfizer
*	Cap 300 mg	56	Pregabalin Pfizer

	Subsidy		Fully	
	(Manufacturer's Pric	,	Subsidised	
	\$	Per		Manufacturer
PRIMIDONE				
* Tab 250 mg	37.35	100	•	Apo-Primidone
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC	52.24	100	✓	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 n	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	•	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail pha	armacy			
Cap 250 mg	509.29	60	•	Diacomit \$29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit \$29

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

A Tele OF man	00	Away Taninamata
▲ Tab 25 mg11.07	60	✓ Arrow-Topiramate
		✓ Topiramate Actavis
26.04		✓ Topamax
▲ Tab 50 mg18.81	60	Arrow-Topiramate
•		✓ Topiramate Actavis
44.26		✓ Topamax
▲ Tab 100 mg31.99	60	✓ Arrow-Topiramate
		✓ Topiramate Actavis
75.25		✓ Topamax
▲ Tab 200 mg55.19	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
129.85		✓ Topamax
▲ Sprinkle cap 15 mg20.84	60	✓ Topamax
▲ Sprinkle cap 25 mg26.04	60	✓ Topamax
VIGABATRIN - Special Authority see SA1997 below - Retail pharmacy		
▲ Tab 500 mg119.30	100	✓ Sabril

⇒SA1997 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

1 Either:

TODIRAMATE

- 1.1 Patient has infantile spasms; or
- 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or



Subsidy		ully Brand or	
(Manufacturer's \$	Price) Subsidis Per	sed Generic Manufacturer	

continued...

1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; and

2 Fither:

- 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
- 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 114

Acute Migraine Treatment

RIZATRIPTAN			
Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
SUMATRIPTAN			
Tab 50 mg	14.41	90	✓ Sumagran
•	24.44	100	✓ Apo-Sumatriptan
Sumagran to be Sole Supply on 1 February 2022			
Tab 100 mg	22.68	90	✓ Sumagran
-	46.23	100	✓ Apo-Sumatriptan
Sumagran to be Sole Supply on 1 February 2022			
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	34.00	2 OP	✓ Imigran
(Apo-Sumatriptan Tab 50 mg to be delisted 1 February 2022)			
(Apo-Sumatriptan Tab 100 mg to be delisted 1 February 2022)			

Prophylaxis of Migraine

*	Tab 500 mcg	23.21	100	✓ Sandomigran
PIZO	OTIFEN			
For	Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM	∕I, page 51		

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
` \$	Per	/	Manufacturer

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8			
APREPITANT - Special Authority see SA0987 below - Retail ph	armacy		
Cap 2 × 80 mg and 1 × 125 mg	30.00	3 OP	Emend Tri-Pack
Emend Tri-Pack to be Principal Supply on 1 December 2	021		

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE * Tab 16 mg	84	✓ Vergo 16
CYCLIZINE HYDROCHLORIDE		
Tab 50 mg	10	✓ Nausicalm
Nausicalm to be Principal Supply on 1 December 2021		
CYCLIZINE LACTATE		
Inj 50 mg per ml, 1 ml21.53	10	✓ <u>Hameln</u>
DOMPERIDONE		
* Tab 10 mg	100	Pharmacy Health
Pharmacy Health to be Principal Supply on 1 February 2022		
HYOSCINE HYDROBROMIDE		
* Inj 400 mcg per ml, 1 ml ampoule93.00	10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1998 below - Retail		
pharmacy14.11	2	Scopoderm TTS
SA1008 Special Authority for Subsidy		

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCI OPRAMIDE HYDROCHI ORIDE

	TOOLOT TO WINDE THE DITOON LOT INDE		
*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50	10	✓ <u>Pfizer</u>
Ν0	IDANSETRON		
*	Tab 4 mg2.68	50	✓ Onrex
*	Tab disp 4 mg – Up to 10 tab available on a PSO	10	✓ Ondansetron
			ODT-DRLA
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ Ondansetron
			ODT-DRLA

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
ROCHLORPERAZINE			
Tab 3 mg buccal	5.97	50	
•	(30.00)		Buccastem
Tab 5 mg - Up to 30 tab available on a PSO	8.00 [°]	250	✓ Nausafix
Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO		10	✓ Stemetil
Antipsychotics			
General			
MISULPRIDE - Safety medicine; prescriber may determine d	lispensing frequency		
Tab 100 mg	5.15	30	✓ Sulprix
•	17.16	100	✓ Amisulpride
			Mylan S29
Tab 200 mg	14 96	60	✓ Sulprix
Tab 400 mg		60	✓ Sulprix
•		00	- Ouiprix
IIPIPRAZOLE – Safety medicine; prescriber may determine	, , ,	00	/ Autologopala Carr
Tab 5 mg		30	✓ Aripiprazole Sand
Tab 10 mg		30	✓ Aripiprazole Sand
Tab 15 mg		30	✓ Aripiprazole Sand
Tab 20 mg		30	✓ Aripiprazole Sand
Tab 30 mg	17.50	30	Aripiprazole Sano
ILORPROMAZINE HYDROCHLORIDE - Safety medicine; p	rescriber may determi	ne dis	spensing frequency
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO	36.73	100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10	✓ Largactil
OZAPINE - Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing freq	Honov		
	•	50	✓ Clozaril
Tab 25 mg		50	
	6.69	400	✓ Clopine
	11.36	100	✓ Clozaril
T-1- 50	13.37		✓ Clopine
Tab 50 mg		50	✓ Clopine
T-1- 400	17.33	100	✓ Clopine
Tab 100 mg		50	✓ Clozaril
	17.33	465	✓ Clopine
	29.45	100	✓ Clozaril
T 000	34.65		✓ Clopine
Tab 200 mg		50	✓ Clopine
	69.30	100	Clopine
Suspension 50 mg per ml		100 m	
	67.62		✓ Versacloz
LOPERIDOL - Safety medicine; prescriber may determine	dispensing frequency		
ILOF ENIDOL – Salety Illeuichie, prescriber may determine i		100	✓ Serenace
	6.23		
Tab 500 mcg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 500 mcg - Up to 30 tab available on a PSO Tab 1.5 mg - Up to 30 tab available on a PSO	9.43	100 50	 ✓ <u>Serenace</u> ✓ Serenace
Tab 500 mcg - Up to 30 tab available on a PSO	9.43 14.86	50	✓ Serenace
Tab 500 mcg - Up to 30 tab available on a PSOTab 1.5 mg - Up to 30 tab available on a PSO	9.43 14.86 29.72		✓ Serenace ✓ Serenace

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Price)	Per	Subsidised	Manufacturer
EVOMEPROMAZINE – Safety medicine; prescriber may o	determine dispensing freg	uenc	<i>I</i>	
Tab 25 mg (33.8 mg as a maleate)		100		Nozinan (Swiss)
Tab 25 mg as a maleate		100	_	Nozinan
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100	_	Nozinan (Owiss)
•				
EVOMEPROMAZINE HYDROCHLORIDE – Safety medici				
Inj 25 mg per ml, 1 ml ampoule	33.50	10	/	<u>Nozinan</u>
THIUM CARBONATE - Safety medicine; prescriber may of	determine dispensing freq	uency	٧	
Tab long-acting 400 mg		100	'	Priadel
Cap 250 mg		100	_	Douglas
LANZAPINE – Safety medicine; prescriber may determine				g
	,	00	./	7.mina
Tab 5 mg		28		Zypine
Tab 5 mg		28		Zypine ODT
Tab orodispersible 5 mg		28	_	Zypine ODT
Tab 10 mg		28		Zypine
Tab orodispersible 10 mg	2.38	28	•	Zypine ODT
ERICYAZINE - Safety medicine; prescriber may determine	e dispensing frequency			
Tab 2.5 mg		84	1	Neulactil
· ••• =•• · · · · · · · · · · · · · · ·	12.49	100		Neulactil
Tab 10 mg	37.34	84		Neulactil
	44.45	100		Neulactil
UICTIADING Cofety was disingly averagible was a data was in a				
OUETIAPINE – Safety medicine; prescriber may determine			,	
Tab 25 mg		90		Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90	_	Quetapel
Tab 300 mg	12.86	90	/	<u>Quetapel</u>
ISPERIDONE - Safety medicine; prescriber may determin	e dispensina frequency			
Tab 0.5 mg		60	/	Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral liq 1 mg per ml		30 m		Risperon
		00 111	, ,	пізрегоп
IPRASIDONE – Safety medicine; prescriber may determin			_	
Cap 20 mg	14.50	60		Zusdone
Cap 40 mg	24.70	60	✓	Zusdone
Cap 60 mg	33.80	60	✓	Zusdone
Cap 80 mg	39.70	60	✓	Zusdone
UCLOPENTHIXOL HYDROCHLORIDE - Safety medicine	: prescriber may determin	e disr	nensina fra	equency
Tab 10 mg	•	100	• .	Clopixol
Tab To Hig	1.40	100	•	Сюріхої
Donat Injections				
Depot Injections				
LUPENTHIXOL DECANOATE - Safety medicine; prescrib	er may determine dispens	sina fr	requency	
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO.		5 III9 II		Fluanxol
ing to mig por mil, i mil op to o mil available on a root.		U	•	·····

✓ Fluanxol	5	ng per mi, 1 mi – Up to 5 inj available on a PSO13.14	Ir
✓ Fluanxol	5	ng per ml, 2 ml - Up to 5 inj available on a PSO20.90	lr
Fluanxol	5	mg per ml, 1 ml - Up to 5 inj available on a PSO40.87	lr

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
HALOPERIDOL DECANOATE - Safety medicine; prescriber ma	ay determine dispensi	ng frequ	ency	
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	✓	Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	✓	Haldol Concentrate
			✓ I	Haldol
				Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail pl	harmacy			
Safety medicine; prescriber may determine dispensing frequ	ency			
Inj 210 mg vial	252.00	1	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	1	Zyprexa Relprevv
Inj 405 mg vial	504.00	1	1	Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

ducty medicine, presented may determine dispersion	ig iroquorioy		
Inj 25 mg syringe	194.25	1	Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe		1	✓ Invega Sustenna
Inj 150 mg syringe		1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing fre	quency		
Inj 25 mg vial	135.98	1	Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	✓ Risperdal Consta

NERVOUS SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	/	Manufacturer

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 ✓ Clopixol

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	20.23	100	✓ Orion
* Tab 10 mg	13.16	100	✓ Orion
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency	/	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine dispen	ising frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Ativan to be Principal Supply on 1 December 2021			
Tab 2.5 mg	12.50	100	✓ Ativan
Ativan to be Principal Supply on 1 December 2021			
OXAZEPAM - Safety medicine; prescriber may determine dispe	nsing frequency		
Tab 10 mg	6.17	100	✓ Ox-Pam
Tab 15 mg	8.53	100	✓ Ox-Pam

Multiple Sclerosis Treatments

⇒SA2051 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic): and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2051 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg	520.00	14	✓ Tecfidera
Cap 240 mg	2,000.00	56	✓ Tecfidera

FINGOLIMOD - Special Authority see SA2051 on the previous page - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 0.5 mg......2,200.00 28 **✓ Gilenya**

GLATIRAMER ACETATE - Special Authority see SA2051 on the previous page - Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

lnj 40 mg prefilled syringe.................................2,275.00 12 **✓ Copaxone**

INTERFERON BETA-1-ALPHA - Special Authority see SA2051 on the previous page - Retail pharmacy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer INTERFERON BETA-1-BETA - Special Authority see SA2051 on page 141 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. NATALIZUMAB - Special Authority see SA2051 on page 141 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Tvsabri OCRELIZUMAB - Special Authority see SA2051 on page 141 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Ocrevus TERIFLUNOMIDE - Special Authority see SA2051 on page 141 - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. Aubagio Sedatives and Hypnotics MELATONIN - Special Authority see SA1666 below - Retail pharmacy Tab modified-release 2 mg - No more than 5 tab per day......11.50 30 ✓ Vigisom Circadin (Circadin Tab modified-release 2 mg to be delisted 1 April 2022) ⇒SA1666 Special Authority for Subsidy Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and 4 Patient is aged 18 years or under*. Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient is aged 18 years or under*; and

- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine	dispensing frequency		
Inj 1 mg per ml, 5 ml ampoule	3.95	10	Mylan Midazolam
, ,	5.50		✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj av	ailable		
on a PSO	14.90	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO mo	ust be endorsed for stati	us epilepticu	ıs use only.
Inj 5 mg per ml, 3 ml ampoule	4.50	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj ava	ilable on		
a PSO	11.90	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO mo	ust be endorsed for stati	us epilepticu	ıs use only.
(Mylan Midazolam Inj 1 mg per ml, 5 ml ampoule to be delis	sted 1 January 2022)		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PHENOBARBITONE SODIUM - Special Authority see SA1386 below - Retail pharmacy				
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓	Max Health S29
■ SA1386 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Both:	d without further renev	wal u	ınless notifi	ed for applications meeting

1 For the treatment of terminal agitation that is unresponsive to other agents; and

2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine di Tab 10 mg		25	✓ <u>Normison</u>
TRIAZOLAM - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 125 mcg	5.10	100	
•	(9.85)		Hypam
Tab 250 mcg	4.10	100	
	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine dis	pensing frequency		
Tab 7.5 mg		500	✓ Zopiclone Actavis
Zopiclone Actavis to be Principal Supply on 1 Februar	y 2022		•

Stimulants/ADHD Treatments

ATOMOXETINE

Cap 10 mg	18.41	28	✓ Generic Partners
, ,	107.03		✓ Strattera
Cap 18 mg	27.06	28	✓ Generic Partners
	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ Generic Partners
Cap 40 mg		28	✓ Generic Partners
•	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ Generic Partners
Cap 80 mg		28	✓ Generic Partners
Cap 100 mg		28	✓ Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see SA114 a) Only on a controlled drug form b) Safety medicine; prescriber may determine dispensing	•	armacy	
Tab 5 mg	' '	100	✓ PSM
PSM to be Principal Supply on 1 January 2022			-

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Fither:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the

Subsidy		Fully	Brand or	
(Manufacturer's Price)	;	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — **(Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg		30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg	7.75	30	✓ Methylphenidate ER
			- Teva
Tab immediate-release 20 mg	7.85	30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg		30	Methylphenidate ER
-			- Teva
Tab extended-release 36 mg	15.50	30	✓ Methylphenidate ER
v			- Teva
Tab extended-release 54 mg	22.25	30	✓ Methylphenidate ER
			- Teva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been



Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine: prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg		30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	✓ Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
- 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmac	у		
Tab 100 mg	29.13	60	Modavigil

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 on the nex	page – Retail pharma	су	
Patch 4.6 mg per 24 hour	38.00	30	Rivastigmine Patch BNM 5
	48.75		 Generic Partners
Generic Partners to be Principal Supply on 1 Februa	ry 2022		
Patch 9.5 mg per 24 hour	35.00	30	Generic Partners
	38.00		 Rivastigmine Patch BNM 10

Generic Partners to be Principal Supply on 1 February 2022

(Generic Partners Patch 4.6 mg per 24 hour to be delisted 1 February 2022)

(Generic Partners Patch 9.5 mg per 24 hour to be delisted 1 February 2022)



Subsidy		ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	/	Manufacturer	

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia: and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate: and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine: prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	.18.37	28	✓ <u>Buprenorphine</u> Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	.53.12	28	✓ <u>Buprenorphine</u> Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health:
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and

NERVOUS SYSTEM

✓ Naltraccord

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	

continued...

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
 - 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see		il pharmacy	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.



Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
 \$	Per		Manufacturer	

NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO18.14 ✓ Habitrol 28 Patch 7 mg for direct distribution only - [Xpharm]......3.94 7 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO19.95 ✓ Habitrol 28 Patch 14 mg for direct distribution only - [Xpharm]......4.52 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO22.86 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......5.18 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......19.18 216 ✓ Habitrol 36 ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......21.02 216 ✓ Habitrol ✓ Habitrol 36 Gum 2 mg (Fruit) - Up to 384 piece available on a PSO38.21 384 ✓ Habitrol Gum 2 mg (Fruit) for direct distribution only - [Xpharm].....8.64 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 384 piece available on a PSO......38.21 384 ✓ Habitrol Gum 2 mg (Mint) for direct distribution only - [Xpharm]......8.64 96 ✓ Habitrol Gum 4 mg (Fruit) - Up to 384 piece available on a PSO44.17 384 ✓ Habitrol

Gum 4 mg (Mint) for direct distribution only – [Xpharm]................10.01

VARENICLINE TARTRATE – Special Authority see SA1845 below – Retail pharmacy

Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17

a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack

96

384

96

✓ Habitrol

✓ Habitrol

✓ Habitrol

- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
	25.64		Champix
Varenicline Pfizer to be Principal Supply on 1 January	2022		
Tab 1 mg	17.62	56	✓ Varenicline Pfizer
·	27.10		✓ Champix

Varenicline Pfizer to be Principal Supply on 1 January 2022

(Champix Tab 0.5 mg x 11 and 1 mg x 42 to be delisted 1 January 2022)

(Champix Tab 1 mg to be delisted 1 January 2022)

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has

NERVOUS SYSTEM

Subsidy (Manufacturer's Price)	Subs	Fully idised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

agreed to this: and

- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see \$A2046 below

Inj 25 mg vial77.00	1	Ribomustin
Inj 100 mg vial308.00	1	✓ Ribomustin
Inj 1 mg for ECP	1 mg	✓ Baxter

⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

Subsidy	Fully		Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

busulfan - Put - Retail pharmacy-specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	✓ DBL Carboplatin
, •	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	1,387.00	1	✓ BiCNU
, ,			✓ Bicnu Heritage S29
Inj 100 mg for ECP	1,387.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial	21.00	1	✓ Cisplatin Ebewe
, 34- ,	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE		-	
Tab 50 mg - PCT - Retail pharmacy-Specialist	79.00	50	✓ Endoxan S29
rab oo mg	145.00	00	✓ Cyclonex
	158.00	100	✓ Procytox S29
Cyclonex to be Principal Supply on 1 January 2022	100.00	100	1 Tooytox
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
, ,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
(Endoxan S29 Tab 50 mg to be delisted 1 January 2022)		-	
(Procytox \$29 Tab 50 mg to be delisted 1 January 2022)			
, 11,11			

	Subsidy (Manufacturer's Price)		Fully Subsidised	I Generic
	\$	Per		Manufacturer
FOSFAMIDE - PCT only - Specialist				
Inj 1 g	96.00	1	1	Holoxan
Inj 2 g	180.00	1	✓	Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓	Baxter
OMUSTINE - PCT - Retail pharmacy-Specialist				
Cap 10 mg	132.59	20	✓	CeeNU
Cap 40 mg	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist		1	1	Alkeran
, , ,			✓	Alkeran S29 S29
	420.00		/	Tillomed S29
Tillomed \$29 Inj 50 mg to be delisted 1 December 2021)				
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	/	Oxaliplatin Actavis
ing 100 mg via	20.01	•	•	100
	110.00		/	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial		1		Oxaliplatin Accord
Inj 1 mg for ECP		1 mg		Baxter
HIOTEPA - PCT only - Specialist		3		
Inj 15 mg vial	CRS	1	ſ	Bedford S29
ing 10 mg viai		'		Max Health \$29
				THIO-TEPA \$29
let 400 mental	000			Tepadina S29
Inj 100 mg vial	CBS	1	_	Max Health S29
			•	Tepadina S29

Antimetabolites

AZACITIDINE – PCT only – Specialist – Special Authority see SA1467 below)W	
Inj 100 mg vial75.0	06 1	 Azacitidine Dr Reddy's
605.0	00	✓ Vidaza
Azacitidine Dr Reddy's to be Principal Supply on 1 December 2021		
Inj 1 mg for ECP0.8	33 1 mg	✓ Baxter

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

CALCIUM FOLINATE		
Tab 15 mg - PCT - Retail pharmacy-Specialist114.69	10	✓ DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialist7.28	1	✓ Calcium Folinate
		<u>Sandoz</u>
		 Calcium Folinate
		Sandoz S29 S29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist9.49	1	 Calcium Folinate
		Sandoz
Inj 100 mg - PCT only - Specialist7.33	1	 Calcium Folinate
		Ebewe
Inj 300 mg – PCT only – Specialist22.51	1	Calcium Folinate Ebewe
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist25.14	1	✓ Calcium Folinate
ing forming per mit, 65 mil viai 1 of only opecialist25.14		Sandoz
		✓ Calcium Folinate
		Sandoz S29 S29
Inj 1 g - PCT only - Specialist67.51	1	✓ Calcium Folinate
ing r g r or only opposition		Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist72.00	1	✓ Calcium Folinate
,g por,	·	Sandoz
Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
CAPECITABINE – Retail pharmacy-Specialist	· ·	
Tab 150 mg	60	✓ Capercit
Tab 500 mg	120	✓ Capercit
CLADRIBINE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml749.96	1	✓ Leustatin
Inj 10 mg for ECP	10 mg OP	✓ Baxter
CYTARABINE	ŭ	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist400.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail	-	
pharmacy-Specialist	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist	10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Specialist80.00	100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE		
Tab 10 mg - PCT - Retail pharmacy-Specialist412.00	20	✓ Fludara Oral
Inj 50 mg vial - PCT only - Specialist576.45	5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist115.29	50 mg OP	✓ Baxter

	Subsidy		Fully	Brand or
	(Manufacturer's Price \$	e) ; Per	Subsidised	Generic Manufacturer
FILLIO DOLUDA OU	Ψ	1 01		Widifuldcturer
FLUOROURACIL	10.51			
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist		1	-	Fluorouracil Accord
	12.00			Fluorouracil Ebewe
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist		1		Fluorouracil Accord
	30.00			Fluorouracil Ebewe
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	y 🗸 I	Baxter
(Fluorouracil Ebewe Inj 50 mg per ml, 20 ml vial to be delisted 1	February 2022)			
(Fluorouracil Ebewe Inj 50 mg per ml, 100 ml vial to be delisted 1	February 2022)			
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62 50	1	✓	DBL Gemcitabine
Inj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	-	Baxter
	0.02	i ilig	• ,	Daxio
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist	F0 F7		,	
Inj 20 mg per ml, 5 ml vial		1		Accord
	71.44		✓	Irinotecan
				Accord S29
			✓	Irinotecan Actavis 100
	100.00		✓	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	✓	Baxter
(Irinotecan Accord S29 Inj 20 mg per ml, 5 ml vial to be delisted		9		
, , ,	i waion zozz)			
MERCAPTOPURINE	07.00	0.5		
Tab 50 mg - PCT - Retail pharmacy-Specialist		25	✓	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist				
Special Authority see SA1725 below	428.00	100 ml C)P 🗸	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

		Subsidy		Fully	Brand or
		(Manufacturer's Price		Subsidised	Generic
_		\$	Per		Manufacturer
ME	THOTREXATE				
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist	9.98	90	1	Trexate
-	Trexate to be Principal Supply on 1 January 2022				
*	Tab 10 mg - PCT - Retail pharmacy-Specialist	33.71	90	1	Trexate
•	Trexate to be Principal Supply on 1 January 2022				
*	Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist	47 50	5	1	Methotrexate DBL
*	Inj 7.5 mg prefilled syringe		1		Methotrexate
*	ing 7.5 mg promied symige	17.01	'	•	Sandoz
*	Ini 10 ma profilled arrings	14.00	1	./	Methotrexate
*	Inj 10 mg prefilled syringe	14.00	ı	V	Sandoz
*	Inj 15 mg prefilled syringe	14.77	1	•	Methotrexate
					Sandoz
*	Inj 20 mg prefilled syringe	14.88	1	✓	Methotrexate
					Sandoz
*	Inj 25 mg prefilled syringe	14.99	1	1	Methotrexate
	, , , ,				Sandoz
*	Inj 30 mg prefilled syringe	15.09	1	/	Methotrexate
-	, g p , g		-		Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Speciali	et 30.00	5	1	Methotrexate DBL
*	mij 25 mg per mi, 2 mi viai - 1 O1 - Hetali phamacy opecial	31	3	•	Onco-Vial
	Ini OF man and OO militial DOT Datail abannasi Casain	.II1 4E 00	4		DBL Methotrexate
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specia	IIIST45.00	1	•	
				_	Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist	t25.00	1	•	Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial - PCT - Retail				
	pharmacy-Specialist		1	✓	Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓	Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist.	4.73	5 mg Ol	P 🗸	Baxter
PF	METREXED - PCT only - Specialist - Special Authority see	SA1679 below			
-	Inj 100 mg vial		1	1	Juno Pemetrexed
	Inj 500 mg vial		1		Juno Pemetrexed
	Inj 1 mg for ECP		1 mg		Baxter
_			1 1119	•	-unioi

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

Subsidy (Manufacturer's Price)	Full Subsidise	' · · · · · ·	
 \$	Per •	Manufacturer	

continued...

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist			
Tab 40 mg	126.31	25	Lanvis

			_
Other	Cvto	tovio	Agents
OHIE	CVIU	LUXIL	AUCIIIO

AMSACRINE – PCT only – Specialist		
Inj 50 mg per ml, 1.5 ml ampoule	6	✓ Amsidine S29
4,736.00		✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mg1,175.87	100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter
BLEOMYCIN SULPHATE - PCT only - Specialist		
Inj 15,000 iu, vial161.01	1	✓ DBL Bleomycin Sulfate
Inj 1,000 iu for ECP12.45	1,000 iu	✓ Baxter
BORTEZOMIB - PCT only - Specialist - Special Authority see SA1889 on the r	next page	
Inj 2.5 mg vial	1	✓ Bortezomib Juno S29 S29
Inj 3.5 mg vial105.00	1	✓ Bortezomib Dr Reddy's S29 S29
		✓ Bortezomib Dr-Reddy's
		✓ Bortezomib Juno \$29
Inj 1 mg for ECP31.20	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	 DBL Dacarbazine
	580.60	10	Dacarbazine
			APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist			
Inj 2 mg per ml, 10 ml	149.50	1	✓ Pfizer
Inj 20 mg for ECP	149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist			
Inj 20 mg	48.75	1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial	26.95	1	✓ Docetaxel
•			Accord S29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	 Doxorubicin Ebewe
	17.00		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial		1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	Arrow-Doxorubicin
	69.99		Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial		1	 Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP	0.50	1 mg	✓ Baxter
ETOPOSIDE			
Cap 50 mg - PCT - Retail pharmacy-Specialist		20	✓ <u>Vepesid</u>
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓ <u>Vepesid</u>
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia	ılist7.90	1	✓ Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	✓ Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist			
Inj 100 mg (of etoposide base)		1	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pharm			_	
Cap 500 mg	23.82	100	/	<u>Devatis</u>
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial - PCT only - Specialist	93.00	1	✓	Zavedos
Inj 10 mg vial - PCT only - Specialist		1	✓	Zavedos
Inj 1 mg for ECP - PCT only - Specialist		1 mg	1	Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority Wastage claimable	see SA2047 below			
Cap 5 mg	5,122.76	28	✓	Revlimid
Cap 10 mg		21	✓	Revlimid
, ,	6,207.00	28	✓	Revlimid
Cap 15 mg	5,429.39	21	✓	Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg	7,627.00	21	✓	Revlimid
0.000				

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 21 | 00
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

	Subsidy		Fully Brand or
	(Manufacturer's Price	A Cuk	osidised Generic
	\$	Per	✓ Manufacturer
MESNA			
Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist		15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 20 mg vial	3,275.00	1	✓ Omegapharm S29
, 3	•		✓ Teva
Inj 1 mg for ECP	470.75	1 mg	✓ Baxter
IITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
DLAPARIB – Retail pharmacy-Specialist – Special Authority se		Ü	
Tab 100 mg		56	✓ Lynparza
Tab 150 mg		56	✓ Lynparza
	-,		, i

⇒SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment: and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist

TOLITABLE TOTOTILY OPCOIDING			
Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
, ,	91.67		Paclitaxel Actavis
Inj 150 mg	26.69	1	Paclitaxel Ebewe
, 0	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
, 0	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter

	(Manufacturer's Price)	Subsidised		,	
	\$	Per	1	Manufacturer	
PEGASPARGASE - PCT only - Special Authority see SA1979 b	elow				
Ini 750 ju per ml. 5 ml vial	3.455.00	1	√ 0	ncaspar LYO S29	

Subeidy

Fully

Brand or

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail			
Cap 50 mg		50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 be			
Cap 5 mg	'	5	✓ Temaccord
Cap 20 mg		5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ Temaccord
•	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ <u>Temaccord</u>
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord \$29
Cap 250 mg		5	✓ Temaccord
•	688.00		✓ Amneal S29

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE	 Retail pharmacy-Specialist – Special Authority see SA1124 belo 	W	
Cap 50 mg.	378.00	28	Thalomid
Cap 100 mg		28	✓ Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRFTINOIN

Vesanoid 100

	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 below			
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OF	•	Venclexta
Tab 10 mg	95.78	14 OF	•	Venclexta
Tab 50 mg	239.44	7 OP	✓	Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓	Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SUI PHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	✓ DBL Vincristine
		Sulfate
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	DBL Vincristine
		Sulfate
Ini 1 mg for ECP - PCT only - Specialist	1 ma	✓ Baxter

	Subsidy (Manufacturer's Price \$) Sı Per	Fully ubsidised	
VINORELBINE - PCT only - Specialist				
Inj 10 mg per ml, 1 ml vial	12.00	1	1	Navelbine
	42.00		/	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	/	Navelbine
	210.00		/	Vinorelbine Ebewe
	328.65		/	Sagent S29
Inj 1 mg for ECP	1.25	1 mg		Baxter
Inj 50 mg for ECP	328.65	0 mg OP	•	Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test: and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

wasiage ciaimable			
Tab 20 mg	3,774.06	60	Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel
3	**		

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

Mostogo alaimable

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
- 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
- 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

⇒SA2000 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive: or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see SA2001 below

Tab 250 mg1,700.00 30 ✓ Iressa

⇒SA2001 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460			
	below	. 2,400.00	60	✓ Glivec
*	Cap 100 mg	58.23	60	✓ Imatinib-Rex
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

 Wastage claimable
 4,680.00
 120
 ✓ Tasigna

 Cap 150 mg
 6,532.00
 120
 ✓ Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either
 - 2.1 Patient has documented CML treatment failure* with imatinib; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

- 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Mostogo eleimeble

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

wasiage cialinable			
Tab 75 mg	4,000.00	21	✓ Ibrance
Tab 100 mg		21	✓ Ibrance
Tab 125 mg	4,000.00	21	✓ Ibrance
Cap 75 mg	4,000.00	21	✓ Ibrance
Cap 100 mg	4,000.00	21	✓ Ibrance
Cap 125 mg	· ·	21	✓ Ibrance

(Ibrance Cap 75 mg to be delisted 1 March 2022)

(Ibrance Cap 100 mg to be delisted 1 March 2022)

(Ibrance Cap 125 mg to be delisted 1 March 2022)

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	✓ ✓	
PAZOPANIB - Special Authority see SA1190 below - Retail pha	rmacy			
Tab 200 mg	1,334.70	30	✓	Votrient
Tab 400 mg	2,669.40	30	✓	Votrient

SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 15 mg	5,000.00	56	Jakavi
Tab 20 mg	5,000.00	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis: and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:

Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
\$	Per	1	Manufacturer

continued...

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2002 below - Retail pharmacy

Cap 12.5 mg2,315.38	28	Sutent
Cap 25 mg4,630.77	28	✓ Sutent
Cap 50 mg	28	✓ Sutent

⇒SA2002 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 86

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2003 below

Wastage claimable

⇒SA2003 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and

(A)	Subsidy fanufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
	\$	Per	✓	Manufacturer

continued...

- 4.2.2 Patient has ECOG performance score of 0-2; and
- 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

BICALUTAMIDE

Tab 50 mg	1.36	10	✓ Calutide-50 S29
· ·	4.21	28	✓ Binarex
(Calutide-50 S29	Tab 50 mg to be delisted 1 January 2022)		
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
	119.50	100	✓ Flutamin
FULVESTRANT -	- Retail pharmacy-Specialist - Special Authority see SA1895 below	1	
Inj 50 mg per	ml, 5 ml prefilled syringe1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	
CTREOTIDE				
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓	Octreotide GH S29
Inj 50 mcg per ml, 1 ml ampoule	30.64	5	✓	Octreotide GH S29
Inj 50 mcg per ml, 1 ml vial	30.64	5	✓	Octreotide
				MaxRx S29
	56.87		✓	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓	DBL Octreotide
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓	Octreotide GH S29
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓	DBL Octreotide
	222.00		•	Octreotide
				(Sun) S29
CTREOTIDE LONG-ACTING - Special Authority see SA2072	below - Retail pharm	асу		
Inj depot 10 mg prefilled syringe	439.97	1	•	Octreotide Depot Teva
	1,772.50		1	Sandostatin LAR
Inj depot 20 mg prefilled syringe	647.03	1	•	Octreotide Depot Teva
	2,358.75		✓	Sandostatin LAR
Inj depot 30 mg prefilled syringe	718.55	1	•	Octreotide Depot Teva
	2,951.25		✓	Sandostatin LAR
Sandostatin LAR Inj depot 10 mg prefilled syringe to be delisted Sandostatin LAR Inj depot 20 mg prefilled syringe to be delisted				

⇒SA2072 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and

(Sandostatin LAR Inj depot 30 mg prefilled syringe to be delisted 1 March 2022)

- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed: and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

ubsidy cturer's Price) Subs	Fully	Brand or Generic
 \$ Per	•	Manufacturer

continued...

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

ΙAΙ	MOXIFEN CITRATE	
*	Tab 10 mg	5.00

Aromatase Inhibitors		
ANASTROZOLE	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE * Tab 2.5 mg5.84	30	✓ Letrole

Letrole to be Principal Supply on 1 January 2022

60

60

✓ <u>Tamoxifen Sandoz</u>✓ <u>Tamoxifen Sandoz</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic S Per ✓ Manufacturer

Immunosuppressants

Cytotoxic Immunosuppressants

ΑZ	ATHI)PI	RINE
110	T - 1-	^-	

*	1au 25 mg7.35	60	▼ <u>Azamun</u>
*	Tab 50 mg	100	✓ Azamun
*	Inj 50 mg vial199.00	1	Imuran

MYCOPHENOLATE MOFETIL

TOOPHENOLATE MOPETIL			
Tab 500 mg	35.90	50	Cellcept
Cap 250 mg	35.90	100	 Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement		165 ml OP	 Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

7 05

Fusion Proteins

ETANERCEPT - Special Authority see SA2048 below -	- Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector		4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2048 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis: or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs. CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right): or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm: Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm: Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less: and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

continued...

- 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

Subsi	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

2 Fither:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for rheumatoid arthritis: or

2 All of the following:

- 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and

-	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
=	<u> </u>				

continued...

- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints;
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plague psoriasis; and
 - - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

ANTITHYMOCYTE GLOBILLIN (FOLLINE) - PCT only - Specialist

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITITIMOOTTE GEODOLIN (EQUINE) - 1 OT OHIJ - Specialist		
Inj 50 mg per ml, 5 ml2,351.25	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only - Specialist		
Subsidised only for bladder cancer.		
Inj 2-8 × 100 million CFU149.37	1	✓ OncoTICE
Inj 40 mg per ml, vial176.90	3	✓ SII-Onco-BCG S29
(SII-Onco-BCG \$29 Inj 40 mg per ml, vial to be delisted 1 April 2022)		

Monoclonal Antibodies

		below – Retail pharmacy	ADALIMUMAB – Special Authority see SA2049 be
Humira	2	1,599.96	Inj 20 mg per 0.4 ml prefilled syringe
✓ HumiraPen	2	1,599.96	Inj 40 mg per 0.8 ml prefilled pen
Humira	2	1,599.96	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2049 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Roth
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Fither:

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 1.2.1 The patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Fither:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the following average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- of the followin 1 Either:
 - 1.1 Applicant is a gastroenterologist; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application; and
- 4 The patient will be assessed for response to treatment after 4 months' adalimumab treatment (see Note).

Note: A maximum of 4 months' adalimumab will be subsidised on an initial Special Authority approval for fistulising Crohn's disease.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or patient has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 The patient has 3 or more active lesions (e.g. inflammatory nodules, abscesses, draining fistulae); and
- 4 The patient has a Dermatology Quality of Life Index of 10 or more and the assessment is no more than 1 month old at time of application; and
- 5 Following the initial loading doses, adalimumab is to be administered at doses no greater than 40mg every 7 days.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for oligoarticular course JIA; or
- 2 All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
` \$	Por 🗸	Manufacturer

continued...

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from etanercept or secukinumab to meet the renewal criteria for etanercept or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (rheumatoid arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept: or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
 - 2.5 Any of the following:
 - 2.5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

- 2.5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
- 2.5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 2.6 Fither:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.7 Either:
 - 2.7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease that is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has not responded adequately to treatment with infliximab (see Notes); or
 - 2.2 The patient has severe ocular, neurological, gastrointestinal, rheumatological, mucocutaneous and/or vasculitic symptoms and has experienced intolerable side effects from treatment with infliximab; and
- 3 The patient is experiencing significant loss of quality of life; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: Behcet's disease diagnosed according to the International Study Group for Behcet's disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al, J Rheumatol. 2004;31:931-7.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from etanercept; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment: and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

- 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from infliximab; or
 - 1.2.2 The patient has received insufficient benefit from infliximab to meet the renewal criteria for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
 - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	Erbitux
Inj 1 mg for ECP	3.82	1 mg	Baxter

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB	- PCT only	- Special	Authority	see	SA2082	helow

Inj 100 mg	806.00	1	Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

continued...

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances: or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab. etanercept or secukinumab for severe chronic plaque psoriasis: or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

	Subsidy		Fully	Brand or
(1	Manufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued...

- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment: and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis: or
 - 2.2 Ankylosing spondylitis: or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis: or
 - 2.10 Severe ulcerative colitis: or
 - 2.11 Plague psoriasis; or
 - 2.12 Neurosarcoidosis: or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
 - 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
(Manuacturer's Frice)		oubsidised		
\$	Per	/	Manufacturer	

continued...

significant response to prior infliximab treatment in the opinion of the treating physician; and

2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - I	Retail pharmacy		
Inj 100 mg prefilled pen	1,638.00	1	Nucala
Ini 100 mg vial	1.638.00	1	✓ Nucala

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5 x 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

Inj 25 mg p	er ml, 40 ml vial.	 	5,910.	.00 1	✓	Gazyva
Inj 1 mg for	· ECP	 	6.	.21 1 r	ng 🗸	Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and</p>
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

		openia riamoni, coo orin ribani priamao,	
Xolair	1	prefilled syringe450.00	Inj 150 mg
✓ Xolair	1	vial450.00	Ini 150 ma

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and

- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses: or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Roth:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 on the next page

Inj 30 mg per ml, 14 ml vial	3,927.00	1	✓	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓	Baxter

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

Inj 100 mg per 10 ml vial	1,075.50	2	Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1.000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2083 below

Inj 100 mg per 10 ml vial	275.33 2	•	' Riximyo
Inj 500 mg per 50 ml vial	688.20 1	•	Riximyo
Inj 1 mg for ECP	1.38 1 mg	•	Baxter (Riximyo)

⇒SA2083 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — **(ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive: or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax: or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following

criteria: Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

375 mg/m2 administered weekly for four weeks; and

- - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks: and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective: or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither

Subsidy	Fı	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
- 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*: and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*: and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

(Ma	Subsidy nufacturer's Price)	Sub	Fully	Brand or Generic
<u> </u>	\$	Per	✓	Manufacturer

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	1	Manufacturer	

continued...

- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of

(Mar	Subsidy nufacturer's Price)	Suk	Fully	Brand or Generic
	\$	Per	•	Manufacturer

continued...

4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

Subsid (Manufacturer	•	Fully Brand of Subsidised Generic		
(Manuacture)	Per	J3iui3eu ✓	Manufacturer	

continued...

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*: and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
 - 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
 - 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

Both:

1 Either:

- 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		osidised	Generic
<u> </u>	Per		Manufacturer

continued...

- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA2078 below

Inj 20 mg per ml, 4 ml vial	220.00	1	Actemra
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
Inj 20 mg per ml, 20 ml vial		1	✓ Actemra
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA2078 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease: or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules: and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy: and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

6 Either:

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Roth:

1.1 Fither:

- 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
- 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule: and

1.2 Fither:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or

2 All of the following:

- 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
- 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or

2 All of the following:

- 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
- 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or

Subsi	idy Fully	Brand or
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient has significantly increased laboratory markers of systemic inflammation (eg CRP, PCT or ferritin); and
- 4 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 5 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose.

Note: Indications marked with * are unapproved indications.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRASTUZUMAB - PCT only - Specialist - Special Authority see	SA1632 below			
Inj 150 mg vial	1,350.00	1	✓	Herceptin
Inj 440 mg vial	3,875.00	1	✓	Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓	Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

continued...

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and

TRACTUZUMAR EMTANCINE ROT only Charielist Chariel Authority and CA1971 below

- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

INASTUZUWADE	ivi fansine – PCT only – specialist – special Authority see	SATO/ I DEIUW	
Inj 100 mg via	2,320.00	1	Kadcyla
Inj 160 mg via	3,712.00	1	✓ Kadcyla
Ini 1 ma for FC	P 23.20	1 ma	✓ Baxter

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

Programmed Cell Death-1 (PD-1) Inhibitors

		NIVOLUMAB – PCT only – Specialist – Special Authority see SA2006 below	N
Opdivo	1	Inj 10 mg per ml, 4 ml vial	
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96	
✓ Baxter	1 mg	Inj 1 mg for ECP27.62	

⇒SA2006 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Either:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
<u> </u>	\$ Pe	er 🗸	Manufacturer

continued...

or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special Au	uthority see SA2007 below	
Inj 25 mg per ml, 4 ml vial	4,680.00 1	Keytruda
Inj 1 mg for ECP	49.14 1 mg	✓ Baxter

⇒SA2007 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Fither:
 - 1.2.1 Response to treatment in target lesions has been determined by radiologic assessment (CT or MRI scan) following the most recent treatment period; or
 - 1.2.2 Both:
 - 1.2.2.1 Patient has measurable disease as defined by RECIST version 1.1; and
 - 1.2.2.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	Neoral
EVEROLIMUS – Special Authority see SA2008 below – Retail ph Wastage claimable	armacy		
Tab 10 mg	6.512.29	30	✓ Afinitor
Tab 5 mg		30	✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

	Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic
	\$	Per	1	Manufacturer
SIROLIMUS – Special Authority see SA2005 below – Retail ph	armacy			
Tab 1 mg	749.99	100	√ R	apamune
Tab 2 mg	1,499.99	100	√ R	apamune
Oral liq 1 mg per ml	449.99	0 ml OP	•	apamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

Cap 0.5 mg	•	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB - Special Authority see SA2079 on the next page - Retail pharmacy ✓ RINVOQ 28

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Por 🗸	Manufacturer	

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

ove - Retail pharmac	у
00 1 OP	✓ VENOX S29
00 1 OP	✓ VENOX S29
00 1 OP	✓ Venomil S29
00 1 OP	✓ Albey
00 1 OP	✓ Hymenoptera S29
<mark>above – Retail pharm</mark>	acy
00 1 OP	✓ Albey
00 1 OP	✓ Hymenoptera S29
	_
00 1 OP	✓ Venomil S29
00 1 OP	✓ Hymenoptera S29
	4 All
00 1 OP	✓ Albey
00 1 00	/ Vanamil oo
UU 1 UP	✓ Venomil S29
	1 OP

	Subsidy		Fully	Brand or
	(Manufacturer's Pri		idised	Generic
	\$	Per	✓	Manufacturer
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1.12	100	✓ Z	ista
* Oral liq 1 mg per ml	2.84	200 ml	√ H	listaclear
Histaclear to be Principal Supply on 1 January 2022				
CHLORPHENIRAMINE MALEATE				
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ H	listafen
		300 1111	• .	iistaicii
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg		40	_	
	(8.40)		P	Polaramine
	1.01	20	_	
	(5.99)		P	Polaramine
* Oral liq 2 mg per 5 ml		100 ml		
	(10.29)		P	Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
ů	(8.23)		Т	elfast
* Tab 120 mg	4.74 [′]	10		
, and a g	(8.23)		Т	elfast
	14.22	30		
	(26.44)		Т	elfast
LORATADINE	(- /			
* Tab 10 mg	1.60	100	./ I	.orafix
· · · · · · · · · · · · · · · · · · ·		100 ml		laylor syrup
	1.43	100 1111	• [iayior syrup
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg		50		Allersoothe
* Tab 25 mg		50	-	Allersoothe
* Oral liq 1 mg per 1 ml		100 ml		Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a	ı PSO 17.87	5	✓ H	lospira
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose		200 dose OP	√ 0	
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OP		Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	√ (
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ E	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ P	ulmicort
• .				Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	√ P	ulmicort
			•	Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	√ □	Pulmicort
1 Officer for inflatation, 400 flicy per dose		_00 003E OF	· F	Tambook alam

Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's		
	\$	Per	✓ Manufacturer
UTICASONE			
Aerosol inhaler, 50 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	 Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	 Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide Accuhaler
nhaled Long-acting Beta-adrenoceptor Agoni	sts		
ORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose de	evice20.64	60 dose	
, 91	(35.80)		Foradil
ORMOTEROL FUMARATE DIHYDRATE	, ,		
Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered do	sa) 10.32	60 dose OP	
(oquivalent to elonnoterol lumarate o moy metered do	(16.90)	ou dose or	Oxis Turbuhaler
DAGATERO	(10.50)		Onio i ulbullalei
DACATEROL Benden (acids delation 450 man	21.25	00 -1- 05	(Out
Powder for inhalation 150 mcg		30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	 Onbrez Breezhaler
ALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	25.00	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated	25.00	60 dose OP	 Serevent Accuhaler
nhaled Corticosteroids with Long-Acting Beta	a-Adrenocept	tor Agonists	
IDEOCNIDE MITH EEODMOTEDOL			
IDESONIDE WITH FEORMOTEROL			
JDESONIDE WITH EFORMOTEROL Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol	with		
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide		120 dose OP	✓ DuoResn Sniromay
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 2		
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 2 82.50	120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 2 82.50 18.23	120 dose OP 120 dose OP	✓ DuoResp Spiromax✓ Vannair
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 2 82.50 18.23	120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)		120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 282.5018.23 6 mcg33.7421.40	120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 282.5018.23 6 mcg33.7421.40	120 dose OP 120 dose OP 120 dose OP	 ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 282.5018.23 6 mcg33.7421.40	120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)		120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	 ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)		120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)		120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	 ✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6
Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide 6 mcg eformoterol fumarate metered dose)	41.50 arate ncg 282.5018.23 6 mcg33.7421.40 6 mcg44.08	120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP 120 dose OP	✓ DuoResp Spiromax ✓ Vannair ✓ Symbicort Turbuhaler 100/6 ✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort

Subsidy (Manufacturer's \$		Fully Brand or dised Generic Manufacturer
25.7932.6033.7444.08	120 dose OP 120 dose OP 60 dose OP	✓ <u>Seretide</u> ✓ <u>Seretide</u> ✓ Seretide Accuhaler ✓ Seretide Accuhaler
40.00	150 ml	✓ Ventolin
118.38	10 5	✓ Ventolin✓ Ventolin
3.80	200 dose OP	✓ Respigen✓ SalAir
(6.00)	20	Ventolin ✓ Asthalin
9.43	20	✓ Asthalin
22.20	120 dose OP	✓ Bricanyl Turbuhaler
16.20	200 dose OP	✓ Atrovent
eb 11.73	20	✓ <u>Univent</u>
holinergic A	Agents	
per 12.19	200 dose OP 20	✓ Duolin HFA ✓ Duolin
	(Manufacturer's \$	(Manufacturer's Price) Subsist

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg......81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg.......81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic
` \$	Per	•	Manufacturer

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note. I interlidorie is not subsidised in combinati	on with subsidised militedamb.		
Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet
Cap 267 mg - Wastage claimable		270	✓ Esbriet
3	-,		

(Esbriet Cap 267 mg to be delisted 1 January 2022)

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
Leukotriene Receptor Antagonists				
MONTELUKAST * Tab 4 mg * Tab 5 mg * Tab 10 mg	4.25	28 28 28	√ √	Montelukast Mylan Montelukast Mylan Montelukast Mylan
Methylxanthines AMINOPHYLLINE * Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a				
PSO		5	✓	DBL Aminophylline
* Tab long-acting 250 mg * Oral liq 80 mg per 15 ml		100 500 m	_	Nuelin-SR Nuelin
Mucolytics				
DORNASE ALFA – Special Authority see SA1978 below – Reta Nebuliser soln, 2.5 mg per 2.5 ml ampoule	, ,	6	✓	Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Autho	rity see SA2017 below		
Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N

	RESPIRAT	TORY SYSTE	IA ME	ND ALLERGIES
	Subsidy (Manufacturer's \$	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer
continued				
and S549R) in the CFTR gene on at least 1 allele 3 Patients must have a sweat chloride value of at least 60 sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly wi 5 Patient must not have an acute upper or lower respirator (including antibiotics) for pulmonary disease in the last 4 6 The dose of ivacaftor will not exceed one tablet or one s 7 Applicant has experience and expertise in the managem	mmol/L by quant th standard thera ry infection, pulma weeks prior to co achet twice daily;	py for this conditionary exacerbations treat and	tion; an	d changes in therapy
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	√ B	iomed
Need Busessians				
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE				
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	_	teroClear
Metered aqueous nasal spray, 100 mcg per dose FLUTICASONE PROPIONATE	2.84	200 dose OP	V <u>5</u>	<u>teroClear</u>
Metered agueous nasal spray, 50 mcg per dose	1.98	120 dose OP	√ F	lixonase Hayfever
3,7				& Allergy
Flixonase Hayfever & Allergy to be Principal Supply on	1 December 202	21		
IPRATROPIUM BROMIDE	T 00	45I OD		
Aqueous nasal spray, 0.03%	5.23	15 ml OP	∨ <u>U</u>	<u>nivent</u>
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
c) Only for children aged six years and under Small	2 20	1	√ e	-chamber Mask
PEAK FLOW METER		•	- 0	onambor maon
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	Low Hange lini-Wright
		•	-	Standard

✓ e-chamber Turbo

✓ e-chamber La Grande

✓ Volumatic

SPACER DEVICE

b) Only on a PSO

a) Up to 50 dev available on a PSO

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml).......15.10 25 ml OP **✓ Biomed**

|--|

				-	н		
Ear	121	er	ЖII	"	11	01	11:
-0.1		Ψï				9	

ACETIC ACID WITH 1, 2- PROPANEDIOL DIACETATE AND BENZETHONIUM For Vosol ear drops with hydrocortisone powder refer Standard Formulae, page 247						
Ear drops 2% with 1, 2-Propanediol diacetate 3% and	a Formulae, pa	19 6 247				
benzethonium chloride 0.02%	6.97	35 ml OP	✓ Vosol			
FLUMETASONE PIVALATE						
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's			
			✓ Locorten-Vioform			
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate	AND NYSTAT	ΓIN				
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb			
Ear/Eye Preparations						
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN						
E /E 500 11 (11 1 1 1						

DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4.50	8 ml OP	
	(9.27)		Sofradex
FRAMYCETIN SULPHATE			
Ear/Eye drops 0.5%	4.13	8 ml OP	
, ,	(8.65)		Soframycin

Eye Preparations

Anti-Infactive Preparations

Eye preparations are only funded for use in the eye, unless explicitly stated otherwise.

Anti-infective r reparations
ACICI OVID

ACICLOVIR	14.88	4.5 g OP	✓ ViruPOS
CHLORAMPHENICOL		-	
Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%	1.54	10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * ar	e unapproved in	dications.	
CIPROFLOXACIN			
Eye drops 0.3% — Subsidy by endorsement	or severe bacterions media (CSOM)	al conjunctivitis	resistant to chloramphenicol; or
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
PROPAMIDINE ISETHIONATE			
* Eye drops 0.1%	2.97	10 ml OP	
	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID]			

5 g OP

✓ Fucithalmic

	Subsidy (Manufacturer's Price) Subs	Fully	Brand or Generic
	` \$	Per	1	Manufacturer
TOBRAMYCIN				
Eye oint 0.3%	10.45	3.5 g OP	√ T	obrex
Eye drops 0.3%	11.48	5 ml OP	✓ T	obrex
Corticosteroids and Other Anti-Inflan	nmatory Preparations			
DEXAMETHASONE				
* Eye oint 0.1%	5.86	3.5 g OP	✓ N	Maxidex
* Eye drops 0.1%		5 ml OP	✓ N	laxidex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has diabetic macular oedema with pseudophakic lens; and

Ocular implant 700 mcg - Special Authority see SA1680 below

- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

*	Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b			
	sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
*	Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DIC	CLOFENAC SODIUM			
	Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha

✓ Ozurdex

Prednisolone

✓ Rexacrom

	Subsidy (Manufacturer's Pr	ice) Sul	Fully	Brand or Generic
	\$	Per	✓	Manufacturer
FLUOROMETHOLONE				
* Eye drops 0.1%	3.09	5 ml OP	√ F	ML
	5.20		✓ F	lucon
KETOROLAC TROMETAMOL - Special Authority see SA1981	below – Retail pha	armacy		
Eye drops 0.5%	9.50 [·]	5 ml OP	✓ A	Acular
(Acular Eye drops 0.5% to be delisted 1 February 2022)				

⇒SA1981 Special Authority for Subsidy

Initial application — (macular oedema) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 The patient has established post-operative or inflammatory (uveitic) cystoid macular oedema; or
- 2 Both:
 - 2.1 The patient is at risk of postoperative macular oedema; and
 - 2.2 The patient has had, or is scheduled to have imminent cataract surgery.

LEVOCABASTINE

LEVOCABASTINE			
Eye drops 0.5 mg per ml	8.71	4 ml OP	
, , ,	(10.34)		Livostin
LODOXAMIDE			
Eye drops 0.1%	8.71	10 ml OP	✓ Lomide
NEPAFENAC			
Eve drops 0.3%	13.80	3 ml OP	✓ Ilevro
(Ilevro Eye drops 0.3% to be delisted 1 February 2022)			
PREDNISOLONE ACETATE			
Eye drops 1%	5.93	10 ml OP	✓ Prednisolone-AFT
As a share of	7.00	5 ml OP	✓ Pred Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority see	SA1715 below	- Retail pharma	acv
Eye drops 0.5%, single dose (preservative free)		20 dose	✓ Minims

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

5 ml OP

SODIUM CROMOGLICATE

(alaucoma Preparations - Beta Blockers		
BE	TAXOLOL		
*	Eye drops 0.25%	5 ml OP	✓ Betoptic S
	Eye drops 0.5%7.50	5 ml OP	✓ Betoptic
TII	MOLOL		•
*	Eye drops 0.25%	5 ml OP	✓ Arrow-Timolol
	Eye drops 0.5%	5 ml OP	✓ Arrow-Timolol
	Eye drops 0.5%, gel forming	2.5 ml OP	✓ Timoptol XE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Bran sidised Gene Man	
Glaucoma Preparations - Carbonic Anhydrase	Inhibitors			
CETAZOLAMIDE Tab 250 mg	17.03	100	✓ Diamos	¢
RINZOLAMIDE Eye drops 1% ORZOLAMIDE HYDROCHLORIDE	7.30	5 ml OP	✓ Azopt	
Eye drops 2%	9.77 (17.44)	5 ml OP	Trusopi	t
ORZOLAMIDE WITH TIMOLOL Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ Dortim	opt
Glaucoma Preparations - Prostaglandin Analo	gues			
IMATOPROST				
€ Eye drops 0.03%	5.95	3 ml OP	✓ Bimato Multi	prost chem
ATANOPROST € Eye drops 0.005% Teva to be Principal Supply on 1 February 2022	1.82	2.5 ml OP	✓ Teva	
RAVOPROST Eye drops 0.004%	7.30 9.75	5 ml OP 2.5 ml OP	✓ Travop ✓ Travata	
Travatan to be Principal Supply on 1 December 2021 Travopt Eye drops 0.004% to be delisted 1 December 2021) Mylan S29 Eye drops 0.004% to be delisted 1 December 202	10.50	5 ml OP	✓ Mylan	S29)
Glaucoma Preparations - Other				
RIMONIDINE TARTRATE Eye drops 0.2%Arrow-Brimonidine to be Principal Supply on 1 January		5 ml OP	✓ Arrow-	Brimonidine
RIMONIDINE TARTRATE WITH TIMOLOL MALEATE Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combi	gan
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ Arrow	- Lattim
Eye drops 1%	4.26	15 ml OP	✓ Isopto	Carpine
€ Eye drops 2%		15 ml OP	✓ Isopto	
Eye drops 4%	7.99	15 ml OP	✓ Isopto	Carpine
Subsidised for oral use pursuant to the Standard Formule Eye drops 2% single dose – Special Authority see SA0895				

continued...

Either:

Subsidy (Manufacturer's Price)	S	Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics	and C	Cyclop	legics
------------	-------	--------	--------

ATROPINE SULPHATE * Eye drops 1%	17.36	15 ml OP	✓ Atropt
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%	8.76	15 ml OP	✓ Cyclogyl
* Eye drops 1%, single dose (preservative free) - Only on a prescription	52.86	20 dose	✓ Minims Cyclopentolate
(Minims Cyclopentolate Eye drops 1%, single dose (preservative fr	ee) to be delis	ted 1 April 2022	, ,
TROPICAMIDE * Eye drops 0.5% * Eye drops 1%		15 ml OP 15 ml OP	✓ Mydriacyl✓ Mydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, page 247

HYPROMELLOSE

*	Eye drops 0.5%	15 ml OP	Methopt

HYPROMELLOSE WITH DEXTRAN

Preservative Free Ocular Lubricants

⇒SA1388 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye: and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eve drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA1388 above – Retail pharmacy
Ophthalmic gel 0.3%, 0.5 g8.25 30 ✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL – Special Authority see SA1388 above – Retail pharmacy

a) Hylo-Fresh has a 6 month expiry after opening. The Pharmacy Procedures Manual restriction allowing one bottle per month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.

- b) Hylo-Fresh to be Principal Supply on 1 January 2022
- ▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

SENSORY ORGANS

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
Other Eye Preparations				
NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	4.15	15 ml OP	✓ N	laphcon Forte
OLOPATADINE Eye drops 0.1%	2.20	5 ml OP	√ <u>0</u>	Diopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.63	3.5 g OP	√ P	oly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	3.80	5 g OP	✓ V	itA-POS

Brand or

Generic

Manufacturer

Fully

Subsidy (Manufacturer's Price) Subsidised Per

Agents Used in the Treatment of Poisonings

Antidotes

ACETYL CYSTEINE

Inj 200 mg per ml, 10 ml ampoule58.76 10 ✓ DBL Acetylcysteine ✓ Martindale Pharma \$29

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO
- ✓ DBL Naloxone 5 Hydrochloride

Removal and Elimination

CHARCOAL

*	Oral liq 50 g per 250 ml	43.50	250 ml OP	✓ Carbosorb-X
---	--------------------------	-------	-----------	---------------

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy Wastage claimable

vvaolage olaimable			
Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exiade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels. liver or cardiac MRI T2*: or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis: or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per µL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE – Special Authority see SA1480 on the nex	t page – Retail pharn	nacy	
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml		250 ml OP	✓ Ferriprox



Su	ubsidy F	ully	Brand or
(Manufac	cturer's Price) Subsid	ised	Generic
	\$ Per	✓	Manufacturer

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE * Inj 500 mg vial	84.53	10	✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
	(156.71)		Calcium Disodium Versenate

Standard Formulae

Standard i Officiale			
ACETYLCYSTEINE EYE DROPS		PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Acetylcysteine inj 200 mg per ml, 10 ml	qs	mg per ml)	
Suitable eye drop base	qs	Phenobarbitone Sodium	400 mg
		Glycerol BP	4 ml
CODEINE LINCTUS (3 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	60 mg		
Glycerol	40 ml	PILOCARPINE ORAL LIQUID	
Preservative	qs	Pilocarpine 4% eye drops	qs
Water	to 100 ml	Preservative	qs
CODEINE LINCTUS (15 mg nor 5 ml)		Water	to 500 ml
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate	300 mg	(Preservative should be used if quantity supplied is	for more
Glycerol	40 ml	than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 100 ml	Methylcellulose	5 g
vvalei	10 100 1111	Preservative	qs
FOLINIC MOUTHWASH		Water	to 500 ml
Calcium folinate 15 mg tab	1 tab	(Preservative should be used if quantity supplied is	
Preservative	qs	than 5 days. Maximum 500 ml per prescription.)	101 111010
Water	to 500 ml	than 5 days. Maximum 500 mi per presemption.	
(Preservative should be used if quantity supplied is	for more	SODIUM CHLORIDE ORAL LIQUID	
than 5 days. Maximum 500 ml per prescription.)		Sodium chloride inj 23.4%, 20 ml	qs
		Water	qs
METHADONE MIXTURE		(Only funded if prescribed for treatment of hyponatra	aemia)
Methadone powder	qs	VANCONAVCINI ODAL COLLITIONI (50 mm mm)	
Glycerol	qs	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	10 vials
Water	to 100 ml	Vancomycin 500 mg injection Glycerol BP	40 ml
METHYL HYDROXYBENZOATE 10% SOLUTION		Water	to 100 ml
Methyl hydroxybenzoate	10 g	(Only funded if prescribed for treatment of Clostridiu	
Propylene glycol	to 100 ml	following metronidazole failure)	iiii uiiiiciie
(Use 1 ml of the 10% solution per 100 ml of oral liqu		lollowing metrorildazole failure)	
(Ose 1 mil of the 10% solution per 100 mil of oral liqu	iiu iiiixtui <i>e)</i>	VOSOL EAR DROPS	
OMEPRAZOLE SUSPENSION		WITH HYDROCORTISONE POWDER 1%	
Omeprazole capules or powder	qs	Hydrocortisone powder	1%
Sodium bicarbonate powder BP	8.4 g	Vosol Ear Drops	to 35 ml
Water	to 100 ml	•	
PHENOBARBITONE ORAL LIQUID			
Phenobarbitone Sodium	1 g		
Glycerol BP	70 ml		
Water	to 100 ml		
TTAIOI	.5 100 1111		

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Pr	rico) Subr	Fully Brand or sidised Generic
	(Manufacturer 5 FT	Per	✓ Manufacturer
Extemporaneously Compounded Preparations	and Galenica	le	
CODEINE PHOSPHATE – Safety medicine; prescriber may dete			
Powder – Only in combination	(90.09)	25 g	Douglas
Only in extemporaneously compounded codeine linctus.	` '		Douglas
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the s determined.	upplier and will b	e delisted fror	
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination			
Only in extemporaneously compounded oral mixtures. Soln	20.00	100 ml	✓ Midwest
	30.00	100 1111	• Iviiuwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE - Only in combination			
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			4
* Liquid – Only in combination Only in extemporaneously compounded oral liquid prepa	3.23	500 ml	✓ healthE Glycerol BP
METHADONE HYDROCHLORIDE	rations.		
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing fre			
d) Extemporaneously compounded methadone will only be in	eimbursed at the	rate of the ch	eapest form available
(methadone powder, not methadone tablets). Powder	7.84	1 g	✓ AFT
METHYL HYDROXYBENZOATE	7.04	' 9	· All
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE		J	
Powder	36.95	100 g	✓ <u>MidWest</u>
Suspension – Only in combination		473 ml	✓ <u>Ora-Plus</u>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH.			4.4
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Onl	•	470	A Ove Bland
Suspension	30.95	473 ml	✓ <u>Ora-Blend</u>
PHENOBARBITONE SODIUM Powder – Only in combination	52 50	10 g	✓ MidWest
1 owder Only in combination	325.00	100 g	✓ MidWest
Only in children up to 12 years		3	
PROPYLENE GLYCOL			
Only in extemporaneously compounded methyl hydroxybenz			
Liq	11.25	500 ml	✓ Midwest
SODIUM BICARBONATE Powder PR Only in combination	10.05	E00 a	✓ Midwoot
Powder BP — Only in combination		500 g	✓ <u>Midwest</u>

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>M</u>	idwest	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price)

Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

400 a OP ✓ Polycal

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	9	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 251

Subsidy		Fully	Brand or
(Manufacturer's Price)	_ 8	Subsidised	Generic
\$	Per		Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	 Special Authority see SA1524 above – Hospital p 	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy Fully (Manufacturer's Price) Subsidised Per

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see	SA1095 above -	- Hospital pharm	acy [HP3]
Liquid	3.75	500 ml OP	✓ Glucerna Select
	7.50	1,000 ml OP	Diason RTH
DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA	1095 above – Ho	spital pharmacy	[HP3]
Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	1.78	237 ml OP	
	(2.10)		Sustagen Diabetic
	(2.10)	200 ml OP	Nutren Diabetes
(Suctagon Diabatic Liquid (vanilla) to be delicted 1 Ephruany 200	22)		

(Sustagen Diabetic Liquid (vanilla) to be delisted 1 February 2022)

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see \$	SA1525 above – Hospital pharma	ıcy [HP3]	
Powder	60.48	400 g OP	Monoger

Subsidy (Manufacturer's Price) Fully Subsidised Per Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Powder54.00 400 g OP

✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

Sub:	•	Fully	Brand or
(Manufactu		sidised	Generic
\$	S Per	•	Manufacturer

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA13 Liquid6.0		e – Hospital pharmacy [HP3] ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1379 Liquid		Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Autho pharmacy [HP3]	rity see SA1379 on the	previous page – Hospital
Liquid	00 500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see SA1379 or Liquid (strawberry)	30 200 ml OP	Hospital pharmacy [HP3] Fortini Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 on Liquid (chocolate) 1.0 Liquid (strawberry) 1.0 Liquid (vanilla) 1.3	200 ml OP 200 ml OP 200 ml OP 200 ml OP	ospital pharmacy [HP3] ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s pharmacy [HP3] Liquid (unflavoured)1.6	60 200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate)	30 200 ml OP	✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the pre- Powder43.6		oharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's P \$	rice) Subsi Per	Fully idised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML — Special Authority see SA1 Liquid		o <mark>us page</mark> – Hos 220 ml OP	✓ N	harmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110		s page – Hospi 237 ml OP	tal pha	rmacy [HP3]
Liquid (apricot) 125 ml Liquid (caramel) 125 ml	(3.31) 11.52	4 OP 4 OP	✓ F	NovaSource Renal Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LiquidLiquid		1,000 ml OP	
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority s	ee SA1377 above	– Hospital phar	macy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see	SA1377 above –	Hospital pharm	acy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special Au	uthority see SA137	7 above – Hosp	oital pharmacy [HP3]
Liquid	12.04	1,000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

(Mar	Subsidy nufacturer's Price)	Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm3); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

ENTERAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 257 - Hospital pharmacy [HP3]

Liquid.......7.00 1,000 ml OP ✓ Nutrison Energy

	Subsidy	F	fully Brand or
	(Manufacturer's		,
	(Маниасинен 3	Per	✓ Manufacturer
	Ψ	1 61	Wallulacturel
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 on	page 257 - Ho	spital pharmacy [H	HP3]
Liquid		250 ml OP	✓ Isosource Standard
Liquid			
	5.29	1,000 ml OP	✓ Nutrison Standard
			RTH
			✓ Osmolite RTH
ENTERN FEED WITH FIRRE COOKON AND CO	044050	0== 11	
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authority			
Liquid	5.29	1,000 ml OP	✓ Nutrison
			800 Complete
			Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority se	e SA1859 on	page 257 – Hospit	al pharmacy [HP3]
Liquid	5.29	1,000 ml OP	✓ Jevity RTH
1		,	✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority s	see SA1859 on	page 257 – Hosp	ital pharmacy [HP3]
Liquid	1.75	250 ml OP	✓ Ensure Plus HN
-1	7.00	1,000 ml OP	✓ Ensure Plus RTH
	7.00	1,000 1111 01	
			✓ Jevity HiCal RTH
			✓ Nutrison Energy
			Multi Fibre
OBAL FEED (DOMDED) Chasiel Authority and CA1050 on nor	o OE7 Hoonit	al nharmanı [LID0	1
ORAL FEED (POWDER) – Special Authority see SA1859 on pag			-
Powder (chocolate)	14.00	840 g OP	 Sustagen Hospital
			Formula Active
	26.00	850 g OP	✓ Ensure
Powder (vanilla)		840 g OP	✓ Sustagen Hospital
rowder (varilla)	14.00	640 y OF	• .
			Formula Active
	26.00	850 g OP	✓ Ensure
ODAL FEED 4 5KOAL/ALL Cresial Authority and CA4050 are no	057 H	:4-1 1 [] []	
ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on pa			
Additional subsidy by endorsement is available for patients be			
epidermolysis bullosa, or as exclusive enteral nutrition in child	Iren under the	age of 18 years fo	r the treatment of Crohn's
disease, or for patients with COPD and hypercapnia, defined			
endorsed accordingly.	uo o o = . u.uo	encocaming commin	.ge p. eeenpalenaet ze
· · · · · · · · · · · · · · · · · · ·			
Liquid (banana) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Limit (shared star) . High an artist of \$4.00 and \$000 art with	(1.20)		i ortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) Lligher subside of \$1.00 per 000 p			rondop
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 n			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
LINOISEMENT		ZUU IIII UF	Fautiaia
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml wi	th		
Endorsement		237 ml OP	
	(1.33)	- " -	Ensure Plus
		000 ml OD	LIIGUIC I IUS
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
	(1.20)		

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 257 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

Elquid (chocolate) Trighter Subsidy of \$1.20 per 200 hill with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — **(Cystic fibrosis)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy (Manufacturer's \$	Price) Subs	Fully idised	Brand or Generic Manufacturer
ENTERAL FEED 2 KCAL/ML – Special Authority see SA1195 o	n the previous p		harma	
	11.00	1,000 ml OP	_	nsure Two Cal HN RTH wo Cal HN RTH
(Two Cal HN RTH Liquid to be delisted 1 February 2022)			• 1	WO Cal HIN H I H
ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the Additional subsidy by endorsement is available for patients lepidermolysis bullosa. The prescription must be endorsed and in the control of the	peing bolus fed t			
Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with Endorsement	0.96 (1.90)	200 ml OP	T	wo Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FOOD THICKENER -	Special Authority see SA1106 above - Hospital pharma	cy [HP3]	
Powder	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	Feed Thickener
			Karicare Antamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

	Subsidy		Fully Brand or
	(Manufacturer's Pr \$	ice) Subsid	dised Generic ✓ Manufacturer
	-		- manaratara
GLUTEN FREE BAKING MIX - Special Authority see SA			harmacy [HP3]
Powder		1,000 g OP	
	(5.15)		Healtheries Simple
			Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA	1729 on the previous pag	je – Hospital ph	armacy [HP3]
Powder	3.93	1,000 g OP	
	(7.32)	•	NZB Low Gluten
	, ,		Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729	on the previous page – I	lospital pharma	acv [HP3]
Powder		2,000 g OP	20, [i ii 0]
1 OWGO!	(18.10)	2,000 g Oi	Horleys Flour
OLLITEN EDEE DAOTA Oracial Authoritana OA4700	(/	La carita La la caraca	*
GLUTEN FREE PASTA – Special Authority see SA1729			acy [HP3]
Buckwheat Spirals		250 g OP	
	(3.11)		Orgran
Corn and Vegetable Shells		250 g OP	
	(2.92)		Orgran
Corn and Vegetable Spirals	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP	
	(3.82)		Orgran
Rice and Corn Macaroni	2.00	250 g OP	
	(2.92)		Orgran
Rice and Corn Penne	2.00	250 g OP	
	(2.92)		Orgran
Rice and Maize Pasta Spirals		250 g OP	
	(2.92)		Orgran
Rice and Millet Spirals	2.00	250 g OP	
	(3.11)		Orgran
Rice and corn spaghetti noodles	2.00	375 g OP	-
	(2.92)	-	Orgran
Vegetable and Rice Spirals	2.00	250 g OP	-
·	(2.92)	-	Orgran
Italian long style spaghetti	2.00	220 g OP	-
, .	(3.11)	-	Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

SPECIAL FOODS Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer Supplements For Homocystinuria AMINOACID FORMULA WITHOUT METHIONINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 500 a OP ✓ XMET Maxamum Powder 461.94 Supplements For MSUD AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] ✓ MSUD Maxamum 500 q OP Supplements For PKU AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 75 OP ✓ Phlexv 10 30 PKU Anamix Junior Orange Powder (chocolate) 36 g sachet......393.00 ✓ PKU Anamix Junior 30 Chocolate Powder (unflavoured) 28 g sachets......936.00 30 ✓ PKU Lophlex Powder ✓ PKU Anamix Junior 30 ✓ PKU Anamix Junior Powder (vanilla) 36 g sachet393.00 30 Vanilla ✓ PKU Anamix Infant 400 g OP ✓ XP Maxamum 500 a OP 500 g OP ✓ XP Maxamum 125 ml OP ✓ PKU Anamix Junior LQ 125 ml OP ✓ PKU Anamix Junior ✓ PKU Anamix Junior 125 ml OP LQ

18 OP

30 OP

36 OP

60 OP

60 OP

60 OP

30 OP

30 OP

✓ Easiphen Liquid

✓ PKU Lophlex Sensation 20

✓ PKU Lophlex LQ 20

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 10

✓ PKU Lophlex LQ 20

✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3] 500 a OP ✓ Loprofin Mix

Liquid (forest berries), 250 ml carton......540.00

Liquid (juicy tropical) 125 ml......936.00

Liquid (juicy berries) 62.5 ml......939.00

Liquid (juicy orange) 125 ml936.00

	Subsidy (Manufacturer's Pr	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
LOW PROTEIN PASTA - Special Authority see SA1108 on page	e 263 – Hospital į	pharmacy [HP:	3]	
Animal shapes	11.91	500 g OP	√ L	.oprofin
Lasagne	5.95	250 g OP	√ L	.oprofin
Low protein rice pasta	11.91	500 g OP	√ L	.oprofin
Macaroni	5.95	250 g OP	√ L	.oprofin
Penne	11.91	500 g OP	√ L	.oprofin
Spaghetti	11.91	500 g OP	√ L	.oprofin
Spirals	11.91	500 g OP	√ L	.oprofin

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FOHMULA – Special Authority see SA1940 below – Hospital phar Powder43.60	400 g OP	✓ Alfamino
Powder (unflavoured)53.00	400 g OP	✓ Alfamino Junio✓ Elecare✓ Elecare LCP
		✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	 ✓ Neocate SYNE ✓ Elecare ✓ Neocate Junior Vanilla

⇒SA1940 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or

continued...

-			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Por 🗸	Manufacturer

- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	osidised	Generic	
\$	Per	1	Manufacturer	

- 2.2.3 Amino acid formula is required for a nutritional deficit; and
- 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis: or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

ENTERAL LIQUID PEPTIDE FORMULA -	- Special Authority see SA1953 below	v – Hospital phari	macy [HP3]
Liquid 1 kcal/ml	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml	15.68	500 ml OP	✓ Nutrini Peptisorb
			Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

continued...



Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Fither:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Sov milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome: or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia: or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued

recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1 ✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Vaccinations** BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm] For infants at increased risk of tuberculosis. Increased risk is defined as: 1) living in a house or family with a person with current or past history of TB; or 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or egual to 40 per 100,000 for 6 months or longer; or 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php. Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin), Danish strain 1331, live attenuated, vial with diluent.................0.00 10 ✓ BCG Vaccine DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm] Funded for any of the following criteria: 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or 5) A single dose for vaccination of patients aged from 65 years old; or 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or 7) For vaccination of previously unimmunised or partially immunised patients: or 8) For revaccination following immunosuppression; or 9) For boosting of patients with tetanus-prone wounds. Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg pertussis toxoid, 8 mcg pertussis filamentous 10 **Boostrix Boostrix** DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm] Funded for any of the following: 1) A single dose for children up to the age of 7 who have completed primary immunisation; or 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive 4) Five doses will be funded for children requiring solid organ transplantation. Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units

Infanrix IPV

Subsidised

Fully

Brand or

Generic

✓ <u>Havrix</u>✓ Havrix Junior

Subsidy

(Manufacturer's Price)

\$ Per	✓	Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUEN	IZAE TYPI	B VACCINE -
[Xpharm]		
Funded for patients meeting any of the following criteria:		
1) Up to four doses for children up to and under the age of 10 for primary immunisation;		
2) An additional four doses (as appropriate) are funded for (re-)immunisation for children		· ·
10 who are patients post haematopoietic stem cell transplantation, or chemotherapy;		splenectomy; pre- or
post solid organ transplant, renal dialysis and other severely immunosuppressive regi		
3) Up to five doses for children up to and under the age of 10 receiving solid organ trans	•	
Note: A course of up-to four vaccines is funded for catch up programmes for children (up to		0 , ,
to complete full primary immunisation. Please refer to the Immunisation Handbook for the	appropriate	e schedule for catch up
programmes.		
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous		
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,		
10 mcg hepatitis B surface antigen in 0.5 ml syringe0.00	✓ Infa	ınrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]	• 11110	IIIIA IICAU
One dose for patients meeting any of the following:		
The dose for patients meeting any or the following. The primary vaccination in children; or		
2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post h.	aematonoi	atic stam call
transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or		
or post cochlear implants, renal dialysis and other severely immunosuppressive regim		z organi iranopiani, pro
3) For use in testing for primary immunodeficiency diseases, on the recommendation of		medicine physician or
paediatrician.		7
·		
Haemophilus Influenzae type B polysaccharide 10 mcg		
conjugated to tetanus toxoid as carrier protein 20-40 mcg;		
prefilled syringe plus vial 0.5 ml0.00 1	✓ Hib	erix
HEPATITIS A VACCINE - [Xpharm]		
Funded for patients meeting any of the following criteria:		
1) Two vaccinations for use in transplant patients; or		
2) Two vaccinations for use in children with chronic liver disease; or		
3) One dose of vaccine for close contacts of known hepatitis A cases.		

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Inj 10 mo	B RECOMBINANT VACCINE - [Xpharm] cg per 0.5 ml prefilled syringe		1	√ E	Engerix-B
1) 2) 3) 4) 5) 6) 7)	ded for patients meeting any of the following criteri for household or sexual contacts of known acute for children born to mothers who are hepatitis B s for children up to and under the age of 18 years i serology and require additional vaccination or rec for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual inte for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC)	hepatitis B patients or h surface antigen (HBsAg nclusive who are considurire a primary course of rcourse; or	, pos derec	itive; or I not to have	e achieved a positive
10)	following needle stick injury.				
	cg per 1 ml prefilled syringeded for patients meeting any of the following criteri		1	√ <u>E</u>	Engerix-B
2) 3) 4) 5) 6) 7) 8) 9) 10) 11)	for household or sexual contacts of known acute for children born to mothers who are hepatitis B s for children up to and under the age of 18 years i serology and require additional vaccination or rec for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sexual inte for patients following immunosuppression; or for solid organ transplant patients; or for post-haematopoietic stem cell transplant (HSC following needle stick injury; or for dialysis patients; or for liver or kidney transplant patients.	surface antigen (HBsAg nclusive who are considurie a primary course of rcourse; or) pos derec	itive; or I not to have	•
Any of th	PILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND ne following:	,	- [Xpl	narm]	
2) Ma 1 2	eximum of two doses for children aged 14 years an eximum of three doses for patients meeting any of the People aged 15 to 26 years inclusive; or either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) patients: eximum of four doses for people aged 9 to 26 years	he following criteria: or	nerap	у	
Inj 270 m	ncg in 0.5 ml syringe	0.00	10	√ <u>(</u>	Gardasil 9

(1)	Subsidy		Fully	Brand or
	Manufacturer's Price)		Subsidised	Generic
	\$	Per	1	Manufacturer

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

- [Xpharm]......9.00 1 ✓ Afluria Quad Junior (2021 Formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

	Subsidy (Manufacturer's Price)	Sı	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer
Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	90.00	10		fluria Quad (2021 Formulation)

- a) Only on a prescription
- b) No patient co-payment payable

C)

A) INFLUENZA VACCINE - people 5 years and over

is available each year for patients aged 5 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (adjuvanted quadrivalent vaccine).......90.00 10 ✓ Fluad Quad (2021 Formulation)

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

- a) Only on a prescription
- b) No patient co-payment payable

c)

- INFLUENZA VACCINE people 65 years and over
 is available each year for patients aged 65 years and over
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine)	-		
[Xpharm]	9.00	1	✓ Influvac Tetra
			(2021 Formulation)

Subsidy	Fully	Brand or	
(Manufacturer's Price	ce) Subsidised	Generic	
\$	Per 🗸	Manufacturer	

A) INFLUENZA VACCINE - people 3 and 4 years of age (inclusive)

is available each year for patients aged 3 and 4 years of age (inclusive) who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 60 mcg in 0.5 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant: or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients; or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C 1 PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B.

7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00

✓ Synflorix

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies: or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note:	please refer to the Im	munisation Han	dbook for the	appropriate	schedule for	catch up	orogrammes
Inj 30.	8 mcg of pneumococc	al polysacchario	de serotypes 1	1, 3, 4,			

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	[Xpharm]			
Up to three doses (as appropriate) for patients with H chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochle All of the following:	tional asplenia, pre- or p	oost-solid o	organ t	ransplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)immuni b) Treatment is for a maximum of two doses; and c) Any of the following: 	sation; and			
i) on immunosuppressive therapy or radiatic immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following orgor vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more the prednisone of 2 mg/kg per day or greater, 20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or failuxii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	or gan transplantation (incl tts; or han two weeks, and wh or children who weigh r asthma treated with hig station; or ure; or	uding haer o are on a nore than	matopo n equiv 10 kg o	oietic stem cell transplant); valent daily dosage of on a total daily dosage of
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)		1	√ <u>P</u>	neumovax 23
For partially vaccinated or previously unvaccinated in 2) For revaccination following immunosuppression. Note: Please refer to the Immunisation Handbook for appr	dividuals; or	tch-up prod	aramm	es.
Inj 80D antigen units in 0.5 ml syringe		1 ' `	´ 🗸 <u>II</u>	
first dose to be administered in infants aged under 14 no vaccination being administered to children aged 2	•			
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	✓ <u>R</u>	<u>Iotarix</u>

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]		
Either:			
 Maximum of one dose for primary vaccination for eith 	ner:		
 a) Any infant born on or after 1 April 2016; or 			
b) For previously unvaccinated children turning 11	years old on or after 1	July 2017, who h	ave not previously had a
varicella infection (chickenpox), or			
2) Maximum of two doses for any of the following:			
a) Any of the following for non-immune patients:			
i) with chronic liver disease who may in futu	re be candidates for tra	nsplantation: or	
ii) with deteriorating renal function before tra		,	
iii) prior to solid organ transplant; or	p		
iv) prior to any elective immunosuppression*	or		

- iv) prior to any elective immunosuppression*, or
- v) for post exposure prophylaxis who are immune competent inpatients.; or
- b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
- c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
- d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
- e) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
- f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
- g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* imm	nosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater tha
28 da	S

✓ Varivax 10 ✓ Varivax

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] - [Xpharm] Funded for patients meeting either of the following criteria:

- 1) One dose for all people aged 65 years; or
- 2) One dose for all people aged between 66 and 80 years inclusive from 1 April 2018 and 31 December 2021.

✓ Zostavax ✓ Zostavax

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] Tubersol

- Symbols -	Agents for Parkinsonism and Related	Amsidine158
UK Synacthen83	Disorders 123	Amzoate31
3TC109	Agents Used in the Treatment of	Anaesthetics125
- A -	Poisonings245	Anagrelide hydrochloride158
A-Scabies70	Agrylin158	Analgesics 126
Abacavir sulphate109	Albendazole92	Anastrozole174
Abacavir sulphate with	Albey230	Anatrole
lamivudine109	Albustix80	Andriol Testocaps83
Abiraterone acetate171	Aldurazyme30	Androderm83
Acarbose11	Alecensa	ANI51
Accarb11	Alectinib	Anoro Ellipta234
Accuretic49	Alendronate sodium116	Antabuse149
Accuretic 1049	Alendronate sodium with	Antacids and Antiflatulents6
Accuretic 2049	colecalciferol116	Anthelmintics92
Acetazolamide242	Alfacalcidol34	Antiacne Preparations63
Acetec	Alfamino265	Antiallergy Preparations230
Acetic acid with 1, 2- propanediol	Alfamino Junior265	Antianaemics38
diacetate and	Alginic acid6	Antiandrogen Oral
benzethonium239	Alglucosidase alfa27	Contraceptives78
Acetic acid with hydroxyquinoline and	Alkeran154	Antiarrhythmics50
ricinoleic acid78	Alkeran S29154	Antibacterials92
Acetylcysteine245	Allersoothe231	Antibacterials Topical63
Aci-Jel78	Allmercap156	Anticholinergic Agents233
Aciclovir	Allopurinol	Anticholinesterases
Infection104	Alpha-Adrenoceptor Blockers48	Antidepressants
Sensory239	Alpha-Keri Lotion	Antidiarrhoeals6
Acidex6	Alphamox95	Antiepilepsy Drugs
Acipimox56	Alphamox 12595	Antifibrinolytics, Haemostatics and
Acitretin70	Alphamox 25095	Local Sclerosants39
Aclasta	Alprolix	Antifibrotics
Aclin	Alu-Tab6	Antifungals 99
Actemra218	Aluminium hydroxide6	Antifungals Topical64
Actinomycin D	Alvogen52	Antihistamines
Actrapid10	Amantadine hydrochloride	Antihypotensives51
Actrapid Penfill10	Ambrisentan59	Antimalarials101
Acular241	Ambrisentan Mylan59	Antimigraine Preparations
Acupan	Amiloride hydrochloride55	Antinausea and Vertigo Agents 137
Adalimumab	Amiloride hydrochloride with	Antipruritic Preparations65
Adapalene63	furosemide56	Antipsychotics
Adenuric	Amiloride hydrochloride with	Antiretrovirals107
ADR Cartridge 1.824	hydrochlorothiazide56	Antirheumatoid Agents115
Adrenaline58	Aminophylline236	Antispasmodics and Other Agents
Advantan66	Amiodarone hydrochloride50	Altering Gut Motility8
Advate42	Amisulpride138	Antithrombotic Agents42
Adynovate42	Amisulpride Mylan138	Antithymocyte globulin
Afinitor	Amitriptyline131	(equine)182
Aflibercept	Amlodipine53	Antitrichomonal Agents102
Afluria Quad	Amneal	Antituberculotics and
(2021 Formulation)	Amorolfine	Antileprotics
Afluria Quad Junior	Amoxicillin95	Antiulcerants8
(2021 Formulation)	Amoxicillin with clavulanic acid95	Antivirals104
AFT-Pyrazinamide103	Amphotericin B33	Anxiolytics141
Agents Affecting the	Amsacrine	Anzatax161
Renin-Angiotensin System 48	AmsaLyo158	Apidra11

Apidra SoloStar	11	Arrow-Timolol	241	Beclazone 100	231
Apo-Azithromycin	93	Arrow-Topiramate		Beclazone 250	
Apo-Bromocriptine	123	Arrow-Tramadol	131	Beclazone 50	231
Apo-Ciclopirox	64	Arsenic trioxide	158	Beclomethasone dipropionate	231
Apo-Clarithromycin		Asacol	7	Bee venom allergy treatment	230
Alimentary	9	Asamax	<mark>7</mark>	Bendamustine hydrochloride	152
Infection	94	Ascorbic acid	34	Bendrofluazide	56
Apo-Clomipramine	131	Aspen Adrenaline	58	Bendroflumethiazide	
Apo-Diclo SR	114	Aspirin		[Bendrofluazide]	56
Apo-Diltiazem CD	54	Blood	42	Benzathine benzylpenicillin	95
Apo-Doxazosin	48	Nervous	126	Benzatropine mesylate	
Apo-Folic Acid	39	Asthalin	233	Benzbromaron AL 100	120
Apo-Furosemide	55	Atazanavir sulphate	109	Benzbromarone	120
Apo-Gabapentin	133	Atenolol	51	Benztrop	124
Apo-Megestrol		Atenolol AFT	51	Benzydamine hydrochloride	33
Apo-Metoprolol	52	Atenolol AFT S29	51	Benzylpenicillin sodium [Penicillin	
Apo-Mirtazapine	132	ATGAM	182	G]	9
Apo-Nadolol		Ativan	141	Beta Cream	
Apo-Oxybutynin		Atomoxetine		Beta Ointment	65
Apo-Pindolol		Atorvastatin	57	Beta Scalp	72
Apo-Prazosin		Atropine sulphate		Beta-Adrenoceptor Agonists	233
Apo-Prazosin S29		Cardiovascular	50	Beta-Adrenoceptor Blockers	
Apo-Prednisone		Sensory	243	Betadine	
Apo-Primidone		Atropt		Betadine Skin Prep	
Apo-Propranolol		Atrovent	233	Betaferon	
Apo-Pyridoxine	34	AU Synacthen		Betahistine dihydrochloride	
Apo-Selegiline S29		Aubagio		Betaine	
Apo-Sumatriptan		Augmentin		Betaloc CR	52
Apo-Temozolomide		Aurorix		Betamethasone dipropionate	65
Apomorphine hydrochloride		AutoSoft 30		Betamethasone dipropionate with	
Aprepitant		AutoSoft 90		calcipotriol	
Apresoline		Avelox	97	Betamethasone sodium phosphate	
Aptamil AllerPro SYNEO 1		Avonex	142	with betamethasone acetate	
Aptamil AllerPro SYNEO 2		Avonex Pen	142	Betamethasone valerate	
Aptamil Gold+ Pepti Junior		Azacitidine	154	Betamethasone valerate with sodi	um
Aqueous cream		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Aratac		Azamun		Betaxolol	
Arava		Azathioprine		Betnovate	
Arginine		Azilect		Betoptic	
Aripiprazole		Azithromycin		Betoptic S	
Aripiprazole Sandoz	138	Azopt		Bexsero	
Aristocort		AZT		Bezafibrate	
Arrow - Lattim		- B -		Bezalip	
Arrow-Amitriptyline		B-D Micro-Fine	16	Bezalip Retard	
Arrow-Bendrofluazide		B-D Ultra Fine		Bicalutamide	
Arrow-Brimonidine		B-D Ultra Fine II		Bicillin LA	
Arrow-Diazepam		Bacillus Calmette-Guerin (BCG)		BiCNU	
Arrow-Doxorubicin		vaccine	182	Bicnu Heritage	
Arrow-Losartan &		Bacillus Calmette-Guerin	102	Bile and Liver Therapy	10
Hydrochlorothiazide	49	vaccine	270	Biltricide	
Arrow-Norfloxacin		Baclofen		Bimatoprost	
Arrow-Ornidazole		Bactroban		Bimatoprost Multichem	
Arrow-Quinapril 10		Balance		Binarex	
Arrow-Quinapril 20		Barrier Creams and Emollients		Binocrit	
Arrow-Quinapril 5		Basic AquaCream		Biodone	
Arrow-Roxithromycin		BCG Vaccine		Biodone Extra Forte	
A WELLOW I TO VIEW OF LINE OF		DOG VACORIO	<i>LI</i> U	DIOGOTIO EXITA I UITE	120

INDEX: Generic Chemicals and Brands

Biodone Forte128	Calci-Tab 50035	Ceftriaxone9
Bisacodyl27	Calcipotriol71	Ceftriaxone-AFT9
Bisoprolol fumarate51	Calcitonin81	Cefuroxime axetil9
Bisoprolol Mylan51	Calcitriol34	Celebrex11
BK Lotion68	Calcitriol-AFT34	Celecoxib11
Bleomycin sulphate158	Calcium carbonate6, 35	Celecoxib Pfizer11
Blood Colony-stimulating	Calcium Channel Blockers53	Celestone Chronodose8
Factors45	Calcium Disodium Versenate246	Cellcept17
Blood glucose diagnostic test	Calcium folinate	Centrally-Acting Agents5
meter	Calcium Folinate Ebewe155	Cephalexin ABM9
Blood glucose diagnostic test	Calcium Folinate Sandoz155	Cetirizine hydrochloride23
strip	Calcium Folinate Sandoz S29 155	Cetomacrogol6
Blood glucose test strips (visually	Calcium gluconate35	Cetomacrogol with glycerol6
impaired)15	Calcium Homeostasis81	Cetuximab19
Blood Ketone Diagnostic Test	Calcium polystyrene sulphonate47	Champix15
·	Calcium Resonium47	Charcoal24
Strip	Calcium-Sandoz Forte35	
Bonjela33		Chemotherapeutic Agents
Boostrix270	Calogen	Chickenpox vaccine28
Bortezomib	Calutide-50	Chlorafast
Bortezomib Dr Reddy's S29 158	Candesartan cilexetil49	Chlorambucil15
Bortezomib Dr-Reddy's	Candestar49	Chloramphenicol23
Bortezomib Juno	Canesten64	Chlorothiazide5
Bortezomib Juno S29158	Capecitabine155	Chlorpheniramine maleate23
Bosentan59	Capercit	Chlorpromazine hydrochloride 13
Bosentan Dr Reddy's59	Capoten48	Chlortalidone [Chlorthalidone]5
Bplex34	Capsaicin	Chlorthalidone5
Breo Ellipta232	Musculoskeletal115	Chlorvescent4
Brevinor 1/2877	Nervous126	Choice Load 3757
Bricanyl Turbuhaler233	Captopril48	Choice TT380 Short7
Brilinta43	Captopril-Mylan48	Choice TT380 Standard7
Brimonidine tartrate242	Carafate10	Choline salicylate with cetalkonium
Brimonidine tartrate with timolol	Carbaccord	chloride3
maleate242	Carbamazepine 133	Ciclopirox olamine6
Brinzolamide242	Carbimazole86	Ciclosporin22
Brolene239	Carbomer243	Cilazapril4
Bromocriptine mesylate123	Carboplatin 153	Cilicaine9
Brufen SR114	Carboplatin Ebewe153	Cilicaine VK9
Buccastem138	Carbosorb-X245	Cinacalcet8
Budesonide	Cardinol LA53	Cipflox9
Alimentary6	Cardizem CD54	Ciprofloxacin
Respiratory231, 237	CareSens Dual14	Infection9
Budesonide with eformoterol232	CareSens N15	Sensory23
Bumetanide55	CareSens N POP15	Ciprofloxacin Teva23
Buprenorphine Naloxone BNM 148	CareSens N Premier15	Circadin14
Buprenorphine with naloxone148	CareSens PRO15	Cisplatin15
Bupropion hydrochloride149	Carmellose sodium with gelatin and	Cisplatin Ebewe15
Burinex55	pectin33	Citalopram hydrobromide13
Burinex S29	Carmustine	Cladribine15
Buscopan8	Carvedilol	Clarithromycin
Buspirone hydrochloride141	Carvedilol Sandoz	Alimentary
Busulfan153		Infection9
- C -	Catapres	Clexane4
_	Cefaclor monohydrate	Clexane Forte4
Cabergoline91		Climara8
Caffeine citrate	Cefalexin92 Cefalexin Sandoz92	Climara8
Calamine	Cefazolin	
Calamine	Gerazoiiri92	Clindamycin9

Clinicians	28	Condyline	73	Daunorubicin	159
Clinicians Renal Vit		Contraceptives - Hormonal		David One Step Cassette Pregna	
Clobazam		Contraceptives - Non-hormonal.		Test	
Clobetasol propionate		Copaxone		DBL Acetylcysteine	
Clobetasone butyrate		Corticosteroids and Related Age		DBL Adrenaline	
		-		DBL Aminophylline	
Clofazimine	102	for Systemic Use		DBL Bleomycin Sulfate	
	64	Corticosteroids Topical			
Dermatological		Contifoam		DBL Carboplatin	
Genito-Urinary		Cosentyx		DBL Cisplatin	
Clomifene citrate		Cosmegen		DBL Dacarbazine	
Clomipramine hydrochloride		Coumadin		DBL Desferrioxamine Mesylate for	,
Clomipramine Teva		Country Life		BP	
Clonazepam		Coversyl		DBL Docetaxel	
Clonidine		Creon 10000		DBL Ergometrine	
Clonidine BNM		Creon 25000		DBL Gemcitabine	
Clonidine hydrochloride		Creon Micro		DBL Gentamicin	
Clopidogrel		Crotamiton		DBL Heparin Sodium	
Clopidogrel Multichem		Crystaderm		DBL Leucovorin Calcium	
Clopine		Curam		DBL Methotrexate Onco-Vial	
Clopixol	. 139, 141	Curam Duo 500/125		DBL Morphine Sulphate	
Clotrimazole		Cvite	34	DBL Naloxone Hydrochloride	245
Dermatological	64	Cyclizine hydrochloride		DBL Octreotide	173
Genito-Urinary	78	Cyclizine lactate	137	DBL Pethidine Hydrochloride	130
Clozapine	138	Cyclogyl	243	DBL Vincristine Sulfate	164
Clozaril	138	Cyclonex		Decozol	
Co-trimoxazole	99	Cyclopentolate hydrochloride	243	Deferasirox	245
Coal tar	71	Cyclophosphamide		Deferiprone	
Coal tar with allantoin, menthol	,	Cyclorin		Denosumab	
phenol and sulphur		Cycloserine		Deolate	101
Coal tar with salicylic acid and		Cyproterone acetate		Deoxycoformycin	162
sulphur	71	Cyproterone acetate with		Depo-Medrol	
Coco-Scalp		ethinyloestradiol	78	Depo-Provera	
Codeine phosphate		Cystadane		Depo-Testosterone	
Extemporaneous	248	Cytarabine		Deprim	
Nervous		Cytotec		Dermol	
Coenzyme Q10		Cytoxan		Desferrioxamine mesilate	
Colchicine		- D -		Desmopressin	
Colecalciferol		D-Penamine	115	Desmopressin acetate	
Colestid		Dabigatran		Desmopressin-PH&T	
Colestipol hydrochloride		Dacarbazine		Desuric	
Colgout		Dacarbazine APP		Detection of Substances in	
Colifoam		Dactinomycin [Actinomycin D]		Urine	80
Colistin sulphomethate		Daivobet		Dexamethasone	
Colistin-Link		Daivonex		Hormone	82
Collodion flexible		Daktarin		Sensory	
Colloidal bismuth subcitrate		Dalacin C		Dexamethasone phosphate	
Colofac		Dantrium		Dexamethasone Phosphate	02
		Dantrium S29		Panpharma	90
Combigan					
Combigan	242	Dantrolene		Dexamethasone with framycetin	
Compound electrolytes		Daonil		gramicidin	239
Compound electrolytes with glu		Dapa-Tabs		Dexamethasone with neomycin	
[Dextrose]		Dapsone		sulphate and polymyxin B	040
Compound hydroxybenzoate		Daraprim		sulphate	
Comtan		Darunavir		Dexamfetamine sulfate	
Concerta		Darunavir Mylan		Dexmethsone	82
Condoms	75	Dasatinib	165	Dextrochlorpheniramine	

INDEX: Generic Chemicals and Brands

maleate231	Dorzolamide hydrochloride242	Eltrombopag	3
Dextrose46–47	Dorzolamide with timolol242	Eltroxin	
DHC Continus128	Dostinex91	EMB Fatol	10
Diabetes10	Dosulepin [Dothiepin]	Emend Tri-Pack	13
Diabetes Management14	hydrochloride131	Emicizumab	4
Diacomit	Dosulepin Mylan131	EMLA	12
Diagnostic Agents281	Dothiepin131	Empagliflozin	1
Diamide Relief6	Doxazosin48	Empagliflozin with metformin	
Diamox242	Doxine96	hydrochloride	1
Diasip253	Doxorubicin Ebewe159	Emtricitabine	
Diason RTH253	Doxorubicin hydrochloride159	Emtricitabine with tenofovir	
Diazepam133, 141	Doxycycline96	disoproxil	10
Diazoxide10	DP Lotion68	Emtriva	
Dibenzyline48	DP Lotn HC66	Emulsifying ointment	6
Diclofenac Sandoz114	DP-Allopurinol120	Emulsifying Ointment ADE	
Diclofenac sodium	Dr Reddy's Omeprazole9	Enalapril maleate	
Musculoskeletal114	Drofate53	Enbrel	
Sensory240	Drugs Affecting Bone	Endocrine Therapy	17
Differin	Metabolism 116	Endoxan	
Difflam33	Dual blood glucose and blood ketone	Engerix-B	
Diflucan99	diagnostic test meter 14	Enlafax XR	
Digestives Including Enzymes25	Dulaglutide13	Enoxaparin sodium	
Digoxin50	Dulcolax SP Drop27	Enstilar	7
Dihydrocodeine tartrate128	Duocal Super Soluble Powder251	Ensure	
Dilantin	Duolin233	Ensure Plus	
Dilantin Infatab134	Duolin HFA233	Ensure Plus HN	
Diltiazem hydrochloride54	DuoResp Spiromax232	Ensure Plus RTH	26
Dilzem54	Duride58	Ensure Two Cal HN RTH	
Dimethicone67–68	-E-	Entacapone	12
Dimethyl fumarate142	e-chamber La Grande237	Entapone	12
Dipentum7	e-chamber Mask237	Entecavir	
Diphtheria, tetanus and pertussis	e-chamber Turbo237	Entecavir Sandoz	
vaccine	E-Mycin94	Entocort CIR	
Diphtheria, tetanus, pertussis and	Ear Preparations239	Entresto 24/26	
polio vaccine270	Ear/Eye Preparations239	Entresto 49/51	4
Diphtheria, tetanus, pertussis, polio,			
Dipininchia, icianas, penassis, pono,	Easiphen Liquid264	Entresto 97/103	4
	Easiphen Liquid264 Econazole nitrate64	Entresto 97/103	4
hepatitis B and haemophilus	Easiphen Liquid 264 Econazole nitrate 64 Efavirenz 108	Epilim	4
hepatitis B and haemophilus influenzae type B vaccine 271	Econazole nitrate 64 Efavirenz 108	Epilim Epilim Crushable	4 13
hepatitis B and haemophilus influenzae type B vaccine271 Diprosone	Econazole nitrate	Epilim Epilim Crushable Epilim IV	13 13 13
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	EpilimEpilim CrushableEpilim IVEpilim IV	4 13 13 13
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim Epilim Crushable Epilim IV Epilim S/F Liquid Epilim Syrup	4 13 13 13
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim	4 13 13 13 15
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim	4 13 13 13 15
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim	4 13 13 13 15 15
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim	4 13 13 13 15 15
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate	Epilim	4 13 13 13 15 15
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29	Epilim	4 13 13 13 15 15 5 5
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124	Epilim	4 13 13 13 15 15 15 5 6
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265	Epilim	43 13 13 15 15 15 15 3 6
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265	Epilim	
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265 Electral 47	Epilim	
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265 Electral 47 Elelyso 32	Epilim	
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265 Electral 47 Elelyso 32 Elemental 028 Extra 256	Epilim	
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265 Electral 47 Elelyso 32 Elemental 028 Extra 256 Elidel 71	Epilim	
hepatitis B and haemophilus influenzae type B vaccine	Econazole nitrate 64 Efavirenz 108 Efavirenz with emtricitabine and tenofovir disoproxil 109 Eformoterol fumarate 232 Eformoterol fumarate dihydrate 232 Eftrenonacog alfa [Recombinant factor IX] 39 Efudix 73 Egopsoryl TA 71 Elaprase 29 Eldepryl 124 Elecare 265 Elecare LCP 265 Electral 47 Elelyso 32 Elemental 028 Extra 256	Epilim	

Esbriet	235	Femme-Tab ED	77	Flutamide	17
Escitalopram	132	Fentanyl	128	Flutamin	17
Escitalopram (Ethics)	132	Fentanyl Sandoz	128	Fluticasone	23
Eskazole		Ferinject	36	Fluticasone furoate with	
Essential Generics	51	Ferodan	36	vilanterol	23
Essential Prednisolone	8	Ferriprox		Fluticasone propionate	
Estradiol TDP Mylan	84	Ferro-F-Tabs		Fluticasone with salmeterol	
Estradot		Ferro-tab		FML	24
Estradot 50 mcg	84	Ferrograd	36	Foban	6
Estrofem	84	Ferrosig	36	Folic acid	39
Etanercept		Ferrous fumarate		Folic Acid multichem	39
Ethambutol hydrochloride	102	Ferrous fumarate with folic acid	l36	Folic Acid Mylan	
Ethics Aspirin		Ferrous sulfate	36	Food Thickeners	
Ethics Aspirin EC		Fexofenadine hydrochloride	231	Foods And Supplements For Ir	ıborn
Ethics Lisinopril		Fibro-vein		Errors Of Metabolism	
Ethics Paracetamol Classic		Filgrastim		Foradil	
Ethinyloestradiol		Finasteride		Forteo	
Ethinyloestradiol with		Fingolimod		Fortini	25
desogestrel	76	Firazyr		Fortini Multi Fibre	
Ethinyloestradiol with		Flagyl		Fortisip	
levonorgestrel	77	Flagyl-S		Fortisip Multi Fibre	
Ethinyloestradiol with		Flamazine		Fosamax	
norethisterone	77	Flecainide acetate		Fosamax Plus	
Ethosuximide		Flecainide BNM		Framycetin sulphate	
Etopophos		Flecainide Controlled Release		Frisium	
Etoposide		Teva	50	Frumil	
Etoposide phosphate		Fleet Phosphate Enema		Frusemide	
Etravirine		Flixonase Hayfever & Allergy		Fucicort	
Eumovate		Flixotide		Fucidin	
Everet		Flixotide Accuhaler		Fucithalmic	
Everolimus		Florinef		Fulvestrant	
Evista		Fluad Quad		Fungilin	
Exemestane		(2021 Formulation)	274	Furosemid-Ratiopharm	
Exjade		Fluanxol		Furosemide [Frusemide]	
Extemporaneously Compounde		Flucil		Furosemide-Baxter	
Preparations and	·u	Flucloxacillin		fusidic acid	
Galenicals	2/18	Flucloxin		Dermatological	64 6
Eye Preparations		Flucon		Infection	
Eylea		Fluconazole		Sensory	
Ezetimibe		Fludara Oral		- G -	20
Ezetimibe Sandoz		Fludarabine Ebewe		Gabapentin	131
Ezetimibe with simvastatin		Fludarabine phosphate		Gacet	
- F -		Fludrocortisone acetate		Galsulfase	
Factor eight inhibitor bypassing		Fluids and Electrolytes		Galvumet	
fraction	//1	Flumetasone pivalate		Galvus	
Famotidine		Fluocortolone caproate with	203	Gardasil 9	
Famotidine Hovid		fluocortolone pivalate and		Gastrodenol	
Faslodex		cinchocaine	0	Gaviscon Double Strength	
Febuxostat		Fluorometholone		Gaviscon Infant	
Febuxostat multichem		Fluorometholone		Gazyva	200
Feed Thickener Karicare	121	Fluorouracil Accord			
	060			Gefitinib	
Aptamil		Fluorouracil Ebewe		GEM Aqueous Cream	
		Fluorouracil sodium		Gemcitabine Ebewe	
Felo 10 ER				Gemcitabine hydrochloride	
Felo 5 ER		Fluoxetine hydrochloride Fluoenthixol decanoate		Genoptic	23
reiodibine	ექ	FINDERITRIXOLUECANOATE	139		

INDEX: Generic Chemicals and Brands

Gentamicin sulphate	Healtheries Simple Baking Mix263	[hydroxycarbamide] 16
Infection97	Hemastix80	Hygroton
Sensory239	Hemlibra40	Hylo-Fresh24
Gilenya142	Heparin sodium45	Hymenoptera23
Ginet78	Heparinised saline45	Hyoscine butylbromide
Glatiramer acetate142	Heparon Junior254	Hyoscine hydrobromide13
Glecaprevir with pibrentasvir 105	Hepatitis A vaccine271	Hypam14
Glibenclamide11	Hepatitis B recombinant	Hyperuricaemia and Antigout 12
Gliclazide12	vaccine 272	Hypromellose24
Glipizide12	Herceptin222	Hypromellose with dextran24
Glivec167	Hiberix271	-1-
Glizide12	Hiprex112	Ibiamox
Glucagen Hypokit10	Histaclear231	Ibrance16
Glucagon hydrochloride10	Histafen231	Ibuprofen11
Glucerna Select253	Holoxan154	Ibuprofen SR BNM11
Glucobay11	Horleys Bread Mix263	Icatibant23
Glucose [Dextrose]46	Horleys Flour263	Idarubicin hydrochloride16
Gluten Free Foods262	Hormone Replacement Therapy -	Idursulfase2
Glycerin with sodium saccharin248	Systemic84	Ifosfamide15
Glycerin with sucrose248	HPV272	Igroton
Glycerol	Humalog11	lkorel
Alimentary27	Humalog Mix 2511	llevro24
Extemporaneous248	Humalog Mix 5011	lloprost
Glyceryl trinitrate	Human papillomavirus (6, 11, 16, 18,	Imatinib mesilate
Alimentary8	31, 33, 45, 52 and 58) vaccine	Imatinib-Rex16
Cardiovascular58	[HPV]272	Imigran13
Glycopyrronium234	Humatin98	Imipramine hydrochloride13
Glycopyrronium bromide8	Humira182	Imiquimod
Glycopyrronium with	HumiraPen182	Immune Modulators11
indacaterol234	Humulin 30/7011	Immunosuppressants17
Go Healthy29	Humulin NPH10	Imuran17
Gold Knight75	Humulin R10	Incruse Ellipta23
Gold Knight XL75–76	Hyaluronic acid243	Indacaterol23
Goserelin90	Hydralazine58	Indapamide
Gutron51	Hydralazine hydrochloride58	Infanrix IPV27
Gynaecological Anti-infectives78	Hydrocortisone	Infanrix-hexa27
- H -	Dermatological66	Infant Formulae
Habitrol 150	Hormone82	Infatrini
Haemophilus influenzae type B	Hydrocortisone (PSM)66	Infliximab19
vaccine271	Hydrocortisone acetate7	Influenza vaccine27
Haldol140	Hydrocortisone acetate with	Influvac Tetra
Haldol Concentrate140	pramoxine hydrochloride7	(2021 Formulation)
Haldol Decanoas140	Hydrocortisone and paraffin liquid	Inhaled Corticosteroids23
Haloperidol138	and lanolin	Inhaled Long-acting
Haloperidol decanoate140	Hydrocortisone butyrate66, 72	Beta-adrenoceptor Agonists 23
Harvoni	Hydrocortisone with cinchocaine8	Inspra
Havrix271	Hydrocortisone with miconazole66	Instillagel Lido12
Havrix Junior271	Hydrocortisone with natamycin and	Insulin aspart1
Haylor syrup231	neomycin	Insulin aspart with insulin aspart
healthE Calamine Aqueous Cream	Hydrogen peroxide63	protamine1
BP65	Hydroxocobalamin34	Insulin glargine1
healthE Dimethicone 10%67	Hydroxocobalamin Mercury	Insulin glulisine
healthE Dimethicone 4% Lotion68	Pharma34	Insulin isophane1
healthE Dimethicone 5%	hydroxycarbamide160	Insulin isophane with insulin
healthE Glycerol BP248	Hydroxychloroquine115	neutral1
healthE Urea Cream	Hydroxyurea	Insulin lispro1
110attile 010a 010a11100	riyaronyaroa	111041111 110p10

Insulin lispro with insulin lispro	Itch-Soothe	65	Lanoxin Paediatric Elixir	
protamine11	Itraconazole	100	Lanoxin PG	
Insulin neutral10	Itrazole	100	Lanoxin S29	
Insulin pen needles16	Ivacaftor	236	Lansoprazole	
Insulin pump 16	Ivermectin	69	Lantus	
Insulin pump cartridge21	- J -		Lantus SoloStar	1
Insulin pump infusion set (steel	Jadelle	77	Lanvis	158
cannula)21	Jakavi	169	Lanzol Relief	
Insulin pump infusion set (steel	Jardiamet	13	Lapatinib ditosylate	16
cannula, straight insertion) 22	Jardiance	13	Largactil	13
Insulin pump infusion set (teflon	Jaydess	85	Laronidase	30
cannula)23	Jevity HiCal RTH	260	Lasix	5
Insulin pump infusion set (teflon	Jevity RTH		Latanoprost	24
cannula, angle insertion with	Juno Pemetrexed		Latanoprost with timolol	
insertion device)24	- K -		Lax-Suppositories	2
Insulin pump infusion set (teflon	Kadcyla	223	Lax-Tab	
cannula, angle insertion) 24	Kaletra	109	Laxatives	20
Insulin pump infusion set (teflon	Kalydeco	236	Laxsol	2
cannula, straight insertion with	Kemadrin		Ledipasvir with sofosbuvir	
insertion device)24	Kenacomb	239	Leflunomide	
Insulin pump infusion set (teflon	Kenacort-A 10	<mark>83</mark>	Lenalidomide	
cannula, straight insertion) 24	Kenacort-A 40		Letrole	
Insulin pump reservoir24	Kenalog in Orabase		Letrozole	
Insulin syringes, disposable with	Kenkay Sorbolene		Leukeran FC	
attached needle	Ketocal 3:1		Leukotriene Receptor	
Intelence	KetoCal 4:1		Antagonists	23
Interferon beta-1-alpha142	Ketoconazole		Leuprorelin	9
Interferon beta-1-beta143	Dermatological	72	Leustatin	
Intra-uterine device76	Infection		Levetiracetam	
Invega Sustenna140	Ketogenic Diet		Levetiracetam-AFT	
IPCA-Frusemide55	Ketoprofen		Levlen ED	
IPCA-Metoprolol52	Ketorolac trometamol		Levocabastine	
IPCA-Propranolol53	KetoSens		Levocarnitine	
IPOL280	Keytruda		Levodopa with benserazide	
Ipratropium bromide233, 237	Kindergen		Levodopa with carbidopa	
Iressa	Kivexa		Levomepromazine	
Irinotecan Accord	Klacid		Levomepromazine	
Irinotecan Actavis 100	Alimentary	9	hydrochloride	139
Irinotecan hydrochloride	Infection		Levonorgestrel	
Irinotecan-Rex	Kliogest		Genito-Urinary	77_7
Iron (as ferric carboxymaltose)36	Kliovance		Hormone	
Iron polymaltose36	Kogenate FS		Levothyroxine	
Isentress	Konakion MM		Lidocaine [Lignocaine]	
Isentress HD110	Konsyl-D		Lidocaine [Lignocaine]	120
Ismo 20	Kuvan		hydrochloride	12
Ismo 40 Retard58	-L-		Lidocaine [Lignocaine] with	120
Isoniazid	Labetalol	52	chlorhexidine	12
Isoniazid with rifampicin103	Lacosamide		Lidocaine [Lignocaine] with	
Isoptin	Lactulose		prilocaine	
Isoptin Retard54	Laevolac		Lidocaine-Baxter	12
Isoptin SR	Lamictal		Lidocaine-Claris	
Isopto Carpine242	Lamivudine		Life Extension	
Isosorbide mononitrate	Lamivudine Alphapharm		Lignocaine	
Isosource Standard260	Lamotrigine		Lioresal Intrathecal	
Isotretinoin	Lamprene		Lipid-Modifying Agents	
Ispaghula (psyllium) husk26	Lamprene		Liquigen	
iopagnula (poyillulli) nuok20	Lanoxiii		Liquigett	404

Lisinopril	48	Martindale Pharma245	Methyldopa Mylan	5
Lithium carbonate	139	Mask for spacer device237	Methyldopa Mylan S29	
Livostin	241	Maviret105	Methylnaltrexone bromide	2
LMX4	125	Maxidex240	Methylphenidate ER - Teva	14
Locacorten-Viaform ED's	239	Maxitrol240	Methylphenidate hydrochloride	
Local preparations for Anal and		MCT oil (Nutricia)252	Methylphenidate hydrochloride	
Rectal Disorders		Measles, mumps and rubella	extended-release	14
Locasol	265	vaccine277	Methylprednisolone	
Locoid	66, 72	Mebendazole92	Methylprednisolone (as sodium	
Locoid Crelo	66	Mebeverine hydrochloride8	succinate)	8
Locoid Lipocream	66	Medco	Methylprednisolone aceponate	6
Locorten-Vioform	239	Dermatological67	Methylprednisolone acetate	8
Lodoxamide	241	Nervous127	Methylxanthines	23
Logem	134	Medrol82	Metoclopramide Actavis 10	13
Lomide	241	Medroxyprogesterone acetate	Metoclopramide hydrochloride	13
Lomustine	154	Genito-Urinary77	Metolazone	5
Loniten	59	Hormone85	Metopirone	
Loperamide hydrochloride	6	Mefenamic acid114	Metoprolol IV Mylan	5
Lopinavir with ritonavir	109	Megestrol acetate 172	Metoprolol succinate	
Lopinavir/Ritonavir Mylan	109	Melatonin143	Metoprolol tartrate	5
Loprofin	265	Melphalan154	Metrogyl	
Loprofin Mix		Menactra277	Metronidazole	10
Lorafix	231	Meningococcal (groups A, C, Y and	Metyrapone	
Loratadine		W-135) conjugate vaccine 277	Mexiletine hydrochloride	
Lorazepam	141	Meningococcal B multicomponent	Mexiletine Hydrochloride USP	5
Lorstat	57	vaccine 278	Miacalcic	8
Losartan Actavis	49	Meningococcal C conjugate	Micolette	2
Losartan potassium	49	vaccine 278	Miconazole	
Losartan potassium with		Menthol65	Miconazole nitrate	
hydrochlorothiazide	49	Mepolizumab201	Dermatological	6
Lovir	104	Mercaptopurine156	Genito-Urinary	7
Loxamine	132	Mercilon 2876	Micreme	7
Lucrin Depot 1-month	91	Mesalazine7	Micreme H	6
Lucrin Depot 3-month	91	Mesna161	Microgynon 20 ED	7
Lyderm	70	Mestinon114	Microgynon 30	<mark>7</mark>
Lynparza	161	Metabolic Disorder Agents27	Microgynon 50 ED	
Lyrica	134	Metformin hydrochloride12	Microlut	
- M -		Metformin Mylan12	Midazolam	14
m-Eslon	129	Methadone hydrochloride	Midazolam-Baxter	14
Mabthera	204	Extemporaneous248	Midodrine	5
Macrobid	113	Nervous128	Mifegyne	8
Macrogol 3350 with potassium		Methatabs128	Mifepristone	8
chloride, sodium bicarbonate	and	Methenamine (hexamine)	Minerals	3
sodium chloride	<mark>27</mark>	hippurate112	Mini-Wright AFS Low Range	23
Macrogol 400 and propylene		Methopt243	Mini-Wright Standard	23
glycol	243	Methotrexate157	Minidiab	1
Madopar 125	123	Methotrexate DBL Onco-Vial 157	MiniMed 1.8 Reservoir	
Madopar 250	123	Methotrexate Ebewe157	MMT-326A	2
Madopar 62.5		Methotrexate Sandoz157	MiniMed 3.0 Reservoir	
Madopar HBS	123	Methyl hydroxybenzoate248	MMT-332A	2
Madopar Rapid		Methylcellulose248	MiniMed 640G	
Magnesium hydroxide	37	Methylcellulose with glycerin and	MiniMed 770G	1
Magnesium sulphate	37	sodium saccharin248	MiniMed Mio MMT-921A	2
Mantoux		Methylcellulose with glycerin and	MiniMed Mio MMT-923A	2
Marevan		sucrose248	MiniMed Mio MMT-925A	
Marine Blue Lotion SPF 50+	72	Methyldopa55	MiniMed Mio MMT-941A	2

MiniMed Mio MMT-943A	23	Mouth and Throat	33	Neocate Junior Unflavoured	26
MiniMed Mio MMT-945A	23	Movapo	123	Neocate Junior Vanilla	26
MiniMed Mio MMT-965A	23	Moxifloxacin		Neocate SYNEO	26
MiniMed Mio MMT-975A	23	MSUD Maxamum	264	Neoral	22
MiniMed Quick-Set MMT-386A		Mucilaginous laxatives with		Neostigmine metilsulfate	
MiniMed Quick-Set MMT-387A		stimulants	26	Nepafenac	
MiniMed Quick-Set MMT-396A	23	Mucolytics	236	Nepro HP (strawberry)	
MiniMed Quick-Set MMT-397A	23	Mucosoothe		Nepro HP (vanilla)	
MiniMed Quick-Set MMT-398A	23	Multiple Sclerosis Treatments		Nepro HP RTH	
MiniMed Quick-Set MMT-399A	23	Multivitamin renal		Neulactil	
MiniMed Silhouette MMT-368A	23	Multivitamins	35	Neulastim	
MiniMed Silhouette MMT-377A	23	Mupirocin	64	NeuroTabs	3
MiniMed Silhouette MMT-378A	23	Muscle Relaxants	122	Nevirapine	10
MiniMed Silhouette MMT-381A	23	Mvite	35	Nevirapine Alphapharm	10
MiniMed Silhouette MMT-382A	23	Myambutol	102	Nicorandil	5
MiniMed Silhouette MMT-383A	23	Mycobutin		Nicotine	
MiniMed Silhouette MMT-384A	23	MycoNail		Nifedipine	5
MiniMed Sure-T MMT-864A	21	Mycophenolate mofetil		Nifuran	
MiniMed Sure-T MMT-866A		Mydriacyl		Nilotinib	
MiniMed Sure-T MMT-874A		Mylan (12 hr release)		Nilstat	
MiniMed Sure-T MMT-876A		Mylan (24 hr release)		Alimentary	3
MiniMed Sure-T MMT-884A		Mylan Atenolol		Genito-Urinary	
MiniMed Sure-T MMT-886A		Mylan Clomiphen		Infection	
Minims Cyclopentolate		Mylan Indapamide		Nintedanib	
Minims Pilocarpine		Mylan Italy (24 hr release)		Nipent	
Minims Prednisolone		Mylan Midazolam		Nitrates	
Minirin		Myleran		Nitroderm TTS	
Minirin Melt		Myometrial and Vaginal Hormo		Nitrofurantoin	
Mino-tabs		Preparations		Nitrolingual Pump Spray	
Minocycline hydrochloride		Myozyme		Nivestim	
Minomycin		- N -		Nivolumab	
Minor Skin Infections		Nadolol	52	Nizoral	
Minoxidil		Nadolol BNM		Nodia	
Mirena		Naglazyme		Noflam 250	
Mirtazapine		Nalcrom		Noflam 500	
Misoprostol		Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	
Mitomycin C		Naltraccord		Drugs	114
Mitozantrone		Naltrexone hydrochloride		Nonacog gamma, [Recombinant	
Mitozantrone Ebewe		Naphazoline hydrochloride		Factor IX]	4
Mixtard 30		Naphcon Forte		Norethisterone	
MMR II		Naprosyn SR 1000		Genito-Urinary	7
Moclobemide		Naprosyn SR 750		Hormone	8
Modafinil		Naproxen		Norflex	
Modavigil		Narcaricin mite		Norfloxacin	
Moduretic		Nasal Preparations		Noriday 28	
Molaxole		Natalizumab		Norimin	
Moments		Natulan		Normacol Plus	
Mometasone furoate		Nausafix		Normison	
Monogen		Nausicalm		Norpress	
Montelukast		Navelbine		Nortriptyline hydrochloride	
Montelukast Mylan		Nefopam hydrochloride		Norvir	
Moroctocog alfa [Recombinant fa		Neisvac-C		Noumed	
VIII]					
	41	Neo-B12	34	Noumed Paracetamol	17
MODULE INDICATIONAL		Neo-B12 Neo-Mercazole		Noumed Paracetamol NovaSource Renal	
Morphine hydrochloride Morphine sulphate	129	Neo-Mercazole	86	NovaSource Renal	25
Morphine sulphate Motetis	129 129		86		250

NovoRapid	11	Ofev	234	Oxycodone Sandoz	130
NovoRapid FlexPen	11	Oil in water emulsion	68	Oxycodone Sandoz S29	130
NovoRapid Penfill	11	Olanzapine	139–140	OxyContin	130
NovoSeven RT	41	Olaparib	161	OxyNorm	130
Noxafil	100	Olbetam	56	Oxytocin	
Nozinan	139	Olbetam S29	56	Oxytocin BNM	78
Nozinan (Swiss)	139	Olopatadine	244	Oxytocin with ergometrine	
Nucala		Olopatadine Teva		maleate	78
Nuelin	236	Olsalazine		Ozurdex	240
Nuelin-SR	236	Omalizumab	202	-P-	
Nupentin	133	Omeprazole	9	Pacifen	122
Nutilis		Omeprazole actavis 10		Pacimol	12
Nutren Diabetes	253	Omeprazole actavis 20	9	Paclitaxel	16 ⁻
Nutrient Modules	250	Omeprazole actavis 40		Paclitaxel Actavis	
Nutrini Energy Multi Fibre	255	Omnitrope		Paclitaxel Ebewe	
Nutrini Energy RTH		Onbrez Breezhaler		Paediatric Seravit	3
Nutrini Low Energy Multi Fibre .		Oncaspar LYO		Palbociclib	
Nutrini Peptisorb		OncoTICE		Paliperidone	
Nutrini Peptisorb Energy		Ondansetron		Pamidronate disodium	
Nutrini RTH		Ondansetron ODT-DRLA		Pamisol	
Nutrison 800 Complete Multi		One-Alpha		Panadol	
Fibre	260	Onrex		Panadol Mini Caps	
Nutrison Concentrated		Opdivo		Pancreatic enzyme	
Nutrison Energy		Ora-Blend		Pantoprazole	
Nutrison Energy Multi Fibre		Ora-Blend SF		Panzop Relief	
Nutrison Multi Fibre		Ora-Plus		Panzytrat	
Nutrison Standard RTH		Ora-Sweet		Papaverine hydrochloride	
Nyefax Retard		Ora-Sweet SF		Para-amino salicylic acid	
		Orabase		Paracare	
Nystatin Alimentary	33	Oral and Enteral Feeds		Paracare Double Strength	
Genito-Urinary		Oratane		Paracetamol	
Infection		Ordine		Paracetamol + Codeine	12
NZB Low Gluten Bread Mix					100
- 0 -	203	Orgran		(Relieve) Paracetamol Pharmacare	10
	60	Ornidazole			
O/W Fatty Emulsion Cream		Orphenadrine citrate		Paracetamol with codeine	
Obstatria Proporations		Ortho-tolidine		Paraffin	
Obstetric Preparations		Oruvail SR		Paraffin liquid with wool fat	
Ocrelizumab		Osmolite RTH		Parasidose	
Ocrevus		Other Endocrine Agents		Parasiticidal Preparations	
Octocog alfa [Recombinant fact		Other Oestrogen Preparation	ns85	Parlodel	
VIII] (Advate)		Other Progestogen	0.5	Parnate	
Octocog alfa [Recombinant fact		Preparations		Parnate S29	
VIII] (Kogenate FS)		Other Skin Preparations	73	Paromomycin	
Octreotide		Ovestin	70	Paroxetine	
Octreotide (Sun)	173	Genito-Urinary		Paser	
Octreotide Depot Teva		Hormone		Paxam	
Octreotide GH		Ox-Pam		Paxtine	
Octreotide long-acting		Oxaliplatin		Pazopanib	
Octreotide MaxRx		Oxaliplatin Accord		Peak flow meter	
Oestradiol		Oxaliplatin Actavis 100		Pedialyte - Bubblegum	
Oestradiol valerate		Oxaliplatin Ebewe		Pediasure	
Oestradiol with norethisterone	85	Oxazepam		Pediasure RTH	
Oestriol		Oxis Turbuhaler	232	Pegaspargase	162
Genito-Urinary	78	Oxpentifylline		Pegasys	110
Hormone	85	Oxybutynin		Pegfilgrastim	46
Oestrogens	84	Oxycodone hydrochloride	130	Pegylated interferon alfa-2a	110

Pembrolizumab	225	PKU Anamix Junior LQ	264	Probenecid-AFT	12
Pemetrexed	157	PKU Anamix Junior Orange	264	Procaine penicillin	9
Penicillamine	115	PKU Anamix Junior Vanilla	264	Procarbazine hydrochloride	16
Penicillin G	95	PKU Lophlex LQ 10	264	Prochlorperazine	
PenMix 30	11	PKU Lophlex LQ 20	264	Proctofoam	
PenMix 40	11	PKU Lophlex Powder	264	Proctosedyl	
PenMix 50	11	PKU Lophlex Sensation 20	264	Procyclidine hydrochloride	12
Pentasa		Plaquenil	115	Procytox	
Pentostatin [Deoxycoformycin]	162	Plendil ER	53	Progesterone	
Pentoxifylline [Oxpentifylline]	59	Pneumococcal (PCV10) conjuga	ate	Proglicem	10
Peptamen Junior		vaccine	278	Proglycem	
Peptisorb	256	Pneumococcal (PCV13) conjuga	ate	Progynova	8
Perhexiline maleate	54	vaccine		Prolia	
Pericyazine	139	Pneumococcal (PPV23)		Promethazine hydrochloride	23
Perindopril		polysaccharide vaccine	280	Propafenone hydrochloride	5
Perjeta		Pneumovax 23	280	Propamidine isethionate	23
Permethrin	70	Podophyllotoxin	73	Propranolol	
Perrigo	72	Polaramine	231	Propylene glycol	24
Pertuzumab		Poliomyelitis vaccine		Propylthiouracil	
Peteha	103	Poloxamer	26	Prostacur	17
Pethidine hydrochloride	130	Poly-Gel	243	Protaphane	
Pevaryl		Poly-Tears	243	Protaphane Penfill	10
Pexsig	54	Poly-Visc		Protifar	
Pfizer Exemestane		Polycal		Protionamide	
Pharmacy Health Sorbolene with		Ponstan		Provera	8
Glycerin	67	Posaconazole	100	Provera HD	8
Pheburane		Postinor-1	78	PSM Citalopram	13
Phenasen	158	Potassium chloride	46-47	Psoriasis and Eczema	
Phenobarbitone	134	Potassium citrate	80	Preparations	70
Phenobarbitone sodium		Potassium iodate		PTU	
Extemporaneous	248	Povidone iodine	68	Pulmicort Turbuhaler	23
Nervous		Pradaxa		Pulmozyme	23
Phenothrin	70	Pramipexole hydrochloride	123	Puri-nethol	150
Phenoxybenzamine		Pravastatin		Puria	
hydrochloride	48	Pravastatin Mylan	57	Pyrazinamide	10
Phenoxymethylpenicillin (Penicillin		Praziquantel	92	Pyridostigmine bromide	
V)		Prazosin		Pyridoxine hydrochloride	
Phenytoin sodium13		Pred Forte	241	Pyridoxine multichem	
Phillips Milk of Magnesia		Prednisolone		Pyrimethamine	
Phlexy 10		Prednisolone acetate	241	Pytazen SR	4
Phosphate Phebra		Prednisolone sodium		- Q -	
Phosphorus		Prednisolone sodium		Quetapel	139
Phytomenadione		phosphate	241	Quetiapine	
Pilocarpine hydrochloride		Prednisolone-AFT		Quick-Set MMT-392	
Pimafucort	67	Prednisone		Quick-Set MMT-393	2
Pimecrolimus	71	Pregabalin	134	Quinapril	
Pindolol	52	Pregabalin Pfizer		Quinapril with	
Pine tar with trolamine laurilsulfate)	Pregnancy Tests - hCG Urine	79	hydrochlorothiazide	49
and fluorescein		Premarin		Qvar	23
Pinetarsol		Prevenar 13		- R -	
Pioglitazone		Priadel		RA-Morph	12
Pirfenidone		Primaquine		Raloxifene hydrochloride	
Pizotifen		Primidone	135	Raltegravir potassium	110
PKU Anamix Infant		Primolut N		Ramipex	
PKU Anamix Junior		Priorix		Ranbaxy-Cefaclor	
PKU Anamix Junior Chocolate		Probenecid		Ranbaxy-Cefaclor S29	9

Rapamune	227	Ropinirole hydrochloride	123	Sildenafil	60
Rasagiline	123	Rotarix	280	Silhouette MMT-373	24
Reandron 1000		Rotavirus oral vaccine	280	Siltuximab	218
Recombinant factor IX	39, 42	Roxane	6	Simvastatin	5
Recombinant factor VIIa	41	Roxane-Propranolol	53	Simvastatin Mylan	5
Recombinant factor VIII	41–42	Roxithromycin	94	Sinemet	
Rectogesic	8	Rubifen		Sinemet CR	
Redipred	83	Rubifen SR	145	Sirolimus	22
Relieve		Rugby Capsaicin Topical		Siterone	8
Relistor	26	Cream	126	Slow-Lopresor	52
Remicade	194	Rulide D	94	Smith BioMed Rapid Pregnancy	
Renilon 7.5	256	Rurioctocog alfa pegol [Reco	mbinant	Test	79
Resonium-A	47	factor VIII]		Sodibic	4
Resource Beneprotein	252	Ruxolitinib		Sodium acid phosphate	2
Respigen		Rythmodan	50	Sodium alginate	(
Respiratory Devices	237	Rytmonorm	51	Sodium benzoate	3
Respiratory Stimulants		· - S -		Sodium bicarbonate	
Retinol palmitate		Sabril	135	Blood	.46-47
ReTrieve		Sacubitril with valsartan	49	Extemporaneous	
Retrovir	109	Sagent	165	Sodium calcium edetate	
Revlimid	160	SalAir		Sodium chloride	
Revolade	39	Salazopyrin		Blood	4
Rexacrom	241	Salazopyrin EN		Respiratory	
Riboflavin	30	Salbutamol		Sodium citrate with sodium lauryl	
Ribomustin	152	Salbutamol with ipratropium		sulphoacetate	2
Ricit	79	bromide	233	Sodium citro-tartrate	
Rifabutin		Salicylic acid		Sodium cromoglicate	
Rifadin		Salmeterol		Alimentary	8
Rifampicin	103	Sandomigran	136	Sensory	
Rifaximin		Sandostatin LAR		Sodium fluoride	
Rifinah		Sanofi Primaquine		Sodium Fusidate [fusidic acid]	
Rilutek	124	Sapropterin dihydrochloride	31	Dermatological	64
Riluzole	124	Scalp Preparations		Infection	
RINVOQ		Scopoderm TTS		Sensory	
Riodine	68	Sebizole		Sodium hyaluronate [Hyaluronic	
Risedronate Sandoz	117	Secukinumab		acid]	243
Risedronate sodium	117	Sedatives and Hypnotics	143	Sodium phenylbutyrate	
Risperdal Consta		Seebri Breezhaler		Sodium picosulfate	
Risperidone		Selegiline hydrochloride		Sodium polystyrene sulphonate	
Risperidone (Teva)		Senna		Sodium tetradecyl sulphate	
Risperon		Senokot	<mark>27</mark>	Sodium valproate	
Ritalin		Sensipar	81	Sofradex	
Ritalin LA	146	SensoCard	15	Soframycin	239
Ritonavir	110	Serenace	138	Solgar	
Rituximab (Mabthera)	204	Seretide	233	Solifenacin Mylan	
Rituximab (Riximyo)	206	Seretide Accuhaler	233	Solifenacin succinate	
Rivaroxaban		Serevent		Solu-Cortef	82
Rivastigmine	147	Serevent Accuhaler	232	Solu-Medrol	82
Rivastigmine Patch BNM 10		Sertraline		Solu-Medrol-Act-O-Vial	82
Rivastigmine Patch BNM 5		Setrona		Somatropin (Omnitrope)	8
Rivotril		Sevredol		Sotalol	5
Riximyo		Sex Hormones Non	-	Spacer device	
RIXUBIS		Contraceptive	83	Span-K	
Rizamelt		Shield XL	75	Spiolto Respimat	
Rizatriptan		shingles vaccine		Spiractin	
Ropin		SII-Onco-BCG		Spiriva	
•				•	

Spiriva Respimat	234	Tamoxifen citrate	174	Tobramycin	
Spironolactone	55	Tamoxifen Sandoz	174	Infection	99
Sporanox	100	Tamsulosin hydrochloride	79	Sensory	240
Sprycel	165	Tamsulosin-Rex	79	Tobramycin BNM	99
Staphlex		Tandem Cartridge		Tobramycin Mylan	99
Stemetil	138	Tandem t:slim X2 with Basal-IQ		Tobrex	
SteroClear		Tap water		Tocilizumab	
Stesolid		Tarceva		Tofranil	
Stimulants/ADHD Treatments	144	Tasigna		Tolcapone	
Stiripentol	135	Tasmar	124	Topamax	135
Stocrin		Taurine		Topical Products for Joint and	
Stomahesive	33	Tecfidera	142	Muscular Pain	115
Strattera		Tegretol		Topiramate	
Strides Shasun		Tegretol CR		Topiramate Actavis	
Stromectol		Telfast		Total parenteral nutrition (TPN)	
Sucralfate		Teligent		TPN	
Sulfadiazine Silver		Temaccord		Tramadol hydrochloride	
Sulfadiazine sodium		Temazepam		Tramal SR 100	
Sulfasalazine		Temozolomide		Tramal SR 150	
Sulindac		Tenofovir disoproxil		Tramal SR 200	
Sulindac Mylan		Tenofovir Disoproxil Teva		Trandate	
Sulphur		Tenoxicam		Tranexamic acid	
Sulprix		Tensipine MR10		Tranylcypromine sulphate	
Sumagran		Tepadina		Trastuzumab	
Sumatriptan		Terbinafine		Trastuzumab emtansine	
Sunitinib				Travatan	
		Terbutaline sulphate			
Sunscreens		Teriflunomide		Travoprost	
Sunscreens, proprietary		Teriparatide		•	
Sure-T MMT-863		Testosterone		Treatments for Dementia	147
Sure-T MMT-873		Testosterone cipionate		Treatments for Substance	4.40
Sustagen Diabetic	253	Testosterone esters		Dependence	
Sustagen Hospital Formula	000	Testosterone undecanoate		Trental 400	59
Active		Tetrabenazine		Tretinoin	
Sustanon Ampoules		Tetrabromophenol		Dermatological	
Sutent		Tetracosactrin		Oncology	
Sylvant		Tetracycline		Trexate	157
Symbicort Turbuhaler 100/6		Thalidomide		Triamcinolone acetonide	
Symbicort Turbuhaler 200/6		Thalomid		Alimentary	
Symbicort Turbuhaler 400/12		Theophylline		Dermatological	
Symmetrel		Thiamine hydrochloride	34	Hormone	83
Sympathomimetics		THIO-TEPA		Triamcinolone acetonide with	
Synacthen		Thioguanine		gramicidin, neomycin and nys	
Synacthen Depot	83	Thiotepa	154	Dermatological	
Synacthene Retard		Thymol glycerin		Sensory	
Synflorix	278	Thyroid and Antithyroid Agents		Triazolam	
Synthroid	86	Ticagrelor	43	Trimethoprim	99
Syntometrine	78	Tilcotil	115	Trimethoprim with	
Syrup (pharmaceutical grade)	249	Tillomed	154	sulphamethoxazole	
Systane Unit Dose	243	Timolol	241	[Co-trimoxazole]	99
-T-		Timoptol XE	241	Trisequens	
Tacrolimus		Tiotropium bromide	234	Trisul	99
Dermatological	72	Tiotropium bromide with		Trophic Hormones	86
Oncology		olodaterol	234	Tropicamide	
Tacrolimus Sandoz		Tivicay		Trulicity	
Taliglucerase alfa		TMP		Trusopt	
Tambacar	50			,	_

TruSteel	22	Versacloz	138	Zaroxolyn	55
Tuberculin PPD [Mantoux] test	281	Vesanoid	163	Zavedos	160
Tubersol	281	Vexazone	12	Zeffix	104
Two Cal HN	262	Vfend	101	Zematop	72
Two Cal HN RTH	262	Viaderm KC	67	Zetlam	104
Tykerb	167	Vidaza	154	Ziagen	109
Tysabri		Vigabatrin	135	Zidovudine [AZT]	109
- U -		Vigisom	143	Zidovudine [AZT] with	
Ultibro Breezhaler	234	Vildagliptin	12	lamivudine	109
Ultraproct	8	Vildagliptin with metformin		Zimybe	57
Umeclidinium	234	hydrochloride	12	Zinc and castor oil	67
Umeclidinium with vilanterol		Vimpat		Zinc sulphate	37
Univent	233, 237	Vinblastine sulphate	164	Zincaps	37
Upadacitinib		Vincristine sulphate		Zinnat	
Ural	80	Vinorelbine	165	Ziprasidone	
Urea	68	Vinorelbine Ebewe	165	Zista	231
Urex Forte	55	Viramune Suspension	108	Zithromax	93
Urinary Agents	79	ViruPOS	239	Zoledronic acid	
Urinary Tract Infections		Vit.D3	34	Hormone	81
Urinorm		Vita-B12	34	Musculoskeletal	
Uromitexan	161	VitA-POS	244	Zoledronic acid Mylan	81
Ursodeoxycholic acid	25	Vitabdeck	35	Zopiclone	
Ursosan		Vital	256	Zopiclone Actavis	144
Utrogestan	85	Vitamin B complex	34	Zostavax	281
- V -		Vitamin B6 25		Zostrix	
Vaccinations	270	Vitamins	34–35	Zostrix HP	126
Vaclovir	104	Vivonex TEN	256	Zuclopenthixol decanoate	141
Valaciclovir	104	Voltaren	114	Zuclopenthixol hydrochloride	139
Valganciclovir	104	Voltaren D	114	Zusdone	
Valganciclovir Mylan	104	Voltaren Ophtha	240	Zyban	149
Vancomycin		Voltaren SR	114	Zypine	139
Vannair		Volumatic		Zypine ODT	139
Varenicline Pfizer	150	Voriconazole	101	Zyprexa Relprevv	
Varenicline tartrate	150	Vosol		Zytiga	
Varicella vaccine [Chickenpox		Votrient		, ,	
vaccine]	281	Vttack	101		
Varicella zoster virus (Oka strair		- W -			
attenuated vaccine [shingles	,	Warfarin sodium	45		
vaccine]	281	Wart Preparations			
Varivax	281	Wasp venom allergy treatme			
Vasodilators	58	Water			
Vasopressin Agonists	91	Blood	46		
Vasorex		Extemporaneous	249		
Vedafil	60	Wool fat with mineral oil			
Veletri	61	- X -			
Venclexta	164	Xarelto	45		
Venetoclax	164	Xifaxan			
Venlafaxine		XMET Maxamum			
Venomil		Xolair			
VENOX		XP Maxamum			
Ventavis		Xylocaine			
Ventolin		Xylocaine 2% Jelly	125		
Vepesid		Xyntha	41		
Verapamil hydrochloride	54	-Z-	*****		
Vergo 16		Zapril	48		
Vermox		Zarontin			