| April 2022 Volume 29 Number 1 | Se |
|---|----|
| Editors: Kaye Wilson & Sophie Molloy | |
| email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington | Se |
| Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays) | |
| Circulation | |
| Accessible in an electronic format at no cost from the Pharmac website www.pharmac.govt.nz/schedule. | |
| You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at | |
| pharmac.govt.nz/subscribe. | |
| Production | |
| Typeset automatically from XML and T _E X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule | |
| Programmers | |
| Anrik Drenth & John Geering | |
| email: texschedule@pharmac.govt.nz | |
| ©Pharmaceutical Management Agency ISSN 1179-3686 | Se |
| This work is licensed under the Creative Commons Attribution 4.0 International licence. | Se |
| In essence, you are free to copy, distribute and adapt it, as long as you attribute the work | |
| to Pharmac and abide by the other licence | Se |
| terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. | - |
| Attribution to Pharmac should be in written | _ |
| form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from. | |
| | |

| Section A | General Rules | 5 |
|-----------|--------------------------------------|-----|
| Section B | Alimentary Tract & Metabolism | 6 |
| | Blood & Blood Forming Organs | 37 |
| | Cardiovascular System | 47 |
| | Dermatologicals | 61 |
| | Genito Urinary System | 71 |
| | Hormone Preparations – Systemic | 78 |
| | Infections – Agents For Systemic Use | 89 |
| | Musculoskeletal System | 111 |
| | Nervous System | 119 |
| | Oncology Agents & Immunosuppressants | 146 |
| | Respiratory System & Allergies | 228 |
| | Sensory Organs | 237 |
| | Various | 242 |
| Section C | Extemporaneous Compounds (ECPs) | 244 |
| Section D | Special Foods | 247 |
| Section I | National Immunisation Schedule | 268 |
| | Index | 278 |
| | | |

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.

The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

| gram g | |
|-----------------------|--|
| kilogram kg | |
| international unit iu | |

Abbreviations

| Capsule Cream Device Dispersible Effervescent Emulsion | Amp Cap Crm Dev Disp Eff Emul EC |
|---|---|
| Enteric Coated | EC |
| | |

| microgram me | cg |
|--------------|----|
| milligramn | ng |
| millilitreı | ml |

| millimole | mmol |
|-----------|------|
| unit | u |

| Gelatinous | Gel | SolutionSoln |
|-------------|------|-----------------------|
| Granules | Gran | SuppositorySupp |
| Infusion | Inf | TabletTab |
| Injection | Inj | Tincture Tinc |
| Liquid | Liq | Trans Dermal Delivery |
| Long Acting | LA | SystemTDDS |
| Ointment | Oint | - |
| Sachet | Sach | |

Read the General Rules : https://www.pharmac.govt.nz/section-a.

SECTION B: ALIMENTARY TRACT AND METABOLISM

| | Subsidy | Full | 5 |
|---|------------------------------|--------------------|---|
| | (Manufacturer's Price) \$ | Subsidise Per 🖌 | d Generic Manufacturer |
| Antacids and Antiflatulents | | | |
| Antacids and Reflux Barrier Agents | | | |
| ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet | | 30 🗸 | Gaviscon Infant |
| SODIUM ALGINATE * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour | 1.80 (8.60) | 60 | Gaviscon Double Strength |
| * Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml | | 500 ml | Acidex |
| Phosphate Binding Agents | | | |
| ALUMINIUM HYDROXIDE * Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement Only when prescribed for patients unable to swallow cal inappropriate and the prescription is endorsed according | | 500 ml 🗸 | ⁷ Alu-Tab ⁷ Roxane cium carbonate tablets are |
| Antidiarrhoeals | | | |
| Agents Which Reduce Motility | | | |
| LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg * Cap 2 mg | | | ´ Nodia ´ <u>Diamide Relief</u> |
| Rectal and Colonic Anti-inflammatories | | | |
| BUDESONIDE Cap 3 mg – Special Authority see SA1886 below – Retail pharmacy | | | Fentocort CIR as for applications meeting |
| Both: 1 Mild to moderate ileal, ileocaecal or proximal Crohn's dise 2 Any of the following: 2.1 Diabetes; or 2.2 Cushingoid habitus; or 2.3 Osteoporosis where there is significant risk of frac | | | |
| | | | continued. |
| | | | |

| Subsidy | Fu | lly Brand or | |
|------------------------|----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth).

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

| Rectal foam 10%, CFC-Free (14 applications) | 15 g OP 21.1 g OP | ✓ Cortifoam ^{S29} ✓ Colifoam |
|--|----------------------|--|
| HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE | | |
| Topical aerosol foam, 1% with pramoxine hydrochloride 1% | 10 g OP | Proctofoam S29 |
| MESALAZINE | | |
| Tab 400 mg49.50 | 100 | Asacol |
| Tab long-acting 500 mg56.10 | 100 | Pentasa |
| Tab 800 mg | 90 | Asacol |
| Modified release granules, 1 g | 100 OP | Pentasa |
| Enema 1 g per 100 ml41.30 | 7 | Pentasa |
| Suppos 500 mg | 20 | Asacol |
| Suppos 1 g | 28 | Pentasa |
| OLSALAZINE | | |
| Tab 500 mg | 100 | Dipentum |
| Cap 250 mg | 100 | Dipentum |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully | Brand or |
|--|------------------------------|------------|-------------|-------------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| REDNISOLONE SODIUM | | | | |
| Rectal foam 20 mg per dose (14 applications) | 74.10 | 1 OP | 1 | Essential Prednisolone S29 |
| ODIUM CROMOGLICATE | | | | |
| Cap 100 mg | 92.91 | 100 | ~ | Nalcrom |
| | 11.00 | 400 | | 0-1 |
| Tab 500 mg Tab EC 500 mg | | 100 100 | | Salazopyrin Salazopyrin EN |
| Local preparations for Anal and Rectal Disorder | | | | <u> </u> |
| Antihaemorrhoidal Preparations | | | | |
| LUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIV | ALATE AND CINCH | OCAII | NE | |
| Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g | | 30 g O | Р 🗸 | Ultraproct |
| Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and | 7.00 | 10 | | 1114 warmen at |
| cinchocaine hydrochloride 1 mg | | 12 | • | Ultraproct |
| YDROCORTISONE WITH CINCHOCAINE Oint 5 mg with cinchocaine hydrochloride 5 mg per g | | 30 g O | Р 🗸 | Proctosedyl |
| Suppos 5 mg with cinchocaine hydrochloride 5 mg per g | | 12 | | Proctosedyl |
| Management of Anal Fissures | | | | |
| LYCERYL TRINITRATE – Special Authority see SA1329 below ← Oint 0.2% | | 30 g O | p 🗸 | Rectogesic |
| • SA1329 Special Authority for Subsidy | | ,o y o | | neologesio |
| itial application from any relevant practitioner. Approvals valid hronic anal fissure that has persisted for longer than three week | | ewal u | nless notif | ied where the patient has |
| Antispasmodics and Other Agents Altering Gut | Motility | | | |
| LYCOPYRRONIUM BROMIDE | | | | |
| Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on | а | | | |
| PSO | 65.45 | 10 | ~ | Max Health |
| YOSCINE BUTYLBROMIDE | 0.05 | | | _ |
| ✓ Tab 10 mg ✓ Inj 20 mg, 1 ml – Up to 5 inj available on a PSO | | 100 5 | | Buscopan Buscopan |
| IEBEVERINE HYDROCHLORIDE | | 5 | · | Duscopan |
| | 9.20 | 90 | 1 | Colofac |
| Tab 135 mg | | | | |
| - | | | | |
| Antiulcerants | | | | |
| - | | | | |
| Antiulcerants | 41.50 | 120 | | Cytotec |

| | Subsidy | | Fully | Brand or |
|--|--|-------------|--------------|--------------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| Helicobacter Pylori Eradication | | | | |
| CLARITHROMYCIN Tab 500 mg - Subsidy by endorsement a) Maximum of 28 tab per prescription b) Subsidised only if prescribed for helicobacter py Note: the prescription is considered endorsed i inhibitor and either amoxicillin or metronidazole | /lori eradication and prescri f clarithromycin is prescribe | | is endorsed | |
| H2 Antagonists | | | | |
| AMOTIDINE – Only on a prescription | | | | |
| K Tab 20 mg | 4.91 | 100 | ✓ F | amotidine Hovid S29 |
| ₭ Tab 40 mg | 8.48 | 100 | ✓ F | amotidine Hovid S29 |
| Inj 10 mg per ml, 4 ml – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients | | 10 of pa | | ylan S29 |
| Proton Pump Inhibitors | | | | |
| ANSOPRAZOLE | | | | |
| € Cap 15 mg | | 100 | | anzol Relief |
| Cap 30 mg | 5.26 | 100 | ✓ <u>L</u> | anzol Relief |
| MEPRAZOLE | | | | |
| For omeprazole suspension refer Standard Formulae, p Cap 10 mg | | 90 | ✓ <u>o</u> | meprazole actavis 10 |
| € Cap 20 mg | 1.86 | 90 | ✓ <u>0</u> | meprazole actavis 20 |
| € Cap 40 mg | 3.11 | 90 | ✓ <u>0</u> | meprazole actavis 40 |
| Powder – Only in combination Only in extemporaneously compounded omeprazo | | 5 g | 🗸 M | lidwest |
| Inj 40 mg ampoule with diluent | | 5 | ✓ <u>D</u> | <u>r Reddy's</u> Omeprazole |
| ANTOPRAZOLE | 0.00 | 100 | . (D | onton Doliof |
| ✓ Tab EC 20 mg ✓ Tab EC 40 mg | | 100 100 | | anzop Relief anzop Relief |
| Site Protective Agents | | 100 | • 1 | |
| - | | | | |
| OLLOIDAL BISMUTH SUBCITRATE | 14 54 | 50 | 1.0 | a atria dan al ana |
| Tab 120 mg | 14.51 | 50 | ✔ G | astrodenol S29 |
| UCRALFATE Tab 1 g | 35.50 (48.28) | 120 | С | arafate |
| Bile and Liver Therapy | | | | |
| RIFAXIMIN – Special Authority see SA1461 on the next pa Tab 550 mg | | 56 | ✓ <u>x</u> | ifaxan |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Pri \$ | ce) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|--|-----------------------|---|--|
| ➤SA1461 Special Authority for Subsidy nitial application only from a gastroenterologist, hepatologist nepatologist. Approvals valid for 6 months where the patient has olerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Practiti nepatologist. Approvals valid without further renewal unless no benefiting from treatment. | as hepatic encephal ioner on the recomm | opathy d | espite an a n of a gastro | dequate trial of maximur penterologist or |
| Diabetes | | | | |
| Hyperglycaemic Agents | | | | |
| DIAZOXIDE – Special Authority see SA1320 below – Retail ph Cap 25 mg Cap 100 mg Oral liq 50 mg per ml ⇒SA1320 Special Authority for Subsidy | | 100 100 30 ml C | ✓I DP ✓I | Proglicem S29 Proglicem S29 Proglycem S29 |
| nitial application from any relevant practitioner. Approvals vanypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid without appropriate and the patient is benefiting from treatment. GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO | ut further renewal ur | | fied where | |
| Insulin - Short-acting Preparations | | | | |
| NSULIN NEUTRAL ▲ Inj human 100 u per ml | | 10 ml C 5 | I I | Actrapid Humulin R Actrapid Penfill |
| Inculing Intermediate entire Dreservations | | | • 1 | Humulin R |
| Insulin - Intermediate-acting Preparations | | | | |
| NSULIN ASPART WITH INSULIN ASPART PROTAMINE ▲ Inj 100 iu per ml, 3 ml prefilled pen NSULIN ISOPHANE | 52.15 | 5 | √ I | NovoMix 30 FlexPen |
| Inj human 100 u per ml | 17.68 | 10 ml C | | Humulin NPH Protaphane |
| ▲ Inj human 100 u per ml, 3 ml | 29.86 | 5 | 🗸 I | Humulin NPH Protaphane Penfill |
| NSULIN ISOPHANE WITH INSULIN NEUTRAL | | | | |
| Inj human with neutral insulin 100 u per ml | 25.26 | 10 ml C | | Humulin 30/70 Mixtard 30 |
| Inj human with neutral insulin 100 u per ml, 3 ml | 42.66 | 5 | ✓ ✓ ✓ | Humulin 30/70 PenMix 30 PenMix 40 PenMix 50 |

| | Subsidy | | Fully | Brand or |
|--|-----------------------|--|--------------------------------------|---|
| | (Manufacturer's Price | e) 5 | Subsidised | Generic |
| | \$ | Per | 1 | Manufacturer |
| | | | | |
| NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE | | | | |
| Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, | , | | | |
| 3 ml | | 5 | ✓ | Humalog Mix 25 |
| | | - | | ······································ |
| | | | | |
| 3 ml | | 5 | ~ | Humalog Mix 50 |
| | | | | |
| Insulin - Long-acting Preparations | | | | |
| inotin _eng toting i toparationo | | | | |
| NSULIN GLARGINE | | | | |
| | 60.00 | 4 | | Lantua |
| Inj 100 u per ml, 10 ml | | 1 | | Lantus |
| Inj 100 u per ml, 3 ml | | 5 | ~ | Lantus |
| Inj 100 u per ml, 3 ml disposable pen | 94.50 | 5 | ✓ | Lantus SoloStar |
| | | | | |
| Insulin - Rapid Acting Preparations | | | | |
| insuint - napiu Acung Freparations | | | | |
| NSULIN ASPART | | | | |
| | ~~~~ | | | |
| Inj 100 u per ml, 10 ml | | 1 | ~ | NovoRapid |
| Inj 100 u per ml, 3 ml | 51.19 | 5 | ✓ | NovoRapid Penfill |
| Inj 100 u per ml, 3 ml syringe | | 5 | 1 | NovoRapid FlexPen |
| | | Ŭ | | novonapia nozi on |
| VSULIN GLULISINE | | | | |
| Inj 100 u per ml, 10 ml | 27.03 | 1 | ✓ | Apidra |
| Inj 100 u per ml, 3 ml | | 5 | 1 | Apidra |
| Inj 100 u per ml, 3 ml disposable pen | | 5 | | • |
| | | 5 | • | Apidra SoloStar |
| NSULIN LISPRO | | | | |
| Inj 100 u per ml, 10 ml | 34 92 | 10 ml OF | - J | Humalog |
| | | | - | • |
| Inj 100 u per ml, 3 ml | | 5 | v | Humalog |
| | | | | |
| Alpha Glucosidase Inhibitors | | | | |
| | | | | |
| ACARBOSE | | | | |
| 🖌 Tab 50 mg | 8.95 | 90 | ✓ | Accarb |
| 🖌 Tab 100 mg | | 90 | 1 | Accarb |
| | | | | |
| Oral Humanluagamia Ananta | | | | |
| | | | | |
| orar riypogiycaeniic Agents | | | | |
| | | | | |
| GLIBENCLAMIDE | 7.50 | 400 | | Descil |
| SLIBENCLAMIDE | 7.50 | 100 | 1 | <u>Daonil</u> |
| GLIBENCLAMIDE K Tab 5 mg | 7.50 | 100 | v | <u>Daonil</u> |
| aLIBENCLAMIDE ∉ Tab 5 mg SLICLAZIDE | | | | |
| GLIBENCLAMIDE ≰ Tab 5 mg GLICLAZIDE | | 100 500 | | <u>Daonil</u> <u>Glizide</u> |
| GLIBENCLAMIDE ≰ Tab 5 mg SLICLAZIDE ≰ Tab 80 mg | | | | |
| GLIBENCLAMIDE ← Tab 5 mg SLICLAZIDE ← Tab 80 mg SLIPIZIDE | 15.18 | 500 | 1 | Glizide |
| GLIBENCLAMIDE | 15.18 | | 1 | |
| GLIBENCLAMIDE | 15.18 4.58 | 500 | 1 | Glizide |
| GLIBENCLAMIDE ← Tab 5 mg SLICLAZIDE ← Tab 80 mg SLIPIZIDE ← Tab 5 mg IETFORMIN HYDROCHLORIDE | 15.18 4.58 | 500 100 | v v | <u>Glizide</u> <u>Minidiab</u> |
| ALIBENCLAMIDE ← Tab 5 mg ALICLAZIDE ← Tab 80 mg ALIPIZIDE ← Tab 5 mg IETFORMIN HYDROCHLORIDE ← Tab immediate-release 500 mg | | 500 100 1,000 | 1 1 1 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan |
| GLIBENCLAMIDE | | 500 100 | 1 1 1 | <u>Glizide</u> <u>Minidiab</u> |
| GLIBENCLAMIDE | | 500 100 1,000 | 1 1 1 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan |
| GLIBENCLAMIDE | | 500 100 1,000 500 | \$ \$ \$ | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan |
| GLIBENCLAMIDE | | 500 100 1,000 500 90 | • • • • | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan <u>Vexazone</u> |
| GLIBENCLAMIDE | | 500 100 1,000 500 90 90 | 5 5 5 5 5 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan <u>Vexazone</u> <u>Vexazone</u> |
| ₭ Tab 5 mg ĜLICLAZIDE ₭ Tab 80 mg ĜLIPIZIDE ₭ Tab 5 mg №TFFORMIN HYDROCHLORIDE ₭ Tab immediate-release 500 mg № Tab immediate-release 850 mg № Tab immediate-release 850 mg № Tab immediate-release 850 mg № Tab 15 mg ₭ Tab 15 mg ₭ Tab 30 mg | | 500 100 1,000 500 90 | 5 5 5 5 5 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan <u>Vexazone</u> |
| GLIBENCLAMIDE | | 500 100 1,000 500 90 90 | 5 5 5 5 5 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan <u>Vexazone</u> <u>Vexazone</u> |
| GLIBENCLAMIDE * Tab 5 mg GLICLAZIDE * Tab 80 mg GLIPIZIDE * Tab 5 mg METFORMIN HYDROCHLORIDE * Tab immediate-release 500 mg * Tab immediate-release 850 mg PIOGLITAZONE * Tab 15 mg * Tab 30 mg | | 500 100 1,000 500 90 90 | 5 5 5 5 5 5 5 5 | <u>Glizide</u> <u>Minidiab</u> Metformin Mylan Metformin Mylan <u>Vexazone</u> <u>Vexazone</u> |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|-------------------------------------|
| VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE Tab 50 mg with 1,000 mg metformin hydrochloride Tab 50 mg with 850 mg metformin hydrochloride | | 60 60 | - | alvumet alvumet |

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 above - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

| * | Inj 1.5mg per 0.5 ml prefille | pen 115.23 | 4 | Trulicity |
|---|-------------------------------|------------|---|-------------------------------|
|---|-------------------------------|------------|---|-------------------------------|

SGLT2 Inhibitors

➡SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
- 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
- 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

| * | Tab 10 mg | 58 56 | 30 | Jardiance |
|---|-----------|-------|----|-------------------------------|
| | | | | ✓ Jardiance |
| * | Tab 25 mg | | 30 | Jardiance |

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see SA2068 on the previous page – Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

| * | Tab 5 mg with 1,000 mg metformin hydrochloride | 60 | Jardiamet |
|---|---|--------|-------------------------------|
| * | Tab 5 mg with 500 mg metformin hydrochloride | 60 | Jardiamet |
| * | Tab 12.5 mg with 1,000 mg metformin hydrochloride | 60 | Jardiamet |
| * | Tab 12.5 mg with 500 mg metformin hydrochloride | 60 | Jardiamet |

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.
 - The prescription must be endorsed accordingly.

.... 15.50 10 strip OP

KetoSens

| | Subsidy (Manufacturer's Price) \$ | | ully Brand or sed Generic Manufacturer |
|---|--|---|---|
| Dual Blood Glucose and Blood Ketone Testing | | | |
| DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A dual blood glucose and blood ketone diagnostic test n type 1 diabetes; or permanent neonatal diabetes; or undergone a pancreatectomy; or cystic fibrosis-related diabetes; or metabolic disease or epilepsy under the care of a The prescription must be endorsed accordingly. Only 1 the avoidance of doubt patients who have previously rec funded CareSens meter. | neter is subsidised for paediatrician, neurolog meter per patient will seived a funded meter | a patient who gist or metabo be subsidised |) has: blic specialist. d (no repeat prescriptions). Fo |
| diagnostic test strips | | 1 OP | ✓ CareSens Dual |
| Blood Glucose Testing | | | |
| BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by a) Maximum of 1 pack per prescription b) Up to 1 pack available on a PSO c) A diagnostic blood glucose test meter is subsidised for a 1) is receiving insulin or sulphonylurea therapy; or 2) is pregnant with diabetes; or 3) is on home TPN at risk of hypoglycaemia or hyperi 4) has a genetic or an acquired disorder of glucose h syndrome. The prescription must be endorsed accordingly. Only or prescriptions). Patients already using the CareSens N F meter, unless they have: 1) type 1 diabetes; or 2) permanent neonatal diabetes; or 3) undergone a pancreatectomy; or 4) cystic fibrosis-related diabetes. For the avoidance of doubt patients who have previously funded CareSens meter. Meter with 50 lancets, a lancing device and 10 diagnostic te strips Note: Only 1 meter available per PSO | a patient who: glycaemia; or omeostasis, excluding ne CareSens meter pr POP meter and CareS y received a funded m | er patient will ens N meter a eter, other tha 1 OP | be subsidised (no repeat are not eligible for a new |

| | Subsidy (Manufacturer's P \$ | rice) Sub Per | Fully sidised | Brand or Generic Manufacturer |
|--|------------------------------------|------------------|------------------|-------------------------------------|
| BLOOD GLUCOSE DIAGNOSTIC TEST STRIP – Up to 50 te | est available on a PS | 60 | | |
| The number of test strips available on a prescription is res | | | | |
| 1) Prescribed for a patient on insulin or a sulphonylure | | | | |
| prescription as endorsed where there exists a record | | | | |
| Prescribed on the same prescription as insulin or a sendorsed; or | suipnonyiurea in whi | ch case the p | rescriptio | on is deemed to be |
| Prescribed for a pregnant woman with diabetes and | | | | |
| 4) Prescribed for a patient on home TPN at risk of hyper | | | | |
| Prescribed for a patient with a genetic or an acquire 2 diabetes and metabolic syndrome and endorsed a | • | e nomeostasis | s exclual | ng type 1 or type |
| | ccordingly. | | | |
| Test strips | | 50 test OP | | <u>areSens N</u> areSens PRO |
| BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED) | | | _ | |
| The number of test strips available on a prescription is res | stricted to 50 unless: | | | |
| 1) Prescribed for a patient on insulin or a sulphonylure | | | | |
| prescription as endorsed where there exists a record | | | | |
| Prescribed on the same prescription as insulin or a sendorsed; or | suipnonyiurea in whi | cn case the p | rescription | on is deemed to be |
| Prescribed for a pregnant woman with diabetes and | endorsed according | ılv: or | | |
| Prescribed for a patient on home TPN at risk of hyperative structure of the st | | | d endors | ed accordingly; or |
| Prescribed for a patient with a genetic or an acquire 2 diabetes and metabolic syndrome and endorsed a | | e homeostasis | s excludi | ng type 1 or type |
| Blood glucose test strips | | 50 test OP | ✓ S | ensoCard |
| Insulin Syringes and Needles | | | | |
| Subsidy is available for disposable insulin syringes, needles, a | and pen needles if p | rescribed on t | he same | form as the one used f |
| ne supply of insulin or when prescribed for an insulin patient a innotate the prescription as endorsed where there exists a re- | and the prescription | is endorsed a | ccording | |
| | | ang or msullfi. | | |
| NSULIN PEN NEEDLES – Maximum of 200 dev per prescrip ₭ 29 g × 12.7 mm | | 100 | ✓ B | -D Micro-Fine |
| k 31 g × 5 mm | | 100 | | -D Micro-Fine |
| | | | | |

🗸 Berpu

100

100

100

- B-D Micro-Fine
 - ✓ B-D Micro-Fine

* 31 g × 6 mm9.50

| | Subsidy (Manufacturer's Price) | Der | Fully Subsidised | Generic |
|--|-----------------------------------|-------|---------------------|-------------------|
| | \$ | Per | <i>✓</i> | Manufacturer |
| INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDL | E – Maximum of 200 | dev j | per prescri | ption |
| * Syringe 0.3 ml with 29 g × 12.7 mm needle | | 100 | 1 | B-D Ultra Fine |
| | 1.36 | 10 | | |
| | (1.99) | | | B-D Ultra Fine |
| * Syringe 0.3 ml with 31 g × 8 mm needle | | 100 | 1 | B-D Ultra Fine II |
| | 1.30 | 10 | | |
| | (1.99) | | | B-D Ultra Fine II |
| * Syringe 0.5 ml with 29 g x 12.7 mm needle | | 100 | 1 | B-D Ultra Fine |
| | 1.36 | 10 | | |
| | (1.99) | | | B-D Ultra Fine |
| * Syringe 0.5 ml with 31 g × 8 mm needle | () | 100 | 1 | B-D Ultra Fine II |
| -, , , , , , , , , , , , , , , , , , , | 1.36 | 10 | | |
| | (1.99) | | | B-D Ultra Fine II |
| * Syringe 1 ml with 29 g × 12.7 mm needle | · / | 100 | 1 | B-D Ultra Fine |
| | 1.36 | 10 | | |
| | (1.99) | | | B-D Ultra Fine |
| * Syringe 1 ml with 31 g × 8 mm needle | () | 100 | 1 | B-D Ultra Fine II |
| | 1.36 | 10 | | |
| | (1.99) | .0 | | B-D Ultra Fine II |
| Insulin Pumps | | | | |

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

| a) | Maximum of 1 dev per prescription |
|----|-----------------------------------|
| b) | Only on a prescription |

| C) | Maximum of T insulin pump per patient each four yea | ar period. | | |
|----|---|------------|---|-----------------------------------|
| Mi | in basal rate 0.025 U/h | | 1 | MiniMed 770G |
| Mi | in basal rate 0.1 U/h | | 1 | Tandem t:slim |
| | | | | X2 with Basal-IQ |

⇒SA1603 Special Authority for Subsidy

data and a factor of the second

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

.

- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

16

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

| Subsidy | Fully | | Brand or |
|------------------------|-------|-----------|--------------|
| (Manufacturer's Price) | Su | lbsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

4 Either:

4.1 Applicant is a relevant specialist; or

4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and

8 Either:

- 8.1 Applicant is a relevant specialist; or
- 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | · · · | Manufacturer | |

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

| Subsidy | Fully | Brand or |
|------------------------|------------|----------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufact |

ic Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes: and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the followina:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application - (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person): and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 vears for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events: and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

| Subs | sidy | Fully | Brand or |
|------------|---------------------|-------|--------------|
| (Manufactu | rer's Price) Subsid | lised | Generic |
| \$ | S Per | 1 | Manufacturer |

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

20

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

| | Subsidy (Manufacturer's Price |) 6 | Fully sidised | Brand or Generic |
|--|----------------------------------|-------------|------------------|----------------------------|
| | (Manulacturer's Frice \$ | Per | | Manufacturer |
| continued | | | | |
| than 80 mmol/mol; and | | | | |
| 2 The patient's HbA1c has not deteriorated more than 5 mr | nol/mol from initial a | oplication; | and | |
| 3 The patient has not had an increase in severe unexplained | ed hypoglycaemic ep | isodes fro | m baseli | ne; and |
| 4 Either: | | | | |
| 4.1 Applicant is a relevant specialist; or | | | | |
| 4.2 Applicant is a nurse practitioner working within the | eir vocational scope. | | | |
| INSULIN PUMP CARTRIDGE - Special Authority see SA1985 | on page 19 – Retail r | harmacy | | |
| a) Maximum of 3 sets per prescription | | , | | |
| b) Only on a prescription | | | | |
| c) Maximum of 13 packs of cartridge sets will be funded pe | r year. | | | |
| Cartridge 300 U, t:lock × 10 | | 1 OP | 🖌 Т | andem Cartridge |
| INSULIN PUMP INFUSION SET (STEEL CANNULA) - Special | Authority see SA198 | 5 on page | 19 – Re | etail pharmacy |
| a) Maximum of 3 sets per prescription | ,, , | 1.0 | | ···· [· ··· ···] |
| b) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | | |
| 10 mm steel needle; 60 cm tubing × 10 | 130.00 | 1 OP | 🗸 N | /iniMed Sure-T |
| | | | | MMT-884A |
| 10 mm steel needle; 80 cm tubing × 10 | | 1 OP | 🗸 N | /iniMed Sure-T |
| | | | | MMT-886A |
| 6 mm steel needle; 60 cm tubing × 10 | 130.00 | 1 OP | 🗸 N | /iniMed Sure-T |
| | | | | MMT-864A |
| 6 mm steel needle; 80 cm tubing × 10 | | 1 OP | ✓ N | AiniMed Sure-T |
| | | | | MMT-866A |
| 8 mm steel needle; 60 cm tubing × 10 | | 1 OP | ✓ N | AiniMed Sure-T |
| 0 mm staal naadlas 00 an tukina s 10 | 100.00 | 1.00 | | MMT-874A |
| 8 mm steel needle; 80 cm tubing × 10 | | 1 OP | • 1 | /iniMed Sure-T MMT-876A |
| 6 mm steel needle; 29 G; manual insertion; 60 cm tubing × | | | | WIWIT-070A |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 10 | Sure-T MMT-863 |
| 8 mm steel needle; 29 G; manual insertion; 60 cm tubing × | | 101 | • 3 | |
| 10 with 10 needles; luer lock | 130.00 | 1 OP | 1 S | Sure-T MMT-873 |
| - | | | - | |
| INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGH Retail pharmacy | 1 INSERTION = 5p | ecial Autri | only see | - 5A 1965 on page 19 - |
| a) Maximum of 3 sets per prescription | | | | |
| b) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. | | | | |
| 6 mm steel cannula; straight insertion; 80 cm line \times 10 with | | | | |
| 10 needles | | 1 OP | 🗸 1 | ruSteel |
| 8 mm steel cannula; straight insertion; 80 cm line × 10 with | | | • | |
| 10 needles | 130.00 | 1 OP | √ T | ruSteel |
| 6 mm steel cannula; straight insertion; 60 cm line \times 10 with | | | | |
| 10 needles | 130.00 | 1 OP | 🗸 I | ruSteel |
| 8 mm steel cannula; straight insertion; 60 cm line \times 10 with | | | | |
| 10 needles | | 1 OP | 🗸 I | ruSteel |
| | | | | |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|-------|---------------------|--------------------------------|
| SULIN PUMP INFUSION SET (TEFLON CANNULA) – Spec | ial Authority see SA19 | 85 on | page 19 - | - Retail pharmacy |
| a) Maximum of 3 set per prescriptionb) Only on a prescription | | | | |
| c) Maximum of 13 infusion sets will be funded per year. 13 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-382A |
| 13 mm teflon needle, 45 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-368A |
| 13 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-381A |
| 13 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-383A |
| 17 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-377A |
| 17 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-378A |
| 17 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Silhouette MMT-384A |
| 6 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-398A |
| 6 mm teflon needle, 45 cm blue tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-941A |
| 6 mm teflon needle, 45 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-921A |
| 6 mm teflon needle, 60 cm blue tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-943A |
| 6 mm teflon needle, 60 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-923A |
| 6 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-399A |
| 6 mm teflon needle, 80 cm blue tubing | 130.00 | 1 OP | 1 | MiniMed Mio MMT-945A |
| 6 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-965A |
| 6 mm teflon needle, 80 cm pink tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-925A |
| 6 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-387A |
| 9 mm teflon needle, 110 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-396A |
| 9 mm teflon needle, 60 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-397A |
| 9 mm teflon needle, 80 cm clear tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Mio MMT-975A |
| 9 mm teflon needle, 80 cm tubing × 10 | 130.00 | 1 OP | 1 | MiniMed Quick-Set MMT-386A |

| | Subsidy (Manufacturer's Price \$ | | Fully Brand or lised Generic Manufacturer |
|--|--|------------------|--|
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. | NSERTION WITH II | NSERTION D | EVICE) – Special Authority see |
| 13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles | | 1 OP | ✓ AutoSoft 30 |
| 13 mm teflon cannula; angle insertion; insertion device; 60 c line x 10 with 10 needles | | 1 OP | ✓ AutoSoft 30 |
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock | | cial Authority | see SA1985 on page 19 – ✓ Silhouette MMT-373 |
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIG see SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; insertion device; | HT INSERTION WIT | 'H INSERTIO | N DEVICE) – Special Authority |
| 110 cm line × 10 with 10 needles 6 mm teflon cannula; straight insertion; insertion device; 60 line × 10 with 10 needles | cm | 1 OP 1 OP | AutoSoft 90 AutoSoft 90 |
| 9 mm teflon cannula; straight insertion; insertion device; 110 cm line × 10 with 10 needles | | 1 OP | ✓ AutoSoft 90 |
| 9 mm teflon cannula; straight insertion; insertion device; 60 line × 10 with 10 needles | | 1 OP | ✓ AutoSoft 90 |
| INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIG Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 w | ith | | |
| 10 needles; luer lock 9 mm teflon cannula; straight insertion; 60 cm tubing × 10 w | ith | 1 OP | ✓ Quick-Set MMT-393 |
| 10 needles; luer lock INSULIN PUMP RESERVOIR – Special Authority see SA1985 (| | 1 OP oharmacy | ✓ Quick-Set MMT-392 |
| a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded pe 10 × luer lock conversion cartridges 1.8 ml for Paradigm pur Cartridge for 5 and 7 series pump; 1.8 ml × 10 | r year. nps50.00 | 1 OP 1 OP | ✓ ADR Cartridge 1.8 ✓ MiniMed 1.8 Reservoir MMT-326A |
| Cartridge for 7 series pump; 3.0 ml × 10 | 50.00 | 1 OP | MiniMed 3.0 Reservoir MMT-332A |

| | Subsidy (Manufacturer's Price \$ |) Per | Fully Subsidised | |
|---|--|------------|---------------------|-------------|
| Digestives Including Enzymes | | | | |
| PANCREATIC ENZYME | | | | |
| Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U) Creon 10000 to be Principal Supply on 1 June 2022 | | 100 | 1 | Creon 10000 |
| Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease)) | | 100 | 1 | Panzytrat |
| Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U) Creon 25000 to be Principal Supply on 1 June 2022 | 94.38 | 100 | ~ | Creon 25000 |
| Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U) | | 20 g Ol | ₽ ✔ | Creon Micro |
| (Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U a | | | | ••••• |
| URSODEOXYCHOLIC ACID – Special Authority see SA1739 be Cap 250 mg. | | icy 100 | 1 | Ursosan |
| SA1739 Special Authority for Subsidy Initial application — (Alagille syndrome or progressive famil Approvals valid without further renewal unless notified for applica Either: | ial intrahepatic cho | lestas | | |

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

24

1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

Parenteral Nutrition (TPN); and

2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

| ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription | | | |
|---|------------------|--------------|---|
| * Powder for oral soln | 6.00 | 250 g OP | Macro Organic Psyllium Husk |
| | 12.20 | 500 g OP | ✓ Konsyl-D |
| MUCILAGINOUS LAXATIVES WITH STIMULANTS | | | |
| * Dry | | 500 g OP | Newserse |
| | (17.32) 2.41 | 200 g OP | Normacol Plus |
| | (8.72) | 200 g 01 | Normacol Plus |
| Faecal Softeners | | | |
| DOCULATE SODIUM Only on a pressintian | | | |
| DOCUSATE SODIUM – Only on a prescription * Tab 50 mg | 2.31 | 100 | Coloxyl |
| * Tab 120 mg | | 100 | ✓ <u>Coloxyl</u> |
| DOCUSATE SODIUM WITH SENNOSIDES | | | |
| * Tab 50 mg with sennosides 8 mg | 4.20 | 200 | Laxsol |
| POLOXAMER – Only on a prescription | | | |
| Not funded for use in the ear. | 2.00 | 30 ml OP | |
| * Oral drops 10% | | 30 MI OP | ✓ <u>Coloxyl</u> |
| Opioid Receptor Antagonists - Peripheral | | | |
| METHYLNALTREXONE BROMIDE - Special Authority see SA | 1691 below - Ret | ail pharmacy | |
| Inj 12 mg per 0.6 ml vial | | 1 | ✓ Relistor |
| | 246.00 | 7 | Relistor |

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| (| Subsidy Manufacturer's Price \$ | e) Subs Per | Fully Brand or idised Generic ✓ Manufacturer |
|---|---------------------------------------|----------------|--|
| continued unless notified for applications meeting the following criteria: | | | |
| Both: | | | |
| The patient is receiving palliative care; and Either: | | | |
| 2.1 Oral and rectal treatments for opioid induced constip2.2 Oral and rectal treatments for opioid induced constip | | | ed. |
| Osmotic Laxatives | | | |
| GLYCEROL | | | |
| * Suppos 3.6 g – Only on a prescription | 9.25 | 20 | ✓ PSM |
| LACTULOSE – Only on a prescription | | | |
| * Oral liq 10 g per 15 ml | 3.33 | 500 ml | Laevolac |
| MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BIC | ARBONATE AND | SODIUM C | HLORIDE |
| Powder for oral soln 13.125 g with potassium chloride 46.6 mg | | | _ |
| sodium bicarbonate 178.5 mg and sodium chloride 350.7 | mg6.70 | 30 | ✓ Molaxole |
| SODIUM ACID PHOSPHATE – Only on a prescription | | | |
| Enema 16% with sodium phosphate 8% | 2.50 | 1 | Fleet Phosphate |
| | . | | Enema |
| SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE | Only on a presc | ription | |
| Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml | 20.09 | 50 | ✓ Micolette |
| 5 111 | | 50 | • MICOIelle |
| Stimulant Laxatives | | | |
| | | | |
| BISACODYL – Only on a prescription | F 00 | 000 | C Dharmaan Uaalth |
| * Tab 5 mg | 5.80 5.99 | 200 | ✓ Pharmacy Health ✓ Lax-Tab |
| Pharmacy Health to be Principal Supply on 1 June 2022 | 5.55 | | |
| * Suppos 10 mg | 3.69 | 10 | Lax-Suppositories |
| (Lax-Tab Tab 5 mg to be delisted 1 June 2022) | | | |
| SENNA – Only on a prescription | | | |
| * Tab, standardised | 2.17 | 100 | |
| | (8.21) | | Senokot |
| | 0.43 | 20 | |
| | (2.06) | | Senokot |
| SODIUM PICOSULFATE - Special Authority see SA2053 below - | | | 4 |
| Oral soln 7.5 mg per ml | 7.40 | 30 ml OP | Dulcolax SP Drop |
| SA2053 Special Authority for Subsidy | | | |

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

| (N | Subsidy lanufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|--|---------------|---------------------|-------------------------------------|
| Metabolic Disorder Agents | | | | |
| ALGLUCOSIDASE ALFA – Special Authority see SA1986 below – Inj 50 mg vial | .1,142.60 | 1 r applic | | yozyme |
| All of the following: 1 The patient is aged up to 24 months at the time of initial appl and | cation and has be | een dia | gnosed with | n infantile Pompe disease; |
| 2 Any of the following: 2.1 Diagnosis confirmed by documented deficiency of aci villus biopsies and/or cultured amniotic cells; or | | | | |
| 2.2 Documented deficiency of acid alpha-glucosidase, an elevation of glucose tetrasaccharides; or | | | • | |
| 2.3 Documented deficiency of acid alpha-glucosidase, an disease-causing mutation in the acid alpha-glucosida2.4 Documented urinary tetrasaccharide testing indicating | se gene (GAA gei | ne); or | • | |
| molecular genetic testing indicating a disease-causing | g mutation in the (| GAA ge | ne; and | |
| 3 Patient has not required long-term invasive ventilation for res (ERT); and | piratory failure pr | ior to st | arting enzy | me replacement therapy |
| Patient does not have another life-threatening or severe dise or might be reasonably expected to compromise a response Alglucosidase alfa to be administered at doses no greater that | to ERT; and | - | - | to be influenced by ERT |
| Renewal only from a metabolic physician. Approvals valid for 12 m | | | | llowing criteria: |
| All of the following: | | | 0 | 0 |
| The treatment remains appropriate for the patient and the pa Alglucosidase alfa to be administered at doses no greater that Patient has not had severe infusion-related adverse reactions and/or adjustment of infusion rates; and | an 20 mg/kg ever | y 2 wee | ks; and | |
| 4 Patient has not developed another life threatening or severe influenced by ERT; and | | - | | - |
| 5 Patient has not developed another medical condition that mig ERT; and | | | | |
| 6 There is no evidence of life threatening progression of respirative ventilation; and7 There is no evidence of new or progressive cardiomyopathy. | | evidenc | ea by the n | eeded for > 14 days of |
| ARGININE - Special Authority see SA2042 below - Retail pharmac | y. | | | |
| Tab 1,000 mg | | 90 | - | linicians |
| Cap 500 mg Powder | | 50 400 q | | olgar iomed |
| ➡SA2042 Special Authority for Subsidy | | | | |
| Initial application only from a metabolic physician. Approvals valid | for 6 months whe | ere pati | ent has a s | uspected inborn error of |
| metabolism that may respond to arginine supplementation. Renewal only from a metabolic physician. Approvals valid for 24 m | onths for applicati | ions me | eting the fo | llowing criteria: |

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment. BETAINE – Special Authority see SA1987 on the next page – Retail pharmacy

180 g OP 🖌 Cystadane

| (Manufacturer's Price) Subsidised Generic Per | Subsidy | | Fully | Brand or | |
|--|------------------------|-----|----------|--------------|--|
| \$ Per ✓ Manufacturer | (Manufacturer's Price) | | bsidised | | |
| | \$ | Per | 1 | Manufacturer | |

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and

3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation. **Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below - Retail pharmacy

| Cap 120 mg | CBŚ | 30 | Solgar |
|------------|---------|----|--------------------------------|
| Cap 160 mg | CBS | 60 | Go Healthy |

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE - Special Authority see SA1988 below - Retail pharmacy

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE - Special Authority see SA1623 on the next page - Retail pharmacy

| (Manufacturer | idy | | Fully | Brand or |
|--|-------------|---------------|-------------|-----------------------------|
| tial application only from a metabolic physician. Approvals valid for 24 we of the following: 1 The patient has been diagnosed with Hunter Syndrome (mucopolysacc 2 Either: 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase a assay in cultured skin fibroblasts; or 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase (idursulfase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. RRONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | er's Price) | Subs Per | idised ✓ | Generic Manufacturer |
| of the following: The patient has been diagnosed with Hunter Syndrome (mucopolysacc Either: Diagnosis confirmed by demonstration of iduronate 2-sulfatase assay in cultured skin fibroblasts; or Detection of a disease causing mutation in the iduronate 2-sulfatase assay in cultured skin fibroblasts; or Detection of a disease causing mutation in the iduronate 2-sulfatase assay in cultured skin fibroblasts; or Detection of a disease causing mutation in the iduronate 2-sulfatase assay in cultured skin fibroblasts; or Detection of a disease causing mutation in the iduronate 2-sulfatase assay in cultured skin fibroblasts; or Patient has not required long-term invasive ventilation for respiratory fat (ERT); and Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. RONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | | | | |
| 2 Either: 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase assay in cultured skin fibroblasts; or 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase (idursulfase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. 3RONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | | | s meeti | ng the following criteria: |
| assay in cultured skin fibroblasts; or 2.2 Detection of a disease causing mutation in the iduronate 2-sulfa 3 Patient is going to proceed with a haematopoietic stem cell transplant (idursulfase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. IRONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | charidosi | s II); and | | |
| 3 Patient is going to proceed with a haematopoietic stem cell transplant (idursulfase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. RONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | | - | blood | cells by either enzyme |
| 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. (RONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | | | next 3 r | nonths and treatment wit |
| 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 greater than 0.5 mg/kg every week. IRONIDASE – Special Authority see SA1695 below – Retail pharmacy Inj 100 U per ml, 5 ml vial | ailure pric | or to startir | ng Enzy | me Replacement Thera |
| Inj 100 U per ml, 5 ml vial | 2 weeks p | re- and 12 | 2 weeks | s post-HSCT) at doses n |
| tial application only from a metabolic physician. Approvals valid for 24 we of the following: The patient has been diagnosed with Hurler Syndrome (mucopolysaccl 2 Either: 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase d assay in cultured skin fibroblasts; or 2.2 Detection of two disease causing mutations in the alpha-L-iduron to have Hurler syndrome; and Patient is going to proceed with a haematopoietic stem cell transplant (laronidase would be bridging treatment to transplant; and Patient has not required long-term invasive ventilation for respiratory fa (ERT); and Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. | ò | 1 | ✓ A | Idurazyme |
| The patient has been diagnosed with Hurler Syndrome (mucopolysacch 2 Either: 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase d assay in cultured skin fibroblasts; or 2.2 Detection of two disease causing mutations in the alpha-L-iduron to have Hurler syndrome; and Patient is going to proceed with a haematopoietic stem cell transplant (laronidase would be bridging treatment to transplant; and Patient has not required long-term invasive ventilation for respiratory fa (ERT); and Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. VOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mgCBS Cap 250 mgCBS Cap 500 mgCBS Oral liq 500 mg per 10 mlCBS | eeks for a | pplication | s meeti | ing the following criteria: |
| assay in cultured skin fibroblasts; or 2.2 Detection of two disease causing mutations in the alpha-L-iduro to have Hurler syndrome; and 3 Patient is going to proceed with a haematopoietic stem cell transplant (laronidase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. EVOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mgCBS Cap 250 mgCBS Cap 500 mgCBS Oral liq 500 mg per 10 mlCBS | chardosis | I-H); and | | |
| to have Hurler syndrome; and 3 Patient is going to proceed with a haematopoietic stem cell transplant (laronidase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. EVOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mgCBS Cap 250 mgCBS Cap 500 mgCBS Oral liq 500 mg per 10 mlCBS | | | | |
| Iaronidase would be bridging treatment to transplant; and 4 Patient has not required long-term invasive ventilation for respiratory fa (ERT); and 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. VOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mg CBS | • | | | - |
| (ERT); and 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 than 100 units/kg every week. VOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mg | . , | | | |
| VOCARNITINE – Special Authority see SA2040 below – Retail pharmacy Tab 500 mg | | | • • | |
| Cap 250 mg CBS Cap 500 mg CBS Oral liq 500 mg per 10 ml CBS SA2040 Special Authority for Subsidy | | | | |
| Cap 500 mgCBS Oral liq 500 mg per 10 mlCBS SA2040 Special Authority for Subsidy | | 30 30 | | olgar olgar |
| SA2040 Special Authority for Subsidy | | 60 | ✓ В | alance |
| | 3 | 00 ml | ✓ B | alance |
| | nths wher | re patient | has a s | uspected inborn error of |
| etabolism that may respond to carnitine supplementation. enewal only from a metabolic physician. Approvals valid for 24 months for a | applicatio | ns meetin | na the fo | ollowing criteria: |

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

| RIBOFLAVIN - Special Authority see SA2041 on the next page - | Retail pharmacy | Y | |
|--|-------------------------------------|-----|----------------------------------|
| Tab 100 mg | CBS | 100 | Country Life |
| Cap 100 mg | CBS | 100 | Solgar |

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) | Full Subsidise | d Generic |
|-----------------------------------|-------------------|--------------|
| þ | Per 🗸 | Manufacturer |

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and

2 The treatment remains appropriate and the patient is benefiting from treatment.

➡SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
 - 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
 - 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
 - 4 Sapropterin to be used alone or in combination with PKU dietary management; and
 - 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy

Soln 100 mg per mlCBS 100 ml 🖌 Amzoate 529

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

| SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - | - Retail pharmacy | / |
|---|-------------------|-----------|
| Grans 483 mg per g2,016.00 | 174 g OP 🔹 | Pheburane |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

■ SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cvcle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

| Cap 500 mg | CBS | 50 | Solgar |
|--------------|-----|-------|------------------------------------|
| Cap 1,000 mg | | 90 | Life Extension |
| Powder | | 300 g | Life Extension |

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA1880 below - Retail pharmacy Inj 200 unit vial......1,072.00 1

Elelvso

■ SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

| The Co-ordinator, Gaucher Treatment Panel | Phone: 04 460 4990 |
|---|-------------------------------------|
| Pharmac PO Box 10 254 | Facsimile: 04 916 7571 |
| Wellington | Email: gaucherpanel@pharmac.govt.nz |

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa: and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 6) 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia: at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

massive symptomatic splenomegaly; or

- 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
- 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
- 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
- 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

| BENZYDAMINE HYDROCHLORIDE | | | |
|--|------------------|-------------------|---------------------------------|
| Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with | | | |
| Endorsement | 9.00 | 500 ml | |
| | (20.31) | | Difflam |
| Additional subsidy by endorsement for a patient who has prescription is endorsed accordingly. | oral mucositis a | s a result of tre | eatment for cancer, and the |
| CARMELLOSE SODIUM WITH GELATIN AND PECTIN | | | |
| Paste | 17.20 | 56 g OP | Stomahesive |
| | 4.55 | 15 g OP | |
| | (7.90) | | Orabase |
| | 1.52 | 5 g OP | |
| | (3.60) | | Orabase |
| Powder | 8.48 | 28 g OP | |
| | (10.95) | | Stomahesive |
| CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE | | | |
| * Adhesive gel 8.7% with cetalkonium chloride 0.01% | 2.06 | 15 g OP | |
| | (6.00) | • | Bonjela |

| | 0 1 1 | | |
|--|-----------------------------|------------|--|
| /A | Subsidy Manufacturer's P | rico) Subo | Fully Brand or idised Generic |
| (r | s | Per | Manufacturer |
| RIAMCINOLONE ACETONIDE | | - | |
| | F 00 | | Kanalay in Orahaaa |
| Paste 0.1% | | 5 g OP | Kenalog in Orabase |
| Oropharyngeal Anti-infectives | | | |
| MPHOTERICIN B | | | |
| Lozenges 10 mg | 5.86 | 20 | Fungilin |
| IICONAZOLE | | | |
| Oral gel 20 mg per g | 4.74 | 40 g OP | Decozol |
| YSTATIN | | Ũ | |
| Oral liq 100,000 u per ml | 1 76 | 24 ml OP | Nilstat |
| | | 24 111 01 | • <u>motat</u> |
| Vitamins | | | |
| Vitamin B | | | |
| YDROXOCOBALAMIN | | | |
| Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO | 1.89 | 3 | ✓ Vita-B12 |
| | 2.84 | | Neo-B12 |
| | 3.15 | 5 | Hydroxocobalamin |
| | | | Mercury Pharma |
| YRIDOXINE HYDROCHLORIDE | | | |
| a) No more than 100 mg per dose | | | |
| b) Only on a prescription | | | |
| Tab 25 mg – No patient co-payment payable | 2.70 | 90 | Vitamin B6 25 |
| Tab 50 mg | | 500 | ✓ Apo-Pyridoxine |
| 5 | 23.45 | | Pyridoxine |
| | | | multichem |
| Apo-Pyridoxine Tab 50 mg to be delisted 1 May 2022) | | | |
| HIAMINE HYDROCHLORIDE – Only on a prescription | | | |
| Tab 50 mg | 7 09 | 100 | 🗸 Max Health |
| - | | 100 | • max nounn |
| ITAMIN B COMPLEX | 715 | 500 | - Paley |
| Tab, strong, BPC | | 500 | Bplex |
| Vitamin C | | | |
| SCORBIC ACID | | | |
| a) No more than 100 mg per dose | | | |
| b) Only on a prescription | | | |
| € Tab 100 mg | 9.90 | 500 | ✓ Cvite |
| | | | <u>•••••</u> |
| Vitamin D | | | |
| LFACALCIDOL | | | |
| Cap 0.25 mcg | 26.32 | 100 | One-Alpha |
| Cap 1 mcg | 87.98 | 100 | One-Alpha |
| Oral drops 2 mcg per ml | | 20 ml OP | One-Alpha |
| ALCITRIOL | | | |
| ← Cap 0.25 mcg | 7.95 | 100 | ✓ Calcitriol-AFT |
| Cap 0.5 mcg | | 100 | ✓ Calcitriol-AFT |
| | | | |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Pr | ice) Subsid | |
|--|-------------------------------|---------------------|--|
| | \$ | Per | Manufacturer |
| COLECALCIFEROL | 0.05 | 10 | |
| Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescript Oral lig 188 mcg per ml (7,500 iu per ml) | | 12 4.8 ml OP | ✓ <u>Vit.D3</u> ✓ Puria |
| | 9.00 | 4.0 IIII OF | • Fulla |
| Multivitamin Preparations | | | |
| MULTIVITAMIN RENAL – Special Authority see SA1546 below - * Cap | | 30 | Clinicians Renal Vit |
| SA1546 Special Authority for Subsidy | 0.43 | 50 | |
| Initial application from any relevant practitioner. Approvals valit the following criteria: Either: | d without further r | enewal unless | notified for applications meeting |
| The patient has chronic kidney disease and is receiving e The patient has chronic kidney disease grade 5, defined a 15 ml/min/1.73 m² body surface area (BSA). | | | |
| MULTIVITAMINS - Special Authority see SA1036 below - Retain | | | |
| * Powder | | 200 g OP | Paediatric Seravit |
| SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali- inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without | | | |
| approval for multivitamins. | | | |
| VITAMINS * Tab (BPC cap strength) | 11.45 | 1,000 | ✓ Mvite |
| * Cap (fat soluble vitamins A, D, E, K) – Special Authority see | | 1,000 | |
| SA1720 below – Retail pharmacy | 23.40 | 60 | Vitabdeck |
| SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valie the following criteria: Any of the following: Patient has cystic fibrosis with pancreatic insufficiency; or Patient is an infant or child with liver disease or short gut s Patient has severe malabsorption syndrome. | | enewal unless | notified for applications meeting |
| Minerals | | | |
| Calcium | | | |
| CALCIUM CARBONATE | | | |
| * Tab 1.25 g (500 mg elemental) | 6.69 | 250 | <u>Calci-Tab 500</u> |
| Tab eff 1.25 g (500 mg elemental) – Subsidy by endorseme Subsidy by endorsement – Only when prescribed for par considered unsuitable. | | 76 5 years) wher | Cacit S29 calcium carbonate oral liquid i |
| CALCIUM GLUCONATE | | | |
| * Inj 10%, 10 ml ampoule | | 10 | Max Health - |
| | 04.00 | 00 | Hamein S29 |
| | 64.00 | 20 | Max Health \$29 |
| | | | |

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Generic |
|---|-----------------------------------|--------|---------------------|--------------------------------|
| | \$ | Per | | Manufacturer |
| CALCIUM LACTATE GLUCONATE WITH CALCIUM CARBONA | | | | |
| Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg element – Subsidy by endorsement | | 20 | | Calcium-Sandoz |
| | | 20 | • | Forte S29 |
| | 260.00 | 100 | 1 | Calcium-Sandoz |
| | 200100 | | | Forte S29 |
| Subsidy by endorsement – Only when prescribed for pae considered unsuitable. | ediatric patients (< 5 | years) | where ca | lcium carbonate oral liquid is |
| Fluoride | | | | |
| SODIUM FLUORIDE | | | | |
| * Tab 1.1 mg (0.5 mg elemental) | | 100 | 1 | PSM |
| | | | | |
| lodine | | | | |
| POTASSIUM IODATE | | | | |
| * Tab 253 mcg (150 mcg elemental iodine) | 4.58 | 90 | ✓ | NeuroTabs |
| Iron | | | | |
| FERROUS FUMARATE | | | | |
| * Tab 200 mg (65 mg elemental) Ferro-tab to be Principal Supply on 1 May 2022 | 3.04 | 100 | ~ | Ferro-tab |
| FERROUS FUMARATE WITH FOLIC ACID | | | | |
| * Tab 310 mg (100 mg elemental) with folic acid 350 mcg | 5.98 | 100 | ✓ | Ferro-F-Tabs |
| FERROUS SULFATE | | | | |
| * Tab long-acting 325 mg (105 mg elemental) | | 30 | | Ferrograd |
| * Oral liq 30 mg (6 mg elemental) per 1 ml | | 500 m | | Ferodan |
| IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority se | | | | Foriniaat |
| Inj 50 mg per ml, 10 ml vial | | 1 | • | Ferinject |
| SA1840 Special Authority for Subsidy | | | | |

SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and

- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
 - 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

| (Man | Subsidy | Fully | | Brand or |
|---------------------------------------|-------------------|------------|---|--------------|
| | Jacturer's Price) | Subsidised | | Generic |
| · · · · · · · · · · · · · · · · · · · | \$ | Per | ✓ | Manufacturer |

continued...

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

| * Inj 50 mg per ml, 2 ml ampoule | 5 | Ferrosig |
|--|--------|------------------------------------|
| Magnesium | | |
| MAGNESIUM HYDROXIDE Suspension 8% | 355 ml | ✓ Phillips Milk of Magnesia S29 |
| MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule25.53 | 10 | ✓ <u>Martindale</u> |
| Zinc | | |
| ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)11.00 | 100 | ✓ Zincaps |
Subsidised

Per

Fully

Subsidy (Manufacturer's Price) \$ Brand or Generic Manufacturer

Antianaemics

Hypoplastic and Haemolytic

➡SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or

3.2 Both:

- 3.2.1 Patient has diabetes mellitus; and
- 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
- 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy. Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following

criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|--------|---------------------|-------------------------|
| EPOETIN ALFA - Special Authority see SA1775 on the previou | is page – Retail pharr | nacy | | |
| Wastage claimable | | | | |
| Inj 1,000 iu in 0.5 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 2,000 iu in 1 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 3,000 iu in 0.3 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 4,000 iu in 0.4 ml, syringe | 96.50 | 6 | ✓ | Binocrit |
| Inj 5,000 iu in 0.5 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 6,000 iu in 0.6 ml, syringe | 145.00 | 6 | ✓ | Binocrit |
| Inj 8,000 iu in 0.8 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 10,000 iu in 1 ml, syringe | | 6 | ✓ | Binocrit |
| Inj 40,000 iu in 1 ml, syringe | | 1 | 1 | Binocrit |
| Megaloblastic | | | | |
| FOLIC ACID | | | | |
| Tab 0.8 mg | 21.84 | 1.000 |) 🗸 | Apo-Folic Acid |
| | 26.60 | ., | | Folic Acid multichem |
| * Tab 5 mg | 5.82 | 100 | ✓ | Folic Acid Mylan |
| Oral liq 50 mcg per ml | | 5 ml C | DP 🗸 | Biomed |
| (Apo-Folic Acid Tab 0.8 mg to be delisted 1 May 2022) | | | | |

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

| Inj 250 iu vial. | | ່ 1 | Alprolix |
|---|----------|-----|------------|
| Inj 500 iu vial | | 1 | ✓ Alprolix |
| Inj 1,000 iu vial | | 1 | Alprolix |
| Inj 2,000 iu vial | | 1 | Alprolix |
| Inj 3,000 iu vial | | 1 | Alprolix |
| ELTROMBOPAG - Special Authority see SA1743 belo | | | |
| Wastage claimable | 4 550 00 | | |
| Tab 25 mg | 1,550.00 | 28 | Revolade |
| Tab 50 mg | | 28 | Revolade |

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and

3 Any of the following:

- 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
- 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
- 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant muccoutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
 - 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

| Inj 30 mg in 1 ml vial | 1 | Hemlibra |
|---------------------------|-------|------------------------------|
| Inj 60 mg in 0.4 ml vial | 1 | Hemlibra |
| Inj 105 mg in 0.7 ml vial | 1 | Hemlibra |
| Inj 150 mg in 1 ml vial | 1 | Hemlibra |

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

continued...

| Subsidy | | Fully | Brand or |
|------------------------|------|---------|--------------|
| (Manufacturer's Price) | Subs | sidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

| Inj 1 mg syringe1,1 | 78.30 1 | NovoSeven RT |
|---------------------|---------|--------------|
| Inj 2 mg syringe | 56.60 1 | NovoSeven RT |
| Inj 5 mg syringe | 91.50 1 | NovoSeven RT |
| Inj 8 mg syringe | 26.40 1 | NovoSeven RT |

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

| Inj 500 U | | 1 | FEIBA NF |
|-------------|----------|---|------------|
| Inj 1,000 U | 2,630.00 | 1 | 🖌 FEIBA NF |
| Inj 2,500 U | 6,575.00 | 1 | 🖌 FEIBA NF |

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

| Inj 250 iu prefilled syringe | 1 | 🗸 Xyntha |
|--------------------------------|-------|----------------------------|
| Inj 500 iu prefilled syringe | 1 | 🗸 Xyntha |
| Inj 1,000 iu prefilled syringe | 1 | Xyntha |
| Inj 2,000 iu prefilled syringe | 1 | 🗸 Xyntha |
| Inj 3,000 iu prefilled syringe | 1 | 🗸 Xyntha |

| | Subsidy (Manufacturaria Driac) | _ | Fully | Brand or |
|--|---|------------------------|----------------------------------|---|
| | (Manufacturer's Price) \$ | Per Su | bsidised | Generic Manufacturer |
| | | 1.01 | - | Manulaotaron |
| IONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpl For patients with haemophilia. Access to funded treatm | | omonhili | a Troate | re Group in conjunction |
| with the National Haemophilia Management Group. | lent is managed by the ha | aemophin | a meate | |
| Inj 500 iu vial | 435.00 | 1 | ~ 1 | RIXUBIS |
| Inj 1,000 iu vial | | 1 | - | RIXUBIS |
| Inj 2,000 iu vial | | 1 | - | RIXUBIS |
| Inj 3,000 iu vial | | 1 | I | RIXUBIS |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATI | E) – [Xpharm] | | | |
| For patients with haemophilia. Preferred Brand of short | | or VIII. A | ccess to | funded treatment is |
| managed by the Haemophilia Treaters Group in conjunc | | | | |
| Inj 250 iu vial | | 1 | | Advate |
| Inj 500 iu vial | | 1 | I | Advate |
| Inj 1,000 iu vial | | 1 | I | Advate |
| lnj 1,500 iu vial | , | 1 | | Advate |
| Inj 2,000 iu vial | | 1 | - | Advate |
| Inj 3,000 iu vial | | 1 | | Advate |
| OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGEN | | | | |
| For patients with haemophilia. Rare Clinical Circumstar | | | | |
| treatment is managed by the Haemophilia Treaters Gro | up in conjunction with the | National | Haemop | philia Management Group |
| subject to criteria. | | | | |
| Inj 250 iu vial | | 1 | | Kogenate FS |
| Inj 500 iu vial | | 1 | | Kogenate FS |
| Inj 1,000 iu vial | | 1 | | Kogenate FS |
| Inj 2,000 iu vial | | 1 | | Kogenate FS |
| Inj 3,000 iu vial | 2,850.00 | 1 | ✓ | Kogenate FS |
| RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR | VIII] – [Xpharm] | | | |
| For patients with haemophilia A receiving prophylaxis tr | eatment. Access to funde | ed treatm | ent is ma | anaged by the Haamonhi |
| Treaters Group in conjunction with the National Usama | | | | anaged by the nachtoph |
| Treaters Group in conjunction with the National Haemor | philia Management group | | | |
| Inj 250 iu vial | bhilia Management group | 1 | • | Adynovate |
| Inj 250 iu vial Inj 500 iu vial | bhilia Management group 300.00 600.00 | 1 1 | I I | Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial | bhilia Management group | 1 1 1 | 5 5 5 | Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial | bhilia Management group | 1 1 | 5 5 5 | Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE | bilia Management group | 1 1 1 1 | 5 5 5 | Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE | billia Management group | 1 1 1 | | Adynovate Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial | bilia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE Inj 3% 2 ml TRANEXAMIC ACID | bilia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE Inj 3% 2 ml | bilia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE | bilia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K | bilia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE Inj 3% 2 ml 'RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE | billia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein <u>Mercury Pharma</u> |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO | billia Management group | 1 1 1 1 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein Mercury Pharma Konakion MM |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE | billia Management group | 1 1 1 5 60 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein <u>Mercury Pharma</u> |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO | billia Management group | 1 1 1 5 60 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein Mercury Pharma Konakion MM |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO Antithrombotic Agents | billia Management group | 1 1 1 5 60 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein Mercury Pharma Konakion MM |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml TRANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO Antithrombotic Agents Antiplatelet Agents | billia Management group | 1 1 1 5 60 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein Mercury Pharma Konakion MM |
| Inj 250 iu vial Inj 500 iu vial Inj 1,000 iu vial Inj 2,000 iu vial SODIUM TETRADECYL SULPHATE k Inj 3% 2 ml RANEXAMIC ACID Tab 500 mg Vitamin K PHYTOMENADIONE Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO Antithrombotic Agents | billia Management group | 1 1 1 5 60 | | Adynovate Adynovate Adynovate Adynovate Fibro-vein Mercury Pharma Konakion MM |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| CLOPIDOGREL * Tab 75 mg | 4.60 | 84 | ✓ <u>c</u> | lopidogrel Multichem |
| DIPYRIDAMOLE * Tab long-acting 150 mg | | 60 | ✓ <u>P</u> | ytazen SR |
| TICAGRELOR – Special Authority see SA1955 below – Retail pha * Tab 90 mg | | 56 | ✔ В | rilinta |

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and

2 Either:

- 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
- 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal - (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for

continued...

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM – Special Authority see SA1646 below – Retail pharmacy

| Inj 20 mg in 0.2 ml syringe | 10 | Clexane |
|------------------------------|--------|-----------------------------------|
| Inj 40 mg in 0.4 ml syringe | 10 | Clexane |
| Inj 60 mg in 0.6 ml syringe | 10 | Clexane |
| Inj 80 mg in 0.8 ml syringe | 10 | Clexane |
| Inj 100 mg in 1 ml syringe | 10 | Clexane |
| Inj 120 mg in 0.8 ml syringe | 10 | Clexane Forte |
| Inj 150 mg in 1 ml syringe | 10 | Clexane Forte |
| | | |

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

| | Subsidy | | Fully | |
|---|------------------------------|-----|------------|-------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| HEPARIN SODIUM | | | | |
| Inj 1,000 iu per ml, 5 ml ampoule | 72.84 | 50 | 1 | Pfizer |
| Inj 5,000 iu per ml, 1 ml | | 5 | 1 | DBL Heparin |
| | | | | Sodium S29 |
| | 70.33 | | ✓ | Hospira |
| Inj 5,000 iu per ml, 5 ml ampoule | | 50 | ✓ | Pfizer |
| Inj 25,000 iu per ml, 0.2 ml | | 5 | ✓ | Hospira |
| | 42.40 | | ✓ | Heparin DBL S29 |
| | 482.20 | 50 | 1 | Heparin DBL S29 |
| HEPARINISED SALINE | | | | |
| Inj 10 iu per ml, 5 ml | 65 48 | 50 | 1 | Pfizer |
| | | 00 | | 1 11201 |
| Oral Anticoagulants | | | | |
| DABIGATRAN | | | | |
| Cap 75 mg – No more than 2 cap per day | | 60 | 1 | Pradaxa |
| Cap 110 mg | | 60 | 1 | Pradaxa |
| Cap 150 mg | | 60 | ✓ | Pradaxa |
| RIVAROXABAN | | | | |
| Tab 10 mg - No more than 1 tab per day | | 30 | 1 | Xarelto |
| Tab 15 mg - Up to 14 tab available on a PSO | | 28 | ✓ | Xarelto |
| Tab 20 mg | | 28 | ✓ | Xarelto |
| WARFARIN SODIUM | | | | |
| Note: Marevan and Coumadin are not interchangeable. | | | | |
| * Tab 1 mg | | 50 | 1 | Coumadin |
| | 6.46 | 100 | 1 | Marevan |
| * Tab 2 mg | 4.31 | 50 | ✓ | Coumadin |
| * Tab 3 mg | | 100 | 1 | Marevan |
| * Tab 5 mg | | 50 | | Coumadin |
| | 11.48 | 100 | 1 | Marevan |

Blood Colony-stimulating Factors

| FILGRASTIM - Special Authority see SA1259 below - Retail phar | macy | | |
|---|--------|----|------------|
| Inj 300 mcg per 0.5 ml prefilled syringe | | 10 | ✓ Nivestim |
| Inj 480 mcg per 0.5 ml prefilled syringe | 148.58 | 10 | ✓ Nivestim |

SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*): or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5×10^{9} /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 on the next page - Retail pharmacy

✓ Neulastim

44

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

| GLUCOSE [DEXTROSE] | | | |
|---|------------------|-----------------|--------------------------------|
| * Inj 50%, 10 ml ampoule - Up to 5 inj available on a PSO | 0.65 | 5 🖌 | Biomed |
| * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO18 | 5.00 | 1 🖌 | Biomed |
| POTASSIUM CHLORIDE | | | |
| * Inj 75 mg per ml, 10 ml65 | 5.00 5 | 50 🗸 | Juno |
| SODIUM BICARBONATE | | | |
| Inj 8.4%, 50 ml2 | 1.40 | 1 🖌 | Biomed |
| a) Up to 5 inj available on a PSO | | | |
| b) Not in combination | | | |
| Inj 8.4%, 100 ml2 | 1.95 | 1 🗸 | Biomed |
| a) Up to 5 inj available on a PSO | | | |
| b) Not in combination | | | |
| SODIUM CHLORIDE | | | |
| Not funded for use as a nasal drop. Not funded for nebuliser use exc | ept when used | I in conjunctio | on with an antibiotic intended |
| for nebuliser use. | | | |
| Inj 0.9%, bag – Up to 2000 ml available on a PSO | | | Baxter |
| | - ,- | | Baxter |
| Only if prescribed on a prescription for renal dialysis, maternity or | r post-natal car | e in the home | e of the patient, or on a PSO |
| for emergency use. (500 ml and 1,000 ml packs) | | | |
| Inj 23.4% (4 mmol/ml), 20 ml ampoule | | 5 🗸 | Biomed |
| For Sodium chloride oral liquid formulation refer Standard Formul | | | |
| Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO | | | Fresenius Kabi |
| Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO | 5.40 5 | 50 🗸 | Fresenius Kabi |

TOTAL PARENTERAL NUTRITION (TPN)

 On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or

20

1 OP

2) On a bulk supply order; or

WATER

3) When used in the extemporaneous compounding of eye drops; or

InfusionCBS

4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

| Inj 10 ml ampoule – Up to 5 inj available on a PSO | 50 | Pfizer |
|--|----|----------------------------|
| Inj 20 ml ampoule – Up to 5 inj available on a PSO | 20 | Fresenius Kabi |
| | | / Marthlada and |

Multichem

Fresenius Kabi

TPN

| | Subsidy (Manufacturer's F \$ | Price) Subsi Per | Fully Brand or idised Generic ✓ Manufacturer |
|--|------------------------------------|---------------------|--|
| Oral Administration | | | |
| CALCIUM POLYSTYRENE SULPHONATE Powder | 169.85 | 300 g OP | Calcium Resonium |
| COMPOUND ELECTROLYTES Powder for oral soln – Up to 5 sach available on a PSO | 9.77 | 50 | ✓ Electral |
| COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml) | 6.55 | 1,000 ml OP | Pedialyte - Bubblegum |
| PHOSPHORUS Tab eff 500 mg (16 mmol) | | 100 | Phosphate Phebra |
| POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq) | | 60 | |
| * Tab long-acting 600 mg (8 mmol) | (11.85) | 200 | Chlorvescent Span-K |
| SODIUM BICARBONATE Cap 840 mg | 8.52 | 100 | ✓ Sodibic |
| SODIUM POLYSTYRENE SULPHONATE | | | ✓ Sodibic |
| Powder | | 454 g OP | Resonium-A |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|---|
| Alpha-Adrenoceptor Blockers | | | | |
| Alpha Adrenoceptor Blockers | | | | |
| DOXAZOSIN ¥ Tab 2 mg | 17.35 | 500 | | Apo-Doxazosin Doxazosin Clinect |
| * Tab 4 mg | 20.94 | 500 | 1 | Apo-Doxazosin Doxazosin Clinect |
| (Apo-Doxazosin Tab 2 mg to be delisted 1 September 2022) (Apo-Doxazosin Tab 4 mg to be delisted 1 September 2022) PHENOXYBENZAMINE HYDROCHLORIDE | | | | |
| * Cap 10 mg | | 30 | 1 | BNM S29 |
| | 216.67 | 100 | 1 | Dibenzyline S29 |
| PRAZOSIN ₩ Tab 1 mg | 5.53 | 100 | | Apo-Prazosin Arrotex-Prazosin S29 S29 |
| * Tab 2 mg | 7.00 | 100 | | Apo-Prazosin Arrotex-Prazosin S29 S29 |
| ₩ Tab 5 mg | 11.70 | 100 | | Apo-Prazosin Arrotex-Prazosin S29 S29 |

(Apo-Prazosin Tab 1 mg to be delisted 1 May 2022) (Apo-Prazosin Tab 2 mg to be delisted 1 May 2022) (Apo-Prazosin Tab 5 mg to be delisted 1 May 2022)

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

| CAPTOPRIL | | | |
|---|------|----------|---------------------------------------|
| * Oral liq 5 mg per ml Oral liquid restricted to children under 12 year | | 95 ml OP | Capoten |
| CILAZAPRIL – Subsidy by endorsement | | | |
| Subsidy by endorsement – Subsidised for patient: endorsed accordingly. Pharmacists may annotate dispensing of cilazapril. | | | |
| * Tab 0.5 mg | 2.09 | 90 | ✓ Zapril |
| * Tab 2.5 mg | 4.80 | 90 | Zapril |
| Tab 5 mg | 8.35 | 90 | ✓ Zapril |
| ENALAPRIL MALEATE | | | |
| * Tab 5 mg | | 100 | Acetec |
| * Tab 10 mg | 2.02 | 100 | ✓ Acetec |
| * Tab 20 mg | | 100 | ✓ Acetec |
| LISINOPRIL | | | |
| * Tab 5 mg | | 90 | Ethics Lisinopril |
| * Tab 10 mg | | 90 | Ethics Lisinopril |
| * Tab 20 mg | | 90 | Ethics Lisinopril |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Outside | | F II. | Durandina |
|---|-----------------------------------|----------|---------------------|------------------------------|
| | Subsidy (Manufacturer's Price) | | Fully Subsidised | |
| | (Manulacialei si lice) | Per | | Manufacturer |
| PERINDOPRIL | | | | |
| Tab 2 mg | 1.58 | 30 | 1 | Coversyl |
| Tab 4 mg | | 30 | | Coversyl |
| QUINAPRIL | | | | <u></u> |
| Tab 5 mg | 5 97 | 90 | 1 | Arrow-Quinapril 5 |
| Tab 10 mg | | 90 | | Arrow-Quinapril 10 |
| Tab 20 mg | | 90 | | Arrow-Quinapril 20 |
| ACE Inhibitors with Diuretics | | | | |
| | | | | |
| QUINAPRIL WITH HYDROCHLOROTHIAZIDE | 4.10 | 30 | | Accuretic 10 |
| Tab 10 mg with hydrochlorothiazide 12.5 mg * Tab 20 mg with hydrochlorothiazide 12.5 mg | | 30 30 | | Accuretic 20 |
| | | 30 | • | Accuretic 20 |
| Angiotensin II Antagonists | | | | |
| CANDESARTAN CILEXETIL | | | _ | |
| * Tab 4 mg | | 90 | | Candestar |
| * Tab 8 mg | | 90 | | Candestar |
| * Tab 16 mg | | 90 | | Candestar Candestar |
| * Tab 32 mg | 5.26 | 90 | • | Candestar |
| LOSARTAN POTASSIUM | | | | |
| Tab 12.5 mg | | 84 | | Losartan Actavis |
| Tab 25 mg | | 84 | | Losartan Actavis |
| Tab 50 mg | | 84 | | Losartan Actavis |
| Tab 100 mg | 3.50 | 84 | • | Losartan Actavis |
| Angiotensin II Antagonists with Diuretics | | | | |
| LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE | | | | |
| Tab 50 mg with hydrochlorothiazide 12.5 mg | 15.25 | 30 | 1 | Arrow-Losartan & |
| | | | | Hydrochlorothiazide |
| Angiotensin II Antagonists with Neprilysin Inhi | bitors | | | |
| | | | | |
| SACUBITRIL WITH VALSARTAN – Special Authority see SA19 Note: Due to the angiotensin II receptor blocking activity of | | | | he co-administered with an |
| ACE inhibitor or another ARB. | | | | |
| Tab 24.3 mg with valsartan 25.7 mg | | 56 | 1 | Entresto 24/26 |
| Tab 48.6 mg with valsartan 51.4 mg | | 56 | | Entresto 49/51 |
| Tab 97.2 mg with valsartan 102.8 mg | | 56 | 1 | Entresto 97/103 |
| ► SA1905 Special Authority for Subsidy | | | | |
| Initial application from any relevant practitioner. Approvals val | id for 12 months for ap | plica | ations mee | ting the following criteria: |
| All of the following: | | | | 0 0 |
| 1 Patient has heart failure; and | | | | |
| 2 Any of the following: | | | | |
| 2.1 Patient is in NYHA/WHO functional class II; or | | | | |
| 2.2 Patient is in NYHA/WHO functional class III; or | | | | |
| 2.3 Patient is in NYHA/WHO functional class IV; and | | | | |
| | | | | |

continued...

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

| For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesth | etics, Local, <mark>pa</mark> | age 119 | |
|---|-------------------------------|---------|--|
| AMIODARONE HYDROCHLORIDE | | | |
| ▲ Tab 100 mg | | 30 | ✓ Aratac |
| ▲ Tab 200 mg | 5.25 | 30 | Aratac |
| Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a | | | _ |
| PSO | 16.37 | 10 | ✓ Max Health |
| ATROPINE SULPHATE | | | |
| * Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a | | | |
| PSO | 15.09 | 10 | Martindale |
| DIGOXIN | | | |
| * Tab 62.5 mcg – Up to 30 tab available on a PSO | 7.00 | 240 | Lanoxin PG |
| * Tab 250 mcg – Up to 30 tab available on a PSO | 15.20 | 240 | <u>Lanoxin</u> |
| * Oral liq 50 mcg per ml | 16.60 | 60 ml | Lanoxin |
| | | | Lanoxin Paediatric |
| | | | Elixir S29 |
| | | | Lanoxin S29 S29 |
| DISOPYRAMIDE PHOSPHATE | | | |
| ▲ Cap 100 mg | 23.87 | 100 | Rythmodan |
| FLECAINIDE ACETATE | | | • |
| ▲ Tab 50 mg | | 60 | Flecainide BNM |
| Cap long-acting 100 mg | | 90 | ✓ Flecainide |
| | | | Controlled |
| | | | Release Teva |
| Cap long-acting 200 mg | 61.06 | 90 | Flecainide |
| | | | Controlled |
| | | | Release Teva |
| Inj 10 mg per ml, 15 ml ampoule | 100.00 | 5 | Tambocor |
| MEXILETINE HYDROCHLORIDE | | | |
| ▲ Cap 150 mg | 162.00 | 100 | Teva S29 |
| ▲ Cap 250 mg | 202.00 | 100 | Teva S29 |
| PROPAFENONE HYDROCHLORIDE | | | |
| Tab 150 mg | 40.90 | 50 | Rytmonorm |
| _ · · · · · · · · · · · · · · · · · · · | | | |
| Antihypotensives | | | |
| MIDODRINE – Special Authority see SA1474 on the next page – R | otail pharmaa | | |
| Tab 2.5 mg | | 100 | ✓ Gutron |
| Tab 5 mg | | 100 | ✓ Gutron |
| 1 uo o 11g | | 100 | |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | F | ully | Brand or |
|------------------------|---------|------|--------------|
| (Manufacturer's Price) | Subsidi | sed | Generic |
| \$ | Per | ✓ | Manufacturer |

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

| ATENOLOL | | | |
|---|---------|-----------|---|
| * Tab 50 mg | 9.33 | 500 | Mylan Atenolol |
| * Tab 100 mg | 14.20 | 500 | Mylan Atenolol |
| * Oral liq 25 mg per 5 ml | 21.25 | 300 ml OP | Atenolol AFT |
| | | | S29 S29 |
| | 38.20 | | Essential |
| | | | Generics S29 |
| | 49.85 | | Atenolol AFT |
| Restricted to children under 12 years of age. | | | |
| BISOPROLOL FUMARATE | | | |
| * Tab 2.5 mg | 1.84 | 90 | Bisoprolol Mylan |
| * Tab 5 mg | | 90 | Bisoprolol Mylan |
| * Tab 10 mg | 3.62 | 90 | Bisoprolol Mylan |
| CARVEDILOL | | | |
| * Tab 6.25 mg | | 60 | Carvedilol Sandoz |
| * Tab 12.5 mg | | 60 | Carvedilol Sandoz |
| * Tab 25 mg | | 60 | Carvedilol Sandoz |
| LABETALOL | | | |
| * Tab 100 mg | 14 50 | 100 | Trandate |
| * Tab 200 mg | | 100 | ✓ Trandate |
| * Inj 5 mg per ml, 20 ml ampoule | | 5 | |
| | (88.60) | | Trandate |
| * inj 5 mg per ml, 20 ml vial | () | 1 | |
| | (48.20) | | Alvogen S29 |
| METOPROLOL SUCCINATE | | | |
| * Tab long-acting 23.75 mg | 1.45 | 30 | Betaloc CR |
| * Tab long-acting 47.5 mg | 1.43 | 30 | Betaloc CR |
| * Tab long-acting 95 mg | 2.15 | 30 | Betaloc CR |
| * Tab long-acting 190 mg | | 30 | Betaloc CR |
| METOPROLOL TARTRATE | | | |
| Tab 50 mg | 5.66 | 100 | IPCA-Metoprolol |
| Tab 100 mg | | 60 | ✓ IPCA-Metoprolol |
| * Tab long-acting 200 mg | 23.40 | 28 | ✓ Slow-Lopresor |
| * Inj 1 mg per ml, 5 ml vial | | 5 | Metoprolol IV Mylan |
| NADOLOL | | | |
| Tab 40 mg | | 100 | Nadolol BNM \$29 |
| Tab 80 mg | | 100 | ✓ Nadolol BNM \$29 |
| · ••• ••• ···y | | 100 | |

✓ <u>Vasorex</u> ✓ Vasorex

✓ Vasorex

✓ Plendil ER✓ Felo 5 ER

✓ Felo 10 ER

90

| | Subsidy (Manufacturer's Price) \$ | | Fully Brand or ised Generic Manufacturer |
|---|---|--------|--|
| PINDOLOL – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who were endorsed accordingly. Pharmacists may annotate the presc dispensing of pindolol. | 01 1 | 0 | |
| * Tab 5 mg | | 100 | Apo-Pindolol |
| * Tab 10 mg | | 100 | Apo-Pindolol |
| * Tab 15 mg | | 100 | Apo-Pindolol |
| (Apo-Pindolol Tab 5 mg to be delisted 1 May 2022) (Apo-Pindolol Tab 10 mg to be delisted 1 May 2022) (Apo-Pindolol Tab 15 mg to be delisted 1 May 2022) | | | |
| PROPRANOLOL | | | |
| Tab 10 mg | 7.04 | 100 | Drofate |
| Tab 40 mg | | 100 | IPCA-Propranolol |
| * Cap long-acting 160 mg | | 100 | Cardinol LA |
| * Oral lig 4 mg per ml - Special Authority see SA1327 below - | | | |
| Retail pharmacy | | 500 ml | Roxane- Propranolol S29 |

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

| * | Tab 80 mg | .32.58 | 500 | ✓ Mylan |
|---|------------|--------|-----|---------|
| * | Tab 160 mg | .10.98 | 100 | 🗸 Mylan |

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

| | | Subsidy (Manufacturaria Brian) | | Fully Subsidised | |
|----------------|--|-----------------------------------|------------|---------------------|--------------------|
| | | (Manufacturer's Price) \$ | Per | Subsidised | Manufacturer |
| IFF | DIPINE | | | | |
| | Fab long-acting 10 mg | | 56 | 1 | Tensipine MR10 S29 |
| (] | Fab long-acting 20 mg | | 50 | 1 | Mylan (12 hr |
| | | | | | release) \$29 |
| | | 17.72 | 100 | ✓ | Nyefax Retard |
| ÷ 1 | Fab long-acting 30 mg | 4.78 | 14 | ✓ | Mylan Italy (24 hr |
| | | | | | release) S29 |
| | | 34.10 | 100 | ~ | Mylan (24 hr |
| | Tables a selice 00 mm | 50.04 | 400 | | release) S29 |
| ŧ | Tab long-acting 60 mg | | 100 | • | Mylan (24 hr |
| | | | | | release) S29 |
| Otl | her Calcium Channel Blockers | | | | |
| ILT | IAZEM HYDROCHLORIDE | | | | |
| (| Cap extended-release 120 mg | | 100 | | Accord S29 |
| | Cap long-acting 120 mg | | 500 | | Apo-Diltiazem CD |
| | Cap long-acting 180 mg | | 30 | | Cardizem CD |
| € (| Cap long-acting 240 mg | 9.30 | 30 | ~ | Cardizem CD |
| | HEXILINE MALEATE | | | | |
| ⊬ 7 | Гаb 100 mg | | 100 | 1 | Pexsig |
| | APAMIL HYDROCHLORIDE | | | | |
| | Гаb 40 mg | | 100 | | Isoptin |
| (] | Гаb 80 mg | 11.74 | 100 | | Isoptin |
| ÷Γ | Tab long-acting 120 mg | | 100 | | Isoptin Retard S29 |
| | | | | | Isoptin SR |
| | Tab long-acting 240 mg | | 30 | ~ | Isoptin SR |
| - | nj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO | 25.00 | 5 | 1 | Isoptin |
| - | 100 | | 5 | • | Isopun |
| Ce | ntrally-Acting Agents | | | | |
| LO | NIDINE | | | | |
| | Patch 2.5 mg, 100 mcg per day – Only on a prescription | | 4 | 1 | Mylan |
| | Patch 5 mg, 200 mcg per day – Only on a prescription | | 4 | | Mylan |
| f F | Patch 7.5 mg, 300 mcg per day – Only on a prescription | | 4 | ~ | <u>Mylan</u> |
| | NIDINE HYDROCHLORIDE | | | | |
| ÷ 1 | Гаb 25 mcg | | 112 | | Clonidine BNM |
| _ | | 36.50 | | | Clonidine Teva |
| | Tab 150 mcg | | 100 | | Catapres |
| | nj 150 mcg per ml, 1 ml ampoule | | 10 | - | Medsurge |
| | HYLDOPA | 15 10 | 100 | | Methyldere Muler |
| 7 | Гаb 250 mg | | 100 500 | | Methyldopa Mylan |
| | | 02.00 | 200 | ~ | Methyldopa Mylan |

| | Cubaidu | | Fully Drand ar |
|--|--------------------------------|----------------|---|
| | Subsidy (Manufacturer's Pri | ce) Sub | Fully Brand or osidised Generic |
| | \$ | Per | Manufacturer |
| Diuretics | | | |
| Loop Diuretics | | | |
| UMETANIDE | | | |
| ← Tab 1 mg | 4.91 | 30 | Burinex S29 S29 |
| | 16.36 | 100 | Burinex |
| ✓ Inj 500 mcg per ml, 4 ml vial | 7.95 | 5 | Burinex |
| UROSEMIDE [FRUSEMIDE] | 0.00 | 1 000 | IDCA Emicomida |
| Tab 40 mg − Up to 30 tab available on a PSO | | 1,000 50 | ✓ <u>IPCA-Frusemide</u> ✓ Urex Forte |
| • 1 db 500 mg | 89.48 | 50 | ✓ Furosemid- |
| | | | Ratiopharm S29 |
| | 169.96 | 100 | Furosemid- Ratiopharm S29 |
| • Oral lig 10 mg per ml | 11.00 | 30 ml OP | |
| Inj 10 mg per ml, 25 ml ampoule | | 50 III OF 6 | ✓ <u>Lasix</u> ✓ Lasix |
| Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a l | | 5 | ✓ Furosemide-Baxter |
| Potassium Sparing Diuretics | | | |
| MILORIDE HYDROCHLORIDE | | | |
| Oral liq 1 mg per ml | 32.10 | 25 ml OP | Biomed |
| PLERENONE - Special Authority see SA1728 below - Retail | pharmacy | | |
| Tab 25 mg | | 30 | Inspra |
| Inspra to be Principal Supply on 1 June 2022 | 25.00 | 30 | ✓ Inspra |
| Tab 50 mg Inspra to be Principal Supply on 1 June 2022 | 25.00 | 30 | |
| •SA1728 Special Authority for Subsidy | | | |
| itial application from any relevant practitioner. Approvals vali ne following criteria: | d without further re | enewal unles | ss notified for applications meeting |
| oth: | | | |
| Patient has heart failure with ejection fraction less than 40 Either: | 0%; and | | |
| 2.1 Patient is intolerant to optimal dosing of spironolac2.2 Patient has experienced a clinically significant adv | | n optimal do | sing of spironolactone. |
| IETOLAZONE | | | |
| Tab 5 mg | CBS | 1 | Metolazone S29 |
| | | 50 | Zaroxolyn S29 |
| PIRONOLACTONE | | | |
| Tab 25 mg | | 100 | ✓ Spiractin |
| Tab 100 mg | | 100 | ✓ Spiractin |
| Oral liq 5 mg per ml | | 25 ml OP | <u>Biomed</u> |
| Potassium Sparing Combination Diuretics | | | |
| MILORIDE HYDROCHLORIDE WITH FUROSEMIDE | | | |
| Tab 5 mg with furosemide 40 mg | 8.63 | 28 | Frumil |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

CARDIOVASCULAR SYSTEM

| * Tab 2.5 mg - Up to 150 tab available on a PSO | | Subsidy | | Fully | Brand or | |
|---|--|---------|---------|------------|-------------------|-----------|
| MULORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE * Tab 5 mg with hydrochlorothiazide 50 mg | | · . | | Subsidised | | |
| ** Tab 5 mg with hydrochlorothiazide 50 mg 5.00 50 ✓ Moduretic Thiazide and Related Diuretics SENDROFLUMETHIAZIDE (BENDROFLUAZIDE) * Arrow-Bendrofluazide ** Tab 2.5 mg - Up to 150 tab available on a PSO .20.00 500 ✓ Arrow-Bendrofluazide May be supplied on a PSO for reasons other than emergency. * Tab 5 mg .34.55 500 ✓ Arrow-Bendrofluazide Oral liq 50 mg per ml .27.82 25 ml OP ✓ Biomed CHLOROTHIAZIDE .25.00 ✓ Hydroton **** Tab 25 mg .390 30 ✓ Igroton ***** Tab 25 mg .3.90 30 ✓ Igroton ***** Yeroton ************************************ | AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZI | | | | manaratarar | |
| SENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO | | | 50 | ✓ | Moduretic | |
| * Tab 2.5 mg - Up to 150 tab available on a PSO | Thiazide and Related Diuretics | | | | | |
| Bendrofluazide May be supplied on a PSO for reasons other than emergency. * Tab 5 mg | BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] | | | | | |
| * Tab 5 mg 34.55 500 ✓ Arrow- Bendrofluazide CHLOROTHIAZIDE Oral liq 50 mg per ml 27.82 25 ml OP ✓ Biomed CHLORTALIDONE [CHLORTHALIDONE] 3.90 30 ✓ Igroton % (Igroton % (Igro | * Tab 2.5 mg – Up to 150 tab available on a PSO | 20.00 | 500 | 1 | | |
| Bendrofluazide CHLOROTHIAZIDE Oral liq 50 mg per ml | | | | | | |
| Oral liq 50 mg per ml 27.82 25 ml OP ✓ Biomed CHLORTALIDONE [CHLORTHALIDONE] 3.90 30 ✓ Igroton Image: Solution of the solution of t | * Tab 5 mg | | 500 | v | | |
| CHLORTALIDONE [CHLORTHALIDONE] Tab 25 mg 30 ✓ Igroton Image: Solution in the second secon | CHLOROTHIAZIDE | | | | | |
| Tab 25 mg 3.90 30 - / Igroton % (Irror) NDAPAMIDE 6.50 50 - / Hygroton * Tab 2.5 mg 10.45 90 - / Dapa-Tabs 11.61 100 - / Mylan Indapamide % Lipid-Modifying Agents - - Mylan SEZAFIBRATE 90 - / Bezalip - * Tab 200 mg 19.46 90 - / Bezalip * Tab long-acting 400 mg 21.21 30 - / Bezalip Other Lipid-Modifying Agents - - - ACIPIMOX - 21.56 30 - / Olbetam * Cap 250 mg 21.56 30 - / Olbetam - COLESTIPOL HYDROCHLORIDE 32.89 30 - / Colestid HMG COA Reductase Inhibitors (Statins) - - - ATORVASTATIN - - - - * Tab 20 mg - - - - - ATORVASTATIN - - - - - * Tab 20 mg - - - - - | | 27.82 | 25 ml C | DP 🗸 | Biomed | |
| 6.50 50 ✓ Hygroton NDAPAMIDE * Tab 2.5 mg 10.45 90 ✓ Dapa-Tabs 11.61 100 ✓ Mylan Indapamide @@ Lipid-Modifying Agents Fibrates SEZAFIBRATE * Tab 200 mg 19.46 90 ✓ Bezalip * Tab 200 mg 19.46 90 ✓ Bezalip Retard Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg 21.21 30 ✓ Olbetam ✓ Olbetam S29 @@ Resins COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g 30 ✓ Colestid HMG CoA Reductase Inhibitors (Statins) ATORVASTATIN * Tab 20 mg 9.24 500 ✓ Lorstat * Tab 40 mg 9.24 500 ✓ Lorstat ✓ * Tab 80 mg 26.54 500 ✓ Lorstat ✓ * Tab 80 mg 26.54 500 ✓ Lorstat ✓ * Tab 80 mg 21.1 28 ✓ ✓ ✓ * Tab 20 mg 21.1 28 ✓ ✓ ✓ | | 2 00 | 20 | | laraton con | |
| NDAPAMIDE ** Tab 2.5 mg 10.45 90 * Dapa-Tabs ** Tab 2.5 mg 11.61 100 * Mylan Indapamide \$29 Lipid-Modifying Agents Fibrates BEZAFIBRATE 90 * Bezalip * Tab 200 mg 19.46 90 * Bezalip * Tab 200 mg 19.46 90 * Bezalip * Tab 200 mg 19.46 90 * Bezalip * Tab 10mg-acting 400 mg 21.21 30 * Bezalip Retard Other Lipid-Modifying Agents 30 * Olbetam * Olbetam ACIPIMOX 21.56 30 * Olbetam * Olbetam * Cap 250 mg 21.56 30 * Olbetam * Olbetam COLESTIPOL HYDROCHLORIDE 32.89 30 * Colestid HMG COA Reductase Inhibitors (Statins) 30 * Colestid ATORVASTATIN * 500 * Lorstat * Tab 20 mg 9.24 500 * Lorstat * Tab 40 mg 26.54 500 * Lorstat * Tab 80 mg 26.54 500 * Lorstat | 1 ab 25 mg | | | | | |
| ** Tab 2.5 mg 10.45 90 ✓ Dapa-Tabs 11.61 100 ✓ Mylan Indapamide same Lipid-Modifying Agents Fibrates BEZAFIBRATE * Tab 200 mg 19.46 90 ✓ Bezalip * Tab 200 mg 19.46 90 ✓ Bezalip Retard Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg 21.56 30 ✓ Olbetam Colspan="2">Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg 21.56 30 ✓ Olbetam COLESTIPOL HYDROCHLORIDE 32.89 30 ✓ Colestid MG CoA Reductase Inhibitors (Statins) ATORVASTATIN * Tab 10 mg 6.16 500 ✓ Lorstat * Tab 40 mg 9.24 500 ✓ Lorstat * Tab 40 mg 26.54 500 ✓ Lorstat * Tab 80 mg 26.54 500 ✓ Lorstat * Tab 20 mg 2.11 28 ✓ Pravastatin Mylan | INDAPAMIDE | 0.00 | | | <u></u> | |
| Indapamide S20 Lipid-Modifying Agents SEZAFIBRATE * Tab long-acting 400 mg * Tab long-acting 400 mg * Tab long-acting 400 mg Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg 21.26 30 • Olbetam • Olbetam <td colspan<="" td=""><td></td><td>10.45</td><td>90</td><td>1</td><td>Dapa-Tabs</td></td> | <td></td> <td>10.45</td> <td>90</td> <td>1</td> <td>Dapa-Tabs</td> | | 10.45 | 90 | 1 | Dapa-Tabs |
| Lipid-Modifying Agents Fibrates BEZAFIBRATE * Tab long-acting 400 mg 19.46 90 • Bezalip * Tab long-acting 400 mg .21.21 30 • Bezalip Retard Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg .21.56 30 • Olbetam • Olbetam • Olbetam S29 @@ Resins COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g .32.89 30 • Colestid HMG CoA Reductase Inhibitors (Statins) ATORVASTATIN * Tab 20 mg .9.24 .500 • Lorstat * Tab 20 mg .26.54 .500 • Lorstat * Tab 20 mg .26.54 .500 • Lorstat * Tab 20 mg .26.54 .500 • Lorstat * Tab 20 mg .21.1 28 • Pravastatin Mylan | | 11.61 | 100 | 1 | • | |
| Fibrates SEZAFIBRATE * Tab long-acting 400 mg 19.46 90 | | | | | Indapamide 529 | |
| BEZAFIBRATE * Tab 200 mg 19.46 90 • Bezalip * Tab long-acting 400 mg 21.21 30 • Bezalip Retard Other Lipid-Modifying Agents ACIPIMOX 21.56 30 • Olbetam * Cap 250 mg 21.56 30 • Colestid PROCLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g 30 • Colestid HMG COA Reductase Inhibitors (Statins) * Colestid • Lorstat * Tab 10 mg 9.24 500 • Lorstat * Tab 20 mg 26.54 500 • Lorstat * Tab 80 mg 26.54 500 • Lorstat | Lipid-Modifying Agents | | | | | |
| ** Tab 200 mg 19.46 90 ✓ Bezalip ** Tab long-acting 400 mg 21.21 30 ✓ Bezalip Retard Other Lipid-Modifying Agents 30 ✓ Olbetam ACIPIMOX 21.26 30 ✓ Olbetam * Cap 250 mg 21.56 30 ✓ Olbetam * Cap 250 mg 21.56 30 ✓ Olbetam S29 529 Resins 21.56 30 ✓ Olbetam S29 529 COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g 32.89 30 ✓ Colestid HMG COA Reductase Inhibitors (Statins) 30 ✓ Lorstat ATORVASTATIN 6.16 500 ✓ Lorstat * Tab 10 mg 9.24 500 ✓ Lorstat * Tab 40 mg 14.92 500 ✓ Lorstat * Tab 80 mg 26.54 500 ✓ Lorstat * Tab 20 mg 26.54 500 ✓ Lorstat * Tab 20 mg 26.54 500 ✓ Lorstat * Tab 20 mg 21.1 28 ✓ Pravastatin Mylan | Fibrates | | | | | |
| * Tab long-acting 400 mg | BEZAFIBRATE | | | - | | |
| Other Lipid-Modifying Agents ACIPIMOX * Cap 250 mg | | | | | | |
| ACIPIMOX * Cap 250 mg | | | 50 | • | Dezanp netaru | |
| * Cap 250 mg 21.56 30 ✓ Olbetam * Olbetam S29 sze Resins COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g 30 ✓ Colestid HMG CoA Reductase Inhibitors (Statins) ATORVASTATIN 6.16 500 ✓ Lorstat * Tab 20 mg 9.24 500 ✓ Lorstat * Tab 40 mg 14.92 500 ✓ Lorstat * Tab 80 mg 26.54 500 ✓ Lorstat * Tab 20 mg 26.54 500 ✓ Lorstat * Tab 20 mg 21.1 28 ✓ Pravastatin Mylan | Other Lipid-Modifying Agents | | | | | |
| ✓ Olbetam S29 529 Resins COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g | ACIPIMOX | 01 56 | 20 | | Olbotom | |
| Resins COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g | * Cap 250 mg | 21.30 | 30 | | | |
| COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g | | | | | | |
| Grans for oral liq 5 g. 32.89 30 ✓ Colestid HMG CoA Reductase Inhibitors (Statins) ATORVASTATIN * Tab 10 mg 6.16 500 ✓ Lorstat * Tab 20 mg 9.24 500 ✓ Lorstat * Tab 40 mg 14.92 500 ✓ Lorstat * Tab 80 mg 26.54 500 ✓ Lorstat PRAVASTATIN * Tab 20 mg 211 28 ✓ Pravastatin Mylan | Resins | | | | | |
| HMG CoA Reductase Inhibitors (Statins) ATORVASTATIN * Tab 10 mg 6.16 500 ✓ Lorstat * Tab 20 mg 9.24 500 ✓ Lorstat * Tab 40 mg 14.92 500 ✓ Lorstat * Tab 80 mg 26.54 500 ✓ Lorstat PRAVASTATIN * 211 28 ✓ Pravastatin Mylan | COLESTIPOL HYDROCHLORIDE | | | | | |
| ATORVASTATIN * Tab 10 mg | Grans for oral liq 5 g | | 30 | ~ | Colestid | |
| ** Tab 10 mg 6.16 500 ✓ Lorstat ** Tab 20 mg 9.24 500 ✓ Lorstat ** Tab 40 mg 14.92 500 ✓ Lorstat ** Tab 80 mg 26.54 500 ✓ Lorstat PRAVASTATIN * Tab 20 mg 211 28 ✓ Pravastatin Mylan | HMG CoA Reductase Inhibitors (Statins) | | | | | |
| ** Tab 20 mg 9.24 500 ✓ Lorstat ** Tab 40 mg 14.92 500 ✓ Lorstat ** Tab 80 mg 26.54 500 ✓ Lorstat PRAVASTATIN ** Tab 20 mg 2.11 28 ✓ Pravastatin Mylan | ATORVASTATIN | | | | | |
| ★ Tab 40 mg 14.92 500 ✓ Lorstat ★ Tab 80 mg 26.54 500 ✓ Lorstat PRAVASTATIN 2.11 28 ✓ Pravastatin Mylan | | | | | | |
| ★ Tab 80 mg | | | | | | |
| * Tab 20 mg | | | | | | |
| | PRAVASTATIN | | | | | |
| ★ Tao 40 mg | | | | | | |
| | * Tab 40 mg | 3.01 | 28 | • | Pravastatin Mylan | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|-----|---------------------|----------------------|
| ROSUVASTATIN - Special Authority see SA2093 below - Reta | il pharmacy | | | |
| Tab 5 mg | | 30 | ✓ | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Sole Supply on 1 May 2022 | | | | |
| Tab 10 mg | 2.42 | 30 | ✓ | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Sole Supply on 1 May 2022 | | | | |
| Tab 20 mg | 3.92 | 30 | 1 | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Sole Supply on 1 May 2022 | | | | |
| Tab 40 mg | 5.28 | 30 | 1 | Rosuvastatin Viatris |
| Rosuvastatin Viatris to be Sole Supply on 1 May 2022 | | | | |

► SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

| * | Tab 10 mg | 90 | Simvastatin Mylan |
|---|---------------|----|---------------------------------------|
| | Tab 20 mg2.03 | 90 | Simvastatin Mylan |
| | Tab 40 mg | 90 | Simvastatin Mylan |
| * | Tab 80 mg7.12 | 90 | Simvastatin Mylan |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Fully Subsidised Per 🖌 | ed Generic |
|--|--|
| | |
| 30 🗸 | Ezetimibe Sandoz ng the following criteria: |
| over 5 years; a | ; and |
| | than $10 \times normal$) when use of the maximal tolerated |
| s. ed after failure high then a rep red that the LD | less potent statin should use e of statin therapy. epeat test should be .DL cholesterol is greater tha opropriate and the patient is |
| rmacy | |
| 30 ✓ 30 ✓ 30 ✓ | / Zimybe / Zimybe / Zimybe / Zimybe |
| ; | |

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| | Subsidy | | Fully Brand or |
|--|-------------------------|-------------------|--|
| | (Manufacturer's F \$ | Price) Sub Per | sidised Generic Manufacturer |
| | Ŧ | | |
| Nitrates | | | |
| GLYCERYL TRINITRATE | | | |
| * Oral pump spray, 400 mcg per dose – Up to 250 dose | | | _ |
| available on a PSO | 6.09 | 250 dose OP | |
| * Patch 25 mg, 5 mg per day | 15 73 | 30 | Spray ✔ Nitroderm TTS |
| * Patch 50 mg, 10 mg per day | | 30 | ✓ Nitroderm TTS |
| ISOSORBIDE MONONITRATE | | | |
| * Tab 20 mg | | 100 | ✓ <u>Ismo 20</u> |
| * Tab long-acting 40 mg | | 30 | Ismo 40 Retard |
| * Tab long-acting 60 mg | 9.25 | 90 | ✓ Duride |
| Sympathomimetics | | | |
| | | | |
| ADRENALINE Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSC | ∩ <u>≬</u> ∩o | 5 | ✓ Aspen Adrenaline |
| ing 1 in 1,000, 1 ini ampoule – Op to 5 ing available on a PSC | 10.76 J | 5 | Aspen Adrenaline DBL Adrenaline |
| Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a F | | 5 | ✓ Hospira |
| | 49.00 | 10 | Aspen Adrenaline |
| Vasodilators | | | |
| | | | |
| HYDRALAZINE HYDROCHLORIDE | | | |
| * Tab 25 mg – Special Authority see SA1321 below – Retail pharmaou | CRS | 1 | - Hudralazina |
| pharmacy | | 56 | Hydralazine Onelink S29 |
| | | 84 | ✓ AMDIPHARM \$29 |
| | | 100 | ✓ Onelink S29 |
| * Inj 20 mg ampoule | 25.90 | 5 | ✓ Apresoline |
| ► SA1321 Special Authority for Subsidy | | | |
| Initial application from any relevant practitioner. Approvals vali | d without further | renewal unles | s notified for applications meeting |
| the following criteria: | | | |
| Either: | | | |
| For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nit | rate, in patients | who are intoler | ant or have not responded to ACF |
| inhibitors and/or angiotensin receptor blockers. | rate, in patiente | | |
| MINOXIDIL | | | |
| ▲ Tab 10 mg | 70.00 | 100 | Loniten |
| NICORANDIL | | | |
| ▲ Tab 10 mg | | 60 | ✓ <u>Ikorel</u> |
| ▲ Tab 20 mg | | 60 | ✓ <u>Ikorel</u> |
| PAPAVERINE HYDROCHLORIDE | 057.40 | - | |
| * Inj 12 mg per ml, 10 ml ampoule | 257.12 | 5 | Hospira |
| PENTOXIFYLLINE [OXPENTIFYLLINE] | 10.00 | 50 | Trantal 400 |
| Tab 400 mg | | 50 | Trental 400 |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|--------------|---------------------|--|
| Endothelin Receptor Antagonists | | | | |
| AMBRISENTAN – Special Authority see SA1702 below – Retail Tab 5 mg Tab 10 mg | 1,550.00 | 30 30 | 1 | <u>Ambrisentan Mylan</u> Ambrisentan Mylan Mylan |
| ► SA1702 Special Authority for Subsidy Special Authority approved by the Pulmonary Arterial Hypertensin Notes: Application details may be obtained from Pharmac's web The Coordinator, PAH Panel Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac | site <u>schedule.pharma</u> | <u>c.gov</u> | t.nz/SAFor | <u>ms</u> or: |
| BOSENTAN – Special Authority see SA1991 below – Retail pha Tab 62.5 mg | | 60 | 1 | <u>Bosentan Dr</u> Reddy's |
| Tab 125 mg | 119.85 | 60 | 1 | Bosentan Dr Reddy's |

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

continued...

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

| Tab 25 mg0.85 | 4 | Vedafil |
|------------------|----|-----------------------------|
| Tab 50 mg | 4 | ✓ Vedafil |
| Tab 100 mg 10.20 | 12 | ✓ Vedafil |

➡SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) | S | Fully ubsidised | Brand or Generic |
|-----------------------------------|-----|--------------------|---------------------|
| \$ | Per | 1 | Manufacturer |

continued...

4 Either:

- 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and

4.1.2 Either:

- 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
- 4.1.2.2 Patient is peri Fontan repair; and
- 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
- 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.
- Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

| EPOPROSTENOL – Special Authority see SA1696 below – Inj 500 mcg vial | | 1 | ✓ Veletri |
|---|-----------------------|------------|------------------------------|
| Inj 1.5 mg vial | | 1 | ✓ Veletri |
| ➡SA1696 Special Authority for Subsidy | | | |
| Special Authority approved by the Pulmonary Arterial Hyper | tension Panel | | |
| Notes: Application details may be obtained from Pharmac's | website schedule.phar | mac.govt.n | z/SAForms or: |
| The Coordinator, PAH Panel | | | |
| Pharmac, PO Box 10-254, WELLINGTON | | | |
| Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pha | irmac.govt.nz | | |
| ILOPROST - Special Authority see SA1705 below - Retail | pharmacy | | |
| Nebuliser soln 10 mcg per ml, 2 ml | | 30 | Ventavis |
| ■ SA1705 Special Authority for Subsidy | | | |
| Special Authority approved by the Pulmonary Arterial Hyper | tension Panel | | |
| Notes: Application details may be obtained from Pharmac's | website schedule.phar | mac.govt.n | z/SAForms or: |
| The Coordinator, PAH Panel | | | |
| Pharmac, PO Box 10-254, WELLINGTON | | | |
| Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pha | irmac.govt.nz | | |

| | Subsidy | | Fully | Brand or |
|---|-----------------------|---------|--------|--------------|
| | (Manufacturer's Price | | idised | Generic |
| | \$ | Per | / | Manufacturer |
| Antiacne Preparations | | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacterials, | page 89 | | | |
| ADAPALENE | - | | | |
| a) Maximum of 30 g per prescription | | | | |
| b) Only on a prescription | | | | |
| Crm 0.1% | | 30 g OP | 🗸 D | lifferin |
| Gel 0.1% | | 30 g OP | 🗸 D | Vifferin |
| ISOTRETINOIN - Special Authority see SA2023 below - Retail p | harmacy | | | |
| Cap 5 mg | | 60 | ✓ 0 | Iratane |
| Cap 10 mg | | 120 | ✓Ō | Iratane |
| Cap 20 mg | | 120 | ✓Ō | Iratane |
| | | | _ | |

SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

| TRETINOIN Crm 0.5 mg per g – Maximum of 50 g per prescription15.57 | 50 g OP | ✓ <u>ReTrieve</u> | |
|---|--------------------|--|--|
| Antibacterials Topical | | | |
| For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89 | | | |
| HYDROGEN PEROXIDE * Crm 1% | 10 g OP 15 g OP | ✓ Crystaderm ✓ Crystaderm | |

| | Subsidy (Manufacturer's F | Price) Subr | Fully Brand or sidised Generic |
|---|--|---|---|
| | (Manulacturers F | Per Subs | Manufacturer |
| IUPIROCIN | | | |
| Oint 2% | 6.60 | 15 g OP | |
| | (11.50) | | Bactroban |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| ODIUM FUSIDATE [FUSIDIC ACID] | | | |
| Crm 2% | 1.59 | 5 g OP | Foban |
| a) Maximum of 5 g per prescription | | | |
| b) Only on a prescription | | | |
| c) Not in combination Oint 2% | 1 50 | 5 a OP | ✓ Foban |
| a) Maximum of 5 g per prescription | 1.59 | 5 g OP | |
| b) Only on a prescription | | | |
| c) Not in combination | | | |
| | | | |
| Crm 1% | 10.80 | 50 g OP | Flamazine |
| a) Up to 250 g available on a PSO | | | |
| b) Not in combination | | | |
| | | | |
| Antifungals Topical | | | |
| ar avatamia antifungala, rafar ta INEECTIONS, Antifungala, r | 000 | | |
| or systemic antifungals, refer to INFECTIONS, Antifungals, p | lage 96 | | |
| MOROLFINE | | | |
| | | | |
| a) Only on a prescription | | | |
| b) Not in combination | 14 03 | 5 ml OP | 🖌 MycoNail |
| b) Not in combination Nail soln 5% | 14.93 | 5 ml OP | ✓ <u>MycoNail</u> |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE | 14.93 | 5 ml OP | ✓ <u>MycoNail</u> |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription | 14.93 | 5 ml OP | ✓ <u>MycoNail</u> |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination | | | |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% | | 5 ml OP 7 ml OP | ✓ <u>MycoNail</u> ✓ Apo-Ciclopirox |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) | | | |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) ICOTRIMAZOLE | 5.72 | 7 ml OP | ✓ Apo-Ciclopirox |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE Crm 1% | 5.72 | | |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE € Crm 1% a) Only on a prescription | 5.72 | 7 ml OP | ✓ Apo-Ciclopirox |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) LOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination | 5.72 | 7 ml OP | ✓ Apo-Ciclopirox |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE € Crm 1% a) Only on a prescription b) Not in combination | 5.72 | 7 ml OP 20 g OP | ✓ Apo-Ciclopirox |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE € Crm 1% a) Only on a prescription b) Not in combination | 5.72 0.77 4.36 | 7 ml OP 20 g OP | ✓ Apo-Ciclopirox ✓ Clomazol |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE € Crm 1% a) Only on a prescription b) Not in combination € Soln 1% | 5.72 0.77 4.36 | 7 ml OP 20 g OP | ✓ Apo-Ciclopirox ✓ Clomazol |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAJOLE NITRATE | 5.72 0.77 4.36 (7.55) | 7 ml OP 20 g OP | ✓ Apo-Ciclopirox ✓ Clomazol |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) LOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% | | 7 ml OP 20 g OP | Apo-Ciclopirox Clomazol Canesten |
| b) Not in combination Nail soln 5% ICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) LOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% | 5.72 0.77 4.36 (7.55) 1.00 | 7 ml OP 20 g OP 20 ml OP | ✓ Apo-Ciclopirox ✓ Clomazol |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Conly on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination | | 7 ml OP 20 g OP 20 ml OP | Apo-Ciclopirox Clomazol Canesten |
| b) Not in combination Nail soln 5% CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CIOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination | 5.72 | 7 ml OP 20 g OP 20 ml OP 20 g OP | Apo-Ciclopirox Clomazol Canesten |
| b) Not in combination Nail soln 5% CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CLOTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription | 5.72 | 7 ml OP 20 g OP 20 ml OP | Apo-Ciclopirox Clomazol Canesten Pevaryl |
| b) Not in combination Nail soln 5% CICLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) CIOTRIMAZOLE Cmn 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination | 5.72 | 7 ml OP 20 g OP 20 ml OP 20 g OP | Apo-Ciclopirox Clomazol Canesten |
| b) Not in combination Nail soln 5% CLOPIROX OLAMINE a) Only on a prescription b) Not in combination Nail-soln 8% Apo-Ciclopirox Nail-soln 8% to be delisted 1 May 2022) COTRIMAZOLE Crm 1% a) Only on a prescription b) Not in combination Soln 1% a) Only on a prescription b) Not in combination CONAZOLE NITRATE Crm 1% a) Only on a prescription b) Not in combination | 5.72 | 7 ml OP 20 g OP 20 ml OP 20 g OP | Apo-Ciclopirox Clomazol Canesten Pevaryl |

| | Subsidy (Manufacturer's Pi \$ | rice) Subs Per | Fully Brand or sidised Generic Manufacturer |
|---|-------------------------------------|-------------------|---|
| CONAZOLE NITRATE | | | |
| Crm 2% | 0.81 | 15 g OP | ✓ Multichem |
| a) Only on a prescription | | | |
| b) Not in combination Lotn 2% | 4 36 | 30 ml OP | |
| LOUT 2 /6 | (10.03) | 50 111 01 | Daktarin |
| a) Only on a prescription | () | | |
| b) Not in combination | | | |
| Tinct 2% | | 30 ml OP | Delsterin |
| a) Only on a prescription | (12.10) | | Daktarin |
| b) Not in combination | | | |
| | | | |
| Antipruritic Preparations | | | |
| ALAMINE | | | |
| a) Only on a prescription | | | |
| b) Not in combination | | | |
| Crm, aqueous, BP | 1.08 1.26 | 100 g | ✓ Calamine-AFT ✓ healthE Calamine |
| | 1.20 | | Aqueous Cream |
| | | | BP |
| Calamine-AFT to be Principal Supply on 1 May 2022 | | | |
| ealthE Calamine Aqueous Cream BP Crm, aqueous, BP to be | delisted 1 May 20 | 022) | |
| ROTAMITON | | | |
| a) Only on a prescription | | | |
| b) Not in combination Crm 10% | 3 29 | 20 g OP | ✓ Itch-Soothe |
| ENTHOL – Only in combination | 0.20 | 20 9 01 | |
| 1) Only in combination with a dermatological base or prop | prietary Topical Co | orticosteriod – | Plain |
| 2) With or without other dermatological galenicals. | notary ropida O | | |
| · | | | |
| Crystals | | 25 g | ✓ MidWest |
| | 29.60 | 100 g | MidWest |
| Corticosteroids Topical | | | |
| | | | |

Corticosteroids - Plain

| BETAMETHASONE DIPROPIONATE | | | |
|-------------------------------------|-------|----------|-----------------------------------|
| Crm 0.05% | 2.96 | 15 g OP | Diprosone |
| | 36.00 | 50 g OP | Diprosone |
| Oint 0.05% | 2.96 | 15 g OP | Diprosone |
| | 36.00 | 50 g OP | Diprosone |
| Oint 0.05% in propylene glycol base | 4.33 | 30 g OP | Diprosone OV |
| BETAMETHASONE VALERATE | | | |
| * Crm 0.1% | 4.53 | 50 g OP | Beta Cream |
| * Oint 0.1% | 5.84 | 50 g OP | Beta Ointment |
| * Lotn 0.1% | 25.00 | 50 ml OP | Betnovate |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully Brand or |
|--|-------------------------|--------------------|--|
| | (Manufacturer's F \$ | Price) Subs Per | sidised Generic Manufacturer |
| | ψ | 1 61 | |
| | 0.10 | 00 × 0D | |
| Crm 0.05% | | 30 g OP | <u>Dermol</u> |
| Oint 0.05% | 2.12 | 30 g OP | ✓ <u>Dermol</u> |
| _OBETASONE BUTYRATE | | | |
| Crm 0.05% | | 30 g OP | _ |
| | (10.00) | | Eumovate |
| YDROCORTISONE | | | |
| Crm 1% – Only on a prescription | 3.70 | 100 g OP | Hydrocortisone |
| | | | <u>(PSM)</u> |
| | 17.15 | 500 g | Hydrocortisone |
| | | | <u>(PSM)</u> |
| Powder – Only in combination | | 25 g | 🗸 ABM |
| Up to 5% in a dermatological base (not proprietary Topi | cal Corticosterio | d – Plain) with c | or without other dermatologica |
| galenicals | | | |
| YDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN | | | |
| Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only | on | | |
| a prescription | | 250 ml | DP Lotn HC |
| YDROCORTISONE BUTYRATE | | | |
| Lipocream 0.1% | | 100 g OP | Locoid Lipocream |
| Oint 0.1% | | 100 g OP | ✓ Locoid |
| Milky emul 0.1% | | 100 ml OP | ✓ Locoid Crelo |
| ETHYLPREDNISOLONE ACEPONATE | | | |
| Crm 0.1% | 4 46 | 15 g OP | Advantan |
| Oint 0.1% | | 15 g OP | ✓ Advantan |
| OMETASONE FUROATE | | .e g e. | |
| Crm 0.1% | 1.05 | 15 g OP | Elocon Alcohol Free |
| 0111 0.1 /6 | 3.10 | 50 g OP | <u>Elocon Alcohol Free</u> |
| Oint 0.1% | •••• | 15 g OP | ✓ Elocon |
| | 2.90 | 50 g OP | ✓ Elocon |
| Lotn 0.1% | | 30 ml OP | ✓ Elocon |
| RIAMCINOLONE ACETONIDE | | | |
| Crm 0.02% | 6 20 | 100 g OP | Aristocort |
| Oint 0.02% | | 100 g OP | ✓ Aristocort |
| Oint 0.02 /8 | 0.05 | 100 g Oi | Alistocon |
| Corticosteroids - Combination | | | |
| | | | |
| ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FL | • | 15 ~ 00 | |
| Crm 0.1% with sodium fusidate (fusidic acid) 2% | | 15 g OP | Eucloart |
| | (10.45) | | Fucicort |
| a) Maximum of 15 g per prescription b) Only on a prescription | | | |
| b) Only on a prescription | | | |
| DROCORTISONE WITH MICONAZOLE - Only on a prescri | | | |
| Crm 1% with miconazole nitrate 2% | 1.89 | 15 g OP | ✓ Micreme H |
| YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - 0 | Only on a prescrip | otion | |
| Crm 1% with natamycin 1% and neomycin sulphate 0.5% | , , , | 15 g OP | Pimafucort |
| Oint 1% with natamycin 1% and neomycin sulphate 0.5% | | 15 g OP | Pimafucort |
| Oline 1 /8 which hatamychin 1 /8 and heomychin Sulphate 0.5 /8 | | | |

| | Subsidy | | Fully Brand or |
|--|-----------------|--------------|---|
| | (Manufacturer's | Price) Subsi | |
| | \$ | Per | Manufacturer |
| TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC | IN AND NYSTA | TIN | |
| Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m | • | | |
| and gramicidin 250 mcg per g - Only on a prescription . | | 15 g OP | |
| | (9.28) | | Viaderm KC |
| Barrier Creams and Emollients | | | |
| | | | |
| Barrier Creams | | | |
| DIMETHICONE | | | |
| * Crm 5% pump bottle | 4.48 | 500 ml OP | ✓ healthE |
| | | | Dimethicone 5% |
| * Crm 10% pump bottle | 4.52 | 500 ml OP | ✓ healthE |
| | | | Dimethicone 10% |
| ZINC AND CASTOR OIL | 4.05 | 500 - | |
| * Oint | 4.65 | 500 g | Boucher |
| Emollients | | | |
| AQUEOUS CREAM | | | |
| * Crm | 1.73 | 500 g | Boucher |
| | | | GEM Aqueous |
| | | | Cream |
| GEM Aqueous Cream to be Principal Supply on 1 July 2 | .022 | | |
| (Boucher Crm to be delisted 1 August 2022) | | | |
| CETOMACROGOL | 4.00 | 500 | (o · · · · · · · · · · · · · · · · · · |
| * Crm BP | | 500 g | Cetomacrogol-AFT healthE |
| Cetomacrogol-AFT to be Principal Supply on 1 May 202 | 2.48 | | |
| (healthE Crm BP to be delisted 1 May 2022) | 2 | | |
| CETOMACROGOL WITH GLYCEROL | | | |
| Crm 90% with glycerol 10% | 2.35 | 500 ml OP | Boucher |
| | 2.00 | 000 01 | ✓ Pharmacy Health |
| | | | Sorbolene with |
| | | | Glycerin |
| | 3.10 | 1,000 ml OP | Boucher |
| EMULSIFYING OINTMENT | | | |
| * Oint BP | 3.40 | 500 g | Emulsifying |
| | | | Ointment ADE |
| OIL IN WATER EMULSION | | | |
| * Crm | | 500 g | Fatty Cream AFT |
| | 2.19 | | ✓ O/W Fatty Emulsion |
| (O/M Fatty Emulsion Cream Crm to be delicted 1 September 200 | 22) | | Cream |
| (O/W Fatty Emulsion Cream Crm to be delisted 1 September 202 | <i>()</i> | | |
| PARAFFIN | 5.25 | 500 ml OP | ✓ healthE |
| Oint liquid paraffin 50% with white soft paraffin 50% | | SUU III UP | |
| UREA | 1 07 | 100 ~ 00 | ✓ healthE Urea Cream |
| * Crm 10% | 1.3/ | 100 g OP | nealthe orea oream |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully Brand or |
|---|--------------------|---------------------|------------------------------------|
| | (Manufacturer's F | | idised Generic |
| | \$ | Per | Manufacturer |
| OOL FAT WITH MINERAL OIL – Only on a prescription | | | |
| Even hydrous 3% with mineral oil | | 1,000 ml | |
| | (11.95) | | DP Lotion |
| | 1.40 | 250 ml OP | |
| | (4.53) | | DP Lotion |
| | 5.60 | 1,000 ml | |
| | (20.53) | | Alpha-Keri Lotion |
| | (23.91) | | BK Lotion |
| | 1.40 | 250 ml OP | |
| | (7.73) | | BK Lotion |
| Other Dermatological Bases | | | |
| ARAFFIN | | | |
| White soft – Only in combination | 4.99 | 450 g | ✓ healthE |
| | 19.99 | 2,500 g | ✓ healthE |
| Only in combination with a dermatological galenical or | as a diluent for a | | |
| | | | |
| Minor Skin Infections | | | |
| OVIDONE IODINE | | | |
| Oint 10% | 7.40 | 65 g OP | Betadine |
| Maximum of 130 g per prescription | | | |
| b) Only on a prescription | | | |
| Antiseptic Solution 10% | 4.15 | 100 ml | ✓ <u>Riodine</u> |
| Antiseptic soln 10% | 3.83 | 15 ml | Riodine |
| | 5.40 | 500 ml | Riodine |
| Skin preparation, povidone iodine 10% with 30% alcohol | 1.63 | 100 ml | |
| | (3.48) | | Betadine Skin Prep |
| Skin preparation, povidone iodine 10% with 70% alcohol | 1.63 | 100 ml | |
| | (7.78) | | Pfizer |
| Parasiticidal Preparations | | | |
| IMETHICONE | | | |
| E Lotn 4% | 4.98 | 200 ml OP | ✓ healthE |
| | | 200 01 | Dimethicone 4% |
| ERMECTIN – Special Authority see SA1225 below – Retail p | harmacy | | Louon |
| Tab 3 mg – Up to 100 tab available on a PSO | | 4 | Stromectol |
| 1) PSO for institutional use only. Must be endorsed | | the institution for | or which the PSO is required an |
| a valid Special Authority for patient of that institut | | anial Authority of | and matterns of the streatment |
| Ivermectin available on BSO provided the BSO in | | , | |
| For the purposes of subsidy of ivermectin, institut | ion means age re | lated residentia | I care tacilities, disability care |
| facilities or prisons. | | | |

► SA1225 Special Authority for Subsidy Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...

| | osidy | Fully | Brand or |
|--|--------------------|--------|--------------|
| | urer's Price) Subs | idised | Generic |
| `````````````````````````````````````` | \$ Per | 1 | Manufacturer |

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

| Crm 5% | 30 g OP | Lyderm |
|---------|----------|--------------------------------------|
| Lotn 5% | 30 ml OP | <u>A-Scabies</u> |

Psoriasis and Eczema Preparations

| ACITRETIN – Special Authority see SA2024 below – Retail pharmacy | | |
|--|----|--------------------------------|
| Cap 10 mg 17.86 | 60 | Novatretin |
| Cap 25 mg | 60 | ✓ Novatretin |

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

| Foam spray 500 mcg with calcipotriol 50 mcg per g Gel 500 mcg with calcipotriol 50 mcg per g Oint 500 mcg with calcipotriol 50 mcg per g | | 60 g OP 60 g OP 30 g OP | ✓ Enstilar ✓ <u>Daivobet</u> ✓ <u>Daivobet</u> |
|--|-------|-------------------------------|--|
| CALCIPOTRIOL Oint 50 mcg per g | 40.00 | 120 g OP | Daivonex |
| COAL TAR Soln BP – Only in combination | | 200 ml | ✓ Midwest |

1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

| | Subsidy (Manufacturer's Pr | riaa) Cuba | Fully Brand or |
|---|-------------------------------|------------------|----------------------------------|
| | (Manulacturer's Pr | Per Subs | idised Generic Manufacturer |
| | * | | |
| COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SUL | | | |
| Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% ar allantoin crm 2.5% | | 75 g OP | |
| | (8.00) | 75 y OF | Egopsoryl TA |
| | 3.43 | 30 g OP | Egopsory |
| | (4.35) | 00 9 01 | Egopsoryl TA |
| COAL TAR WITH SALICYLIC ACID AND SULPHUR | () | | |
| Soln 12% with salicylic acid 2% and sulphur 4% oint | 4 97 | 25 g OP | ✓ Coco-Scalp |
| | 7.95 | 40 g OP | ✓ Coco-Scalp |
| PIMECROLIMUS - Special Authority see SA1970 below - Reta | | 10 9 01 | e eee eeup |
| a) Maximum of 15 g per prescription | an phannacy | | |
| b) Note: a maximum of 15 g per prescription and no more | than one prescript | tion per 12 wee | ake |
| Cream 1% | | 15 g OP | ✓ Elidel |
| SA1970 Special Authority for Subsidy | 20.00 | le g el | |
| Initial application only from a dermatologist, paediatrician, oph | thalmologist or any | v relevant prac | titioner on the recommendation |
| of a dermatologist, paediatrician or ophthalmologist. Approvals | | | |
| meeting the following criteria: | | | |
| Both: | | | |
| 1 Patient has atopic dermatitis on the eyelid; and | | | |
| 2 Patient has at least one of the following contraindications | to topical corticos | steroids: perio | rificial dermatitis, rosacea, |
| documented epidermal atrophy, documented allergy to to | | | |
| pressure. | | | |
| PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORE | ESCEIN – Only on | n a prescriptior | 1 |
| * Soln 2.3% with trolamine laurilsulfate and fluorescein sodiur | | 500 ml | ✓ Pinetarsol |
| SALICYLIC ACID | | | |
| Powder – Only in combination | 18 88 | 250 g | ✓ Midwest |
| | 10.00 | 200 g | ✓ PSM |
| 1) Only in combination with a dermatological base or | r proprietary Topic | al Corticostero | id – Plain or collodion flexible |
| With or without other dermatological galenicals. | propriotary ropio | | |
| , | | | |
| (PSM Powder to be delisted 1 May 2022) | | | |
| SULPHUR | | | |
| Precipitated – Only in combination | 6.35 | 100 g | ✓ Midwest |
| 1) Only in combination with a dermatological base of | | 0 | id – Plain |
| 2) With or without other dermatological galenicals. | propriotal j ropio | | |
| _, | | | |
| TACROLIMUS | | | |
| Oint 0.1% – Special Authority see SA2074 below – Retail | | | |
| pharmacy | 33.00 | 30 g OP | Zematop |
| a) Maximum of 30 g per prescription | | 00 9 01 | <u> Zematop</u> |
| b) Note: a maximum of 30 g per prescription and no r | ore than one pres | cription per 12 | weeks |
| SA2074 Special Authority for Subsidy | | | |
| Initial application only from a dermatologist, paediatrician or ar | w relevant practitic | oner on the rec | commendation of a dermatologist |
| paediatrician, . Approvals valid without further renewal unless n | | | |
| Both: | eou ioi uppiloui | | ie ielienning enterlai |
| 1 Patient has atopic dermatitis on the face; and | | | |
| 2 Patient has at least one of the following contraindications | to topical corticos | steroids: perio | rificial dermatitis, rosacea. |
| documented epidermal atrophy or documented allergy to | | | |
| | | | |

| | Subsidy (Manufacturer's F \$ | Price) Subs Per | Fully Brand or idised Generic ✓ Manufacturer |
|---|------------------------------------|--------------------|---|
| Scalp Preparations | | | |
| BETAMETHASONE VALERATE | | | |
| ₭ Scalp app 0.1% | 9.84 | 100 ml OP | Beta Scalp |
| CLOBETASOL PROPIONATE ₭ Scalp app 0.05% | 5.69 | 30 ml OP | Dermol |
| IYDROCORTISONE BUTYRATE | | | |
| Scalp lotn 0.1% | 6.57 | 100 ml OP | ✓ Locoid |
| ETOCONAZOLE | | | |
| Shampoo 2% | 3.23 | 100 ml OP | ✓ <u>Sebizole</u> ✓ Sebizole |
| a) Maximum of 100 ml per prescription | | | |
| b) Only on a prescription | | | |
| Sunscreens | | | |
| | | | |
| UNSCREENS, PROPRIETARY – Subsidy by endorsement Only if prescribed for a patient with severe photosensitivity | recordant to a de | afined clinical co | ndition and the prescription is |
| endorsed accordingly. | | | |
| Lotn, | 5.10 | 200 g OP | ✓ <u>Marine Blue Lotion</u> |
| | | | <u>SPF 50+</u> |
| Wart Preparations | | | |
| or salicylic acid preparations refer to PSORIASIS AND ECZE | MA PREPARATIC | NS page 68 | |
| | | nie, page ee | |
| MIQUIMOD | | nie, page co | |
| Crm 5%, 250 mg sachet | 21.72 | 24 | ✓ Perrigo |
| Crm 5%, 250 mg sachet ODOPHYLLOTOXIN | | 24 | Ū |
| Crm 5%, 250 mg sachet ODOPHYLLOTOXIN Soln 0.5% | | | ✓ Perrigo✓ Condyline |
| Crm 5%, 250 mg sachet ODOPHYLLOTOXIN | | 24 | Ū |
| Crm 5%, 250 mg sachet ODOPHYLLOTOXIN Soln 0.5% a) Maximum of 3.5 ml per prescription | | 24 | Ū |
| Crm 5%, 250 mg sachet ODOPHYLLOTOXIN Soln 0.5% a) Maximum of 3.5 ml per prescription b) Only on a prescription Other Skin Preparations | | 24 | Ū |
| Crm 5%, 250 mg sachet PODOPHYLLOTOXIN Soln 0.5% a) Maximum of 3.5 ml per prescription b) Only on a prescription | | 24 | Ū |

GENITO-URINARY SYSTEM

| Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per 🖌 Manufacturer | |
|--|--|
|--|--|

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----------|--|-------------------------------------|
| Contraceptives - Non-hormonal | | | | |
| Condoms | | | | |
| ONDOMS | | | | |
| 49 mm – Up to 144 dev available on a PSO | | 144 | √ | Moments |
| 53 mm | | 10 | | Moments |
| | 11.64 | 144 | ✓] | Moments |
| a) Maximum of 60 dev per prescription | | | | |
| b) Up to 60 dev available on a PSO | | | | |
| 53 mm, 0.05 mm thickness | 0.95 | 10 | 🗸 I | Moments |
| | 11.42 | 144 | ✓] | Moments |
| a) Up to 60 dev available on a PSO | | | | |
| b) Maximum of 60 dev per prescription | | | | |
| 53 mm, chocolate, brown | | 10 | - | Moments |
| | 11.64 | 144 | ✓ [| Moments |
| a) Up to 60 dev available on a PSO | | | | |
| b) Maximum of 60 dev per prescription | | | - | - |
| 53 mm, strawberry, red | | 10 | - | Moments |
| | 11.64 | 144 | ✓ [| Moments |
| a) Up to 60 dev available on a PSO | | | | |
| b) Maximum of 60 dev per prescription 56 mm. | 0.07 | | | |
| 56 mm | | 10 | - | Moments |
| | 11.64 | 144 | • [| Moments |
| a) Maximum of 60 dev per prescription | | | | |
| b) Up to 60 dev available on a PSO 56 mm, 0.05 mm thickness. | 1.00 | 10 | | Cald Kulaht |
| 56 mm, 0.05 mm thickness | 1.30 15.57 | 12 144 | | <u>Gold Knight</u> Gold Knight |
| a) Un to 60 day available on a DCO | 15.57 | 144 | • | |
| a) Up to 60 dev available on a PSO | | | | |
| b) Maximum of 60 dev per prescription 56 mm, 0.05mm thickness (bulk pack) | 1/ 61 | 144 | 1 | Gold Knight |
| a) Maximum of 60 dev per prescription | | 144 | • • | |
| b) Up to 60 dev available on a PSO | | | | |
| 56 mm, 0.08 mm thickness | 0.97 | 10 | ~ 1 | Moments |
| | 0.97 11.64 | 144 | - | Moments |
| a) Up to 60 dev available on a PSO | TINT | 1 7 7 | • ! | |
| b) Maximum of 60 dev per prescription | | | | |
| 56 mm, 0.08 mm thickness, red | 0.97 | 10 | 🗸 I | Moments |
| | 11.64 | 144 | - | Noments |
| a) Up to 60 dev available on a PSO | | | = | |
| b) Maximum of 60 dev per prescription | | | | |
| 56 mm, chocolate | 1.30 | 12 | ✓ (| Gold Knight |
| | 15.57 | 144 | | Gold Knight |
| a) Up to 60 dev available on a PSO | | | - | |
| b) Maximum of 60 dev per prescription | | | | |
| 56 mm, strawberry | 1.30 | 12 | 1 | Gold Knight |
| - | 15.57 | 144 | | Gold Knight |
| a) Up to 60 dev available on a PSO | | | - | |
| b) Maximum of 60 dev per prescription | | | | |
| 60 mm | 1.42 | 12 | v | Gold Knight XL |
| | 14.87 | 144 | Image: A second s | Shield XL |
| | 17.02 | | ✓ (| Gold Knight XL |

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply
GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|--|-------------------------------------|
| 60 mm (bulk pack) a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO | 14.87 | 144 | ✓ <u>(</u> | Gold Knight XL |
| Contraceptive Devices | | | | |
| NTRA-UTERINE DEVICE a) Up to 40 dev available on a PSO b) Only on a PSO | | | | |
| # IUD 29.1 mm length × 23.2 mm width | | 1 | Image: A second s | Choice TT380 Short |
| # IUD 33.6 mm length × 29.9 mm width | | 1 | 1 | Choice TT380 Standard |
| IUD 35.5 mm length × 19.6 mm width | | 1 | √ | Choice Load 375 |

Contraceptives - Hormonal

Combined Oral Contraceptives

➡SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 Either:

1.1 Patient is on a Social Welfare benefit; or

1.2 Patient has an income no greater than the benefit; and

2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

84

✓ Mercilon 28

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab – Up to 84 tab available on a PSO......10.00

GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|---|---|-----|---------------------|----------------------------------|
| ETHINYLOESTRADIOL WITH LEVONORGESTREL | | | | |
| * Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets · | - | | | |
| Up to 112 tab available on a PSO | 2.18 | 84 | ✓ | Microgynon 20 ED |
| | 6.45 | 112 | ✓ | Femme-Tab ED |
| * Tab 30 mcg with levonorgestrel 150 mcg | 6.62 | 63 | | |
| | (16.50) | | | Microgynon 30 |
| a) Higher subsidy of \$15.00 per 63 tab with Special Auth b) Up to 63 tab available on a PSO Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets Up to 112 tab available on a PSO | - | 84 | · · · | age Levlen ED Femme-Tab ED |
| ETHINYLOESTRADIOL WITH NORETHISTERONE | 0.10 | | | |
| Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO | 6.95 | 84 | 1 | Brevinor 1/28 |
| Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – U to 84 tab available on a PSO | | 84 | 1 | Norimin |
| Progestogen-only Contraceptives | | | | |

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

| * Tab 30 mcg – Up to 84 tab available on a PSO | | 84 | ✓ <u>Microlut</u> |
|---|--------|-----|-------------------|
| | 22.00 | 112 | ✓ Microlut |
| Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO | 106.92 | 1 | ✓ Jadelle |
| MEDROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSC |)7.98 | 1 | ✓ Depo-Provera |
| NORETHISTERONE Tab 350 mcg – Up to 84 tab available on a PSO | 12.25 | 84 | ✓ Noriday 28 |

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer \$ Emergency Contraceptives LEVONORGESTREL 1 Postinor-1 a) Maximum of 2 tab per prescription b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Antiandrogen Oral Contraceptives Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows: • \$5.00 prescription charge (patient co-payment) will apply. • prescription may be written for up to six months supply. Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply. CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL * Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO 4.98 168 Ginet Gynaecological Anti-infectives ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator 8.43 100 g OP Aci-Jel (24.00)CLOTRIMAZOLE * Vaginal crm 1% with applicators......2.50 Clomazol 35 a OP 20 g OP Clomazol MICONAZOLE NITRATE 40 g OP Micreme NYSTATIN Vaginal crm 100,000 u per 5 g with applicator(s)4.00 Nilstat 75 g OP **Myometrial and Vaginal Hormone Preparations** ERGOMETRINE MALEATE Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a PSO......160.00 5 ✓ DBL Ergometrine OFSTRIO 15 g OP Ovestin Ovestin 15 OXYTOCIN - Up to 5 inj available on a PSO 5 Oxvtocin BNM Inj 10 iu per ml, 1 ml ampoule4.98 5 Oxytocin BNM OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj available on a PSO 5 Syntometrine

GENITO-URINARY SYSTEM

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

GENITO-URINARY SYSTEM

| | Subsidy (Manufacturer's P \$ | Price) Sub Per | Fully Brand or sidised Generic ✓ Manufacturer |
|---|--|-------------------|---|
| Pregnancy Tests - hCG Urine | | - | |
| REGNANCY TESTS - HCG URINE | | | |
| a) Up to 200 test available on a PSO | | | |
| b) Only on a PSO | | | |
| Cassette | 12.00 | 40 test OP | ✓ David One Step |
| | | | Cassette Pregnancy Test |
| | | | Smith BioMed Rapid |
| | | | Pregnancy Test |
| Urinary Agents | | | |
| or urinary tract Infections refer to INFECTIONS, Antibacter | als, page 107 | | |
| 5-Alpha Reductase Inhibitors | | | |
| INASTERIDE – Special Authority see SA0928 below – Re | tail pharmacy | | |
| Tab 5 mg | | 100 | ✓ <u>Ricit</u> |
| SA0928 Special Authority for Subsidy | valid without further | | a natified for annliastions mosting |
| itial application from any relevant practitioner. Approvals e following criteria: | s valid without iurther | renewal unles | s noulled for applications meeting |
| oth: | | | |
| 1 Patient has symptomatic benign prostatic hyperplasia | a; and | | |
| 2 Either: 2.1 The patient is intolerant of non-selective alpha | blockers or these ar | o contraindica | ted: or |
| 2.2 Symptoms are not adequately controlled with | | | ieu, oi |
| lote: Patients with enlarged prostates are the appropriate of | andidates for therapy | / with finasteri | de. |
| Alpha-1A Adrenoreceptor Blockers | | | |
| AMSULOSIN HYDROCHLORIDE - Special Authority see | SA1032 below – Reta | ail pharmacy | |
| Cap 400 mcg | 17.73 | 100 | <u>Tamsulosin-Rex</u> |
| »SA1032 Special Authority for Subsidy | and the state of t | | and the state of the |
| itial application from any relevant practitioner. Approvals e following criteria: | valid without further | renewal unles | s notified for applications meeting |
| oth: | | | |
| 1 Patient has symptomatic benign prostatic hyperplasia | | | |
| 2 The patient is intolerant of non-selective alpha blocke | ers or these are contra | aindicated. | |
| Other Urinary Agents | | | |
| XYBUTYNIN – Subsidy by endorsement | | | |
| Subsidy by endorsement – Subsidised for patients who | were taking oxybutyn | in prior to 1 Ju | ine 2021 and the prescription is |
| endorsed accordingly. Pharmacists may annotate the p dispensing of oxybutynin. | rescription as endors | eu where ther | e exists a record of prior |
| Tab 5 mg | 5.42 | 100 | Alchemy |
| | | | Oxybutynin S29 |
| Crallia 5 ma por 5 ml | 11.70 | 500 472 ml | Apo-Oxybutynin Apo-Oxybutynin |
| ← Oral liq 5 mg per 5 ml Apo-Oxybutynin Tab 5 mg to be delisted 1 May 2022) | 00.40 | 473 ml | Apo-Oxybutynin |
| Apo-Oxybutynin Oral liq 5 mg per 5 ml to be delisted 1 May | 2022) | | |
| fully subsidized | 000 11- | | aunaliad under Ca-ti 00 |
| fully subsidised | -229 UNADD | novea medicine | supplied under Section 29 |

| 1 The patient has recurrent calcium oxalate urolithiasis; and 2 The patient has had more than two renal calculi in the two years prior to the application. enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is meliting from the treatment. DDIUM CITRO-TARTRATE Grans eff 4 g sachets DIFENACIN SUCCINATE Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Data 5 mg 2.05 30 ✓ Solifenacin Mylan Detection of Substances in Urine 3.72 30 ✓ Solifenacin Mylan RTHO-TOLIDINE 7.50 50 test OP 68.25) Hemastix ETRABROMOPHENOL Blue diagnostic strips 7.02 100 test OP (13.92) Albustix | | | | | |
|--|--|----------------------|------------------|---------------------------------|------------------|
| § Per ✓ Manufacturer OTASSIUM CITRATE Oral lig 3 mmol per ml – Special Authority see SA1083 below – Retail pharmacy | | | | | |
| DTASSIUM CITRATE Oral lig 3 mmol per ml - Special Authority see SA1083 below - Retail pharmacy | | · · · | | | uror |
| Oral liq 3 mmol per ml - Special Authority see SA1083 below - 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed SPA1083 Special Authority for Subsidy 31.80 200 ml OP ✓ Biomed Stationary relevant practitioner. Approvals valid for 12 months for application. means appropriate and the patient is inefitting from the treatment. DDUM CITRO-TARTRATE Grans eff 4 g sachets 2.22 28 ✓ Ural DLIFENACIN SUCCINATE 2.05 30 ✓ Solifenacin Mylan Detection of Substances in Urine (8.25) Hemastix RTHO-TOLIDINE (8.25) Hemastix Compound diagnostic stricks 7.02 <td></td> <td>φ</td> <td>rei</td> <td>• Iviariulaci</td> <td>ulei</td> | | φ | rei | • Iviariulaci | ulei |
| Retail pharmacy | | | | | |
| SA1083 Special Authority for Subsidy titial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: 1 The patient has recurrent calcium oxalate urolithiasis; and 2 The patient has no ore than two renal calculi in the two years prior to the application. enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is newal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is newal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is DDUIFENCTORTO-TARTRATE 2.05 30 ✓ Solifenacin Mylan Tab 50 mg | | | | | |
| tital application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: 1 The patient has recurrent calcium oxalate urolithiasis; and 2 The patient has had more than two renal calculi in the two years prior to the application. nerwal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is inefitting from the treatment. DDIUM CITRO-TARTRATE 2.22 28 ✓ Ural OLIFENACIN SUCCINATE 2.05 30 ✓ Solifenacin Mylan Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Tab 10 mg 3.72 30 ✓ Solifenacin Mylan Compound diagnostic sticks 7.50 50 test OP (8.25) Compound diagnostic strips 7.02 100 test OP (13.92) Albustix Obstetric Preparations Albustix Obstetric Preparations 60.00 1 ✓ Mifegyne Tab 200 mg 60.00 1 ✓ Mifegyne | | 31.80 | 200 ml OP | Biomed | |
| attraction attraction attraction attraction bit: attraction attraction attraction bit: attractin | SA1083 Special Authority for Subsidy | | | | |
| 1 The patient has recurrent calcium oxalate urolithiasis; and 2 The patient has had more than two renal calculi in the two years prior to the application. enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is inefitting from the treatment. DDIUR CITRO-TARTRATE Grans eff 4 g sachets Tab 5 mg 2.22 28 ✓ Ural DUIFENACIN SUCCINATE Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Tab 10 mg 3.72 30 ✓ Solifenacin Mylan Octection of Substances in Urine 8.205 Hemastix RTHO-TOLIDINE 7.50 50 test OP (8.25) Compound diagnostic sticks 7.02 100 test OP (13.92) Albustix Dotstetric Preparations 7.02 100 test OP 1 Mifegyne Tab 200 mg 60.00 1 ✓ Mifegyne 180.00 3 ✓ Mifegyne | nitial application from any relevant practitioner. Approvals v | alid for 12 months | for applications | meeting the follo | wing criteria: |
| 2 The patient has had more than two renal calculi in the two years prior to the application. enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is inefitting from the treatment. DDIUM CITRO-TARTRATE Grans eff 4 g sachets DLIFENACIN SUCCINATE Tab 5 mg Tab 5 mg Quire Objectetion of Substances in Urine RTHO-TOLIDINE Compound diagnostic sticks To 200 (8.25) Hemastix Objectetic Preparations Antiprogesterones IFEPRISTONE Tab 200 mg Tab 200 mg 180.00 3 | Both: | | | | |
| enewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is unefitting from the treatment. DDIUM CITRO-TARTRATE Grans eff 4 g sachets 2.22 28 ✓ Ural DLIFENACIN SUCCINATE Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Tab 10 mg 3.72 30 ✓ Solifenacin Mylan Detection of Substances in Urine 3.72 30 ✓ Solifenacin Mylan Detection of Substances in Urine (8.25) Hemastix ETRABROMOPHENOL 8lue diagnostic sticks 7.02 100 test OP Blue diagnostic strips .7.02 100 test OP Albustix Dbstetric Preparations | 1 The patient has recurrent calcium oxalate urolithiasis; a | ind | | | |
| Interfitting from the treatment. Interfitting from the treatment. DDIUM CITRO-TARTRATE Grans eff 4 g sachets Grans eff 4 g sachets 2.22 Tab 5 mg 2.05 Tab 5 mg 2.05 Tab 10 mg 3.72 Octection of Substances in Urine RTHO-TOLIDINE Compound diagnostic sticks 7.50 (8.25) Hemastix ETRABROMOPHENOL Blue diagnostic strips 7.02 (13.92) Albustix | 2 The patient has had more than two renal calculi in the t | wo years prior to th | ne application. | | |
| DDIUM CITRO-TARTRATE Grans eff 4 g sachets 2.22 28 ✓ Ural DLIFENACIN SUCCINATE 2.05 30 ✓ Solifenacin Mylan Tab 5 mg 3.72 30 ✓ Solifenacin Mylan Detection of Substances in Urine 3.72 30 ✓ Solifenacin Mylan Detection of Substances in Urine (8.25) Hemastix RTHO-TOLIDINE (8.25) Hemastix Compound diagnostic sticks 7.02 100 test OP Blue diagnostic strips 7.02 100 test OP (13.92) Albustix Destetric Preparations Ibustix Destervice Preparations 60.00 1 ✓ Mifegyne Tab 200 mg 60.00 1 ✓ Mifegyne | | years where the tre | eatment remains | s appropriate and | I the patient is |
| Grans eff 4 g sachets 2.22 28 ✓ Ural DLIFENACIN SUCCINATE 30 ✓ Solifenacin Mylan Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Tab 10 mg 3.72 30 ✓ Solifenacin Mylan Octection of Substances in Urine 3.72 30 ✓ Solifenacin Mylan Octection of Substances in Urine 8 8 9 RTHO-TOLIDINE 7.50 50 test OP 9 Compound diagnostic sticks 7.02 100 test OP 100 test OP Blue diagnostic strips 7.02 100 test OP Albustix Obstetric Preparations 7.02 100 test OP 100 test OP IFEPRISTONE 60.00 1 ✓ Mifegyne Tab 200 mg 60.00 1 ✓ Mifegyne | penefitting from the treatment. | | | | |
| DLIFENACIO SUCCINATE Tab 5 mg .2.05 30 ✓ Solifenacin Mylan Tab 10 mg .3.72 30 ✓ Solifenacin Mylan Oetection of Substances in Urine | SODIUM CITRO-TARTRATE | | | | |
| Tab 5 mg 2.05 30 ✓ Solifenacin Mylan Tab 10 mg 3.72 30 ✓ Solifenacin Mylan Detection of Substances in Urine 3.72 30 ✓ Solifenacin Mylan RTHO-TOLIDINE 7.50 50 test OP 60.00 1 Compound diagnostic strips 7.02 100 test OP 100 test OP Blue diagnostic strips 7.02 100 test OP Albustix Obstetric Preparations 7.02 100 test OP 100 test OP IFEPRISTONE 60.00 1 ✓ Mifegyne Tab 200 mg 180.00 3 ✓ Mifegyne | & Grans eff 4 g sachets | 2.22 | 28 | ✓ Ural | |
| Tab 10 mg | SOLIFENACIN SUCCINATE | | | | |
| Detection of Substances in Urine RTHO-TOLIDINE Compound diagnostic sticks Compound diagnostic sticks (8.25) Hemastix ETRABROMOPHENOL Blue diagnostic strips (13.92) Albustix Destetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg 180.00 3 Vifegyne | Tab 5 mg | 2.05 | 30 | Solifenacia | n Mylan |
| RTHO-TOLIDINE 7.50 50 test OP Compound diagnostic sticks (8.25) Hemastix ETRABROMOPHENOL 7.02 100 test OP Blue diagnostic strips 7.02 100 test OP (13.92) Albustix Obstetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg 60.00 1 ✓ Mifegyne 180.00 3 ✓ Mifegyne | Tab 10 mg | 3.72 | 30 | Solifenacia | n Mylan |
| RTHO-TOLIDINE 7.50 50 test OP Compound diagnostic sticks 7.50 50 test OP (8.25) Hemastix ETRABROMOPHENOL 7.02 100 test OP Blue diagnostic strips 7.02 100 test OP (13.92) Albustix Obstetric Preparations Image: Compound diagnostic strips Antiprogesterones Image: Compound diagnostic strips IFEPRISTONE 60.00 1 ✓ Mifegyne Tab 200 mg 3 ✓ Mifegyne | | | | | |
| Compound diagnostic sticks 7.50 50 test OP (8.25) Hemastix ETRABROMOPHENOL 100 test OP Blue diagnostic strips 7.02 100 test OP (13.92) Albustix Obstetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg 60.00 1 ✓ Mifegyne 180.00 3 ✓ Mifegyne | Detection of Substances in Urine | | | | |
| Compound diagnostic sticks 7.50 50 test OP (8.25) Hemastix ETRABROMOPHENOL 100 test OP Blue diagnostic strips 7.02 100 test OP (13.92) Albustix Obstetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg 60.00 1 ✓ Mifegyne 180.00 3 ✓ Mifegyne | | | | | |
| (8.25) Hemastix ETRABROMOPHENOL 7.02 100 test OP Blue diagnostic strips | | 7.50 | 50 test OP | | |
| ETRABROMOPHENOL | | | | Hemastix | |
| Blue diagnostic strips | | () | | | |
| (13.92) Albustix Destetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg | | 7 02 | 100 test OP | | |
| Dbstetric Preparations Antiprogesterones IFEPRISTONE Tab 200 mg 1 ✓ Mifegyne 180.00 3 ✓ Mifegyne | | | 100 1001 01 | Albustix | |
| Antiprogesterones IFEPRISTONE Tab 200 mg | | (10102) | | , abdoux | |
| Antiprogesterones IFEPRISTONE Tab 200 mg | Obstetric Preparations | | | | |
| IFEPRISTONE | • | | | | |
| IFEPRISTONE 60.00 1 ✓ Mifegyne Tab 200 mg 180.00 3 ✓ Mifegyne | Antiprogesterones | | | | |
| Tab 200 mg | | | | | |
| 180.00 3 🖌 Mifegyne | | 00.00 | | | |
| | 1 ad 200 mg | | | | |
| aj up lu 15 lau avaliable UI a PSU | a) Up to 15 tab available on a BSO | 180.00 | 3 | • whegyne | |
| b) Only on a PSO | , , | | | | |

GENITO-URINARY SYSTEM

b) Only on a PSO

| | Subsidy | F | Fully Brand or |
|--|------------------------|---------------|---------------------------------------|
| | (Manufacturer's Price) | Subsid | |
| | \$ | Per | Manufacturer |
| Calcium Homeostasis | | | |
| CALCITONIN | | | |
| * Inj 100 iu per ml, 1 ml ampoule | 121.00 | 5 | ✓ Miacalcic |
| CINACALCET - Special Authority see SA1618 below - Retail pha | rmacy | | |
| Tab 30 mg | | 28 | Cinacalet Devatis |
| a) Brand switch fee payable (Pharmacode 2634120) - se | | ls | |
| b) Wastage claimable | | | |
| Tab 60 mg – Wastage claimable | | 28 | Cinacalet Devatis |
| ➡SA1618 Special Authority for Subsidy | | | |
| Initial application only from a nephrologist or endocrinologist. Ap following criteria: Either: | provals valid for 6 m | nonths for ap | oplications meeting the |
| 1 All of the following: | | | |
| 1.1 The patient has been diagnosed with a parathyroid of 1.2 The patient has persistent hypercalcaemia (serum c first-line treatments including sodium thiosulfate (wh 1.3 The patient is symptomatic; or 2 All of the following: | alcium greater than | or equal to | , , , |
| 2.1 The patient has been diagnosed with calciphylaxis (2.2 The patient has symptomatic (e.g. painful skin ulcer 3 mmol/L); and | rs) hypercalcaemia (| (serum calci | um greater than or equal to |
| 2.3 The patient's condition has not responded to previou thiosulfate. | is first-line treatmen | ts including | bisphosphonates and sodium |
| Renewal only from a nephrologist or endocrinologist. Approvals v meeting the following criteria: | alid without further r | enewal unle | ss notified for applications |
| Both: | | | |
| 1 The patient's serum calcium level has fallen to < 3mmol/L; a | | | |
| 2 The patient has experienced clinically significant symptom i | • | | |
| Note: This does not include parathyroid adenomas unless these h | ave become malign | ant. | |
| ZOLEDRONIC ACID | | | |
| Inj 4 mg per 5 ml, vial – Special Authority see SA2109 below - | | | |
| Retail pharmacy | 18.00 | 1 | Zoledronic acid |
| | | | <u>Mylan</u> |
| SA2109 Special Authority for Subsidy | | | |
| Initial application — (bone metastases) from any relevant pract | itioner. Approvals v | alid without | further renewal unless notified |
| for applications meeting the following criteria: | | | |
| Any of the following: | | | |
| Patient has hypercalcaemia of malignancy; or Both: | | | |
| 2.1 Patient has bone metastases or involvement; and | | | |
| 2.2 Patient has severe bone pain resistant to standard fi | irst-line treatments; | or | |
| 3 Both: | | | |
| 3.1 Patient has bone metastases or involvement; and | | | |
| 3.2 Patient is at risk of skeletal-related events pathologic | cal fracture, spinal c | ord compres | ssion, radiation to bone or |
| suraerv to bone. | | | |

Initial application - (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia.

Note: Indications marked with * are unapproved indications.

| Corticosteroids and Related Agents for Systemic Use | | | | | | |
|--|---|--|--|--|--|--|
| BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE * Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20 5 (36.96) | Celestone | | | | | |
| DEXAMETHASONE | Chronodose | | | | | |
| * Tab 0.5 mg - Up to 60 tab available on a PSO | <u></u> | | | | | |
| DEXAMETHASONE PHOSPHATE Dexamethasone phosphate injection will not be funded for oral use. | | | | | | |
| Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25 | ✓ <u>Dexamethasone</u> <u>Phosphate</u> Panpharma | | | | | |
| Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 16.37 | Dexamethasone Phosphate Panpharma | | | | | |
| FLUDROCORTISONE ACETATE | | | | | | |
| * Tab 100 mcg | Florinef | | | | | |
| * Tab 5 mg | Douglas | | | | | |
| * Tab 20 mg20.32 100 | - J | | | | | |
| * Inj 100 mg vial4.38 a) Up to 5 inj available on a PSO b) Only on a PSO | ✓ <u>Solu-Cortef</u> | | | | | |
| METHYLPREDNISOLONE | | | | | | |
| * Tab 4 mg 112.00 100 | | | | | | |
| * Tab 100 mg | Medrol | | | | | |

| | Subsidy (Manufacturer's Pric | e) Subs | | rand or Generic |
|---|---------------------------------|----------|-----------------------|---|
| | \$ | Per | M | lanufacturer |
| METHYLPREDNISOLONE (AS SODIUM SUCCINATE) | | | | |
| Inj 40 mg vial | 22.30 | 1 | | I-Medrol-Act- Vial |
| Inj 125 mg vial | 34.10 | 1 | | I-Medrol-Act- Vial |
| Inj 500 mg vial | 26.88 | 1 | | I-Medrol-Act- Vial |
| Inj 1 g vial | | 1 | 🗸 Solu | I-Medrol |
| METHYLPREDNISOLONE ACETATE | | | | |
| Inj 40 mg per ml, 1 ml vial | 47.06 | 5 | 🗸 Dep | o-Medrol |
| PREDNISOLONE | | | _ | |
| * Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age. | 6.00 | 30 ml OP | ✓ <u>Red</u> | ipred |
| PREDNISONE | | | | |
| * Tab 1 mg | | 500 | | -Prednisone |
| * Tab 2.5 mg | 21 04 | 500 | | Inisone Clinect -Prednisone |
| - τως 2.0 mg | | 500 | | Inisone Clinect |
| * Tab 5 mg – Up to 30 tab available on a PSO | | 500 | | -Prednisone |
| | | | 🗸 Prec | Inisone Clinect |
| * Tab 20 mg – Up to 30 tab available on a PSO | 50.51 | 500 | • | -Prednisone Inisone Clinect |
| (Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) | | | | |
| TETRACOSACTRIN | | | | |
| * Inj 250 mcg per ml, 1 ml ampoule | 75.00 | 1 | 🗸 AU | Synacthen S29 Synacthen |
| * Inj 1 mg per ml, 1 ml ampoule | 690.00 | 1 | 🗸 Syn | acthen acthen Depot acthene etard \$29 |
| TRIAMCINOLONE ACETONIDE | | | | |
| Inj 10 mg per ml, 1 ml ampoule | | 5 | 🗸 Ken | acort-A 10 |
| Inj 40 mg per ml, 1 ml ampoule | | 5 | | acort-A 40 |
| Sex Hormones Non Contraceptive | | | | |
| Androgen Agonists and Antagonists | | | | |
| CYPROTERONE ACETATE | | | | |
| Tab 50 mg | | 50 | ✓ Site | rone |
| Tab 100 mg | | 50 | ✓ Site | |
| TESTOSTERONE | | | | |
| Patch 5 mg per day | 90.00 | 30 | 🗸 And | roderm |
| TESTOSTERONE CIPIONATE | | | | |
| Inj 100 mg per ml, 10 ml vial | | 1 | 🗸 Dep | o-Testosterone |
| | | | | |
| 80 Fully subsidised | S29 Unappro | | supplied und | ler Section 29 |

Principal Supply

Sole Subsidised Supply

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|------------|---------------------|--|
| TESTOSTERONE ESTERS Inj 250 mg per ml, 1 ml | 12 98 | 1 | 1 | Sustanon Ampoules |
| TESTOSTERONE UNDECANOATE | | | | · |
| Cap 40 mg – Subsidy by endorsement Subsidy by endorsement – subsidised for patients who w | | 60 ne u | | Andriol Testocaps e cap 40mg prior to |
| November 2021 and the prescription is endorsed accor where there exists a record of prior dispensing of testoste Inj 250 mg per ml, 4 ml vial | erone undecanoate o | | 0 mg in the | |

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

Oestrogens

| | TRADIOL – See prescribing guideline above | | |
|-------|--|-------|-------------------------------------|
| * | Tab 1 mg4.12 | 28 OP | |
| | (11.10) | | Estrofem |
| * | Tab 2 mg4.12 | 28 OP | |
| | (11.10) | _ | Estrofem |
| | Patch 25 mcg per day6.12 | 8 | Estradot |
| | 7.85 | | Estradiol TDP |
| | | | Mylan S29 |
| | a) No more than 2 patch per week | | |
| | b) Only on a prescription | | |
| I | Patch 50 mcg per day7.04 | 8 | Estradot 50 mcg |
| | 9.22 | | Estradiol TDP |
| | | | Mylan S29 |
| | a) No more than 2 patch per week | | - |
| | b) Only on a prescription | | |
| I | Patch 75 mcg per day7.91 | 8 | Estradot |
| | 10.60 | | Estradiol TDP |
| | | | Mylan S29 |
| | a) No more than 2 patch per week | | , |
| | b) Only on a prescription | | |
| 1 | Patch 100 mcg per day7.91 | 8 | Estradot |
| | a) No more than 2 patch per week | Ũ | Echladot |
| | b) Only on a prescription | | |
| /Ectr | adiol TDP Mylan (s29) Patch 25 mcg per day to be delisted 1 May 2022) | | |
| • | | | |
| • | adiol TDP Mylan ³²⁹ Patch 50 mcg per day to be delisted 1 May 2022) | | |
| , | adiol TDP Mylan 2010 Patch 75 mcg per day to be delisted 1 May 2022) | | |
| OES | TRADIOL VALERATE – See prescribing guideline above | | |
| * | Tab 1 mg | 84 | Progynova |
| * | Tab 2 mg | 84 | Progynova |
| OES | TROGENS – See prescribing guideline above | | |
| * (| Conjugated, equine tab 300 mcg | 28 | |
| | (17.50) | | Premarin |
| * (| Conjugated, equine tab 625 mcg4.12 | 28 | |
| | (17.50) | | Premarin |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | | Subsidy (Manufacturer's Price) | | Fully Subsidised | |
|-----|---|-----------------------------------|-----------|---------------------|---------------------|
| _ | | \$ | Per | | Manufacturer |
| Ρ | rogestogens | | | | |
| | DROXYPROGESTERONE ACETATE - See prescribing guid | | s page | | |
| | Tab 2.5 mg | | 30 | | Provera |
| | Tab 5 mg Tab 10 mg | | 100 30 | | Provera Provera |
| - | č | | 00 | • | Tiovera |
| P | rogestogen and Oestrogen Combined Prepara | tions | | | |
| | STRADIOL WITH NORETHISTERONE – See prescribing gui | | us pag | je | |
| * | Tab 1 mg with 0.5 mg norethisterone acetate | 5.40 | 28 OF | D | |
| | | (18.10) | ~~~~ | | Kliovance |
| * | Tab 2 mg with 1 mg norethisterone acetate | 5.40 (18.10) | 28 OF | • | Kliogest |
| * | Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg | (10.10) | | | Riogest |
| | oestradiol tab (12) and 1 mg oestradiol tab (6) | 5.40 | 28 OF | þ | |
| | | (18.10) | | | Trisequens |
| 0 | other Oestrogen Preparations | | | | |
| | HINYLOESTRADIOL – Subsidy by endorsement | | | | |
| | Subsidy by endorsement – Subsidised for patients who were prescription is endorsed accordingly. Pharmacists may anno prior dispensing of ethinyloestradiol. Tab 10 mcg | tate the prescription | | dorsed wh | |
| | · · | | | | Scientific |
| • | Z Medical and Scientific Tab 10 mcg to be delisted 1 February | 2023) | | | |
| | STRIOL | 7.00 | ~~ | | o |
| * | Tab 2 mg | 7.00 | 30 | ~ | Ovestin |
| С | ther Progestogen Preparations | | | | |
| I F | VONOBGESTBEL | | | | |
| | Intra-uterine device 52 mg | | 1 | 1 | Mirena |
| | Intra-uterine device 13.5 mg | | 1 | 1 | Jaydess |
| ME | DROXYPROGESTERONE ACETATE | | | | |
| | Tab 100 mg | 116.15 | 100 | 1 | Provera HD |
| | RETHISTERONE | | | | |
| * | Tab 5 mg – Up to 30 tab available on a PSO | 5.49 | 30 | 1 | Primolut N |
| PR | OGESTERONE | | | | |
| | Cap 100 mg - Special Authority see SA1609 below - Retail | | | | |
| _ | pharmacy | 16.50 | 30 | - | Utrogestan |
| | SA1609 Special Authority for Subsidy | annovala valid for 10 | manth | o for on-l | actions mosting the |
| | tial application only from an obstetrician or gynaecologist. Ap owing criteria: | provais valid for 12 | month | is for appli | cations meeting the |
| Ro | 0 | | | | |

Both:

82

1 For the prevention of pre-term labour*; and

2 Either:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
- 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 For the prevention of pre-term labour*; and

Thuroid and Antithuroid Agonte

2 Treatment is required for second or subsequent pregnancy; and

3 Either:

- 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
- 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

| | myrolu allu Allilliyrolu Agenis | | | |
|---|---------------------------------|-------|-------|------------------------------------|
| | ARBIMAZOLE Tab 5 mg | | 100 | ✓ Neo-Mercazole |
| | EVOTHYROXINE Tab 25 mcg | 5.55 | 90 | ✓ Synthroid |
| | Tab 50 mcg | | 28 | ✓ Mercury Pharma |
| | ũ | 5.79 | 90 | Synthroid |
| | | 64.28 | 1,000 | Eltroxin |
| * | Tab 100 mcg | | 28 | Mercury Pharma |
| | | 6.01 | 90 | Synthroid |
| | | 66.78 | 1,000 | Eltroxin |
| | | | | |

PROPYLTHIOURACIL – Special Authority see SA1199 below – Retail pharmacy

Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

| Tab 50 mg35.00 | 100 | PTU \$29 |
|----------------|-----|----------|
|----------------|-----|----------|

SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 The patient has hyperthyroidism; and

2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

| SO | MATROPIN (OMNITROPE) – Special Authority see S | A2032 below – Retail pharr | nacy | |
|----|--|----------------------------|------|-------------------------------|
| * | Inj 5 mg cartridge | | 1 | Omnitrope |
| * | Inj 10 mg cartridge | | 1 | Omnitrope |
| * | lnj 15 mg cartridge | 139.50 | 1 | ✓ Omnitrope |

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: Either:

| | Subsidy | F | ully | Brand or |
|-------|--------------------|----------|------|--------------|
| (Manu | Ifacturer's Price) | Subsidis | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or
- 2 All of the following:
 - Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and</p>
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under ; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and

continued...

84

| Subsidy | | Fully | Brand or |
|------------------------|--------|-------|--------------|
| (Manufacturer's Price) | Subsic | dised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

| Subsidy | | Fully | Brand or | |
|--------------------|--------|------------|--------------|--|
| (Manufacturer's Pr | ice) S | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight eximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

| Subsidy | | Fully | Brand or | |
|-------------------|-------|------------|--------------|--|
| (Manufacturer's P | rice) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and

2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or 3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN

| Implant 3.6 mg, syringe | 65.68 | 1 | ✓ Teva |
|--------------------------|-------|---|--------------------------|
| Implant 10.8 mg, syringe | | 1 | Teva |

| | Subsidy (Manufacturer's Prio \$ | ce) Per | Fully Subsidised | |
|---|---|---------------------|---------------------------|---------------------------------------|
| EUPRORELIN | | | | |
| Additional subsidy by endorsement where the patient is a chi | ld or adolescent a | nd is una | able to tole | rate administration of |
| goserelin and the prescription is endorsed accordingly. | | | | |
| Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy | of | | | |
| \$221.60 per 1 inj with Endorsement | 66.48 | 1 | | |
| | (221.60) | | | Lucrin Depot 1-month |
| Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy | | | | |
| of \$591.68 per 1 inj with Endorsement | 177.50 | 1 | | |
| | (591.68) | | | Lucrin Depot 3-month |
| Vasopressin Agonists | | | | |
| ESMOPRESSIN | | | | |
| Wafer 120 mcg | 47 00 | 30 | 1 | Minirin Melt |
| ESMOPRESSIN ACETATE | | 00 | • | |
| | 25.00 | 30 | 1 | Minirin |
| Tab 100 mcg Tab 200 mcg | | 30 | | Minirin |
| Nasal spray 10 mcg per dose | | 6 ml O | | Desmopressin- |
| | | 011110 | | <u>PH&T</u> |
| Inj 4 mcg per ml, 1 ml | 67.18 | 10 | 1 | Minirin |
| Other Endocrine Agents | | | | |
| | | | | |
| ABERGOLINE | | | | |
| Tab 0.5 mg – Maximum of 2 tab per prescription; can be | | | | |
| waived by Special Authority see SA2070 below | | 2 | | Dostinex |
| | 15.20 | 8 | ~ | Dostinex |
| SA2070 Special Authority for Waiver of Rule | | | | |
| itial application from any relevant practitioner. Approvals valid | d without further re | enewal u | nless notif | ed for applications meeting |
| e following criteria: | | | | |
| ny of the following: | | | | |
| 1 Hyperprolactinemia; or | | | | |
| 2 Acromegaly*; or | | | | |
| 3 Inhibition of lactation. | | | | |
| | | | CA100 | (1) from any relevant |
| enewal — (for patients who have previously been funded u | | | | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notifie | ed where the patie | nt has p | reviously h | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notified hich has expired and the treatment remains appropriate and the | ed where the patie | nt has p | reviously h | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notified hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. | ed where the patie | nt has p | reviously h | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notifie hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. LOMIFENE CITRATE | ed where the patie patient is benefiti | nt has p | reviously h | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notified hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. | ed where the patie patient is benefiti | nt has p | reviously f treatment. | |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notifie hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. LOMIFENE CITRATE | ed where the patie patient is benefiti | nt has p ng from | reviously f treatment. | ield a valid Special Author |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notifie hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. LOMIFENE CITRATE | ed where the patie patient is benefiti | nt has p ng from | reviously f treatment. | ield a valid Special Authori Mylan |
| enewal — (for patients who have previously been funded un actitioner. Approvals valid without further renewal unless notified hich has expired and the treatment remains appropriate and the ote: Indication marked with * is an unapproved indication. LOMIFENE CITRATE Tab 50 mg | ed where the patie patient is benefiti | nt has p ng from | reviously f treatment. | eÍd a valid Special Authori Mylan |

| | Subsidy | | Fully Brand or | |
|--|-------------------------|-----------|---------------------------------------|---------|
| | (Manufacturer's Price) | S | Subsidised Generic | |
| | \$ | Per | Manufacturer | |
| | | | | |
| Anthelmintics | | | | |
| ALRENDAZOLE Special Authority and SA1218 below Batai | hormooy | | | |
| ALBENDAZOLE – Special Authority see SA1318 below – Retai | | ~~ | | |
| Tab 400 mg | | 60 | Eskazole S29 | |
| SA1318 Special Authority for Subsidy | | | | |
| Initial application only from an infectious disease specialist or o | linical microbiologist. | Appro | vals valid for 6 months whe | ere the |
| patient has hydatids. | innear merebiere gieti | | | |
| | arabialagiat Approve | lo volid | for 6 months whore the tre | otmont |
| Renewal only from an infectious disease specialist or clinical mi | | lis valiu | | aimeni |
| remains appropriate and the patient is benefitting from the treatm | nent. | | | |
| MEBENDAZOLE – Only on a prescription | | | | |
| Tab 100 mg | 7.97 | 6 | Vermox | |
| Oral liq 100 mg per 5 ml | 2.18 | 15 ml | | |
| | (7.53) | | Vermox | |
| | (7.50) | | Verniex | |
| PRAZIQUANTEL | | | | |
| Tab 600 mg | | 8 | Biltricide | |
| - | | | | |
| Antibacterials | | | | |
| | | | | |
| a) For topical antibacterials, refer to DERMATOLOGICALS, page | 10 61 | | | |
| | | | | |
| b) For anti-infective eye preparations, refer to SENSORY ORG/ | AINS, page 237 | | | |
| Cephalosporins and Cephamycins | | | | |
| Cephalospornis and Cephalnycins | | | | |
| CEFACLOR MONOHYDRATE | | | | |
| | 04 70 | 100 | Panhavy Cafaal | |
| Cap 250 mg | 24.70 | 100 | Ranbaxy-Cefaclo | |
| | | | Ranbaxy-Cefaclo | or |
| | | | S29 S29 | |
| Grans for oral lig 125 mg per 5 ml - Wastage claimable | 3.53 | 100 ml | Ranbaxy-Cefaclo | or |
| | | | Ranbaxy-Cefaclo | |
| | | | - | |
| | | | S29 S29 | |
| CEFALEXIN | | | | |
| Cap 250 mg | 3.33 | 20 | Cephalexin ABM | |
| Cap 500 mg | | 20 | Cephalexin ABM | |
| Grans for oral liq 25 mg per ml – Wastage claimable | | 100 ml | ✓ Cefalexin Sando | |
| | | | | |
| Grans for oral liq 50 mg per ml – Wastage claimable | 11./5 | 100 ml | Cefalexin Sando | Z |
| CEFAZOLIN – Subsidy by endorsement | | | | |
| Only if prescribed for dialysis or cellulitis in accordance with | a DHB approved pro- | tocol an | nd the prescription is endors | sed |
| accordingly. | | | · · · · · · · · · · · · · · · · · · · | |
| Inj 500 mg vial | 3 30 | 5 | 🗸 AFT | |
| | | 5 | | |
| Inj 1 g vial | | 5 | ✓ <u>AFT</u> | |
| CEFTRIAXONE – Subsidy by endorsement | | | | |
| a) Up to 10 inj available on a PSO | | | | |
| b) Subsidised only if prescribed for a dialysis or cystic fibros | sis nationt or the trea | tmont o | of apporrhoad or the treatm | ont of |
| | | | | |
| pelvic inflammatory disease, or the treatment of suspect | eu meningococcal dis | ease, a | and the prescription of PSO | 19 |
| endorsed accordingly. | _ | | | |
| Inj 500 mg vial | 0.89 | 1 | Ceftriaxone-AFT | |
| Inj 1 g vial | 3.99 | 5 | Ceftriaxone-AFT | |
| CEFUROXIME AXETIL – Subsidy by endorsement | | | | |
| Only if properihed for prophylogic of and coordina and the and | porintion is and aread | 0000- | lingh | |
| Only if prescribed for prophylaxis of endocarditis and the pre | | | | |
| Tab 250 mg | | 50 | Zinnat | |
| | | | | |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

 $\ensuremath{\textbf{\#}}$ Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Sub: Per | Fully sidised | Brand or Generic Manufacturer |
|--|---|-------------|------------------|-------------------------------------|
| Macrolides | | | | |
| AZITHROMYCIN – Maximum of 5 days treatment per prescriptio A maximum of 24 months of azithromycin treatment for non-c Authority. | | | | |
| Tab 250 mg | 8.19 | 30 | 🗸 A | po-Azithromycin |
| Tab 500 mg – Up to 8 tab available on a PSO | 2.57 | 2 | ✓ Z | lithromax |
| Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage claimable | | 15 ml | ✓ Z | lithromax |

► SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

| CLARITHROMYCIN - Maximum of 500 mg per prescription; ca | n be waived by Spe | ecial Authori | ty see SA1857 below |
|---|--------------------|---------------|----------------------------|
| Tab 250 mg | 8.53 | 14 | Klacid |
| Grans for oral lig 250 mg per 5 ml - Wastage claimable | | 50 ml | ✓ Klacid |

➡SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

continued...

90

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

1 Atypical mycobacterial infection; or

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and

2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

| ERYTHROMYCIN (AS LACTOBIONATE) Inj 1 g vial | | 1 | Erythrocin IV |
|---|---------|--------|---|
| ERYTHROMYCIN ETHYL SUCCINATE | | | |
| Tab 400 mg | | 100 | E-Mycin |
| a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP | | | |
| Grans for oral liq 200 mg per 5 ml | 5.00 | 100 ml | E-Mycin |
| a) Up to 300 ml available on a PSO | | | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | |
| c) Wastage claimable | 6 77 | 100 ml | E Musin |
| Grans for oral liq 400 mg per 5 ml | 6.// | 100 ml | E-Mycin |
| a) Up to 200 ml available on a PSOb) Wastage claimable | | | |
| ERYTHROMYCIN STEARATE | | | |
| Tab 500 mg | 29.90 | 100 | |
| | (44.58) | 100 | ERA |
| (ERA Tab 500 mg to be delisted 1 September 2022) | (1.100) | | |
| ROXITHROMYCIN | | | |
| Tab disp 50 mg | 8.29 | 10 | Rulide D |
| Restricted to children under 12 years of age. | | | |
| Tab 150 mg | 8.28 | 50 | <u>Arrow-</u> <u>Roxithromycin</u> |
| Tab 300 mg | | 50 | ✓ <u>Arrow-</u> <u>Roxithromycin</u> |

| Penicillins AMOXICILLIN Cap 250 mg | | Subsidy (Manufacturer's Price \$ |) Subs Per | Fully idised | Brand or Generic Manufacturer |
|--|---|--|---------------|-----------------|-------------------------------------|
| Cap 250 mg. 22.50 500 ✓ Alphamox a) Up to 30 cap available on a PSO 36.98 500 ✓ Alphamox b) Up to 10 x the maximum PSO quantity for RFPP 36.98 500 ✓ Alphamox a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP 500 ✓ Alphamox a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP 100 ml ✓ Alphamox 125 a) Up to 200 ml available on a PSO b) Wastage claimable 1.40 100 ml ✓ Alphamox 250 b) Wastage claimable 1.73 100 ml ✓ Alphamox 250 500 ✓ Alphamox 250 b) Up to 300 ml available on a PSO 1.73 100 ml ✓ Alphamox 250 500 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP 15.97 10 ✓ Ibiamox inj 500 mg vial 17.43 10 ✓ Ibiamox inj 1 g vial – Up to 5 inj available on a PSO 21.64 10 ✓ Ibiamox AMOXICILLIN WITH CLAVULANIC ACID 74.00 ✓ Ibiamox 10 ✓ Ibiamox Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab available on a PSO 0.89 10 ✓ Curam Duo 500/125 | illins | | | | |
| a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg | ILLIN | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg | 5 | 22.50 | 500 | ✓ <u>A</u> | lphamox |
| Cap 500 mg | | | | | |
| a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml | , | 26.09 | 500 | ۷ ۸ | Inhamov |
| b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml | 5 | | 500 | • <u>A</u> | iphaniox |
| Grans for oral liq 125 mg per 5 ml | | | | | |
| a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 250 mg per 5 ml | | | 100 ml | 🗸 A | lphamox 125 |
| Grans for oral liq 250 mg per 5 ml | | | | _ | • |
| a) Up to 300 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable Inj 250 mg vial | b) Wastage claimable | | | | |
| b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable Inj 250 mg vial | ns for oral liq 250 mg per 5 ml | 1.73 | 100 ml | ✓ <u>A</u> | Iphamox 250 |
| c) Wastage claimable Inj 250 mg vial | , , | | | | |
| Inj 250 mg vial 15.97 10 ✓ Ibiamox Inj 500 mg vial 17.43 10 ✓ Ibiamox Inj 1 g vial – Up to 5 inj available on a PSO 21.64 10 ✓ Ibiamox AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab 0.89 10 ✓ Curam Duo 500/125 Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg 5.00 100 ml ✓ Augmentin | | | | | |
| Inj 500 mg vial | c) Wastage claimable | 45.07 | | | |
| In j 1 g vial – Up to 5 inj available on a PSO | | | | | |
| AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab available on a PSO | | | | | |
| Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab 0.89 10 ✓ Curam Duo 500/125 Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg 5.00 100 ml ✓ Augmentin | • • • | 21.04 | 10 | • 10 | hamox |
| available on a PSO0.89 10 Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml | | | | | |
| Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml | | 0.90 | 10 | | uram Dua 500/125 |
| per ml | | | 10 | • • | uram Duo 500/125 |
| , g | , | • | 100 ml | ۸ 🗸 | uamentin |
| | a) Up to 200 ml available on a PSO | | 100 111 | • • | agmentin |
| b) Wastage claimable | , i | | | | |
| Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg | , 0 | mg | | | |
| per ml – Up to 200 ml available on a PSO | | | 00 ml OP | ✓ C | uram |
| BENZATHINE BENZYLPENICILLIN | | | | | |
| Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj | | | | | |
| available on a PSO | | | 10 | 🗸 В | icillin LA |
| BENZYLPENICILLIN SODIUM (PENICILLIN G) | PENICILLIN SODIUM [PENICILLIN G] | | | | |
| Inj 600 mg (1 million units) vial – Up to 5 inj available on a PSO11.09 10 ✓ Sandoz | | SO 11.09 | 10 | ✓ <u>S</u> | andoz |

| | Subsidy | | Fully | Brand or |
|--|---------------------------------|---------|---------|--------------------------------|
| | (Manufacturer's Price | | sidised | |
| | \$ | Per | / | Manufacturer |
| FLUCLOXACILLIN | 15 70 | 050 | | |
| Cap 250 mg – Up to 30 cap available on a PSO | 15.79 | 250 | | Flucloxacillin-AFT Staphlex |
| Flucloxacillin-AFT to be Principal Supply on 1 May 2022 | | | | |
| Cap 500 mg – Up to 30 cap available on a PSO | 52.99 | 500 | | Flucloxacillin-AFT Staphlex |
| Flucloxacillin-AFT to be Principal Supply on 1 May 2022 | | | | - |
| Grans for oral liq 25 mg per ml | 3.29 | 100 ml | 1 | AFT |
| a) Up to 200 ml available on a PSOb) Wastage claimable | | | | |
| Grans for oral liq 50 mg per ml | 3.68 | 100 ml | 1 | AFT |
| a) Up to 200 ml available on a PSO b) Wastage claimable | | | | |
| Inj 250 mg vial | | 10 | 1 | Flucloxin |
| Inj 500 mg vial | | 10 | | Flucloxin |
| Inj 1 g vial – Up to 5 inj available on a PSO | 5.70 | 5 | 1 | Flucil |
| (Staphlex Cap 250 mg to be delisted 1 May 2022) (Staphlex Cap 500 mg to be delisted 1 May 2022) | | | | |
| PHENOXYMETHYLPENICILLIN (PENICILLIN V) | | | | |
| Cap 250 mg – Up to 30 cap available on a PSO | 3.84 | 50 | 1 | Cilicaine VK |
| Cap 500 mg | | 50 | 1 | Cilicaine VK |
| a) Up to 20 cap available on a PSO | | | | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | | |
| Grans for oral liq 125 mg per 5 ml. | 2.99 | 100 ml | 1 | <u>AFT</u> |
| a) Up to 200 ml available on a PSOb) Wastage claimable | | | | |
| Grans for oral lig 250 mg per 5 ml | 3 00 | 100 ml | 1 | AFT |
| a) Up to 300 ml available on a PSO | | 100 111 | • | |
| b) Up to 2 x the maximum PSO quantity for RFPP | | | | |
| c) Wastage claimable | | | | |
| PROCAINE PENICILLIN | | | | |
| Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO | 123 50 | 5 | 1 | Cilicaine |
| nij 1.5 g in 3.4 nii synnge – op to 5 nij avaliable on a PSO. | | 5 | • | Chicalite |
| Tetracyclines | | | | |
| DOXYCYCLINE | | | | |
| * Tab 100 mg – Up to 30 tab available on a PSO | 64.43 | 500 | 1 | Doxine |
| MINOCYCLINE HYDROCHLORIDE | | | | |
| * Tab 50 mg – Additional subsidy by Special Authority see | | | | |
| SA1355 below - Retail pharmacy | 5.79 | 60 | | |
| | (12.05) | | | Mino-tabs |
| * Cap 100 mg | | 100 | | |
| | (52.04) | | | Minomycin |
| SA1355 Special Authority for Manufacturers Price | | | | |
| Initial application from any relevant practitioner. Approvals val rosacea. | | | s notif | ied where the patient has |
| TETRACYCLINE – Special Authority see SA1332 on the next p | <mark>age –</mark> Retail pharm | асу | | |
| Tab 250 mg | 21.42 | 28 | 1 | Accord S29 |
| | | | | |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|----------|---------------------|---------------------------|
| SA1332 Special Authority for Subsidy itial application from any relevant practitioner. Approvals va oth: | lid for 3 months for app | olicati | ons meetir | ng the following criteria |
| For the eradication of helicobacter pylori following unsuc For use only in combination with bismuth as part of a quite | | | opriate first | t-line therapy; and |
| Other Antibiotics | | | | |
| or topical antibiotics, refer to DERMATOLOGICALS, page 61 IPROFLOXACIN Recommended for patients with any of the following: i) microbiologically confirmed and clinically significant part ii) prostatitis; or iii) pyelonephritis; or iv) gonorrhoea. | seudomonas infection; | or | | |
| Tab 250 mg – Up to 5 tab available on a PSO | 2.42 | 28 | 1 | Cipflox |
| Tab 500 mg – Up to 5 tab available on a PSO | 3.40 | 28 | | Cipflox |
| Tab 750 mg | 5.95 | 28 | ~ | Cipflox |
| | | | | |
| Cap hydrochloride 150 mg Inj phosphate 150 mg per ml, 4 ml ampoule | | 24 10 | | Dalacin C Dalacin C |
| | | | • | |
| OLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – Only if prescribed for dialysis or cystic fibrosis patient and the | | | accordinal | u . |
| Inj 150 mg | | 1360 | | v. Colistin-Link |
| ENTAMICIN SULPHATE | | | | |
| Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement | | 5 | 1 | DBL Gentamicin |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | t or complicated urinary | y trac | t infection | and the prescription is |
| Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement | 91.00 | 5 | 1 | Wockhardt S29 |
| | 182.00 | 10 | | Teligent S29 |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | t or complicated urinary | y trac | t infection | and the prescription is |
| Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement | | 10 | | Pfizer |
| Only if a second and for a state of a second of the second s | 87.50 | 50 | | Pfizer |
| Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly. | t or complicated urinary | y traci | Infection : | and the prescription is |
| OXIFLOXACIN – Special Authority see SA1740 below – Reta No patient co-payment payable | | | | |
| Tab 400 mg | | | | Avelox |

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

94

1.1 Active tuberculosis*; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.
- Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Either:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and
- 3 Treatment is only for 7 days.

Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.

Note: Indications marked with * are unapproved indications.

PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy

Cap 250 mg......126.00 16 🖌 Humatin 😒

► SA1689 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

1 Patient has confirmed cryptosporidium infection; or

2 For the eradication of Entamoeba histolyica carriage.

Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:

Either:

- 1 Patient has confirmed cryptosporidium infection; or
- 2 For the eradication of Entamoeba histolyica carriage.

PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy

➡SA1328 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or
- 2 For pregnant patients for the term of the pregnancy; or
- 3 For infants with congenital toxoplasmosis until 12 months of age.

SODIUM FUSIDATE [FUSIDIC ACID]

| Tab 250 mg | 67.85 | 36 | Fucidin |
|------------|-------|----|-----------------------------|
|------------|-------|----|-----------------------------|

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Fully | |
|---|--------------------------|-----------------|--------------|------------------------------|
| | (Manufacturer's Pr \$ | rice) Per | Subsidised | Generic Manufacturer |
| SULFADIAZINE SODIUM - Special Authority see SA1331 below | / – Retail pharma | су | | |
| Tab 500 mg | | 56 | ~ | Wockhardt S29 |
| SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months | a period of 3 mc | | nless notif | ied for applications meeting |
| TOBRAMYCIN | | | | |
| Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient an | | 5 Lio andorr | | Tobramycin Mylan |
| Solution for inhalation 60 mg per ml, 5 ml – Subsidy by | u ille prescription | | eu accon | lingiy. |
| endorsement | | 56 dos | e 🗸 | Tobramycin BNM |
| a) Wastage claimable | | | | |
| b) Only if prescribed for a cystic fibrosis patient and the | prescription is en | ndorsed ad | ccordingly | |
| TRIMETHOPRIM * Tab 300 mg – Up to 30 tab available on a PSO | 18 55 | 50 | 1 | ТМР |
| TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX. | | 50 | • | <u></u> |
| Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – L | - | | | |
| to 30 tab available on a PSO | 64.80 | 500 | 1 | <u>Trisul</u> |
| Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 n available on a PSO | | 100 m | · • | Deprim |
| VANCOMYCIN - Subsidy by endorsement | | | | |
| Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is | | | s or for tre | eatment of Clostridium |
| Inj 500 mg vial | | ungiy. 1 | 1 | Mylan |
| | | | | |
| Antifungals | | | | |
| a) For topical antifungals refer to DERMATOLOGICALS, page 6 | 2 | | | |
| b) For topical antifungals refer to GENITO URINARY, page 75 | | | | |
| FLUCONAZOLE | | | | |
| Cap 50 mg Cap 150 mg | | 28 1 | | <u>Mylan</u> Mylan |
| Cap 200 mg | | 28 | | Mylan |
| Powder for oral suspension 10 mg per ml - Special Authority | | | | |
| see SA1359 below – Retail pharmacy Wastage claimable | 109.34 | 35 ml | 1 | Diflucan |
| ■ SA1359 Special Authority for Subsidy Initial application — (Systemic candidiasis) from any relevant meeting the following criteria: Both: | practitioner. Ap | provals va | alid for 6 w | eeks for applications |
| Patient requires prophylaxis for, or treatment of systemic of Patient is unable to swallow capsules. | candidiasis; and | | | |

Initial application - (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications

continued...

96

| INF | ECTIONS - | AGE | IN 1 3 | | R SYSTEMIC USE |
|--|-------------------------------------|----------|-----------|---------------------------|------------------------------------|
| (M: | Subsidy anufacturer's Prio \$ | | Sul er | Full <u>y</u> osidised | d Generic |
| continued meeting the following criteria: All of the following: | | | | | |
| Patient is immunocompromised; and Patient is at moderate to high risk of invasive fungal infection; Patient is unable to swallow capsules. | and | | | | |
| Renewal — (Systemic candidiasis) from any relevant practitioner. following criteria: Both: | Approvals va | lid for | 6 wee | ks for | applications meeting the |
| Patient requires prophylaxis for, or treatment of systemic cand Patient is unable to swallow capsules. | didiasis; and | | | | |
| Renewal — (Immunocompromised) from any relevant practitioner following criteria: All of the following: | . Approvals va | alid for | 6 mo | nths fo | or applications meeting the |
| Patient remains immunocompromised; and Patient remains at moderate to high risk of invasive fungal info Patient is unable to swallow capsules. | ection; and | | | | |
| ITRACONAZOLE | | | | | |
| Cap 100 mg | 4.27 | 1 | 5 | - | Itrazole |
| Oral liq 10 mg per ml – Special Authority see SA1322 below – Retail pharmacy | 141.80 | 150 m | nl OP | 1 | Sporanox |
| ▶ SA1322 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinical practitioner on the recommendation of a infectious disease physician valid for 6 months where the patient has a congenital immune deficie Renewal from any relevant practitioner. Approvals valid for 6 month benefitting from the treatment. | , clinical micro ncy. | biologi | ist or | clinica | l immunologist. Approvals |
| KETOCONAZOLE | | | | | |
| Tab 200 mg - PCT | CBS | 30 | 0 | | Link Healthcare S29 Nizoral S29 |
| | | 10 | 0 | 1 | Strides Shasun S29 |
| NYSTATIN | | | | | |
| Tab 500,000 u | 14.16 (17.09) | 50 | 0 | | Nilstat |
| Cap 500,000 u | 12.81 | 50 | 0 | | |
| POSACONAZOLE - Special Authority see SA1285 below - Retail p | (15.47) harmaov | | | | Nilstat |
| Tab modified-release 100 mg | | 24 | 4 | - | Noxafil |
| Oral liq 40 mg per ml | | 105 m | nl OP | ~ | Noxafil |
| ■ SA1285 Special Authority for Subsidy Initial application only from a haematologist or infectious disease sp meeting the following criteria: Either: | pecialist. Appr | rovals | valid 1 | ior 6 w | eeks for applications |

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or |
|-----------------------|--------|-------|--------------|
| Manufacturer's Price) | Subsic | lised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

| * Tab 250 mg8 | 3.15 | 84 | Deolate |
|---|------|-------|-----------------------------|
| VORICONAZOLE - Special Authority see SA1273 below - Retail pharma | асу | | |
| Tab 50 mg9* | 1.00 | 56 | Vttack |
| Tab 200 mg | 00.0 | 56 | Vttack |
| Powder for oral suspension 40 mg per ml – Wastage | | | |
| claimable1,523 | 3.22 | 70 ml | Vfend |

► SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

98

PRIMAQUINE – Special Authority see SA1684 on the next page – Retail pharmacy Tab 15 mg400.00 100

Sanofi
 Primaguine S29

■SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

. -avalo molori

| The patient has relapsed vivax or ovale malaria; and Primaquine is to be given for a maximum of 21 days. | | | |
|---|--------------------|----------------|---|
| Antitrichomonal Agents | | | |
| METRONIDAZOLE | | | |
| Tab 200 mg – Up to 30 tab available on a PSO Tab 400 mg – Up to 15 tab available on a PSO | | 250 21 | ✓ <u>Metrogyl</u> ✓ Metrogyl |
| Oral lig benzoate 200 mg per 5 ml | | 100 ml | ✓ <u>Metrogyr</u> ✓ Flagyl-S |
| Suppos 500 mg | | 10 | ✓ Flagyl |
| ORNIDAZOLE | | | |
| Tab 500 mg | | 10 | ✓ <u>Arrow-Ornidazole</u> |
| Antituberculotics and Antileprotics | | | |
| Note: There is no co-payment charge for all pharmaceuticals liste immigration status. | d in the Antitube | erculotics and | Antileprotics group regardless of |
| CLOFAZIMINE – Retail pharmacy-Specialist | | | |
| a) No patient co-payment payable | | | |
| b) Prescriptions must be written by, or on the recommendation dermatologist. | on of, an infectio | us disease ph | nysician, clinical microbiologist or |
| * Cap 50 mg | | 100 | Lamprene S29 |
| CYCLOSERINE – Retail pharmacy-Specialist | | | |
| a) No patient co-payment payable | | | |
| b) Prescriptions must be written by, or on the recommendation requirements of the provision | on of, an infectio | us disease pr | vysician, clinical microbiologist or |
| respiratory physician. Cap 250 mg | 344 00 | 60 | Cyclorin S29 |
| DAPSONE – Retail pharmacy-Specialist | | 00 | e eyelenin eze |
| a) No patient co-payment payable | | | |
| b) Prescriptions must be written by, or on the recommendation dermatologist | on of, an infectio | us disease ph | nysician, clinical microbiologist or |
| Tab 25 mg | | 100 | Dapsone |
| Tab 100 mg | 329.50 | 100 | ✓ Dapsone |
| ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist | | | |
| a) No patient co-payment payable | | | |
| b) Prescriptions must be written by, or on the recommendation respiratory physician | on of, an infectio | us disease ph | nysician, clinical microbiologist or |
| Tab 100 mg | | 100 | EMB Fatol S29 |
| Tab 400 mg | | 56 | Myambutol S29 |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|---|--|----------------|----------------------------------|--|
| SONIAZID – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician | ion of, an internal me | edicine | physician | , paediatrician, clinical |
| * Tab 100 mg | | 100 | ~ | PSM |
| SONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat microbiologist, dermatologist or public health physician | | dicine | physician | , paediatrician, clinical |
| * Tab 100 mg with rifampicin 150 mg | | 100 | | Rifinah |
| * Tab 150 mg with rifampicin 300 mg | 179.13 | 100 | / | Rifinah |
| PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician Grans for oral lig 4 g sachet | | disease 30 | | t, clinical microbiologist or Paser ^{\$29} |
| PROTIONAMIDE – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician Tab 250 mg | | disease 100 | • | t, clinical microbiologist or |
| PYRAZINAMIDE – Retail pharmacy-Specialist | | | | |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat respiratory physician * Tab 500 mg | | disease 100 | | n, clinical microbiologist of |
| RIFABUTIN – Retail pharmacy-Specialist | | | | • |
| a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendat gastroenterologist | | | | |
| ₭ Cap 150 mg | | 30 | ~ | Mycobutin |
| RIFAMPICIN – Subsidy by endorsement a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection antimicrobial based on susceptibilities and the prescriptic Retail pharmacy - Specialist. Specialist must be an inter paediatrician, or public health physician. Cap 150 mg Cap 300 mg Yoral liq 100 mg per 5 ml | on is endorsed accord nal medicine physicia | dingly; d | can be wa cal micro ✓ ✓ | aived by endorsement - |
| Antivirals | | | | |
| For eye preparations refer to Eye Preparations, Anti-Infective Pre | eparations, page 237 | | | |
| Hepatitis B Treatment | · - | | | |
| ENTECAVIR ¥ Tab 0.5 mg | | 30 | 1 | Entecavir Sandoz |

| | Subsidy | | Fully | Brand or |
|--|----------------------------|-----------------|------------|------------------------------|
| (N | lanufacturer's Price \$ | e) Subsi Per | dised | Generic Manufacturer |
| | nacy | | | |
| Tab 100 mg | | 28 | 1 | Zetlam |
| Oral liq 5 mg per ml | | 240 ml OP | 1 | Zeffix |
| ⇒SA1685 Special Authority for Subsidy | | | | |
| nitial application only from a relevant specialist or medical practition | oner on the reco | mmendation | of a re | elevant specialist. |
| Approvals valid for 1 year where used for the treatment or prevention | | | | · |
| Renewal from any relevant practitioner. Approvals valid for 2 years | where used for | the treatment | t or pr | revention of hepatitis B. |
| TENOFOVIR DISOPROXIL | | | | |
| Tenofovir disoproxil prescribed under endorsement for the treat | | cluded in the | coun | t of up to 4 subsidised |
| antiretrovirals for the purposes of Special Authority SA1651., pa | | | | |
| * Tab 245 mg (300.6 mg as a succinate) | 38.10 | 30 | • | Tenofovir Disoproxil Teva |
| | | | | Teva |
| Herpesvirus Treatments | | | | |
| ACICLOVIR | | | | |
| Tab dispersible 200 mg | 1.60 | 25 | ~ 1 | Lovir |
| Tab dispersible 400 mg | | 56 | | Lovir |
| * Tab dispersible 800 mg | | 35 | - | Lovir |
| ALACICLOVIR | | | - | |
| Tab 500 mg | 6.50 | 30 | 1 | Vaclovir |
| Tab 1,000 mg | | 30 | 1 | Vaclovir |
| VALGANCICLOVIR - Special Authority see SA1993 below - Retail | | | - | |
| Tab 450 mg | | 60 | 1 | Valganciclovir |
| - | | | - | Mylan |

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
- 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

| GLECAPREVIR WITH PIBRENTASVIR – [Xpharm] Note the supply of treatment is via Pharmac's approved direct | t distribution supp | ly. Further d | etails can be found on Pharmac's |
|---|---------------------|----------------|----------------------------------|
| website https://pharmac.govt.nz/maviret | | | |
| Tab 100 mg with pibrentasvir 40 mg | 24,750.00 | 84 OP | Maviret |
| LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authori | ty see SA1605 be | low | |
| No patient co-payment payable | | | |
| Tab 90 mg with sofosbuvir 400 mg | 24,363.46 | 28 | Harvoni |
| SA1605 Special Authority for Subsidy | | | |
| Special Authority approved by the Hepatitis C Treatment Panel (H | lepCTP) | | |
| Notes: By application to the Hepatitis C Treatment Panel (HepCT | ſP). | | |
| Applications will be considered by HepCTP and approved subject | to confirmation of | f eligibility. | |
| Application details may be obtained from Pharmac's website http: | //www.pharmac.g | ovt.nz/mavire | t or: |
| The Coordinator, Hepatitis C Treatment Panel | | | |
| Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990, | | | |
| Email: hepcpanel@pharmac.govt.nz | | | |

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Su | ubsidised | Generic |
| \$ | Per | 1 | Manufacturer |

30

🗸 Teva

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see SA1994 below

below Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

➡SA1994 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Either:
 - 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months; or
 - 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | : | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

those risks: and

5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and

6 Fither:

- 6.1 All of the following:
 - 6.1.1 Patient is male or transgender; and
 - 6.1.2 Patient has sex with men; and
 - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months: and
 - 6.1.4 Any of the following:
 - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months: or
 - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
 - 6.1.4.3 Patient has used methamphetamine in the last three months: or
- 6.2 All of the following:
 - 6.2.1 Patient has a regular partner who has HIV infection; and
 - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
 - 6.2.3 Condoms have not been consistently used.

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability. Lagevrio 40

Cap 200 mg......0.00

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Tab 150 mg with ritonavir 100 mg0.00 30 Paxlovid

Antiretrovirals

SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application - (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

| EFAVIRENZ - Special Authority see SA1651 on the previous page | e – Retail pharm | lacy | |
|---|--------------------------------|------|-------------------------------|
| Tab 200 mg | 190.15 | 90 | Stocrin |
| Tab 600 mg | 63.38 | 30 | Stocrin |
| ETRAVIRINE - Special Authority see SA1651 on the previous pa | <mark>ge</mark> – Retail pharr | nacy | |
| Tab 200 mg | 770.00 | 60 | Intelence |

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's I \$ | Price) Subsi Per | Fully Brand or dised Generic Manufacturer |
|---|------------------------------------|---------------------|--|
| NEVIRAPINE – Special Authority see SA1651 on page 104 – Re | * | | |
| Tab 200 mg | | 60 | ✓ <u>Nevirapine</u> Alphapharm |
| Oral suspension 10 mg per ml | 203.55 | 240 ml | ✓ Viramune Suspension |
| Nucleosides Reverse Transcriptase Inhibitors | | | |
| ABACAVIR SULPHATE – Special Authority see SA1651 on pag | e 104 – Retail p | harmacy | |
| Tab 300 mg Oral lig 20 mg per ml | | 60 240 ml OP | ✓ <u>Ziagen</u> ✓ Ziagen |
| ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority. | v see SA1651 or | | tail pharmacy |
| Tab 600 mg with lamivudine 300 mg | 63.00 | 30 | ✓ Kivexa |
| EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPI | | I Authority see S | SA1651 on page 104 – Retail |
| pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil of anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopro | xil | | |
| 245 mg (300 mg as a maleate) | | 30 | ✓ <u>Mylan</u> |
| EMTRICITABINE – Special Authority see SA1651 on page 104 Cap 200 mg | | су 30 | ✓ Emtriva |
| LAMIVUDINE – Special Authority see SA1651 on page 104 – Re Tab 150 mg | | 60 | ✓ <u>Lamivudine</u> Alphapharm |
| Oral lig 10 mg per ml | | 240 ml OP | ✓ 3TC |
| ZIDOVUDINE [AZT] – Special Authority see SA1651 on page 10 | | | |
| Cap 100 mg | | 100 | Retrovir |
| Oral liq 10 mg per ml | 30.45 | 200 ml OP | Retrovir |
| ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority. | | | |
| Tab 300 mg with lamivudine 150 mg | | 60 | Alphapharm |
| Protease Inhibitors | | | |
| ATAZANAVIR SULPHATE – Special Authority see SA1651 on p | age 104 – Retai | l pharmacy | |
| Cap 150 mg | • | 60 | ✓ <u>Teva</u> |
| Cap 200 mg | | 60 | ✓ Teva |
| DARUNAVIR – Special Authority see SA1651 on page 104 – Re | etail pharmacy | | |
| Tab 400 mg | | 60 | Darunavir Mylan |
| Tab 600 mg | | 60 | Darunavir Mylan |
| LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 | | Retail pharmacy | |
| Tab 100 mg with ritonavir 25 mg – Brand switch fee payable (Pharmacode 2621959) - see page 242 for details | | 60 | ✓ Lopinavir/Ritonavir Mylan |
| Tab 200 mg with ritonavir 50 mg – Brand switch fee payable | | | |
| (Pharmacode 2621959) - see page 242 for details | | 120 | <u>Lopinavir/Ritonavir</u> <u>Mylan</u> |
| Oral liq 80 mg with ritonavir 20 mg per ml | 735.00 | 300 ml OP | ✓ Kaletra |
| | | | |

fully subsidised
 Principal Supply

106

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

| | Subsidy (Manufacturer's Price) \$ | Subsic Per | Fully dised | Brand or Generic Manufacturer |
|---|---|---------------------|----------------|-------------------------------------|
| RITONAVIR – Special Authority see SA1651 on page 104 – Reta Tab 100 mg | | 30 | ✓ <u>N</u> | orvir |
| Strand Transfer Inhibitors | | | | |
| DOLUTEGRAVIR – Special Authority see SA1651 on page 104 - Tab 50 mg | | 30 | ✔ Т | ivicay |
| RALTEGRAVIR POTASSIUM – Special Authority see SA1651 of Tab 400 mg Tab 600 mg | 1,090.00 | harmacy 60 60 | | entress sentress HD |

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects. Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - · PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0×10^9) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

1 Any of the following:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and

2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C. genotype 1: and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
INFECTIONS - AGENTS FOR SYSTEMIC USE

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or

3 Both:

- 3.1 Patient has a myeloproliferative disorder; and
- 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and

3.2.2 Either:

- 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
- 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE

100

Hiprex

INFECTIONS - AGENTS FOR SYSTEMIC USE

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|-----|---------------------|-------------------------------------|
| NITROFURANTOIN | | | | |
| * Tab 50 mg – Up to 30 tab available on a PSO | | 100 | ✓ | Nifuran |
| * Tab 100 mg | | 100 | ✓ | Nifuran |
| * Cap modified-release 100 mg - Wastage claimable | | 100 | ✓ | Macrobid |
| NORFLOXACIN | | | | |
| Tab 400 mg – Subsidy by endorsement | 245.00 | 100 | ✓ | Arrow-Norfloxacin |

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

| | Subsidy (Manufacturer's Price) | | Fully Brand or Subsidised Generic |
|---|-----------------------------------|--------|---------------------------------------|
| | (Manulactuler's Frice) | Per | Manufacturer |
| | | | |
| Anticholinesterases | | | |
| NEOSTIGMINE METILSULFATE | | | |
| Inj 2.5 mg per ml, 1 ml ampoule | | 10 | ✓ Max Health |
| PYRIDOSTIGMINE BROMIDE | | | |
| ▲ Tab 60 mg | 45.79 | 100 | Mestinon |
| Non-Steroidal Anti-Inflammatory Drugs | | | |
| DICLOFENAC SODIUM | | | |
| * Tab EC 25 mg | | 50 | Diclofenac Sandoz |
| * Tab 50 mg dispersible | | 20 | ✓ Voltaren D |
| * Tab EC 50 mg | | 50 | Diclofenac Sandoz |
| * Tab long-acting 75 mg | | 100 | ✓ Voltaren SR |
| | 22.80 | 500 | Apo-Diclo SR |
| * Tab long-acting 100 mg | 25.15 | 500 | Apo-Diclo SR |
| * Inj 25 mg per ml, 3 ml ampoule - Up to 5 inj available on a R | | 5 | ✓ Voltaren |
| * Suppos 12.5 mg | 2.04 | 10 | Voltaren |
| * Suppos 25 mg | 2.44 | 10 | Voltaren |
| * Suppos 50 mg – Up to 10 supp available on a PSO | 4.22 | 10 | Voltaren |
| * Suppos 100 mg | 7.00 | 10 | Voltaren |
| (Apo-Diclo SR Tab long-acting 75 mg to be delisted 1 May 2022) |) | | |
| (Apo-Diclo SR Tab long-acting 100 mg to be delisted 1 May 202 | 2) | | |
| IBUPROFEN | | | |
| * Tab 200 mg | | 1,000 | ✓ Relieve |
| * Tab long-acting 800 mg | | 30 | ✓ Brufen SR |
| * Oral liq 20 mg per ml | | 200 ml | ✓ Ethics |
| KETOPROFEN | | | |
| * Cap long-acting 200 mg | 12.07 | 28 | Oruvail SR |
| | 12.07 | 20 | |
| MEFENAMIC ACID | 4.05 | | |
| * Cap 250 mg | | 50 | |
| | (9.16) | ~~ | Ponstan |
| | 0.50 | 20 | |
| | (5.60) | | Ponstan |
| NAPROXEN | | | _ |
| * Tab 250 mg | | 500 | Noflam 250 |
| * Tab 500 mg | | 250 | ✓ Noflam 500 |
| * Tab long-acting 750 mg | | 28 | ✓ <u>Naprosyn SR 750</u> |
| * Tab long-acting 1 g | 8.62 | 28 | Naprosyn SR 1000 |
| SULINDAC | | | |
| * Tab 100 mg | 9.57 | 56 | 🗸 Mylan S29 |
| (Mylan 🖘 Tab 100 mg to be delisted 1 May 2022) | | | - |
| TENOXICAM | | | |
| * Tab 20 mg | 9 15 | 100 | Tilcotil |
| * Inj 20 mg vial | | 100 | ✓ <u>Incoun</u> ✓ AFT |
| | | | - 111 |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | | Full | |
|--|-----------------------------|----------|------------|----------------------------|
| | (Manufacturer's Price \$ |) Per | Subsidise | d Generic Manufacturer |
| | • | | | manadotaron |
| NSAIDs Other | | | | |
| ELECOXIB | | | | _ |
| Cap 100 mg | | 60 | | Celecoxib Pfizer |
| Cap 200 mg | 2.30 3.30 | 30 | | Celebrex |
| | 5.50 | | • | Celecoxib F lizer |
| Topical Products for Joint and Muscular Pain | | | | |
| APSAICIN | | | | |
| Crm 0.025% - Special Authority see SA1289 below - Retail | | | | |
| pharmacy | 9.75 | 45 g Ol | ⊳ √ | Zostrix |
| SA1289 Special Authority for Subsidy | | | | |
| nitial application from any relevant practitioner. Approvals valid | | | | |
| steoarthritis that is not responsive to paracetamol and oral non- | steroidal anti-inflamr | natorie | s are cor | traindicated. |
| Antirheumatoid Agents | | | | |
| | | | | |
| YDROXYCHLOROQUINE – Subsidy by endorsement | | | | |
| Subsidised only if prescribed for rheumatoid arthritis, system | | , | , | |
| suppression, relevant dermatological conditions (cutaneous f | • | | | |
| mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmo | | | | |
| Pharmacists may annotate the prescription as endorsed whe | | | orior disp | ensing of |
| hydroxychloroquine. Note: Indication marked with a * is an | | | | Diamanil |
| ★ Tab 200 mg | 8.78 | 100 | v | Plaquenil |
| EFLUNOMIDE Tab 10 mg | 6.00 | 20 | | ´ Arava |
| Tab 10 mg Tab 20 mg | | 30 30 | | Arava Arava |
| 8 | 0.00 | 50 | • | Alava |
| 'ENICILLAMINE Tab 125 mg | 67 23 | 100 | 1 | D-Penamine |
| Tab 250 mg | | 100 | | D-Penamine |
| · ~ | | | | 2 |
| Drugs Affecting Bone Metabolism | | | | |
| Alendronate for Osteoporosis | | | | |
| • | | | | |
| ALENDRONATE SODIUM | 0.44 | 4 | | Facamay |
| ₭ Tab 70 mg | 2.44 | 4 | • | <u>Fosamax</u> |
| ALENDRONATE SODIUM WITH COLECALCIFEROL | 4 54 | | | |
| * Tab 70 mg with colecalciferol 5,600 iu | 1.51 | 4 | • | Fosamax Plus |
| Other Treatments | | | | |
| ENOSUMAB – Special Authority see SA1777 below – Retail ph | | | | |
| Inj 60 mg prefilled syringe | | 1 | ✓ | Prolia |
| SA1777 Special Authority for Subsidy | | | | |
| nitial application from any relevant practitioner. Approvals valid | d without further ren | ewal ur | nless noti | fied for applications meet |
| he following criteria: | | | | |

the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
 - 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

| Inj 3 mg per ml, 10 ml vial | | 1 | Pamisol |
|--|--------------------|----------|-----------------------------|
| Inj 6 mg per ml, 10 ml vial | 74.67 | 1 | Pamisol |
| Inj 9 mg per ml, 10 ml vial | | 1 | Pamisol |
| RALOXIFENE HYDROCHLORIDE - Special Authority see SA177 | 9 on the next page | – Retail | pharmacy |
| * Tab 60 mg | 53.76 | 28 | Evista |

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Price) | Ful Subsidise | | |
|-----------------------------------|------------------|----------------------------------|--|
| \$ | Per • | Manufacturer | |

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or
- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

| Tab 35 mg3. | 10 4 | 4 | Risedronate Sandoz |
|---|------|---|--|
| TERIPARATIDE - Special Authority see SA1139 below - Retail pharmacy | | | |
| Inj 250 mcg per ml, 2.4 ml | 00 | 1 | Forteo |

SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily;

continued...

zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.

- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

ZOLEDRONIC ACID

100 ml OP 🖌 Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Su | bsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

- 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
- 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
 - 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
 - 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

| Subsidy | Fu | lly Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

| ALLOPURINOL | | | |
|---|--|-----|---|
| * Tab 100 mg | | 500 | DP-Allopurinol |
| * Tab 300 mg | | 500 | DP-Allopurinol |
| BENZBROMARONE - Special Authority see SA196 | <mark>3 below</mark> – Retail pharmacy | | |
| Tab 50 mg | | 100 | Narcaricin mite S29 |
| Tab 100 mg | | 30 | Desuric S29 |
| | | | Urinorm S29 |
| | 45.00 | 100 | Benzbromaron AL |
| | | | 100 S29 |

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

| * Tab 500 mcg6.00 | 100 | Colgout |
|--|-----|--------------------------------|
| FEBUXOSTAT - Special Authority see SA2054 on the next page - Retail pharmacy | | |
| Tab 80 mg | 28 | Febuxostat |
| | | multichem |
| Tab 120 mg20.00 | 28 | Febuxostat |
| | | multichem |

| Subsidy | e) | Fully | Brand or |
|-----------------------|-----|------------|--------------|
| (Manufacturer's Price | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
 - 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout..

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

| * Tab 500 mg66.95 | 100 | ✓ Probenecid-AFT |
|--|-----|--|
| Muscle Relaxants | | |
| BACLOFEN | | |
| * Tab 10 mg4.20 | 100 | Pacifen |
| Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorsement11.55 | 1 | Lioresal Intrathecal |
| Subsidised only for use in a programmable pump in patients where oral anti- caused intolerable side effects and the prescription is endorsed accordingly. | | ents have been ineffective or have |
| Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement | 5 | ✓ <u>Medsurge</u> |
| Subsidised only for use in a programmable pump in patients where oral anti- caused intolerable side effects and the prescription is endorsed accordingly. | | ents have been ineffective or have |
| DANTROLENE | | |
| Cap 25 mg97.50 | 100 | Dantrium |
| | | Dantrium S29 S29 |
| Cap 50 mg77.00 | 100 | Dantrium |
| ORPHENADRINE CITRATE | | |
| Tab 100 mg20.76 | 100 | ✓ <u>Norflex</u> |

| | Subsidy (Manufacturer's Price) \$ | Per | Subsidised (| Brand or Generic Manufacturer |
|--|---|--------|-------------------------|-------------------------------------|
| Agents for Parkinsonism and Related Disore | ders | | | |
| Dopamine Agonists and Related Agents | | | | |
| MANTADINE HYDROCHLORIDE | | | | |
| Cap 100 mg | | 60 | Syn | nmetrel |
| POMORPHINE HYDROCHLORIDE | 50.50 | - | <i>.</i> | |
| Inj 10 mg per ml, 2 ml ampoule | | 5 5 | | |
| Inj 10 mg per ml, 5 ml ampoule | 121.04 | Э | ✓ Mov | rapo |
| ROMOCRIPTINE MESYLATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who prescription is endorsed accordingly. Pharmacists may prior dispensing of bromocriptine mesylate. | annotate the prescription | as er | dorsed where | there exists a record of |
| Tab 2.5 mg | | 30 | Parl | odel S29 |
| Parlodel S29 Tab 2.5 mg to be delisted 1 September 2022 |) | | | |
| NTACAPONE – Brand switch fee payable (Pharmacode 2 | 634139) - see <mark>page 242</mark> fe | or det | ails | |
| Tab 200 mg | | 100 | ✓ <u>Con</u> | ntan |
| EVODOPA WITH BENSERAZIDE | | | | |
| Tab dispersible 50 mg with benserazide 12.5 mg | 13.25 | 100 | | lopar Rapid |
| Cap 50 mg with benserazide 12.5 mg | | 100 | | lopar 62.5 |
| Cap 100 mg with benserazide 25 mg | | 100 | | lopar 125 |
| Cap long-acting 100 mg with benserazide 25 mg | | 100 | | lopar HBS |
| Cap 200 mg with benserazide 50 mg | | 100 | Mac | lopar 250 |
| EVODOPA WITH CARBIDOPA | | | | |
| Tab 100 mg with carbidopa 25 mg | | 100 | ✓ Sine | |
| Tab long-acting 200 mg with carbidopa 50 mg | | 100 | | emet CR |
| Tab 250 mg with carbidopa 25 mg | | 100 | ✓ Sine | emet |
| RAMIPEXOLE HYDROCHLORIDE | | | <i></i> | |
| Tab 0.25 mg | | 100 | ✓ <u>Ran</u> | |
| Tab 1 mg | | 100 | ✓ Ran | nipex |
| ASAGILINE | | | | |
| fab 1 mg | 53.50 | 30 | 🗸 Azil | ect S29 |
| OPINIROLE HYDROCHLORIDE | | | | |
| Tab 0.25 mg | 2.85 | 84 | ✓ Rop | bin |
| | 3.39 | 100 | 🖌 Myl | an S29 |
| Tab 1 mg | 3.95 | 84 | ✓ Rop | bin |
| | 4.70 | 100 | 🖌 Myl | an S29 |
| Tab 2 mg | | 84 | ✓ Rop | |
| Tab 5 mg | 12.50 | 84 | ✓ <u>Rop</u> | <u>bin</u> |
| ELEGILINE HYDROCHLORIDE - Subsidy by endorsement | | | | |
| Subsidy by endorsement – Subsidised for patients who prescription is endorsed accordingly. Pharmacists may prior dispensing of selegiline hydrochloride. | | | • | • |
| Tab 5 mg | | 100 | 🗸 Elde | epryl S29 |
| OLCAPONE | | | | |
| | 150.00 | | / - | |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

✓ Tasmar

100

| Anticholinergics BENZATROPINE MESYLATE 9.59 60 ✓ Benztrop Tab 2 mg 9.59 60 ✓ Phebra a) Up to 10 inj available on a PSO 95.00 5 ✓ Phebra a) Up to 10 inj available on a PSO 90.00 ✓ Mebra ✓ Phebra a) Up to 10 inj available on a PSO 7.40 100 ✓ Kemadrin Agents for Essential Tremor, Chorea and Related Disorders RILUZOLE – Special Authority see SA1403 below – Retail pharmacy Wastage claimable Tab 5 mg | or C Icturer |
|---|------------------------|
| Tab 2 mg 9.59 60 ✓ Benztrop In 1 mg per mi, 2 mi 95.00 5 ✓ Phetra a) Up to 10 inj available on a PSO 95.00 5 ✓ Phetra b) Only on a PSO PROCYCLIDINE HYDROCHLORIDE 7.40 100 ✓ Kemadrin Agents for Essential Tremor, Chorea and Related Disorders RUUZOLE – Special Authority see SA1403 below – Retail pharmacy Wastage claimable 7.40 100 ✓ Kemadrin Tab 5 mg | |
| Tab 5 mg 7.40 100 ✓ Kemadrin Agents for Essential Tremor, Chorea and Related Disorders RILUZOLE - Special Authority see SA1403 below - Retail pharmacy Wastage claimable Tab 50 mg 130.00 56 ✓ Rilutek Special Authority for Subsidy Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications m following criteria: All of the following: 1 The patient has anot undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 3 The patient has not undergone a tracheostomy; and 4 The patient is able to use upper limbs; or 5.3 The patient is able to use upper limbs; or 5.3 The patient has not experienced respiratory failure; and 3 Any of the following: 1 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or </td <td></td> | |
| BLUZOLE - Special Authority see SA1403 below - Retail pharmacy Wastage claimable Tab 50 mg | ı |
| Wastage claimable Tab 50 mg 130.00 56 ✓ Rilutek ■SA1403 Special Authority for Subsidy initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications m ollowing criteria: All of the following: 1 The patient has anyotrophic lateral sclerosis with disease duration of 5 years or less; and 2 The patient has not undergone a tracheostomy; and 5 Any of the following: 5.1 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is able to use upper limbs; or 5.2 The patient is able to use upper limbs; or 5.2 The patient has not undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 3 Any of the following: 1 The patient has not undergone a tracheostomy; and 3 2 The patient has not undergone a tracheostomy; and 3 3 Any of the following: 1 1 3.1 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. FETRABENAZINE The patient is able to swallow. 91.10 112 ✓ Motetis Anae | |
| Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications m In the patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and 1 The patient has anyotrophic lateral sclerosis with disease duration of 5 years or less; and 2 The patient has not undergone a tracheostomy; and 4 The patient has not experienced respiratory failure; and 5 Any of the following: 5.1 The patient is able to use upper limbs; or 5.3 The patient is able to use upper limbs; or 5.3 The patient has not undergone a tracheostomy; and Prepatient has not undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 3 Any of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is able to use upper limbs; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. TETRABENAZINE Tab 25 mg 91.10 112 V Motetis Anaesthetics LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement. <td></td> | |
| 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial applic 3 The patient has not undergone a tracheostomy; and 4 The patient has not undergone a tracheostomy; and 5 Any of the following: 5.1 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteri All of the following: 1 The patient has not undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 2 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is able to use upper limbs; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.3 The patient is able to use upper limbs; or 3.4 The patient is able to use upper limbs; or 3.5 The patient is able to use upper limbs; or 3.6 The patient is able to use upper limbs; or 3.7 The patient is able to swallow. TETRABENAZINE Tab 25 mg | s meeting the |
| 1 The patient has not undergone a tracheostomy; and 2 The patient has not experienced respiratory failure; and 3 Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. IETRABENAZINE Tab 25 mg | |
| Tab 25 mg | |
| LIDOCAINE [LIGNOCAINE] Gel 2%, tube – Subsidy by endorsement | |
| IDOCAINE [LIGNOCAINE] Gel 2%, tube - Subsidy by endorsement | |
| Gel 2%, tube – Subsidy by endorsement | |
| fully subsidised (\$29) Unapproved medicine supplied under Section | accordingly. I Lido |
| 120 Driveling County | ction 29 |

| | Subsidy | | Fully Brand or |
|--|---|--|--|
| | | | , |
| | (Manufacturer's P | | sidised Generic |
| | \$ | Per | Manufacturer |
| IDOCAINE [LIGNOCAINE] HYDROCHLORIDE | | | |
| Oral (gel) soln 2% | 38.00 | 200 ml | Mucosoothe |
| Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO | | 25 | ✓ Lidocaine-Baxter |
| | 17.50 | 23 50 | |
| | | 50 | Vulaasias |
| | (35.00) | | Xylocaine |
| Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO | | 25 | Lidocaine-Baxter |
| Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO | | 5 | |
| | (20.00) | | Xylocaine |
| Inj 1%, 20 ml vial – Up to 5 inj available on a PSO | 6.20 | 5 | Lidocaine-Claris |
| Inj 2%, 20 ml vial – Up to 5 inj available on a PSO | 6.45 | 5 | Lidocaine-Baxter |
| | | | Lidocaine-Claris |
| (Lidocaine-Claris Inj 2%, 20 ml vial to be delisted 1 June 2022) | | | |
| | | | |
| IDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE | | | |
| Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes | - | | |
| Subsidy by endorsement | 103.32 | 10 | Pfizer |
| a) Up to 5 each available on a PSO | | | |
| b) Subsidised only if prescribed for urethral or cervica | l administration and | d the prescript | ion is endorsed accordingly |
| | | | |
| Topical Local Anaesthetics | | | |
| • | | | |
| condition requiring frequent injections or venepuncture. Renewal from any relevant practitioner. Approvals valid for 2 y penefiting from treatment | years where the tre | atment remair | ns appropriate and the patient is |
| Renewal from any relevant practitioner. Approvals valid for 2 poenefiting from treatment. | bove – Retail phar 5.40 | macy 5 g OP | ✓ LMX4 |
| Renewal from any relevant practitioner. Approvals valid for 2 penefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a | bove – Retail phar | macy | |
| Renewal from any relevant practitioner. Approvals valid for 2 yo penefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phan 5.40 27.00 | macy 5 g OP 30 g OP | ✓ LMX4 ✓ LMX4 |
| Renewal from any relevant practitioner. Approvals valid for 2 young from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phan 5.40 27.00 thority see SA0906 | macy 5 g OP 30 g OP above – Reta | LMX4 LMX4 iil pharmacy |
| Renewal from any relevant practitioner. Approvals valid for 2 y penefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 y enefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP above – Reta | LMX4 LMX4 iil pharmacy |
| Renewal from any relevant practitioner. Approvals valid for 2 years IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Au Crm 2.5% with prilocaine 2.5% Crm 2.5% with prilocaine 2.5% (5 g tubes) | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Itenewal from any relevant practitioner. Approvals valid for 2 yenefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Au Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Au Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 your predicting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 years benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Au Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 page 111 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 | ✓ LMX4 ✓ LMX4 ✓ LMX4 will pharmacy ✓ EMLA ✓ EMLA ✓ EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autor 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN * Tab dispersible 300 mg – Up to 30 tab available on a PSC | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 page 111 | macy 5 g OP 30 g OP 5 above – Reta 30 g OP | LMX4 LMX4 LMX4 ill pharmacy EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 y benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phan 5.40 27.00 thority see SA0906 45.00 page 111 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 | ✓ LMX4 ✓ LMX4 ✓ LMX4 will pharmacy ✓ EMLA ✓ EMLA ✓ EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 year genefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5% Crm 2.5% with prilocaine 2.5% Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics SPIRIN ≰ Tab dispersible 300 mg – Up to 30 tab available on a PSC | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 | LMX4 LMX4 LMX4 MX4 MLA EMLA EMLA EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 yean benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5% IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5% Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics NSPIRIN ★ Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 | LMX4 LMX4 LMX4 MX4 MX4 EMLA EMLA EMLA EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 y benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Au Crm 2.5% with prilocaine 2.5% Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN * Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or accordingly. | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 | LMX4 LMX4 LMX4 MX4 MLA EMLA EMLA EMLA |
| Renewal from any relevant practitioner. Approvals valid for 2 years benefiting from treatment. LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority see SA0906 a Crm 2.5% with prilocaine 2.5%. LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Autority See SA0906 a Crm 2.5% with prilocaine 2.5%. Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN * Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral 11.95 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 100 1 neuropathy a 45 g OP | LMX4 LMX4 LMX4 MX4 MLA EMLA EMLA EMLA Ethics Aspirin nd the prescription is endorsed <u>Zostrix HP</u> |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN ★ Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or accordingly. | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 | LMX4 LMX4 LMX4 LMX4 MLA EMLA EMLA EMLA EMLA Ethics Aspirin and the prescription is endorsed <u>Zostrix HP</u> Rugby Capsaicin |
| Renewal from any relevant practitioner. Approvals valid for 2 y benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral 11.95 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 100 1 neuropathy a 45 g OP | LMX4 LMX4 LMX4 Image: LMX4 EMLA EMLA EMLA EMLA EMLA Ethics Aspirin Ind the prescription is endorsed <u>Zostrix HP</u> Rugby Capsaicin Topical |
| Renewal from any relevant practitioner. Approvals valid for 2 y benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4% | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral 11.95 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 100 1 neuropathy a 45 g OP | LMX4 LMX4 LMX4 LMX4 MLA EMLA EMLA EMLA EMLA Ethics Aspirin nd the prescription is endorsed <u>Zostrix HP</u> Rugby Capsaicin |
| Renewal from any relevant practitioner. Approvals valid for 2 yean benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5%. Crm 2.5% with prilocaine 2.5% (5 g tubes) Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics SPIRIN ★ Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or accordingly. Crm 0.075%. | bove – Retail phar 5.40 27.00 thority see SA0906 45.00 page 111 04.50 diabetic peripheral 11.95 | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 100 1 neuropathy a 45 g OP | LMX4 LMX4 LMX4 Image: LMX4 Image: LMX4 EMLA EmmLA E |
| Renewal from any relevant practitioner. Approvals valid for 2 year benefiting from treatment. IDOCAINE [LIGNOCAINE] – Special Authority see SA0906 a Crm 4%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. IDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 a Crm 2.5% with prilocaine 2.5%. Analgesics For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, Non-opioid Analgesics ASPIRIN ★ Tab dispersible 300 mg – Up to 30 tab available on a PSC CAPSAICIN – Subsidy by endorsement Subsidised only if prescribed for post-herpetic neuralgia or accordingly. | bove – Retail phar | macy 5 g OP 30 g OP 6 above – Reta 30 g OP 5 100 100 1 neuropathy a 45 g OP | LMX4 LMX4 LMX4 Image: LMX4 Image: LMX4 EMLA EmmLA E |

| | Subsidy | | Fully Brand or |
|--|--|------------------|--------------------------------------|
| | (Manufacturer's P | rice) Subs | sidised Generic |
| | \$ | Per | Manufacturer |
| PARACETAMOL | | | |
| Tab 500 mg - blister pack | 19 75 | 1,000 | Pacimol |
| a) Maximum of 300 tab per prescription; can be waiv | | | |
| b) Up to 30 tab available on a PSO | red by endorsement | | |
| , , | | | |
| C) | a ia availabla far nai | tionto with long | torm conditions who require |
| Subsidy by endorsement for higher quantitie regular daily dosing for one month or greate | | | |
| annotate the prescription as endorsed where | | | 0, |
| Maximum of 100 tab per dispensing for non- | | | |
| (for non-endorsed patients), then dispense i | | | |
| | in repeat dispensings | | g 100 tab per dispensing. |
| Tab 500 mg - bottle pack – Maximum of 300 tab per | 17.00 | 1 000 | Maumad |
| prescription; can be waived by endorsement | 17.92 | 1,000 | ✓ <u>Noumed</u> |
| | | | Paracetamol |
| Subsidy by endorsement for higher quantities is | | • | |
| daily dosing for one month or greater, and the p | | | |
| prescription as endorsed where dispensing hist | , ,, , , , , , , , , , , , , , , , , , , | | |
| Maximum of 100 tab per dispensing for non-end | | | |
| non-endorsed patients), then dispense in repea | t dispensings not ex | ceeding 100 ta | ab per dispensing. |
| | | | |
| * Oral liq 120 mg per 5 ml | 5.45 | 1,000 ml | Paracare |
| a) Up to 200 ml available on a PSO | | | |
| b) Not in combination | | | |
| * Oral líq 250 mg per 5 ml | 6.25 | 1,000 ml | Paracare Double |
| | | | Strength |
| a) Up to 100 ml available on a PSO | | | v |
| b) Not in combination | | | |
| * Suppos 125 mg | 3.59 | 10 | ✓ Gacet |
| * Suppos 250 mg | | 10 | ✓ Gacet |
| * Suppos 500 mg | | 50 | ✓ Gacet |
| | | | |
| Opioid Analgesics | | | |
| opiola Analgeoloo | | | |
| CODEINE PHOSPHATE - Safety medicine; prescriber may of | determine dispensing | a frequency | |
| Tab 15 mg | | 100 | 🗸 PSM |
| Tab 30 mg | 7.45 | 100 | ✓ PSM |
| Tab 60 mg | | 100 | ✓ PSM |
| | | | |
| | 0.00 | 60 | A DHC Continue |
| Tab long-acting 60 mg | 8.00 | 60 | DHC Continus |
| FENTANYL | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing | g frequency | | |
| Inj 50 mcg per ml, 2 ml ampoule | 3.75 | 10 | Boucher and Muir |
| Inj 50 mcg per ml, 10 ml ampoule | 9.41 | 10 | Boucher and Muir |
| Patch 12.5 mcg per hour | 6.99 | 5 | Fentanyl Sandoz |
| Patch 25 mcg per hour | 7.99 | 5 | Fentanyl Sandoz |
| Patch 50 mcg per hour | 9.49 | 5 | Fentanyl Sandoz |
| Patch 75 mcg per hour | | 5 | Fentanyl Sandoz |
| Patch 100 mcg per hour | | 5 | ✓ Fentanyl Sandoz |
| | | - | |

| | Subsidy (Manufacturer's P \$ | rice) Sub Per | Fully Brand or sidised Generic Manufacturer |
|--|------------------------------------|------------------|--|
| METHADONE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing | requency | | |
| d) Extemporaneously compounded methadone will only be | e reimbursed at the | e rate of the ch | neapest form available |
| (methadone powder, not methadone tablets). | | | |
| e) For methadone hydrochloride oral liquid refer Standard | Formulae, page 24 | 44 | |
| Tab 5 mg | 1.40 | 10 | Methatabs |
| Oral liq 2 mg per ml | | 200 ml | Biodone |
| Oral liq 5 mg per ml | 6.40 | 200 ml | Biodone Forte |
| Oral liq 10 mg per ml | 7.50 | 200 ml | Biodone Extra Forte |
| Inj 10 mg per ml, 1 ml | 61.00 | 10 | 🖌 AFT |
| MORPHINE HYDROCHLORIDE | | | |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing to | requency | | |
| Oral lig 1 mg per ml | | 200 ml | RA-Morph |
| Oral lig 2 mg per ml | | 200 ml | ✓ RA-Morph |
| Oral lig 5 mg per ml | | 200 ml | ✓ Ordine S29 |
| | | 200 111 | ✓ RA-Morph |
| Oral lig 10 mg per ml | 27 74 | 200 ml | ✓ Ordine S29 |
| | | 200 111 | ✓ RA-Morph |
| IORPHINE SULPHATE | | | • |
| a) Only on a controlled drug form | | | |
| b) No patient co-payment payable | | | |
| c) Safety medicine; prescriber may determine dispensing | requency | | |
| Tab immediate-release 10 mg | | 10 | Sevredol |
| Tab immediate-release 20 mg | 5.52 | 10 | Sevredol |
| Cap long-acting 10 mg | 2.05 | 10 | ✓ <u>m-Eslon</u> |
| Cap long-acting 30 mg | 3.00 | 10 | ✓ <u>m-Eslon</u> |
| Cap long-acting 60 mg | 6.12 | 10 | ✓ <u>m-Eslon</u> |
| Cap long-acting 100 mg | | 10 | ✓ <u>m-Eslon</u> |
| Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a l | PSO6.99 | 5 | DBL Morphine |
| | | _ | Sulphate |
| Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a | PSO 5.61 | 5 | DBL Morphine |
| | | | Sulphate |
| Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a | PSO 7.08 | 5 | DBL Morphine Sulphate |
| lui 00 ma navuel duel amenada Un to Elizi avaitable ava | DCO 7 00 | - | Sulphate |
| Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a | PSU 7.28 | 5 | DBL Morphine Sulphoto |
| | | | Sulphate |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|-------|---------------------|------------------|
| YCODONE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing free | uency | | | |
| Tab controlled-release 5 mg. | | 20 | 1 | Oxycodone Sandoz |
| ů | 3.01 | 28 | 1 | Oxycodone Sandoz |
| | | | | S29 S29 |
| Oxycodone Sandoz to be Principal Supply on 1 June 2022 | > | | | |
| Tab controlled-release 10 mg. | | 20 | 1 | Oxycodone Sandoz |
| | 3.23 | 30 | | Oxycodone Sandoz |
| | 0.20 | | | S29 S29 |
| | 5.38 | 50 | 1 | Oxycodone Sandoz |
| | 5.50 | 50 | • | S29 S29 |
| | 10.75 | 100 | | |
| | 10.75 | 100 | v | Oxycodone Sandoz |
| | | | - | S29 S29 |
| | 11.50 | 28 | ~ | OxyContin |
| Oxycodone Sandoz to be Principal Supply on 1 June 2022 | | | - | |
| Tab controlled-release 20 mg | | 20 | | Oxycodone Sandoz |
| | 5.38 | 50 | ~ | Oxycodone Sandoz |
| | | | | S29 S29 |
| | 10.75 | 100 | 1 | Oxycodone Sandoz |
| | | | | S29 S29 |
| | 13.25 | 28 | 1 | OxyContin |
| Oxycodone Sandoz to be Principal Supply on 1 June 2022 | 2 | | | |
| Tab controlled-release 40 mg | | 20 | 1 | Oxycodone Sandoz |
| Oxycodone Sandoz to be Principal Supply on 1 June 2022 | | | | |
| Tab controlled-release 80 mg | | 20 | 1 | Oxycodone Sandoz |
| Oxycodone Sandoz to be Principal Supply on 1 June 2022 | | | | • |
| Cap immediate-release 5 mg | | 20 | 1 | OxyNorm |
| Cap immediate-release 10 mg | | 20 | ✓ | OxyNorm |
| Cap immediate-release 20 mg | | 20 | ✓ | OxyNorm |
| Oral liq 5 mg per 5 ml | | 250 n | nl 🖌 | OxyNorm |
| Inj 10 mg per ml, 1 ml ampoule | | 5 | ✓ | Hameln |
| · | 7.28 | | ✓ | OxyNorm |
| Hameln to be Principal Supply on 1 July 2022 | | | | |
| Inj 10 mg per ml, 2 ml ampoule | 11.49 | 5 | 1 | Hameln |
| · | 14.36 | | ✓ | OxyNorm |
| Hameln to be Principal Supply on 1 July 2022 | | | | |
| Inj 50 mg per ml, 1 ml ampoule | | 5 | 1 | Hameln |
| | 30.60 | | ✓ | OxyNorm |

| | | | - " | |
|---|-----------------------------------|------------|------------------|---|
| | Subsidy (Manufacturer's Price) |) Sub | Fully sidised | Brand or Generic |
| | (Manulaciulei S Flice) | Per | | Manufacturer |
| Hameln to be Principal Supply on 1 July 2022 | | | | |
| Oxycodone Sandoz S29 © Tab controlled-release 5 mg to b | e delisted 1 June 202 | 22) | | |
| Oxycodone Sandoz S29 1 Tab controlled-release 10 mg to | | | | |
| Oxycodone Sandoz S29 (S29) Tab controlled-release 10 mg to | | | | |
| Oxycodone Sandoz S29 s29 Tab controlled-release 10 mg to | | ' | | |
| OxyContin Tab controlled-release 10 mg to be delisted 1 June 2 | | //// | | |
| Oxycodone Sandoz S29 S29 Tab controlled-release 20 mg to | be delisted 1 June 20 |)22) | | |
| (Oxycodone Sandoz S29 S29) Tab controlled-release 20 mg to | be delisted 1 June 20 |)22) | | |
| OxyContin Tab controlled-release 20 mg to be delisted 1 June 2 | | | | |
| OxyNorm Inj 10 mg per ml, 1 ml ampoule to be delisted 1 July 2 | | | | |
| OxyNorm Inj 10 mg per ml, 2 ml ampoule to be delisted 1 July 2 | | | | |
| (OxyNorm Inj 50 mg per ml, 1 ml ampoule to be delisted 1 July 2 | 2022) | | | |
| PARACETAMOL WITH CODEINE – Safety medicine; prescribe | | ensing fre | | |
| * Tab paracetamol 500 mg with codeine phosphate 8 mg | | 1,000 | | racetamol + |
| | | | | Codeine (Relieve) |
| PETHIDINE HYDROCHLORIDE | | | | |
| a) Only on a controlled drug form | | | | |
| b) No patient co-payment payable | | | | |
| c) Safety medicine; prescriber may determine dispensing fr | requency | | | |
| Tab 50 mg | | 10 | ✓ PS | |
| Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a | PSO29.88 | 5 | | 3L Pethidine |
| | | _ | | Hydrochloride |
| Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a | PSO 30.72 | 5 | | 3L Pethidine |
| | | | | Hydrochloride |
| TRAMADOL HYDROCHLORIDE | | | | |
| Tab sustained-release 100 mg | | 20 | | amal SR 100 |
| Tab sustained-release 150 mg | | 20 | | amal SR 150 |
| Tab sustained-release 200 mg | | 20 | | amal SR 200 |
| Cap 50 mg | 2.80 | 100 | ✓ <u>AI</u> | row-Tramadol |
| Antidepressants | | | | |
| Annaoprocounto | | | | |
| Cyclic and Related Agents | | | | |
| AMITRIPTYLINE - Safety medicine; prescriber may determine | dispensing frequency | , | | |
| Tab 10 mg | | 100 | 🗸 🗸 | row-Amitriptyline |
| Tab 25 mg | 1.51 | 100 | 🖌 🗸 | row-Amitriptyline |
| Tab 50 mg | 2.51 | 100 | ✓ <u>A</u> | row-Amitriptyline |
| CLOMIPRAMINE HYDROCHLORIDE | | | | |
| a) Brand switch fee payable (Pharmacode 2630915) - see | | | | |
| | page 242 for details | | | |
| b) Safety medicine; prescriber may determine dispensing fr | requency | | | |
| | requency | 30 | | <u>omipramine Teva</u> omipramine Teva |

| | Subsidy | | Fully | Brand or |
|---|-------------------------------|-------------|---------------|--------------------------|
| | (Manufacturer's Price \$ |) Su Per | bsidised ✓ | Generic Manufacturer |
| DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Suba | sidy by endorsement | | | |
| a) Safety medicine; prescriber may determine disp | | | | |
| b) Subsidy by endorsement – Subsidised for patie | | | | |
| 2019 and the prescription is endorsed according | | te the pre | scription | as endorsed where there |
| exists a record of prior dispensing of dosulepin Tab 75 mg | | 30 | . г | Dosulepin Mylan |
| Cap 25 mg | | 50 | | Dosulepin |
| oop _og | | | - | Mylan S29 |
| MIPRAMINE HYDROCHLORIDE – Safety medicine; p | prescriber may determine disp | ensing fre | equency | • |
| Tab 10 mg | 5.48 | 50 | | ofranil |
| | 10.96 | 100 | | ofranil |
| Tab 25 mg | | 50 | | ofranil |
| VORTRIPTYLINE HYDROCHLORIDE – Safety medici | | | | |
| Tab 10 mg | | 100 | | lorpress |
| Tab 25 mg | 5.98 | 180 | ✓ № | lorpress |
| Monoamine-Oxidase Inhibitors (MAOIs) | - Non Selective | | | |
| FRANYLCYPROMINE SULPHATE | | | | |
| Tab 10 mg | | 28 | 🗸 F | Parnate S29 S29 |
| | 22.94 | 50 | 🗸 F | Parnate |
| | 45.88 | 100 | 🗸 F | Parnate S29 S29 |
| | 96.00 | | 🗸 F | Parnate S29 S29 |
| Monoamine-Oxidase Type A Inhibitors | | | | |
| MOCLOBEMIDE | | | | |
| * Tab 150 mg | 11.80 | 60 | 🗸 🖌 | Aurorix |
| * Tab 300 mg | | 60 | ✓ <u> </u> | Aurorix |
| Selective Serotonin Reuptake Inhibitors | | | | |
| CITALOPRAM HYDROBROMIDE | | | | |
| * Tab 20 mg | 1.91 | 84 | ✓ <u>F</u> | SM Citalopram |
| ESCITALOPRAM | | | | |
| * Tab 10 mg | 1.07 | 28 | ✓ <u>E</u> | scitalopram |
| | | | | (Ethics) |
| * Tab 20 mg | 1.92 | 28 | ✓ <u>E</u> | Escitalopram (Ethica) |
| | | | | (Ethics) |
| FLUOXETINE HYDROCHLORIDE * Tab dispersible 20 mg, scored – Subsidy by endor | reamont 1.98 | 30 | / = | luox |
| Subsidised by endorsement | Joint I 1.30 | 50 | • ٢ | IUUA |
| When prescribed for a patient who cannot | t swallow whole tablets or ca | osules and | d the pre | scription is endorsed |
| accordingly; or | | | | |
| When prescribed in a daily dose that is n endorsed. Note: Tablets should be com | | | | |
| Cap 20 mg | 2 01 | 84 | . | luox |
| | | 04 | • ٢ | |
| PAROXETINE 卷 Tab 20 mg | 3 61 | 90 | . / I | .oxamine |
| * 100 20 mg | | 30 | • ⊑ | |
| | | | | |
| 4 · · · · · · · · | | | | |

| | Subsidy | | Fully | Brand or |
|--|------------------------------|--------|-----------------------|--------------|
| | (Manufacturer's Price) | | Subsidised | |
| | (Manalaotaror o F 100) \$ | Per | Cubbiai000 | Manufacturer |
| | Ψ | 1.01 | • | Manufacturei |
| SERTRALINE | | | | |
| * Tab 50 mg | 0.02 | 30 | 1 | Setrona |
| * Tab 50 mg | 0.92 | 30 | | |
| | | | ~ | Setrona AU |
| * Tab 100 mg | 1 61 | 30 | ✓ | Setrona |
| | | 00 | - | octiona |
| · · · · · | | | | |
| Other Antidepressants | | | | |
| | | | | |
| MIRTAZAPINE | | | | |
| | 0.00 | ~~ | | N |
| Tab 30 mg | 2.60 | 28 | ~ | Noumed |
| Tab 45 mg | 3.45 | 28 | ✓ | Noumed |
| · | | | | |
| VENLAFAXINE | | | | |
| * Cap 37.5 mg | 6.38 | 84 | ✓ | Enlafax XR |
| | | | - | |
| * Cap 75 mg | | 84 | - | Enlafax XR |
| * Cap 150 mg | | 84 | ~ | Enlafax XR |
| • | | | | |
| Antionilonov Drugo | | | | |
| Antiepilepsy Drugs | | | | |
| | | | | |
| Agents for Control of Status Epilepticus | | | | |
| Agents for Control of Status Ephepticus | | | | |
| | | | | |
| DIAZEPAM – Safety medicine; prescriber may determine disp | ensing frequency | | | |
| Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement | 23.66 | 5 | ✓ | Hospira |
| | 20.00 | Ũ | | |
| a) Up to 5 inj available on a PSO | | | | |
| b) Only on a PSO | | | | |
| c) PSO must be endorsed "not for anaesthetic proce | duros" | | | |
| | | _ | | |
| Rectal tubes 5 mg – Up to 5 tube available on a PSO | | 5 | ~ | Stesolid |
| | | | | |
| PHENYTOIN SODIUM | | | | |
| * Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on | а | | | |
| | | E | | Heenire |
| PSO | | 5 | • | Hospira |
| * Inj 50 mg per ml, 5 ml ampoule - Up to 5 inj available on | а | | | |
| PSO | | 5 | 1 | Hospira |
| 1.50 | | 5 | • | поэрна |
| | | | | |
| Control of Epilepsy | | | | |
| | | | | |
| | | | | |
| CARBAMAZEPINE | | | | |
| * Tab 200 mg | 14.53 | 100 | ✓ | Tegretol |
| * Tab long-acting 200 mg | | 100 | - | Tegretol CR |
| 5 5 5 | | | - | • |
| * Tab 400 mg | | 100 | | Tegretol |
| * Tab long-acting 400 mg | | 100 | ✓ | Tegretol CR |
| | | 250 n | | Tegretol |
| * Oral liq 20 mg per ml | | 200 II | | regretor |
| CLOBAZAM - Safety medicine; prescriber may determine dis | pensing frequency | | | |
| 3 7 1 3 | | 50 | ./ | Frigium |
| Tab 10 mg | 9.12 | 50 | • | Frisium |
| CLONAZEPAM - Safety medicine; prescriber may determine | dispensing frequency | | | |
| | | | | Discotal! |
| Oral drops 2.5 mg per ml | | 0 ml (| א א | Rivotril |
| ETHOSUXIMIDE | | | | |
| | | | - | |
| Cap 250 mg | | 100 | ✓ | Zarontin |
| Oral lig 250 mg per 5 ml | | 200 m | nl 🗸 | Zarontin |
| 1 01 | | | 2 | |
| GABAPENTIN | | | | |
| Note: Not subsidised in combination with subsidised prec | ahalin | | | |
| | | | | |
| * Cap 100 mg | | 100 | | Nupentin |
| * Cap 300 mg | | 100 | ✓ | Nupentin |
| | | 100 | | Nupentin |
| * Cap 400 mg | 10.20 | 100 | v | nupentin |
| | | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully bsidised | Brand or Generic Manufacturer | |
|--|---|-----------|-------------------|-------------------------------------|--|
| LACOSAMIDE – Special Authority see SA1125 below – Retail | pharmacy | | | | |
| ▲ Tab 50 mg | | 14 | 🗸 V | impat | |
| ▲ Tab 100 mg | | 14 | 🗸 V | impat | |
| • | 200.24 | 56 | 🗸 V | impat | |
| ▲ Tab 150 mg | 75.10 | 14 | 🗸 V | impat | |
| v | 300.40 | 56 | 🗸 V | impat | |
| ▲ Tab 200 mg | 400.55 | 56 | 🗸 V | impat | |

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

| ▲ ▲ * | Tab dispersible 2 mg 55.00 Tab dispersible 5 mg 50.00 Tab dispersible 25 mg 2.76 | 30 30 56 | ~ | Lamictal Lamictal Logem |
|-------------|--|----------------|---|-------------------------------|
| * | Tab dispersible 50 mg | 56 | | Logem |
| * | Tab dispersible 100 mg4.40 | 56 | | Logem |
| LE | VETIBACETAM | | | |
| | Tab 250 mg | 60 | 1 | Everet |
| | Tab 500 mg | 60 | 1 | Everet |
| | Tab 750 mg14.39 | 60 | 1 | Everet |
| | Tab 1,000 mg | 60 | 1 | Everet |
| | Oral liq 100 mg per ml44.78 | 300 ml OP | 1 | Levetiracetam-AFT |
| PH | ENOBARBITONE For phenobarbitone oral liquid refer Standard Formulae, page 244 | | | |
| * | Tab 15 mg40.00 | 500 | 1 | PSM |
| * | Tab 30 mg40.00 | 500 | 1 | PSM |
| PH | ENYTOIN SODIUM | | | |
| * | Tab 50 mg75.00 | 200 | 1 | Dilantin Infatab |
| | Cap 30 mg74.00 | 200 | 1 | Dilantin |
| | Cap 100 mg | 200 | 1 | Dilantin |
| * | Oral liq 30 mg per 5 ml22.03 | 500 ml | 1 | Dilantin |
| PF | EGABALIN | | | |
| | Note: Not subsidised in combination with subsidised gabapentin | | | |
| | Cap 25 mg2.25 | 56 | 1 | Pregabalin Pfizer |
| * | Cap 75 mg2.65 | 56 | 1 | Pregabalin Pfizer |
| | Cap 150 mg4.01 | 56 | 1 | Lyrica |
| | | | 1 | Pregabalin Pfizer |
| | Cap 300 mg7.38 | 56 | 1 | Pregabalin Pfizer |
| | | | | |

| | Subsidy | | Fully | Brand or |
|---|----------------------|-------|------------|-------------------|
| | Manufacturer's Price |) | Subsidised | Generic |
| | \$ | Per | 1 | Manufacturer |
| PRIMIDONE | | | | |
| * Tab 250 mg | 37.35 | 100 | ✓ | Apo-Primidone |
| SODIUM VALPROATE | | | | |
| Tab 100 mg | 13.65 | 100 | 1 | Epilim Crushable |
| Tab 200 mg EC | 27.44 | 100 | ✓ | Epilim |
| Tab 500 mg EC | | 100 | ✓ | Epilim |
| * Oral liq 200 mg per 5 ml | 20.48 | 300 m | nl 🗸 | Epilim S/F Liquid |
| | | | 1 | Epilim Syrup |
| * Inj 100 mg per ml, 4 ml | 41.50 | 1 | 1 | Epilim IV |
| STIRIPENTOL - Special Authority see SA1330 below - Retail pha | irmacy | | | |
| Cap 250 mg | 509.29 | 60 | 1 | Diacomit S29 |
| Powder for oral liq 250 mg sachet | 509.29 | 60 | ✓ | Diacomit S29 |
| | | | | |

SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

| Tab 25 mg | | 60 | Arrow-Topiramate |
|--|--------------|-----|--|
| Ŭ | | | Topiramate Actavis |
| | 26.04 | | Topamax |
| Tab 50 mg | | 60 | Arrow-Topiramate |
| J. J | | | Topiramate Actavis |
| | 44.26 | | Topamax |
| Tab 100 mg | | 60 | Arrow-Topiramate |
| - | | | Topiramate Actavis |
| | 75.25 | | Topamax |
| Tab 200 mg | | 60 | Arrow-Topiramate |
| 0 | | | Topiramate Actavis |
| | 129.85 | | Topamax |
| Sprinkle cap 15 mg | 20.84 | 60 | Topamax |
| Sprinkle cap 25 mg | | 60 | Topamax |
| /IGABATRIN - Special Authority see SA2088 below - Re | ail pharmacy | | - |
| Tab 500 mg | | 100 | Sabril |

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:

1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages. **Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 111

Acute Migraine Treatment

| RIZATRIPTAN | | |
|--|------|-------------------------------------|
| Tab orodispersible 10 mg3.65 | 30 | <u>Rizamelt</u> |
| SUMATRIPTAN | | |
| Tab 50 mg14.41 | 90 | Sumagran |
| Tab 100 mg | 90 | Sumagran |
| Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per | 0.00 | |
| prescription | 2 OP | Imigran |
| Prophylaxis of Migraine | | |
| For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50 | | |
| PIZOTIFEN | | |
| * Tab 500 mcg23.21 | 100 | Sandomigran |
| | 100 | • Cundonnigram |
| Antinausea and Vertigo Agents | | |
| For Antispasmodics refer to ALIMENTARY TRACT, page 8 | | |
| APREPITANT - Special Authority see SA0987 on the next page - Retail pharmacy | | |
| Cap 2 × 80 mg and 1 × 125 mg | 3 OP | Emend Tri-Pack |
| | | |

| | Subsidy | | Fully | Brand or |
|---|--|---------------------|---------------------------|------------------------------|
| | (Manufacturer's Price) \$ | Per | Subsidised | Generic Manufacturer |
| ► SA0987 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid emetogenic chemotherapy and/or anthracycline-based chemother Renewal from any relevant practitioner. Approvals valid for 12 mc chemotherapy and/or anthracycline-based chemotherapy for the tr BETAHISTINE DIHYDROCHLORIDE | apy for the treatmen onths where the patie | t of ma ent is u | alignancy. | 0 0 0 7 |
| * Tab 16 mg | | 84 | 🗸 V | ergo 16 |
| | 4.62 | 100 | ✓ <u>s</u> | • |
| (Vergo 16 Tab 16 mg to be delisted 1 July 2022) | | | | |
| CYCLIZINE HYDROCHLORIDE | | | | |
| Tab 50 mg | 0.49 | 10 | ✓ <u>N</u> | lausicalm |
| CYCLIZINE LACTATE | | | | |
| Inj 50 mg per ml, 1 ml | 21.53 | 10 | ✓ H | lameln |
| DOMPERIDONE | | | | |
| * Tab 10 mg | 2.85 | 100 | ✓ <u>P</u> | harmacy Health |
| HYOSCINE HYDROBROMIDE | | | | |
| * Inj 400 mcg per ml, 1 ml ampoule | 93.00 | 10 | 🗸 N | lartindale S29 |
| Patch 1.5 mg – Special Authority see SA1998 below – Retail pharmacy | 14.11 | 2 | ✓ s | copoderm TTS |
| SA1998 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid Either: Control of intractable nausea, vomiting, or inability to swalld where the patient cannot tolerate or does not adequately re Control of clozapine-induced hypersalivation where trials of ineffective. | ow saliva in the treat espond to oral anti-na | ment o ausea | of malignan agents; or | cy or chronic disease |
| Renewal from any relevant practitioner. Approvals valid for 1 year benefiting from treatment. | r where the treatmer | nt rema | ains approp | priate and the patient is |
| METOCLOPRAMIDE HYDROCHLORIDE | | | | |
| * Tab 10 mg – Up to 30 tab available on a PSO | 1.30 | 100 | ✓ <u>N</u> | letoclopramide Actavis 10 |
| * Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PS ONDANSETRON | O9.50 | 10 | ✓ <u>P</u> | fizer |
| * Tab 4 mg | | 50 | | nrex |
| * Tab disp 4 mg – Up to 10 tab available on a PSO | 0.76 | 10 | ✓ <u>C</u> | Ondansetron ODT-DRLA |
| * Tab 8 mg | | 50 | | nrex |
| * Tab disp 8 mg – Up to 10 tab available on a PSO | 1.13 | 10 | ✓ <u>0</u> | Indansetron |

| PROCHL | ORPERAZINE |
|--------|------------|
|--------|------------|

| * | Tab 3 mg buccal | 5.97 | 50 | |
|---|---|---------|-----|------------------------------|
| | Ĵ | (30.00) | | Buccastem |
| * | Tab 5 mg - Up to 30 tab available on a PSO | | 250 | Nausafix |
| * | Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO | 25.81 | 10 | ✓ Stemetil |

ODT-DRLA

| | Subsidy | | Fully | Brand or |
|--|------------------------|---------|-------------------------|-----------------------------|
| | (Manufacturer's Price) | | sidised | Generic |
| | \$ | Per | 1 | Manufacturer |
| Antipsychotics | | | | |
| | | | | |
| General | | | | |
| MISULPRIDE - Safety medicine; prescriber may determine dis | | | | |
| Tab 100 mg | | 30 | | Sulprix |
| | 17.16 | 100 | v | Amisulpride |
| | | | | Mylan S29 |
| Tab 200 mg | 14.96 | 60 | ✓ § | Sulprix |
| Tab 400 mg | | 60 | | Sulprix |
| misulpride Mylan 329 Tab 100 mg to be delisted 1 August 20. | | | - | _ - |
| RIPIPRAZOLE – Safety medicine; prescriber may determine d | ispensing frequency | | | |
| Tab 5 mg | | 30 | I | Aripiprazole Sandoz |
| Tab 10 mg | | 30 | - | Aripiprazole Sandoz |
| Tab 15 mg | | 30 | | Aripiprazole Sandoz |
| Tab 20 mg | | 30 | | Aripiprazole Sandoz |
| Tab 30 mg | | 30 | | Aripiprazole Sandoz |
| 0 | | | | •• |
| ILORPROMAZINE HYDROCHLORIDE – Safety medicine; pro | | | | |
| Tab 10 mg – Up to 30 tab available on a PSO | | 100 | | argactil |
| Tab 25 mg – Up to 30 tab available on a PSO | | 100 | | Largactil |
| Tab 100 mg – Up to 30 tab available on a PSO | | 100 | | Largactil |
| Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO | | 10 | ✓ I | Largactil |
| LOZAPINE – Hospital pharmacy [HP4] | | | | |
| Safety medicine; prescriber may determine dispensing frequ | encv | | | |
| Tab 25 mg | | 50 | ✓ (| Clozaril |
| · «» =• …g | 6.69 | | | Clopine |
| | 11.36 | 100 | - | Clozaril |
| | 13.37 | 100 | | Clopine |
| Tab 50 mg | | 50 | | |
| Tab 50 mg | | 50 | | Clopine |
| T 1 (10) | 17.33 | 100 | | Clopine |
| Tab 100 mg | | 50 | | Clozaril |
| | 17.33 | | | Clopine |
| | 29.45 | 100 | | Clozaril |
| | 34.65 | | | Clopine |
| Tab 200 mg | | 50 | ✓ (| Clopine |
| | 69.30 | 100 | ✓ (| Clopine |
| Suspension 50 mg per ml | 67.62 | 100 ml | • | Versacloz |
| ALOPERIDOL – Safety medicine; prescriber may determine di | spensing frequency | | | |
| Tab 500 mcg - Up to 30 tab available on a PSO | | 100 | ✓ 9 | Serenace |
| Tab 1.5 mg – Up to 30 tab available on a PSO | | 100 | | Serenace |
| Tab 5 mg – Up to 30 tab available on a PSO | | 50 | | Serenace |
| | 29.72 | 100 | - | Serenace |
| Oral liq 2 mg per ml – Up to 200 ml available on a PSO | | 100 ml | - | Serenace |
| Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO | | 100 111 | | Serenace |
| VOMEPROMAZINE – Safety medicine; prescriber may deter | | MARCY | - | |
| Tab 25 mg (33.8 mg as a maleate) | 16 10 | | | Nozinan (Swice) |
| | | 100 | • I | Nozinan (Swiss) |
| Tab 25 mg as a maleate | | 100 | | Nozinan Nazinan (Ouriss) |
| | /1 76 | 100 | ✓ [| Nozinan (Swiss) |
| Tab 100 mg (135 mg as a maleate) Tab 100 mg as a maleate | | 100 | | Nozinan |

| | Subsidy | | Fully | Brand or |
|--|------------------------|---------|------------|--------------------|
| | (Manufacturer's Price) | | Subsidised | |
| | \$ | Per | 1 | Manufacturer |
| LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine; | prescriber may detern | nine d | spensing | frequency |
| Inj 25 mg per ml, 1 ml ampoule | | 10 | | Nozinan |
| LITHIUM CARBONATE - Safety medicine; prescriber may dete | | uency | , | |
| Tab long-acting 400 mg | | 100 | | Priadel |
| Cap 250 mg | | 100 | - | Douglas |
| OLANZAPINE – Safety medicine; prescriber may determine dis | | | | 2019.00 |
| Tab 2.5 mg | | 28 | 1 | Zypine |
| Tab 5 mg | | 28 | | Zypine |
| Tab orodispersible 5 mg | | 28 | | Zypine ODT |
| Tab 10 mg | | 28 | | Zypine |
| Tab orodispersible 10 mg | | 28 | | Zypine ODT |
| | | 20 | • | Zypine OD1 |
| PERICYAZINE – Safety medicine; prescriber may determine di | | | | |
| Tab 2.5 mg | | 84 | | Neulactil |
| | 12.49 | 100 | | Neulactil |
| Tab 10 mg | | 84 | | Neulactil |
| | 44.45 | 100 | ~ | Neulactil |
| QUETIAPINE – Safety medicine; prescriber may determine dis | pensing frequency | | | |
| Tab 25 mg | 2.15 | 90 | ✓ | Quetapel |
| Tab 100 mg | 5.06 | 90 | ✓ | Quetapel |
| Tab 200 mg | 8.90 | 90 | ✓ | Quetapel |
| Tab 300 mg | 12.86 | 90 | ✓ | Quetapel |
| RISPERIDONE - Safety medicine; prescriber may determine d | lispensing frequency | | | |
| Tab 0.5 mg | 1 0 1 7 | 60 | 1 | Risperidone (Teva) |
| Tab 1 mg | | 60 | 1 | Risperidone (Teva) |
| Tab 2 mg | | 60 | | Risperidone (Teva) |
| Tab 3 mg | | 60 | | Risperidone (Teva) |
| Tab 4 mg | | 60 | | Risperidone (Teva) |
| Oral liq 1 mg per ml | | 30 ml | - | Risperon |
| ZIPRASIDONE – Safety medicine; prescriber may determine d | | | | |
| Cap 20 mg | | 60 | 1 | Zusdone |
| Cap 40 mg | | 60 | | Zusdone |
| Cap 40 mg | | 60 | | Zusdone |
| Cap 80 mg | | 60 | | Zusdone |
| | | | | |
| ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; pr | , | | 0 | |
| Tab 10 mg | | 100 | ~ | Clopixol |
| Depot Injections | | | | |
| FLUPENTHIXOL DECANOATE – Safety medicine; prescriber | may determine dispon | sina fr | auonov | |
| Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 5 | | Fluanxol |
| Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO | | э 5 | | Fluanxol |
| Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO | | э 5 | | Fluanxol |
| HALOPERIDOL DECANOATE - Safety medicine: prescriber m | | - | | |

HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO......28.39 5 5

- Inj 100 mg per ml, 1 ml Up to 5 inj available on a PSO......55.90
- ✓ Haldol
- ✓ Haldol Concentrate
- ✓ Haldol
 - Decanoas S29

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|-------------------------------------|
| OLANZAPINE - Special Authority see SA1428 below - Retail ph Safety medicine; prescriber may determine dispensing freque | , | | | |
| Inj 210 mg vial | , | 1 | ✓ Z | yprexa Relprevv |
| Inj 300 mg vial | | 1 | 🗸 Z | yprexa Relprevv |
| Inj 405 mg vial | | 1 | ✓ Z | yprexa Relprevv |

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

| Inj 25 mg syringe | 1 | Invega Sustenna |
|--------------------|-------|-------------------|
| Inj 50 mg syringe | 1 | Invega Sustenna |
| Inj 75 mg syringe | 1 | Invega Sustenna |
| Inj 100 mg syringe | 1 | Invega Sustenna |
| Inj 150 mg syringe | 1 | ✓ Invega Sustenna |
| | | ga easterina |

SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

| Inj 25 mg vial | 1 | Risperdal Consta |
|------------------|---|--------------------------------------|
| Inj 37.5 mg vial | 1 | Risperdal Consta |
| Inj 50 mg vial | 1 | Risperdal Consta |

| Subsidy (Manufacturer's Price) | Subsi | Fully | Brand or Generic |
|---------------------------------------|-------|-------|---------------------|
| \$ | Per | ✓ | Manufacturer |

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or

- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency Inj 200 mg per ml, 1 ml - Up to 5 inj available on a PSO......19.80 5 Clopixol

Anxiolytics

BUSPIBONE HYDROCHLOBIDE 100 Buspirone Viatris 20.23 Orion Buspirone Viatris to be Principal Supply on 1 May 2022 100 Buspirone Viatris * 13.16 ✓ Orion Buspirone Viatris to be Principal Supply on 1 May 2022 (Orion Tab 5 mg to be delisted 1 May 2022) (Orion Tab 10 mg to be delisted 1 May 2022) CLONAZEPAM - Safety medicine: prescriber may determine dispensing frequency Paxam 100 100 Paxam Tab 2 mg 10.78 DIAZEPAM - Safety medicine: prescriber may determine dispensing frequency Tab 2 mg61.07 500 Arrow-Diazepam ✓ Arrow-Diazepam Tab 5 mg73.60 500 LORAZEPAM - Safety medicine; prescriber may determine dispensing frequency 250 Ativan Ativan 100

Multiple Sclerosis Treatments

⇒SA2051 Special Authority for Subsidy

Initial application - (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and

| Subsidy (Manufacturer's Price) | Sub | Fully sidised | Brand or Generic | |
|-----------------------------------|-----|------------------|---------------------|--|
| \$ | Per | Juiscu ✓ | Manufacturer | |

- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
- 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. **Renewal — (Multiple sclerosis)** only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2051 on the previous page - Retail pharmacy

| a) Wastage claimable | | | |
|---|--|---------------------------|--------------------------------|
| b) Note: Treatment on two or more funded multiple so | clerosis treatments simul | taneously is | s not permitted. |
| Cap 120 mg | | 14 | Tecfidera |
| Cap 240 mg | 2,000.00 | 56 | Tecfidera |
| FINGOLIMOD - Special Authority see SA2051 on the prev | <mark>/ious page</mark> – Retail phari | nacy | |
| a) Wastage claimable | | | |
| b) Note: Treatment on two or more funded multiple so | clerosis treatments simul | taneously is | s not permitted. |
| Cap 0.5 mg | 2,200.00 | 28 | Gilenya |
| GLATIRAMER ACETATE - Special Authority see SA2051 | on the previous page - | Retail pharr | nacy |
| Note: Treatment on two or more funded multiple sclere | osis treatments simultan | eously is no | t permitted. |
| Inj 40 mg prefilled syringe | 2,275.00 | 12 | Copaxone |
| INTERFERON BETA-1-ALPHA - Special Authority see SA | 2051 on the previous pa | <mark>age</mark> – Retail | pharmacy |
| Note: Treatment on two or more funded multiple sclere | osis treatments simultan | eously is no | t permitted. |
| Inj 6 million iu prefilled syringe | 1,170.00 | 4 | Avonex |
| Injection 6 million iu per 0.5 ml pen injector | 1,170.00 | 4 | Avonex Pen |
| | | | |

S29 Unapproved medicine supplied under Section 29 Sole Subsidised Supply

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer INTERFERON BETA-1-BETA - Special Authority see SA2051 on page 135 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Betaferon 15 NATALIZUMAB - Special Authority see SA2051 on page 135 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 1 Tvsabri OCRELIZUMAB - Special Authority see SA2051 on page 135 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 1 ✓ Ocrevus TERIFLUNOMIDE - Special Authority see SA2051 on page 135 - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 28 Aubagio Sedatives and Hypnotics MELATONIN - Special Authority see SA1666 below - Retail pharmacy Brand switch fee payable (Pharmacode 2634112) - see page 242 for details Tab modified-release 2 mg - No more than 5 tab per day 11.50 30 Viaisom SA1666 Special Authority for Subsidy Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*: and 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and 4 Patient is aged 18 years or under*. Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following: 1 Patient is aged 18 years or under*; and 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day. Note: Indications marked with * are unapproved indications. MIDAZOLAM - Safety medicine: prescriber may determine dispensing frequency 10 Midazolam-Baxter Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available on a PSO......17.28 10 Pfizer On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. Midazolam-Baxter Ini 5 mg per ml. 3 ml ampoule4.50 5 Ini 5 mg per ml. 3 ml plastic ampoule - Up to 5 ini available on a PSO......13.09 5 Pfizer On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only. PHENOBARBITONE SODIUM - Special Authority see SA1386 on the next page - Retail pharmacy 10 ✓ Max Health S29

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|-----|---------------------|--------------------------------------|
| SA1386 Special Authority for Subsidy nitial application from any relevant practitioner. Approval ne following criteria: Noth: | | | nless notifie | ed for applications meetir |
| For the treatment of terminal agitation that is unrespondent The applicant is part of a multidisciplinary team work | | 1 | | |
| EMAZEPAM – Safety medicine; prescriber may determine Tab 10 mg | | 25 | √ <u>i</u> | Normison |
| RIAZOLAM – Safety medicine; prescriber may determine Tab 125 mcg | | 100 | | |
| Tab 250 mcg | (9.85) 4.10 (11.20) | 100 | | Hypam Hypam |
| OPICLONE – Safety medicine; prescriber may determine Tab 7.5 mg | dispensing frequency | 500 | | Zopiclone Actavis |
| Stimulants/ADHD Treatments | | | - | |
| TOMOXETINE | | | | |
| Cap 10 mg | 18.41 | 28 | | APO- Atomoxetine S29 |
| | 107.03 | | - | <u>Generic Partners</u> Strattera |
| Cap 18 mg | | 28 | | Generic Partners |
| | 107.03 | | | Strattera |
| Cap 25 mg | | 28 | | Generic Partners |
| Cap 40 mg | | 28 | - | Generic Partners |
| 0 | 107.03 | ~~ | | Strattera |
| Cap 60 mg | | 28 | | <u>Generic Partners</u> APO- |
| Cap 80 mg | | 28 | • / | APO- Atomoxetine S29 |
| 0 | 50.40 | ~~ | | Generic Partners |
| Cap 100 mg | | 28 | • | APO- Atomoxetine S29 |
| | | | ✓ <u>(</u> | Generic Partners |
| EXAMFETAMINE SULFATE – Special Authority see SA1 a) Only on a controlled drug form | 149 below – Retail pharma | асу | | |
| b) Safety medicine; prescriber may determine dispensi | na frequency | | | |
| Tab 5 mg | | 100 | ✓] | PSM |

recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

138

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a) Only on a controlled drug form

| b) Sa | fety medicine; | prescriber m | av determine | dispensing | frequency |
|-------|----------------|--------------|--------------|------------|-----------|
|-------|----------------|--------------|--------------|------------|-----------|

| Tab immediate-release 5 mg | | 30 | Rubifen |
|-----------------------------|-------|----|--|
| Tab immediate-release 10 mg | | 30 | Ritalin |
| ů | | | Rubifen |
| Tab extended-release 18 mg | 7.75 | 30 | Methylphenidate ER Teva |
| Tab immediate-release 20 mg | 7.85 | 30 | Rubifen |
| Tab sustained-release 20 mg | | 30 | Rubifen SR |
| Tab extended-release 27 mg | 11.45 | 30 | Methylphenidate ER Teva |
| Tab extended-release 36 mg | 15.50 | 30 | Methylphenidate ER Teva |
| Tab extended-release 54 mg | 22.25 | 30 | Methylphenidate ER Teva |

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: All of the following:

| S | Subsidy | Fully | Brand or |
|----------|-----------------------|---------|--------------|
| (Manufac | acturer's Price) Subs | sidised | Generic |
| | \$ Per | ~ | Manufacturer |

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER - Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

| Tab extended-release 18 mg | | 30 | Concerta |
|----------------------------|-------|----|--------------------------------|
| Tab extended-release 27 mg | | 30 | Concerta |
| Tab extended-release 36 mg | 71.93 | 30 | Concerta |
| Tab extended-release 54 mg | | 30 | Concerta |
| Cap modified-release 10 mg | 15.60 | 30 | Ritalin LA |
| Cap modified-release 20 mg | 20.40 | 30 | Ritalin LA |
| Cap modified-release 30 mg | | 30 | Ritalin LA |
| Cap modified-release 40 mg | | 30 | Ritalin LA |

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

- 3.1 Applicant is a paediatrician or psychiatrist; or
- 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and

4 Either:

- 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
- 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hvdrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

| Tab 100 mg | | 60 | Modavigil |
|------------|--|----|-----------|
|------------|--|----|-----------|

■ SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the followina:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and

3 Either:

- 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects: or
- 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

| * Tab 5 mg | | 90 | ✓ <u>Donepezil-Rex</u> |
|---|------------------------------------|----|------------------------|
| * Tab 10 mg | 6.64 | 90 | Donepezil-Rex |
| RIVASTIGMINE - Special Authority see SA1488 on the next p | o <mark>age</mark> – Retail pharma | су | |
| Patch 4.6 mg per 24 hour | | 30 | Rivastigmine Patch |
| | | | <u>BNM 5</u> |
| Patch 9.5 mg per 24 hour | | 30 | Rivastigmine Patch |
| | | | BNM 10 |

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic |
|-----------------------------------|-----|-----------------------|---------------------|
| \$ | Per | ✓ | Manufacturer |

➡SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

a) No patient co-payment payable

| b) Safety medicine; prescriber may determine dispensing free | requency |
|---|----------|
|---|----------|

| Tab sublingual 2 mg with naloxone 0.5 mg | 28 | <u>Buprenorphine</u> |
|---|----|--|
| | | Naloxone BNM |
| Tab sublingual 8 mg with naloxone 2 mg53.12 | 28 | Buprenorphine |
| | | Naloxone BNM |

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | Ibsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from

any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

| Tab modified-release 150 mg | 0 🖌 <u>Zyban</u> |
|---|----------------------|
| DISULFIRAM Tab 200 mg236.40 10 | 00 ✓ <u>Antabuse</u> |
| NALTREXONE HYDROCHLORIDE - Special Authority see SA1408 below - Retail phan | macy |
| Tab 50 mg 133.33 30 | 0 Valtraccord |

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|---|---|-------|---------------------|--------------|
| | Ψ | 1.61 | • | Manulacturei |
| NICOTINE | | | | |
| a) Nicotine will not be funded in amounts less than 4 weeks | of treatment. | | | |
| b) Note: Direct Provision by a pharmacist permitted under the | he provisions in Part I | of Se | ection A. | |
| Patch 7 mg – Up to 28 patch available on a PSO | | 28 | - | Habitrol |
| Patch 7 mg for direct distribution only - [Xpharm] | | 7 | 1 | Habitrol |
| Patch 14 mg – Up to 28 patch available on a PSO | | 28 | 1 | Habitrol |
| Patch 14 mg for direct distribution only - [Xpharm] | | 7 | 1 | Habitrol |
| Patch 21 mg – Up to 28 patch available on a PSO | | 28 | 1 | Habitrol |
| Patch 21 mg for direct distribution only - [Xpharm] | | 7 | 1 | Habitrol |
| Lozenge 1 mg - Up to 216 loz available on a PSO | | 216 | 1 | Habitrol |
| Lozenge 1 mg for direct distribution only - [Xpharm] | | 36 | 1 | Habitrol |
| Lozenge 2 mg - Up to 216 loz available on a PSO | | 216 | 1 | Habitrol |
| Lozenge 2 mg for direct distribution only - [Xpharm] | | 36 | 1 | Habitrol |
| Gum 2 mg (Fruit) - Up to 384 piece available on a PSO | | 384 | 1 | Habitrol |
| Gum 2 mg (Fruit) for direct distribution only – [Xpharm] | | 96 | 1 | Habitrol |
| Gum 2 mg (Mint) – Up to 384 piece available on a PSO | | 384 | 1 | Habitrol |
| Gum 2 mg (Mint) for direct distribution only - [Xpharm] | | 96 | 1 | Habitrol |
| Gum 4 mg (Fruit) – Up to 384 piece available on a PSO | | 384 | 1 | Habitrol |
| Gum 4 mg (Fruit) for direct distribution only – [Xpharm] | | 96 | 1 | Habitrol |
| Gum 4 mg (Mint) – Up to 384 piece available on a PSO | | 384 | | Habitrol |
| Gum 4 mg (Mint) for direct distribution only – [Xpharm] | | 96 | | Habitrol |
| 2 () · · · · · · · · · · · · · · · · · · | | | | |

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack

b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

| Tab 0.5 mg × 11 and 1 mg × 4216.67 | 53 OP | Varenicline Pfizer |
|------------------------------------|-------|--|
| Tab 1 mg17.62 | 56 | ✓ Varenicline Pfizer |

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;
| Subs | sidy Full | / Brand or |
|-------------|------------------------|--------------|
| (Manufactur | rer's Price) Subsidise | d Generic |
| \$ | 6 Per 🖌 | Manufacturer |

continued...

and

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

| | Subsidy (Manufacturer's Price) | Su | Fully | Brand or Generic |
|--|-----------------------------------|-------------|---------|-------------------------|
| | \$ | Per | 1 | Manufacturer |
| Chemotherapeutic Agents | | | | |
| Alkylating Agents | | | | |
| BENDAMUSTINE HYDROCHLORIDE - PCT only - Special | st – Special Authority se | e SA204 | 6 below | |
| Inj 25 mg vial | | 1 | 🗸 F | Ribomustin |
| Inj 100 mg vial | | 1 | 🗸 F | Ribomustin |
| Inj 1 mg for ECP | | 1 mg | 🗸 E | Baxter |
| SA2046 Special Authority for Subsidy | | • | | |
| Initial application — (treatment naive CLL) only from a relevant | evant specialist or medic | al practiti | oner on | the recommendation of a |
| relevant specialist. Approvals valid for 12 months for application | | | | |
| All of the following: | J | J | | |

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Either:

3.2.3.1 Both:

- 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

| Su | ubsidy | - ully | Brand or |
|----------|------------------------|-----------|--------------|
| (Manufac | cturer's Price) Subsid | ised | Generic |
| | \$ Per | 1 | Manufacturer |

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

| Tab 2 mg | | 100 | Myleran |
|---|----------|-----------|---------------------------------------|
| CARBOPLATIN – PCT only – Specialist | | | |
| Inj 10 mg per ml, 45 ml vial | | 1 | DBL Carboplatin |
| | 45.20 | | Carboplatin Ebewe |
| | 48.50 | | Carbaccord |
| Inj 1 mg for ECP | 0.10 | 1 mg | Baxter |
| CARMUSTINE – PCT only – Specialist | | | |
| Inj 100 mg vial | 710.00 | 1 | BiCNU |
| | 1,387.00 | | ✓ Bicnu Heritage S29 |
| Inj 100 mg for ECP | | 100 mg OP | ✓ Baxter |
| (Bicnu Heritage S29 Inj 100 mg vial to be delisted 1 Septembe | | - | |
| CHLORAMBUCIL – PCT – Retail pharmacy-Specialist | | | |
| Tab 2 mg | | 25 | Leukeran FC |
| CISPLATIN – PCT only – Specialist | | | |
| Inj 1 mg per ml, 50 ml vial | | 1 | Cisplatin Ebewe |
| Inj 1 mg per ml, 100 ml vial | | 1 | Cisplatin Ebewe |
| ······································ | 29.66 | | ✓ DBL Cisplatin |
| Inj 1 mg for ECP | 0.31 | 1 mg | ✓ Baxter |
| CYCLOPHOSPHAMIDE | | 5 | |
| Tab 50 mg – PCT – Retail pharmacy-Specialist | 145 00 | 50 | Cyclonex |
| Inj 1 g vial – PCT – Retail pharmacy-Specialist | | 1 | ✓ Endoxan |
| | 127.80 | 6 | ✓ Cytoxan |
| Inj 2 g vial – PCT only – Specialist | | 1 | ✓ Endoxan |
| Inj 1 mg for ECP - PCT only - Specialist | | 1 mg | ✓ Baxter |
| IFOSFAMIDE – PCT only – Specialist | | 5 | |
| lnj 1 g | 96.00 | 1 | Holoxan |
| Inj 2 g | | 1 | ✓ Holoxan |
| Inj 1 mg for ECP | | 1 mg | ✓ Baxter |
| LOMUSTINE – PCT – Retail pharmacy-Specialist | | 5 | |
| Cap 10 mg | 132 59 | 20 | ✓ CeeNU |
| Cap 40 mg | | 20 | ✓ CeeNU |
| oopg | | _0 | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| (N | Subsidy /anufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|---|--|------|---------------------|----------------------------|
| MELPHALAN | | | | |
| Tab 2 mg – PCT – Retail pharmacy-Specialist | 40.70 | 25 | 1 | Alkeran |
| Inj 50 mg – PCT only – Specialist | 67.80 | 1 | ✓ | Alkeran |
| | | | 1 | Alkeran S29 S29 |
| OXALIPLATIN – PCT only – Specialist | | | | |
| Inj 100 mg vial | 25.01 | 1 | 1 | Oxaliplatin Actavis |
| , | | | | 100 |
| | 110.00 | | 1 | Oxaliplatin Ebewe |
| Inj 5 mg per ml, 20 ml vial | 46.32 | 1 | | Oxaliplatin Accord |
| Inj 1 mg for ECP | | 1 mg | 1 | Baxter |
| THIOTEPA – PCT only – Specialist | | | | |
| Inj 15 mg vial | CBS | 1 | 1 | Bedford S29 |
| , , | | | 1 | Max Health S29 |
| | | | 1 | THIO-TEPA S29 |
| | | | | Tepadina S29 |
| Ini 100 ma vial | CDC | 1 | | Max Health S29 |
| Inj 100 mg vial | | I | | |
| | | | • | Tepadina S29 |
| Antimetabolites | | | | |
| AZACITIDINE - PCT only - Specialist - Special Authority see SA1 | 467 below | | | |
| Inj 100 mg vial | | 1 | 1 | Azacitidine Dr |
| | | • | | Reddy's |
| | 605.00 | | 1 | Vidaza |
| Inj 1 mg for ECP | | 1 mc | | Baxter |
| | | | | |

⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
 - 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
 - 2 The patient has performance status (WHO/ECOG) grade 0-2; and
 - 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
 - 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

| Subsid | y | Fully Brand or |
|--|-----------|--|
| (Manufacturer | | sidised Generic |
| \$ | Per | Manufacturer |
| CALCIUM FOLINATE | | |
| Tab 15 mg – PCT – Retail pharmacy-Specialist114.69 | 10 | DBL Leucovorin |
| | | Calcium |
| Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist | 5 | Hospira |
| Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist7.28 | 1 | Calcium Folinate |
| | | <u>Sandoz</u> |
| | | Calcium Folinate |
| | | Sandoz S29 S29 |
| Inj 50 mg – PCT – Retail pharmacy-Specialist72.80 | 10 | Leucovorin |
| Jee Sterring Photos and Photos an | | Pharmacia S29 |
| Inj 10 mg per ml, 10 ml vial – PCT only – Specialist9.49 | 1 | ✓ Calcium Folinate |
| | 1 | Sandoz |
| Inj 100 mg – PCT only – Specialist7.33 | 1 | ✓ Calcium Folinate |
| | I | Ebewe |
| 04.00 | 10 | |
| 94.90 | 10 | Leucovorin |
| | | Pharmacia S29 |
| Inj 300 mg - PCT only - Specialist22.51 | 1 | Calcium Folinate |
| | | Ebewe |
| 25.14 | | Leucovorin DBL §29 |
| | | |
| Inj 10 mg per ml, 35 ml vial – PCT only – Specialist | 1 | Calcium Folinate |
| | | Sandoz |
| | | Calcium Folinate |
| | | Sandoz S29 S29 |
| Inj 1 g – PCT only – Specialist67.51 | 1 | Calcium Folinate |
| | | Ebewe |
| Inj 10 mg per ml, 100 ml vial – PCT only – Specialist | 1 | Calcium Folinate |
| | | Sandoz |
| Inj 1 mg for ECP – PCT only – Specialist0.06 | 1 mg | Baxter |
| APECITABINE – Retail pharmacy-Specialist | | |
| Tab 150 mg | 60 | Capercit |
| Tab 500 mg | 120 | ✓ Capercit |
| - | | |
| LADRIBINE – PCT only – Specialist Inj 1 mg per ml, 10 ml749.96 | 1 | Leustatin |
| Inj 10 mg for ECP | 10 mg OP | ✓ Baxter |
| | TO THE OF | • Daxlei |
| YTARABINE | - | |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist400.00 | 5 | Pfizer |
| Inj 100 mg per ml, 20 ml vial – PCT – Retail | | _ |
| pharmacy-Specialist | 1 | ✓ Pfizer |
| Inj 1 mg for ECP – PCT only – Specialist0.25 | 10 mg | Baxter |
| Inj 100 mg intrathecal syringe for ECP – PCT only – Specialist80.00 | 100 mg OP | Baxter |
| LUDARABINE PHOSPHATE | | |
| Tab 10 mg - PCT - Retail pharmacy-Specialist412.00 | 20 | Fludara Oral |
| Inj 50 mg vial - PCT only - Specialist | 5 | Fludarabine Ebewe |
| Inj 50 mg for ECP – PCT only – Specialist115.29 | 50 mg OP | Baxter |
| LUOROURACIL | - | |
| Inj 50 mg per ml, 20 ml vial – PCT only – Specialist | 1 | Fluorouracil Accord |
| Inj 50 mg per ml, 100 ml vial – PCT only – Specialist | 1 | Fluorouracil Accord Fluorouracil Accord |
| Inj 1 mg for ECP – PCT only – Specialist | 100 mg | ✓ Pluorouracii Accoru ✓ Baxter |
| | roonig | - DUALCI |

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | | Subsidy | | Fully | |
|---------------------|--|---|--|---|--|
| | | (Manufacturer's Pr \$ | rice) Sub Per | sidised | I Generic Manufacturer |
| | | φ | Fei | • | Manulacturer |
| iE | MCITABINE HYDROCHLORIDE – PCT only – Specialist | CO 50 | | | DDI Comeltatine |
| | lnj 1 g, 26.3 ml vial | | 1 | | DBL Gemcitabine |
| | lnj 1 g | | 1 | | Gemcitabine Ebewe |
| | Inj 1 mg for ECP | 0.02 | 1 mg | ~ | Baxter |
| R | NOTECAN HYDROCHLORIDE - PCT only - Specialist | | | | |
| | Inj 20 mg per ml, 5 ml vial | 52.57 | 1 | - | Accord |
| | | 71.44 | | 1 | Irinotecan Actavis 100 |
| | | 100.00 | | 1 | Irinotecan-Rex |
| | Inj 1 mg for ECP | 0.54 | 1 mg | 1 | Baxter |
| | RCAPTOPURINE | | 5 | | |
| | Tab 50 mg – PCT – Retail pharmacy-Specialist | 27.00 | 25 | 1 | Puri-nethol |
| | | | 25 | • | Full-fieldo |
| | Oral suspension 20 mg per ml – Retail pharmacy-Specialist Special Authority see SA1725 below | | 100 ml OP | 1 | Allmercap |
| | SA1725 Special Authority for Subsidy ial application only from a paediatric haematologist or paedia | atric oncologist. | Approvals vali | d for 1 | 12 months where the patie |
| | uires a total dose of less than one full 50 mg tablet per day. | | rr | | |
| | newal only from a paediatric haematologist or paediatric onco | logist Approvals | valid for 12 n | nonthe | s where patient still requir |
| | otal dose of less than one full 50 mg tablet per day. | | | iona i | |
| | 0 1 1 | | | | |
| | THOTREXATE | 0.00 | 00 | | T |
| | Tab 2.5 mg – PCT – Retail pharmacy-Specialist | | 90 | | Trexate |
| | Tab 10 mg – PCT – Retail pharmacy-Specialist | | 90 | | Trexate |
| | Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist . | | 5 | | Methotrexate DBL |
| ÷ | Inj 7.5 mg prefilled syringe | 14.61 | 1 | | Methotrexate |
| | , , , , , | | | • | |
| | | | | | Sandoz |
| ÷ | Inj 10 mg prefilled syringe | | 1 | | Sandoz Methotrexate |
| ŧ | Inj 10 mg prefilled syringe | 14.66 | 1 | ~ | Sandoz Methotrexate Sandoz |
| | | 14.66 | | ~ | Sandoz Methotrexate Sandoz Methotrexate |
| | Inj 10 mg prefilled syringe | 14.66 | 1 | ~ | Sandoz Methotrexate Sandoz |
| ŧ | Inj 10 mg prefilled syringe | 14.66 14.77 | 1 | | Sandoz Methotrexate Sandoz Methotrexate |
| ŧ | Inj 10 mg prefilled syringe | 14.66 14.77 | 1 | | Sandoz Methotrexate Sandoz Methotrexate Sandoz |
| * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe | | 1 | 1 1 1 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate |
| * | Inj 10 mg prefilled syringe | | 1 1 1 | 1 1 1 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate |
| * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe | 14.66 14.77 14.88 14.99 | 1 1 1 1 | \$ \$ \$ | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz |
| * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe | 14.66 14.77 14.88 14.99 | 1 1 1 | \$ \$ \$ | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate |
| * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe | 14.66 14.77 14.88 14.99 15.09 | 1 1 1 1 1 | י י י | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz |
| * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe | 14.66 14.77 14.88 14.99 15.09 | 1 1 1 1 | י י י | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL |
| * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special | 14.66 14.77 14.88 14.99 15.09 list30.00 | 1 1 1 1 1 5 | 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial |
| * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe | 14.66 14.77 14.88 14.99 15.09 list30.00 | 1 1 1 1 1 | 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate |
| * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special | 14.66 14.77 14.88 14.99 15.09 list30.00 alist45.00 | 1 1 1 1 1 5 1 | 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial |
| * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special | 14.66 14.77 14.88 14.99 15.09 list30.00 alist45.00 | 1 1 1 1 1 5 | 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate |
| * * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special | 14.66 14.77 14.88 14.99 15.09 list30.00 alist45.00 | 1 1 1 1 1 5 1 | 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial |
| * * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Special | 14.66 | 1 1 1 1 1 5 1 | 5 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial |
| * * * * * * ** | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail | 14.66 14.77 14.88 14.99 15.09 list30.00 alist45.00 st25.00 | 1 1 1 1 1 5 1 1 | 5 5 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe |
| * * * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail Inj 100 mg per ml, 50 ml vial – PCT – Retail Inj 100 mg per ml, 50 ml vial – PCT – Retail Inj 100 mg per ml, 50 ml vial – PCT – Retail | 14.66 14.77 14.88 14.99 15.09 list30.00 alist45.00 st25.00 | 1 1 1 1 1 5 1 1 1 1 | | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe <u>Methotrexate Ebewe</u> |
| * * * * * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist Inj 1 mg for ECP – PCT only – Specialist Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist | 14.66 | 1 1 1 1 5 1 1 1 1 1 1 mg 5 mg OP | | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe Methotrexate Ebewe Baxter |
| * * * * * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist Inj 1 mg for ECP – PCT only – Specialist Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist METREXED – PCT only – Specialist – Special Authority see | | 1 1 1 1 1 5 1 1 1 1 1 1 5 mg OP ext page | 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 5 | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe Baxter Baxter Baxter |
| ** ** * * * * * * | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist Inj 1 mg for ECP – PCT only – Specialist Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist METREXED – PCT only – Specialist – Special Authority see Inj 100 mg vial | | 1 1 1 1 1 5 1 1 1 1 mg 5 mg OP ext page 1 | \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe Methotrexate Ebewe Baxter Baxter Baxter |
| | Inj 10 mg prefilled syringe Inj 15 mg prefilled syringe Inj 20 mg prefilled syringe Inj 25 mg prefilled syringe Inj 30 mg prefilled syringe Inj 30 mg prefilled syringe Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Special Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Special Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist Inj 1 mg for ECP – PCT only – Specialist Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist METREXED – PCT only – Specialist – Special Authority see | | 1 1 1 1 1 5 1 1 1 1 1 1 5 mg OP ext page | \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ \$ | Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate Sandoz Methotrexate DBL Onco-Vial DBL Methotrexate Onco-Vial Methotrexate Ebewe Methotrexate Ebewe Baxter Baxter |

| Subsidy (Manufacturer's Price) | Su | Fully | Brand or Generic |
|-----------------------------------|-----|-------|---------------------|
| `\$ | Per | 1 | Manufacturer |

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and

2 Either:

- 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
- 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

THIOGUANINE - PCT - Retail pharmacy-Specialist

| Tab 40 mg126.31 | 25 | Lanvis |
|---|----------|----------------------------------|
| Other Cytotoxic Agents | | |
| AMSACRINE – PCT only – Specialist | | |
| Inj 50 mg per ml, 1.5 ml ampoule1,500.00 | 6 | Amsidine S29 |
| 4,736.00 | | Amsidine S29 |
| Inj 75 mg1,250.00 | 5 | AmsaLyo S29 |
| ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist | | |
| Cap 0.5 mg1,175.87 | 100 | 🗸 Agrylin |
| ARSENIC TRIOXIDE – PCT only – Specialist | | |
| Inj 1 mg per ml, 10 ml vial4,817.00 | 10 | Phenasen |
| Inj 10 mg for ECP | 10 mg OP | Baxter |
| | | |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy (Manufacturer's Pric \$ | e) Subs Per | Fully sidised | Brand or Generic Manufacturer |
|---------------------------------------|--|---|--|
| | 1 | √ [| DBL Bleomycin Sulfate |
| 14.32 | 1,000 iu | 🗸 E | Baxter |
| A1889 below 75.00 | 1 | ✓ E | Sortezomib Juno |
| 105.00 | 1 | ✓ E | Bortezomib Dr Reddy's S29 S29 Bortezomib Dr-Reddy's Bortezomib |
| | 1 mg | _ | Juno ^(S29) Baxter |
| | (Manufacturer's Pric \$185.1614.32 A1889 below75.00105.00 | (Manufacturer's Price) Sub- <u>Per</u> 185.16 1 14.32 1,000 iu A1889 below 75.00 1 | (Manufacturer's Price) Subsidised Per ✓ |

(Bortezomib Juno S29 Inj 3.5 mg vial to be delisted 1 August 2022)

► SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.
- Note: Indications marked with * are unapproved indications.
- DACARBAZINE PCT only Specialist

| Inj 200 mg vial | 62.70 | 1 | DBL Dacarbazine |
|--|--------|-----------|--------------------------------------|
| | 580.60 | 10 | Dacarbazine |
| | | | APP S29 |
| Inj 200 mg for ECP | 62.70 | 200 mg OP | Baxter |
| DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist | | | |
| Inj 0.5 mg vial | 255.00 | 1 | Cosmegen |
| Inj 0.5 mg for ECP | | 0.5 mg OP | Baxter |
| DAUNORUBICIN – PCT only – Specialist | | | |
| Inj 2 mg per ml, 10 ml | 149.50 | 1 | Pfizer |
| Inj 20 mg for ECP | 149.50 | 20 mg OP | Baxter |
| DOCETAXEL – PCT only – Specialist | | | |
| Inj 20 mg | | 1 | Docetaxel Sandoz |
| Inj 10 mg per ml, 8 ml vial | 46.89 | 1 | DBL Docetaxel |
| Inj 20 mg per ml, 4 ml vial | 26.95 | 1 | Docetaxel |
| | | | Accord S29 |
| Inj 80 mg | 195.00 | 1 | Docetaxel Sandoz |
| Inj 1 mg for ECP | | 1 mg | Baxter |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Brand or Subsidised Generic ✓ Manufacturer |
|--|---|--------|--|
| DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist | | | |
| Inj 2 mg per ml, 5 ml vial | | 1 | Doxorubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | 11.50 | 1 | Doxorubicin Ebewe |
| | 17.00 | | Arrow-Doxorubicin |
| Inj 2 mg per ml, 50 ml vial | | 1 | Doxorubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | 65.00 | 1 | Arrow-Doxorubicin |
| | 69.99 | | Doxorubicin Ebewe |
| Inj 1 mg for ECP | 0.35 | 1 mg | Baxter |
| EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist | | | |
| Inj 2 mg per ml, 5 ml vial | | 1 | Epirubicin Ebewe |
| Inj 2 mg per ml, 25 ml vial | | 1 | Epirubicin Ebewe |
| Inj 2 mg per ml, 100 ml vial | | 1 | Epirubicin Ebewe |
| Inj 1 mg for ECP | 0.50 | 1 mg | Baxter |
| ETOPOSIDE | | - | |
| Cap 50 mg – PCT – Retail pharmacy-Specialist | 340.73 | 20 | ✓ Vepesid |
| Cap 100 mg – PCT – Retail pharmacy-Specialist | | 10 | ✓ Vepesid |
| Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia | | 1 | ✓ Rex Medical |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | _ |
| ETOPOSIDE PHOSPHATE – PCT only – Specialist | | | |
| Inj 100 mg (of etoposide base) | 40.00 | 1 | Etopophos |
| Inj 1 mg (of etoposide base) for ECP | 0.47 | 1 mg | • • |
| HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pha | | 1 1119 | Builton |
| | | 100 | Devatis |
| Cap 500 mg | 23.02 | 100 | • Devalis |
| DARUBICIN HYDROCHLORIDE | 100 74 | | |
| Inj 5 mg vial – PCT only – Specialist | | 1 | Zavedos |
| Inj 10 mg vial – PCT only – Specialist | | 1 | ✓ Zavedos |
| Inj 1 mg for ECP – PCT only – Specialist | | 1 mg | Baxter |
| ENALIDOMIDE – Retail pharmacy-Specialist – Special Author Wastage claimable | rity see SA2047 below | | |
| Cap 5 mg | 5,122.76 | 28 | Revlimid |
| Cap 10 mg | 4,655.25 | 21 | Revlimid |
| | 6,207.00 | 28 | Revlimid |
| Cap 15 mg | 5,429.39 | 21 | Revlimid |
| | 7,239.18 | 28 | Revlimid |
| Cap 25 mg | 7,627.00 | 21 | Revlimid |

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

| Inj 100 mg per ml, 10 ml ampoule – PCT only – Specialist | Tab 400 mg – PCT – Retail pharmacy-Specialist Tab 600 mg – PCT – Retail pharmacy-Specialist Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist | .448.50 .177.45 | 50 50 15 | <u>Uromitexan</u> <u>Uromitexan</u> <u>Uromitexan</u> |
|--|---|--------------------|----------------|---|
| MITOMYCIN C – PCT only – Specialist Inj 5 mg vial | , , , , , , , | | 15 100 mg | Uromitexan Baxter |
| Inj 20 mg vial | , | 2.00 | roo mg | Daxiel |
| ✓ Teva ✓ Teva ✓ Inj 1 mg for ECP ✓ Mitozantrone Ebewe ✓ Mitozantrone Ebewe ✓ Mitozantrone Ebewe ✓ Baxter ✓ Mitozantrone Ebewe ✓ Baxter | Inj 5 mg vial | .641.70 | 1 | Accord S29 |
| MITOZANTRONE – PCT only – Specialist Inj 2 mg per ml, 10 ml vial | Inj 20 mg vial | 8,275.00 | 1 | |
| Inj 2 mg per ml, 10 ml vial | Inj 1 mg for ECP | .470.75 | 1 mg | Baxter |
| Inj 1 mg for ECP 5.51 1 mg ✓ Baxter OLAPARIB – Retail pharmacy-Specialist – Special Authority see SA1883 below ✓ Yunparza Tab 100 mg 56 ✓ Lynparza | MITOZANTRONE – PCT only – Specialist | | | |
| OLAPARIB – Retail pharmacy-Specialist – Special Authority see SA1883 below Tab 100 mg | Inj 2 mg per ml, 10 ml vial | 97.50 | 1 | Mitozantrone Ebewe |
| Tab 100 mg | Inj 1 mg for ECP | 5.51 | 1 mg | Baxter |
| 5 | OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1 | 883 below | | |
| Tab 150 mg | Tab 100 mg | 3,701.00 | 56 | Lynparza |
| | | 3,701.00 | 56 | Lynparza |

➡SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist

| Inj 30 mg | | 5 | Paclitaxel Ebewe |
|--|----------------|------|--------------------------------------|
| Inj 100 mg | | 1 | Paclitaxel Ebewe |
| | 91.67 | | Paclitaxel Actavis |
| Inj 150 mg | | 1 | Paclitaxel Ebewe |
| | 137.50 | | Anzatax |
| | | | Paclitaxel Actavis |
| Inj 300 mg | | 1 | Paclitaxel Ebewe |
| | 275.00 | | Anzatax |
| | | | Paclitaxel Actavis |
| Inj 1 mg for ECP | 0.20 | 1 mg | Baxter |
| PEGASPARGASE - PCT only - Special Authority se | e SA1979 below | - | |
| Inj 750 iu per ml, 5 ml vial | 3,455.00 | 1 | Oncaspar LYO S29 |

➡SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has newly diagnosed acute lymphoblastic leukaemia; and

2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

| PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist | t | | |
|---|-----|---|--------------------------------|
| Inj 10 mg | CBS | 1 | Nipent S29 |

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|---|---|-----|---------------------|------------------|
| PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharm | acy-Specialist | | | |
| Cap 50 mg | | 50 | 1 | Natulan S29 |
| FEMOZOLOMIDE - Special Authority see SA1741 below - I | Retail pharmacy | | | |
| Cap 5 mg | | 5 | 1 | Temaccord |
| Cap 20 mg | | 5 | ✓ | Temaccord |
| | 18.30 | | ✓ | Apo-Temozolomide |
| | 136.00 | 14 | 1 | Accord S29 |
| Cap 100 mg | | 5 | 1 | Temaccord |
| | 40.20 | | ✓ | Apo-Temozolomide |
| | 532.00 | 14 | 1 | Accord S29 |
| Cap 140 mg | | 5 | 1 | Temaccord |
| | 400.00 | | 1 | Amneal S29 |
| Cap 180 mg | 620.00 | 14 | 1 | Accord S29 |
| Cap 250 mg | | 5 | 1 | Temaccord |
| · - | 688.00 | | 1 | Amneal S29 |

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | F | ully | Brand or |
|------------------------|--------|------|--------------|
| (Manufacturer's Price) | Subsid | ised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

Both:

1 No evidence of disease progression; and

2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 No evidence of disease progression; and

2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special Authority see SA1124 below

| Cap 50 mg | 378.00 | 28 | Thalomid |
|------------|--------|----|------------------------------|
| Cap 100 mg | 756.00 | 28 | Thalomid |

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

| Cap 10 mg – PCT – Retail pharmacy-Specialist | | 100 | Vesanoid |
|--|----------------------|-------|-------------------------------|
| VENETOCLAX - Retail pharmacy-Specialist - Special Author | rity see SA1868 belo | w | |
| Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg | | 42 OP | Venclexta |
| Tab 10 mg | | 14 OP | Venclexta |
| Tab 50 mg | | 7 OP | Venclexta |
| Tab 100 mg - Wastage claimable | 8,209.41 | 120 | Venclexta |
| | | | |

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal --- (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application --- (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

| Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37 Inj 1 mg for ECP – PCT only – Specialist6.00 | 5 1 mg | ✓ Hospira✓ Baxter |
|--|-----------|---|
| VINCRISTINE SULPHATE | | |
| Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist74.52 | 5 | DBL Vincristine Sulfate |
| Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist 102.73 | 5 | DBL Vincristine Sulfate |
| Inj 1 mg for ECP – PCT only – Specialist12.60 | 1 mg | Baxter |
| VINORELBINE – PCT only – Specialist | | |
| Inj 10 mg per ml, 1 ml vial | 1 | Navelbine |
| 42.00 | | Vinorelbine Ebewe |
| Inj 10 mg per ml, 5 ml vial56.00 | 1 | Navelbine |
| 210.00 | | Vinorelbine Ebewe |
| 328.65 | | Sagent S29 |
| Inj 1 mg for ECP 1.25 | 1 mg | Baxter |
| Inj 50 mg for ECP | 50 mg OP | Baxter (Sagent) |

Protein-tyrosine Kinase Inhibitors

| ALECTINIB - Retail pharmacy-Specialist - Special Author | ity see SA1870 below | |
|---|----------------------|-----|
| Wastage claimable | - | |
| Cap 150 mg | 7,935,00 | 224 |

| Cap 150 mg7,9 | 935.00 |
|---------------|--------|
|---------------|--------|

SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate

✓ Alecensa

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

continued...

ALK test; and

3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB – Special Authority see SA1805 below – Retail pharmacy

| Wastage claimable | |
|-------------------|--|
|-------------------|--|

| Tab 20 mg | 60 | Sprycel |
|-------------------|----|-----------------------------|
| Tab 50 mg6,214.20 | 60 | Sprycel |
| Tab 70 mg7,692.58 | 60 | Sprycel |

➡SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 Both:

- 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
- 1.2 Maximum dose of 140 mg/day; or

2 Both:

- 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
- 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

| ERLOTINIB - Retail pharmacy-Speci | alist – Special Authority see SA2115 below |
|-----------------------------------|--|
|-----------------------------------|--|

| Tab 100 mg | 30 | Tarceva |
|--------------------|----|-----------------------------|
| Tab 150 mg1,146.00 | 30 | Tarceva |

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|------|---------|--------------|--|
| (Manufacturer's Price) | Subs | sidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see SA2116 below

| Tab 250 mg1,700.00 |) 30 | Iressa |
|--------------------|------|----------------------------|
|--------------------|------|----------------------------|

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

| | Tab 100 mg - [Xpharm] - Special Authority see SA1460 o | n the | | |
|---|--|-------|----|----------------------------------|
| | next page | | 60 | Glivec |
| * | Cap 100 mg | | 60 | Imatinib-Rex |
| | Cap 400 mg | | 30 | Imatinib-Rex |

| | | Subsidy (Manufacturer's Price) \$ | Subs Per | Fully sidised | Brand or Generic Manufacturer |
|--|--|---|--------------|------------------|-------------------------------------|
| ■ SA1460 Special Authority for Special Authority approved by the Notes: Application details may be should be sent to: | | ite <u>schedule.pharm</u> | ac.govt.nz/ | SAForm | 18, and prescriptions |
| The CML/GIST Co-ordinator | Phone: (04) 460 4990 | | | | |
| Pharmac | Facsimile: (04) 916 7571 | | | | |
| PO Box 10 254 | Email: cmlgistcoordinator@pha | armac.govt.nz | | | |
| Wellington | ······································ | ······································ | | | |
| Special Authority criteria for G Funded for patients: | IST – access by application | | | | |
| | ed by an oncologist) of unresecta | ble and/or metastat | ic maligna | nt gastro | pintestinal stromal tumou |
| d) Initial and subsequent ap | g/day. and subsequent prescriptions can plications are valid for one year. b (prescriber determined). | | | an adeq | uate clinical response to |
| Note – no new patients to be | ecial Authority see SA2035 below e initiated on lapatinib ditosylate. | | 70 | | |
| ° | • · · · | 1,899.00 | 70 | ✓ T | ykerb |
| SA2035 Special Authority for Benewal — (metastatic breast) | or Subsidy cancer) only from a relevant spe | cialist or medical p | actitioner (| on the re | commendation of a |
| | alid for 12 months for applications | | | | |
| 5 | c breast cancer expressing HER- | 2 IHC 3+ or ISH+ (i | ncluding F | ISH or o | ther current technology); |
| 1 0 | essed at any time point during the in combination with trastuzumab; red at disease progression. | | ns whilst or | ı lapatin | ib; and |
| NILOTINIB – Special Authority s Wastage claimable | see SA1489 below - Retail pharm | асу | | | |
| | | | 120 120 | | asigna asigna |
| SA1489 Special Authority for Initial application only from a har All of the following: | or Subsidy aematologist. Approvals valid for | 6 months for applic | ations mee | eting the | following criteria: |
| 0 | f chronic myeloid leukaemia (CML |) in blast crisis, acc | elerated p | hase, oi | r in chronic phase; and |
| | nented CML treatment failure* with | , | | | |
| | enced treatment limiting toxicity w | vith imatinib preclud | ing further | treatme | nt with imatinib; and |
| 3 Maximum nilotinib dose o | 0,00 | | | | |
| 4 Subsidised for use as mo | | | | | |
| | ed by Leukaemia Net Guidelines. gist. Approvals valid for 6 months | | eeting the f | ollowing | j criteria: |
| 1 Lack of treatment failure | while on nilotinib as defined by Le ns appropriate and the patient is b of 800 mg/day: and | | | | |

- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|---|--|------|---------------------|--------------------|
| PALBOCICLIB – Retail pharmacy-Specialist – Wastage claimable | Special Authority see SA1894 below | | | |
| Tab 75 mg | | 21 | 1 | Ibrance |
| Tab 100 mg | 4,000.00 | 21 | 1 | Ibrance |
| Tab 125 mg | 4,000.00 | 21 | 1 | Ibrance |
| SA1894 Special Authority for Subsidy | ict or modical practitionar on the recom | mond | ation of a | Madical appalagist |

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:
 - second or subsequent line setting
 - 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

| Tab 200 mg | | 30 | Votrient |
|------------|----------|----|------------------------------|
| Tab 400 mg | 2,669.40 | 30 | Votrient |

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal: or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70: or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable

| Tab 5 mg | 2,500.00 | 56 | 🖌 Jakavi |
|-----------|----------|----|----------|
| Tab 10mg | | 56 | 🗸 Jakavi |
| Tab 15 mg | | 56 | 🗸 Jakavi |
| Tab 20 mg | | 56 | 🖌 Jakavi |

■ SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or 2.2 Both:
 - - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment: and
- 2 A maximum dose of 20 mg twice daily is to be given.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | |
|--|---|-----|---------------------|------------------|
| SUNITINIB – Special Authority see SA2117 below – Retail phan | macy | | | |
| Cap 12.5 mg | | 28 | ✓ | Sunitinib Pfizer |
| | 2,315.38 | | ✓ | Sutent |
| Sunitinib Pfizer to be Principal Supply on 1 July 2022 | | | | |
| Cap 25 mg | | 28 | 1 | Sunitinib Pfizer |
| | 4,630.77 | | 1 | Sutent |
| Sunitinib Pfizer to be Principal Supply on 1 July 2022 | | | | |
| Cap 50 mg | | 28 | ✓ | Sunitinib Pfizer |
| | 9.261.54 | | 1 | Sutent |
| Subitinih Bfizer to be Bringing! Supply on 1 July 2022 | -, | | | |

Sunitinib Pfizer to be Principal Supply on 1 July 2022

(Sutent Cap 12.5 mg to be delisted 1 July 2022)

(Sutent Cap 25 mg to be delisted 1 July 2022) (Sutent Cap 50 mg to be delisted 1 July 2022)

► SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1. 1 The r
- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- Both:
 - 1 No evidence of disease progression; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | ; | Subsidised | Generic | |
| \$ | Per | ~ | Manufacturer | |

continued...

2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

Tab 250 mg4,276.19

.4,276.19 120

Zytiga

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and

4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy | F | ully | Brand or |
|-----|---------------------|--------|------|--------------|
| (Ma | nufacturer's Price) | Subsid | ised | Generic |
| | \$ | Per | ✓ | Manufacturer |

continued...

- 4.1.3 Patient has ECOG performance score of 0-1; and
- 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICALUTAMIDE

| Tab 50 mg | 4.21 | 28 | Binarex |
|---|-----------------------|-----|------------------------------|
| FLUTAMIDE | | | |
| Tab 250 mg | | 90 | Prostacur S29 |
| - | 119.50 | 100 | Flutamin |
| FULVESTRANT - Retail pharmacy-Specialist - Special Auth | nority see SA1895 bel | ow | |
| Inj 50 mg per ml, 5 ml prefilled syringe | | 2 | Faslodex |

► SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|----------|---------------------|---|
| MEGESTROL ACETATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who we prescription is endorsed accordingly. Pharmacists may an prior dispensing of megestrol acetate. | | | | |
| Tab 160 mg | 48.80 63.53 | 30 | | Megace S29 Apo-Megestrol |
| (Megace Sean Tab 160 mg to be delisted 1 February 2023) (Apo-Megestrol Tab 160 mg to be delisted 1 May 2022) | | | | |
| OCTREOTIDE | | _ | | |
| Inj 50 mcg per ml, 1 ml vial | | 5 | ✓ (| Octreotide |
| | | | | MaxRx S29 |
| | 56.87 | - | | DBL Octreotide |
| Inj 100 mcg per ml, 1 ml vial | | 5 | | DBL Octreotide |
| Inj 500 mcg per ml, 1 ml vial | 145.00 222.00 | 5 | | DBL Octreotide Octreotide (Sun) 829 |
| Inj 50 mcg per ml, 1 ml ampoule | 27.58 | 5 | √ 1 | Max Health |
| | 30.64 | Ŭ | | Octreotide GH S29 |
| Max Health to be Principal Supply on 1 June 2022 | 00.04 | | | |
| Inj 100 mcg per ml, 1 ml ampoule | 32.71 | 5 | √ 1 | Max Health |
| | | Ŭ | | Octreotide GH S29 |
| Max Health to be Principal Supply on 1 June 2022 | | | | |
| Inj 500 mcg per ml, 1 ml ampoule | | 5 | √ 1 | Max Health |
| | | | | Octreotide GH S29 |
| Max Health to be Principal Supply on 1 June 2022 | | | | |
| (Octreotide MaxRx S29 Inj 50 mcg per ml, 1 ml vial to be delis | sted 1 June 2022) | | | |
| (DBL Octreotide Inj 50 mcg per ml, 1 ml vial to be delisted 1 Ju | | | | |
| (DBL Octreotide Inj 100 mcg per ml, 1 ml vial to be delisted 1 J | | | | |
| (DBL Octreotide Inj 500 mcg per ml, 1 ml vial to be delisted 1 J | lune 2022) | | | |
| (Octreotide (Sun) 529 Inj 500 mcg per ml, 1 ml vial to be delis | ted 1 June 2022) | | | |
| (Octreotide GH S29 Inj 50 mcg per ml, 1 ml ampoule to be de | listed 1 June 2022) | | | |
| (Octreotide GH S29 Inj 100 mcg per ml, 1 ml ampoule to be d | | | | |
| (Octreotide GH 529 Inj 500 mcg per ml, 1 ml ampoule to be d | | | | |
| OCTREOTIDE LONG-ACTING – Special Authority see SA211 | , | | | |
| Inj depot 10 mg prefilled syringe | | acy 1 | ✓ <u>(</u> | <u>Octreotide Depot</u> Teva |
| Inj depot 20 mg prefilled syringe | 647.03 | 1 | ✓ <u>(</u> | Octreotide Depot Teva |
| Inj depot 30 mg prefilled syringe | 718.55 | 1 | ✓ <u>(</u> | Octreotide Depot Teva |
| - CA0110 Createl Authority for Cubaidy | | | | |

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | : | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|------------------------------|-------------------------------|--|
| continued Renewal — (Other Indications) only from a relevant special specialist. Approvals valid for 2 years where the treatment renitial application — (pre-operative acromegaly) only from of a relevant specialist. Approvals valid for 12 months for application the following: Patient has a cromegaly; and Patient has a large pituitary tumour, greater than 10 m | mains appropriate and th n a relevant specialist or i plications meeting the fol nm at its widest; and | ne pat medic | ient is bene al practition | fiting from treatment. |
| 3 Patient is scheduled to undergo pituitary surgery in the AMOXIFEN CITRATE k Tab 10 mg k Tab 20 mg | 15.00 | 60 60 | | amoxifen Sandoz amoxifen Sandoz |
| Aromatase Inhibitors | | | | |
| ANASTROZOLE * Tab 1 mg EXEMESTANE | 4.55 | 30 | ✓ A | natrole |
| ₭ Tab 25 mg | 14.50 | 30 | ✓ P | fizer Exemestane |
| ETROZOLE ≰ Tab 2.5 mg | 5.84 | 30 | ✓ <u>L</u> | .etrole |
| Immunosuppressants Cytotoxic Immunosuppressants | | | | |
| AZATHIOPRINE ★ Tab 25 mg | 7.60 | 60 100 1 | A | Izamun Izamun muran |
| MYCOPHENOLATE MOFETIL Tab 500 mg Cap 250 mg Powder for oral liq 1 g per 5 ml – Subsidy by endorseme Mycophenolate powder for oral liquid is subsidised o the prescription is endorsed accordingly. | 35.90 ent187.25 16 | 50 100 5 ml (swall | ✓C DP ✓C | Celicept Celicept Celicept and capsules, and when |
| Fusion Proteins | | | | |
| TANERCEPT – Special Authority see SA2103 below – Ret Inj 25 mg Inj 25 mg autoinjector Inj 50 mg autoinjector Inj 50 mg prefilled syringe >> SA2103 Special Authority for Subsidy | | 4 4 4 4 | ✓ E | inbrel inbrel inbrel inbrel |

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Either:

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

| Sub | osidy F | ully | Brand or |
|------------|---------|------|--------------|
| (Manufactu | | sed | Generic |
| \$ | \$Per | 1 | Manufacturer |

continued...

2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | : | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and

1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and

2.4 Either:

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g.
 - prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price | e) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- diagnosis; or
- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Renewal — (severe chronic plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|----------|--------------|
| (Manufacturer's Price) | Su | bsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

| ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Spec Inj 50 mg per ml, 5 ml | | 5 | ✔ ATGAM |
|--|-----------------|---|-------------------------------|
| BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT on Subsidised only for bladder cancer. | ly – Specialist | | |
| Inj 2-8 × 100 million CFU | 149.37 | 1 | ✓ OncoTICE |
| lnj 40 mg per ml, vial | 176.90 | 3 | ✓ SII-Onco-BCG ^{S29} |

Monoclonal Antibodies

| ADALIMUMAB (AMGEVITA) - Special Authority see SA2102 below | - Retail pharm | acy | |
|--|----------------|-----|------------------------------|
| Inj 20 mg per 0.4 ml prefilled syringe | 190.00 | 1 | Amgevita |
| Inj 40 mg per 0.8 ml prefilled pen | 375.00 | 2 | Amgevita |
| Inj 40 mg per 0.8 ml prefilled syringe | 375.00 | 2 | Amgevita |

⇒SA2102 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

1 Both:

1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.
- Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and

| Subsidy | | Fully | Brand or | |
|--------------------|------|------------|--------------|--|
| (Manufacturer's Pr | ice) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or

2 Both:

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
 - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI
| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | S | Subsidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

| Subsidy | | Fully | Brand or |
|------------------------|------|--------|--------------|
| (Manufacturer's Price) | Subs | idised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and

2.6 Either:

- 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and

1.2 Either:

- 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
- 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) 5 | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|---------|--------------|
| (Manufacturer's Price) | Sub | sidised | Generic |
| \$ | Per | ~ | Manufacturer |

continued...

- 3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA) - Special Authority see SA2101 below - Retail pharmacy

| Inj 20 mg per 0.4 ml prefilled syringe | | 2 | 🗸 Humira |
|--|----------|---|-------------------------------|
| Inj 40 mg per 0.8 ml prefilled pen | 1,599.96 | 2 | HumiraPen |
| Inj 40 mg per 0.8 ml prefilled syringe | 1,599.96 | 2 | 🗸 Humira |

⇒SA2101 Special Authority for Subsidy

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | . : | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

| Subsidy (Manufacturer's Price) | Sub | Fully osidised | Brand or Generic |
|-----------------------------------|-----|-------------------|---------------------|
| \$ | Per | ~ | Manufacturer |

continued...

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient reported pain.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Fither:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

3 Either:

- 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Fither:

- 1.1 Applicant is a dermatologist; or
- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

| Subsidy | | Fully | Brand or | |
|----------------------|------|------------|--------------|--|
| (Manufacturer's Pric | e) : | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

2.2 Both:

- 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2.2 Either:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial......1,250.00 1 🖌 🗸 Eylea

➡SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

1 All of the following:

- 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
- 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
- 1.3 There is no structural damage to the central fovea of the treated eye; and
- 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or

2 Either:

continued...

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

- 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
- 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

- Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg
 - per ml imdevimab, 11.1 ml vial (1).....0.00 1 OP 🗸 Ronapreve

➡SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

| Inj 5 mg per ml, 20 ml vial | | 1 | Erbitux |
|------------------------------|----------|------|-----------------------------|
| Inj 5 mg per ml, 100 ml vial | 1,820.00 | 1 | 🗸 Erbitux |
| Inj 1 mg for ECP | | 1 mg | Baxter |

| Subsidy (Manufacturer's Price) | Fully Subsidised | | Brand or Generic |
|-----------------------------------|---------------------|---|---------------------|
| \$ | Per | 1 | Manufacturer |
| | | | |

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA2082 below

| Inj 100 mg | 1 | Remicade |
|------------------|----------|------------------------------|
| Inj 1 mg for ECP | 1 mg | Baxter |

⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 Paediatric patient has severe active Crohn's disease; and
 - 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
 - 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and

| Subsidy | | Fully | Brand or | |
|--------------------|-----|-----------------------|--------------|--|
| (Manufacturer's Pr | ce) | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

| Subsidy (Manufacturer's Price) | ç | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | ~ | Manufacturer | |

continued...

- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and

continued...

AThree months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

4 Either:

- 4.1 IV cyclophosphamide has been tried; or
- 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Renewal — (plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

1.1 Both:

| Subsidy | | Fully | Brand or |
|--------------------|----------|----------|--------------|
| (Manufacturer's Pr | ice) Sul | bsidised | Generic |
| \$ | Per | ~ | Manufacturer |

continued...

- 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation; or
 - 2.5 Chronic ocular inflammation; or
 - 2.6 Crohn's disease (adults); or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease; or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis; or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

| Subsidy | Ful | ly Brand or | |
|-------------------|------------------|----------------------------------|--|
| (Manufacturer's F | Price) Subsidise | d Generic | |
| \$ | Per | Manufacturer | |

continued...

- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or

2 Both:

- 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
- 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

- Any of the following:
 - 1 The patient has had a good clinical response following 3 initial doses; or
 - 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and

| Subsidy | Fu | ly Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per | Manufacturer | |

continued...

corticosteroids; and

4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g.
- prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

1 Patient has shown clinical improvement: and

- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

| | | see SA1896 below – Retail pharmacy | MEPOLIZUMAB – Special Authority see SA |
|----------------------------|---|------------------------------------|--|
| Nucala | 1 | | Inj 100 mg prefilled pen |
| Nucala | 1 | | Inj 100 mg vial |

► SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^{9} cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and

6 Either:

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| \$ | Per 🗸 | Manufacturer | |

continued...

- 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Authority see SA1627 below

| Inj 25 mg per ml, 40 ml vial | 5,910.00 | 1 | 🗸 Gazyva |
|------------------------------|----------|------|----------------------------|
| Inj 1 mg for ECP | 6.21 | 1 mg | Baxter |

➡SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^{9} /L and platelets greater than or equal to 75×10^{9} /L.

OMALIZUMAB – Special Authority see SA1744 below – Retail pharmacy

| Inj 150 mg prefilled syringe | 1 | 🗸 Xolair |
|------------------------------|-------|----------|
| Inj 150 mg vial | 1 | 🗸 Xolair |

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | S | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

Patient has previously adequately responded* to 6 doses of omalizumab; or

2 Both:

- 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
- 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab.

| | Subsidy (Manufacturer's Price) | | Fully idised | Brand or Generic |
|---|-----------------------------------|-------------|-----------------|-------------------------|
| | \$ | Per | | Manufacturer |
| continued | | | | |
| Complete response is defined as UAS7 less than or equal to 6 au chronic urticaria on stopping prednisone/ciclosporin does not just | | | or UC1 | F of 16. Relapse of |
| PERTUZUMAB - PCT only - Specialist - Special Authority see | SA1606 below | | | |
| Inj 30 mg per ml, 14 ml vial | 3,927.00 | 1 | ✓ P | erjeta |
| Inj 420 mg for ECP | | 20 mg OP | 🗸 В | axter |
| ► SA1606 Special Authority for Subsidy Initial application — (metastatic breast cancer) only from a re | elevant specialist or r | nedical pra | ctitione | r on the recommendation |

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

2 Either:

- 2.1 Patient is chemotherapy treatment naïve; or
- 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

| Inj 100 mg per 10 ml vial1,075.5 | 0 2 | Mabthera |
|----------------------------------|--------|------------------------------|
| Inj 500 mg per 50 ml vial2,688.3 | 0 1 | Mabthera |
| Inj 1 mg for ECP | 4 1 mg | 🗸 Baxter (Mabthera) |

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of

| Subsidy | | Fully | Brand or | |
|-----------------------|-----|------------|--------------|--|
| (Manufacturer's Price |) | Subsidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

continued...

leflunomide alone or in combination with oral or parenteral methotrexate; and

- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and

8 Either:

- 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2114 below

| ······································ | | | |
|--|-----------|---|------------------|
| Inj 100 mg per 10 ml vial | | ✓ | Riximyo |
| Inj 500 mg per 50 ml vial | | ✓ | Riximyo |
| Inj 1 mg for ECP | 1.38 1 mg | ✓ | Baxter (Riximyo) |

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*. Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

2 Any of the following:

- 2.1 The patient is rituximab treatment naive; or
- 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
- 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2. **Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.
- Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is

| Subsidy | Ful | y Brand or |
|-----------------|------------------|--------------|
| (Manufacturer's | Price) Subsidise | d Generic |
| \$ | Per • | Manufacturer |

continued...

considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and

2 Either:

- 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
- 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and

2 Either:

2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or

2.2 Both:

- 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
- 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal - (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of

| | ubsidy | Fully | Brand or |
|----|--------------------|-----------|--------------|
| | cturer's Price) Su | Ibsidised | Generic |
| Υ. | \$ Per | 1 | Manufacturer |

continued...

a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

- All of the following:
 - 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
 - 2 An initial response lasting at least 12 months was demonstrated; and
 - 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS))

only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.
- Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

| Subsidy | Ful | ly Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

continued

All of the followina:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months. but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application - (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or

2 Both:

- 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
- 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.
- Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe concenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20.000 platelets per microlitre: or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:

| Subsidy | | Fully | Brand or |
|------------------------|-------|-------|--------------|
| (Manufacturer's Price) | Subsi | dised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2.1 Treatment with steroids and splenectomy have been ineffective; or
- 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
- 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

- Either:
 - 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
 - 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
- 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.
- Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

| Subsidy | Full | y Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application - (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
 - 2 The patient has not received rituximab in the previous 6 months; and

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | | Fully | | |
|---------------------|------|------------|-----------|-----|
| (Manufacturer's Pri | ice) | Subsidised | I Generic | |
| \$ | Pe | er 🗸 | Manufactu | rer |

continued...

- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
- 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria: All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

| Subsidy | Full | y Brand or | |
|------------------------|-----------|--------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per 🖌 | Manufacturer | |

continued...

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.
- Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

| Cosentyx | 1 | Inj 150 mg per ml, 1 ml prefilled syringe799.50 |
|------------------------------|---|---|
| Cosentyx | 2 | 1,599.00 |

➡SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and

| Subsidy | | Fully | Brand or |
|------------------------|-----|----------|--------------|
| (Manufacturer's Price) | Sub | osidised | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal - (ankylosing spondylitis - second-line biologic) only from a rheumatologist or medical practitioner on the

| Subsidy | Ful | ly Brand or | |
|------------------------|-------------|----------------------------------|--|
| (Manufacturer's Price) |) Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

continued...

recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
 Secukinumab to be administered at doses no greater than 150 mg monthly.
- Initial application (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

| Note: Siltuximab is to be administered at doses no g | reater than 11 mg/kg every | 3 weeks. | |
|--|----------------------------|----------|-----------------------------|
| Inj 100 mg vial | | 1 | Sylvant |
| Inj 400 mg vial | | 1 | Sylvant |

*Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Ful | ly Brand or | |
|------------------------|-----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidise | d Generic | |
| \$ | Per • | Manufacturer | |

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

| TOCILIZUMAB – PCT only – Special Authority see SA2100 below | | |
|---|------|-------------------------------------|
| Inj 20 mg per ml, 4 ml vial | 1 | Actemra |
| | | Actemra S29 S29 |
| | | RoActemra S29 S29 |
| 880.00 | 4 | RoActemra S29 S29 |
| Inj 20 mg per ml, 10 ml vial550.00 | 1 | Actemra |
| | | Actemra S29 S29 |
| | | RoActemra S29 S29 |
| Inj 20 mg per ml, 20 ml vial1,100.00 | 1 | Actemra |
| | | Actemra S29 S29 |
| | | RoActemra S29 S29 |
| 4,400.00 | 4 | RoActemra S29 S29 |
| Inj 1 mg for ECP | 1 mg | Baxter |

⇒SA2100 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
| Subsidy | | Fully | Brand or | |
|-----------------------|------|------------|--------------|--|
| (Manufacturer's Price | e) 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

continued...

- 2.4 polyarticular juvenile idiopathic arthritis; or
- 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

Initial application — (moderate to severe COVID-19*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Note: Indications marked with * are unapproved indications.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

| Inj 150 mg vial | | 1 | Herceptin |
|------------------|------|------|-------------------------------|
| Inj 440 mg vial | | 1 | Herceptin |
| Inj 1 mg for ECP | 9.36 | 1 mg | Baxter |

➡SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | S | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

continued...

- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

continued...

3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and

4 Either:

- 4.1 Trastuzumab will not be given in combination with pertuzumab; or
- 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

| TRASTUZUMAB EMTANSINE - PCT only - Specialist | - Special Authority see SA18 | 371 below | |
|---|------------------------------|-----------|-----------------------------|
| Inj 100 mg vial | 2,320.00 | 1 | 🗸 Kadcyla |
| Inj 160 mg vial | 3,712.00 | 1 | Kadcyla |
| Inj 1 mg for ECP | 24.52 | 1 mg | Baxter |

⇒SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and

5 Either:

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

| | ext page | /OLUMAB – PCT only – Specialist – Special Authority see SA2120 on the n | NIVOLUMAB – PC |
|----------------------------|----------|---|------------------|
| Opdivo | 1 | Inj 10 mg per ml, 4 ml vial | Inj 10 mg per m |
| Opdivo | 1 | Inj 10 mg per ml, 10 ml vial2,629.96 | Inj 10 mg per m |
| Baxter | 1 mg | Inj 1 mg for ECP | Inj 1 mg for ECI |

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | Su | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

- Either:
 - 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

| | Subsidy (Manufacturer's Price) \$ | S Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|---|----------|---------------------|-------------------------------------|
| PEMBROLIZUMAB – PCT only – Specialist – Special Authority s Inj 25 mg per ml, 4 ml vial | | 1 | ✓ К | eytruda |
| Inj 1 mg for ECP | 49.14 | 1 mg | 🗸 В | axter |

➡SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

| CICLOSPORIN | | | |
|---|-----------------|----------|------------------------------|
| Cap 25 mg | | 50 | Neoral |
| Cap 50 mg | | 50 | Neoral |
| Cap 100 mg | | 50 | Neoral |
| Oral liq 100 mg per ml | 198.13 | 50 ml OP | Neoral |
| EVEROLIMUS - Special Authority see SA2008 below - | Retail pharmacy | | |
| Wastage claimable | | | |
| Tab 10 mg | 6,512.29 | 30 | Afinitor |
| Tab 5 mg | 4,555.76 | 30 | Afinitor |

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has tuberous sclerosis; and

2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS – Special Authority see SA2005 below – Retail pharmacy

| Tab 1 mg | .749.99 | 100 | Rapamune |
|----------------------|---------|----------|------------------------------|
| Tab 2 mg1 | ,499.99 | 100 | Rapamune |
| Oral liq 1 mg per ml | .449.99 | 60 ml OP | Rapamune |

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min; or
- · Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- · HUS or TTP; or
- Leukoencepthalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - $2.3\$ Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| Subsidy | Fu | lly Brand or | |
|-------------------|----------------|----------------------------------|--|
| (Manufacturer's P | rice) Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

continued...

- 2.2.1 Vigabatrin is contraindicated; and
- 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

| Cap 0.5 mg | 100 | Tacrolimus Sandoz |
|-------------|-----|---------------------------------------|
| Cap 0.75 mg | 100 | Tacrolimus Sandoz |
| Cap 1 mg | 100 | Tacrolimus Sandoz |
| Cap 5 mg | 50 | Tacrolimus Sandoz |

➡SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
 - 1 Patient requires long-term systemic immunosuppression; and
 - 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB - Special Authority see SA2079 below - Retail pharmacy

Tab 15 mg1,271.00

SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and

2 Either:

- 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
- 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and

3 Either:

3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or

continued...

✓ RINVOQ

28

| Subsidy | F | ully | Brand or |
|------------------------|----------|------|--------------|
| (Manufacturer's Price) | Subsidis | sed | Generic |
| \$ | Per | 1 | Manufacturer |

continued...

- 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and

3.2.2 Either:

- 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
- 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|--|--|-----------------------------|--|-------------------------------------|
| Antiallergy Preparations | | | | |
| Allergic Emergencies | | | | |
| ICATIBANT - Special Authority see SA1558 below - Retail phylin 10 mg per ml, 3 ml prefilled syringe | 2,668.00 specialist. Approvals l/oro-pharyngeal or sev s of C1-esterase inhibit eed upon an action pla | ere al or def n for s | for 12 month odominal att iciency; and self-adminis | acks of acute hereditary |
| Allergy Desensitisation | | | | |
| ■ SA1367 Special Authority for Subsidy | alid for 2 years for anal | icatio | as mosting t | he following criteria: |

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see SA1367 above - Retail pharmacy

| riotan phanna | oy . |
|------------------|--|
| 1 OP | VENOX S29 |
| 1 OP | VENOX S29 |
| | |
| 1 OP | Venomil S29 |
| | |
| 1 OP | Albey |
| 1 OP | Hymenoptera S29 |
| e – Retail pharn | nacy |
| | |
| 1 OP | Albey |
| | |
| 1 OP | Hymenoptera S29 |
| | |
| 1 OP | Venomil S29 |
| | |
| 1 OP | Hymenoptera S29 |
| | |
| 1 OP | Albey |
| | |
| 1 OP | Venomil S29 |
| | 1 OP 1 OP 1 OP 1 OP 1 OP e – Retail pharm 1 OP 1 OP 1 OP |

| | Subsidy | | Fully | Brand or |
|---|-------------------|-------------|---------|---------------|
| | (Manufacturer's F | Price) Sub | sidised | |
| | \$ | Per | 1 | Manufacturer |
| Antikistowing | | | | |
| Antihistamines | | | | |
| CETIRIZINE HYDROCHLORIDE | | | | |
| * Tab 10 mg | 1.12 | 100 | 1 | Zista |
| * Oral liq 1 mg per ml | 2.84 | 200 ml | 1 | Histaclear |
| CHLORPHENIRAMINE MALEATE | | | | |
| * Oral lig 2 mg per 5 ml | 9.37 | 500 ml | 1 | Histafen |
| | | | | |
| * Tab 2 mg | 2.02 | 40 | | |
| 4. Tao 2 mg | (8.40) | 10 | | Polaramine |
| | 1.01 | 20 | | |
| | (5.99) | | | Polaramine |
| * Oral liq 2 mg per 5 ml | | 100 ml | | |
| | (10.29) | | | Polaramine |
| FEXOFENADINE HYDROCHLORIDE | , , | | | |
| * Tab 60 mg | 4.34 | 20 | | |
| | (8.23) | 20 | | Telfast |
| * Tab 120 mg | | 10 | | londot |
| · · · · · · · · · · · · · · · · · · · | (8.23) | | | Telfast |
| | 14.22 | 30 | | |
| | (26.44) | | | Telfast |
| LORATADINE | , , | | | |
| * Tab 10 mg | 1 69 | 100 | 1 | Lorafix |
| * Oral liq 1 mg per ml | | 100 ml | | Haylor syrup |
| PROMETHAZINE HYDROCHLORIDE | | | | |
| * Tab 10 mg | 1 30 | 50 | 1 | Allersoothe |
| * Tab 10 mg | | 50 | | Allersoothe |
| * Oral liq 1 mg per 1 ml | | 100 ml | | Allersoothe |
| Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a P | | 5 | | Hospira |
| | | ů. | | |
| Inhaled Corticosteroids | | | | |
| BECLOMETHASONE DIPROPIONATE | | | | |
| Aerosol inhaler, 50 mcg per dose | 14 01 | 200 dose OP | 1 | Qvar |
| Aerosol inhaler, 50 mcg per dose CFC-free | | 200 dose OP | | Beclazone 50 |
| Aerosol inhaler, 100 mcg per dose | | 200 dose OP | | Qvar |
| Aerosol inhaler, 100 mcg per dose CFC-free | | 200 dose OP | 1 | Beclazone 100 |
| Aerosol inhaler, 250 mcg per dose CFC-free | | 200 dose OP | 1 | Beclazone 250 |
| BUDESONIDE | | | | |
| Powder for inhalation, 100 mcg per dose | 17 00 | 200 dose OP | 1 | Pulmicort |
| Towaci for initialation, foo fileg per dose | | 200 0030 01 | • | Turbuhaler |
| Powder for inhalation, 200 mcg per dose | 10.00 | 200 dose OP | 1 | Pulmicort |
| r owder for initialation, 200 meg per dose | | | • | Turbuhaler |
| Powder for inhalation, 400 mcg per dose | 32 00 | 200 dose OP | 1 | Pulmicort |
| r owder for initialation, 400 mby per dose | 02.00 | | • | Turbuhaler |
| | | | | |
| | | | | |

| | <u> </u> | | | |
|---|----------------------------|--------------|---------------------|---------------------|
| | Subsidy (Manufacturer's | Price) | Fully Subsidised | |
| | (Manufacturer's | Price) Pe | | |
| FLUTICASONE | | | | |
| Aerosol inhaler, 50 mcg per dose | 7 10 | 120 dos | | Flixotide |
| Powder for inhalation, 50 mcg per dose | | 60 dos | | Flixotide Accuhaler |
| Powder for inhalation, 30 mcg per dose | | 60 dos | | Flixotide Accuhaler |
| Aerosol inhaler, 125 mcg per dose | | 120 dos | | Flixotide |
| Aerosol inhaler, 725 mcg per dose | | 120 dos | | Flixotide |
| Powder for inhalation, 250 mcg per dose | | 60 dos | | Flixotide Accuhaler |
| Inhaled Long-acting Beta-adrenoceptor Agonist | s | | | |
| innared zong deting beta dareneoopter Ageniet | | | | |
| EFORMOTEROL FUMARATE | | | | |
| Powder for inhalation, 12 mcg per dose, and monodose device | ce20.64 | 60 do | ose | |
| | (35.80) | | | Foradil |
| EFORMOTEROL FUMARATE DIHYDRATE | | | | |
| Powder for inhalation 4.5 mcg per dose, breath activated | | | | |
| (equivalent to eformoterol fumarate 6 mcg metered dose |) 10.32 | 60 dos | ≏ OP | |
| | (16.90) | 00 000 | | Oxis Turbuhaler |
| | (10.00) | | | |
| NDACATEROL | C1 00 | 00 dee | | Onbras Brassbalar |
| Powder for inhalation 150 mcg | | 30 dos | | Onbrez Breezhaler |
| Powder for inhalation 300 mcg | 61.00 | 30 dos | e OP 🗸 | Onbrez Breezhaler |
| SALMETEROL | | | | |
| Aerosol inhaler CFC-free, 25 mcg per dose | | 120 dos | | Serevent |
| Powder for inhalation, 50 mcg per dose, breath activated | | 60 dos | e OP 🗸 | Serevent Accuhaler |
| Inhaled Corticosteroids with Long-Acting Beta- | Adrenocept | tor Ago | nists | |
| BUDESONIDE WITH EFORMOTEROL | | | | |
| | | | | |
| Powder for inhalation 160 mcg with 4.5 mcg eformoterol | ith | | | |
| fumarate per dose (equivalent to 200 mcg budesonide w | | 120 dos | | Due Deen Chiremey |
| 6 mcg eformoterol fumarate metered dose) | | 120 008 | se OP 🔮 | DuoResp Spiromax |
| Powder for inhalation 320 mcg with 9 mcg eformoterol fumara | | | | |
| per dose (equivalent to 400 mcg budesonide with 12 mcg | 9 | | | |
| eformoterol fumarate metered dose) - No more than 2 | | 100 1 | <u></u> | |
| dose per day | | 120 dos | | DuoResp Spiromax |
| Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg | | 120 dos | | Vannair |
| Powder for inhalation 100 mcg with eformoterol fumarate 6 m | icg33.74 | 120 dos | se OP 🗸 | Symbicort |
| | | | | Turbuhaler 100/6 |
| Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg | | 120 dos | | Vannair |
| Powder for inhalation 200 mcg with eformoterol fumarate 6 m | icg44.08 | 120 dos | se OP 🗸 | Symbicort |
| | | | | Turbuhaler 200/6 |
| Powder for inhalation 400 mcg with eformoterol fumarate | | | | |
| 12 mcg - No more than 2 dose per day | | 60 dos | e OP 🛛 🗸 | Symbicort |
| | | | | Turbuhaler 400/12 |
| FLUTICASONE FUROATE WITH VILANTEROL | | | | |
| Powder for inhalation 100 mcg with vilanterol 25 mcg | | 30 dos | e OP 🖌 | Breo Ellipta |
| · ···································· | | 00 000 | | milete |

| | Subsidy | | Fully | Brand or |
|--|-----------------|-----------|------------|---------------------|
| | (Manufacturer's | Price) | Subsidised | |
| | \$ | Per | ~ | Manufacturer |
| FLUTICASONE WITH SALMETEROL | | | | |
| Aerosol inhaler 50 mcg with salmeterol 25 mcg | 25 70 | 120 dose | op 🖌 | Seretide |
| | | | - | |
| Aerosol inhaler 125 mcg with salmeterol 25 mcg | | 120 dose | UP V | <u>Seretide</u> |
| Powder for inhalation 100 mcg with salmeterol 50 mcg - No | | | _ | |
| more than 2 dose per day | | 60 dose (| DP 🗸 | Seretide Accuhaler |
| Powder for inhalation 250 mcg with salmeterol 50 mcg - No | | | | |
| more than 2 dose per day | | 60 dose (| DP 🗸 | Seretide Accuhaler |
| , , | | | | |
| Beta-Adrenoceptor Agonists | | | | |
| SALBUTAMOL | | | | |
| Oral lig 400 mcg per ml | 40.00 | 150 ml | 1 | Ventolin |
| Infusion 1 mg per ml, 5 ml | | 10 | | Ventolin |
| Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO | | 5 | | Ventolin |
| ing 500 meg per mi, 1 mi – Op to 5 mij avaliable on a PSO | | 5 | • | ventonn |
| Inhaled Beta-Adrenoceptor Agonists | | | | |
| SALBUTAMOL | | | | |
| | | | | |
| Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO | 0.00 | 000 daaa | | Deenland |
| | | 200 dose | - · | Respigen |
| | (2.22) | | ~ | SalAir |
| | (6.20) | | | Ventolin |
| Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb | | | | |
| available on a PSO | 8.96 | 20 | ✓ | Asthalin |
| Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb | | | | |
| available on a PSO | 9.43 | 20 | 1 | Asthalin |
| TERBUTALINE SULPHATE | | | | |
| | | | | |
| Powder for inhalation, 200 mcg per dose (equivalent to | ~~~~ | 100 1 | <u></u> | <u>.</u> |
| 250 mcg metered dose), breath activated | | 120 dose | OP 🗸 | Bricanyl Turbuhaler |
| Antichalizzania Anonto | | | | |
| Anticholinergic Agents | | | | |
| IPRATROPIUM BROMIDE | | | | |
| | - | | | |
| Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose | | 000 | | A |
| available on a PSO | | 200 dose | OP 🗸 | Atrovent |
| Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne | | | _ | |
| available on a PSO | 11.73 | 20 | 1 | Univent |
| Inhaled Beta-Adrenoceptor Agonists with Anticl | holinergic / | Agents | | |
| | • | - | | |
| SALBUTAMOL WITH IPRATROPIUM BROMIDE | | | | |
| Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p | | | | |
| dose CFC-free | 12.19 | 200 dose | ор 🗸 | Duolin HFA |
| Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per | | | | |
| vial, 2.5 ml ampoule - Up to 20 neb available on a PSO | 11.04 | 20 | 1 | Duolin |
| | | | | |

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|------------------------------|---|---|
| Long-Acting Muscarinic Antagonists | | | | |
| GLYCOPYRRONIUM – Subsidy by endorsement a) Inhaled glycopyrronium treatment will not be subsidised umeclidinium. b) Glycopyrronium powder for inhalation 50 mcg per dose is having COPD using spirometry if spirometry is possible, Powder for inhalation 50 mcg per dose | s subsidised only for and the prescription 61.00 3 | patient is endo 0 dose | s who have orsed accord OP ✓ S | e been diagnosed as dingly. Seebri Breezhaler |
| umeclidinium. b) Tiotropium bromide is subsidised only for patients who h spirometry is possible, and the prescription is endorsed a 1 October 2018 with a valid Special Authority are deeme Powder for inhalation, 18 mcg per dose | accordingly. Patient d endorsed. 50.37 | s who h 30 dos | iad tiotropiu e 🖌 🖌 S | m dispensed before |
| Soln for inhalation 2.5 mcg per dose UMECLIDINIUM – Subsidy by endorsement a) Umeclidinium will not be subsidised if patient is also recertiotropium bromide. b) Umeclidinium powder for inhalation 62.5 mcg per dose is COPD using spirometry if spirometry is possible, and the Powder for inhalation 62.5 mcg per dose | iving treatment with s subsidised only for prescription is endo | patient orsed ad | ised inhaled s who have ccordingly. | |
| Long-Acting Muscarinic Antagonists with Long | -Acting Beta-A | drend | ceptor A | Agonists |

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

| Initial application from any relevant practitioner. | Approvals valid for 2 years for applications meeting the following criteria: |
|---|--|
| Both: | |

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

| GLYCOPYRRONIUM WITH INDACATEROL - Special Authority se | | | , |
|---|------------|--------------------|--|
| Powder for Inhalation 50 mcg with indacaterol 110 mcg | 81.00 | 30 dose OP | Ultibro Breezhaler |
| TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority | see SA1584 | 4 above – Retail p | pharmacy |
| Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg | 81.00 | 60 dose OP | Spiolto Respimat |
| UMECLIDINIUM WITH VILANTEROL - Special Authority see SA15 | 84 above – | Retail pharmacy | |
| Powder for inhalation 62.5 mcg with vilanterol 25 mcg | 77.00 | 30 dose OP | Anoro Ellipta |

Antifibrotics

| NINTEDANIB – Special Authority see SA2012 on the next page – Retail pharmacy | | | | | |
|--|------------------------|-------|--------------------------|--|--|
| Note: Nintedanib not subsidised in combination with s | ubsidised pirfenidone. | | | | |
| Cap 100 mg | 2,554.00 | 60 OP | Ofev | | |
| Cap 150 mg | | 60 OP | Ofev | | |

| | Subsidy | F | ully | Brand or |
|----|----------------------|---------|------|--------------|
| (M | anufacturer's Price) | Subsidi | sed | Generic |
| | \$ | Per | ✓ | Manufacturer |

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

| Note: Pirfenidone is not subsidised in combination with su | bsidised nintedanib. | | |
|--|----------------------|----|-----------------------------|
| Tab 801 mg | | 90 | Esbriet |
| Tab 267 mg | 1,215.00 | 90 | 🗸 Esbriet |

➡SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

| | Subsidy (Manufacturer's Price) \$ | Su Per | Fully bsidised | Brand or Generic Manufacturer |
|---|--|--------------------------|-------------------|---|
| Leukotriene Receptor Antagonists | | | | |
| MONTELUKAST * Tab 4 mg * Tab 5 mg * Tab 10 mg | 4.25 | 28 28 28 | ✓ | Montelukast Mylan Montelukast Mylan Montelukast Mylan |
| Methylxanthines | | | | |
| AMINOPHYLLINE * Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on PSO | | 5 | 1 | DBL Aminophylline |
| THEOPHYLLINE * Tab long-acting 250 mg * Oral liq 80 mg per 15 ml | | 100 500 ml | | <u>Nuelin-SR</u> Nuelin |
| Mucolytics | | | | |
| DORNASE ALFA – Special Authority see SA1978 below – Rei Nebuliser soln, 2.5 mg per 2.5 ml ampoule | 250.00 | 6 cian. App | | Pulmozyme valid for 12 months for |
| applications meeting the following criteria: All of the following: Patient has a confirmed diagnosis of cystic fibrosis; and Patient has previously undergone a trial with, or is curre Any of the following: | | , hypertor | nic salin | e; and |
| 3.1 Patient has required one or more hospital inpatie 3.2 Patient has had 3 exacerbations due to CF, requireriod; or 3.3 Patient has had 1 exacerbation due to CF, requirering Brasfield score of < 22/25; or | iring oral or intravenouring oral or IV antibiotic | is (IV) an s in the p | tibiotics | in the previous 12 month |
| 3.4 Patient has a diagnosis of allergic bronchopulmo Renewal — (cystic fibrosis) only from a respiratory physiciar notified where the treatment remains appropriate and the patien | or paediatrician. App | rovals va | | out further renewal unless |
| IVACAFTOR – PCT only – Specialist – Special Authority see S Tab 150 mg Oral granules 50 mg, sachet Oral granules 75 mg, sachet | 29,386.00 29,386.00 | 56 56 56 | ✓ | Kalydeco Kalydeco Kalydeco |
| SA2017 Special Authority for Subsidy Initial application only from a respiratory specialist or paediatr applications meeting the following criteria: All of the following: | ician. Approvals valid | without f | urther re | enewal unless notified for |
| Patient has been diagnosed with cystic fibrosis; and Either: 2.1 Patient must have G551D mutation in the cystic | fibrosis transmembran | e conduc | tance re | egulator (CFTR) gene on at |
| least 1 allele; or 2.2 Patient must have other gating (class III) mutatic | n (G1244F, G1349D, | G178B (| 3551S. | S1251N S1255P S549N |

| | ATONT STSTEM AND ALLENGIES |
|---|--|
| Subsid (Manufacturer \$ | |
| continued | |
| and S549R) in the CFTR gene on at least 1 allele; and Patients must have a sweat chloride value of at least 60 mmol/L by qua sweat collection system; and Treatment with ivacaftor must be given concomitantly with standard the Patient must not have an acute upper or lower respiratory infection, pul (including antibiotics) for pulmonary disease in the last 4 weeks prior to The dose of ivacaftor will not exceed one tablet or one sachet twice dai Applicant has experience and expertise in the management of cystic fib | erapy for this condition; and Imonary exacerbation, or changes in therapy o commencing treatment with ivacaftor; and ily; and |
| SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7%24.50 | 90 ml OP ✓ <u>Biomed</u> |
| Nasal Preparations | |
| Allergy Prophylactics | |
| BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose | |
| FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose1.98 | 120 dose OP ✓ <u>Flixonase Hayfever</u> <u>& Allergy</u> |
| IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%5.23 | 15 ml OP 🖌 <u>Univent</u> |
| Respiratory Devices | |
| MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under Small | 1 ✔ e-chamber Mask |
| PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO | |
| Low range | 1 ✓ Mini-Wright AFS Low Range |
| Normal range9.54 | • |
| SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO | |
| 220 ml (single patient)2.95510 ml (single patient)5.12 | |
| 800 ml6.50 | |

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|---------|-----------------------|-------------------------------------|
| Respiratory Stimulants | | | | |
| CAFFEINE CITRATE Oral liq 20 mg per ml (10 mg base per ml) | | 5 ml Ol | ⊳ √ <u>B</u> i | iomed |

SENSORY ORGANS

| | Cubaidu | | Fully Brand or |
|--|------------------------------|------------|---|
| | Subsidy (Manufacturer's P | rica) Subs | Fully Brand or sidised Generic |
| | (Manulacialer ST | Per | ✓ Manufacturer |
| | * | - | |
| Ear Preparations | | | |
| FLUMETASONE PIVALATE | | | |
| Ear drops 0.02% with clioquinol 1% | 4.46 | 7.5 ml OP | Locacorten-Viaform ED's |
| | | | Locorten-Vioform |
| TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI | | INI | |
| | N AND N ISTAT | IIN | |
| Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g | 5 16 | 7.5 ml OP | ✓ Kenacomb |
| | | 7.5 III OF | • Reliacollib |
| Ear/Eye Preparations | | | |
| DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN | | | |
| Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and | | | |
| gramicidin 50 mcg per ml | 4.50 | 8 ml OP | |
| gramicium so meg per mi | | 0 IIII OF | Sofradex |
| | (9.27) | | Solladex |
| FRAMYCETIN SULPHATE | 4.40 | | |
| Ear/Eye drops 0.5% | | 8 ml OP | Catramusia |
| | (8.65) | | Soframycin |
| Eye Preparations | | | |
| | | | |
| Eye preparations are only funded for use in the eye, unless explic | itly stated otherw | vise. | |
| Anti-Infective Preparations | | | |
| Anti-Intective Treparations | | | |
| ACICLOVIR | | | |
| * Eye oint 3% | 14.88 | 4.5 g OP | ✓ <u>ViruPOS</u> |
| CHLORAMPHENICOL | | | |
| Eve oint 1% | | 5 g OP | Devatis |
| Eve drops 0.5% | | 10 ml OP | Chlorafast |
| Funded for use in the ear*. Indications marked with * are | | dications. | |
| CIPROFLOXACIN | | | |
| Eye drops 0.3% – Subsidy by endorsement | | 5 ml OP | Ciprofloxacin Teva |
| When prescribed for the treatment of bacterial keratitis of | | | |
| for the second line treatment of chronic suppurative otitis | | , | 1 / |
| Note: Indication marked with a * is an unapproved indica | () | | , |
| GENTAMICIN SULPHATE | | | |
| Eye drops 0.3% | 11.40 | 5 ml OP | ✓ Genoptic |
| | | | |
| PROPAMIDINE ISETHIONATE * Eye drops 0.1% | 2.07 | 10 ml OP | |
| - x ⊂ye ui∪µs ∪. i % | | | Brolene |
| | (14.55) | | DIOIEITE |
| SODIUM FUSIDATE [FUSIDIC ACID] | F 00 | 5 00 | |
| Eye drops 1% | 5.29 | 5 g OP | Fucithalmic |
| TOBRAMYCIN | | | |
| Eye oint 0.3% | | 3.5 g OP | Tobrex |
| Eye drops 0.3% | 11.48 | 5 ml OP | Tobrex |
| | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

| | Subsidy (Manufacturer's Pri | ne) Suba | | rand or Generic |
|--|--------------------------------|----------------|-------------------------|-------------------------|
| | | Per | | lanufacturer |
| Corticosteroids and Other Anti-Inflammatory Pr | eparations | | | |
| DEXAMETHASONE | | | | |
| Eye oint 0.1% | | 3.5 g OP | Max | |
| ₭ Eye drops 0.1% | | 5 ml OP | 🗸 Max | Idex |
| Ocular implant 700 mcg – Special Authority see SA1680 belo – Retail pharmacy | | 1 | 🗸 Ozu | rdex |
| ⇒SA1680 Special Authority for Subsidy | - | | | |
| nitial application — (Diabetic macular oedema) only from an neeting the following criteria: All of the following: | ophthalmologist. | Approvals va | llid for 12 n | nonths for applications |
| Patient has diabetic macular oedema with pseudophakic le | ans: and | | | |
| Patient has reduced visual acuity of between 6/9 - 6/48 with Either: | | eness of redu | uction in vis | sion; and |
| 3.1 Patient's disease has progressed despite 3 injectio3.2 Patient is unsuitable or contraindicated to treatmen | | | | |
| 4 Dexamethasone implants are to be administered not more maximum of 3 implants per eye per year. | frequently than o | nce every 4 r | months into | each eye, and up to a |
| Renewal — (Diabetic macular oedema) only from an ophthalm | ologist. Approval | s valid for 12 | months for | r applications meeting |
| he following criteria: 3oth: | | | | |
| 1 Patient's vision is stable or has improved (prescriber deter | mined): and | | | |
| Dexamethasone implants are to be administered not more maximum of 3 implants per eye per year. | | nce every 4 r | months into | each eye, and up to a |
| nitial application — (Women of child bearing age with diabet | ic macular oeder | ma) only fror | n an ophth | almologist. Approvals |
| ralid for 12 months for applications meeting the following criteria: All of the following: | | | | |
| 1 Patient has diabetic macular oedema; and | | | | |
| 2 Patient has reduced visual acuity of between 6/9 - 6/48 with | | eness of redu | uction in vis | sion; and |
| 3 Patient is of child bearing potential and has not yet comple 4 Dexamethasone implants are to be administered not more maximum of 3 implants per eye per year. | | nce every 4 r | months into | each eye, and up to a |
| Renewal — (Women of child bearing age with diabetic macul | ar oedema) onlv | from an opht | halmologis | t. Approvals valid for |
| 2 months for applications meeting the following criteria: All of the following: | | | | |
| 1 Patient's vision is stable or has improved (prescriber deter | mined); and | | | |
| 2 Patient is of child bearing potential and has not yet comple | | | | |
| 3 Dexamethasone implants are to be administered not more maximum of 3 implants per eye per year. | frequently than o | nce every 4 r | months into | each eye, and up to |
| DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYN | IYXIN B SULPHA | TE | | |
| Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin sulphate 6,000 u per g | | 3.5 g OP | 🗸 Max | itrol |
| ₭ Eye drops 0.1% with neomycin sulphate 0.35% and polymyxi | | 5.5 y Oi | - widk | |
| b sulphate 6,000 u per ml | | 5 ml OP | 🗸 Max | itrol |
| DICLOFENAC SODIUM | | | | |
| Eye drops 0.1% | 8.80 | 5 ml OP | ✓ Volt | aren Ophtha |
| LUOROMETHOLONE | | | | |
| ₭ Eye drops 0.1% | | 5 ml OP | 🖌 FML | |
| | 5 20 | | Flue | on |

Flucon

SENSORY ORGANS

| | Subsidy | | Fully Brand or |
|--|---------------------|-----------------|------------------------------------|
| | (Manufacturer's F | | idised Generic |
| | \$ | Per | Manufacturer |
| LEVOCABASTINE Eye drops 0.5 mg per ml | 8 71 | 4 ml OP | |
| | (10.34) | 4 111 01 | Livostin |
| LODOXAMIDE | (1000) | | |
| Eye drops 0.1% | | 10 ml OP | ✓ Lomide |
| PREDNISOLONE ACETATE | | | |
| Eye drops 1% | 5.93 | 10 ml OP | Prednisolone-AFT |
| | 7.00 | 5 ml OP | Pred Forte |
| PREDNISOLONE SODIUM PHOSPHATE – Special Authority | see SA1715 below | – Retail pharr | nacy |
| Eye drops 0.5%, single dose (preservative free) | | 20 dose | Minims |
| | | | Prednisolone |
| SA1715 Special Authority for Subsidy | | o | |
| Initial application only from an ophthalmologist or optometrist following criteria: | . Approvals valid f | or 6 months for | r applications meeting the |
| Both: | | | |
| 1 Patient has severe inflammation; and | | | |
| 2 Patient has a confirmed allergic reaction to preservative | e in eye drops. | | |
| Renewal from any relevant practitioner. Approvals valid for 6 | months where the | reatment rema | ins appropriate and the patient is |
| penefiting from treatment. | | | |
| SODIUM CROMOGLICATE | | | |
| Eye drops 2% | 1.79 | 5 ml OP | Rexacrom |
| Glaucoma Preparations - Beta Blockers | | | |
| BETAXOLOL | | | |
| ■ TAXOLOL ¥ Eye drops 0.25% | 11.80 | 5 ml OP | Betoptic S |
| * Eye drops 0.5% | | 5 ml OP | ✓ Betoptic |
| TIMOLOL | | | |
| * Eye drops 0.25% | 1.81 | 5 ml OP | Arrow-Timolol |
| * Eye drops 0.5% | | 5 ml OP | ✓ Arrow-Timolol |
| * Eye drops 0.5%, gel forming | 3.78 | 2.5 ml OP | Timoptol XE |
| Glaucoma Preparations - Carbonic Anhydrase | e Inhibitors | | |
| ACETAZOLAMIDE | | | |
| * Tab 250 mg | | 100 | Diamox |
| BRINZOLAMIDE | | | |
| * Eye drops 1% | 7.30 | 5 ml OP | ✓ Azopt |
| DORZOLAMIDE HYDROCHLORIDE | | | |
| * Eye drops 2% | 9.77 | 5 ml OP | |
| | (17.44) | | Trusopt |
| DORZOLAMIDE WITH TIMOLOL | | | |
| * Eye drops 2% with timolol 0.5% | 2.73 | 5 ml OP | Dortimopt |
| Glaucoma Preparations - Prostaglandin Analo | ogues | | |
| BIMATOPROST | | | |
| * Eye drops 0.03% | 5.95 | 3 ml OP | Bimatoprost |
| | | | Multichem |
| | | | |

A Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

SENSORY ORGANS

| (M | Subsidy Ianufacturer's P \$ | rice) Subsi Per | Fully Brand or idised Generic ✓ Manufacturer |
|--|-----------------------------------|--------------------|--|
| ATANOPROST | | | |
| * Eye drops 0.005% | 1.82 | 2.5 ml OP | ✓ <u>Teva</u> |
| TRAVOPROST | | | |
| * Eye drops 0.004% | 9.75 | 2.5 ml OP | <u>Travatan</u> |
| Glaucoma Preparations - Other | | | |
| BRIMONIDINE TARTRATE | | | |
| ₭ Eye drops 0.2% | 4.29 | 5 ml OP | Arrow-Brimonidine |
| BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE | | | |
| ₭ Eye drops 0.2% with timolol maleate 0.5% | 18.50 | 5 ml OP | Combigan |
| ATANOPROST WITH TIMOLOL | | | |
| Eye drops 0.005% with timolol 0.5% | 2.49 | 2.5 ml OP | Arrow - Lattim |
| PILOCARPINE HYDROCHLORIDE | | | |
| Eye drops 1% | 4.26 | 15 ml OP | Isopto Carpine |
| Eye drops 2% | 5.35 | 15 ml OP | Isopto Carpine |
| * Eye drops 4% | | 15 ml OP | Isopto Carpine |
| Subsidised for oral use pursuant to the Standard Formulae. | | | |
| ₭ Eye drops 2% single dose – Special Authority see SA0895 | | | * |
| below – Retail pharmacy | 31.95 | 20 dose | Minims Pilocarpine |

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Patient has to use an unpreserved solution due to an allergy to the preservative; or

2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

| ATROPINE SULPHATE * Eye drops 1% | 15 ml OP | ✔ Atront |
|-------------------------------------|-----------|-------------------------------|
| CYCLOPENTOLATE HYDROCHLORIDE | 13 111 01 | • <u>Auopi</u> |
| * Eye drops 1% | 15 ml OP | Cyclogyl |
| TBOPICAMIDE | | e jele gji |
| * Eye drops 0.5% | 15 ml OP | Mydriacyl |
| * Eye drops 1%8.66 | | Mydriacyl |
| | | |

Preparations for Tear Deficiency

| For acetylcysteine eye drops refer Standard Formulae, page 244 | | | |
|--|-------|----------|--------------------------------|
| HYPROMELLOSE | | | |
| * Eye drops 0.5% | 19.50 | 15 ml OP | Methopt |
| HYPROMELLOSE WITH DEXTRAN | | | |
| * Eye drops 0.3% with dextran 0.1% | 2.30 | 15 ml OP | Poly-Tears |

| | | SEN | SORY ORGANS |
|---|---|------------------------------|-------------------------------------|
| | Subsidy (Manufacturer's Price) \$ | Fully Subsidised Per ✓ | Brand or Generic Manufacturer |
| Preservative Free Ocular Lubricants | φ | | |
| SA1388 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals vali Both: | d for 12 months for ap | oplications meetir | ng the following criteria: |
| Confirmed diagnosis by slit lamp of severe secretory dry 2 Either: | eye; and | | |
| 2.1 Patient is using eye drops more than four times da 2.2 Patient has had a confirmed allergic reaction to pr | , , | | |
| Renewal from any relevant practitioner. Approvals valid for 24 r drops and has benefited from treatment. | nonths where the patie | ent continues to | require lubricating eye |
| CARBOMER – Special Authority see SA1388 above – Retail ph Ophthalmic gel 0.3%, 0.5 g | | 30 🖌 F | Poly-Gel |

| 1 6 7 6 | | |
|---|---------------------|---------------------------------------|
| MACROGOL 400 AND PROPYLENE GLYCOL - Special Authority see SA1 | 388 above - Retail | pharmacy |
| Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml4.3 | 0 24 | Systane Unit Dose |
| SODIUM HYALURONATE [HYALURONIC ACID] - Special Authority see S | A1388 above – Reta | il pharmacy |
| Eye drops 1 mg per ml13.8 | | |
| Hylo-Fresh has a 6 month expiry after opening. The Pharmacy Pro | cedures Manual rest | triction allowing one bottle per |
| month is not relevant and therefore only the prescribed dosage to the | ie nearest OP may b | e claimed. |

Other Eye Preparations

| NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1% 4.15 | 15 ml OP | Naphcon Forte |
|---|----------|-----------------------------------|
| OLOPATADINE Eye drops 0.1%2.20 | 5 ml OP | ✓ Olopatadine Teva |
| PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3% | 3.5 g OP | Poly-Visc |
| RETINOL PALMITATE Eye oint 138 mcg per g | 5 g OP | ✓ VitA-POS |

| | Subsidy | | Fully | Brand or |
|--|----------------------------------|-----------|----------|-------------------|
| | (Manufacturer's Price | | bsidised | Generic |
| | \$ | Per | | Manufacturer |
| Various | | | | |
| PHARMACY SERVICES | | | | |
| May only be claimed once per patient. | | | | |
| Brand switch fee | 4 50 | 1 fee | ✓ B | SF Cinacalcet |
| | 4.50 | Tiee | • 0 | Devatis |
| | | | | |
| | | | V B | SF Clomipramine |
| | | | | Teva |
| | | | | SF Comtan |
| | | | ✓ B | SF Lopinavir/ |
| | | | | Ritonavir Mylan |
| | | | 🗸 В | SF Vigisom |
| a) The Pharmacode for BSF Lopinavir/Ritonavir Mylan | is 2621959 - see als | o page 10 |)6 | |
| b) The Pharmacode for BSF Clomipramine Teva is 263 | 0915 - see also <mark>pag</mark> | e 125 | | |
| c) The Pharmacode for BSF Vigisom is 2634112 - see | | | | |
| d) The Pharmacode for BSF Cinacalcet Devatis is 2634 | 120 - see also page | 78 | | |
| e) The Pharmacode for BSF Comtan is 2634139 - see | also page 119 | | | |
| (BSF Cinacalcet Devatis Brand switch fee to be delisted 1 July 2 | 022) | | | |
| (BSF Clomipramine Teva Brand switch fee to be delisted 1 May | , | | | |
| (BSF Comtan Brand switch fee to be delisted 1 July 2022) | / | | | |
| (BSF Lopinavir/Ritonavir Mylan Brand switch fee to be delisted 1 | May 2022) | | | |
| (BSF Vigisom Brand switch fee to be delisted 1 July 2022) | may 2022/ | | | |
| | | | | |
| Agents Used in the Treatment of Poisonings | | | | |
| Antidotes | | | | |
| ACETYLCYSTEINE | | | | |
| Inj 200 mg per ml, 10 ml ampoule | | 10 | 🗸 D | BL Acetylcysteine |
| j j , , , | | | | lartindale |
| | | | | Pharma S29 |
| | | | | T Humu 👄 |
| NALOXONE HYDROCHLORIDE | | | | |
| a) Up to 5 inj available on a PSO | | | | |
| b) Only on a PSO | | _ | | |
| * Inj 400 mcg per ml, 1 ml ampoule | 22.60 | 5 | ✓ D | BL Naloxone |
| | | | | Hydrochloride |
| Removal and Elimination | | | | |
| CHARCOAL | | | | |
| * Oral liq 50 g per 250 ml | 13 50 0 | 50 ml OP | 10 | arbosorb-X |
| | | JU III UP | • 0 | ai 00301 D-V |
| a) Up to 250 ml available on a PSO | | | | |
| b) Only on a PSO | | | | |
| DEFERASIROX – Special Authority see SA1492 on the next page | <mark>ge – Retail pharmac</mark> | у | | |
| Wastage claimable | | | | |
| Tab 125 mg dispersible | 276.00 | 28 | 🖌 E | xjade |
| Tab 250 mg dispersible | | 28 | 🖌 E | xjade |
| Tab 500 mg dispersible | | 28 | 🗸 E | xjade |
| | | | | |

| Subsidy (Manufacturer's Price) | Full Subsidise | | |
|-----------------------------------|-------------------|--------------|--|
| \$ | Per 🖌 | Manufacturer | |

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - Retail pharmacy

| Tab 500 mg | | 7 100 | Ferriprox |
|--------------------|---------|-----------------|-------------------------------|
| Oral liq 100 mg pe | ər 1 ml |) 250 ml OP | Ferriprox |

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or

2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE

| * Inj 500 mg vial | 151.31 | 10 | ✓ DBL Desferrioxamine Mesylate for Inj BP |
|---------------------------|----------|----|--|
| SODIUM CALCIUM EDETATE | | | |
| * Inj 200 mg per ml, 5 ml | 53.31 | 6 | |
| | (156.71) | | Calcium Disodium Versenate |

Standard Formulae

| ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base | qs qs | PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water | 1 g 70 ml to 100 ml |
|--|---|---|--|
| CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water | 60 mg 40 ml qs to 100 ml | PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium Glycerol BP Water | LIQUID (10 400 mg 4 ml to 40 ml |
| CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water FOLINIC MOUTHWASH Calcium folinate 15 mg tab | 300 mg 40 ml qs to 100 ml 1 tab | PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water (Preservative should be used if quantity supplied is than 5 days.) | qs qs to 500 ml for more |
| Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) METHADONE MIXTURE Methadone powder | qs to 500 ml for more qs | SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water (Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) | 5 g qs to 500 ml for more |
| Glycerol Water METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol | qs to 100 ml 10 g to 100 ml | SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water (Only funded if prescribed for treatment of hyponatr VANCOMYCIN ORAL SOLUTION (50 mg per ml) | qs qs raemia) |
| (Use 1 ml of the 10% solution per 100 ml of oral liqu OMEPRAZOLE SUSPENSION Omeprazole capules or powder Sodium bicarbonate powder BP Water | | Vancomycin 500 mg injection Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure) | 10 vials 40 ml to 100 ml um difficile |

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | <u> </u> | | | | |
|---|----------------------------------|-----------------|-------------------|---------------------|-------|
| | Subsidy (Manufacturer's Price | -) <u></u> | Fully bsidised | | |
| | (Manulacturer's Frice | Per | | Manufacturer | |
| | Ť | - | | | |
| Extemporaneously Compounded Preparations | and Galenicals | | | | |
| CODEINE PHOSPHATE - Safety medicine; prescriber may dete | ermine dispensing f | requency | | | |
| Powder – Only in combination | 63.09 | 25 g | | | |
| | (90.09) | | | Douglas | |
| Only in extemporaneously compounded codeine linctus. | | | | | |
| COLLODION FLEXIBLE | | | | | |
| Note: This product is no longer being manufactured by the s | supplier and will be | delisted fr | om the S | Schedule at a date | to be |
| determined. | | | | | |
| Collodion flexible | 19.30 | 100 ml | ~ | PSM | |
| COMPOUND HYDROXYBENZOATE - Only in combination | | | | | |
| Only in extemporaneously compounded oral mixtures. | | | | | |
| Soln | | 100 ml | ~ | Midwest | |
| GLYCERIN WITH SODIUM SACCHARIN - Only in combination | | | | | |
| Only in combination with Ora-Plus. | | | | | |
| Suspension | | 473 ml | ✓ | Ora-Sweet SF | |
| GLYCERIN WITH SUCROSE - Only in combination | | | | | |
| Only in combination with Ora-Plus. | | | | | |
| Suspension | | 473 ml | 1 | Ora-Sweet | |
| GLYCEROL | | | | | |
| * Liquid – Only in combination | 3 23 | 500 ml | 1 | healthE Glycerol I | RP |
| Only in extemporaneously compounded oral liquid prepa | | 000 1111 | - | | |
| METHADONE HYDROCHLORIDE | | | | | |
| a) Only on a controlled drug form | | | | | |
| b) No patient co-payment payable | | | | | |
| c) Safety medicine; prescriber may determine dispensing fr | equency | | | | |
| d) Extemporaneously compounded methadone will only be | | ate of the | cheanes | st form available | |
| (methadone powder, not methadone tablets). | | | onoupoe | | |
| Powder | 7.84 | 1 g | 1 | AFT | |
| METHYL HYDROXYBENZOATE | | 0 | | | |
| Powder | 8 98 | 25 g | 1 | Midwest | |
| | | 20 g | - | | |
| METHYLCELLULOSE Powder | 26.05 | 100 a | | MidWoot | |
| Suspension – Only in combination | | 100 g 473 ml | - | MidWest Ora-Plus | |
| | | | • | <u>Ula-Flus</u> | |
| METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCH | | | | Over Diamed CE | |
| Suspension | | 473 ml | • | Ora-Blend SF | |
| METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - On | | | | | |
| Suspension | | 473 ml | ~ | Ora-Blend | |
| PHENOBARBITONE SODIUM | | | | | |
| Powder – Only in combination | 52.50 | 10 g | ~ | MidWest | |
| | 325.00 | 100 g | ~ | MidWest | |
| Only in children up to 12 years | | | | | |
| PROPYLENE GLYCOL | | | | | |
| Only in extemporaneously compounded methyl hydroxybenz | zoate 10% solution. | | | | |
| Liq | 11.25 | 500 ml | ✓ | Midwest | |
| SODIUM BICARBONATE | | | | | |
| Powder BP – Only in combination | | 500 g | 1 | Midwest | |
| Only in extemporaneously compounded omeprazole and | d lansoprazole susp | ension. | | | |
| | | | | | |

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

| | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Brand or Generic Manufacturer |
|---|---|--------|---------------------|-------------------------------------|
| SYRUP (PHARMACEUTICAL GRADE) – Only in combination | | | | |
| Only in extemporaneously compounded oral liquid preparation | ns. | | | |
| Liq | 14.95 | 500 ml | 🖌 М | idwest |
| WATER | | | | |
| Tap – Only in combination | 0.00 | 1 ml | 🗸 Ta | ap water |

SECTION D: SPECIAL FOODS

Fully

Subsidy (Manufacturer's Price)

\$

Subsidised Per ✓ Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Either:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Initial application — (Inborn errors of metabolism)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. **Renewal — (Cystic fibrosis or renal failure)** only from a dietitian, relevant specialist, vocationally registered general

practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

. Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

| Powder5.29 | 400 g OP | Polycal |
|------------|----------|-----------------------------|
|------------|----------|-----------------------------|

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

| Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic | |
|-----------------------------------|-----|---------------------|---------------------|--|
| \$ | Per | | Manufacturer | |

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| CARBOHYDRATE AND FAT SU | JPPLEMENT - Special Autho | rity see SA1376 on | the previous page | ge - | Hospital pharmacy [HP3] |
|-------------------------|---------------------------|--------------------|-------------------|------|-------------------------|
| Powder (neutral) | | | 400 g OP | 1 | Duocal Super |
| | | | - | | Soluble Powder |

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

continued...

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

| Emulsion (neutral) | | 200 ml OP | ✓ Calogen |
|-----------------------|-------|-----------|--|
| | 30.75 | 500 ml OP | Calogen |
| Emulsion (strawberry) | | 200 ml OP | Calogen |
| Oil | | 500 ml OP | MCT oil (Nutricia) |
| Oil, 250 ml | | 4 OP | Liquigen |

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PROTEIN SUPPLEMENT - S | Special Authority see SA1524 above – Hospital pl | armacy [HP3] | |
|------------------------|--|--------------|-------|
| Powder | | 225 g OP | 🗸 Pro |
| | 8.95 | 227 g OP | 🗸 Re |
| | | 0 | |

 Protifar
 Resource Beneprotein

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🗸 | Manufacturer |

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority se | ee SA1095 above - | - Hospital pharm | nacy [HP3] |
|---|-------------------|------------------|-------------------------------------|
| Liquid | 3.75 | 500 ml OP | Glucerna Select |
| | 7.50 | 1,000 ml OP | Diason RTH |
| DIABETIC ORAL FEED 1KCAL/ML - Special Authority see S | A1095 above – Ho | spital pharmacy | [HP3] |
| Liquid (strawberry) | 1.50 | 200 ml OP | Diasip |
| Liquid (vanilla) | 1.50 | 200 ml OP | Diasip |
| | 2.10 | | Nutren Diabetes |

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| FAT MODIFIED FEED – Special Authority s | ee SA1525 above – Hospital pharm | acy [HP3] | |
|---|----------------------------------|-----------|-----------|
| Powder | | 400 g OP | 🗸 Monogen |

| Subsidy | Fully | Brand or | |
|------------------------|------------|--------------|--|
| (Manufacturer's Price) | Subsidised | Generic | |
| `\$ | Per 🗸 | Manufacturer | |

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL FEED 1KCAL/ML - Special Authority se | e SA1099 above - Ho | ospital pharmacy | / [HP3] |
|---|---------------------|------------------|-------------------------------|
| Powder | | 400 g OP | Kindergen |

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

| | Subsidy (Manufacturer's Price \$ | Fully e) Subsidised Per ✓ | Brand or Generic Manufacturer |
|---|--|--|--|
| continued applications meeting the following criteria: Both: | | | |
| The treatment remains appropriate and the patient is bene General Practitioners must include the name of the dietitia practitioner and date contacted. | | | registered general |
| PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority Liquid | | | Hospital pharmacy [HP3] Nutrini Energy RTH |
| PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority se Liquid | | 00 ml OP 🖌 🗸 | ospital pharmacy [HP3] Nutrini RTH Pediasure RTH |
| PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Spe pharmacy [HP3] | ecial Authority see S | SA1379 on the pre | evious page – Hospital |
| Liquid | 6.00 5 | 500 ml OP 🗸 | Nutrini Energy Multi Fibre |
| PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see Liquid (strawberry) Liquid (vanilla) | 1.60 2 | 200 ml OP 🖌 🗸 | pital pharmacy [HP3] Fortini Fortini |
| PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see S Liquid (chocolate) Liquid (strawberry) Liquid (vanilla) | 1.07 2 1.07 2 1.07 2 | 200 ml OP 200 ml OP 200 ml OP 200 ml OP 200 ml OP 200 ml OP | tal pharmacy [HP3] Pediasure Pediasure Pediasure Pediasure |
| PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special pharmacy [HP3] | | | |
| Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla) | 1.60 2 1.60 2 | 200 ml OP 200 ml OP | Fortini Multi Fibre Fortini Multi Fibre Fortini Multi Fibre Fortini Multi Fibre |
| PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 Powder | | | rmacy [HP3] Peptamen Junior |

Renal Products

► SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see SA110 | 1 above - | Hospital pharmacy | y [HP3] |
|--|-----------|-------------------|--------------|
| Liquid | 6.08 | 500 ml OP | Nepro HP RTH |
| | Subsidy | | Fully | Brand or |
|---|-------------------------|--------------------|--|--|
| | (Manufacturer's F \$ | Price) Subs Per | idised ✓ | Generic Manufacturer |
| RENAL ORAL FEED 1.8 KCAL/ML - Special Authority see SA1 | 101 on the previ | ous page – Hos | pital p | harmacy [HP3] |
| Liquid | | 220 ml OP | Image: A second s | Vepro HP (strawberry) Vepro HP (vanilla) |
| ENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110 | | | tal pha | ırmacy [HP3] |
| Liquid | 2.88 (3.31) | 237 ml OP | ١ | NovaSource Renal |
| Liquid, 200 ml bottle | | 4 OP | | NovaSource Renal |
| | (13.24) | | I | NUVASUUILE HEIIAI |
| Liquid (apricot) 125 ml | · · · | 4 OP | | Renilon 7.5 |

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Any of the following:
 - 1 malabsorption; or
 - 2 short bowel syndrome; or
 - 3 enterocutaneous fistulas; or
 - 4 eosinophilic oesophagitis; or
 - 5 inflammatory bowel disease; or
 - 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Spe Liquid | | e SA1377 abov 1,000 ml OP | |
|---|----------------|------------------------------|---|
| ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see | SA1377 above - | – Hospital phar | macy [HP3] |
| Liquid (grapefruit), 250 ml carton | 171.00 | 18 OP | Elemental 028 Extra |
| Liquid (pineapple & orange), 250 ml carton | 171.00 | 18 OP | Elemental 028 Extra |
| Liquid (summer fruits), 250 ml carton | 171.00 | 18 OP | Elemental 028 Extra |
| ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see S | A1377 above – | Hospital pharm | acy [HP3] |
| Powder (unflavoured) | 4.50 | 80 g OP | Vivonex TEN |
| SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Auth Liquid | | | , ,, , |

| Subsidy | | Fully | Brand or |
|------------------------|-----|-----------|--------------|
| (Manufacturer's Price) | S | ubsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML | - Special Authority | see SA1196 | above - | - Hospital pharmacy [HP3] |
|---|---------------------|------------|---------|---------------------------|
| Liquid | 4.00 | 500 ml OP | ✓ | Nutrini Low Energy |
| | | | | Multi Fibre |

Standard Supplements

► SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - Patient is Malnourished
 - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
 - Patient has not responded to first-line dietary measures over a 4 week period by:
 - 2.1 Increasing their food intake frequency (eg snacks between meals); or
 - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
 - 2.3 Using over the counter supplements (e.g. Complan); and

3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
 - Patient is Malnourished
 - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

| Subsidy | | Fully | Brand or | |
|------------------------------|--------|----------------|-------------------------|--|
| (Manufacturer's Price) \$ | Per St | ubsidised ✓ | Generic Manufacturer | |

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or</p>
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority

forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

SPECIAL FOODS

| | Subsidy (Manufacturer's \$ | | Fully Brand or idised Generic ✓ Manufacturer |
|--|----------------------------------|--|--|
| ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 o Liquid | 1 0 | Hospital pharmac 250 ml OP 1,000 ml OP | y [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Nutrison Energy |
| ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on Liquid | | spital pharmacy 250 ml OP 1,000 ml OP | [HP3] ✓ Isosource Standard ✓ Nutrison Standard RTH ✓ Osmolite RTH |
| ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority Liquid | | on page 254 – He 1,000 ml OP | ospital pharmacy [HP3] Vutrison 800 Complete Multi Fibre |
| ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority se Liquid | | page 254 – Hosp 1,000 ml OP | bital pharmacy [HP3] Jevity RTH Nutrison Multi Fibre |
| ENTERAL FEED WITH FIBRE 1.2KCAL/ML – Special Authority s Liquid | | n <mark>page 254</mark> – Hos 1,000 ml OP | spital pharmacy [HP3] ✓ Jevity Plus |
| ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority s Liquid | | n page 254 – Hos 1,000 ml OP | spital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre |
| ORAL FEED (POWDER) – Special Authority see SA1859 on pag Powder (chocolate) | | al pharmacy [HP 840 g OP | P3] ✓ Sustagen Hospital |
| Powder (vanilla) | 26.00 14.00 | 850 g OP 840 g OP | Formula ✓ Ensure ✓ Sustagen Hospital Formula Active |
| | 26.00 | 850 g OP | ✓ Ensure |

| | Subsidy (Manufacturer's Pi \$ | | Fully Brand or ised Generic Manufacturer |
|---|-------------------------------------|-----------------------------------|--|
| DRAL FEED 1.5KCAL/ML – Special Authority see SA1859 on pa Additional subsidy by endorsement is available for patients b epidermolysis bullosa, or as exclusive enteral nutrition in chilk disease, or for patients with COPD and hypercapnia, defined endorsed accordingly. | eing bolus fed the | rough a feeding ge of 18 years fo | tube, who have severe or the treatment of Crohn's |
| Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Ensure Plus |
| | (1.26) | | Fortisip |
| Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Ensure Plus |
| | (1.26) | | Fortisip |
| Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 r | | | |
| with Endorsement | | 200 ml OP | |
| | (1.26) | | Ensure Plus |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with | า | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Fortisip |
| Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml w | ith | | |
| Endorsement | 0.85 | 237 ml OP | |
| | (1.33) | | Ensure Plus |
| | 0.72 | 200 ml OP | |
| | (1.26) | | Ensure Plus |
| | (1.26) | | Fortisip |
| DRAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see Additional subsidy by endorsement is available for patients be epidermolysis bullosa. The prescription must be endorsed as Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with | eing bolus fed the | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| Liquid (vanilla) – Higher subsidy of \$1.26 per 200 ml with | | | |
| Endorsement | 0.72 | 200 ml OP | |
| | (1.26) | | Fortisip Multi Fibre |
| | | | |

SA1195 Special Authority for Subsidy Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

| Subsidy (Manufacturer's | Price) | Fully Subsidised | Brand or Generic | |
|----------------------------|--------|---------------------|---------------------|--|
| \$ | Per | ✓ | Manufacturer | |

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- All of the following:
 - 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
 - 2 other lower calorie products have been tried; and
 - 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195 on th | e previous p | <mark>age</mark> – Hospital p | harmacy [HP3] |
|--|----------------|-------------------------------|---|
| Liquid | 5.50 | 500 ml OP | Nutrison |
| | | | Concentrated |
| | 11.00 | 1,000 ml OP | Ensure Two Cal HN RTH |
| ORAL FEED 2 KCAL/ML – Special Authority see SA1195 on the pre Additional subsidy by endorsement is available for patients being epidermolysis bullosa. The prescription must be endorsed acco Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with | g bolus fed t | | |
| Endorsement | 0.96 (1.90) | 200 ml OP | Two Cal HN |

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| | Subsidy (Manufacturer's Prio \$ | ce) Per | Fully Subsidised | |
|---|---------------------------------------|---------------------------------|---------------------|---|
| FOOD THICKENER – Special Authority see SA1106 on the prev Powder | rious page – Hosp 6.53 7.25 | ital phar 300 g (380 g (| DP 🍈 🗸 | 3] Nutilis Feed Thickener Karicare Aptamil |

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

| GLUTEN FREE BAKING MIX – Special Authority see SA1729 above – Hospital Powder | pharmacy [HP3] 1,000 g OP | |
|--|------------------------------|----------------------------------|
| (5.15) | | Healtheries Simple Baking Mix |
| GLUTEN FREE BREAD MIX - Special Authority see SA1729 above - Hospital | pharmacy [HP3] | |
| Powder | 1,000 g OP | |
| (7.32) | - | NZB Low Gluten Bread Mix |
| 3.51 | | |
| (10.87) | | Horleys Bread Mix |
| GLUTEN FREE FLOUR - Special Authority see SA1729 above - Hospital phan | macy [HP3] | |
| Powder | 2,000 g OP | |
| (18.10) | - | Horleys Flour |

| | Subsidy | | Fully | Brand or |
|---|---------------------|---------------|----------|--------------|
| | (Manufacturer's Pri | | osidised | Generic |
| | \$ | Per | | Manufacturer |
| GLUTEN FREE PASTA - Special Authority see SA1729 on the | previous page - H | lospital phar | macy [H | P3] |
| Buckwheat Spirals | 2.00 | 250 g OP | | |
| | (3.11) | | C | Drgran |
| Corn and Vegetable Shells | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Corn and Vegetable Spirals | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Rice and Corn Lasagne Sheets | 1.60 | 200 g OP | | |
| | (3.82) | | C | Drgran |
| Rice and Corn Macaroni | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Rice and Corn Penne | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Rice and Maize Pasta Spirals | 2.00 | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Rice and Millet Spirals | 2.00 | 250 g OP | | |
| | (3.11) | | C | Drgran |
| Rice and corn spaghetti noodles | | 375 g OP | | |
| | (2.92) | | C | Drgran |
| Vegetable and Rice Spirals | | 250 g OP | | |
| | (2.92) | | C | Drgran |
| Italian long style spaghetti | | 220 g OP | | |
| | (3.11) | | C | Drgran |

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

| AMINOACID FORMULA WITHOUT METHIONINE - Special | Authority see SA110 | 8 above – Hos | pital pharmacy [HP3] |
|--|---------------------|---------------|----------------------------------|
| Powder | | 500 g OP | XMET Maxamum |

Supplements For MSUD

| Powder 437.22 | 500 a OP | MSUD Maxamum |
|--|--------------------|---------------------------|
| pharmacy [HP3] | | |
| AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Spi | ecial Authority se | e SA1108 above – Hospital |

| | Subsidy (Manufacturer's Price \$ | e) Per | Fully Subsidised | |
|--|--|-----------|---------------------|--------------------------------|
| Supplements For PKU | | | | |
| MINOACID FORMULA WITHOUT PHENYLALANINE – Spec Iarmacy [HP3] | | | | |
| Tabs | | 75 OF | | Phlexy 10 |
| Powder (orange) 36 g sachet | | 30 | 1 | PKU Anamix Junior Orange |
| Powder (berry) 28 g sachets | 936.00 | 30 | 1 | PKU Lophlex Powder |
| Powder (chocolate) 36 g sachet | | 30 | 1 | PKU Anamix Junior Chocolate |
| Powder (orange) 28 g sachets | 936.00 | 30 | ~ | PKU Lophlex Powder |
| Powder (unflavoured) 28 g sachets | 936.00 | 30 | 1 | PKU Lophlex Powder |
| Powder (unflavoured) 36 g sachets | | 30 | 1 | PKU Anamix Junior |
| Powder (vanilla) 36 g sachet | | 30 | 1 | PKU Anamix Junior Vanilla |
| Infant formula | | 400 g C | P 🗸 | PKU Anamix Infant |
| Powder (orange) | | 500 g C | | XP Maxamum |
| Powder (unflavoured) | | 500 g C | | XP Maxamum |
| Liquid (berry) | | 125 ml (| | PKU Anamix Junior |
| Liquid (orange) | 13.10 · | 125 ml (| OP 🗸 | PKU Anamix Junior LQ |
| Liquid (unflavoured) | 13.10 · | 125 ml (| OP 🗸 | PKU Anamix Junior LQ |
| Liquid (forest berries), 250 ml carton | | 18 OF | · · | Easiphen Liquid |
| Liquid (juicy tropical) 125 ml | | 30 OF | | PKU Lophlex LQ 20 |
| Oral semi-solid (berries) 109 g | | 36 OF | | PKU Lophlex Sensation 20 |
| Liquid (juicy berries) 62.5 ml | | 60 OF | · · | PKU Lophlex LQ 10 |
| Liquid (juicy citrus) 62.5 ml | | 60 OF | | PKU Lophlex LQ 10 |
| Liquid (juicy orange) 62.5 ml | | 60 OF | | PKU Lophlex LQ 10 |
| Liquid (juicy berries) 125 ml. | | 30 OF | | PKU Lophlex LQ 20 |
| Liquid (juicy orange) 125 ml | | 30 OF | | PKU Lophlex LQ 20 |

Foods

| LOW PROTEIN BAKING MIX – Special Authority see SA1108 Powder | | | oharmacy [HP3] |
|---|-------------------|----------------|------------------------------|
| LOW PROTEIN PASTA - Special Authority see SA1108 on th | e previous page - | Hospital pharm | acy [HP3] |
| Animal shapes | | 500 g OP | Loprofin |
| Lasagne | 5.95 | 250 g OP | Loprofin |
| Low protein rice pasta | 11.91 | 500 g OP | Loprofin |
| Macaroni | 5.95 | 250 g OP | Loprofin |
| Penne | 11.91 | 500 g OP | Loprofin |
| Spaghetti | 11.91 | 500 g OP | Loprofin |
| Spirals | 11.91 | 500 g OP | Loprofin |

| Subsidy | Fully | | Brand or |
|------------------------|------------|---|----------|
| (Manufacturer's Price) | Subsidised | | Generic |
| \$ | Per | 1 | |

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the

recommendation of a dietitian, relevant specialist, vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

| LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 |) above – Hos | spital pharmac | y [HP3] | |
|---|---------------|----------------|-----------------------------|--|
| Powder | 44.40 | 400 g OP | Locasol | |

Gastrointestinal and Other Malabsorptive Problems

| AMINO ACID FORMULA - Special Authority see SA2092 below - Hospital phar | macy [HP3] | |
|---|------------|--|
| Powder | 400 g OP | ✓ Alfamino ✓ Alfamino Junior |
| Powder (unflavoured)53.00 | 400 g OP | Anamino dunior Elecare Elecare LCP Neocate Gold Neocate Junior Unflavoured |
| Powder (vanilla)53.00 | 400 g OP | ✓ Neocate SYNEO ✓ Elecare ✓ Neocate Junior Vanilla |

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

| Subsidy | F | ully | Brand or |
|------------------------|--------|------|----------|
| (Manufacturer's Price) | Subsid | ised | Generic |
| \$ | Per | ✓ | |

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Fither:

- 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

1 Either:

1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | Su | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

| ENTERAL LIQUID PEPTIDE FORMULA - | Special Authority see SA1953 below - | Hospital phar | macy [HP3] |
|----------------------------------|--------------------------------------|---------------|----------------|
| Liquid 1 kcal/ml | 10.45 | 500 ml OP | Nutrini Pentis |

| | 0.45 500 mi OP | Nutrini Peptisorb |
|---------------------|----------------|---------------------------------------|
| Liquid 1.5 kcal/ml1 | 5.68 500 ml OP | Nutrini Peptisorb |
| | | Energy |

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or

2.10 Both:

2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

.

| Subsidy | | Fully | Brand or | |
|------------------------|-----|----------|--------------|--|
| (Manufacturer's Price) | Sub | osidised | Generic | |
| \$ | Per | 1 | Manufacturer | |

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

| EXTENSIVELY HYDROLYSED FORMULA - Special Authority see 5 | SA1557 belo | w – Hospital ph | armacy [HP3] |
|--|-------------|-----------------|--|
| Powder | 15.21 | 450 g OP | Aptamil Gold+ Pepti Junior |
| | 30.42 | 900 g OP | Aptamil AllerPro SYNEO 1 |
| | | | Aptamil AllerPro SYNEO 2 |

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

| Subsidy (Manufacturer's Price) | Subsid | Fully | Brand or Generic | |
|-----------------------------------|--------|-----------|---------------------|--|
| (Manulactuler's Flice) \$ | Per | iseu ✓ | Manufacturer | |

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML - Special Authority see SA1698 below - Hospital pharmacy [HP3]

| Liquid | 2.35 | 125 ml OP | 🗸 Infatrini |
|--------|------|-----------|-------------|
|--------|------|-----------|-------------|

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

| HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority s | see SA1197 a | above – Retail p | oharmacy |
|---|--------------|------------------|---------------------------------|
| Powder (unflavoured) | 35.50 | 300 g OP | KetoCal 4:1 |
| | | | Ketocal 3:1 |
| Powder (vanilla) | 35.50 | 300 g OP | KetoCal 4:1 |

| | | Subsidy | | Fully | Brand or |
|--|---|---------------------------|------------|-----------|----------------------------|
| | | (Manufacturer's Price) | Sub | sidised | Generic |
| | | \$ | Per | 1 | Manufacturer |
| accinations | | | | | |
| | GUERIN VACCINE – [Xpharm] sed risk of tuberculosis. Increased risk | is defined as: | | | |
| | | | | | |
| , 0 | se or family with a person with current or | | | | |
| , 0 | more household members or carers who | o within the last 5 years | s lived in | a count | ry with a rate of TB > or |
| | r 100,000 for 6 months or longer; or | | | | |
| during their first | st 5 years will be living 3 months or long | er in a country with a ra | ate of TB | > or eq | ual to 40 per 100,000 |
| Note a list of countri | ies with high rates of TB are available at | www.health.govt.nz/tu | iberculos | is (seard | ch for downloads) or |
| www.bcgatlas.org/ir | idex.php. | | | | |
| Ini Mycobacterium b | ovis BCG (Bacillus Calmette-Guerin), | | | | |
| | 331, live attenuated, vial with diluent | 0.00 | 10 | 🗸 В | CG Vaccine |
| PHTHERIA, TETANU Funded for any of th | S AND PERTUSSIS VACCINE – [Xpha e following criteria: | rm] | | _ | |
| 1) A single dose | for pregnant women in the second or thi | rd trimester of each pr | eanancy. | or | |
| | for parents or primary caregivers of infai | | | | re Unit or Specialist Care |
| , 0 | nore than 3 days, who had not been exp | | | | • |
| | | | | | |
| 3) A course of up primary immur | to four doses is funded for children from nisation; or | n age / up to the age (| лоуеа | IS INCIUS | ive to complete full |

- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients: or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

| haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe0.00 | 10 | Boostrix |
|--|----|------------------------------|
| | 1 | Boostrix |

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

Vaccinatio

BACILLUS CA For infants 1) living 2) havir equa 3) durin Note a list www.bcga

Danisl DIPHTHERIA.

- A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation: or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

| pertussis toxoid, 25 mcg pertussis filamentous | |
|--|--|
| haemagglutinin, 8 mcg pertactin and 80 D-antigen units | |

| poliomyelitis virus in 0.5ml syringe | 0.00 | 10 | Infanrix IPV |
|--------------------------------------|------|----|--------------|
| | 0.00 | 10 | |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|------------|--------------|--|
| (Manufacturer's Price) | 5 | Subsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]

Funded for patients meeting any of the following criteria:

- 1) Up to four doses for children up to and under the age of 10 for primary immunisation; or
- 2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 3) Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.

Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe0.00 10 ✓ Infanrix-hexa

HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]

One dose for patients meeting any of the following:

- 1) For primary vaccination in children; or
- 2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, preor post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

| Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml0.00 | 1 | ✓ Hiberix |
|---|---|-----------------------------------|
| HEPATITIS A VACCINE – [Xpharm] Funded for patients meeting any of the following criteria: | | |
| Two vaccinations for use in transplant patients; or | | |
| 2) Two vaccinations for use in children with chronic liver disease; or | | |
| 3) One dose of vaccine for close contacts of known hepatitis A cases. | | |
| Inj 1440 ELISA units in 1 ml syringe0.00 | 1 | ✓ <u>Havrix</u> |
| Inj 720 ELISA units in 0.5 ml syringe0.00 | 1 | Havrix Junior |

| | Subsidy (Manufacturer's Price) | | Fully Subsidised | Brand or Generic |
|--|-----------------------------------|--------|---------------------|---------------------|
| | (Manulaciuler's Flice) \$ | Per | | Manufacturer |
| EPATITIS B RECOMBINANT VACCINE – [Xpharm] | | | | |
| Inj 10 mcg per 0.5 ml prefilled syringe | | 1 | ✓ E | ngerix-B |
| Funded for patients meeting any of the following criteri | | | | |
| 1) for household or sexual contacts of known acute | | | | s; or |
| 2) for children born to mothers who are hepatitis B | | | | |
| for children up to and under the age of 18 years in the second sec | | | | achieved a positive |
| serology and require additional vaccination or rec 4) for HIV positive patients; or | quire a primary course c | of vac | ccination; or | |
| 5) for hepatitis C positive patients; or | | | | |
| 6) for patients following non-consensual sexual inte | rcourse: or | | | |
| 7) for patients following immunosuppression; or | | | | |
| 8) for solid organ transplant patients; or | | | | |
| 9) for post-haematopoietic stem cell transplant (HS | CT) patients: or | | | |
| 10) following needle stick injury. | · , , . | | | |
| lei 00 mees neu 1 mi nuefilled eurigen | 0.00 | | | |
| Inj 20 mcg per 1 ml prefilled syringe Funded for patients meeting any of the following criteri | | 1 | • - | ngerix-B |
| | | | litic D corrier | |
| for household or sexual contacts of known acute for children born to mothers who are hepatitis B s | | | | s; or |
| 3) for children up to and under the age of 18 years i | | | | achieved a nositive |
| serology and require additional vaccination or rec | | | | |
| 4) for HIV positive patients; or | | | | |
| 5) for hepatitis C positive patients; or | | | | |
| 6) for patients following non-consensual sexual inte | rcourse; or | | | |
| 7) for patients following immunosuppression; or | | | | |
| for solid organ transplant patients; or | | | | |
| for post-haematopoietic stem cell transplant (HS) | CT) patients; or | | | |
| following needle stick injury; or | | | | |
| 11) for dialysis patients; or | | | | |
| 12) for liver or kidney transplant patients. | | | | |
| UMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND | 58) VACCINE [HPV] - | - [Xp | harml | |
| Any of the following: | | L | | |
| 1) Maximum of two doses for children aged 14 years an | d under; or | | | |
| 2) Maximum of three doses for patients meeting any of t | | | | |
| 1) People aged 15 to 26 years inclusive; or | C C | | | |
| 2) Either: | | | | |
| _/ | | | | |
| People aged 9 to 26 years inclusive | | | | |
| | | | | |
| People aged 9 to 26 years inclusive | or | | | |

| | Inj 270 mcg in 0.5 ml syringe0.00 | 10 |
|--|-----------------------------------|----|
|--|-----------------------------------|----|

✓ Gardasil 9

| FLUENZA VACCINE Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) [Xpharm] | | Subsidy (Manufacturer's Price) \$ | Per | Fully Subsidised | Generic |
|--|--|---|--------|---------------------|----------------------------|
| - [Xpharm] | FLUENZA VACCINE | | | | |
| A) INFLUENZA VACCINE – child aged 6 months to 35 months is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharm have any of the following cardiovascular diseases ischaemic heart disease, or congestive heart failure, or rheumatic heart disease, or congenital heart disease, or congenital heart disease, or congenital heart disease; or congenital heart disease; or have either of the following chronic respiratory diseases: asthma, if on a regular preventative therapy, or other chronic respiratory disease with impaired lung function; or have diabets; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or have any of the following other conditions: autoimmune disease, or immune suppression or immune deficiency, or HIV, or transplant recipients, or on long term aspirin, or have a cochlear implant, or errors of metabolism at risk of major metabolic decompensation, or pre and post splenectomy, or down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: asthma not requiring regular preventative therapy, hypertension and/or dyslipidaemia without evidence of end-organ disease. | | | 1 | ~ | |
| i) have any of the following cardiovascular diseases a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) have a cochlear implant, or h) have a cochlear implant, or h) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. | | | | | . , |
| a) ischaemic heart disease, or b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) have a cochlear implant, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. | is available each year for patients aged 6 months to | o 35 months who mee | et the | e following | criteria, as set by Pharma |
| b) congestive heart failure, or c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic respiratory disease with impaired lung function; or iv) have diabetes; or v) have diabetes; or v) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or v) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have acoblear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 meg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | have any of the following cardiovascular disea | ases | | | |
| c) rheumatic heart disease, or d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | a) ischaemic heart disease, or | | | | |
| d) congenital heart disease, or e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) astma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) have a cochlear implant, or i) have a cochlear implant, or i) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. | , 5 | | | | |
| e) cerebo-vascular disease; or ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. | , | | | | |
| ii) have either of the following chronic respiratory diseases: a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , 0 | | | | |
| a) asthma, if on a regular preventative therapy, or b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , | | | | |
| b) other chronic respiratory disease with impaired lung function; or iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , | • | | | |
| iii) have diabetes; or iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , | 1.2.1 | | | |
| iv) have chronic renal disease; or v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , , , , | npaired lung function; | or | | |
| v) have any cancer, excluding basal and squamous skin cancers if not invasive; or vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | | | |
| vi) have any of the following other conditions: a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | oue ekin concore if n | at inv | acivo: or | |
| a) autoimmune disease, or b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | ous skill calicels if he | JUIIIV | asive, 01 | |
| b) immune suppression or immune deficiency, or c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | | | |
| c) HIV, or d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | ncv or | | | |
| d) transplant recipients, or e) neuromuscular and CNS diseases/disorders, or f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | loy, ol | | | |
| f) haemoglobinopathies, or g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , , , , , , , , , , , , , , , , , , , | | | | |
| g) on long term aspirin, or h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | e) neuromuscular and CNS diseases/disor | ders, or | | | |
| h) have a cochlear implant, or i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | f) haemoglobinopathies, or | | | | |
| i) errors of metabolism at risk of major metabolic decompensation, or j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | | | |
| j) pre and post splenectomy, or k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | | | |
| k) down syndrome, or vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | , | tabolic decompensat | ion, c | or | |
| vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness; Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | | | |
| Unless meeting the criteria set out above, the following conditions are excluded from funding: a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | ar have a history of a | :~~:fi | aant raanir | aton (illnaad) |
| a) asthma not requiring regular preventative therapy, b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | - | | |
| b) hypertension and/or dyslipidaemia without evidence of end-organ disease. B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | 0 | 0 | xciuu | | nung. |
| B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine in 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | | dicoa | 20 | |
| 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutic | | - | | | ly of influenze veccine in |
| | 30 mcg in 0.25 ml syringe (paediatric quadrivalent subsidised immunisation and they may only do so i | vaccine) to patients e | ligibl | e under the | above criteria for |
| | | | | | |
| | | | | | |

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)......110.00

10

 Afluria Quad (2022 formulation)

| Subsidy | | Fully | |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ~ | Manufacturer |

a) Only on a prescription

- b) No patient co-payment payable
- C)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Maori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

| Subsidy | Fu | Illy Brand or | |
|------------------------|----------|----------------------------------|--|
| (Manufacturer's Price) | Subsidis | ed Generic | |
| \$ | Per | Manufacturer | |

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- C)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml.

| diluent 0.5 ml 112.50 | 5 | 🗸 MMR II |
|-----------------------|----|-----------|
| 250.00 | 10 | ✓ Priorix |

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. Inj 4 mcg of each meningococcal polysaccharide conjugated to

| a total of approximately 48 mcg of diphtheria toxoid c | arrier | | |
|--|--------|---|------------------------------|
| per 0.5 ml vial | 0.00 | 1 | Menactra |

| Subsidy | | Fully | Brand or | |
|------------------------|-----|-----------|--------------|--|
| (Manufacturer's Price) | S | ubsidised | Generic | |
| \$ | Per | ✓ | Manufacturer | |

MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm]

Either:

- A) Both:
 - 1) Child is under one year of age; and
 - 2) Any of the following:
 - i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to three doses for close contacts of meningococcal cases of any group; or
 - iii) up to three doses for child who has previously had meningococcal disease of any group; or
 - iv) up to three doses for bone marrow transplant patients; or
 - v) up to three doses for child pre- and post-immunosuppression*; or
- B) Both:
 - 1) Person is one year of age or over; and
 - 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

| Inj 175 mcg per 0.5 ml prefilled syringe0.00 | 1 | Bexsero | |
|--|---|-----------------------------|--|
| IENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm] | | | |
| Both: | | | |
| 1) The child is under 9 months of age; and | | | |
| 2) Any of the following: | | | |

- 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
- 2) Two doses for close contacts of meningococcal cases of any group; or
- 3) Two doses for child who has previously had meningococcal disease of any group; or
- 4) A maximum of two doses for bone marrow transplant patients; or
- 5) A maximum of two doses for child pre- and post-immunosuppression*.

Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe......0.00 1 Veisvac-C

PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm]

1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B,

| 7F, 9V, 14 and 23F; 3 mcg of pneumococcal | | |
|---|----|-------------------------------|
| polysaccharide serotypes 4, 18C and 19F in 0.5 ml | | |
| prefilled syringe0.00 | 10 | Synflorix |

| Subsidy | | Fully | Brand or |
|------------------------|-----|------------|--------------|
| (Manufacturer's Price) | | Subsidised | Generic |
| \$ | Per | ✓ | Manufacturer |

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously
 received two doses of the primary course of PCV10; or
- Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

| Ini 30.8 mcg of pneumococcal | polysaccharide serotypes 1, 3, 4, |
|-------------------------------|-----------------------------------|
| ing oolo mog of priodinoooodd | |

| 5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml | | |
|---|----|-------------|
| syringe0.00 | 10 | Prevenar 13 |
| | 1 | Prevenar 13 |

| | Subsidy (Manufacturer's Price) \$ | Subsid Per | Fully lised | Brand or Generic Manufacturer |
|---|---|---------------|----------------|-------------------------------------|
| PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – [X Either: | (pharm] | | | |
| Up to three doses (as appropriate) for patients with HIV. | for natients nost had | ematonoieti | c stem | cell transplant or |
| chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochlear All of the following: | nal asplenia, pre- or p | post-solid o | rgan tr | ansplant, renal dialysis, |
| a) Patient is a child under 18 years for (re-)immunisa b) Treatment is for a maximum of two doses; and c) Any of the following: | tion; and | | | |
| any of the following. an immunosuppressive therapy or radiation t | herany vaccinate wł | non thoro is | avnar | ted to be a sufficient |
| immune response; or | nerapy, vaccinate wi | | expec | |
| ii) with primary immune deficiencies; or | | | | |
| iii) with HIV infection; or | | | | |
| iv) with renal failure, or nephrotic syndrome; orv) who are immune-suppressed following organ | transplantation (incl | udina haan | natono | iatic stam call transplant). |
| or | | ualing nach | latopo | |
| vi) with cochlear implants or intracranial shunts; | or | | | |
| vii) with cerebrospinal fluid leaks; or | | | | - Level als the also see as f |
| viii) receiving corticosteroid therapy for more that prednisone of 2 mg/kg per day or greater, or | | | | |
| 20 mg or greater; or | children who weight | | lo ky o | Tha total daily dosage of |
| ix) with chronic pulmonary disease (including as | thma treated with high | gh-dose cor | ticoste | roid therapy); or |
| x) pre term infants, born before 28 weeks gesta | | | | |
| xi) with cardiac disease, with cyanosis or failurexii) with diabetes; or | ; or | | | |
| xii) with Down syndrome; or | | | | |
| xiv) who are pre-or post-splenectomy, or with fur | ctional asplenia. | | | |
| hi 575 maa in 0.5 ml andillad aviana (05 maa af aach | | | | |
| Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype) | 0.00 | 1 | 🖌 Pi | neumovax 23 |
| POLIOMYELITIS VACCINE – [Xpharm] | | | • 😐 | |
| Up to three doses for patients meeting either of the following: | | | | |
| 1) For partially vaccinated or previously unvaccinated indiv | riduals; or | | | |
| 2) For revaccination following immunosuppression. | | | | |
| Note: Please refer to the Immunisation Handbook for approp | | | | |
| Inj 80D antigen units in 0.5 ml syringe | 0.00 | 1 | ✓ IP | |
| ROTAVIRUS ORAL VACCINE – [Xpharm] | | | | |
| Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 14 w | eeks of age: and | | | |
| no vaccination being administered to children aged 24 v | | | | |
| Oral susp live attenuated human rotavirus | | | | |
| 1,000,000 CCID50 per dose, prefilled oral applicator | 0.00 | 10 | ✓ <u>R</u> | otarix |

| Subsidy | Fully | Brand or |
|------------------------|------------|--------------|
| (Manufacturer's Price) | Subsidised | Generic |
| \$ | Per 🖌 | Manufacturer |

VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]

Either:

- 1) Maximum of one dose for primary vaccination for either:
 - a) Any infant born on or after 1 April 2016; or
 - b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or
- 2) Maximum of two doses for any of the following:
 - a) Any of the following for non-immune patients:
 - i) with chronic liver disease who may in future be candidates for transplantation; or
 - ii) with deteriorating renal function before transplantation; or
 - iii) prior to solid organ transplant; or
 - iv) prior to any elective immunosuppression*, or
 - v) for post exposure prophylaxis who are immune competent inpatients.; or
 - b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or
 - c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or
 - d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or
 - e) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or
 - f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or
 - g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.

* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days

| Inj 1350 PFU prefilled syringe | 0.00 | · . | <u>Varivax</u> | |
|--------------------------------|------|------|----------------|---|
| | 1 | 10 🗸 | Varivax | |
| | | | | - |

VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm] Funded for patients meeting the following criteria:

1) One dose for all people aged 65 years

| Inj 19,400 PFU prefilled syringe plus vial0.00 | 1 | Zostavax |
|--|----|------------------------------|
| | 10 | Zostavax |

Diagnostic Agents

| TUBERCULIN PPD [MANTOUX] TEST - [Xpharm] | | | |
|--|------|---|----------|
| Inj 5 TU per 0.1 ml, 1 ml vial | 0.00 | 1 | Tubersol |

į

- Symbols -

| UK Synacthen |
|---|
| ЗТС 106 - Δ - |
| A-Scabies |
| Abacavir sulphate |
| Abacavir sulphate with |
| lamivudine 106 |
| Abiraterone acetate |
| Additate offe acetate |
| Accarb |
| Accuretic 10 |
| Accuretic 20 |
| Acetazolamide |
| |
| Acetec47 Acetic acid with hydroxyquinoline and |
| ricinoleic acid |
| Acetylcysteine |
| Acetylcystelle |
| Aciclovir |
| Infection 101 |
| Sensory |
| Acidex |
| Acipimox |
| Acipiniox |
| Aclasta |
| Actemra |
| Actemra S29216 |
| Actinomycin D |
| Actrapid |
| Actrapid Penfill10 |
| Acupan |
| Adalimumab (Amgevita) 176 |
| Adalimumab (Humira)185 |
| Adapalene |
| ADR Cartridge 1.8 |
| Adrenaline |
| Advantan64 |
| Advate |
| Adynovate41 |
| Afinitor |
| Aflibercept189 |
| Afluria Quad |
| (2022 formulation) 271 |
| Afluria Quad Junior |
| (2022 formulation) 271 |
| AFT-Pyrazinamide |
| Agents Affecting the |
| Renin-Angiotensin System 47 |
| Agents for Parkinsonism and Related |
| Disorders 119 |
| Agents Used in the Treatment of |
| Poisonings242 |
| Agrylin151 |

| Albendazole | |
|-------------------------------------|----------|
| Albey | |
| Albustix | .77 |
| Alchemy Oxybutynin | .76 |
| Aldurazyme | .29 |
| Alecensa | 158 |
| Alectinib | 158 |
| Alendronate sodium | 112 |
| Alendronate sodium with | |
| colecalciferol | 112 |
| Alfacalcidol | .33 |
| Alfamino | |
| Alfamino Junior | |
| Alginic acid | |
| Alglucosidase alfa | |
| Alkeran | |
| Alkeran S29 | 140 |
| Allersoothe | |
| | |
| Allmercap | |
| Allopurinol | 117 |
| Alpha-Adrenoceptor Blockers | .4/ |
| Alpha-Keri Lotion | . 66 |
| Alphamox | . 92 |
| Alphamox 125 | . 92 |
| Alphamox 250 | . 92 |
| Alprolix | . 38 |
| Alu-Tab | 6 |
| Aluminium hydroxide | 6 |
| Alvogen Amantadine hydrochloride | . 50 |
| Amantadine hydrochloride | 119 |
| Ambrisentan | |
| Ambrisentan Mylan | . 58 |
| Amgevita | 176 |
| Amiloride hydrochloride | .53 |
| Amiloride hydrochloride with | |
| furosemide | . 53 |
| Amiloride hydrochloride with | |
| hydrochlorothiazide | . 54 |
| Aminophylline | |
| Amiodarone hydrochloride | .49 |
| Amisulpride | |
| Amisulpride Mylan | 132 |
| Amitriptyline | 125 |
| Amlodipine | |
| Amneal | |
| Amorolfine | |
| Amoxicillin | |
| Amoxicillin with clavulanic acid | 20 00 |
| Amphotericin B | |
| | |
| Amsacrine | |
| AmsaLyo | |
| Amsidine | |
| Amzoate | |
| Anaesthetics | 120 |

| Anagrelide hydrochloride 1 | 51 |
|-------------------------------------|-----|
| Analgesics1 | |
| Anastrozole1 | 69 |
| Anatrole1 | |
| Andriol Testocaps | |
| Androderm | 80 |
| Anoro Ellipta2 | |
| Antabuse 1 | |
| Antacids and Antiflatulents | 6 |
| Anthelmintics | 89 |
| Antiacne Preparations | 61 |
| Antiallergy Preparations | |
| Antianaemics | 37 |
| Antiandrogen Oral | · · |
| Contraceptives | 75 |
| Antiarrhythmics | 10 |
| Antibacterials | |
| Antibacterials Topical | 61 |
| Anticholinergic Agents | 21 |
| Anticholinesterases1 | |
| Antidepressants1 | |
| Antidiarrhoeals | |
| Antiepilepsy Drugs1 | |
| Antifibrinolytics, Haemostatics and | 21 |
| Local Sclerosants | 20 |
| Local Scierosarius | 30 |
| Antifibrotics | |
| Antifungals | |
| Antifungals Topical | |
| Antihistamines | |
| Antihypotensives | |
| Antimalarials | 98 |
| Antimigraine Preparations1 | 30 |
| Antinausea and Vertigo Agents 1 | 30 |
| Antipruritic Preparations | |
| Antipsychotics1 | 32 |
| Antiretrovirals1 | 04 |
| Antirheumatoid Agents 1 | 12 |
| Antispasmodics and Other Agents | |
| Altering Gut Motility | . 8 |
| Antithrombotic Agents | 41 |
| Antithymocyte globulin | |
| (equine) 1 | |
| Antitrichomonal Agents | 99 |
| Antituberculotics and | |
| Antileprotics | |
| Antiulcerants | |
| Antivirals 1 | |
| Anxiolytics1 | 35 |
| Anzatax1 | 55 |
| Apidra | |
| Apidra SoloStar | 11 |
| APO-Atomoxetine 1 | 38 |
| Apo-Azithromycin | 90 |
| Apo-Ciclopirox | 62 |

| Apo-Diclo SR 111 |
|-------------------------------|
| Apo-Diltiazem CD |
| Apo-Doxazosin47 |
| Apo-Folic Acid |
| Apo-Megestrol |
| Apo-Oxybutynin |
| Apo-Pindolol |
| |
| Apo-Prazosin |
| Apo-Prednisone |
| Apo-Primidone129 |
| Apo-Pyridoxine |
| Apo-Temozolomide156 |
| Apomorphine hydrochloride119 |
| Aprepitant 130 |
| Apresoline57 |
| Aptamil AllerPro SYNEO 1 |
| Aptamil AllerPro SYNEO 2 |
| Aptamil Gold+ Pepti Junior266 |
| Aqueous cream |
| Aratac |
| Arava |
| Arginine |
| Aripiprazole |
| Aripiprazole Sandoz |
| |
| Aristocort |
| Arrotex-Prazosin S29 |
| Arrow - Lattim |
| Arrow-Amitriptyline |
| Arrow-Bendrofluazide |
| Arrow-Brimonidine |
| Arrow-Diazepam |
| Arrow-Doxorubicin 153 |
| Arrow-Losartan & |
| Hydrochlorothiazide 48 |
| Arrow-Norfloxacin 110 |
| Arrow-Ornidazole99 |
| Arrow-Quinapril 1048 |
| Arrow-Quinapril 2048 |
| Arrow-Quinapril 548 |
| Arrow-Roxithromycin91 |
| Arrow-Timolol |
| Arrow-Topiramate129 |
| Arrow-Tramadol125 |
| Arsenic trioxide151 |
| Asacol7 |
| Ascorbic acid |
| Aspen Adrenaline |
| Aspirin |
| Blood41 |
| Nervous121 |
| Asthalin |
| Atazanavir sulphate |
| Atenolol |
| Atenolol AFT50 |
| Atenolol AFT S29 |
| ATGAM |
| A I GAWI |

| Ativan135 |
|--------------------------------|
| Atomoxetine138 |
| Atorvastatin54 |
| Atropine sulphate |
| Cardiovascular |
| Sensory |
| Atropt |
| Atrovent |
| Autovent |
| |
| Aubagio |
| Augmentin |
| Aurorix |
| AutoSoft 3023 |
| AutoSoft 9023 |
| Avelox |
| Avonex |
| Avonex Pen 136 |
| Azacitidine148 |
| Azacitidine Dr Reddy's148 |
| Azamun 169 |
| Azathioprine169 |
| Azilect |
| Azithromycin90 |
| Azopt |
| AZT106 |
| -B- |
| B-D Micro-Fine |
| B-D Ultra Fine |
| B-D Ultra Fine II |
| Bacillus Calmette-Guerin (BCG) |
| vaccine |
| Bacillus Calmette-Guerin |
| |
| vaccine |
| Baclofen |
| Bactroban |
| Balance |
| Barrier Creams and Emollients |
| BCG Vaccine |
| Beclazone 100229 |
| Beclazone 250229 |
| Beclazone 50229 |
| Beclomethasone dipropionate |
| |
| Bee venom allergy treatment |

| Beta Ointment63 |
|-------------------------------------|
| Beta Scalp70 |
| Beta-Adrenoceptor Agonists231 |
| Beta-Adrenoceptor Blockers50 |
| Betadine66 |
| Betadine Skin Prep66 |
| Betaferon 137 |
| Betahistine dihydrochloride131 |
| Betaine27 |
| Betaloc CR50 |
| Betamethasone dipropionate63 |
| Betamethasone dipropionate with |
| calcipotriol68 |
| Betamethasone sodium phosphate |
| with betamethasone acetate 79 |
| Betamethasone valerate63, 70 |
| Betamethasone valerate with sodium |
| fusidate [fusidic acid] 64 |
| Betaxolol239 |
| Betnovate63 |
| Betoptic239 |
| Betoptic S 239 |
| Bexsero274 |
| Bezafibrate54 |
| Bezalip54 |
| Bezalip Retard54 |
| Bicalutamide 166 |
| Bicillin LA92 |
| BiCNU147 |
| Bicnu Heritage 147 |
| Bile and Liver Therapy9 |
| Biltricide |
| Bimatoprost239 |
| Bimatoprost Multichem 239 |
| Binarex 166 |
| Binocrit |
| Biodone 123 |
| Biodone Extra Forte123 |
| Biodone Forte 123 |
| Bisacodyl |
| Bisoprolol fumarate |
| Bisoprolol Mylan |
| BK Lotion |
| Bleomycin sulphate 152 |
| Blood Colony-stimulating |
| Factors |
| Blood glucose diagnostic test |
| meter |
| Blood glucose diagnostic test |
| strip |
| Blood glucose test strips (visually |
| impaired)15 |
| Blood Ketone Diagnostic Test |
| Strip |
| Bonjela |
| Boostrix |

| Bortozomih 1 | 50 |
|---|-----|
| | 52 |
| Bortezomib1 Bortezomib Dr Reddy's S291 | 52 |
| Bortezomib Dr-Reddy's1 | 52 |
| Bortezomib Juno1 | |
| Bortezomib Juno S291 | 50 |
| | |
| Bosentan | 58 |
| Bosentan Dr Reddy's | 58 |
| Bplex | 33 |
| Breo Ellipta2 | |
| | |
| Brevinor 1/28 | 74 |
| Bricanyl Turbuhaler2 | |
| Brilinta | 42 |
| Brimonidine tartrate2 | 40 |
| Brimonidine tartrate with timolol | 10 |
| | |
| maleate2 | 40 |
| Brinzolamide2 | 39 |
| Brolene2 | 37 |
| Bromocriptine mesylate1 | |
| | 13 |
| Brufen SR 1 | 11 |
| BSF Clomipramine Teva2 | |
| BSF Comtan2 | 42 |
| BSF Lopinavir/Ritonavir Mylan2 | 42 |
| BSF Vigisom | |
| | 42 |
| Buccastem1 | 31 |
| Budesonide | |
| Alimentary | .6 |
| Respiratory229, 2 | 35 |
| Budesonide with eformoterol2 | 30 |
| Bumetanide | |
| | |
| Buprenorphine Naloxone BNM 1 | |
| Buprenorphine with naloxone1 | |
| Bupropion hydrochloride1 | 43 |
| Burinex | |
| Burinex S29 | |
| | |
| Buscopan | .8 |
| Buspirone hydrochloride1 | 35 |
| Buspirone Viatris1 | 35 |
| Busulfan1 | 47 |
| - C - | |
| Cabergoline | 00 |
| | |
| Cacit | 34 |
| Caffeine citrate2 | 36 |
| Calamine | 63 |
| Calamine-AFT | 63 |
| Calci-Tab 500 | 21 |
| | |
| Calcipotriol | 68 |
| Calcitonin | 78 |
| Calcitriol | |
| Calcitriol-AFT | 33 |
| Calcium carbonate6, | |
| Calcium Channel Blockers | C 1 |
| | |
| Calcium Disodium Versenate2 | |
| Calcium folinate 1 | 49 |
| Calcium Folinate Ebewe1 | 49 |
| Calcium Folinate Sandoz1 | 49 |
| Calcium Folinate Sandoz S29 1 | |
| | |

| Calcium gluconate | 34 |
|--|--|
| Calcium Homeostasis7 | 78 |
| Calcium lactate gluconate with | |
| calcium carbonate | 35 |
| Calcium polystyrene sulphonate | 16 |
| Calcium Resonium4 | 16 |
| Calcium-Sandoz Forte | 25 |
| Calogen | |
| Candesartan cilexetil | 19 |
| Candestar | |
| Canesten | |
| Capecitabine14 | |
| Capercit | +9 10 |
| Capercil | +9 17 |
| Capoten | +/ |
| Capsaicin Musculoskeletal11 | |
| | |
| Nervous12 | |
| Captopril | |
| Carafate | |
| Carbaccord 14 | |
| Carbamazepine 12 | 27 |
| Carbimazole | |
| Carbomer24 | |
| Carboplatin14 | 17 |
| Carboplatin Ebewe 14 | |
| Carbosorb-X | 12 |
| Cardinol LA | |
| Cardizem CD | 52 |
| | |
| CareSens Dual 1 | |
| CareSens N14-1 | 15 |
| CareSens N | 5 4 |
| CareSens N14-1 | 5 4 |
| CareSens N14-1 CareSens N POP14-1 CareSens N Premier | 5 4 4 |
| CareSens N | 5 4 4 |
| CareSens N | 15 14 14 15 |
| CareSens N | 15 14 15 32 |
| CareSens N | 15 14 15 32 |
| CareSens N | 15 14 15 32 17 50 |
| CareSens N | 15 14 15 32 17 50 50 |
| CareSens N | 15 14 15 32 17 50 50 |
| CareSens N | 15 14 15 32 17 50 50 50 50 |
| CareSens N | 15 14 15 32 17 50 50 50 52 17 |
| CareSens N | 15 14 15 32 750 50 50 50 50 52 17 39 |
| CareSens N | 15 14 15 32 7 50 50 52 17 39 39 |
| CareSens N | 15 14 15 37 50 50 52 17 39 39 39 |
| CareSens N | 15 14 15 32 750 50 52 739 39 39 39 |
| CareSens N | 15 14 15 27 50 05 27 39 39 39 39 39 39 |
| CareSens N | 15 14 15 327 50 50 52 7 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 15 27 50 90 52 7 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 15 27 50 05 27 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 15 27 50 02 27 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 15 27 50 00 27 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 14 15 32 17 50 50 50 50 50 50 50 50 50 50 50 50 50 |
| CareSens N | 15 14 14 15 32 17 50 00 52 27 17 39 90 90 90 90 90 90 90 90 90 90 90 90 90 |
| CareSens N | 15 14 15 32 17 50 00 22 17 75 00 00 52 17 75 39 39 39 39 39 39 39 39 39 39 39 39 39 |
| CareSens N | 15 14 14 15 32 17 50 00 50 50 00 55 21 77 93 93 93 93 93 93 93 93 93 93 93 93 93 |

| A | |
|-------------------------------------|------------|
| Cetomacrogol | 65 |
| Cetomacrogol with glycerol | 65 |
| Cetomacrogol-AFT | 65 |
| Cetuximab | 190 |
| Charcoal | 242 |
| Chemotherapeutic Agents | 146 |
| | |
| Chickenpox vaccine | |
| Chlorafast | 237 |
| Chlorambucil | |
| Chloramphenicol | |
| Chlorothiazide | 54 |
| Chlorpheniramine maleate | 229 |
| Chlorpromazine hydrochloride | . 132 |
| Chlortalidone [Chlorthalidone] | 54 |
| Chlorthalidone | |
| | |
| Chlorvescent | 40 |
| Choice Load 375 | 73 |
| Choice TT380 Short | |
| Choice TT380 Standard | 73 |
| Choline salicylate with cetalkonium | |
| chloride | 32 |
| Ciclopirox olamine | 62 |
| Ciclosporin | 224 |
| Cilazapril | 224 |
| | |
| Cilicaine | |
| Cilicaine VK | |
| Cinacalcet | |
| Cipflox | 94 |
| Ciprofloxacin | |
| Infection | 94 |
| Sensory | |
| Ciprofloxacin Teva | 237 |
| Cisplatin | |
| Cisplatin Ebewe | |
| | |
| Citalopram hydrobromide | 120 |
| Cladribine | 149 |
| Clarithromycin | |
| Alimentary | 9 |
| Infection | 90 |
| Clexane | 43 |
| Clexane Forte | 43 |
| Clindamycin | |
| Clinicians | |
| Clinicians Renal Vit | <u>2</u> 7 |
| | |
| Clobazam | . 127 |
| Clobetasol propionate6 | |
| Clobetasone butyrate | |
| Clofazimine | 99 |
| Clomazol | |
| Dermatological | 62 |
| Genito-Urinary | |
| Clomifene citrate | |
| Clomipramine hydrochloride | 105 |
| | . 120 |
| Clomipramine Teva | . 125 |
| Clonazepam 127 | |
| Clonidine | 52 |

| Clonidine BNM |
|------------------------------------|
| Clonidine hydrochloride52 |
| Clonidine Teva52 |
| Clopidogrel42 |
| Clopidogrel Multichem 42 |
| Clopine |
| Clopixol |
| Clotrimazole |
| Dermatological 62 |
| Genito-Urinary75 |
| Clozapine |
| Clozaril |
| Co-trimoxazole96 |
| Coal tar |
| Coal tar with allantoin, menthol, |
| phenol and sulphur 69 |
| Coal tar with salicylic acid and |
| sulphur |
| Coco-Scalp |
| Codeine phosphate |
| Extemporaneous245 |
| Nervous |
| Coenzyme Q10 |
| Colchicine |
| Colecalciferol |
| Colestid |
| |
| Colestipol hydrochloride |
| Colgout |
| Colifoam |
| Colistin sulphomethate |
| Colistin-Link |
| Collodion flexible |
| Colloidal bismuth subcitrate9 |
| Colofac8 |
| Coloxyl25 |
| Combigan |
| Compound electrolytes46 |
| Compound electrolytes with glucose |
| [Dextrose]46 |
| Compound hydroxybenzoate245 |
| Comtan 119 |
| Concerta 140 |
| Condoms |
| Condyline70 |
| Contraceptives - Hormonal73 |
| Contraceptives - Non-hormonal71 |
| Copaxone 136 |
| Corticosteroids and Related Agents |
| for Systemic Use79 |
| Corticosteroids Topical63 |
| Cortifoam7 |
| Cosentyx213 |
| |
| Cosmegen |
| Cosmegen |
| Cosmegen 152 |

| Creon 10000 | 24 |
|--|--|
| Creon 25000 | 24 |
| Creon Micro | |
| Crotamiton | |
| Crystaderm | |
| Curam | |
| Curam Duo 500/125 | 92 |
| Cvite | 33 |
| Cyclizine hydrochloride | |
| Cyclizine lactate | |
| Cyclogyl | 240 |
| Cyclonex | |
| Cyclopentolate hydrochloride | 240 |
| Cyclophosphamide | |
| Cyclorin | 00 |
| Cycloserine | oo |
| Cyproterone acetate | |
| Cyproterone acetate with | 00 |
| ethinyloestradiol | 75 |
| Cystadane | 75 |
| Cytarabine | 1/0 |
| Cytotec | . 149 0 |
| Cytoxan | 0 1 / 7 |
| - D - | 147 |
| D-Penamine | 110 |
| Dabigatran | |
| Dacarbazine | 150 |
| Dacarbazine APP | |
| Dataidazine AFF | 102 |
| Dactinomycin [Actinomycin D] | 150 |
| Dactinomycin [Actinomycin D] | |
| Daivobet | 68 |
| Daivobet Daivonex | 68 68 |
| Daivobet Daivonex Daktarin | 68 68 63 |
| Daivobet Daivonex Daktarin Dalacin C | 68 68 63 94 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium | 68 63 94 .118 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 | 68 63 94 .118 .118 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene | 68 63 94 . 118 . 118 . 118 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil | 68 63 94 . 118 . 118 . 118 11 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil Dapa-Tabs | 68 63 94 .118 .118 .118 11 54 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil Dapa-Tabs Dapsone | 68 63 94 .118 .118 .118 11 54 99 |
| Daivobet Daivonex Daktarin Dalacin C Dantrium Dantrium S29 Dantrolene Daonil Dapa-Tabs Dapasone Daraprim | 68 63 94 .118 .118 .118 11 54 99 95 |
| Daivobet | 68 63 94 .118 .118 .118 11 54 99 95 .106 |
| Daivobet | 68 63 94 . 118 . 118 . 118 . 118 . 118 11 54 99 95 .106 . 106 |
| Daivobet | 68 63 94 . 118 . 118 . 118 11 54 99 95 . 106 . 106 . 159 |
| Daivobet | 68 68 94 .118 .118 .118 .118 11 54 99 95 .106 .159 .152 |
| Daivobet | 68 63 94 .118 .118 .118 .118 11 54 99 95 .106 .159 .152 y |
| Daivobet | 68 68 94 .118 .118 .118 .118 11 .118 99 99 .106 .106 .159 .152 y 76 |
| Daivobet | 68 68 94 .118 .118 .118 11 54 99 95 .106 .106 .159 76 .242 |
| Daivobet | 68 68 94 .118 .118 .118 11 54 99 95 .152 y 76 .242 57 |
| Daivobet | 68 68 94 .118 .118 .118 .118 11 54 99 95 .106 .159 .152 y 76 .242 57 .234 |
| Daivobet | 68 63 94 .118 .118 .118 11 54 99 95 .106 .159 .152 y 76 .242 57 .234 .152 |
| Daivobet | 68 63 94 .118 .118 .118 11 54 95 .106 .159 .152 y 76 .242 57 .234 .152 .147 |
| Daivobet | 68 68 94 .118 .118 .118 11 54 99 95 .106 .106 .159 .152 y 76 .242 57 .234 .152 .147 .147 |
| Daivobet | 68 68 94 .118 .118 .118 11 54 99 95 .106 .106 .159 .152 y 76 .242 57 .234 .152 .147 .152 |
| Daivobet | 68 68 94 .118 .118 .118 .118 11 54 .106 .106 .159 .152 y 76 .242 57 .234 .152 .147 .152 .147 .152 |
| Daivobet | 68 68 94 .118 .118 .118 .118 11 54 99 99 .106 .106 .159 .152 y y 76 .242 57 .234 .152 .147 .152 .147 .152 .147 .152 |

| DBL Ergometrine | .75 |
|-----------------------------------|------|
| DBL Gemcitabine | 150 |
| DBL Gentamicin | . 94 |
| DBL Heparin Sodium | .44 |
| DBL Leucovorin Calcium | 149 |
| DBL Methotrexate Onco-Vial | 150 |
| DBL Morphine Sulphate | 123 |
| DBL Naloxone Hydrochloride | 242 |
| DBL Octreotide | 167 |
| DBL Pethidine Hydrochloride | 125 |
| DBL Vincristine Sulfate | 158 |
| Decozol | . 33 |
| Deferasirox | 242 |
| Deferiprone | 243 |
| Denosumab | 112 |
| Deolate | . 98 |
| Deoxycoformycin | 155 |
| Depo-Medrol | . 80 |
| Depo-Provera | .74 |
| Depo-Testosterone | . 80 |
| Deprim | . 96 |
| Dermol64 | , 70 |
| Desferrioxamine mesilate | 243 |
| Desmopressin | . 88 |
| Desmopressin acetate | |
| Desmopressin-PH&T | . 88 |
| Desuric | 117 |
| Detection of Substances in | |
| Urine | . 77 |
| Dexamethasone | |
| Hormone | |
| Sensory | 238 |
| Dexamethasone phosphate | . 79 |
| Dexamethasone Phosphate | |
| Panpharma | . 79 |
| Dexamethasone with framycetin and | |
| gramicidin | 237 |
| Dexamethasone with neomycin | |
| sulphate and polymyxin B | |
| sulphate | 238 |
| Dexamfetamine sulfate | 138 |
| Dexmethsone | .79 |
| Dextrochlorpheniramine | |
| maleate | 229 |
| Dextrose45 | -46 |
| DHC Continus | 122 |
| Diabetes | |
| Diabetes Management | |
| Diacomit | |
| Diagnostic Agents | |
| Diamide Relief | |
| Diamox | |
| Diasip | |
| Diason RTH | |
| Diazepam 127, | 135 |
| Diazoxide | |

| Dibenzyline |
|--|
| Diclofenac Sandoz111 |
| Diclofenac sodium |
| Musculoskeletal 111 |
| Sensory238 |
| Differin61 |
| Difflam |
| Diflucan96 |
| Digestives Including Enzymes24 |
| Digoxin49 |
| Dihydrocodeine tartrate 122 |
| Dilantin128 |
| Dilantin Infatab128 |
| Diltiazem hydrochloride52 |
| Dimethicone65–66 |
| Dimethyl fumarate 136 |
| Dipentum7 |
| Diphtheria, tetanus and pertussis |
| vaccine |
| Diphtheria, tetanus, pertussis and |
| polio vaccine268 |
| Diphtheria, tetanus, pertussis, polio, |
| hepatitis B and haemophilus |
| influenzae type B vaccine |
| Diprosone |
| Diprosone OV63 |
| Dipyridamole |
| Disopyramide phosphate |
| Disulfiram143 |
| Diuretics |
| Docetaxel152 |
| Docetaxel Accord 152 |
| Docetaxel Sandoz 152 |
| Docusate sodium |
| Docusate sodium with |
| sennosides |
| Dolutegravir |
| Domperidone |
| Donepezil hydrochloride141 |
| Donepezil-Rex141 |
| Dornase alfa |
| Dortimopt |
| Dorzolamide hydrochloride |
| Dorzolamide with timolol |
| Dostinex |
| Dosulepin [Dothiepin] |
| hydrochloride |
| Dosulepin Mylan |
| Dothiepin |
| Doxazosin |
| Doxazosin Clinect |
| Doxazosin Clinect |
| Doxorubicin Ebewe |
| Doxorubicin Ebewe |
| Doxorubicin nyurochionae |
| Doxycycline |
| DP Lotion |

| DP Lotn HC64 |
|---|
| DP-Allopurinol117 |
| Dr Reddy's Omeprazole9 |
| Drofate51 |
| Drugs Affecting Bone |
| Metabolism |
| Dual blood glucose and blood ketone |
| diagnostic test meter 14 |
| Dulaglutide12 |
| Dulcolax SP Drop |
| Duocal Super Soluble Powder248 |
| Duolin |
| Duolin HFA231 |
| DuoResp Spiromax230 |
| Duride |
| -E- |
| e-chamber La Grande235 |
| e-chamber Mask235 |
| e-chamber Turbo235 |
| E-Mycin |
| Ear Preparations |
| Ear/Eye Preparations |
| Easiphen Liquid |
| Econazole nitrate |
| Efavirenz |
| Efavirenz with emtricitabine and |
| tenofovir disoproxil |
| Eformoterol fumarate |
| |
| |
| Eformoterol fumarate dihydrate 230 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant |
| Eformoterol fumarate dihydrate230 Eftrenonacog alfa [Recombinant factor IX] |
| Eformoterol fumarate dihydrate |
| Eformoterol fumarate dihydrate230 Eftrenonacog alfa [Recombinant factor IX] |
| Eformoterol fumarate dihydrate |
| Eformoterol fumarate dihydrate230 Eftrenonacog alfa [Recombinant factor IX] |
| Eformoterol fumarate dihydrate |
| Eformoterol fumarate dihydrate |
| Eformoterol fumarate dihydrate |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon 64 Eltornbopag 38 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant 38 factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl. 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant 38 factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Eltrombopag 38 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant 38 factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Eltrombopag 38 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl. 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel. 69 Elocon 64 Elromobopag 38 Eltroxin 83 BMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 EMLA 121 Empagliflozin 13 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 EMLA 121 Empagliflozin 13 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elmental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 EMLA 121 Empagliflozin 13 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 EMLA 121 Empagliflozin 13 Empagliflozin with metformin hydrochloride hydrochloride 13 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elevos 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Eltorxin 83 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Empagliflozin 13 Empagliflozin with metformin hydrochloride hydrochloride 13 Emtricitabine 106 |
| Eformoterol fumarate dihydrate 230 Eftrenonacog alfa [Recombinant factor IX] factor IX] 38 Efudix 70 Egopsoryl TA 69 Elaprase 28 Eldepryl 119 Elecare 263 Electral 46 Elelyso 31 Elemental 028 Extra 253 Elidel 69 Elocon 64 Elocon Alcohol Free 64 Eltroxin 83 EMB Fatol 99 Emend Tri-Pack 130 Emicizumab 39 EMLA 121 Empagliflozin 13 Empagliflozin with metformin hydrochloride hydrochloride 13 |

| Emulsifying ointment | 65 |
|----------------------------------|-----|
| Emulsifying Ointment ADE | 60 |
| Endusiying Ontinent ADE | 00 |
| Enalapril maleate | 47 |
| Enbrel 1 | 69 |
| Endocrine Therapy 1 | 65 |
| Endoxan 1 | 47 |
| Engerix-B | 270 |
| Enlafax XR1 | 27 |
| Enoxaparin sodium | 19 |
| Enstilar | |
| | |
| Ensure | :57 |
| Ensure Plus | :58 |
| Ensure Plus HN | 257 |
| Ensure Plus RTH | 257 |
| Ensure Two Cal HN RTH | 259 |
| Entacapone1 | 19 |
| Entecavir1 | |
| Entecavir Sandoz1 | 00 |
| | |
| Entocort CIR | |
| Entresto 24/26 | |
| Entresto 49/51 | 48 |
| Entresto 97/103 | 48 |
| Epilim1 | 29 |
| Epilim Crushable 1 | 29 |
| Epilim IV1 | |
| Epilim S/F Liquid1 | 20 |
| Epilim Syrup 1 | 20 |
| Epiliili Sylup | 28 |
| Epirubicin Ebewe1 | 53 |
| Epirubicin hydrochloride1 | |
| Eplerenone | 53 |
| Epoetin alfa | |
| Epoprostenol | 60 |
| Eptacog alfa [Recombinant factor | |
| VIIa] | 40 |
| ERA | 01 |
| Erbitux 1 | 00 |
| | |
| Ergometrine maleate | |
| Erlotinib1 | |
| Erythrocin IV | 91 |
| Erythromycin (as lactobionate) | 91 |
| Erythromycin ethyl succinate | 91 |
| Erythromycin stearate | |
| Esbriet | |
| Escitalopram1 | 26 |
| Escitalopram (Ethics)1 | 26 |
| Eskazole | 20 |
| | 09 |
| Essential Generics | |
| Essential Prednisolone | |
| Estradiol TDP Mylan | 81 |
| Estradot | 81 |
| Estradot 50 mcg | |
| Estrofem | |
| Etanercept1 | |
| Ethambutol hydrochloride | |
| | |
| Ethics Aspirin1 | 21 |
| Ethics Aspirin EC | 41 |

| Ethics Lisinopril47 |
|--|
| Ethinyloestradiol |
| Ethinvloestradiol with |
| desogestrel73 |
| Ethinyloestradiol with |
| levonorgestrel |
| Ethinyloestradiol with |
| norethisterone |
| Ethosuximide |
| Etopophos |
| Etoposide |
| Etoposide phosphate |
| Etravirine |
| Eumovate |
| Everet |
| Everolimus |
| |
| Evista |
| Exemestane |
| Exjade |
| Extemporaneously Compounded |
| Preparations and |
| Galenicals |
| Eye Preparations |
| Eylea |
| Ezetimibe |
| Ezetimibe Sandoz56 |
| |
| Ezetimibe with simvastatin56 |
| - F - |
| - F - Factor eight inhibitor bypassing |
| - F - |
| - F - Factor eight inhibitor bypassing fraction |
| - F - Factor eight inhibitor bypassing fraction |
| - F - Factor eight inhibitor bypassing fraction |
| - F - Factor eight inhibitor bypassing fraction |
| - F - Factor eight inhibitor bypassing fraction |
| - F - Factor eight inhibitor bypassing fraction40 Famotidine |
| - F - Factor eight inhibitor bypassing fraction40 Famotidine |
| - F - Factor eight inhibitor bypassing fraction40 Famotidine |
| - F - Factor eight inhibitor bypassing fraction40 Famotidine |
| - F - Factor eight inhibitor bypassing fraction40 Famotidine |

| Fibro-vein Filgrastim Finasteride | |
|--|--|
| Filgrastim Finasteride | - 41 |
| Finasteride | |
| | |
| | |
| Fingolimod | |
| Firazyr | 228 |
| Flagy | 99 |
| FlagyI-S | |
| Flamazine | |
| Flecainide acetate | |
| Flecainide BNM | |
| | 49 |
| Flecainide Controlled Release | |
| Teva | 49 |
| Fleet Phosphate Enema | |
| Flixonase Hayfever & Allergy | 235 |
| Flixotide | 230 |
| Flixotide Accuhaler | 230 |
| Florinef | |
| Fluanxol | |
| | |
| Flucil | |
| Flucloxacillin | |
| Flucloxacillin-AFT | |
| Flucloxin | |
| Flucon | 238 |
| Fluconazole | 96 |
| Fludara Oral | |
| Fludarabine Ebewe | |
| Fludarabine phosphate | |
| Fludrocortisone acetate | |
| Fluids and Electrolytes | |
| | |
| | |
| Flumetasone pivalate | |
| Flumetasone pivalate Fluocortolone caproate with | |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and | 237 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine | 8 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine | 8 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone | 8 .238 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil | 237 8 238 149 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord | |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium | |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium | |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluox Fluoxetine hydrochloride | |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluox Fluoxetine hydrochloride Fluopenthixol decanoate | 237 8 .238 .149 .149 70 .126 .126 .133 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluox Fluoxetine hydrochloride Flupenthixol decanoate Flutamide | 237 8 238 149 149 70 126 126 133 166 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluoxtine hydrochloride Flupenthixol decanoate Flutamide Flutamin | 237 8 238 149 149 70 126 126 133 166 166 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluoxtine hydrochloride Flupenthixol decanoate Flutamide Flutamin | 237 8 238 149 149 70 126 126 133 166 166 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil Sodium Fluorouracil sodium Fluorouracil sodium Fluoretine hydrochloride Flupenthixol decanoate Flutamin Flutamin Flutasone | 237 8 238 149 149 70 126 126 133 166 166 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluorouracil Fluorouracil sodium Fluorouracil sodium | 237 8 238 149 149 126 126 133 166 133 166 230 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil | 237 8 238 149 149 70 126 133 166 230 230 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluox Fluorouracil sodium Fluox Fluoxetine hydrochloride Fluoxetine hydrochloride Fluoxetine hydrochloride Flutorasone fluotae with vilanterol Fluticasone propionate | 237 8 238 149 149 70 126 133 166 230 230 235 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sod | 237 8 238 149 149 70 126 133 166 230 235 231 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sod | 237 8 238 149 149 70 126 133 166 133 166 230 235 231 238 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord. Fluorouracil sodium Fluox Fluoxetine hydrochloride Fluoxetine hydrochloride | 237 8 238 149 149 126 126 133 166 230 230 235 231 238 62 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluorouracil sodium Fluoretine hydrochloride Fluoretine hydrochl | 237 8 238 149 149 70 126 133 166 166 230 235 231 238 62 38 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord. Fluorouracil sodium Fluox Fluoxetine hydrochloride Fluoxetine hydrochloride | 237 8 238 149 149 70 126 133 166 166 230 235 231 238 62 38 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil so | 237 8 238 149 149 70 126 133 166 230 235 231 238 62 38 38 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluoxetine hydrochloride Fluoxetine hydrochloride Fluoxet | 237 8 238 149 149 70 126 133 166 133 166 230 235 231 238 62 38 38 38 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Fluorethinkol decanoate Fluotenine hydrochloride Fluotenine hydrochloride Flutamide Flutamide Flutamin Fluticasone furoate with vilanterol Fluticasone propionate Fluticasone propionate Fluticasone with salmeterol FML Foban Folic acid Folic Acid multichem Folic Acid Mylan Food Thickeners | 237 8 238 149 149 70 126 133 166 133 166 230 235 231 238 62 38 38 38 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Accord Fluorouracil sodium Fluorouracil sodium Folic acid Food Sudi Mylan Food Sudi Supplements For Inborn | 237 8 238 149 149 70 126 133 166 230 235 231 238 62 38 38 259 |
| Flumetasone pivalate Fluocortolone caproate with fluocortolone pivalate and cinchocaine Fluorometholone Fluorouracil Fluorouracil Accord Fluorouracil sodium Fluoretine hydrochloride Fluoretine hydrochloride Fluor | 237 8 238 149 149 126 126 126 133 166 230 235 231 238 62 38 38 259 261 |

| _ | |
|-------------------------------|------|
| Forteo | |
| Fortini | |
| Fortini Multi Fibre | |
| Fortisip | 258 |
| Fortisip Multi Fibre | 258 |
| Fosamax | 112 |
| Fosamax Plus | 112 |
| Framycetin sulphate | 237 |
| Frisium | |
| Frumil | |
| Frusemide | |
| Fucicort | |
| Fucidin | |
| Fucithalmic | |
| Fulvestrant | |
| Fungilin | |
| Furosemid-Ratiopharm | .00 |
| | .00 |
| Furosemide [Frusemide] | . 53 |
| Furosemide-Baxter | . 53 |
| fusidic acid | ~ . |
| Dermatological 62 | |
| Infection | |
| Sensory | 237 |
| - G - | |
| Gabapentin | 127 |
| Gacet | |
| Galsulfase | |
| Galvumet | |
| Galvus | |
| Gardasil 9 | 270 |
| Gastrodenol | 9 |
| Gaviscon Double Strength | 6 |
| Gaviscon Infant | 6 |
| Gazyva | 199 |
| Gefitinib | |
| GEM Aqueous Cream | |
| Gemcitabine Ebewe | |
| Gemcitabine hydrochloride | 150 |
| Genoptic | |
| Gentamicin sulphate | |
| Infection | 94 |
| Sensory | |
| Gilenya | 136 |
| Ginet | |
| Glatiramer acetate | |
| Glecaprevir with pibrentasvir | 100 |
| Glibenclamide | 102 |
| | |
| Gliclazide | |
| Glipizide | |
| Glivec | |
| Glizide | .11 |
| Glucagen Hypokit | .10 |
| Glucagon hydrochloride | |
| Glucerna Select | |
| Glucose [Dextrose] | |
| Gluten Free Foods | 260 |

| Glycerin with sodium saccharin 245 |
|------------------------------------|
| Glycerin with sucrose245 |
| Glycerol |
| Alimentary |
| Extemporaneous245 |
| Glyceryl trinitrate |
| Alimentary |
| Cardiovascular |
| Glycopyrronium |
| Glycopyrronium bromide |
| Glycopyrronium with |
| indacaterol |
| Go Healthy |
| Gold Knight |
| Gold Knight XL |
| Goserelin |
| Gutron |
| Gynaecological Anti-infectives |
| - H - |
| наbitrol 144 |
| Haemophilus influenzae type B |
| |
| vaccine |
| Haldol |
| Haldol Concentrate |
| Haldol Decanoas |
| Haloperidol |
| Harvoni |
| Harvoni |
| Havrix Junior |
| |
| Haylor syrup |
| BP |
| healthE Dimethicone 10%65 |
| healthE Dimethicone 4% Lotion |
| healthE Dimethicone 5% |
| healthE Glycerol BP245 |
| healthE Urea Cream |
| Healtheries Simple Baking Mix |
| Hemastix |
| Hemlibra |
| Heparin sodium |
| Heparinised saline |
| Heparinised saline |
| Hepatitis A vaccine |
| Hepatitis B recombinant |
| vaccine |
| Herceptin |
| Hiberix |
| Hiprex |
| Histaclear |
| Histafen |
| Holoxan |
| Horleys Bread Mix |
| Horleys Flour |
| Hormone Replacement Therapy - |
| |

| Systemic 81 |
|--|
| HPV |
| Humalog |
| Humalog Mix 2511 |
| Humalog Mix 5011 Human papillomavirus (6, 11, 16, 18, |
| 31, 33, 45, 52 and 58) vaccine |
| [HPV] 270 |
| umatin |
| Humira |
| HumiraPen |
| Humulin 30/70 10 |
| Humulin NPH10 |
| Humulin R10 |
| Hyaluronic acid241 |
| Hydralazine |
| Hydralazine hydrochloride57 |
| Hydrocortisone |
| Dermatological |
| Hormone |
| Hydrocortisone (PSM)64 |
| Hydrocortisone acetate7 |
| Hydrocortisone acetate with |
| pramoxine hydrochloride7 |
| Hydrocortisone and paraffin liquid |
| and lanolin 64 |
| Hydrocortisone butyrate64, 70 |
| Hydrocortisone with cinchocaine8 |
| Hydrocortisone with miconazole 64 |
| Hydrocortisone with natamycin and |
| neomycin 64 |
| Hydrogen peroxide61 |
| Hydroxocobalamin33 |
| Hydroxocobalamin Mercury |
| Pharma33 |
| nydroxycarbamide153 |
| Hydroxychloroquine112 |
| Hydroxyurea |
| [hydroxycarbamide] 153 |
| Hygroton54 Hylo-Fresh241 |
| Tymenoptera228 |
| Hyoscine butylbromide8 |
| Hyoscine hydrobromide |
| Hypam |
| Hyperuricaemia and Antigout |
| Hypromellose |
| Hypromellose with dextran |
| -1- |
| biamox92 |
| brance |
| buprofen111 |
| catibant 228 |
| darubicin hydrochloride153 |
| dursulfase |
| fosfamide147 |
| |

| Igroton5 | 4 |
|------------------------------------|----|
| Ikorel5 | |
| lloprost6 | 0 |
| Imatinib mesilate16 | 0 |
| Imatinib-Rex16 | |
| Imigran13 | |
| Imipramine hydrochloride 12 | |
| Imiquimod7 | 0 |
| Immune Modulators10 | 17 |
| Immunosuppressants16 | |
| Imuran | 9 |
| Incruse Ellipta23 | 2 |
| Indacaterol23 | |
| Indapamide5 | |
| Infanrix IPV26 | 8 |
| Infanrix-hexa26 | |
| Infant Formulae26 | |
| Infatrini26 | |
| Infliximab19 | |
| Influenza vaccine27 | |
| Inhaled Corticosteroids22 | |
| Inhaled Long-acting | |
| Beta-adrenoceptor Agonists 23 | 0 |
| Inspra5 | |
| Instillagel Lido12 | |
| Insulin aspart 1 | |
| Insulin aspart with insulin aspart | |
| protamine1 | 0 |
| Insulin glargine1 | 1 |
| Insulin glulisine 1 | 1 |
| Insulin isophane1 | 0 |
| Insulin isophane with insulin | |
| neutral1 | 0 |
| Insulin lispro1 | |
| Insulin lispro with insulin lispro | |
| protamine1 | 1 |
| Insulin neutral1 | |
| Insulin pen needles1 | |
| Insulin pump 1 | 6 |
| Insulin pump cartridge2 | |
| Insulin pump infusion set (steel | |
| cannula)2 | 1 |
| Insulin pump infusion set (steel | |
| cannula, straight insertion) 2 | 1 |
| Insulin pump infusion set (teflon | |
| cannula)2 | 2 |
| Insulin pump infusion set (teflon | |
| cannula, angle insertion with | |
| insertion device) 2 | 3 |
| Insulin pump infusion set (teflon | |
| cannula, angle insertion) 2 | 3 |
| Insulin pump infusion set (teflon | |
| cannula, straight insertion with | |
| insertion device) 2 | 3 |
| Insulin pump infusion set (teflon | |
| cannula, straight insertion) 2 | 3 |
| | |

| Insulin pump reservoir | 00 |
|--|------------|
| | . 23 |
| Insulin syringes, disposable with | |
| attached needle | . 16 |
| Intelence | 105 |
| Interferon beta-1-alpha | 136 |
| Interferon beta-1-beta | |
| Intra-uterine device | |
| Invega Sustenna | |
| IPCA-Frusemide | 53 |
| IPCA-Metoprolol | |
| | .00 |
| IPCA-Propranolol | |
| IPOL | |
| Ipratropium bromide231, 2 | 235 |
| Iressa | 160 |
| Irinotecan Actavis 100 | 150 |
| Irinotecan hydrochloride | |
| Irinotecan-Rex | 150 |
| Iron (as ferric carboxymaltose) | . 35 |
| Iron polymaltose | |
| Isentress | |
| Isentress HD | 107 |
| Ismo 20 | |
| Ismo 40 Retard | . 57 |
| | |
| Isoniazid Isoniazid with rifampicin | 100 |
| | |
| Isoptin | . 52 |
| Isoptin Retard | . 52 |
| Isoptin SR | |
| Isopto Carpine | |
| Isosorbide mononitrate | |
| Isosource Standard | |
| Isotretinoin | |
| Ispaghula (psyllium) husk | .25 |
| Itch-Soothe | .63 |
| Itraconazole | .97 |
| Itrazole | .97 |
| Ivacaftor | |
| Ivermectin | .66 |
| - J - | |
| Jadelle | .74 |
| Jakavi | 163 |
| Jardiamet | |
| Jardiance | 13 |
| Jaydess | 00 |
| Jevity HiCal RTH | .02 057 |
| | 207 |
| Jevity Plus | |
| Jevity RTH | 257 |
| Juno Pemetrexed | 150 |
| - K - | |
| Kadcyla | |
| Kaletra | |
| Kalydeco | |
| Kemadrin | 120 |
| Kenacomb | |
| Kenacort-A 10 | . 80 |
| Kenacort-A 40 | . 80 |

| Kenalog in Orabase33 |
|-------------------------------|
| Ketocal 3:1 |
| KetoCal 4:1 |
| Ketoconazole |
| Dermatological |
| Infection |
| Ketogenic Diet |
| |
| Ketoprofen 111 |
| KetoSens13 |
| Keytruda |
| Kindergen251 |
| Kivexa106 |
| Klacid |
| Alimentary9 |
| Infection90 |
| Kliogest82 |
| Kliovance82 |
| Kogenate FS41 |
| Konakion MM |
| Konsyl-D |
| Kuvan |
| - L - |
| - L - Labetalol |
| |
| Lacosamide |
| Lactulose |
| Laevolac |
| Lagevrio104 |
| Lamictal 128 |
| Lamivudine 101, 106 |
| Lamivudine Alphapharm106 |
| Lamotrigine128 |
| Lamprene99 |
| Lanoxin |
| Lanoxin Paediatric Elixir49 |
| Lanoxin PG49 |
| Lanoxin S29 |
| Lansoprazole |
| Lantus11 |
| Lantus SoloStar11 |
| Lanvis |
| Lanzol Relief |
| Lanzon Hener |
| Largactil |
| Largacui |
| Laronidase |
| Lasix |
| Latanoprost240 |
| Latanoprost with timolol240 |
| Lax-Suppositories26 |
| Lax-Tab26 |
| Laxatives25 |
| Laxsol25 |
| Ledipasvir with sofosbuvir102 |
| Leflunomide112 |
| Lenalidomide |
| Letrole |
| Letrozole |
| |

| Leukeran FC 147 |
|---------------------------------|
| Leukotriene Receptor |
| Antagonists 234 |
| Leuprorelin |
| Leustatin149 |
| Levetiracetam 128 |
| Levetiracetam-AFT128 |
| Levlen ED74 |
| Levocabastine |
| Levocarnitine |
| Levodopa with benserazide119 |
| Levodopa with carbidopa119 |
| Levomepromazine 132 |
| Levomepromazine |
| hydrochloride 133 |
| Levonorgestrel |
| Genito-Urinary74-75 |
| Hormone |
| Levothyroxine83 |
| Lidocaine [Lignocaine] 120-121 |
| Lidocaine [Lignocaine] |
| hydrochloride 121 |
| Lidocaine [Lignocaine] with |
| chlorhexidine 121 |
| Lidocaine [Lignocaine] with |
| prilocaine 121 |
| Lidocaine-Baxter121 |
| Lidocaine-Claris121 |
| Life Extension |
| Lignocaine 120–121 |
| Lioresal Intrathecal 118 |
| Lipid-Modifying Agents54 |
| Liquigen249 |
| Lisinopril47 |
| Lithium carbonate 133 |
| Livostin |
| LMX4 |
| Locacorten-Viaform ED's237 |
| Local preparations for Anal and |
| Rectal Disorders 8 |
| Locasol |
| Locoid |
| Locoid Crelo64 |
| Locoid Lipocream |
| Locorten-Vioform |
| Lodoxamide |
| Logem |
| Lomide |
| Lomustine |
| Loniten |
| Loperamide hydrochloride |
| Lopinavir with ritonavir |
| Lopinavir/Ritonavir Mylan 106 |
| Loprofin |
| Loprofin Mix |
| Lorofix |

| Loratadine |
|--|
| Lorazepam135 |
| Lorstat |
| Losartan Actavis |
| Losartan potassium |
| Losartan potassium with |
| hydrochlorothiazide |
| Lovir |
| |
| Loxamine |
| Lucrin Depot 1-month |
| Lucrin Depot 3-month |
| Lyderm |
| Lynparza154 |
| Lyrica |
| - M - |
| m-Eslon |
| Mabthera |
| Macro Organic Psyllium Husk25 |
| Macrobid110 |
| Macrogol 3350 with potassium |
| chloride, sodium bicarbonate and |
| sodium chloride26 |
| Macrogol 400 and propylene |
| glycol 241 |
| Madopar 125 119 |
| Madopar 250 119 |
| Madopar 62.5 119 |
| Madopar HBS 119 |
| Madopar Rapid119 |
| Magnesium hydroxide36 |
| Magnesium sulphate36 |
| Mantoux277 |
| Marevan44 |
| Marine Blue Lotion SPF 50+70 |
| Martindale Pharma 242 |
| Mask for spacer device235 |
| Maviret |
| Maxidex |
| Maxitrol |
| MCT oil (Nutricia)249 |
| Measles, mumps and rubella |
| vaccine 273 |
| Mebendazole89 |
| Mebeverine hydrochloride8 |
| Medrol79 |
| Medroxyprogesterone acetate |
| Genito-Urinary74 |
| Hormone 82 |
| Mefenamic acid 111 |
| Megace |
| Megestrol acetate |
| Melatonin |
| Melphalan |
| Menactra |
| Meningococcal (groups A, C, Y and |
| W-135) conjugate vaccine |
| ······································ |

| Meningococcal B multicomponent |
|---|
| vaccine 274 |
| Meningococcal C conjugate |
| vaccine 274 |
| Menthol63 |
| Mepolizumab 198 |
| Mercaptopurine150 |
| Mercilon 2873 |
| Mesalazine7 |
| Mesna154 |
| Mestinon 111 |
| Metabolic Disorder Agents27 |
| Metformin hydrochloride 11 |
| Metformin Mylan 11 |
| Methadone hydrochloride |
| Extemporaneous245 |
| Nervous123 |
| Methatabs123 |
| Methenamine (hexamine) |
| hippurate 109 |
| Methopt240 |
| Methotrexate150 |
| Methotrexate DBL Onco-Vial150 |
| Methotrexate Ebewe150 |
| Methotrexate Sandoz 150 |
| Methyl hydroxybenzoate245 |
| Methylcellulose245 |
| Methylcellulose with glycerin and |
| |
| sodium saccharin245 |
| sodium saccharin |
| Methylcellulose with glycerin and sucrose |
| Methylcellulose with glycerin and sucrose |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 135 Methylphenidate hydrochloride 135 Methylphenidate hydrochloride 140 Methylphenidate 1 140 |
| Methylcellulose with glycerin and sucrose |
| Methylcellulose with glycerin and sucrose |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate Provide 25 Methylphenidate hydrochloride 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 140 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 136 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 140 Methylprednisolone 76 Methylprednisolone (as sodium succinate) 80 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 136 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 140 Methylprednisolone 76 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 140 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 26 Methylprednisolone acetate 27 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 136 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 140 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone aceponate 84 Methylprednisolone acetate 84 Methylprednisolone acetate 84 Methylprednisolone acetate 84 Methylprednisolone acetate 84 Methylprednisolone acetate |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 140 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 30 Methylprednisolone acetate 31 Metoclopramide Actavis 10 131 Metoclopramide Actavis 10 131 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 136 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 140 Methylprenidate hydrochloride 75 Methylprednisolone 76 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone aceponate 80 Methylprednisolone aceponate 81 Methylprednisolone aceponate 81 Methylprednisolone aceponate 82 Methylprednisolone aceponate 81 Methylprednisolone aceponate 82 Methylprednisolone aceponate 82 Methylprednisolone aceponate 82 Methylprednisolone aceponate 82 Methylprednisolone acetate 82 Metoclopramide Actavis 10 131 Metoclopramide hydrochloride 131 Metolazone 52 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphonidate S29 52 Methylphenidate ER - Teva 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 140 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 81 Metolopramide Actavis 10 131 Metolopramide hydrochloride 131 Metolopramide hydrochloride 53 Metopirone 86 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 135 Methylphenidate Hydrochloride 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 136 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone aceponate 64 Methylprednisolone aceponate 64 Methylprednisolone aceponate 36 Methylprednisolone aceponate 37 Methylprednisolone aceponate 36 Methylprednisolone aceponate 36 Methylprednisolone aceponate 37 Metolopramide Actavis 10 31 Metolopramide hydrochloride 33 Metopirone 36 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 135 Methylphenidate Phylochloride 135 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 137 Methylphenidate hydrochloride 136 Methylphenidate hydrochloride 137 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone aceponate 64 Methylprednisolone aceponate 64 Methylprednisolone aceponate 56 Methylprednisolone aceponate 56 Methylprednisolone aceponate 56 Methylprednisolone aceponate 56 Metopirone 38 Metopirone 36 Metoprolol IV Mylan 50 Metoprolol IV Mylan 50 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 138 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone aceponate 64 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 50 Metoclopramide Actavis 10 131 Metoclopramide hydrochloride 131 Metoprolol IV Mylan 56 Metoprolol V Mylan 56 Metoprolol V Mylan 56 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 138 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 50 Methylprednisolone 50 Metoprolol V Mylan 50 Metoprolol V Mylan 50 |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 138 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 138 Methylphenidate hydrochloride 138 Methylprednisolone 76 Methylprednisolone 76 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 50 Metoclopramide Actavis 10 131 Metoclopramide hydrochloride 131 Metoprolol IV Mylan 50 Metoprolol V |
| Methylcellulose with glycerin and sucrose 245 Methyldopa 52 Methyldopa Mylan 52 Methyldopa Mylan S29 52 Methyldopa Mylan S29 52 Methylphenidate ER - Teva 138 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylphenidate hydrochloride 139 Methylprednisolone 75 Methylprednisolone 75 Methylprednisolone (as sodium succinate) 80 Methylprednisolone aceponate 64 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 80 Methylprednisolone acetate 50 Methylprednisolone 50 Metoprolol V Mylan 50 Metoprolol V Mylan 50 |

| Miacalcic | |
|--|-----|
| Micolette | 26 |
| Miconazole | 33 |
| Miconazole nitrate | |
| Dermatological | 63 |
| Genito-Urinary | 75 |
| Micreme | |
| Micreme H | |
| Microgynon 20 ED | 74 |
| Microgynon 30 | 74 |
| Microlut | 74 |
| Midazolam1 | 37 |
| Midazolam-Baxter | 37 |
| Midodrine | |
| Mifegyne | |
| Mifepristone | |
| Minerals | |
| Mini-Wright AFS Low Range | 25 |
| Mini-Wright Standard | 235 |
| Minidiab | 11 |
| MiniMed 1.8 Reservoir | |
| MMT-326A | 23 |
| MiniMed 3.0 Reservoir | 20 |
| MMT-332A | 22 |
| MiniMed 770G | 16 |
| MiniMed Mio MMT-921A | 20 |
| MiniMed Mio MMT-923A | |
| MiniMed Mio MMT-925A | |
| MiniMed Mio MMT-925A | |
| MiniMed Mio MMT-943A | |
| MiniMed Mio MMT-945A | |
| MiniMed Mio MMT-965A | 22 |
| | |
| MiniMed Mio MMT-975A MiniMed Quick-Set MMT-386A | 22 |
| MiniMed Quick-Set MMT-387A | 22 |
| MiniMed Quick-Set MMT-396A | 22 |
| MiniMed Quick-Set MMT-396A | 22 |
| MiniMed Quick-Set MMT-397A | 22 |
| MiniMed Quick-Set MMT-398A | 22 |
| MiniMed Quick-Set MMT-399A | 22 |
| MiniMed Silhouette MMT-368A MiniMed Silhouette MMT-377A | 22 |
| MiniMed Silhouette MMT-377A | 22 |
| MiniMed Silhouette MMT-378A | 22 |
| MiniMed Silhouette MMT-381A | 22 |
| MiniMed Silhouette MMT-382A | 22 |
| MiniMed Silhouette MMT-383A | 22 |
| MiniMed Silhouette MMT-384A | |
| MiniMed Sure-T MMT-864A | 21 |
| MiniMed Sure-T MMT-866A | |
| MiniMed Sure-T MMT-874A | |
| MiniMed Sure-T MMT-876A | |
| MiniMed Sure-T MMT-884A | |
| MiniMed Sure-T MMT-886A | |
| Minims Pilocarpine | 240 |
| Minims Prednisolone | |
| Minirin | |
| Minirin Melt | 88 |

| Mino-tabs Q3 |
|--|
| Mino-tabs |
| Minomycin |
| Minor Skin Infections |
| |
| Minoxidil |
| Mirena |
| Mirtazapine |
| Misoprostol |
| Mitomycin C154 |
| Mitozantrone154 |
| Mitozantrone Ebewe154 |
| Mixtard 3010 |
| MMR II273 |
| Moclobemide 126 |
| Modafinil 141 |
| Modavigil141 |
| Moduretic54 |
| Molaxole |
| Molnupiravir 104 |
| Moments |
| Mometasone furoate |
| Monogen |
| Montelukast |
| Montelukast Mylan |
| Moroctocog alfa [Recombinant factor |
| |
| VIII] |
| Morphine hydrochloride 123 |
| Morphine sulphate 123 |
| Motetis 120 |
| Mouth and Throat32 |
| Movapo119 |
| Moxifloxacin94 |
| MSUD Maxamum261 |
| Mucilaginous laxatives with |
| |
| stimulants 25 |
| stimulants |
| Mucolytics234 |
| Mucolytics |
| Mucolytics |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 114 Mvite 34 Myambutol 99 Mycobutin 100 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 MycoNail 62 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycobutin 62 Mycophenolate mofetil 169 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 62 Mycophenolate mofetil 169 Mydriacyl 240 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobhenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 Mylan (24 hr release) 52 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 Mylan (24 hr release) 52 Mylan (24 hr release) 50 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 Mylan (24 hr release) 52 Mylan Clomiphen 88 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 Mylan (24 hr release) 52 Mylan Clomiphen 88 Mylan Indapamide 54 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal. 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mylan (12 hr release) 52 Mylan (12 hr release) 52 Mylan Atenolol 52 Mylan Indapamide 54 Mylan Italy (24 hr release) 52 |
| Mucolytics 234 Mucosoothe 121 Multiple Sclerosis Treatments 135 Multivitamin renal 34 Multivitamins 34 Mupirocin 62 Muscle Relaxants 118 Mvite 34 Myambutol 99 Mycobutin 100 Mycophenolate mofetil 169 Mydriacyl 240 Mylan (12 hr release) 52 Mylan (24 hr release) 52 Mylan Clomiphen 88 Mylan Indapamide 54 |

| Preparations75 |
|---------------------------------|
| Myozyme |
| - N - |
| Nadolol |
| Nadolol BNM50 |
| Naglazyme28 |
| Nalcrom8 |
| Naloxone hydrochloride242 |
| Naltraccord 143 |
| Naltrexone hydrochloride143 |
| Naphazoline hydrochloride |
| Naphcon Forte |
| Naprosyn SR 1000 111 |
| Naprosyn SR 750 111 |
| Naproxen |
| Narcaricin mite |
| Natalizumab137 |
| Natulan |
| Nausafix |
| Nausicalm |
| Navelbine158 |
| Nefopam hydrochloride |
| Neisvac-C |
| Neo-B12 |
| Neo-Mercazole |
| Neocate Gold263 |
| Neocate Junior Unflavoured |
| Neocate Junior Vanilla263 |
| Neocate SYNEO263 |
| Neoral |
| Neostigmine metilsulfate111 |
| Nepro HP (strawberry)253 |
| Nepro HP (vanilla) |
| Nepro HP RTH 252 |
| Neulactil133 |
| Neulastim |
| NeuroTabs |
| Nevirapine106 |
| Nevirapine Alphapharm |
| Nicorandil |
| Nicotine |
| Nifedipine52 Nifuran110 |
| Nilotinib |
| Nilstat |
| |
| Alimentary 33 Genito-Urinary |
| Infection |
| Nintedanib |
| Nipent |
| Nirmatrelvir with ritonavir |
| Nitrates |
| Nitroderm TTS |
| Nitrofurantoin 110 |
| Nitrolingual Pump Spray |
| |

| Nivestim | 44 |
|---------------------------------|-----|
| Nivolumab | 221 |
| Nizoral | |
| Nodia | |
| Noflam 250 | 111 |
| Noflam 5001 | 11 |
| Non-Steroidal Anti-Inflammatory | |
| Drugs1 | 11 |
| Nonacog gamma, [Recombinant | |
| Factor IX] | 41 |
| Norothistorono | |
| Genito-Urinary | .74 |
| Hormone | 82 |
| Norflex1 | 18 |
| Norfloxacin1 | 10 |
| Noriday 28 | 74 |
| Norimin | 74 |
| Normacol Plus | 25 |
| Normison1 | 38 |
| Norpress1 | 26 |
| Nortriptyline hydrochloride1 | 26 |
| Norvir 1 | 07 |
| Noumed1 | 27 |
| Noumed Paracetamol1 | 22 |
| NovaSource Renal | 253 |
| Novatretin | 68 |
| NovoMix 30 FlexPen | 10 |
| NovoRapid | 11 |
| NovoRapid FlexPen | 11 |
| NovoRapid Penfill | 11 |
| NovoSeven RT | 40 |
| Noxafil | 97 |
| Nozinan 132-1 | 33 |
| Nozinan (Swiss)1 | 32 |
| Nucala1 | 98 |
| Nuelin | 234 |
| Nuelin-SR | 234 |
| Nupentin 1 | 27 |
| Nutilis | 260 |
| Nutren Diabetes | |
| Nutrient Modules | 247 |
| Nutrini Energy Multi Fibre | 252 |
| Nutrini Energy RTH | 252 |
| Nutrini Low Energy Multi Fibre | 254 |
| Nutrini Peptisorb | 265 |
| Nutrini Peptisorb Energy | 265 |
| Nutrini RTH | 252 |
| Nutrison 800 Complete Multi | |
| Fibre | 257 |
| Nutrison Concentrated | 259 |
| Nutrison Energy | 257 |
| Nutrison Energy Multi Fibre | 257 |
| Nutrison Multi Fibre | 257 |
| | |
| Nyefax Retard | 52 |

Nvstatin Genito-Urinary......75 Infection97 NZB Low Gluten Bread Mix......260 -0-O/W Fatty Emulsion Cream......65 Obinutuzumab 199 Obstetric Preparations......77 Ocrelizumab137 Octocog alfa [Recombinant factor VIII] (Advate)..... 41 Octocog alfa [Recombinant factor VIII] (Kogenate FS)..... 41 Octreotide167 Octreotide (Sun) 167 Octreotide Depot Teva 167 Octreotide GH.....167 Octreotide long-acting167 Octreotide MaxRx......167 Oestradiol81 Oestradiol valerate81 Oestradiol with norethisterone......82 Oestriol Genito-Urinary.....75 Hormone82 Oestrogens......81 Oil in water emulsion65 Olanzapine 133-134 Olaparib.....154 Olbetam......54 Olbetam S29.....54 Olopatadine Teva241 Omalizumab199 Omeprazole......9 Omeprazole actavis 10.....9 Omeprazole actavis 20.....9 Omeprazole actavis 40.....9 Omnitrope......83 Oncaspar LYO.....155 OncoTICE.....176 Ondansetron.....131 Ondansetron ODT-DRLA 131 Onrex.....131 Ora-Blend SF......245 Ora-Plus245

| Orabase3 | 2 |
|---------------------------------------|----|
| Oral and Enteral Feeds25 | 0 |
| Oratane | 1 |
| Ordine | 0 |
| | |
| Orgran | 1 |
| Ornidazole9 | 9 |
| Orphenadrine citrate11 | 8 |
| Ortho-tolidine7 | 7 |
| Oruvail SR11 | 1 |
| Osmolite RTH25 | 7 |
| Other Endocrine Agents8 | 8 |
| Other Oestrogen Preparations | 2 |
| Other Data and the second | |
| Other Progestogen Preparations | 2 |
| Other Skin Preparations7 | 'n |
| Ovestin | 0 |
| Genito-Urinary7 | - |
| | |
| Hormone8 | |
| Oxaliplatin14 | 8 |
| Oxaliplatin Accord14 | |
| Oxaliplatin Actavis 10014 | 8 |
| | 8 |
| Oxis Turbuhaler23 | 0 |
| Oxpentifylline5 | 7 |
| Oxybutynin7 | 6 |
| Oxycodone hydrochloride12 | 4 |
| Oxycodone Sandoz 12 | 4 |
| Oxycodone Sandoz S2912 | 4 |
| DxyContin | 4 |
| OxyNorm | 4 |
| Oxytocin | |
| Oxytocin BNM7 | |
| | 9 |
| Oxytocin with ergometrine maleate7 | _ |
| | |
| Ozurdex23 | 8 |
| - P - | |
| Pacifen11 | 8 |
| Pacimol12 | 2 |
| Paclitaxel15 | 5 |
| Paclitaxel Actavis15 | 5 |
| Paclitaxel Ebewe15 | 5 |
| Paediatric Seravit3 | 4 |
| Palbociclib16 | 2 |
| Paliperidone13 | |
| Pamidronate disodium11 | 3 |
| Pamisol11 | |
| Pancreatic enzyme2 | |
| Pantonrazolo | 0 |
| Pantoprazole | 3 |
| Panzop Relief | 9 |
| Panzytrat2 | |
| Papaverine hydrochloride5 | |
| Para-amino salicylic acid 10 | |
| Paracare 12 | |
| Paracare Double Strength 12 | |
| Paracetamol12 | 2 |
| Paracetamol + Codeine | |

| (Relieve) | 125 |
|---------------------------------|-------|
| Paracetamol with codeine | .125 |
| Paraffin | |
| Paraffin liquid with wool fat | |
| Parasiticidal Preparations | |
| | 00 |
| Parlodel | . 119 |
| Parnate | . 126 |
| Parnate S29 | . 126 |
| Paromomycin | 95 |
| Paroxetine | . 126 |
| Paser | . 100 |
| Paxam | .135 |
| Paxlovid | .104 |
| Pazopanib | 162 |
| Peak flow meter | 235 |
| Pedialyte - Bubblegum | |
| Pediagura | 050 |
| Pediasure | . 252 |
| Pediasure RTH | |
| Pegaspargase | . 155 |
| Pegasys | . 107 |
| Pegfilgrastim | 44 |
| Pegylated interferon alfa-2a | |
| Pembrolizumab | .223 |
| Pemetrexed | . 150 |
| Penicillamine | |
| Penicillin G | |
| PenMix 30 | 10 |
| PenMix 40 | 10 |
| PenMix 50 | 10 |
| | |
| Pentasa | |
| Pentostatin [Deoxycoformycin] | . 155 |
| Pentoxifylline [Oxpentifylline] | |
| Peptamen Junior | .252 |
| Peptisorb | |
| Perhexiline maleate | 52 |
| Pericyazine | .133 |
| Perindopril | 48 |
| Perjeta | .201 |
| Permethrin | |
| Perrigo | |
| Pertuzumab | 201 |
| Peteha | 100 |
| Pethidine hydrochloride | 100 |
| Petiniume nyurochionue | . 120 |
| Pevaryl | |
| Pexsig | |
| Pfizer Exemestane | . 169 |
| Pharmacy Health Sorbolene with | |
| Glycerin | 65 |
| Pharmacy Services | .242 |
| Pheburane | 30 |
| Phenasen | .151 |
| Phenobarbitone | |
| Phenobarbitone sodium | 0 |
| Extemporaneous | 245 |
| Nervous | 127 |
| Phenoxybenzamine | . 137 |
| FILEHUXVDEHZällille | |
INDEX: Generic Chemicals and Brands

| hydrochloride | 47 |
|---------------------------------------|-------|
| Phenoxymethylpenicillin (Penicillin | 47 |
| | 02 |
| V) | |
| Phenytoin sodium 127 | -128 |
| Phillips Milk of Magnesia | 36 |
| Phlexy 10 | . 262 |
| Phosphate Phebra | |
| Phosphorus | |
| Phytomenadione | |
| Pilocarpine hydrochloride | |
| Pimafucort | 64 |
| Pimecrolimus | |
| Pindolol | |
| Pine tar with trolamine laurilsulfate | |
| and fluorescein | 69 |
| Pinetarsol | |
| Pioglitazone | |
| Pirfenidone | |
| Pizotifen | |
| | |
| PKU Anamix Infant | |
| PKU Anamix Junior | |
| PKU Anamix Junior Chocolate | . 262 |
| PKU Anamix Junior LQ | |
| PKU Anamix Junior Orange | |
| PKU Anamix Junior Vanilla | |
| PKU Lophlex LQ 10 | .262 |
| PKU Lophlex LQ 20 | |
| PKU Lophlex Powder | |
| PKU Lophlex Sensation 20 | 262 |
| Plaquenil | 112 |
| Plendil ER | 51 |
| Pneumococcal (PCV10) conjugate | |
| vaccine | 074 |
| Pneumococcal (PCV13) conjugate | . 214 |
| | 075 |
| vaccine | . 275 |
| Pneumococcal (PPV23) | |
| polysaccharide vaccine | |
| Pneumovax 23 | |
| Podophyllotoxin | |
| Polaramine | |
| Poliomyelitis vaccine | |
| Poloxamer | 25 |
| Polv-Gel | .241 |
| Poly-Tears | .240 |
| Poly-Visc | 241 |
| Polycal | |
| Ponstan | |
| Posaconazole | |
| Postinor-1 | |
| | |
| Potassium chloride4 | |
| Potassium citrate | |
| Potassium iodate | |
| Povidone iodine | |
| Pradaxa | |
| Pramipexole hydrochloride | |
| Pravastatin | 54 |

| Pravastatin Mylan | F 4 |
|---|-----------------|
| | |
| Praziquantel | |
| Prazosin | 47 |
| Pred Forte | 239 |
| Prednisolone | 80 |
| Prednisolone acetate | 239 |
| Prednisolone sodium | |
| Prednisolone sodium | |
| phosphate | 220 |
| Prednisolone-AFT | 000 |
| | |
| Prednisone | |
| Prednisone Clinect | |
| Pregabalin | |
| Pregabalin Pfizer | 128 |
| Pregnancy Tests - hCG Urine | 76 |
| Premarin | 81 |
| Prevenar 13 | |
| Priadel | |
| Primaguine | |
| Primidone | |
| Primolut N | . 123 |
| | |
| Priorix | |
| Probenecid | |
| Probenecid-AFT | 118 |
| Procaine penicillin | |
| Procarbazine hydrochloride | |
| Prochlorperazine | 131 |
| Proctofoam | |
| Proctosedyl | |
| Procyclidine hydrochloride | 120 |
| Progesterone | 82 |
| Proglicem | 10 |
| Proglycem | |
| | |
| Progynova | |
| Prolia | |
| Promethazine hydrochloride | |
| Propafenone hydrochloride | 49 |
| Propamidine isethionate | 237 |
| Propranolol | 51 |
| Propylene glycol | 245 |
| Propylthiouracil | 83 |
| Prostacur | |
| Protaphane | |
| Protaphane Penfill | 10 |
| Protifar | |
| Protionamide | 100 |
| Provera | |
| | |
| Provera HD | |
| PSM Citalopram | 126 |
| Psoriasis and Eczema | |
| Preparations | <mark>68</mark> |
| PTU | |
| Pulmicort Turbuhaler | 229 |
| Pulmozyme | 234 |
| Puri-nethol | |
| Puria | |
| · • • • • • • • • • • • • • • • • • • • | |

| Pyrazinamide1 | 00 |
|---|--|
| Pyridostigmine bromide 1 | 11 |
| Pyridoxine hydrochloride | |
| Pyridoxine multichem | 33 |
| Pyrimethamine | 95 |
| Pytazen SR | 42 |
| Pytazen SR | |
| Quetapel1 | 33 |
| Ouetianine | 22 |
| Quetiapine 1 Quick-Set MMT-392 | 22 |
| Quick-Set MMT-393 | 20 |
| Quinapril | 10 |
| Quinapril with | 40 |
| Quinapril with hydrochlorothiazide | 40 |
| | 40 |
| Qvar2 | 29 |
| -K- | ~~ |
| RA-Morph 1 | 23 |
| Raloxifene hydrochloride 1 | 13 |
| Raltegravir potassium1 | 07 |
| Ramipex 1 | 19 |
| Ranbaxy-Cefaclor | 89 |
| Ranbaxy-Cefaclor Ranbaxy-Cefaclor S29 | 89 |
| Rapamune | 24 |
| Rasagiline1 | 19 |
| Reandron 1000 | 81 |
| Recombinant factor IX | 41 |
| Recombinant factor VIIa | 40 |
| Recombinant factor VIII 40- | 41 |
| Rectogesic | 8 |
| D III I | |
| Redipred | 80 |
| Redipred1 | 80 |
| Relieve1 | 80 11 |
| Relieve1 Relistor | 80 11 25 |
| Relieve | 80 11 25 91 |
| Relieve1 Relistor Remicade1 Renilon 7.5 | 80 11 25 91 53 |
| Relieve1 Relistor Remicade1 Renilon 7.52 Resonium-A. | 80 11 25 91 53 46 |
| Relieve1 Relistor | 80 11 25 91 53 46 49 |
| Relieve1 Relistor | 80 11 25 91 53 46 49 31 |
| Relieve1 Relistor | 80 11 25 91 53 46 249 31 35 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 | 80 11 25 91 53 46 49 31 35 36 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 Retinol palmitate 2 | 80 11 25 91 53 46 49 31 35 36 241 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 Retinol palmitate 2 ReTrieve 2 | 80 11 25 91 53 46 49 31 35 36 241 61 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A. 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 Retinol palmitate 2 Retrovir 1 | 80 11 25 91 53 46 49 31 35 36 41 61 06 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 Retinol palmitate 2 Retrovir 1 Revimid 1 | 80 11 25 91 53 46 49 31 35 36 41 61 06 53 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A. 2 Respiratory Devices 2 Respiratory Devices 2 Retroiv Stimulants 2 Retroive 2 Retroivir 1 Revolade 1 | 80 11 25 91 53 46 249 31 35 36 41 61 06 53 38 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respigen 2 Respiratory Devices 2 Retinol palmitate 2 Retrovir 1 Revlimid 1 Revolade 2 | 80 11 25 91 53 46 249 35 36 41 61 53 38 39 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respiratory Devices 2 Retinol palmitate 2 Retroir 1 Revolade 1 Revolade 2 Retactor 2 Retactor 2 Retroir 1 Revolade 2 Retactor 2 Riboflavin 2 | 80 11 25 91 53 46 249 31 35 36 41 61 06 53 38 29 29 |
| Relieve 1 Relistor 1 Renicade 1 Renilon 7.5 2 Resonium-A 2 Respigen 2 Respiratory Devices 2 Retinol palmitate 2 Retroir 1 Revolade 1 Revolade 2 Resacrom 2 Riboflavin 1 | 80 11 25 91 25 46 249 31 35 241 61 06 53 38 29 46 |
| Relieve 1 Relistor 1 Renicade 1 Renilon 7.5 2 Resonium-A 2 Respiratory Devices 2 Respiratory Devices 2 Respiratory Devices 2 Retinol palmitate 2 Retrive 1 Revolade 1 Revacrom 2 Riboflavin 1 Ribomustin 1 | 80 11 25 91 53 46 29 31 35 36 41 61 06 53 38 29 46 76 |
| Relieve 1 Relistor 1 Renicade 1 Renilon 7.5 2 Resonium-A 2 Respiratory Devices 2 Respiratory Devices 2 Respiratory Stimulants 2 Retriol palmitate 2 Retrovir 1 Revolade 2 Riboflavin 1 Riot 1 Riot 1 Riott 1 Riott 1 Riott 1 Restration 2 | 80 11 25 91 53 46 249 35 249 35 241 61 06 53 38 29 46 76 00 |
| Relieve 1 Relistor 1 Relistor 1 Renicade 1 Renicon 7.5 2 Resonium-A 2 Respiratory Devices 2 Respiratory Devices 2 Respiratory Stimulants 2 Retriol palmitate 2 Retrovir 1 Revolade 2 Revolade 2 Riboflavin 1 Riboflavin 1 Ridoutin 1 Rifabutin 1 | 80 11 25 91 253 46 249 31 35 241 61 06 53 38 29 46 76 00 00 |
| Relieve 1 Relistor 1 Renicade 1 Renicon 7.5 2 Resonium-A 2 Respired 2 Respired 2 Respiratory Devices 2 Respiratory Stimulants 2 Retrive 2 Retrive 2 Retrovir 1 Revolade 2 Ribomustin 1 Ridatin 1 Rifadin 1 Rifagin 1 | 80 11 25 91 53 46 249 31 35 36 41 61 06 53 38 29 46 76 00 00 00 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respired 2 Respired 2 Respiratory Devices 2 Respiratory Stimulants 2 Retrivel palmitate 2 Retrive 2 Retrovir 1 Revolade 2 Riboflavin 1 Ritodlautin 1 Ridatin 1 Rifadin 1 Rifarpicin 1 Rifarmin 1 | 80 11 25 91 25 46 249 35 36 41 61 53 38 39 29 46 76 00 00 9 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A. 2 Respigen 2 Respiratory Devices 2 Respiratory Stimulants 2 Retrovir 1 Revolade 2 Revolade 2 Riboflavin 1 Ridoflavin 1 Rifabutin 1 Rifabutin 1 Rifanjicin 1 Rifamin 1 Rifamin 1 Rifanin 1 Rifanin 1 Rifanin 1 | 80 11 25 91 53 46 249 35 36 41 61 53 8 39 29 46 76 00 00 00 00 00 00 00 00 00 00 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A. 2 Respigen 2 Respiratory Devices 2 Retinol palmitate 2 Retrovir 1 Revolade 2 Reiboflavin 1 Ribonustin 1 Rifabutin 1 Rifadin 1 Rifadin 1 Rifanpicin 1 Rifanh 1 Rifuek 1 | 80 11 25 91 53 46 49 31 35 36 41 61 65 38 39 46 76 00 00 00 00 00 20 |
| Relieve 1 Relistor 1 Remicade 1 Renilon 7.5 2 Resonium-A 2 Respired 2 Respired 2 Respiratory Devices 2 Respiratory Stimulants 2 Retrivel palmitate 2 Retrive 2 Retrovir 1 Revolade 2 Riboflavin 1 Ritodlautin 1 Ridatin 1 Rifadin 1 Rifarpicin 1 Rifarmin 1 | 80 11 25 91 53 46 49 31 35 36 41 61 65 38 39 46 76 00 00 00 00 00 20 |

INDEX: Generic Chemicals and Brands

| Riodine |
|-------------------------------------|
| Risedronate Sandoz114 |
| |
| Risedronate sodium114 |
| Risperdal Consta |
| Risperidone 133–134 |
| Risperidone (Teva)133 |
| Risperon 133 |
| Ritalin139 |
| Ritalin LA 140 |
| Ritonavir107 |
| Rituximab (Mabthera)201 |
| Rituximab (Riximyo) |
| Rivaroxaban |
| Rivastigmine |
| Rivastigmine Patch BNM 10141 |
| Rivastigmine Patch BNM 5141 |
| |
| Rivotril |
| Riximyo203 |
| RIXUBIS41 |
| Rizamelt130 |
| Rizatriptan 130 |
| RoActemra S29216 |
| Ronapreve 190 |
| Ropin |
| Ropinirole hydrochloride119 |
| Rosuvastatin55 |
| Rosuvastatin Viatris55 |
| Rotarix |
| Rotavirus oral vaccine |
| Roxane |
| Roxane-Propranolol |
| Roxithromycin |
| Rubifen |
| Rubifen SR |
| Rublien SR |
| Rugby Capsaicin Topical Cream |
| Cream |
| Rulide D91 |
| Rurioctocog alfa pegol [Recombinant |
| factor VIII] 41 |
| Ruxolitinib |
| Rythmodan 49 |
| Rytmonorm49 |
| - S - |
| Sabril 129 |
| Sacubitril with valsartan48 |
| Sagent 158 |
| SalAir231 |
| Salazopyrin8 |
| Salazopyrin EN8 |
| Salbutamol |
| Salbutamol with ipratropium |
| bromide |
| Salicylic acid |
| Salmeterol |
| Sandomigran |
| Sanofi Primaguine |
| Junon I Innaquino |

| Sapropterin dihydrochloride | 30 |
|-----------------------------------|-----------------|
| Scalp Preparations | 70 |
| Scopoderm TTS | |
| Sebizole | 70 |
| Secukinumab | |
| Sedatives and Hypnotics | 137 |
| Seebri Breezhaler | |
| Selegiline hydrochloride | 119 |
| Senna | 26 |
| Senokot | 26 |
| SensoCard | |
| Serc | |
| Serenace | |
| Seretide | 231 |
| Seretide Accuhaler | 231 |
| Serevent | 230 |
| Serevent Accuhaler | 230 |
| Sertraline | |
| Setrona | |
| Setrona AU | |
| Sevredol | |
| Sex Hormones Non | 120 |
| Contraceptive | 00 |
| Shield XL | 00 |
| | |
| shingles vaccine | 2// |
| SII-Onco-BCG | 1/6 |
| Sildenafil | 59 |
| Silhouette MMT-373 | 23 |
| Siltuximab | 215 |
| Simvastatin | |
| Simvastatin Mylan | 55 |
| Sinemet | 119 |
| Sinemet CR | 119 |
| Sirolimus | 224 |
| Siterone | 80 |
| Slow-Lopresor | 50 |
| Smith BioMed Rapid Pregnancy | |
| Test | |
| Sodibic | |
| Sodium acid phosphate | <mark>26</mark> |
| Sodium alginate | |
| Sodium benzoate | 30 |
| Sodium bicarbonate | |
| Blood | 45–46 |
| Extemporaneous | 245 |
| Sodium calcium edetate | 243 |
| Sodium chloride | |
| Blood | 45 |
| Respiratory | 235 |
| Sodium citrate with sodium lauryl | |
| sulphoacetate | 26 |
| Sodium citro-tartrate | |
| Sodium cromoglicate | |
| Alimentary | |
| Sensory | |
| Sodium fluoride | |
| | |

| Sodium Fusidate [fusidic acid] | |
|--------------------------------------|-------------|
| Dermatological | . 62 |
| Infection | . 95 |
| Sensory | 237 |
| Sodium hyaluronate [Hyaluronic acid] | |
| Sodium phenylbutyrate | 20 |
| Sodium picosulfate | 26 |
| Sodium polystyrene sulphonate | |
| Sodium tetradecyl sulphate | |
| Sodium valproate | .41 120 |
| Sofradex | 123 227 |
| Soframycin | |
| Solgar | 21 |
| Solifenacin Mylan | 77 |
| Solifenacin succinate | 77 |
| Solu-Cortef | 70 |
| Solu-Medrol | |
| Solu-Medrol-Act-O-Vial | .00. |
| Somatropin (Omnitrope) | |
| Sotalol | 51 |
| Spacer device | 235 |
| Span-K | |
| Spiolto Respimat | 232 |
| Spiractin | .53 |
| Spiriva | 232 |
| Spiriva Respimat | 232 |
| Spironolactone | 53 |
| Sporanox | . 97 |
| Sprycel | 159 |
| Staphlex | . 93 |
| Stemetil | 131 |
| SteroClear | 235 |
| Stesolid | 127 |
| Stimulants/ADHD Treatments | |
| Stiripentol | 129 |
| Stocrin | |
| Stomahesive | . 32 |
| Strattera | 138 |
| Strides Shasun | . 97 |
| Stromectol | . 66 |
| Sucralfate | |
| Sulfadiazine Silver | |
| Sulfadiazine sodium | .96 |
| Sulfasalazine | 8 |
| Sulindac | 111 |
| Sulphur | |
| Sulprix | 132 |
| Sumagran | 130 |
| Sumatriptan | |
| Sunitinib Pfizer | |
| Sunitiniti Prizer | |
| Sunscreens, proprietary | |
| Sure-T MMT-863 | 21 |
| Sure-T MMT-873 | . 2 I 21 |
| | ا ک |

| Sustagen Hospital Formula |
|--|
| Sustagen Hospital Formula |
| Active |
| Sustanon Ampoules81 |
| Sutent |
| Sylvant215 |
| Symbicort Turbuhaler 100/6 230 |
| Symbicort Turbuhaler 200/6 230 |
| Symbicort Turbuhaler 400/12 230 |
| Symmetrel 119 |
| Sympathomimetics |
| Synacthen80 |
| Synacthen Depot80 |
| Synacthene Retard80 |
| Synflorix |
| Synthroid83 |
| Syntometrine75 |
| Syrup (pharmaceutical grade) |
| Systane Unit Dose |
| -T- |
| Tacrolimus |
| Dermatological |
| Oncology |
| Tacrolimus Sandoz |
| Taliglucerase alfa |
| Tambocor |
| Tamoxifen citrate 169 |
| Tamoxifen Sandoz |
| T |
| l amsulosin hydrochloride |
| Tamsulosin hydrochloride76 Tamsulosin-Rex |
| Tamsulosin-Rex76 |
| Tamsulosin-Rex |

INDEX: Generic Chemicals and Brands

| 0 1 |
|---|
| Oncology157 |
| Trexate 150 |
| Triamcinolone acetonide |
| Alimentary |
| Annientaly |
| Dermatological64 |
| Hormone |
| Triamcinolone acetonide with |
| gramicidin, neomycin and nystatin |
| Dermatological |
| |
| Sensory |
| Triazolam138 |
| Trimethoprim96 |
| Trimethoprim with |
| sulphamethoxazole |
| |
| [Co-trimoxazole] |
| Trisequens82 |
| Trisul96 |
| Trophic Hormones83 |
| Tropicamide240 |
| |
| Trulicity12 |
| Trusopt239 |
| TruSteel |
| Tuberculin PPD [Mantoux] test277 |
| Tubersol |
| |
| Two Cal HN259 |
| Tykerb161 |
| Tysabri137 |
| |
| - 11 - |
| -U- |
| Ultibro Breezhaler |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural. 77 Urea. 65 Urex Forte 53 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urex Forte 53 Urinary Agents 76 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urex Forte 53 Urinary Agents 76 Urinary Tract Infections 109 |
| Ultibro Breezhaler |
| Ultibro Breezhaler |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urak Forte 53 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urany Agents 76 Urinary Agents 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Ursosan 24 Utrogestan 82 |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinorm 109 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 24 Urogestan 82 |
| Ultibro Breezhaler. 232 Ultraproct. 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinorm 109 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 24 Urogestan 82 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 82 - V - Vaccinations |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinorm 109 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 82 - V - Vaccinations 268 Vaclovir 101 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urex Forte 53 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Urogestan 82 - V - Vaccinations Valaciclovir 101 Valaciclovir 101 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urixer Forte 53 Urinary Agents 76 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursosan 24 Urogestan 82 - V - Vaccinations Valganciclovir 101 Valganciclovir 101 Valganciclovir Mylan 101 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinorm 117 Uromitexan 154 Ursosan 24 Utrogestan 82 - V - Vaccinations Valganciclovir 101 Valganciclovir 101 Valganciclovir 101 Vancomycin 96 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 Valaciclovir 101 Valagnciclovir Mylan 101 Vancomycin 96 Vannair 230 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 Valaciclovir 101 Valagnciclovir Mylan 101 Vancomycin 96 Vanair 230 Varenicline Pfizer 144 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 Valaganciclovir Mylan 101 Vanarir 230 Varenicline Pfizer 144 |
| Ultibro Breezhaler 232 Ultraproct 8 Umeclidinium 232 Umeclidinium with vilanterol 232 Univent 231, 235 Upadacitinib 226 Ural 77 Urea 65 Urinary Agents 76 Urinary Tract Infections 109 Urinorm 117 Uromitexan 154 Ursodeoxycholic acid 24 Utrogestan 82 - V - Vaccinations Valaciclovir 101 Valaciclovir 101 Valagnciclovir Mylan 101 Vancomycin 96 Vanair 230 Varenicline Pfizer 144 |

INDEX: Generic Chemicals and Brands

| Varicella zoster virus (Oka strain) live |
|--|
| attenuated vaccine [shingles |
| vaccine] 277 |
| Various |
| Varivax277 |
| Vasodilators57 |
| Vasopressin Agonists |
| Vasorex51 |
| Vedafil59 |
| Veletri60 |
| Venclexta157 |
| Venetoclax157 |
| Venlafaxine127 |
| Venomil228 |
| VENOX |
| Ventavis60 |
| Ventolin231 |
| Vepesid153 |
| Verapamil hydrochloride52 |
| Vergo 16131 |
| Vermox |
| Versacloz132 |
| Vesanoid157 |
| Vexazone11 |
| Vfend |
| Viaderm KC65 |
| Vidaza148 |
| Vigabatrin 129 |
| Vigisom137 |
| Vildagliptin11 |
| Vildagliptin with metformin |
| hydrochloride 12 |
| Vimpat 128 |
| Vinblastine sulphate 158 |
| Vincristine sulphate 158 |
| Vinorelbine158 |
| Vinorelbine Ebewe158 |
| Viramune Suspension 106 |
| ViruPOS237 |
| Vit.D3 |
| Vita-B12 |
| VitA-POS |
| Vitabdeck |
| Vital253 |
| Vitamin B complex |
| Vitamin B6 25 |
| Vitamins |
| Vivonex TEN253 |
| Voltaren |
| Voltaren D111 |
| Voltaren Ophtha |
| Voltaren SR 111 |
| Volumatic |
| Voriconazole |
| Votrient |
| Vttack98 |

| - W - |
|-------------------------------------|
| Warfarin sodium44 |
| Wart Preparations70 |
| Wasp venom allergy treatment228 |
| Water |
| Blood45 |
| Extemporaneous246 |
| Wool fat with mineral oil66 |
| - X - |
| Xarelto 44 |
| Xifaxan9 |
| XMET Maxamum261 |
| Xolair 199 |
| XP Maxamum262 |
| Xylocaine 121 |
| Xylocaine 2% Jelly 120 |
| Xyntha40 |
| - Z - |
| Zapril47 |
| Zarontin127 |
| Zaroxolyn53 |
| Zavedos153 |
| Zeffix101 |
| Zematop |
| Zetlam101 |
| Ziagen |
| Zidovudine [AZT] 106 |
| Zidovudine [AZT] with lamivudine |
| |
| Zimybe |
| Zinc and castor oil |
| Zinc sulphate |
| Zincaps |
| Ziprasidone |
| Zista |
| Zista |
| Zoledronic acid |
| Hormone |
| Musculoskeletal 115 |
| Zoledronic acid Mylan |
| Zopiclone |
| Zopiclone Actavis |
| Zostavax |
| Zostrix |
| Zostrix HP121 |
| Zuclopenthixol decanoate135 |
| Zuclopenthixol hydrochloride133 |
| Zusdone133 |
| Zyban143 |
| Zypine133 |
| Zypine ODT 133 |
| Zyprexa Relprevv 134 |
| Zytiga165 |











