### May 2022 Volume 29 Number 1

#### **Editors:**

Kaye Wilson, Doris Chong, & Sophie Molloy email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

PO Box 10 254 Wellington

### Circulation

Accessible in an electronic format at no cost from the Pharmac website <a href="https://www.pharmac.govt.nz/schedule">www.pharmac.govt.nz/schedule</a>.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

#### Production

Typeset automatically from XML and T<sub>E</sub>X. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

### **Programmers**

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be

liable for any consequences arising there from.

		L
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	61
	Genito Urinary System	71
	Hormone Preparations – Systemic	78
	Infections – Agents For Systemic Use	89
	Musculoskeletal System	110
	Nervous System	118
	Oncology Agents & Immunosuppressants	145
	Respiratory System & Allergies	227
	Sensory Organs	236
	Various	241
Section C	Extemporaneous Compounds (ECPs)	243
Section D	Special Foods	246
Section I	National Immunisation Schedule	267
	la day.	077

Introducing Pharmac

Index

### **Introducing Pharmac**

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

#### Pharmac's role:

"Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided."

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <a href="https://www.pharmac.govt.nz/about">https://www.pharmac.govt.nz/about</a>.

### **Purpose of the Pharmaceutical Schedule**

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply:
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that that can be used in DHB hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

### **Explaining pharmaceutical entries**

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

### Example



### Glossary

### **Units of Measure**

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

### SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price \$	) Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID  Sodium alginate 225 mg and magnesium alginate 87.5 mg sachet	•	30	<b>√</b> G	aviscon Infant
SODIUM ALGINATE  * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour		60	G	aviscon Double Strength
Oral liq 500 mg with sodium bicarbonate 267 mg and calcicarbonate 160 mg per 10 ml		500 ml	А	cidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE  * Tab 600 mg  CALCIUM CARBONATE	12.56	100	<b>✓</b> A	.lu-Tab
Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) — Subsidy by endorsement Only when prescribed for patients unable to swallow c inappropriate and the prescription is endorsed accordi	alcium carbonate table	500 ml ets or whe		loxane m carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE - Up to 30 cap available o  * Tab 2 mg*  * Cap 2 mg	10.75	400 400		lodia liamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE  Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy  SA1886 Special Authority for Subsidy nitial application — (Crohn's disease) from any relevant profile following criteria:  3oth:		90 valid for 6		intocort CIR or applications meeting

continued...

2 Any of the following:2.1 Diabetes; or2.2 Cushingoid habitus; or

1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and

2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation\*.

Note: Indication marked with \* is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis\*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
  - 3.1 Diabetes; or
  - 3.2 Cushingoid habitus; or
  - 3.3 Osteoporosis where there is significant risk of fracture; or
  - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
  - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
  - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
  - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
  - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with \* is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

#### HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	<ul><li>✓ Cortifoam S29</li><li>✓ Colifoam</li></ul>
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	✓	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mg	92.91	100	✓	Nalcrom
SULFASALAZINE	44.00	400		
* Tab 500 mg Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

### Local preparations for Anal and Rectal Disorders

### **Antihaemorrhoidal Preparations**

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g11.06	30 g OP	✓ Ultraproct		
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg7.30	12	✓ Ultraproct		
HYDROCORTISONE WITH CINCHOCAINE  Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00  Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	30 g OP 12	✓ Proctosedyl ✓ Proctosedyl		

### **Management of Anal Fissures**

### ⇒SA1329 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

### **Antispasmodics and Other Agents Altering Gut Motility**

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a			
PSO	65.45	10	Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	<ul><li>Buscopan</li></ul>
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

### **Antiulcerants**

### **Antisecretory and Cytoprotective**

MIS	SOPROSTOL STORE ST		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	<ul><li>Cytotec</li></ul>

	ALIMENTARY	TRACT ANI	D METABOLISM
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
Helicobacter Pylori Eradication			
CLARITHROMYCIN  Tab 500 mg — Subsidy by endorsement	eradication and prescri	iption is endorse	0,
H2 Antagonists			
FAMOTIDINE - Only on a prescription			

✓ Famotidine

Hovid S29

100

*	Tab 40 mg	8.48	100	✓ Famotidine  Hovid S29
*	Inj 10 mg per ml, 4 ml — Subsidy by endorsement Subsidy by endorsement — Subsidised for patients received			•
Р	roton Pump Inhibitors			

Subsidy by endorsement – Subsidised for patients receiving treatment as	part of pallia	itive care.
Proton Pump Inhibitors		
LANSOPRAZOLE  * Cap 15 mg	100 100	✓ <u>Lanzol Relief</u> ✓ <u>Lanzol Relief</u>
For omeprazole suspension refer Standard Formulae, page 243  * Cap 10 mg1.94	90	✓ Omeprazole actavis 10
* Cap 20 mg	90	✓ Omeprazole actavis 20
* Cap 40 mg	90	✓ Omeprazole actavis 40
Powder – Only in combination42.50 Only in extemporaneously compounded omeprazole suspension.	5 g	✓ Midwest
* Inj 40 mg ampoule with diluent	5	✓ <u>Dr Reddy's</u> <u>Omeprazole</u>
PANTOPRAZOLE         * Tab EC 20 mg       2.02         * Tab EC 40 mg       2.85	100 100	✓ Panzop Relief ✓ Panzop Relief
Site Protective Agents		
COLLOIDAL BISMUTH SUBCITRATE  Tab 120 mg14.51 SUCRALFATE	50	✓ Gastrodenol S29
Tab 1 g	120	Carafate
Bile and Liver Therapy		

RIFAXIMIN – Special Authority see SA1461 on the next page – Ret	ail pharmacy		
Tab 550 mg	625.00	56	✓ Xifaxan

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	lised	Generic	
\$	Per	1	Manufacturer	

### ⇒SA1461 Special Authority for Subsidy

**Initial application** only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

**Renewal** only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

### **Diabetes**

Hvp	eral	ycaemi	ic Aa	ents
,	~	,		•••••

DIAZOXIDE - Special Authority see SA1320 below -	- Retail pharmacy		
Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral lig 50 mg per ml	620.00	30 ml OP	✓ Proglycem S29

### **⇒SA1320** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

#### GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit -	Up to 5 kit available on a PSO	32.00	1	Glucagen Hypokit
------------------------	--------------------------------	-------	---	------------------

### **Insulin - Short-acting Preparations**

INS	SULIN NEUTRAL		
•	Inj human 100 u per ml25.20	6 10 ml OP	<ul><li>✓ Actrapid</li><li>✓ Humulin R</li></ul>
<b>A</b>	Inj human 100 u per ml, 3 ml	5 5	✓ Actrapid Penfill ✓ Humulin B

# Insulin - Intermediate-acting Preparations INSULIN ASPART WITH INSULIN ASPART PROTAMINE

▲ Inj 100 iu per ml, 3 ml prefilled pen52.15	5	✓ NovoMix 30 FlexPen
INSULIN ISOPHANE		
▲ Inj human 100 u per ml17.68	10 ml OP	✓ Humulin NPH
		✓ Protaphane
▲ Inj human 100 u per ml, 3 ml29.86	5	✓ Humulin NPH
		Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL		
▲ Inj human with neutral insulin 100 u per ml25.26	10 ml OP	✓ Humulin 30/70
,		✓ Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml42.66	5	✓ Humulin 30/70
•		✓ PenMix 30
		✓ PenMix 40
		✓ PenMix 50

(PenMix 40 Inj human with neutral insulin 100 u per ml, 3 ml to be delisted 1 December 2022)

Sole Subsidised Supply

	Subsidy (Manufacturer's Price	e) Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,	,			
3 ml		5	✓	Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,	ı		_	
3 ml	42.66	5	<b>✓</b>	Humalog Mix 50
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
Inj 100 u per ml, 10 ml	63.00	1	✓	Lantus
Inj 100 u per ml, 3 ml		5	✓	Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓	Lantus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
Inj 100 u per ml, 10 ml	30.03	1	<b>✓</b>	NovoRapid
Inj 100 u per ml, 3 ml		5		NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	<b>✓</b>	NovoRapid FlexPen
NSULIN GLULISINE				
Inj 100 u per ml, 10 ml	27.03	1	1	Apidra
Inj 100 u per ml, 3 ml	46.07	5	1	Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5		Apidra SoloStar
NSULIN LISPRO				
Inj 100 u per ml, 10 ml		10 ml OP	✓	Humalog
Inj 100 u per ml, 3 ml	59.52	5	<b>√</b>	Humalog
Alpha Glucosidase Inhibitors				
CARBOSE				
★ Tab 50 mg	8.95	90	1	Accarb
₭ Tab 100 mg	15.29	90	1	<u>Accarb</u>
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
₹ Tab 5 mg	7.50	100	<b>✓</b>	Daonil
GLICLAZIDE			-	<del></del>
₹ Tab 80 mg	15.18	500	1	Glizide
GLIPIZIDE		*	-	
₭ Tab 5 mg	4.58	100	<b>✓</b> I	Minidiab
METFORMIN HYDROCHLORIDE				
F Tab immediate-release 500 mg	14 74	1.000	<b>√</b> 1	Metformin Mylan
★ Tab immediate-release 500 mg		500		Metformin Mylan
PIOGLITAZONE				
Tab 15 mg	6.80	90	1	Vexazone
· · · · · · · · · · · · · · · · · · ·		90	1	Vexazone Vexazone
€ Tab 30 mg				
≮ Tab 30 mg ≮ Tab 45 mg	12.25	90		Vexazone
· · · · · · · · · · · · · · · · · · ·	12.25	90		Vexazone

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE  Tab 50 mg with 1,000 mg metformin hydrochloride  Tab 50 mg with 850 mg metformin hydrochloride		60 60	-	Galvumet Galvumet

### **GLP-1 Agonists**

### ⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or
    - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
    - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
    - 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 above - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

#### SGLT2 Inhibitors

### ⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
  - 2.1 Patient has type 2 diabetes; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is Māori or any Pacific ethnicity\*; or
    - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)\*; or

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	<b>✓</b>	Manufacturer

#### continued...

- 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator\*; or
- 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult\*; or
- 2.2.5 Patient has diabetic kidney disease (see note b)\*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: \* Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN – Special Authority see SA2068 on the previous page – Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg	58.56	30	<ul><li>Jardiance</li></ul>
*	Tab 25 mg	58.56	30	<ul><li>Jardiance</li></ul>

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	<ul><li>Jardiamet</li></ul>
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	✓ Jardiamet

### **Diabetes Management**

### **Ketone Testing**

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes; or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### **Dual Blood Glucose and Blood Ketone Testing**

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
  - 1) type 1 diabetes; or
  - 2) permanent neonatal diabetes: or
  - 3) undergone a pancreatectomy; or
  - 4) cystic fibrosis-related diabetes; or
  - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.......20.00 1 OP ✓ CareSens Dual

### **Blood Glucose Testing**

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
  - 1) is receiving insulin or sulphonylurea therapy; or
  - 2) is pregnant with diabetes; or
  - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
  - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

rips.......10.00 1 OP ✓ <u>CareSens N</u>

✓ CareSens N POP

20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
, , , , , , , , , , , , , , , , , , , ,	Per 🗸	Manufacturer

### BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	•	CareSens N
		1	CareSens PRO

#### BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
  prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	.20 50 te	st OP	SensoCard
---------------------------	-----------	-------	-----------

### Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

### INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10.95	100	✓ B-D Micro-Fine
	31 g × 5 mm12.26	100	✓ B-D Micro-Fine
	31 g × 6 mm9.50	100	✓ Berpu
	31 g × 8 mm	100	✓ B-D Micro-Fine
	32 g x 4 mm 10.95	100	✓ B-D Micro-Fine

		Subsidy		Fully	Brand or
		(Manufacturer's Price)		Subsidised	Generic
		\$	Per		Manufacturer
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE	E – Maximum of 200	dev p	oer prescrip	otion
*	Syringe 0.3 ml with 29 g × 12.7 mm needle	13.56	100	1	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.30	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
		1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
	, ,	1.36	10		
		(1.99)			B-D Ultra Fine
*	Syringe 1 ml with 31 g x 8 mm needle	13.56	100	✓	B-D Ultra Fine II
		1.36	10		
		(1.99)			B-D Ultra Fine II

### **Insulin Pumps**

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

### ⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic	
	Per	•	Manufacturer	

continued...

- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Fither:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (HbA1c)** only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
  - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 3.2 The pump is due for replacement; and
- 4 Either:
  - 4.1 Applicant is a relevant specialist; or
  - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 8.2 The pump is due for replacement; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
  - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
  - 4.2 The pump is due for replacement; and
- 5 Either:
  - 5.1 Applicant is a relevant specialist; or
  - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

## Insulin Pump Consumables

### ⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
  - 6.1 Applicant is a relevant specialist; or
  - 6.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal — (permanent neonatal diabetes)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
  - 8.1 Applicant is a relevant specialist: or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
  - 3.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
  - 9.1 Applicant is a relevant specialist; or
  - 9.2 Applicant is a nurse practitioner working within their vocational scope.

**Renewal** — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
  - 3.1 Applicant is a relevant specialist; or
  - 3.2 Applicant is a nurse practitioner working within their vocational scope.

**Initial application** — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
  - 8.1 Applicant is a relevant specialist; or
  - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

✓ TruSteel

	Subsidy (Manufacturer's Price) \$	Subsid Per	Fully ised ✓	Brand or Generic Manufacturer
continued than 80 mmol/mol; and The patient's HbA1c has not deteriorated more than 5 mm The patient has not had an increase in severe unexplained Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their	d hypoglycaemic epis			ne; and
INSULIN PUMP CARTRIDGE — Special Authority see SA1985 or     a) Maximum of 3 sets per prescription     b) Only on a prescription     c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U. t:lock × 10	year.	narmacy	<b>✓</b> T	andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special A  a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	Authority see SA1985		9 – Re	etail pharmacy
10 mm steel needle; 60 cm tubing × 10		1 OP 1 OP		liniMed Sure-T MMT-884A liniMed Sure-T
6 mm steel needle; 60 cm tubing × 10		1 OP	✓ M	MMT-886A liniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10		1 OP 1 OP		liniMed Sure-T MMT-866A liniMed Sure-T
8 mm steel needle; 80 cm tubing × 10		1 OP		MMT-874A liniMed Sure-T MMT-876A
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	<b>√</b> S	ure-T MMT-863
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock		1 OP	_	ure-T MMT-873
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT Retail pharmacy  a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles		cial Authorii 1 OP		SA1985 on page 19 –
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	130.00	1 OP	<b>✓</b> T	ruSteel
10 needles	130.00	1 OP	<b>✓</b> T	ruSteel

8 mm steel cannula; straight insertion; 60 cm line × 10 with

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

\$ Per 

Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette  MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	MMT-384A  ✓ MiniMed Quick-Set
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	MMT-398A  ✓ MiniMed Mio
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	MMT-941A  ✓ MiniMed Mio
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	MMT-921A  ✓ MiniMed Mio
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	MMT-943A ✓ MiniMed Mio
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	MMT-923A ✓ MiniMed Quick-Set
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	MMT-399A ✓ MiniMed Mio
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	MMT-945A ✓ MiniMed Mio
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	MMT-965A ✓ MiniMed Mio
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	MMT-925A ✓ MiniMed Quick-Set
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	MMT-387A ✓ MiniMed Quick-Set
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	MMT-396A  ✓ MiniMed Quick-Set
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	MMT-397A ✓ MiniMed Mio
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	MMT-975A ✓ MiniMed Quick-Set
			MMT-386A

	Subsidy (Manufacturer's Price) \$	) Sub Per	Fully osidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH IN	SERTION	DEVICE	E) - Special Authority see
SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c	:m			
line x 10 with 10 needles		1 OP	✓ Aı	utoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cn line x 10 with 10 needles		1 OP	✓ A	utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SERTION) - Spec	cial Author	ity see S/	A1985 on page 19 –
Retail pharmacy	, ,		•	
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	120.00	1 OP	√ Si	ilhouette MMT-373
•		-		
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH see SA1985 on page 19 – Retail pharmacy	I INSERTION WITH	H INSEKI	ION DEV	/ICE) - Special Authority
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles	140.00	1 OP	✓ Aı	utoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cr				
line × 10 with 10 needles	140.00	1 OP	✓ Ai	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device;	140.00	4 OD		
110 cm line × 10 with 10 needles		1 OP	<b>♥</b> Al	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr		1 OP	✓ A	utoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH				
Retail pharmacy	I INSERTION) - 3	ppeciai Au	Honly Se	e 3A 1905 on page 19 -
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing $\times$ 10 with	h			
10 needles; luer lock		1 OP	✓ Q	uick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with		4.00		
10 needles; luer lock		1 OP	<b>₽</b> Q	uick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 or	n page 19 – Retail p	harmacy		
A) Maximum of 3 sets per prescription				
<ul><li>b) Only on a prescription</li><li>c) Maximum of 13 packs of reservoir sets will be funded per</li></ul>	voor			
10 x luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP	<b>✓</b> Δ1	DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		iniMed
1 17				1.8 Reservoir
				MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		iniMed
				3.0 Reservoir
				MMT-332A

Subsidy (Manufacturer's Price)		Fully ubsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

### **Digestives Including Enzymes**

Р	Δ	١	J	CI	R	F	Δ	Т	IC.	F	N	7	1	И	F

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Creon 10000 to be Principal Supply on 1 June 2022			
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase,			
1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase			
25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Creon 25000 to be Principal Supply on 1 June 2022			
Modified release granules pancreatin 60.12 mg (amylase			
3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	✓ Creon Micro
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amy	lase, 1,250 U p	protease)) to	be delisted 1 June 2023)
URSODEOXYCHOLIC ACID - Special Authority see SA1739 below	v – Retail pharı	macy	
Cap 250 mg		100	✓ <u>Ursosan</u>

### ⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury: and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

**Initial application** — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total

(M.	Subsidy		Fully	Brand or
	(Manufacturer's Price)		ubsidised	Generic
	\$	Per	1	Manufacturer

continued...

Lavatives

Parenteral Nutrition (TPN); and

2 Liver function has not improved with modifying the TPN composition.

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives		
Bulk-forming Agents		
ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription		
* Powder for oral soln	250 g OP	<ul><li>Macro Organic</li><li>Psyllium Husk</li></ul>
12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS		
* Dry	500 g OP	N 181
(17.32) 2.41	200 g OP	Normacol Plus
(8.72)	200 y OP	Normacol Plus
Faecal Softeners		
DOCUSATE SODIUM – Only on a prescription	100	<b>Colored</b>
* Tab 50 mg2.31 * Tab 120 mg3.13	100 100	✓ <u>Coloxyl</u> ✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES	100	- OJIONYI
* Tab 50 mg with sennosides 8 mg4.20	200	✓ Laxsol
POLOXAMER - Only on a prescription	_00	
Not funded for use in the ear.	00 I OD	✓ Oalanul
* Oral drops 10%	30 ml OP	✓ <u>Coloxyl</u>
Opioid Receptor Antagonists - Peripheral		
METHYLNALTREXONE BROMIDE - Special Authority see SA1691 below - Re	tail pharmacy	

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal

246.00

continued...

✓ Relistor

✓ Relistor

7

Inj 12 mg per 0.6 ml vial ......36.00

Subsidy (Manufacturer's Price)			Brand or Generic
 \$	Per	✓	Manufacturer

continued...

unless notified for applications meeting the following criteria:

#### Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
  - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
  - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

### **Osmotic Laxatives**

GLYCEROL			
* Suppos 3.6 g - Only on a prescription	9.25	20	✓ PSM
LACTULOSE - Only on a prescription			
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ <u>Laevolac</u>
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICAL	RBONATE A	ND SODIUM C	CHLORIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 mg,			
sodium bicarbonate 178.5 mg and sodium chloride 350.7 m	g6.70	30	✓ <u>Molaxole</u>
SODIUM ACID PHOSPHATE - Only on a prescription			
Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate
			Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE -	Only on a pro	escription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml,			
5 ml	29.98	50	✓ <u>Micolette</u>
Chimoslant I assetissa			
Stimulant Laxatives			
BISACODYL - Only on a prescription			
* Tab 5 mg	5.80	200	✓ Pharmacy Health
	5.99		✓ Lax-Tab
Pharmacy Health to be Principal Supply on 1 June 2022	0.00	40	41 0 "
* Suppos 10 mg	3.69	10	✓ <u>Lax-Suppositories</u>
(Lax-Tab Tab 5 mg to be delisted 1 June 2022)			
SENNA – Only on a prescription  * Tab, standardised	0.17	100	
* Tab, standardised	(8.21)	100	Senokot
	0.43	20	Jenokot
	(2.06)	_0	Senokot
SODIUM PICOSULFATE - Special Authority see \$A2053 below - F	, ,	acv	
Oral soln 7.5 mg per ml		30 ml OP	✓ Dulcolax SP Drop
			•

### ⇒SA2053 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

### **Metabolic Disorder Agents**

ALGLUCOSIDASE ALFA - Special Authority see SA1986 below - Retail pharmacy
Inj 50 mg vial ......1,142.60 1 ✓ Myozyme

### ⇒SA1986 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
  - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells: or
  - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
  - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
  - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authority see SA2042 below – Retail pharmacy

Tab 1,000 mg	CBS	90	<ul><li>Clinicians</li></ul>
Cap 500 mg		50	✓ Solgar
Powder		400 a	✓ Biomed

#### ⇒SA2042 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA1987 on the next page - Retail pharmacy

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### ⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
  - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
  - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
  - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - S	pecial Authority see SA2039 below – Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

### ⇒SA2039 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE − Special Authority see SA1988 below − Retail pharmacy
Inj 1 mg per ml, 5 ml vial......2,234.00 1 ✓ Naglazyme

### ⇒SA1988 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

**Renewal** only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### ⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts; or
  - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene, and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – Special Authority see SA1695 below – Retail pharmacy
Inj 100 U per ml, 5 ml vial......1,335.16

✓ Aldurazyme

### ⇒SA1695 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
  - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
  - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

### ⇒SA2040 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the next page - Retail pharmacy
Tab 100 mg .......CBS 100
Cap 100 mg ......CBS 100
✓ Country Life
✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

### **⇒SA2041** Special Authority for Subsidy

**Initial application** only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

#### Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

### **⇒SA1989** Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

**Renewal** only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Either:
  - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
  - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
  - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
  - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
  - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy
Soln 100 mg per ml ......CBS

SODIUM BENZOATE – Special Authority see SA1599 below – Retail pharmacy

#### ⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ Pheburane

100 ml

✓ Amzoate S29

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

### ⇒SA1990 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

**Renewal** only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	<ul> <li>Life Extension</li> </ul>

### ⇒SA2043 Special Authority for Subsidy

**Initial application** only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

**Renewal** only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### Gaucher's Disease

### ⇒SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990 Pharmac PO Box 10 254 Facsimile: 04 916 7571

Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

#### **Access Criteria**

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- The patient has a diagnosis of symptomatic type 1 or type 3\* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

massive symptomatic splenomegaly; or

- 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
- 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
- Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
- 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

#### \*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

### Mouth and Throat

### **Agents Used in Mouth Ulceration**

#### BENZYDAMINE HYDROCHLORIDE

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

#### CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste	17.20	56 g OP	Stomahesive
	4.55	15 g OP	
	(7.90)	-	Orabase
	1.52	5 g OP	
	(3.60)	•	Orabase
Powder	8.48	28 g OP	
	(10.95)	-	Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
-	(6.00)	-	Bonjela

	Subsidy		Fully Brand or
(	(Manufacturer's P \$	rice) Subs Per	idised Generic  ✓ Manufacturer
TRIAMCINOLONE ACETONIDE	Ψ		That land tall of
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
1 4010 011 /0			- Itonaiog in Orabaoo
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN	4.70	04 100	<b>4</b> NIII
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN			
★ Inj 1 mg per ml, 1 ml ampoule -Up to 6 inj available on a PS0	D1.89	3	✓ Vita-B12
	2.84	•	✓ Neo-B12
	3.15	5	✓ Hydroxocobalamin
			Mercury Pharma
PYRIDOXINE HYDROCHLORIDE			
a) No more than 100 mg per dose			
b) Only on a prescription  * Tab 25 mg - No patient co-payment payable	2.70	90	✓ Vitamin B6 25
Tab 50 mg		500	✓ Pyridoxine
1 ab 00 mg		000	multichem
THIAMINE HYDROCHLORIDE - Only on a prescription			
* Tab 50 mg	7.09	100	✓ Max Health
VITAMIN B COMPLEX			
* Tab, strong, BPC	7.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID			
a) No more than 100 mg per dose			
b) Only on a prescription  * Tab 100 mg	9 90	500	✓ Cvite
		000	- <u>01110</u>
Vitamin D			
ALFACALCIDOL			
* Cap 0.25 mcg		100	✓ One-Alpha
* Cap 1 mcg		100	✓ One-Alpha
* Oral drops 2 mcg per ml	60.68	20 ml OP	✓ One-Alpha
CALCITRIOL	7.05	400	Contained AFT
* Cap 0.25 mcg		100	✓ Calcitriol-AFT ✓ Calcitriol-AFT
* Cap 0.5 mcg	13./5	100	✓ <u>Calcitriol-AFT</u>
COLECALCIFEROL * Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescriptic	n 295	12	✓ Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)		4.8 ml OP	✓ Puria

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

### **Multivitamin Preparations**

MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pharmacy

### ⇒SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Fither:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).</p>

MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy

### ⇒SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

#### **VITAMINS**

★ Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy .......23.40 60 ✓ Vitabdeck

### ⇒SA1720 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

### **Minerals**

### Calcium

#### CALCIUM CARBONATE

*	Tab 1.25 g (500 mg elemental)	6.69	250	Caici-Tab 500
*	Tab off 1 25 g (500 mg elemental) - Subsidy by endorsement	54.60	76	✓ Cacit S29

Tableff 1.25 g (500 mg elemental) − Subsidy by endorsement.......54.60 76 Cacit S29 260.00 100 Calcium 500 mg

Hexal S29

Subsidy by endorsement – Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable.

(Cacit S29) Tab eff 1.25 g (500 mg elemental) to be delisted 1 July 2022)

#### CALCIUM GLUCONATE

64.00 20 **Max Health** \$29

	Subsidy (Manufacturer's Price) \$	Sı Per	Fully ubsidised	Brand or Generic Manufacturer
CALCIUM LACTATE GLUCONATE WITH CALCIUM CARBONA Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg element  - Subsidy by endorsement	tal)	20	<b>√</b> 0	alcium-Sandoz
Cabbay by chaorotheric	260.00	100	_	Forte \$29 Calcium-Sandoz
				Forte S29

Subsidy by endorsement – Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable.

(Calcium-Sandoz Forte 29 Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg elemental) to be delisted 1 July 2022) (Calcium-Sandoz Forte 29 Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg elemental) to be delisted 1 July 2022)

Fluoride		
SODIUM FLUORIDE  * Tab 1.1 mg (0.5 mg elemental)	100	✓ PSM
lodine		
POTASSIUM IODATE  * Tab 253 mcg (150 mcg elemental iodine)4.58	90	✓ <u>NeuroTabs</u>
Iron		
FERROUS FUMARATE  * Tab 200 mg (65 mg elemental)	100	✓ <u>Ferro-tab</u>
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	100	✓ Ferro-F-Tabs
# Tab long-acting 325 mg (105 mg elemental)	30 500 ml	<ul><li>✓ Ferrograd</li><li>✓ Ferodan</li></ul>
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority see SA1840 below Inj 50 mg per ml, 10 ml vial	w – Retail phari 1	macy <b>✓ Ferinject</b>

### SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist,

Subsidy	0.1	Fully	Brand or	
(Manufacturer's Price) \$	Per	sidised •	Generic Manufacturer	

continued...

anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
  - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
  - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
  - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
  - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

### 

Magnesium		
MAGNESIUM HYDROXIDE Suspension 8%33.60	355 ml	✓ Phillips Milk of Magnesia S29
MAGNESIUM SULPHATE  * Inj 2 mmol per ml, 5 ml ampoule25.53	10	✓ Martindale
Zinc		
ZINC SULPHATE  * Cap 137 4 mg (50 mg elemental) 11 00	100	✓ 7incans

✓ Ferrosia

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

## Antianaemics

## Hypoplastic and Haemolytic

### ⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
  - 3.1 Both:
    - 3.1.1 Patient does not have diabetes mellitus: and
    - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
  - 3.2 Both:
    - 3.2.1 Patient has diabetes mellitus: and
    - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
  - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)\*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

**Renewal — (chronic renal failure)** from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with \* is an unapproved indication

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
EPOETIN ALFA - Special Authority see SA1775 on the previous	s page – Retail pharm	асу		
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	✓	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	Binocrit
Megaloblastic				
FOLIC ACID				

## Antifibrinolytics, Haemostatics and Local Sclerosants

#### EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

1.000

100

25 ml OP

✓ Folic Acid multichem

✓ Folic Acid Mylan ✓ Riomed

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial		1	✓ Alprolix
Inj 2,000 iu vial		1	✓ Alprolix
Inj 3,000 iu vial		1	✓ Alprolix
ELTROMBOPAG - Special Authority see SA1743 be	elow – Retail pharmacy		

# Wastage claimable

Tractage claimable			
Tab 25 mg	1.550.00	28	Revolade
Tab 50 mg		28	✓ Revolade

#### ⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
  - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
  - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
  - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

continued...

Subsidy	Subsid	-ully	Brand or
(Manufacturer's Price)		ised	Generic
\$	Per	1	Manufacturer

continued...

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Fither:
  - 3.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microliter: or
  - 3.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
  - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
  - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

#### EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

✓ Hemlibra	1	3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	7,138.00	Inj 60 mg in 0.4 ml vial
✓ Hemlibra	1	12,492.00	Inj 105 mg in 0.7 ml vial
✓ Hemlibra	1	17.846.00	lni 150 mg in 1 ml vial

⇒SA1969 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 2 Either:
  - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
  - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more: and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
  - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
  - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

**Renewal** only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

### EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe		1	✓ NovoSeven RT
Inj 5 mg syringe		1	✓ NovoSeven RT
Inj 8 mg syringe		1	✓ NovoSeven RT

### FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	5.00 1	FEIBA NF
Inj 1,000 U2,630	0.00 1	✓ FEIBA NF
Inj 2,500 U	5.00 1	✓ FEIBA NF

#### MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	✓ Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

#### NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	0 1	✓ RIXUBIS
Inj 1,000 iu vial870.0	0 1	✓ RIXUBIS
Inj 2,000 iu vial	0 1	✓ RIXUBIS
Inj 3,000 iu vial2,610.0	0 1	✓ RIXUBIS

	0.1		- ·	
	Subsidy (Manufacturer's Price)	Su	Fully bsidised	
	\$	Per	<b>√</b>	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	[Xpharm]			
For patients with haemophilia. Preferred Brand of short half	-life recombinant factor	r VIII. A	ccess to	o funded treatment is
managed by the Haemophilia Treaters Group in conjunction				
Inj 250 iu vial		1		Advate
Inj 500 iu vial	420.00	1	✓	Advate
Inj 1,000 iu vial	840.00	1	✓	Advate
Inj 1,500 iu vial	1,260.00	1	✓	Advate
Inj 2,000 iu vial		1		Advate
Inj 3,000 iu vial	,	1	/	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances				
treatment is managed by the Haemophilia Treaters Group in	conjunction with the I	Vational	Haemo	philia Management Group,
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Kogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Kogenate FS
Inj 3,000 iu vial		1	•	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatm		d treatm	ent is n	nanaged by the Haemophilia
Treaters Group in conjunction with the National Haemophilia			_	
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial	· ·	1		Adynovate
Inj 2,000 iu vial	2,400.00	1	•	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(73.00)			Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓	Mercury Pharma
Vitamin K				
PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO	8.00	5	1	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Konakion MM
ing to mg por mi, i mi op to o mg available on a room		Ŭ		Tronauton iiiii
Antithrombotic Agents				
- Tillian Omsons rigonic				
Antiplatelet Agents				
ASPIRIN				
* Tab 100 mg	10.80	990	•	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	1	Clopidogrel
				<u>Multichem</u>
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	1	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next pag				-
* Tab 90 mg		56	1	Brilinta

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
\$	Per	✓	Manufacturer

### ⇒SA1955 Special Authority for Subsidy

**Initial application** — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient has had a neurological stenting procedure\* in the last 60 days; or
  - 1.2 Patient is about to have a neurological stenting procedure performed\*; and
- 2 Either:
  - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
  - 2.2 Either:
    - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
    - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

**Initial application** — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic\*\*.

Notes: indications marked with \* are unapproved indications.

Note: \*\* Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	(Manufacturer's Price)	Subsi Per	dised	Generic Manufacturer	
Hamanin and Antononiat Dromonations					

## **Heparin and Antagonist Preparations**

•			
ENOXAPARIN SODIUM - Special Authority see SA1646	below – Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	<ul><li>Clexane</li></ul>
Inj 60 mg in 0.6 ml syringe	60.67	10	<ul><li>Clexane</li></ul>
Inj 80 mg in 0.8 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 100 mg in 1 ml syringe		10	<ul><li>Clexane</li></ul>
Inj 120 mg in 0.8 ml syringe		10	Clexane Forte
Inj 150 mg in 1 ml syringe		10	Clexane Forte

#### ⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

#### HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule	72.84	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml	32.66	5	DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	289.05	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓ Hospira
	42.40		✓ Heparin DBL S29
	482.20	50	✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg	76.36	60	1	Pradaxa
Cap 150 mg	76.36	60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	1	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	77.56	28	1	Xarelto
Tab 20 mg	77.56	28	1	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
	6.46	100		Marevan
* Tab 2 mg		50	_	Coumadin
* Tab 3 mg		100		Marevan
* Tab 5 mg		50		Coumadin
	11.48	100	•	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail p	harmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	1	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓	Nivestim

#### ⇒SA1259 Special Authority for Subsidy

**Initial application** only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%\*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10<sup>9</sup>/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC  $< 0.5 \times 10^9$ /L).

Note: \*Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

#### ⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%\*). Note: \*Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	BLOOD AN	D BLOOD	FUN	WIING ONGANS
	Subsidy (Manufacturer's Pric	ce) Subs Per	Fully idised	Brand or Generic Manufacturer
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE]  * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO  * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO  POTASSIUM CHLORIDE		5 1	_	Biomed Biomed
* Inj 75 mg per ml, 10 ml	65.00	50	<b>√</b> J	uno
SODIUM BICARBONATE Inj 8.4%, 50 ml	21.40	1	<b>✓</b> E	Biomed
b) Not in combination Inj 8.4%, 100 ml	21.95	1	<b>✓</b> E	Biomed
SODIUM CHLORIDE  Not funded for use as a nasal drop. Not funded for nebulise for nebuliser use.	er use except when	used in conju	ınction	with an antibiotic intende
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.26	500 ml 1,000 ml	<b>✓</b> E	Baxter Baxter
Only if prescribed on a prescription for renal dialysis, ma for emergency use. (500 ml and 1,000 ml packs)	aternity or post-nata	al care in the	home o	of the patient, or on a PSC
Inj 23.4% (4 mmol/ml), 20 ml ampoule		5 243	<b>✓</b> E	Biomed
Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO		20	<b>√</b> F	resenius Kabi
Inj 0.9%, 10 ml ampoule — Up to 5 inj available on a PSO Inj 0.9%, 20 ml ampoule	5.40	50 20	_	resenius Kabi resenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)		1 OP	- ✓ T	
WATER		TOF	• 1	riv
<ol> <li>On a prescription or Practitioner's Supply Order only w Schedule requiring a solvent or diluent; or</li> <li>On a bulk supply order; or</li> <li>When used in the extemporaneous compounding of ey</li> <li>When used for the dilution of sodium chloride soln 7%</li> </ol>	ye drops; or	·	ection li	sted in the Pharmaceutic
Inj 10 ml ampoule – Up to 5 inj available on a PSOInj 20 ml ampoule – Up to 5 inj available on a PSO		50 20	<b>√</b> F	Pfizer Fresenius Kabi Multichem
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	<b>√</b> (	Calcium Resonium
COMPOUND ELECTROLYTES  Powder for oral soln - Up to 5 sach available on a PSO	9.77	50	<b>√</b> E	:lectral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]		000 ml OB	<b>√</b> □	andialuta -

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

1,000 ml OP

✓ Pedialyte -Bubblegum

	Subsidy (Manufacturer's Price \$	) Per	Fully Subsidised	Brand or Generic Manufacturer
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	<b>✓</b> F	Phosphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60		
	(11.85)			Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90	200	<b>√</b> 9	Span-K
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	<b>√</b> 9	Sodibic
			<b>√</b> 9	Sodibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84 65	154 a C	)P 🗸 F	Resonium-A

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	/	Manufacturer

Almba Advanasantas Dissis			
Alpha Adrenoceptor Blocke	ers		
OXAZOSIN			
Fab 2 mg	17.35	500	✓ Apo-Doxazosin
- Tab 4 mg	20.94	500	✓ Doxazosin Clinect ✓ Apo-Doxazosin
Apo-Doxazosin Tab 2 mg to be delisi Apo-Doxazosin Tab 4 mg to be delisi	. ,		✓ Doxazosin Clinect
HENOXYBENZAMINE HYDROCHL	ORIDE		
Cap 10 mg	65.00	30	✓ BNM S29
	216.67	100	✓ Dibenzyline S29
RAZOSIN			
Tab 1 mg	5.53	100	✓ Arrotex-Prazosin S29 S29
Tab 2 mg	7.00	100	✓ Arrotex-Prazosin
3			<b>S29</b> S29
Tab 5 mg	11.70	100	✓ Arrotex-Prazosin S29 S29
ACF Inhibitors			
APTOPRIL	94 99	95 ml OP	✓ Canoten
		95 ml OP	✓ Capoten
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre	en under 12 years of age.	95 ml OP	✓ Capoten
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL – Subsidy by endorsem	en under 12 years of age. ent		
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL – Subsidy by endorsem Subsidy by endorsement – Subsi	en under 12 years of age. ent dised for patients who were taking cilazap.	ril prior to 1 May	2021 and the prescription
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL – Subsidy by endorsem Subsidy by endorsement – Subsi	en under 12 years of age. ent	ril prior to 1 May	2021 and the prescription
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL – Subsidy by endorsem Subsidy by endorsement – Subsi endorsed accordingly. Pharmacidispensing of cilazapril.	en under 12 years of age. ent dised for patients who were taking cilazap.	ril prior to 1 May	2021 and the prescription
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL – Subsidy by endorsem Subsidy by endorsement – Subsi endorsed accordingly. Pharmacidispensing of cilazapril. Tab 0.5 mg	en under 12 years of age.  ent dised for patients who were taking cilazap sts may annotate the prescription as endo	ril prior to 1 May rsed where there	2021 and the prescription e exists a record of prior
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childred to chil	en under 12 years of age.  ent dised for patients who were taking cilazap sts may annotate the prescription as endo	ril prior to 1 May rsed where there	2021 and the prescription exists a record of prior
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childred to chil	en under 12 years of age.  ent dised for patients who were taking cilazap sts may annotate the prescription as endo2.094.80	ril prior to 1 May rsed where there 90 90	2021 and the prescription exists a record of prior  Zapril Zapril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childred to chi	en under 12 years of age.  ent dised for patients who were taking cilazap sts may annotate the prescription as endo  2.09 4.80 8.35	ril prior to 1 May rsed where there 90 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Zapril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL — Subsidy by endorsem Subsidy by endorsement — Subsi endorsed accordingly. Pharmaci dispensing of cilazapril. Tab 0.5 mg Tab 2.5 mg Tab 5 mg	en under 12 years of age. In the dised for patients who were taking cilazapests may annotate the prescription as endo  2.09 4.80 8.35	ril prior to 1 May rsed where there 90 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre LAZAPRIL — Subsidy by endorsement — Subsidy by endorsement — Subsidendorsed accordingly. Pharmacic dispensing of cilazapril. Tab 0.5 mg Tab 2.5 mg Tab 5 mg VALAPRIL MALEATE Tab 5 mg Tab 10 mg	en under 12 years of age.  ent dised for patients who were taking cilazap sts may annotate the prescription as endo  2.09 4.80 8.35  1.82	ril prior to 1 May rsed where there 90 90 90 90	2021 and the prescription exists a record of prior Zapril Zapril Zapril Acetec Acetec
APTOPRIL Oral liq 5 mg per ml Oral liq 5 mg per ml	en under 12 years of age. In the dised for patients who were taking cilazapests may annotate the prescription as endo  2.09 4.80 8.35	ril prior to 1 May rsed where there 90 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec
APTOPRIL  Oral liq 5 mg per ml Oral liquid restricted to childre ilLAZAPRIL — Subsidy by endorsem Subsidy by endorsem — Subsidy by endorsed accordingly. Pharmacidispensing of cilazapril.  Tab 0.5 mg Tab 2.5 mg Tab 5 mg  NALAPRIL MALEATE  Tab 5 mg	en under 12 years of age. Itent dised for patients who were taking cilazap sts may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42	ril prior to 1 May rsed where there 90 90 90 100 100 100	2021 and the prescription e exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec
APTOPRIL  Oral liq 5 mg per ml  Oral liquid restricted to childre ilLAZAPRIL — Subsidy by endorsem Subsidy by endorsement — Subsidy by endorsed accordingly. Pharmacidispensing of cilazapril.  Tab 0.5 mg  Tab 2.5 mg  TAB 5 mg  NALAPRIL MALEATE  Tab 5 mg  Tab 10 mg  Tab 20 mg  SINOPRIL  Tab 5 mg	en under 12 years of age. In the dised for patients who were taking cilazapsets may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42	ril prior to 1 May rsed where there 90 90 90 100 100 100	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre in Carlon in Car	en under 12 years of age. In the dised for patients who were taking cilazapsets may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42 11.07 11.67	ril prior to 1 May rsed where there 90 90 90 100 100 100 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml	en under 12 years of age. In the dised for patients who were taking cilazapsets may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42	ril prior to 1 May rsed where there 90 90 90 100 100 100	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml	en under 12 years of age. In the dised for patients who were taking cilazapsets may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42 11.07 11.67	ril prior to 1 May rsed where there 90 90 90 100 100 100 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril
APTOPRIL Oral liq 5 mg per ml Oral liquid restricted to childre in Carlor of the	en under 12 years of age. In the dised for patients who were taking cilazapsets may annotate the prescription as endo  2.09 4.80 8.35  1.82 2.02 2.42 11.07 11.67	ril prior to 1 May rsed where there 90 90 90 100 100 100 90 90	2021 and the prescription exists a record of prior  Zapril Zapril Zapril Acetec Acetec Acetec Ethics Lisinopril Ethics Lisinopril

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
QUINAPRIL				
Tab 5 mg	5.97	90	✓ .	Arrow-Quinapril 5
Tab 10 mg	5.18	90	✓ .	Arrow-Quinapril 10
Tab 20 mg	7.95	90	✓	Arrow-Quinapril 20

### **ACE Inhibitors with Diuretics**

QUINAPRIL WITH HYDROCHLOROTHIAZIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide.

✓ Accuretic 10	30	g4.10	Tab 10 mg with hydrochlorothiazide 12.5 mg
✓ Accuretic 20	30	g5.25	Tab 20 mg with hydrochlorothiazide 12.5 mg

## **Angiotensin II Antagonists**

* Tab 4 mg	2.00	90	<ul><li>Candestar</li></ul>
* Tab 8 mg		90	✓ Candestar
* Tab 16 mg		90	✓ Candestar
* Tab 32 mg	5.26	90	✓ Candestar
LOSARTAN POTASSIUM			
Tab 12.5 mg	1.56	84	✓ Losartan Actavis
Tab 25 mg	1.84	84	✓ Losartan Actavis
Tab 50 mg	2.25	84	✓ Losartan Actavis
Tab 100 mg	3.50	84	✓ Losartan Actavis

## Angiotensin II Antagonists with Diuretics

Tab 50 mg with hydrochlorothiazide 12.5 mg	15.25	30	✓ Arrow-Losartan &
			Hydrochlorothiazide

## Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

#### ⇒SA1905 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
  - 2.1 Patient is in NYHA/WHO functional class II; or
  - 2.2 Patient is in NYHA/WHO functional class III: or
  - 2.3 Patient is in NYHA/WHO functional class IV; and

continued...

Sub:	•	Fully	Brand or
(Manufactu		sidised	Generic
\$	S Per	•	Manufacturer

continued...

- 3 Either:
  - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
  - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
  - 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, p	age 118	
AMIODARONE HYDROCHLORIDE		
▲ Tab 100 mg	30	✓ Aratac
▲ Tab 200 mg5.25	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule - Up to 10 inj available on a		
PSO16.37	10	✓ Max Health
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule - Up to 5 inj available on a		
PSO15.09	10	✓ <u>Martindale</u>
DIGOXIN		
* Tab 62.5 mcg – Up to 30 tab available on a PSO	240	✓ <u>Lanoxin PG</u>
* Tab 250 mcg – Up to 30 tab available on a PSO	240	Lanoxin
* Oral liq 50 mcg per ml	60 ml	<ul><li>✓ Lanoxin</li><li>✓ Lanoxin Paediatric</li></ul>
		Elixir S29
		✓ Lanoxin S29 S29
DIOODYDAMIDE DIOODIATE		Lanoxin 529 529
DISOPYRAMIDE PHOSPHATE  A Cap 100 mg23.87	100	✓ Rythmodan
	100	• Hydiiilouali
FLECAINIDE ACETATE <b>A</b> Tab 50 mg19.95	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	90	✓ Flecainide BNM
<b>2</b> Out to fig accuracy footing	00	Controlled
		Release Teva
▲ Cap long-acting 200 mg61.06	90	✓ Flecainide
		Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.00	100	✓ Teva S29
▲ Cap 250 mg202.00	100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg40.90	50	✓ Rytmonorm

## **Antihypotensives**

MIDODRINE - Special Authority see SA1474 on the next	page – Retail pharmacy		
Tab 2.5 mg	53.00	100	<ul><li>Gutron</li></ul>
Tab 5 mg	79.00	100	<ul><li>Gutron</li></ul>

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

### ⇒SA1474 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## **Beta-Adrenoceptor Blockers**

## **Beta Adrenoceptor Blockers**

ATENOI OI

ATENOLOL			
* Tab 50 mg	9.33	500	✓ Mylan Atenolol
* Tab 100 mg	14.20	500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
			<b>S29</b> S29
	38.20		✓ Essential
	00.20		Generics \$29
	49.85		✓ Atenolol AFT
Restricted to children under 12 years of age.	49.00		▼ Ateriolol AFT
, ,			
BISOPROLOL FUMARATE			
* Tab 2.5 mg		90	✓ Bisoprolol Mylan
* Tab 5 mg		90	✓ Bisoprolol Mylan
* Tab 10 mg	3.62	90	✓ Bisoprolol Mylan
CARVEDILOL			
* Tab 6.25 mg	2.24	60	<ul><li>Carvedilol Sandoz</li></ul>
* Tab 12.5 mg	2.30	60	<ul><li>Carvedilol Sandoz</li></ul>
* Tab 25 mg	2.95	60	<ul><li>Carvedilol Sandoz</li></ul>
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
* Tab 200 mg		100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	
, 01	(88.60)		Trandate
* inj 5 mg per ml, 20 ml vial	42.29 <sup>′</sup>	1	
, 01	(48.20)		Alvogen S29
METOPROLOL SUCCINATE	(10.00)		9
* Tab long-acting 23.75 mg	1 //5	30	✓ Betaloc CR
* Tab long-acting 25.75 mg		30	✓ Betaloc CR
* Tab long-acting 47.5 mg		30	✓ Betaloc CR
* Tab long-acting 95 mg		30	✓ Betaloc CR
	4.21	30	• Detailor Cit
METOPROLOL TARTRATE	F 00	100	/ IDOA Matarasalal
Tab 50 mg		100	✓ IPCA-Metoprolol
Tab 100 mg		60	✓ IPCA-Metoprolol
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	Metoprolol IV Mylan
NADOLOL			
Tab 40 mg	19.19	100	✓ Nadolol BNM S29
Tab 80 mg	30.39	100	✓ Nadolol BNM S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPRANOLOL				
Tab 10 mg	7.04	100	✓ [	Profate
Tab 40 mg	8.75	100	<b>√</b>	PCA-Propranolol
* Cap long-acting 160 mg		100	✓ (	Cardinol LA
* Oral lig 4 mg per ml - Special Authority see SA1327 be				
Retail pharmacy		500 m	ıl <b>√</b> F	Roxane- Propranolol 829

#### ⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

#### SOTALOL

*	Tab 80 mg	32.58	500	✓ Mylan
*		10.00	100	✓ Mylan

## **Calcium Channel Blockers**

## **Dihydropyridine Calcium Channel Blockers**

AM	LODIPINE		
*	Tab 2.5 mg1.08	90	✓ Vasorex
*	Tab 5 mg	90	✓ Vasorex
*	Tab 10 mg1.19	90	✓ Vasorex
FE	LODIPINE		
*	Tab long-acting 2.5 mg1.45	30	✓ Plendil ER
*	Tab long-acting 5 mg4.07	90	✓ Felo 5 ER
*	Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
NIF	EDIPINE		
*	Tab long-acting 10 mg	56	✓ Tensipine MR10 S29
*	Tab long-acting 20 mg9.12	50	✓ Mylan (12 hr release) S29
	17.72	100	✓ Nyefax Retard
*	Tab long-acting 30 mg4.78	14	Mylan Italy (24 hr
			release) \$29
	34.10	100	✓ Mylan (24 hr
			release) \$29
*	Tab long-acting 60 mg52.81	100	Mylan (24 hr
			release) S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44 40	100	1	Accord S29
★ Cap long-acting 120 mg		500		Apo-Diltiazem CD
★ Cap long-acting 180 mg		30		Cardizem CD
K Cap long-acting 240 mg		30		Cardizem CD
PERHEXILINE MALEATE			_	
★ Tab 100 mg	62 90	100	<b>√</b> 1	Pexsig
· ·	02.30	100	• !	CASIG
/ERAPAMIL HYDROCHLORIDE	7.04	100		
≮ Tab 40 mg ≮ Tab 80 mg		100 100	_	soptin
				soptin
* Tab long-acting 120 mg	36.02	100		soptin Retard \$29
L. Top long acting 040 mg	15 10	20		soptin SR
* Tab long-acting 240 mg	15.12	30	<b>V</b> 1	soptin SR
★ Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a	05.00	_		
PSO	25.00	5	<b>V</b> 1	soptin
Centrally-Acting Agents				
CLONIDINE				
★ Patch 2.5 mg, 100 mcg per day — Only on a prescription		4		<u> Mylan</u>
★ Patch 5 mg, 200 mcg per day – Only on a prescription		4		<u>Mylan</u>
★ Patch 7.5 mg, 300 mcg per day — Only on a prescription	16.93	4	<b>✓</b> [	<u> Mylan</u>
CLONIDINE HYDROCHLORIDE				
<b>₭</b> Tab 25 mcg	8.75	112		Clonidine BNM
	36.50			Clonidine Teva
★ Tab 150 mcg		100	-	Catapres Participation
★ Inj 150 mcg per ml, 1 ml ampoule	29.68	10	<b>√</b> <u>i</u>	<u> Medsurge</u>
METHYLDOPA				
★ Tab 250 mg	15.10	100	<b>✓</b> [	Methyldopa Mylan
	52.85	500	<b>✓</b> [	Methyldopa Mylan
				<b>S29</b> S29
Diuretics				
Loop Diuretics				
BUMETANIDE				
<b>₭</b> Tab 1 mg	4.91	30	<b>√</b> [	Burinex S29 S29
•	16.36	100		Burinex
₭ Inj 500 mcg per ml, 4 ml vial		5	,	Burinex

Subs (Manufactur \$	er's Price)	Fully Subsidised	d Generic
FUROSEMIDE (FRUSEMIDE)			
Tab 40 mg - Up to 30 tab available on a PSO8.0	0 1,00	00 🗸	IPCA-Frusemide
* Tab 500 mg25.0	0 50	· •	Urex Forte
89.4	8	✓	Furosemid-
			Ratiopharm S29
169.9	6 100	0 🗸	Furosemid-
			Ratiopharm S29
* Oral liq 10 mg per ml11.2	0 30 ml	OP 🗸	Lasix
* Inj 10 mg per ml, 25 ml ampoule		•	Lasix
* Inj 10 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO 1.1	5 5	•	Furosemide-Baxter
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Oral liq 1 mg per ml32.1	0 25 ml	OP 🗸	Biomed
EPLERENONE – Special Authority see SA1728 below – Retail pharmacy			
Tab 25 mg18.5	0 30	•	Inspra
Inspra to be Principal Supply on 1 June 2022			-
Tab 50 mg25.0	0 30	•	Inspra
Inspra to be Principal Supply on 1 June 2022			

## **⇒SA1728** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Both:

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Fither
  - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
  - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

M	ΕT	О	LAZ	ZΟ	NI	Ξ

	Tab 5 mg	CBS	1	1	Metolazone S29
			50	✓	Zaroxolyn S29
SP	PIRONOLACTONE				
*	Tab 25 mg	3.68	100	1	Spiractin
	Tab 100 mg		100	1	Spiractin
	Oral liq 5 mg per ml		25 ml OP		Biomed

# **Potassium Sparing Combination Diuretics**

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE		
* Tab 5 mg with furosemide 40 mg8.63	28	✓ Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE		
* Tab 5 mg with hydrochlorothiazide 50 mg 5 00	50	✓ Moduretic

	Subsidy (Manufacturer's Pri	ce) Subs	Fully sidised	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]  * Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ <u>A</u>	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than eme  * Tab 5 mg	• .	500	✓ <u>A</u>	rrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	27.82	25 ml OP	<b>√</b> B	iomed
CHLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	3.90 6.50	30 50		groton \$29 ygroton
INDAPAMIDE  * Tab 2.5 mg	10.45 11.61	90 100		<u>apa-Tabs</u> Iylan Indapamide (\$29)
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE  * Tab 200 mg  * Tab long-acting 400 mg		90 30	_	ezalip ezalip Retard
Other Lipid-Modifying Agents				
ACIPIMOX * Cap 250 mg	21.56	30	-	olbetam olbetam S29 S29
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	<b>✓</b> C	olestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN  * Tab 10 mg  * Tab 20 mg  * Tab 40 mg  * Tab 80 mg  PRAVASTATIN	9.24 14.92	500 500 500 500	✓ <u>L</u>	orstat orstat orstat orstat
* Tab 20 mg*  * Tab 40 mg		28 28	_	ravastatin Mylan ravastatin Mylan

	Subsidy (Manufacturer's Price) \$		Fully ubsidised	Brand or Generic Manufacturer
ROSUVASTATIN - Special Authority see SA2093 below - Retail	pharmacy			
Tab 5 mg	1.70	30	✓ F	Rosuvastatin Viatris
Tab 10 mg	2.42	30	✓ F	Rosuvastatin Viatris
Tab 20 mg	3.92	30	✓ F	Rosuvastatin Viatris
Tab 40 mg	5.28	30	<b>✓</b> <u>F</u>	Rosuvastatin Viatris

#### ⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

#### . . .

- 1 Both:
  - 1.1 Patient is considered to be at risk of cardiovascular disease; and
  - 1.2 Patient is Māori or any Pacific ethnicity: or
- 2 Both:
  - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
  - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 Patient has proven coronary artery disease (CAD); or
  - 1.2 Patient has proven peripheral artery disease (PAD); or
  - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	//VASTATIN			
*	Tab 10 mg	1.23	90	<ul><li>Simvastatin Mylan</li></ul>
	Tab 20 mg		90	✓ Simvastatin Mylan
*	Tab 40 mg	3.58	90	✓ Simvastatin Mylan
	Tab 80 mg		90	✓ Simvastatin Mylan

## **Selective Cholesterol Absorption Inhibitors**

EZ	ETIMIBE - Special Authority see SA1045 on the next page - Retail pharmacy		
*	Tab 10 mg1.95	30	✓ Ezetimibe Sandoz

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	idised	Generic
 \$	Per	1	Manufacturer

## **⇒SA1045** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
  - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
  - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
  - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

### EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

#### ⇒SA1046 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

#### **Nitrates**

### GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Sprav
	Patch 25 mg, 5 mg per day Patch 50 mg, 10 mg per day		30 30	✓ Nitroderm TTS ✓ Nitroderm TTS

	C	AKD	IUVAS	CULAR SYSTEM
	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
ISOSORBIDE MONONITRATE				
* Tab 20 mg		100		Ismo 20
* Tab long-acting 40 mg		30		Ismo 40 Retard
* Tab long-acting 60 mg	9.25	90	•	<u>Duride</u>
Sympathomimetics				
ADRENALINE				
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a F	SO4.98	5		Aspen Adrenaline
	10.76	_		DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a		5		Hospira
	49.00	10		Aspen Adrenaline
Vasodilators				
HYDRALAZINE HYDROCHLORIDE				
* Tab 25 mg - Special Authority see SA1321 below - Reta	il			
pharmacy	CBS	1	1	Hydralazine
		56	•	Onelink S29
		84	1	AMDIPHARM \$29
		100	1	Onelink \$29
* Inj 20 mg ampoule	25.90	5	•	Apresoline
■ SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals we the following criteria: Either:  1 For the treatment of refractory hypertension; or 2 For the treatment of heart failure in combination with a				
inhibitors and/or angiotensin receptor blockers.	milato, in patiente une c	210 11110	,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	That's not responded to he
MINOXIDIL  A Tab 10 mg	70.00	100	1	Loniten
NICORANDIL			-	
▲ Tab 10 mg	25 57	60	/	Ikorel
▲ Tab 20 mg		60		Ikorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257 12	5	/	Hospira
		J	•	
PENTOXIFYLLINE [OXPENTIFYLLINE] Tab 400 mg	42.26	50	1	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN - Special Authority see SA1702 below - Ret	ail pharmacy			
Tab 5 mg		30	1	Ambrisentan Mylan
Toh 10 mg	1 550.00	20		Ambricanton Mulan

MDRISENTAN - Special Authority see SAT702 below -	- Retail pharmacy		
Tab 5 mg	1,550.00	30	Ambrisentan Mylan
Tab 10 mg	·	30	✓ Ambrisentan Mylan
·	,		✓ Mylan

## **⇒SA1702** Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic	
BOSENTAN - Special Authority see SA1991 below - Retail phar	macy				
Tab 62.5 mg	119.85	60	•	Bosentan Dr Reddy's	
Tab 125 mg	119.85	60	•	Bosentan Dr Reddy's	

#### ⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II. III. or IV: and
- 4 Any of the following:
  - 4.1 Both:
    - 4.1.1 Bosentan is to be used as PAH monotherapy; and
    - 1.1.2 Either
      - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
      - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
  - 4.2 Both:
    - 4.2.1 Bosentan is to be used as PAH dual therapy; and
    - 4.2.2 Either:
      - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
      - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
  - 4.3 Both:
    - 4.3.1 Bosentan is to be used as PAH triple therapy; and
    - 4.3.2 Any of the following:
      - 4.3.2.1 Patient is on the lung transplant list; or
      - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
      - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
      - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

**Renewal** only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Bosentan is to be used as PAH monotherapy; and
  - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
  - 2.1 Bosentan is to be used as PAH dual therapy; and
  - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
  - 3.1 Bosentan is to be used as PAH triple therapy; and
  - 3.2 Any of the following:
    - 3.2.1 Patient is on the lung transplant list; or
    - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or

continued...

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subs	idised	Generic
	\$	Per	1	Manufacturer

continued...

- 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
- 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

## **Phosphodiesterase Type 5 Inhibitors**

SILDENAFIL - Special Authority see SA1992 below - Retail pharmacy		
Tab 25 mg	4	✓ Vedafil
Tab 50 mg1.70	4	✓ Vedafil
Tab 100 mg10.20	12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon\*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon\*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs): and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension\*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)\*; and
- 2 Any of the following:
  - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
  - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
  - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
  - 3.1 PAH is in NYHA/WHO functional class II: or
  - 3.2 PAH is in NYHA/WHO functional class III: or
  - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
    - 4.1.2 Either:
      - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
      - 4.1.2.2 Patient is peri Fontan repair; and
    - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
  - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with \* are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

continued...

Subsid (Manufacturer		ılly Brand or ed Generic	
	Per	✓ Manufacturer	

continued...

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

## **Prostacyclin Analogues**

EPOPROSTENOL – Special Authority see SA1696 below – Retail pharmacy		
Inj 500 mcg vial36.61	1	✓ Veletri
Inj 1.5 mg vial73.21	1	✓ Veletri

## **⇒SA1696** Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml ......740.10 30 ✓ Ventavis

#### ⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

### **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

# **Antiacne Preparations**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

#### **ADAPALENE**

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	<ul><li>Differin</li></ul>
ISOTRETINOIN - Special Authority see SA2023 below - R	letail pharmacy		
Cap 5 mg	11.26	60	<ul><li>Oratane</li></ul>
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

#### ⇒SA2023 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
  - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

**TRFTINOIN** 

Crm 0.5 mg per q − Maximum of 50 g per prescription......15.57 50 g OP ✓ ReTrieve

## **Antibacterials Topical**

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

## **DERMATOLOGICALS**

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer
MUPIROCIN			
Oint 2%		15 g OP	
a). Oak an a marandation	(11.50)		Bactroban
<ul><li>a) Only on a prescription</li><li>b) Not in combination</li></ul>			
ODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2%	1.59	5 g OP	✓ Foban
<ul> <li>a) Maximum of 5 g per prescription</li> </ul>			
b) Only on a prescription			
c) Not in combination	1.50	E ~ OD	√ Cabo∷
Oint 2%	1.59	5 g OP	✓ <u>Foban</u>
<ul><li>a) Maximum of 5 g per prescription</li><li>b) Only on a prescription</li></ul>			
c) Not in combination			
SULFADIAZINE SILVER			
Crm 1%	10.80	50 g OP	✓ Flamazine
a) Up to 250 g available on a PSO	10.00	30 g Oi	• I lalliazille
b) Not in combination			
Antitungals Tonical			
Antifungals Topical			
	ale page 96		
or systemic antifungals, refer to INFECTIONS, Antifunga	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga MOROLFINE  a) Only on a prescription	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga MOROLFINE  a) Only on a prescription b) Not in combination		5 ml OP	✓ MycoNail
for systemic antifungals, refer to INFECTIONS, Antifungal MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ <u>MycoNail</u>
For systemic antifungals, refer to INFECTIONS, Antifunga AMOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%	14.93	5 ml OP 20 g OP	✓ <u>MycoNail</u> ✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%	14.93	20 g OP	
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
ior systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol  Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol  Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol  Canesten
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP 20 g OP	✓ Clomazol  Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE  a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP 20 g OP	✓ Clomazol  Canesten  Pevaryl

	Subsidy (Manufacturer's P		Fully sidised	Brand or Generic
	<u> </u>	Per		Manufacturer
IICONAZOLE NITRATE				
€ Crm 2%	0.81	15 g OP	✓ N	lultichem
<ul> <li>a) Only on a prescription</li> </ul>		•	· <u></u>	
b) Not in combination				
<b>k</b> Lotn 2%	4.36	30 ml OP		
	(10.03)		D	)aktarin
a) Only on a prescription	,			
b) Not in combination				
F Tinct 2%	4.36	30 ml OP		
	(12.10)		D	aktarin
a) Only on a prescription	( - /			
b) Not in combination				
-,				
Antipruritic Preparations				
CALAMINE				

CALAMINE		
a) Only on a prescription		
b) Not in combination		
Crm, aqueous, BP1.08	3 100 g	✓ Calamine-AFT
CROTAMITON		
a) Only on a prescription		
b) Not in combination		
Ćrm 10%	9 20 g OP	✓ <u>Itch-Soothe</u>
MENTHOL - Only in combination		

Only in combination with a dermatological base or proprietary Topical Corticosteriod – Plain

2) With or without other dermatological galenicals.

5 g <b>✓ MidWest</b>	25 g	Crystals6.92
00 g ✓ MidWest	100 g	29.60

# **Corticosteroids Topical**

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

### Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE	
Crm 0.05%2.96	S 15 g OP ✓ <u>Diprosone</u>
36.00	50 g OP ✓ <u>Diprosone</u>
Oint 0.05%2.96	6 15 g OP ✓ <u>Diprosone</u>
36.00	0 50 g OP ✓ <u>Diprosone</u>
Oint 0.05% in propylene glycol base4.33	30 g OP ✓ Diprosone OV
BETAMETHASONE VALERATE	
* Crm 0.1%4.53	B 50 g OP ✓ Beta Cream
* Oint 0.1%	
* Lotn 0.1%	50 ml OP ✓ Betnovate
CLOBETASOL PROPIONATE	
* Crm 0.05%2.18	30 g OP ✓ Dermol
<b>*</b> Oint 0.05%2.12	2 30 g OP <b>✓ Dermol</b>

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully	Brand or
	(Manufacturer's F	rice) Subs	idised	Generic
	\$	Per	1	Manufacturer
CLOBETASONE BUTYRATE				
Crm 0.05%	5.38	30 g OP		
	(10.00)	J	Е	Eumovate
HYDROCORTISONE	,			
* Crm 1% – Only on a prescription	3 70	100 g OP	<b>✓</b> L	lydrocortisone
Total 178 - Only on a prescription		100 g O1	• <u>1</u>	(PSM)
	17.15	500 g	<b>✓</b> ⊦	lydrocortisone
	17.10	300 g	• 1	(PSM)
* Powder – Only in combination	40.05	25 g	<b>✓</b> A	<del></del>
Up to 5% in a dermatological base (not proprietary Topic				
galenicals	ai Corticosterioc	ı – Fiairi) Willi C	) WILIO	ut other dermatological
· ·				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - Only				
a prescription	10.57	250 ml	✓ [	OP Lotn HC
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%	4.85	100 g OP	<b>√</b> L	ocoid Lipocream
Oint 0.1%		100 g OP	<b>√</b> L	.ocoid
Milky emul 0.1%	12.33	100 ml OP	<b>√</b> L	ocoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	<b>✓</b> A	Advantan
Oint 0.1%		15 g OP	_	Advantan
MOMETASONE FUROATE		- 3 -	_	
Crm 0.1%	1 95	15 g OP	<b>√</b> F	locon Alcohol Free
0111 0.1 /0	3.10	50 g OP	_	locon Alcohol Free
Oint 0.1%		15 g OP		locon
Ont 0.1 /0	2.90	50 g OP		locon
Lotn 0.1%		30 ml OP	_	locon
TRIAMCINOLONE ACETONIDE			_	
Crm 0.02%	6 20	100 g OP	1	Aristocort
Oint 0.02%		100 g OP	_	Aristocort
Olit 0.02 /6	0.03	100 g O1	· -	anstocort
Corticosteroids - Combination				
Cornocaterolas Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FU	SIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49	15 g OP		
	(10.45)		F	ucicort
<ul> <li>a) Maximum of 15 g per prescription</li> </ul>				
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescrip	ntion			
* Crm 1% with miconazole nitrate 2%		15 g OP	✓ N	Nicreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - C		•		
			<b>√</b> n	Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP	<b>▼</b> P	iiiidlucult
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC		IN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 m	•			
and gramicidin 250 mcg per g - Only on a prescription.		15 g OP		
	(9.28)		V	iaderm KC

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Barrier Creams and Emollients		
Barrier Creams		
DIMETHICONE  * Crm 5% pump bottle4.48	500 ml OP	✓ healthE
* OIII 5 % pump bottle4.40	300 IIII OF	Dimethicone 5%
* Crm 10% pump bottle4.52	500 ml OP	✓ healthE Dimethicone 10%
ZINC AND CASTOR OIL		
* Oint	500 g	✓ Boucher
Emollients		
AQUEOUS CREAM		
* Crm	500 g	<ul><li>✓ Boucher</li><li>✓ GEM Aqueous</li><li>Cream</li></ul>
GEM Aqueous Cream to be Principal Supply on 1 July 2022 (Boucher Crm to be delisted 1 August 2022)		
CETOMACROGOL  * Crm BP	E00 a	✓ Cetomacrogol-AFT
* CITI BF1.99 CETOMACROGOL WITH GLYCEROL	500 g	▼ Cetomacrogor-AF1
Crm 90% with glycerol 10%2.35	500 ml OP	✓ <u>Boucher</u> ✓ Pharmacy Health Sorbolene with Glycerin
3.10	1,000 ml OP	✓ Boucher
EMULSIFYING OINTMENT	•	
* Oint BP	500 g	<ul><li>Emulsifying     Ointment ADE</li></ul>
OIL IN WATER EMULSION		
* Crm	9	<ul> <li>✓ Fatty Cream AFT</li> <li>✓ O/W Fatty Emulsion</li> <li>Cream</li> </ul>
(O/W Fatty Emulsion Cream Crm to be delisted 1 September 2022)		Orcam
PARAFFIN		
Oint liquid paraffin 50% with white soft paraffin 50%5.35 UREA	500 ml OP	✓ healthE
* Crm 10%1.37	100 g OP	✓ healthE Urea Cream
WOOL FAT WITH MINERAL OIL - Only on a prescription		
* Lotn hydrous 3% with mineral oil	,	DB Lation
(11.95 1.40	,	DP Lotion
(4.53		DP Lotion
5.60		
(20.53 (23.91	, )	Alpha-Keri Lotion BK Lotion
4.40	050 100	

(23.91) 1.40

(7.73)

250 ml OP

**BK** Lotion

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. \*Three months or six months, as applicable, dispensed all-at-once

## **DERMATOLOGICALS**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

## Other Dermatological Bases

#### **PARAFFIN**

450 g ✓ healthE 2.500 a ✓ healthE 19.99

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid - Plain.

## **Minor Skin Infections**

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
<ul> <li>a) Maximum of 130 g per prescription</li> </ul>			
b) Only on a prescription			
Antiseptic Solution 10%	4.15	100 ml	✓ Riodine
Antiseptic soln 10%		15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

## **Parasiticidal Preparations**

#### DIMETHICONE

200 ml OP ✓ healthE Dimethicone 4% Lotion

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy Tab 3 mg - Up to 100 tab available on a PSO.......17.20

1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and

- a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

#### ⇒SA1225 Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

## Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist: and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or

continued...

✓ Stromectol

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
  - 2.2.1 The Patient is a resident in an institution; and
  - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
  - 2.2.3 Any of the following:
    - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
    - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
    - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 The patient is in the community; and
    - 2.1.2 Any of the following:
      - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy: or
      - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
  - 2.2 All of the following:
    - 2.2.1 The Patient is a resident in an institution; and
    - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
    - 2.2.3 Any of the following:
      - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
      - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
      - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- Strongyloidiasis.

#### **PERMETHRIN**

Crm 5%5.75	30 g OP	✓ Lyderm
Lotn 5%	30 ml OP	✓ A-Scables

Subsidy	Fully	Brand or	П
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

## **Psoriasis and Eczema Preparations**

ACITRETIN - Special Authority see SA2024 below - Retail pharmac	у		
Cap 10 mg	17.86	60	✓ <u>Novatretin</u>
Cap 25 mg	41.36	60	✓ Novatretin

#### ⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
  - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
  - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment;
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL	
DETAINETHASONE DIFTOFTONATE WITH CALCIFOTHIOL	

Foam spray 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP 60 g OP 30 g OP	<ul><li>✓ Enstilar</li><li>✓ <u>Daivobet</u></li><li>✓ <u>Daivobet</u></li></ul>
CALCIPOTRIOL Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex
COAL TAR Soln BP - Only in combination	36.25	200 ml	✓ Midwest
Up to 10% only in combination with a dermatologica		ietary Topical C	Corticosteriod – Pla

ain

2) With or without other dermatological galenicals.

### COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and

allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	J	Egopsoryl TA
	3.43	30 g OP	
	(4.35)	Ü	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 on the next page - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

Cream 1%	15 g OP	✓ Elidel
----------	---------	----------

### **DERMATOLOGICALS**

	Subsidy		Fully	Brand or
(M	lanufacturer's Price)	Sı	ubsidised	Generic
	\$	Per	1	Manufacturer

### ⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

## PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

**★** Soln 2.3% with trolamine laurilsulfate and fluorescein sodium..........4.44 500 ml ✓ Pinetarsol

## SALICYLIC ACID

Powder − Only in combination......18.88 250 g ✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
- 2) With or without other dermatological galenicals.

### **SULPHUR**

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

#### **TACROLIMUS**

Oint 0.1% − Special Authority see SA2074 below − Retail pharmacy.......33.00 30 g OP ✓ Zematop

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

#### ⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

## **Scalp Preparations**

BETAMETHASONE VALERATE  * Scalp app 0.1%	9.84	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE  * Scalp app 0.05%	5 69	30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%		100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%		100 ml OP	✓ Sebizole
Grampoo 276	0.20	100 1111 01	✓ <u>Sebizole</u>

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

### **DERMATOLOGICALS**

Subsidy (Manufacturer's Price) Fully Subsidised Brand or Generic Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

200 g OP

Per

✓ Marine Blue Lotion SPF 50+

**Wart Preparations** 

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

**IMIQUIMOD** 

Crm 5%, 250 mg sachet .......21.72 24 **✓ Perrigo** 

**PODOPHYLLOTOXIN** 

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

**Other Skin Preparations** 

Antineoplastics

FLUOROURACIL SODIUM

20 g OP

✓ Efudix

## **GENITO-URINARY SYSTEM**

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	1	Manufacturer	

# **Contraceptives - Non-hormonal**

## **Condoms**

_	a) Maximum of 60 dev per prescription			supplied under Section 20
		17.02		✓ Gold Knight XL
		14.87	144	✓ Shield XL
ĸ	60 mm	1.42	12	✓ Gold Knight XL
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
	, - <del></del>	15.57	144	✓ Gold Knight
<del>:</del>	56 mm, strawberry	1.30	12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	10.07	177	- word rangist
•	00 mm, 01000late	15.57	144	✓ Gold Knight
:	56 mm, chocolate	1.30	12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	11.01		-
	55 mm, 5.56 mm unordioco, 164	11.64	144	✓ Moments
:	56 mm, 0.08 mm thickness, red	n 97	10	✓ Moments
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	11.04	177	MOMENTS
+	50 mm, 0.00 mm unokness	11.64	144	✓ Moments
	56 mm, 0.08 mm thickness	0.07	10	✓ Moments
	<ul><li>a) Maximum of 60 dev per prescription</li><li>b) Up to 60 dev available on a PSO</li></ul>			
	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	b) Maximum of 60 dev per prescription	44.64	111	Cold Valabi
	a) Up to 60 dev available on a PSO			
	a) The to CO day available are a DOO	15.57	144	✓ Gold Knight
	56 mm, 0.05 mm thickness		12	✓ Gold Knight
	b) Up to 60 dev available on a PSO	4.00	10	. Cold V. Lubi
	a) Maximum of 60 dev per prescription			
	A Mariana (20 day)	11.64	144	✓ Moments
	56 mm		10	✓ Moments
	b) Maximum of 60 dev per prescription	0.07	10	/ Namenta
	a) Up to 60 dev available on a PSO			
		11.64	144	✓ <u>Moments</u>
+	53 mm, strawberry, red		10	Moments .
	b) Maximum of 60 dev per prescription	0.05	40	
	a) Up to 60 dev available on a PSO			
		11.64	144	✓ Moments
+	53 mm, chocolate, brown		10	✓ <u>Moments</u>
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		11.42	144	✓ <u>Moments</u>
+	53 mm, 0.05 mm thickness		10	✓ Moments
	b) Up to 60 dev available on a PSO			_
	a) Maximum of 60 dev per prescription			
		11.64	144	✓ Moments
	53 mm		10	✓ Moments
	49 mm - Up to 144 dev available on a PSO	11.42	144	✓ Moments

a) Maximumosidisedev per prescription b) Hindopassdayiqvailable on a PSO

#### GENITO-URINARY SYSTEI

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
*	60 mm (bulk pack)	14.87	144	•	Gold Knight XL
C	ontracentive Devices				

## ontraceptive Devices

#### INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO

	b) Only on a PSO			
*	IUD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓ Choice
	-			TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	Choice Load 375

# Contraceptives - Hormonal

## **Combined Oral Contraceptives**

## ⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- - 1 Patient is on a Social Welfare benefit: or
  - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - U	p to		
	84 tab available on a PSO	10.00	84	Mercilon 28

## **GENITO-URINARY SYSTEM**

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	✓	Microgynon 20 ED
·	6.45	112	✓	Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
ů ů	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Auth	nority see SA0500 on	the r	orevious n	age
b) Up to 63 tab available on a PSO	ionly dod on loods on		oroviouo p	ago
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO		84	1	Levlen ED
Op to 112 tab available on a 1 30	6.45	112		Femme-Tab ED
	0.45	112	•	remine-rab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to				
84 tab available on a PSO	6.95	84	1	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U				
to 84 tab available on a PSO	•	84	1	Norimin
10 0 1 tab aranabio on a 1 00	29.32	112		Norimin
	20.02	112	•	1401111111

# **Progestogen-only Contraceptives**

## ⇒SA0500 Special Authority for Alternate Subsidy

**Initial application** from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Fither:
  - 1.1 Patient is on a Social Welfare benefit: or
  - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

#### LEVONORGESTREL

*	Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	✓ Microlut
		22.00	112	✓ <u>Microlut</u>
*	Subdermal implant (2 × 75 mg rods) - Up to 3 pack available			
	on a PSO	.106.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE			
	Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PSO	7.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	✓	Noriday 28	
<b>Emergency Contraceptives</b>					
LEVONORGESTREL  * Tab 1.5 mg  a) Maximum of 2 tab per prescription b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted un		1 Part I		Postinor-1 A.	

# **Antiandrogen Oral Contraceptives**

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

#### CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

# **Gynaecological Anti-infectives**

Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel
CLOTRIMAZOLE (24.00)		Aci-Jei
<ul><li>* Vaginal crm 1% with applicators</li></ul>	35 g OP 20 g OP	✓ Clomazol ✓ Clomazol
MICONAZOLE NITRATE  * Vaginal crm 2% with applicator	40 g OP	✓ <u>Micreme</u>
NYSTATIN Vaginal crm 100.000 u per 5 g with applicator(s)4.00	75 a OP	✓ Nilstat

# **Myometrial and Vaginal Hormone Preparations**

ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule - Up to 5 inj available on a			
PSO	160.00	5	<ul> <li>DBL Ergometrine</li> </ul>
OESTRIOL			
* Crm 1 mg per g with applicator	6.62	15 g OP	✓ Ovestin
* Pessaries 500 mcg	6.86	15	✓ Ovestin
OXYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	<ul><li>Oxytocin BNM</li></ul>
Inj 10 iu per ml, 1 ml ampoule		5	<ul> <li>Oxytocin BNM</li> </ul>
OXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj available	le on a PSO		
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml		5	✓ Syntometrine

## GENITO-URINARY SYSTEM

	Subsidy		ully	Brand or
(Ma	nufacturer's Price)	Subsid	ised	Generic
	\$	Per	✓	Manufacturer

# **Pregnancy Tests - hCG Urine**

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

✓ Smith BioMed Rapid
Pregnancy Test

# **Urinary Agents**

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

## 5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

## ⇒SA0928 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
  - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
  - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

# **Alpha-1A Adrenoreceptor Blockers**

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

#### ⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

# **Other Urinary Agents**

OXYBUTYNIN

#### POTASSIUM CITRATE

Oral liq 3 mmol per ml – Special Authority see SA1083 on the

# **GENITO-URINARY SYSTEM**

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

## ⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

**Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	<ul><li>Solifenacin Mylan</li></ul>
Tab 10 mg	3.72	30	✓ Solifenacin Mylan

# **Detection of Substances in Urine**

ORTHO-TOLIDINE	<u>.                                      </u>	_	_
* Compound diagnostic sticks	7.50	50 test OP	
, , , , , , ,	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
	(13.92)		Albustix

# **Obstetric Preparations**

# **Antiprogesterones**

MIFEPRISTONE			
Tab 200 mg	60.00	1	✓ Mifegyne
-	180.00	3	✓ Mifegyne

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
· · · · ·	Por 🗸	Manufacturer

# Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	✓ Miacalcic
CINACALCET - Special Authority see SA1618 below -	- Retail pharmacy		
Tab 30 mg	42.06	28	✓ Cinacalet Devatis
a) Brand switch fee payable (Pharmacode 26	34120) - see page 241 for d	etails	
b) Wastage claimable			

## ⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Fither:

- 1 All of the following:
  - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and

Tab 60 mg - Wastage claimable ......84.12

1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and

28

✓ Cinacalet Devatis

- 1.3 The patient is symptomatic; or
- 2 All of the following:
  - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
  - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
  - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

**Renewal** only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

#### ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA2109 below −
Retail pharmacy.......18.00 1

✓ Zoledronic acid
Mylan

## ⇒SA2109 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
  - 2.1 Patient has bone metastases or involvement: and
  - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
  - 3.1 Patient has bone metastases or involvement; and
  - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer\*) from any relevant practitioner. Approvals valid for 3 years for applications meeting

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with \* are unapproved indications.

**Initial application — (symptomatic hypercalcaemia\*)** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia .

5

100

100

1

100

20

✓ Douglas✓ Douglas

✓ Medrol

✓ Medrol

✓ Solu-Cortef

Note: Indications marked with \* are unapproved indications.

Corticosteroids and Related Agents for Systemic Use

a) Up to 5 ini available on a PSO

b) Only on a PSO METHYLPREDNISOLONE

(36.96)	v	Celestone Chronodose
DEXAMETHASONE		
* Tab 0.5 mg - Up to 60 tab available on a PSO	30	✓ Dexmethsone
* Tab 4 mg - Up to 30 tab available on a PSO2.65	30	✓ Dexmethsone
Oral liq 1 mg per ml48.15	25 ml OP	✓ Biomed
DEXAMETHASONE PHOSPHATE  Dexamethasone phosphate injection will not be funded for oral use.		
* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO16.37	10	✓ <u>Dexamethasone</u> <u>Phosphate</u> <u>Panpharma</u>
FLUDROCORTISONE ACETATE		
* Tab 100 mcg14.32	100	✓ Florinef
HYDROCORTISONE		

79

	Subsidy (Manufacturer's Price \$	e) Su Per	Fully Brand or ubsidised Generic ✓ Manufacturer
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)			
Inj 40 mg vial	22.30	1	✓ Solu-Medrol-Act- O-Vial
Inj 125 mg vial	34.10	1	✓ Solu-Medrol-Act- O-Vial
Inj 500 mg vial	26.88	1	✓ Solu-Medrol-Act- O-Vial
Inj 1 g vial METHYLPREDNISOLONE ACETATE	32.84	1	✓ Solu-Medrol
Inj 40 mg per ml, 1 ml vial	47.06	5	✓ Depo-Medrol
Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓ <u>Redipred</u>
PREDNISONE  * Tab 1 mg	18.58	500	<ul><li>✓ Apo-Prednisone</li><li>✓ Prednisone Clinect</li></ul>
* Tab 2.5 mg	21.04	500	✓ Apo-Prednisone ✓ Prednisone Clinect
* Tab 5 mg - Up to 30 tab available on a PSO		500	<ul><li>✓ Apo-Prednisone</li><li>✓ Prednisone Clinect</li></ul>
* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	<ul><li>✓ Apo-Prednisone</li><li>✓ Prednisone Clinect</li></ul>
Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 20 mg to be delisted 1 November 2022)			
FETRACOSACTRIN			
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	✓ UK Synacthen S29 ✓ AU Synacthen ✓ Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ Synacthen Depot ✓ Synacthene Retard S29
FRIAMCINOLONE ACETONIDE			_
Inj 10 mg per ml, 1 ml ampoule Inj 40 mg per ml, 1 ml ampoule		5 5	✓ Kenacort-A 10 ✓ Kenacort-A 40
Sex Hormones Non Contraceptive			
Androgen Agonists and Antagonists			
CYPROTERONE ACETATE			_
Tab 50 mg		50	✓ <u>Siterone</u>
Tab 100 mg	28.03	50	✓ <u>Siterone</u>
FESTOSTERONE  Retch 5 mg per dev	00.00	20	Androde
Patch 5 mg per day  FESTOSTERONE CIPIONATE	90.00	30	✓ Androderm
Inj 100 mg per ml, 10 ml vial	85.00	1	✓ Depo-Testosterone

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	<u> </u>	Per		Manufacturer
TESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12.98	1	•	Sustanon Ampoules
TESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement	21.00	60	1	Andriol Testocaps
Subsidy by endorsement – subsidised for patients who w				
1 November 2021 and the prescription is endorsed acco	0,	,		
where there exists a record of prior dispensing of testost		ap 40 i		
Inj 250 mg per ml, 4 ml vial	86.00	1	•	Reandron 1000

# **Hormone Replacement Therapy - Systemic**

# Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

0	estrogens			
	STRADIOL - See prescribing guideline above			
*	Tab 1 mg	4.12 (11.10)	28 OP	Estrofem
*	Tab 2 mg	4.12 <sup>′</sup>	28 OP	
	Datab OF man new days	(11.10)	0	Estrofem
	Patch 25 mcg per day	6.12	8	✓ Estradot
	<ul><li>a) No more than 2 patch per week</li><li>b) Only on a prescription</li></ul>			
	Patch 50 mcg per day	7.04	8	✓ Estradot 50 mcg
	a) No more than 2 patch per week	7.04	O	- Estrador 30 meg
	b) Only on a prescription			
	Patch 75 mcg per day	7.91	8	✓ Estradot
	a) No more than 2 patch per week		•	
	b) Only on a prescription			
	Patch 100 mcg per day	7.91	8	✓ Estradot
	a) No more than 2 patch per week			
	b) Only on a prescription			
OF	STRADIOL VALERATE – See prescribing guideline above			
*	Tab 1 mg	12.36	84	✓ Progynova
*	Tab 2 mg	12.36	84	✓ Progynova
ΩF	STROGENS - See prescribing guideline above			
*	Conjugated, equine tab 300 mcg	3.01	28	
		(17.50)		Premarin
*	Conjugated, equine tab 625 mcg	4.12 <sup>′</sup>	28	
		(17.50)		Premarin
P	rogestogens			
ME	DROXYPROGESTERONE ACETATE - See prescribing gui	deline above		
*	Tab 2.5 mg		30	✓ Provera
*	Tab 5 mg	17.50	100	✓ Provera
*	Tab 10 mg	8.94	30	✓ Provera

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy

Fully

Brand or

	(Manufacturer's Price) \$	Per	sidised •	Generic Manufacturer	
Progestogen and Oestrogen Combined Prepara	ations				
OESTRADIOL WITH NORETHISTERONE – See prescribing gu  * Tab 1 mg with 0.5 mg norethisterone acetate		us page 28 OP			

OESTRADIOL WITH NORETHISTER	RONE – See prescribing gui	deline on the pre	vious page	
* Tab 1 mg with 0.5 mg norethister	one acetate	5.40	28 OP	
		(18.10)		Kliovance
* Tab 2 mg with 1 mg norethisteror	ne acetate	5.40	28 OP	
		(18.10)		Kliogest
* Tab 2 mg with 1 mg norethisteror	ne acetate (10), and 2 mg			
oestradiol tab (12) and 1 mg	oestradiol tab (6)	5.40	28 OP	
		(18.10)		Trisequens

# Other Oestrogen Preparations

ETHINYLOESTRADIOL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking ethinyloestradiol prior to 1 March 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ethinyloestradiol.

Tab 10 mcg	17.60	100	<ul> <li>NZ Medical and Scientific</li> </ul>
(NZ Medical and Scientific Tab 10 mcg to be delisted 1 Febr	uary 2023)		Scientific

OESTRIOL

# Other Progestogen Preparations

ena
dess
vera HD
nolut N
ogestan
1

## ⇒SA1609 Special Authority for Subsidy

**Initial application** only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour\*; and
- 2 Fither:
  - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

**Renewal** only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour\*; and
- 2 Treatment is required for second or subsequent pregnancy; and

(Mar	Subsidy nufacturer's Price)	F Subsid	ully	Brand or Generic
<u> </u>	\$	Per	1	Manufacturer

continued...

- 3 Either:
  - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
  - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with \* are unapproved indications.

# **Thyroid and Antithyroid Agents**

CA	RBIMAZOLE			
*	Tab 5 mg7	.56	100	✓ Neo-Mercazole
LE	VOTHYROXINE			
*	Tab 25 mcg5	.55	90	✓ Synthroid
*	Tab 50 mcg1	.71	28	Mercury Pharma
	5	.79	90	Synthroid
	64	.28	1,000	✓ Eltroxin
*	Tab 100 mcg1	.78	28	Mercury Pharma
	6	.01	90	Synthroid
	66	.78	1,000	✓ Eltroxin

PROPYLTHIOURACIL - Special Authority see SA1199 below - Retail pharmacy

Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

# ⇒SA1199 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

# **Trophic Hormones**

#### **Growth Hormones**

	harmacy	see SA2032 below – Retail ph	OMATROPIN (OMNITROPE) - Special Authority	SO
✓ Omnitrope	ĺ	69.75	Inj 5 mg cartridge	*
✓ Omnitrope	1	69.75	Inj 10 mg cartridge	*
✓ Omnitrope	1	139.50	Inj 15 mg cartridge	*

### ⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:

Subsidy (Manufacturer's Price)	F Subsidi	ully	Brand or Generic
 \$	Per	<b>√</b>	Manufacturer

continued...

- 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
- 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay: and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer	
	. 0.		mananananan	

continued...

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

**Initial application** — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
  - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m<sup>2</sup> as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m<sup>2</sup> in a child who may or may not be receiving dialysis; or
  - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	<b>✓</b>	Manufacturer

continued...

and/or ENT surgeon; and

- 5 Either:
  - 5.1 Both:
    - 5.1.1 The patient is aged two years or older; and
    - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
  - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

**Renewal — (Prader-Willi syndrome)** only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

- All of the following:
  - 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
  - 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
  - 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
  - 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
  - 5 No malignancy has developed after growth hormone therapy was commenced; and
  - 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

**Initial application** — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
  - 1.1 The patient has been treated with somatropin for < 12 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
- 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
  - 2.1 The patient has been treated with somatropin for more than 12 months; and
  - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
  - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
  - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
  - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
  - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
  - 3.3 The patient has severe growth hormone deficiency (see notes); and
  - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
  - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

# **GnRH Analogues**

GOSERELIN		
Implant 3.6 mg, syringe65.68	1	✓ Teva
Implant 10.8 mg, syringe122.37	1	✓ Teva
LEUPRORELIN		
Additional subsidy by endorsement where the patient is a child or adolescent and goserelin and the prescription is endorsed accordingly.	d is unable	to tolerate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy of		
\$221.60 per 1 inj with Endorsement	1	
(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy		
of \$591.68 per 1 inj with Endorsement177.50	1	
(591.68)		Lucrin Depot 3-month

# Vasopressin Agonists

DESMOPRESSIN			
Wafer 120 mcg	47.00	30	Minirin Melt

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	linirin
Tab 200 mcg	54.45	30	✓ N	linirin
▲ Nasal spray 10 mcg per dose		6 ml OP	<b>✓</b> <u>D</u>	esmopressin-
				PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	<b>✓</b> N	linirin

# **Other Endocrine Agents**

#### **CABERGOLINE**

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
<ul><li>Dostinex</li></ul>	2	waived by Special Authority see SA2070 below
✓ Dostinex	8	15.20

# ⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly\*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with \* is an unapproved indication.

CLOMIFENE CITRATE	
-------------------	--

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

**INFECTIONS - AGENTS FOR SYSTEMIC USE** Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 Fskazole S29 ⇒SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription Tab 100 mg .......7.97 6 Vermox 15 ml (7.53)Vermox PRAZIQUANTFI 8 Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 61 b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 236 Cephalosporins and Cephamycins CEFACLOR MONOHYDRATE Cap 250 mg......24.70 ✓ Ranbaxy-Cefaclor 100 ✓ Ranbaxy-Cefaclor S29 S29 ✓ Ranbaxy-Cefaclor 100 ml ✓ Ranbaxy-Cefaclor S29 S29 **CEFALEXIN** 20 ✓ Cephalexin ABM ✓ Cephalexin ABM 20 Grans for oral lig 25 mg per ml - Wastage claimable.....8.75 100 ml ✓ Cefalexin Sandoz ✓ Cefalexin Sandoz Grans for oral lig 50 mg per ml - Wastage claimable......11.75 100 ml CEFAZOLIN - Subsidy by endorsement Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly. CEFTRIAXONE - Subsidy by endorsement a) Up to 10 ini available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is

endorsed accordingly.

✓ Ceftriaxone-AFT ✓ Ceftriaxone-AFT 

CEFUROXIME AXETIL - Subsidy by endorsement

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

Zinnat 

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

#### **Macrolides**

AZITHROMYCIN - Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special

Tab 500 mg - Up to 8 tab available on a PSO ......2.57 Zithromax Grans for oral lig 200 mg per 5 ml (40 mg per ml) - Wastage Zithromax

#### ⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome\*: or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome\*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms\*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with \* are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis\*; and
- 2 Patient is aged 18 and under: and
- 3 Fither:
  - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
  - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis\*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with \* are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

Tab 250 mg ......8.53 14 ✓ Klacid Grans for oral lig 250 mg per 5 ml - Wastage claimable......192.00 ✓ Klacid 50 ml

#### ⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria: Either:

1 Atypical mycobacterial infection; or

Cubaide		F. III.	Drondor	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**ERYTHROMYCIN (AS LACTOBIONATE)** 

- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents. Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:
  - 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
  - 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	<ul><li>E-Mycin</li></ul>
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO			
<ul><li>b) Up to 2 x the maximum PSO quantity for RFPP</li><li>c) Wastage claimable</li></ul>			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO		100 1111	- L myom
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 500 mg	29.90	100	
	(44.58)		ERA
(ERA Tab 500 mg to be delisted 1 September 2022)			
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	Arrow- Roxithromycin
Tab 300 mg	16.33	50	✓ Arrow-
			Roxithromycin

(Rulide D Tab disp 50 mg to be delisted 1 September 2022)

	Subsidy (Manufacturer's Price \$	) S Per	Fully ubsidised	Brand or Generic Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500	1	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500	✓	<u>Alphamox</u>
<ul> <li>a) Up to 30 cap available on a PSO</li> </ul>				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	•	Alphamox 125
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>				
b) Wastage claimable			_	
Grans for oral liq 250 mg per 5 ml	1.73	100 ml	/	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable	45.07	40	,	U. I
Inj 250 mg vial		10		Ibiamox Ibiamox
Inj 500 mg vial Inj 1 g vial – Up to 5 inj available on a PSO		10 10		Ibiamox
	21.04	10	•	IDIAIIIOX
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab			_	
available on a PSO		10	•	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r		400 1		
per ml	5.00	100 ml	•	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r		00 1 01		O
per ml – Up to 200 ml available on a PSO	2.20 1	00 ml Ol	•	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj			_	
available on a PSO	375.97	10	•	Bicillin LA
BENZYLPENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10	1	Sandoz
FLUCLOXACILLIN				
Cap 250 mg - Up to 30 cap available on a PSO		250	1	Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO		500	✓	Flucloxacillin-AFT
Grans for oral liq 25 mg per ml	3.29	100 ml	1	<u>AFT</u>
<ul> <li>a) Up to 200 ml available on a PSO</li> </ul>				
b) Wastage claimable			_	
Grans for oral liq 50 mg per ml	3.68	100 ml	/	<u>AFT</u>
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial		10		Flucioxin
Inj 500 mg vial		10		Flucion
Inj 1 g vial – Up to 5 inj available on a PSO	5./0	5	•	Flucil

	Subsidy		Fully	
(I	Manufacturer's Price)	Subsic Per	dised •	
PHENOXYMETHYLPENICILLIN (PENICILLIN V)	<u> </u>			
Cap 250 mg – Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK
Cap 500 mg		50		Cilicaine VK
a) Up to 20 cap available on a PSO		00	Ť	Omounic VIX
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral lig 125 mg per 5 ml	2.99	100 ml	1	AFT
a) Up to 200 ml available on a PSO				<u></u>
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	1	AFT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe - Up to 5 inj available on a PSO	123.50	5	1	Cilicaine
Tetracyclines				
DOXYCYCLINE				
* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	1	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
,	(12.05)			Mino-tabs
* Cap 100 mg	19.32	100		
	(52.04)			Minomycin
OA4055 One stall Asalth with fan Manuel athur and Date				

#### ⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 $\label{temperature} \textbf{TETRACYCLINE} \ - \ \textbf{Special Authority see} \ \frac{\textbf{SA1332 below}}{\textbf{SA1332 below}} - \ \textbf{Retail pharmacy}$ 

### ⇒SA1332 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

#### Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

#### **CIPROFLOXACIN**

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	<b>✓</b>	Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.61	24	<b>√</b> [	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	<b>√</b> [	Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	ent		
Only if prescribed for dialysis or cystic fibrosis patient and the	e prescription is endor	rsed	accordingly.	
Inj 150 mg	65.00	1	✓ (	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule - Subsidy by endorsement	95.00	5	✓ [	OBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection a	nd the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	<b>✓</b> \	Wockhardt S29
	182.00	10	✓ 1	Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection a	nd the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓	Pfizer
	87.50	50	✓	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	t infection a	nd the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retai No patient co-payment payable	l pharmacy			
Tab 400 mg	42.00	5	1	<u>Avelox</u>

#### ⇒SA1740 Special Authority for Subsidy

**Initial application** — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Active tuberculosis\*: and
  - 1.2 Any of the following:
    - 1.2.1 Documented resistance to one or more first-line medications: or
    - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
    - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
    - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
    - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.\*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with \* are unapproved indications.

**Renewal** only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium\* and is symptomatic; and
- 2 Fither:
  - 2.1 Has tried and failed to clear infection using azithromycin; or
  - 2.2 Has laboratory confirmed azithromycin resistance; and

	Subsidy (Manufacturer's Price) \$	Subsid Per	ised	Brand or Generic Manufacturer	
continued 3 Treatment is only for 7 days.					
Initial application — (Penetrating eye injury) only from an opht requires prophylaxis following a penetrating eye injury and treatment Note: Indications marked with * are unapproved indications.			1 mont	th where the patient	ſ
PAROMOMYCIN - Special Authority see SA1689 below - Retail	pharmacy				
Cap 250 mg	126.00	16	✓ Hu	matin S29	
■ SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	cal microbiologist or g	gastroentero	ologist.	Approvals valid for	1
1 Patient has confirmed cryptosporidium infection; or					
2 For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microbapplications meeting the following criteria: Either:	piologist or gastroente	erologist. A	pprova	ls valid for 1 month	for
<ol> <li>Patient has confirmed cryptosporidium infection; or</li> <li>For the eradication of Entamoeba histolyica carriage.</li> </ol>					
PYRIMETHAMINE – Special Authority see SA1328 below – Reta Tab 25 mg	'	30	✓ Dai	raprim S29	
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:  1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	a period of 3 months		ouned	тог аррисацото тнее	zung
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg	67.85	36	✓ Fu	cidin	
SULFADIAZINE SODIUM - Special Authority see SA1331 below Tab 500 mg	- Retail pharmacy	56	✓ Wo	ockhardt §29	
⇒SA1331 Special Authority for Subsidy		50	• WO	CKIIaiul	
initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:			otified	for applications mee	eting
For the treatment of toxoplasmosis in patients with HIV for For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months of Formatter in the congenital toxoplasmosis until 12 months of Formatter in the congenital toxoplasmosis until 12 months of Formatter in the congenital toxoplasmosis until 12 months of Formatter in the congenitation in the conge	•	; or			
TOBRAMYCIN		_	4		
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement  Only if prescribed for dialysis or cystic fibrosis patient and		5 ndorsed acc		<b>bramycin Mylan</b> ly.	
Solution for inhalation 60 mg per ml, 5 ml - Subsidy by endorsement	395.00 56	3 dose	<b>✓</b> <u>Tol</u>	bramycin BNM	
b) Only if prescribed for a cystic fibrosis patient and the p	prescription is endors	ed accordir	ıgly.		
TRIMETHOPRIM  * Tab 300 mg – Up to 30 tab available on a PSO	18.55	50	✓ <u>TM</u>	<u> P</u>	

		Subsidy (Manufacturer's Price \$	) Su Per	Fully bsidised	Brand or Generic Manufacturer	
TRIME	THOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXA	AZOLE]				
* Tal	b trimethoprim 80 mg and sulphamethoxazole 400 mg - L to 30 tab available on a PSO		500	<b>✓</b> <u>Tı</u>	<u>risul</u>	
* Ora	al liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 r available on a PSO		100 ml	✓ D	eprim	
On	DMYCIN – Subsidy by endorsement ly if prescribed for a dialysis or cystic fibrosis patient or for icile following metronidazole failure and the prescription is			or for treat	ment of Clostridiun	n
	500 mg vial		1	✓ M	<u>ylan</u>	

# **Antifungals**

- a) For topical antifungals refer to DERMATOLOGICALS, page 62
- b) For topical antifungals refer to GENITO URINARY, page 75

#### FLUCONAZOLE

Cap 50 mg	28	✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg12.89	28	✓ Mylan
Powder for oral suspension 10 mg per ml – Special Authority		
see SA1359 below – Retail pharmacy109.34	35 ml	Diflucan
Wastage claimable		

### ⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Initial application — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

**Renewal** — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

**Renewal — (Immunocompromised)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

#### **ITRACONAZOLE**

Cap 100 mg	4.27	15	✓ <u>Itrazole</u>
Oral liq 10 mg per ml - Special Authority see SA1322 on the			
next page – Retail pharmacy. 14	1.80	150 ml OP	✓ Sporanox

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ⇒SA1322 Special Authority for Subsidy

**Initial application** only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

**Renewal** from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

#### KETOCONAZOI E

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - I	Retail pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml	761.13	105 ml OP	✓ Noxafil

#### ⇒SA1285 Special Authority for Subsidy

**Initial application** only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy\*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

#### Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression\* and requires on going posaconazole treatment.

Note: \* Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

#### **TERRINAFINE**

* Tab 250 mg	8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the ne	ext page - Retail phar	macy	
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml – Wastage			
claimable	1,523.22	70 ml	✓ Vfend

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	•	Manufacturer	

## **⇒SA1273** Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient has proven or probable invasive aspergillus infection; or
  - 3.2 Patient has possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
  - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
  - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
  - 3.3 Patient has fluconazole resistant candidiasis; or
  - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

## **Antimalarials**

PRIMAQUINE - Special Authority see SA1684 below -	- Retail pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaguine \$29

### ⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

**Renewal** only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

## **Antitrichomonal Agents**

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ <u>Arrow-Ornidazole</u>

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

# **Antituberculotics and Antileprotics**

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

#### CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

#### CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.
- Cap 250 mg.......344.00 60 **✓ Cyclorin** 529

#### DAPSONE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Dapsone	100	Tab 25 mg268.50	Т
Dapsone	100	Tab 100 mg329.50	Т

#### ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician

Tab 100 mg	85.73	100	✓ EMB Fatol S29
Tab 400 mg	49.34	56	✓ Myambutol S29

### ISONIAZID - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician
- **\*** Tab 100 mg ......23.00 100 ✓ <u>PSM</u>

#### ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician

*	Tab 100 mg with rifampicin 150 mg	89.82	100	✓ Rifinah
*	Tab 150 mg with rifampicin 300 mg	179.13	100	Rifinah

## PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

#### PROTIONAMIDE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

		Subsidy	, 0	Fully	Brand or
		(Manufacturer's Price \$	e) Si Per	ubsidised •	Generic Manufacturer
PY	RAZINAMIDE - Retail pharmacy-Specialist	<u> </u>			
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation	on of, an infectious	disease p	hysician,	clinical microbiologist or
×	respiratory physician Tab 500 mg	64.05	100	1	AFT-Pyrazinamide
	ABUTIN - Retail pharmacy-Specialist	04.93	100	• •	AF 1-F y I azılla illi üe
ПП	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation	on of, an infectious	disease p	hysician,	respiratory physician or
	gastroenterologist				
*	Cap 150 mg	299.75	30	✓ N	/lycobutin
RIF	AMPICIN – Subsidy by endorsement				
	a) No patient co-payment payable	in combination with	athar off		i atanbulananal
	<ul> <li>For confirmed recurrent Staphylococcus aureus infection i antimicrobial based on susceptibilities and the prescription</li> </ul>				
	Retail pharmacy - Specialist. Specialist must be an intern				
	paediatrician, or public health physician.			_	
	Cap 150 mg		100	_	Rifadin
*	Cap 300 mg Oral liq 100 mg per 5 ml		100 60 ml	_	<u>Rifadin</u> Rifadin
-	oral ng 100 mg par 0 mi	12.00	00 1111	· ·	<u>maam</u>
A	ntivirals				
Foi	eye preparations refer to Eye Preparations, Anti-Infective Prep	narations nage 236	3		
		parationo, pago 200	•		
Н	epatitis B Treatment				
	TECAVIR				
	Tab 0.5 mg		30	<b>✓</b> E	Entecavir Sandoz
LA	MIVUDINE – Special Authority see SA1685 below – Retail pha Tab 100 mg		28	17	Zetlam
	Oral liq 5 mg per ml		40 ml OF		Zeffix
<b>&gt;</b>	SA1685 Special Authority for Subsidy				
	ial application only from a relevant specialist or medical pract	titioner on the recor	nmendati	on of a re	elevant specialist.
	provals valid for 1 year where used for the treatment or prevent				
	newal from any relevant practitioner. Approvals valid for 2 year	ars where used for t	he treatm	nent or pre	evention of hepatitis B.
ΙĽ	NOFOVIR DISOPROXIL  Tenofovir disoproxil prescribed under endorsement for the tre	eatment of HIV is in	cluded in	the count	t of up to 4 subsidised
	antiretrovirals for the purposes of Special Authority SA1651.,		oluucu III	tile dodin	to up to 4 substation
*	Tab 245 mg (300.6 mg as a succinate)	38.10	30	<b>✓</b> T	enofovir Disoproxil
					Teva
Н	erpesvirus Treatments				
ΑU **	CLOVIR Tab dispersible 200 mg	1.60	25	<b>√</b> 1	.ovir
*	Tab dispersible 400 mg		56		.ovir
*	Tab dispersible 800 mg	5.98	35	<b>✓</b> <u>L</u>	.ovir
۷A	LACICLOVIR				
	Tab 500 mg		30		/aclovir
	Tab 1,000 mg	13./6	30	<b>✓</b> <u>\</u>	/aclovir

	Subsidy (Manufacturer's Price)	Sul	Fully bsidised	Brand or Generic	
	\$	Per	•	Manufacturer	
VALGANCICLOVIR - Special Authority see SA1993 below - R	Retail pharmacy				
Tab 450 mg	132.00	60	_	alganciclovir	
				Mylan	

## ⇒SA1993 Special Authority for Subsidy

**Initial application** — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
  - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Roth:
  - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
  - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Fither:
  - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
  - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for a months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
  - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
  - 2.3 Patient has cytomegalovirus retinitis.

**Renewal — (Cytomegalovirus in immunocompromised patients)** only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
  - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

## **Hepatitis C Treatment**

#### GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg ......24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg.......24,363.46

28 

Harvoni

#### **⇒SA1605** Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

# **HIV Prophylaxis and Treatment**

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA1994 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

#### ⇒SA1994 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment: and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

#### continued...

- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

**Renewal** from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or https://ashm.org.au/HIV/PrEP/ for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and
- 5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and
- 6 Fither:
  - 6.1 All of the following:
    - 6.1.1 Patient is male or transgender; and
    - 6.1.2 Patient has sex with men; and
    - 6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and
    - 6.1.4 Any of the following:
      - 6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or
      - 6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or
      - 6.1.4.3 Patient has used methamphetamine in the last three months; or
  - 6.2 All of the following:
    - 6.2.1 Patient has a regular partner who has HIV infection; and
    - 6.2.2 Partner is either not on treatment or has a detectable viral load; and
    - 6.2.3 Condoms have not been consistently used.

### **COVID-19 Treatments**

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Paxlovid

# Antiretrovirals

#### ⇒SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Prevention of maternal foetal transmission: or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
  - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
  - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
  - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria: Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:

|--|

continued...

- 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

# Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ - Special Authority see SA1651 on the previous	us page – Retail phari	macy	
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE - Special Authority see SA1651 on the previous	ous page – Retail pha	rmacy	
Tab 200 mg	770.00	60	✓ Intelence
NEVIRAPINE - Special Authority see SA1651 on the previous	ous page – Retail pha	rmacy	
Tab 200 mg	84.00	60	✓ Nevirapine
			<u>Alphapharm</u>
Oral suspension 10 mg per ml	203.55	240 ml	✓ Viramune
			Suspension

# Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the Tab 300 mg  Oral liq 20 mg per ml	180.00	Retail pharmac 60 240 ml OP	y ✔ <u>Ziagen</u> ✔ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority Note: abacavir with lamivudine (combination tablets) counts anti-retroviral Special Authority.	as two anti-retro	oviral medication	ns for the purposes of the
Tab 600 mg with lamivudine 300 mg		30	✓ <u>Kivexa</u>
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOPI Retail pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil coanti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disopro 245 mg (300 mg as a maleate)	ounts as three a	-	
			<i>y</i>
EMTRICITABINE – Special Authority see SA1651 on the previous Cap 200 mg		pnarmacy 30	✓ <u>Emtriva</u>
LAMIVUDINE – Special Authority see SA1651 on the previous p Tab 150 mg	•	irmacy 60	✓ <u>Lamivudine</u> <u>Alphapharm</u>

240 ml OP

100

200 ml OP

✓ 3TC

✓ Retrovir ✓ Retrovir

ZIDOVUDINE [AZT] - Special Authority see SA1651 on the previous page - Retail pharmacy Cap 100 mg.......152.25

	Subsidy (Manufacturer's Price) \$	F Subsid Per	ully ised	Brand or Generic Manufacturer
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see Note: zidovudine [AZT] with lamivudine (combination tablets the anti-retroviral Special Authority.	s) counts as two anti-r	etroviral me	dicat	ions for the purposes of
Tab 300 mg with lamivudine 150 mg	33.00	60	•	Alphapharm
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA1651 on p Cap 150 mg Cap 200 mg	141.68	rmacy 60 60		Teva Teva
DARUNAVIR – Special Authority see SA1651 on page 104 – Re Tab 400 mg Tab 600 mg	132.00	60 60	_	Darunavir Mylan Darunavir Mylan
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 Tab 100 mg with ritonavir 25 mg	1 0	pharmacy 60	<b>✓</b> <u>I</u>	<u>Lopinavir/Ritonavir</u> Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	<b>✓</b> <u>L</u>	Lopinavir/Ritonavir Mylan
Oral liq 80 mg with ritonavir 20 mg per ml	735.00 30	0 ml OP	<b>✓</b> ł	Kaletra
RITONAVIR – Special Authority see SA1651 on page 104 – Ret Tab 100 mg	, ,	30	<b>✓</b> <u>I</u>	Norvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA1651 on page 104 Tab 50 mg		30	<b>√</b> 1	Гіvісау
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 o Tab 400 mg	1,090.00	60		sentress
Tab 600 mg	1,090.00	60	<b>√</b>	sentress HD

## Immune Modulators

#### Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

#### **Criteria for Treatment**

- 1) Diagnosis
  - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
  - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
  - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

#### **Exclusion Criteria**

 Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).

Subsidy (Manufacturer's Price)	Sul	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10<sup>9</sup>) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

#### **Exit Criteria**

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline on the previous page
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.
- Inj 180 mcg prefilled syringe......500.00

✓ Pegasys

## ⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
  - 1.2 Patient has chronic hepatitis C and is co-infected with HIV: or
  - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

#### Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- . Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400.000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 Patient has chronic hepatitis C. genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
  - 3.1 Patient has responder relapsed; or
  - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
  - 3.1 Patient has responder relapsed; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 3.2 Patient was a partial responder; or
- 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
  - 5.1 HBeAg positive; or
  - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease: and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma\*; or
- 2 All of the following:
  - 2.1 Patient has a myeloproliferative disorder\*; and
  - 2.2 Patient is intolerant of hydroxyurea; and
  - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
  - 3.1 Patient has a myeloproliferative disorder; and
  - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
  - 3.1 Patient has a cutaneous T cell lymphoma\*; or
  - 3.2 Both:
    - 3.2.1 Patient has a myeloproliferative disorder\*; and
    - 3.2.2 Fither:

### **INFECTIONS - AGENTS FOR SYSTEMIC USE**

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

- 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
- 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with \* are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant\* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with \* are unapproved indications.

### **Urinary Tract Infections**

ME	THENAMINE (HEXAMINE) HIPPURATE			
*	Tab 1 g	40.01	100	✓ Hiprex
NIT	ROFURANTOIN			
*	Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
*	Tab 100 mg	37.50	100	✓ Nifuran
*	Cap modified-release 100 mg - Wastage claimable	86.40	100	✓ <u>Macrobid</u>
NO	RFLOXACIN			
	Tab 400 mg - Subsidy by endorsement	245.00	100	✓ Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
Anticholinesterases			
Anticholinesterases			
NEOSTIGMINE METILSULFATE			
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓ Max Health
PYRIDOSTIGMINE BROMIDE			
▲ Tab 60 mg	45 79	100	✓ Mestinon
_ 100 00 mg		100	<u> </u>
Non-Steroidal Anti-Inflammatory Drugs			
DICLOFENAC SODIUM	4.00		/ Distatores Constan
* Tab EC 25 mg		50	✓ <u>Diclofenac Sandoz</u>
* Tab 50 mg dispersible		20	✓ Voltaren D
* Tab EC 50 mg		50	✓ <u>Diclofenac Sandoz</u>
* Tab long-acting 75 mg		100	✓ Voltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a F		5	✓ Voltaren
* Suppos 12.5 mg		10	✓ Voltaren
* Suppos 25 mg		10	✓ Voltaren
* Suppos 50 mg – Up to 10 supp available on a PSO		10	✓ Voltaren
* Suppos 100 mg	7.00	10	✓ Voltaren
IBUPROFEN			
* Tab 200 mg	21.40	1,000	✓ Relieve
* Tab long-acting 800 mg	3.05	30	✓ Brufen SR
* Oral liq 20 mg per ml	2.25	200 ml	✓ Ethics
	11.29		<ul><li>Fenpaed 100 mg per</li></ul>
			5 ml \$29
KETOPROFEN			
* Cap long-acting 200 mg	12.07	28	✓ Oruvail SR
			0.0.0
MEFENAMIC ACID	4.05		
* Cap 250 mg		50	Danatan
	(9.16)	00	Ponstan
	0.50	20	Damatan
	(5.60)		Ponstan
NAPROXEN			
* Tab 250 mg	32.69	500	✓ <u>Noflam 250</u>
* Tab 500 mg	28.71	250	✓ Noflam 500
* Tab long-acting 750 mg	6.47	28	✓ Naprosyn SR 750
* Tab long-acting 1 g	8.62	28	✓ Naprosyn SR 1000
TENOXICAM			
* Tab 20 mg	9.15	100	✓ Tilcotil
* Inj 20 mg vial		1	✓ AFT
, ,			
NSAIDs Other			
CELECOXIB			
Cap 100 mg	5.80	60	✓ Celecoxib Pfizer
Cap 200 mg		30	✓ Celebrex
σαρ 200 mg	3.30	00	✓ Celecoxib Pfizer
	0.00		- OCICCOAID FIIZGI

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# (Manufacturer's Pi

### Topical Products for Joint and Muscular Pain

**CAPSAICIN** 

Crm 0.025% - Special Authority see SA1289 below - Retail

#### ⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

### **Antirheumatoid Agents**

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)\*, sarcoidosis (pulmonary and non-pulmonary)\*, and the prescription is endorsed accordingly.

Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a \* is an unapproved indication.

* Tab 200 mg	8.78	100	Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg		30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	✓ D-Penamine
Tab 250 mg		100	✓ D-Penamine

### **Drugs Affecting Bone Metabolism**

### **Alendronate for Osteoporosis**

* Tab 70 mg2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECALCIFEROL  * Tab 70 mg with colecalciferol 5,600 iu	4	✓ Fosamax Plus
Tub 70 mg with colocatolic of 5,000 ta	7	1 Countar i luo

### **Other Treatments**

ALENDOONATE CODILIM

DENOSUMAB − Special Authority see SA1777 below − Retail pharmacy
Inj 60 mg prefilled syringe.......326.00 1 ✓ Prolia

#### ⇒SA1777 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither
  - 2.1 The patient is female and postmenopausal; or
  - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
  - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

#### continued...

- (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
- 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

#### PAMIDRONATE DISODIUM

inj 3 mg per mi, 10 mi viai	27.53	1	Pamisoi
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	79.95	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA177	79 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	<ul><li>Evista</li></ul>

#### ⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or

Sub	bsidy F	ully Br	and or
(Manufactu	urer's Price) Subsidis	sed Ge	eneric
•	\$ Per	✓ Ma	anufacturer

#### continued...

- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

#### Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
   Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

#### 

#### **⇒SA1139** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

#### Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

**70I FDRONIC ACID** 

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

SA2110 below – Retail pharmacy .......60.00 100 ml OP 🗸 Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
  - 2.1 Bone or articular pain; or
  - 2.2 Bone deformity; or
  - 2.3 Bone, articular or neurological complications; or
  - 2.4 Asymptomatic disease, but risk of complications; or
  - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

**Initial application — (Underlying cause - glucocorticosteroid therapy)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
  - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
  - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
  - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note): or
  - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
  - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
  - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
  - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
  - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury\*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with \* are unapproved indications.

Renewal — (spinal cord injury\*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with \* are unapproved indications.

a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

### Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg		500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA196	3 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	<ul><li>Benzbromaron AL</li></ul>
			100 S29

#### **⇒SA1963** Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

### COLCHICINE

* Tab 500 mcg	6.00	100	Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Reta	il pharmacy		
Tab 80 mg	20.00	28	✓ <u>Febuxostat</u> multichem
Tab 120 mg	20.00	28	✓ <u>Febuxostat</u> multichem

#### ⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
  - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
  - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required

	MUS	CULOSKEL	ETAL SYSTEM
Subsid (Manufacturer \$	s Price)	Fully Subsidised Per	Brand or Generic Manufacturer
continued			
and serum urate remains greater than 0.36 mmol/l despite use of maximum tolerated dose; or			, ,,
2.3 The patient has renal impairment such that probenecid is contra			
remains greater than 0.36 mmol/l despite optimal treatment with 2.4 The patient has previously had an initial Special Authority appro			
Initial application — (Tumour lysis syndrome) only from a haematologist o			•
applications meeting the following criteria: Both:	J		
<ol> <li>Patient is scheduled to receive cancer therapy carrying an intermediate</li> <li>Patient has a documented history of allopurinol intolerance.</li> </ol>	or high ris	sk of tumour lys	sis syndrome; and
<b>Renewal</b> — <b>(Gout)</b> from any relevant practitioner. Approvals valid for 2 years patient is benefitting from treatment.	s where the	e treatment ren	nains appropriate and the
Renewal — (Tumour lysis syndrome) only from a haematologist or oncolog treatment remains appropriate and the patient is benefitting from treatment.	ist. Approv	vals valid for 6	weeks where the
Note: In chronic renal insufficiency, particularly when the glomerular filtration reflective. The efficacy and safety of febuxostat have not been fully evaluated in clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary impairment. Optimal treatment with allopurinol in patients with renal impairment clearance-adjusted dose of allopurinol then, if serum urate remains greater that allopurinol to 600 mg or the maximum tolerated dose.	in patients cessary in p nt is define	with severe repartients with medias treatment	nal impairment (creatinine ild or moderate renal to the creatinine
PROBENECID			
* Tab 500 mg	10	00 <b>~</b> P	robenecid-AFT
Muscle Relaxants			
BACLOFEN			
* Tab 10 mg4.20	10		acifen
Inj 0.05 mg per ml, 1 ml ampoule — Subsidy by endorsement11.55 Subsidised only for use in a programmable pump in patients where or caused intolerable side effects and the prescription is endorsed accor		_	ioresal Intrathecal e been ineffective or have
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement306.82	uirigiy. 5	5 <b>✓</b> <u>N</u>	ledsurge

#### caused intolerable side effects and the prescription is endorsed accordingly. DANITROI ENE

Cap 25 mg	97.50	100	✓ Dantrium ✓ Dantrium S29 S29
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE  Tab 100 mg	20.76	100	✓ Norflex

Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have

AMANTA DINIC LIVEDOCLII ODIDE

LEVODOPA WITH CARBIDOPA

**RASAGII INF** 

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised Per Manufacturer

### **Agents for Parkinsonism and Related Disorders**

### **Dopamine Agonists and Related Agents**

24 (	60 🗸	Symmetrel
50	5	Movapo
34	5 🗸	Movapo
		to 1 March 2021 and the here there exists a record of
70 3	30 <b>•</b>	Parlodel S29
age 241 for	details	
04 1	00 🗸	<u>Comtan</u>
25 1	00 🗸	Madopar Rapid
75 1	00 🗸	Madopar 62.5
30 1	00 🗸	Madopar 125
	00 🗸	Madopar HBS
	mocriptine m rescription as 70 3 3 1 2 2 5 1 1 7 5 1 1 3 0 1 1	50 5 7 34 5 7 mocriptine mesylate prior rescription as endorsed w 70 30 7 page 241 for details 100 7 25 100 7 30 7 30 7

*	Tab 100 mg with carbidopa 25 mg	21.11	100
*	Tab long-acting 200 mg with carbidopa 50 mg	43.65	100
*	Tab 250 mg with carbidopa 25 mg	38.39	100
PR	AMIPEXOLE HYDROCHLORIDE		

✓ Sinemet ▲ Tab 0.25 mg ......6.12 100 ✓ Ramipex ✓ Ramipex ▲ Tab 1 mg .......20.73 100

> 84 84

100

* Tab 1 mg	53.50	30
ROPINIROLE HYDROCHLORIDE		
▲ Tab 0.25 mg	2.85	84
	0.00	400

Tab 0.25 mg2.85	84	
3.39	100	
Tab 1 mg	84	
4.70	100	

v	<u>Kopin</u>
✓	Mylan S29
✓	Ropin

✓ Azilect \$29

✓ Madopar 250

Sinemet ✓ Sinemet CR

$\blacktriangle$	Tab 2 mg	5.48
	Tab 5 mg	
SF	I FGII INF HYDROCHI ORIDF – Subsidy by endorsement	

1	Mylan S29
✓	Ropin
1	Ropin

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the
prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of
prior dispensing of selegiline hydrochloride.

*	Tab 5 mg	48.00	100	✓ Eldepryl S29
TO	LCAPONE			
$\blacktriangle$	Tab 100 mg	152.38	100	Tasmar

	0		F. II.	Durantan
	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
	\$	Per	•	Manufacturer
Anticholinergics				
BENZATROPINE MESYLATE				
Tab 2 mg	9.59	60	1	Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	1	<u>Phebra</u>
<ul> <li>a) Up to 10 inj available on a PSO</li> </ul>				
b) Only on a PSO				
PROCYCLIDINE HYDROCHLORIDE			_	
Tab 5 mg	7.40	100	/	Kemadrin
Agents for Essential Tremor, Chorea and Relat	ted Disorders			
RILUZOLE - Special Authority see SA1403 below - Retail pha	rmacv			
Wastage claimable	,			
Tab 50 mg	130.00	56	✓	Rilutek
⇒SA1403 Special Authority for Subsidy				
Initial application only from a neurologist or respiratory special	ist. Approvals valid fo	r 6 mc	onths for a	pplications meeting the
following criteria:				
All of the following:				
1 The patient has amyotrophic lateral sclerosis with diseas				
2 The patient has at least 60 percent of predicted forced vi	tal capacity within 2 m	onths	prior to the	e initial application; and
<ul><li>3 The patient has not undergone a tracheostomy; and</li><li>4 The patient has not experienced respiratory failure; and</li></ul>				
5 Any of the following:				
5.1 The patient is ambulatory; or				
5.2 The patient is able to use upper limbs; or				
5.3 The patient is able to swallow.				
Renewal from any relevant practitioner. Approvals valid for 18 All of the following:	months for applications	s mee	ting the fol	llowing criteria:
The patient has not undergone a tracheostomy; and				
2 The patient has not experienced respiratory failure; and				
3 Any of the following:				
3.1 The patient is ambulatory; or				
3.2 The patient is able to use upper limbs; or				
3.3 The patient is able to swallow.				
TETRABENAZINE				
Tab 25 mg	91.10	112	✓	Motetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE]				
Gel 2%, tube - Subsidy by endorsement	14.50	30 ml	1	Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO				
b) Subsidised only if prescribed for urethral or cervical			•	٠,
Gel 2%, 11 ml urethral syringe – Subsidy by endorsement.	42.00	10	•	Instillagel Lido
a) Up to 5 each available on a PSO	vental administration	onel H		tion is and are
b) Subsidised only if prescribed for urethral, cervical or	rectal administration a	and th	e prescrip	lion is endorsed

accordingly.

	Subsidy (Manufacturer's Price	) Sı	Fully ubsidised	
	\$	Per	1	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	•	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	1	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	/	Lidocaine-Baxter
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO		5	/	Lidocaine-Claris
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO	6.45	5	1	<u>Lidocaine-Baxter</u>
			1	Lidocaine-Claris
(Lidocaine-Claris Inj 2%, 20 ml vial to be delisted 1 June 2022)				
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement	103.32	10	1	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical a	administration and th	e prescri	ption is	endorsed accordingly.

### **Topical Local Anaesthetics**

#### ⇒SA0906 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] - Special Authority se	ee SA0906 above – Retail pharr	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE -	- Special Authority see SA0906	above - Reta	ail pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

### **Analgesics**

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid	<b>Analgesics</b>
------------	-------------------

ASPIRIN  * Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or dial accordingly.	betic periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream S29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

	<u> </u>	VETTV OOD OTOTEW
		Fully Brand or
	(Manufacturer's Price) Subsid	lised Generic  ✓ Manufacturer
PARACETAMOL	¥ 101	
Tab 500 mg - blister pack	19.75 1,000	✓ Pacimol
a) Maximum of 300 tab per prescription; can b		<u>r dominor</u>
b) Up to 30 tab available on a PSO	•	
c)		per l
Subsidy by endorsement for higher queries the regular daily design for one month or an end of the regular daily daily design for one month or an end of the regular daily d	antities is available for patients with long t reater, and the prescription is annotated a	
	where dispensing history supports a long-	
2) Maximum of 100 tab per dispensing for		
, , , , ,	nse in repeat dispensings not exceeding	100 tab per dispensing.
Tab 500 mg - bottle pack - Maximum of 300 tab pe		
prescription; can be waived by endorsement	17.92 1,000	✓ <u>Noumed</u>
		<u>Paracetamol</u>
Subsidy by endorsement for higher quant  doily design for one month or greater, and		
daily dosing for one month or greater, and prescription as endorsed where dispensin	1 1	. Pharmacists may annotate th
Maximum of 100 tab per dispensing for no	, , , ,	hed for more than 100 tabs (for
non-endorsed patients), then dispense in		•
2 рания другий и други		F
* Oral liq 120 mg per 5 ml	5.45 1,000 ml	✓ Paracare
a) Up to 200 ml available on a PSO		
b) Not in combination		_
* Oral liq 250 mg per 5 ml	6.25 1,000 ml	✓ Paracare Double
) II + 400 I - 11 II - B00		<u>Strength</u>
<ul><li>a) Up to 100 ml available on a PSO</li><li>b) Not in combination</li></ul>		
* Suppos 125 mg	3.59 10	✓ Gacet
* Suppos 250 mg		✓ Gacet
* Suppos 500 mg		✓ Gacet
., .		
Opioid Analgesics		
CODEINE PHOSPHATE - Safety medicine; prescriber	may determine dispensing frequency	
Tab 15 mg		✓ PSM
Tab 30 mg		✓ PSM
Tab 60 mg	14.25 100	✓ PSM
DIHYDROCODEINE TARTRATE		
Tab long-acting 60 mg	8.60 60	✓ DHC Continus
FENTANYL		
a) Only on a controlled drug form		
b) No patient co-payment payable		
<ul> <li>c) Safety medicine; prescriber may determine dispersion</li> </ul>	. ,	
Inj 50 mcg per ml, 2 ml ampoule		Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule		Boucher and Muir
Patch 12.5 mcg per hour		✓ Fentanyl Sandoz
Patch 50 mag per hour		✓ <u>Fentanyl Sandoz</u> ✓ Fentanyl Sandoz
Patch 50 mcg per hour Patch 75 mcg per hour		✓ Fentanyi Sandoz ✓ Fentanyi Sandoz
Patch 100 mcg per hour		✓ Fentanyl Sandoz
r aton 100 may per noul	10.00	- I citaliyi Januoz

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	quency			
d) Extemporaneously compounded methadone will only be r	eimbursed at the rat	e of th	ne cheapes	t form available
(methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard Fo	ormulae, page 243			
Tab 5 mg	1.40	10	✓	Methatabs
Oral liq 2 mg per ml	6.40	200 m		<u>Biodone</u>
Oral liq 5 mg per ml		200 m		Biodone Forte
Oral liq 10 mg per ml		200 m		Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	✓.	AFT
ORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	quency			
Oral liq 1 mg per ml	9.28	200 m	nl 🗸 l	RA-Morph
Oral liq 2 mg per ml	16.24	200 m	nl 🗸 l	RA-Morph
Oral liq 5 mg per ml	19.44	200 m		Ordine \$29
,			✓	RA-Morph
Oral liq 10 mg per ml	27.74	200 m		Ordine S29
1 01			✓	RA-Morph
MORPHINE SULPHATE				•
a) Only on a controlled drug form				
b) No patient co-payment payable				
<ul><li>c) Safety medicine; prescriber may determine dispensing fre</li></ul>	guency			
Tab immediate-release 10 mg		10	1	Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10	✓	m-Eslon
Cap long-acting 60 mg	6.12	10	✓	m-Eslon
Cap long-acting 100 mg	7.13	10	1	m-Eslon
Inj 5 mg per ml, 1 ml ampoule - Up to 5 inj available on a PS	6.99 io	5	1	DBL Morphine
				Sulphate

Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO ......5.61

Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO ......7.08

Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO ......7.28

5

5

5

✓ DBL Morphine Sulphate

**✓** DBL Morphine

Sulphate

DBL Morphine

Sulphate

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	` \$	Per	•	Manufacturer
YCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	equency			
Tab controlled-release 5 mg		20	/	Oxycodone Sando
Ç	3.01	28	✓	Oxycodone Sando
				<b>S29</b> S29
Oxycodone Sandoz to be Principal Supply on 1 June 20	22			
Tab controlled-release 10 mg		20	1	Oxycodone Sando
<b>g</b>	3.23	30		Oxycodone Sando
				S29 S29
	5.38	50	1	Oxycodone Sando
	0.00	•		S29 S29
	10.75	100	1	Oxycodone Sando
	10.75	100	•	S29 S29
	44.50	00		
Oversedens Condo- to be Dringing County on 1 June 00	11.50	28	•	OxyContin
Oxycodone Sandoz to be Principal Supply on 1 June 20		20	./	Oxycodone Sando
Tab controlled-release 20 mg	5.38	50		Oxycodone Sando
	5.56	50	•	S29 S29
	10.75	100		
	10.75	100	•	Oxycodone Sando
			_	S29 S29
	13.25	28	•	OxyContin
Oxycodone Sandoz to be Principal Supply on 1 June 20			,	
Tab controlled-release 40 mg		20	•	Oxycodone Sando
Oxycodone Sandoz to be Principal Supply on 1 June 20.		20		O dana Canda
Tab controlled-release 80 mg Oxycodone Sandoz to be Principal Supply on 1 June 20		20	•	Oxycodone Sando
Cap immediate-release 5 mg		20	1	OxyNorm
Cap immediate-release 3 mg		20		OxyNorm
Cap immediate-release 20 mg		20		OxyNorm
Oral lig 5 mg per 5 ml		250 m		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		Hameln
,g po, apoa.o	7.28	Ū		OxyNorm
Hameln to be Principal Supply on 1 July 2022	•			. ,
Inj 10 mg per ml, 2 ml ampoule	11.49	5	1	Hameln
, 31 ,	14.36	-		OxyNorm
Hameln to be Principal Supply on 1 July 2022				•
Inj 50 mg per ml, 1 ml ampoule	22.92	5	✓	Hameln
	30.60		/	OxyNorm

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	Sul	osidised	Generic
	\$	Per	J0101000	Manufacturer
		. 0.		manada o
Hameln to be Principal Supply on 1 July 2022				
(Oxycodone Sandoz S29 S29 Tab controlled-release 5 mg to be	delisted 1 June 202	22)		
(Oxycodone Sandoz S29 S29 Tab controlled-release 10 mg to b	e delisted 1 June 20	122)		
(Oxycodone Sandoz S29 S29 Tab controlled-release 10 mg to b				
, ,		,		
(Oxycodone Sandoz S29 S29 Tab controlled-release 10 mg to b		)22)		
(OxyContin Tab controlled-release 10 mg to be delisted 1 June 20	)22)			
(Oxycodone Sandoz S29 S29 Tab controlled-release 20 mg to b	e delisted 1 June 20	022)		
(Oxycodone Sandoz S29 S29 Tab controlled-release 20 mg to b				
(OxyContin Tab controlled-release 20 mg to be delisted 1 June 20		,,		
(OxyNorm Inj 10 mg per ml, 1 ml ampoule to be delisted 1 July 20				
(OxyNorm Inj 10 mg per ml, 2 ml ampoule to be delisted 1 July 20				
(OxyNorm Inj 50 mg per ml, 1 ml ampoule to be delisted 1 July 20	122)			
PARACETAMOL WITH CODEINE - Safety medicine; prescriber	may determine disc	ensina fre	eauencv	
* Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
Tab paracolamor oco mg war ocacino pricopriate e mg		1,000		Codeine (Relieve)
				oodelile (Helleve)
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	auencv			
Tab 50 mg		10	<b>✓</b> P	SM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a P	SO 29.88	5	_	OBL Pethidine
ing 50 mg per mi, 1 mi ampoule op to 5 mg available on a r	0020.00	J		Hydrochloride
1:50	00 00 70	_		•
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a P	SO30.72	5	✓ L	DBL Pethidine
				Hydrochloride
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	<b>√</b> T	ramal SR 100
Tab sustained-release 150 mg		20	_	ramal SR 150
Tab sustained-release 200 mg		20	_	ramal SR 200
Cap 50 mg		100	_	Arrow-Tramadol
Cap 50 mg	2.00	100	· <u>-</u>	ATTOW-TTAITIAUUI
Autidonycoconto				
Antidepressants				
Cyclic and Related Agents				
AMITRIPTYLINE - Safety medicine; prescriber may determine di			_	
Tab 10 mg	2.49	100		Arrow-Amitriptyline
Tab 25 mg	1.51	100	✓ <u>A</u>	Arrow-Amitriptyline
Tab 50 mg	2.51	100	<b>✓</b> A	Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE - Safety medicine; prescri	har may datarmina	dienonein	a fraguer	nov
Tab 10 mg		30		Clomipramine Teva
Tab 25 mg	11.99	30	• [	Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by en-	dorsement			
a) Safety medicine; prescriber may determine dispensing fre	quency			
b) Subsidy by endorsement – Subsidised for patients who we		[dothiani	nl hydro	chloride prior to 1 lune
2019 and the prescription is endorsed accordingly. Pharr				
		e ine pre	scription	as endoised where there
exists a record of prior dispensing of dosulepin [dothiepin]		00	, -	Naardanin Meise
Tab 75 mg		30		Oosulepin Mylan
Cap 25 mg	7.83	50	✓ [	)osulepin
				Mylan S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic  Manufacturer
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine dispe	nsing	g frequency
Tab 10 mg	5.48	50	✓ Tofranil
	10.96	100	
Tab 25 mg		50	✓ Tofranil
NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; presci			
Tab 10 mg		100	
Tab 25 mg	5.98	180	✓ <u>Norpress</u>
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	elective		
TRANYLCYPROMINE SULPHATE			
Tab 10 mg	12.85	28	✓ Parnate S29 S29
	22.94	50	✓ Parnate
	45.88	100	✓ Parnate S29 S29
	96.00		✓ Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg	11.80	60	✓ Aurorix
* Tab 300 mg		60	✓ Aurorix
			- Marotix
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	1.91	84	✓ PSM Citalopram
ESCITALOPRAM			_
* Tab 10 mg	1.07	28	✓ <u>Escitalopram</u>
W. Tab 00	1.00	00	(Ethics)
* Tab 20 mg	1.92	28	✓ <u>Escitalopram</u> (Ethics)
FLUOXETINE HYDROCHLORIDE			(Etillos)
* Tab dispersible 20 mg, scored – Subsidy by endorsement	1.98	30	✓ Fluox
Subsidised by endorsement		•	
When prescribed for a patient who cannot swallow accordingly; or	whole tablets or caps	sules	and the prescription is endorsed
<ul><li>When prescribed in a daily dose that is not a multiple</li></ul>	ole of 20 mg in which	case	the prescription is deemed to be
endorsed. Note: Tablets should be combined with			
000	0.04		
Cap 20 mg	2.91	84	✓ Fluox
PAROXETINE	0.04	00	./ Lavamina
* Tab 20 mg	3.01	90	✓ <u>Loxamine</u>
SERTRALINE No. Tal. 50 and	0.00	00	1 Octobria
* Tab 50 mg	0.92	30	<ul> <li>✓ <u>Setrona</u></li> <li>✓ Setrona AU</li> </ul>
* Tab 100 mg	1.61	30	✓ Setrona AU ✓ Setrona
	••••••		✓ Setrona AU

	Subsidy (Manufacturer's Price	) Subs	Fully Brand sidised Gene	
	\$	Per		facturer
Other Antidepressants				
MIRTAZAPINE				
Tab 30 mg		28	Noumed	-
Tab 45 mg	3.45	28	✓ <u>Noumed</u>	<u>d</u>
VENLAFAXINE	6.00	84	✓ Enlafax	VD
* Cap 75 mg		84	✓ Enlarax	
* Cap 150 mg		84	✓ Enlafax	
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
DIAZEPAM – Safety medicine; prescriber may determine dispen	ising frequency			
Inj 5 mg per ml, 2 ml ampoule - Subsidy by endorsement	. ,	5	✓ Hospira	I
a) Up to 5 inj available on a PSO				
b) Only on a PSO	"			
<ul> <li>c) PSO must be endorsed "not for anaesthetic procedules."</li> <li>Rectal tubes 5 mg – Up to 5 tube available on a PSO</li></ul>		5	✓ Stesolic	ı
PHENYTOIN SODIUM		3	• Otesone	•
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a				
PSO	104.58	5	✓ Hospira	
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a				
PSO	154.01	5	Hospira	I
Control of Epilepsy				
CARBAMAZEPINE				
* Tab 200 mg		100	Tegreto	
* Tab long-acting 200 mg		100	✓ Tegreto	
* Tab long acting 400 mg		100 100	✓ Tegreto	
* Tab long-acting 400 mg * Oral liq 20 mg per ml		250 ml	<ul><li>✓ Tegreto</li><li>✓ Tegreto</li></ul>	
CLOBAZAM – Safety medicine; prescriber may determine dispe		230 1111	• regicto	•
Tab 10 mg	. ,	50	✓ Frisium	
CLONAZEPAM – Safety medicine; prescriber may determine dis		00	- Triolani	
Oral drops 2.5 mg per ml		0 ml OP	✓ Rivotril	
ETHOSUXIMIDE		· · ·		
Cap 250 mg	140.88	100	✓ Zarontir	1
Oral liq 250 mg per 5 ml		200 ml	✓ Zarontir	-
GABAPENTIN				
Note: Not subsidised in combination with subsidised pregab	alin			
* Cap 100 mg		100	✓ Nupenti	_
* Cap 300 mg		100	✓ Nupenti	
* Cap 400 mg	10.26	100	✓ Nupenti	<u>n</u>

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
ACOSAMIDE - Special Authority see SA1125 below	- Retail pharmacy			
▲ Tab 50 mg	25.04	14	✓ V	impat
▲ Tab 100 mg	50.06	14	✓ V	impat
· ·	200.24	56	✓ V	impat
Tab 150 mg	75.10	14	✓ V	impat
Ŭ	300.40	56	✓ V	impat
▲ Tab 200 mg	400.55	56		impat

### **⇒SA1125** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

**Renewal** from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

#### LAMOTRIGINE

LA	MOTRIGINE			
$\blacktriangle$	Tab dispersible 2 mg	55.00	30	✓ Lamictal
$\blacktriangle$	Tab dispersible 5 mg	50.00	30	✓ Lamictal
*	Tab dispersible 25 mg	2.76	56	✓ Logem
*	Tab dispersible 50 mg	3.31	56	Logem
*	Tab dispersible 100 mg		56	Logem
۱F	VETIRACETAM			<del></del>
	Tab 250 mg	4 99	60	✓ Everet
	Tab 500 mg		60	✓ Everet
	Tab 750 mg		60	✓ Everet
	Tab 1,000 mg		60	✓ Everet
	Oral liq 100 mg per ml		300 ml OP	✓ Levetiracetam-AFT
PH	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formul	ae, page 243		
*	Tab 15 mg		500	✓ PSM
*	Tab 30 mg		500	✓ PSM
DI I	ENYTOIN SODIUM			
*		75.00	200	✓ Dilantin Infatab
不	Tab 50 mg		200	✓ Dilantin
	Cap 100 mg		200	✓ Dilantin
*	Cap 100 mg		200 500 ml	✓ Dilantin
-	Oral liq 30 mg per 5 ml	22.03	300 1111	Dilantin
PR	EGABALIN			
	Note: Not subsidised in combination with subsidised	• •		
	Cap 25 mg		56	✓ Pregabalin Pfizer
*			56	Pregabalin Pfizer
	Cap 150 mg	4.01	56	✓ Lyrica
				✓ Pregabalin Pfizer
	Cap 300 mg	7.38	56	Pregabalin Pfizer

### **NERVOUS SYSTEM**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PRIMIDONE  * Tab 250 mg	37.35	100	/	Apo-Primidone
(Apo-Primidone Tab 250 mg to be delisted 1 January 2023)			•	Primidone Clinect
SODIUM VALPROATE				
Tab 100 mg	13.65	100	1	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC	52.24	100	✓	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 m	nl 🗸	Epilim S/F Liquid
			✓	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail pl	harmacy			
Cap 250 mg	509.29	60	✓	Diacomit S29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit S29

#### ⇒SA1330 Special Authority for Subsidy

TODIRAMATE

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

**Renewal** from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

▲ Tab 25 mg11.07	60	✓ Arrow-Topiramate
v		✓ Topiramate Actavis
26.04		✓ Topamax
Tab 50 mg18.81	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
44.26		✓ Topamax
Tab 100 mg31.99	60	✓ Arrow-Topiramate
•		✓ Topiramate Actavis
75.25		✓ Topamax
Tab 200 mg55.19	60	✓ Arrow-Topiramate
•		✓ Topiramate Actavis
129.85		✓ Topamax
Sprinkle cap 15 mg20.84	60	✓ Topamax
Sprinkle cap 25 mg26.04	60	✓ Topamax
GABATRIN – Special Authority see SA2088 below – Retail pharmacy		
Tab 500 mg119.30	100	✓ Sabril

#### ⇒SA2088 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
  - 1.1 Patient has infantile spasms; or
  - 1.2 Both:
    - 1.2.1 Patient has epilepsy; and

Subsidy	S	Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

#### 1.2.2 Either:

- 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
- 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
  - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

### Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Tab orodispersible 10 mg3.65	5 30	✓ Rizamelt
SUMATRIPTAN		
Tab 50 mg14.41	90	✓ Sumagran
Tab 100 mg22.68	90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per		_
prescription34.00	2 OP	✓ Imigran
·		

### **Prophylaxis of Migraine**

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

**PIZOTIFEN** 

\* Tab 500 mcg......23.21 100 ✓ Sandomigran

### Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 on the next page - Retail pharmacy

3 OP

✓ Emend Tri-Pack

Subsi	dy Fully Brand	or
(Manufacture	er's Price) Subsidised Gener	ric
· · · · · · · · · · · · · · · · · · ·	Por / Manus	facturor

#### ⇒SA0987 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

#### BETAHISTINE DIHYDROCHLORIDE

4.62 100 ✓ <u>Serc</u> (Vergo 16 Tab 16 mg to be delisted 1 July 2022)	
CYCLIZINE HYDROCHLORIDE	
Tab 50 mg0.49 10 ✓ <u>Nausicalm</u>	
CYCLIZINE LACTATE	
Inj 50 mg per ml, 1 ml21.53 10 <b>✓ <u>Hamein</u></b>	
DOMPERIDONE	
<b>※</b> Tab 10 mg2.85 100 ✓ <u>Pharmacy Hea</u>	<u>lth</u>
HYOSCINE HYDROBROMIDE	
★ Inj 400 mcg per ml, 1 ml ampoule	)
Patch 1.5 mg - Special Authority see SA1998 below - Retail	
pharmacy	S

#### ⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

**Renewal** from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

#### METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO9.50	10	✓ <u>Pfizer</u>
ON	DANSETRON		
*	Tab 4 mg2.68	50	✓ Onrex
*	Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ Ondansetron ODT-DRLA
*	Tab 8 mg4.57	50	✓ Onrex
*	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ Ondansetron ODT-DRLA
PR	OCHLORPERAZINE		
*	Tab 3 mg buccal5.97	50	
	(30.00)		Buccastem
*	Tab 5 mg - Up to 30 tab available on a PSO8.00	250	✓ Nausafix
*	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

## **Antipsychotics**

### General

AMISULPRIDE - Safety medicine; prescriber may determine disper		•	
Tab 100 mg		30	✓ <u>Sulprix</u>
	17.16	100	Amisulpride
			Mylan S29
Tab 200 mg	14.96	60	✓ Sulprix
Tab 400 mg	29.78	60	✓ Sulprix
(Amisulpride Mylan S29) Tab 100 mg to be delisted 1 August 2022)			<del></del>
ARIPIPRAZOLE – Safety medicine; prescriber may determine dispe		201	
Tab 5 mg		30	✓ Aripiprazole Sandoz
· · · · · · · · · · · · · · · · · · ·		30	✓ Aripiprazole Sandoz
Tab 10 mg			
Tab 15 mg		30	✓ Aripiprazole Sandoz
Tab 20 mg		30	✓ Aripiprazole Sandoz
Tab 30 mg		30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; presc		ermine dispen	sing frequency
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	30.79	10	✓ Largactil
CLOZAPINE - Hospital pharmacy [HP4]			
Safety medicine; prescriber may determine dispensing frequency	cv		
Tab 25 mg	•	50	✓ Clopine
			✓ Clozaril
	13.37	100	✓ Clopine
			✓ Clozaril
Tab 50 mg	8.67	50	✓ Clopine
J	17.33	100	✓ Clopine
Tab 100 mg	17.33	50	✓ Clopine
<b>v</b>			✓ Clozaril
	34.65	100	✓ Clopine
			✓ Clozaril
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml	67.62	100 ml	✓ Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine dispe	nsina fraguer	nev	
Tab 500 mcg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg - Up to 30 tab available on a PSO		100	✓ Serenace
Tab 5 mg — Up to 30 tab available on a PSO		50	✓ Serenace
Tab 5 mg Op to 50 tab available on a 1 50	29.72	100	✓ Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 ml	✓ Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO		100 1111	✓ Serenace
, ,			- <del>Octoriado</del>
LEVOMEPROMAZINE – Safety medicine; prescriber may determin			/ Namhan (Outer)
Tab 25 mg (33.8 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 25 mg as a maleate		100	✓ <u>Nozinan</u>
Tab 100 mg (135 mg as a maleate)		100	✓ Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	✓ <u>Nozinan</u>

	Subsidy		Fully	
	(Manufacturer's Price) \$	Per	Subsidised	I Generic Manufacturer
EVOMEPROMAZINE HYDROCHLORIDE – Safety medicine	o: prescriber may determ	ina d	ienoneina	
Inj 25 mg per ml, 1 ml ampoule		10		Nozinan
ITHIUM CARBONATE - Safety medicine; prescriber may de		. •		
		100		Priadel
Tab long-acting 400 mg		100	_	Douglas
Cap 250 mg		100	•	Douglas
LANZAPINE – Safety medicine; prescriber may determine d				
Tab 2.5 mg		28		Zypine
Tab 5 mg		28		Zypine
Tab orodispersible 5 mg		28		Zypine ODT
Tab 10 mg		28		Zypine
Tab orodispersible 10 mg	2.38	28	•	Zypine ODT
ERICYAZINE - Safety medicine; prescriber may determine	dispensing frequency			
Tab 2.5 mg	10.49	84	✓	Neulactil
	12.49	100	•	Neulactil
Tab 10 mg	37.34	84	1	Neulactil
-	44.45	100	1	Neulactil
UETIAPINE - Safety medicine; prescriber may determine di	spensing frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	_	Quetapel
•		00		<u>uuotupoi</u>
ISPERIDONE – Safety medicine; prescriber may determine		60	./	Dianavidana (Taya)
Tab 0.5 mg		60		Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60	_	Risperidone (Teva)
Oral liq 1 mg per ml		30 ml	•	Risperon
IPRASIDONE - Safety medicine; prescriber may determine			_	
Cap 20 mg		60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg	46.55	60	/	Zusdone
UCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; p	rescriber may determine	e disp	ensing fre	equency
Tab 10 mg	31.45	100	<b>'</b>	Clopixol
				•
Depot Injections				
LUDENTUYOU DECANOATE Cofety modified and the		to a f		
LUPENTHIXOL DECANOATE – Safety medicine; prescriber		•		Fluencel
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO		5		Fluanxol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO		5		Fluanxol
ALOPERIDOL DECANOATE - Safety medicine; prescriber	•	•		
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	•	Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	•	<b>Haldol Concentrate</b>
			✓	Haldol
				Decanoas \$29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
OLANZAPINE – Special Authority see SA1428 below – Retail p Safety medicine; prescriber may determine dispensing frequ				
Inj 210 mg vial	,	1	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	1	Zyprexa Relprevv
Inj 405 mg vial	504.00	1	•	Zyprexa Relprevv
SA1/28 Special Authority for Subsidy				

#### SA1428 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia; and
  - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

### PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensi	ng trequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna
			=

#### ⇒SA1429 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

### RISPERIDONE - Special Authority see SA1427 on the next page - Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency	Jency		
Inj 25 mg vial	135.98	1	Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial		1	✓ Risperdal Consta



Subsidy (Manufacturer's Price)	Subsid	Fully dised	Brand or Generic
 \$	Per	•	Manufacturer

### **⇒SA1427** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
  - 2.1 The patient has schizophrenia or other psychotic disorder; and
  - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
  - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 **Clopixol** 

### **Anxiolytics**

BUSPIRONE HYDROCHLORIDE  * Tab 5 mg  * Tab 10 mg		100 100	✓ Buspirone Viatris ✓ Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine	dispensing frequency	,	
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine d	ispensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ Ativan

### **Multiple Sclerosis Treatments**

#### **⇒SA2051** Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
  - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not

			NEF	RVOUS SYSTEM
		Subsidy	Fully	Brand or
		(Manufacturer's Price) \$	Subsidised Per 🗸	Generic Manufacturer
continued				
	necessarily have been seen by them during the re	elapse, but the neurolo	gist/physician m	ust be satisfied that the
5.2	clinical features were characteristic); and Each significant relapse is associated with characteristic	taristic naw symptom(	el/eian(e) or eub	stantially worsening of
5.2	previously experienced symptoms(s)/sign(s); and	densite new symptom	s)/sign(s) or sub	startially worsering or
5.3	Each significant relapse has lasted at least one w	eek and has started at	least one month	after the onset of a
	previous relapse; and			
5.4	Each significant relapse can be distinguished from	n the effects of genera	I fatigue; and is i	not associated with a fever
	(T> 37.5°C); and			
5.5	Either:			***
	5.5.1 Each significant relapse is severe enough	•	DSS or at least	one of the Kurtze
	Functional System scores by at least 1 poi 5.5.2 Each significant relapse is a recurrent parc		ultinla eclaracie	Itonic caizurae/enaeme
	trigeminal neuralgia, Lhermitte's symptom)		iditipie scierosis	(toriic scizures/spasiris,
6 Evider	nce of new inflammatory activity on an MR scan wit		s: and	
	f the following:		-,	
7.1	A sign of that new inflammatory activity is a gadol	inium enhancing lesior	n; or	
	A sign of that new inflammatory activity is a lesion			
	A sign of that new inflammatory is a T2 lesion with			
7.4	A sign of that new inflammatory activity is a promi		arly is responsib	le for the clinical features of
7.5	a recent relapse that occurred within the last 2 years A sign of that new inflammatory activity is new T2		a a provious MD	coon
	rumab can only be dispensed from a pharmacy req	•	•	
	he supplier. Treatment on two or more funded mul			
	Multiple sclerosis) only from a neurologist or ger			
had an EDSS	score of 0 to 6.0 (inclusive) with or without the use	e of unilateral or bilater	al aids at any tin	ne in the last six months
	ent has walked 100 metres or more with or without			
	rumab can only be dispensed from a pharmacy reg			
. ,	he supplier. Treatment on two or more funded mul	•		sly is not permitted.
	TUMARATE – Special Authority see SA2051 on the	e previous page – Reta	ail pharmacy	
	age claimable	da tua atua auta alue di e		: 1.1 a al
	Treatment on two or more funded multiple scleros			rmitted. <b>Tecfidera</b>
Cap 120	· ·			reclidera Footidera

Cap 240 mg	2,000.00	56	✓ Tecfidera
FINGOLIMOD - Special Authority see SA2051 on the	previous page - Retail pharr	nacy	
a) Wastage claimable			
b) Note: Treatment on two or more funded multip		taneously is	•
Cap 0.5 mg	2,200.00	28	<ul><li>Gilenya</li></ul>
GLATIRAMER ACETATE - Special Authority see SA2			,
Note: Treatment on two or more funded multiple s		eously is no	' -
Inj 40 mg prefilled syringe	1,137.48	12	Copaxone
INTERFERON BETA-1-ALPHA - Special Authority se		•	, ,
Note: Treatment on two or more funded multiple s		,	•
Inj 6 million iu prefilled syringe	· · · · · · · · · · · · · · · · · · ·	4	Avonex
Injection 6 million iu per 0.5 ml pen injector	1,170.00	4	Avonex Pen
INTERFERON BETA-1-BETA – Special Authority see			
Note: Treatment on two or more funded multiple s		,	' -
Inj 8 million iu per 1 ml	1,322.89	15	Betaferon



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer NATALIZUMAB - Special Authority see SA2051 on page 134 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Tvsabri OCRELIZUMAB - Special Authority see SA2051 on page 134 - Retail pharmacy Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. ✓ Ocrevus TERIFLUNOMIDE - Special Authority see SA2051 on page 134 - Retail pharmacy a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted. 28 ✓ Aubagio Sedatives and Hypnotics MELATONIN - Special Authority see SA1666 below - Retail pharmacy Brand switch fee payable (Pharmacode 2634112) - see page 241 for details Tab modified-release 2 mg - No more than 5 tab per day ......11.50 30 Vigisom

#### ⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)\*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under\*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient is aged 18 years or under\*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with \* are unapproved indications.

MIDAZOLAM - Safety medicine; prescriber may determine dispe	nsing frequency		
Inj 1 mg per ml, 5 ml ampoule	5.50	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	17.28	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for stat	us epilepticu	is use only.
Inj 5 mg per ml, 3 ml ampoule	4.50	5	<ul><li>Midazolam-Baxter</li></ul>
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available	on		
a PSO	13.09	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for stat	us epilepticu	is use only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 o	n the next page -	- Retail phar	macy
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓ Max Health S29

Subsidy (Manufacturer's Price)	Subsi	Fully dised	Brand or Generic	
\$	Per	•	Manufacturer	

#### ⇒SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM - Safety medicine; prescriber may determine of	dispensing frequency		
Tab 10 mg	1.33	25	✓ Normison
TRIAZOLAM - Safety medicine; prescriber may determine di	ispensing frequency		
Tab 125 mcg	5.10	100	
	(9.85)		Hypam
Tab 250 mcg	4.10	100	
,	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine di	ispensing frequency		
Tab 7.5 mg	10.80	500	✓ Zopiclone Actavis

### **Stimulants/ADHD Treatments**

**ATOMOXETINE** 

Cap 10 mg	18.41	28	✓ APO- Atomoxetine S29
	107.03		✓ Generic Partners ✓ Strattera
Cap 18 mg		28	✓ Generic Partners
	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ Generic Partners
Cap 40 mg		28	✓ Generic Partners
	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ APO-
			Atomoxetine S29
			✓ Generic Partners
Cap 80 mg	56.45	28	✓ APO-
			Atomoxetine S29
			✓ Generic Partners
Cap 100 mg	58.48	28	✓ APO-
			Atomoxetine \$29
			✓ Generic Partners
EXAMFETAMINE SULFATE - Special Authority see SA114	9 on the next page -	Retail phar	macy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency 100 ✓ PSM



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### ⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application — (Narcolepsy)** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

**Renewal** — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 on the next page - Retail pharmacy

a)	Only	on	а	controlled	drug	form
----	------	----	---	------------	------	------

b) Safety medicine; prescriber may determine dispensing frequency		
Tab immediate-release 5 mg3.20	30	✓ Rubifen
Tab immediate-release 10 mg3.00	30	✓ Ritalin
•		✓ Rubifen
Tab extended-release 18 mg7.75	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab immediate-release 20 mg7.85	30	✓ Rubifen
Tab sustained-release 20 mg	30	✓ Rubifen SR
Tab extended-release 27 mg11.45	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab extended-release 36 mg	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>
Tab extended-release 54 mg	30	<ul><li>Methylphenidate ER</li><li>Teva</li></ul>

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

#### ⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

**Initial application** — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

**Renewal** — **(ADHD in patients under 5)** only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy\*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: \*narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine: prescriber may determine dispensing frequency

by carety meaning, processes may actermine dispersing			
Tab extended-release 18 mg	58.96	30	<ul><li>Concerta</li></ul>
Tab extended-release 27 mg	65.44	30	<ul><li>Concerta</li></ul>
Tab extended-release 36 mg	71.93	30	<ul><li>Concerta</li></ul>
Tab extended-release 54 mg		30	<ul><li>Concerta</li></ul>
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	✓ Ritalin LA
Cap modified-release 40 mg		30	✓ Ritalin LA

#### ⇒SA1965 Special Authority for Subsidy

**Initial application** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid



(Ma	Subsidy nufacturer's Price)	Sul	Fully bsidised	Brand or Generic
	\$	Per	1	Manufacturer

continued...

for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
  - 3.1 Applicant is a paediatrician or psychiatrist; or
  - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
  - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
  - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

**Renewal** only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

1 Tho

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
  - 2.1 Applicant is a paediatrician or psychiatrist; or
  - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

#### ⇒SA1999 Special Authority for Subsidy

**Initial application** only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Fither:
  - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
  - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Fither:
  - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
  - 3.2 Methylphenidate and dexamfetamine are contraindicated.

**Renewal** only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for	Dementia
DONEDEZII LIVDDO	

טט	NEFEZIL HTDNOCHLONIDE		
*	Tab 5 mg4.34	90	✓ Donepezil-Rex
*	Tab 10 mg	90	✓ Donepezil-Rex

	Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
RIVASTIGMINE – Special Authority see SA1488 below – Retail p Patch 4.6 mg per 24 hour	•	30	<b>√</b> <u>R</u>	Rivastigmine Patch
Patch 9.5 mg per 24 hour	38.00	30	<b>√</b> <u>R</u>	BNM 5 Rivastigmine Patch BNM 10

#### ⇒SA1488 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

### Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency
- Tab sublingual 2 mg with naloxone 0.5 mg .......18.37
- ✓ Buprenorphine Naloxone BNM
- Tab sublingual 8 mg with naloxone 2 mg ......53.12
- ✓ Buprenorphine Naloxone BNM

### ⇒SA1203 Special Authority for Subsidy

**Initial application — (Detoxification)** from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

**Initial application — (Maintenance treatment)** from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone: and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the



Subsidy (Manufacturer's Price)	Fu Subsidise	,
 \$	Per	Manufacturer

continued...

following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone);
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

#### **BUPROPION HYDROCHLORIDE**

Tab modified-release 150 mg	11.00	30	✓ Zyban
DISULFIRAM Tab 200 mg	236.40	100	✓ Antabuse
NALTREXONE HYDROCHLORIDE - Special Authority see	SA1408 below - Retai	l pharmacy	
Tab 50 mg	133.33	30	✓ Naltraccord

#### ⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

**Renewal** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
  - 2.1 Patient is still unstable and requires further treatment; or
  - 2.2 Patient achieved significant improvement but requires further treatment; or
  - 2.3 Patient is well controlled but requires maintenance therapy.

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

#### NICOTINE

- a) Nicotine will not be funded in amounts less than 4 weeks of treatment.
- b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A. Patch 7 mg - Up to 28 patch available on a PSO ......18.14 ✓ Habitrol 28 Patch 7 mg for direct distribution only - [Xpharm]......3.94 7 ✓ Habitrol Patch 14 mg - Up to 28 patch available on a PSO ......19.95 28 ✓ Habitrol Patch 14 mg for direct distribution only - [Xpharm]......4.52 7 ✓ Habitrol Patch 21 mg - Up to 28 patch available on a PSO ......22.86 28 ✓ Habitrol Patch 21 mg for direct distribution only - [Xpharm]......5.18 7 ✓ Habitrol Lozenge 1 mg - Up to 216 loz available on a PSO......19.18 216 ✓ Habitrol 36 ✓ Habitrol Lozenge 2 mg - Up to 216 loz available on a PSO......21.02 216 ✓ Habitrol 36 ✓ Habitrol ✓ Habitrol Gum 2 mg (Fruit) - Up to 384 piece available on a PSO ......38.21 384 96 ✓ Habitrol Gum 2 mg (Mint) - Up to 384 piece available on a PSO......38.21 ✓ Habitrol 384 Gum 2 mg (Mint) for direct distribution only - [Xpharm]......8.64 96 ✓ Habitrol ✓ Habitrol Gum 4 mg (Fruit) - Up to 384 piece available on a PSO ......44.17 384 96 ✓ Habitrol Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17 ✓ Habitrol 384 Gum 4 mg (Mint) for direct distribution only - [Xpharm]......10.01 ✓ Habitrol 96

VARENICLINE TARTRATE - Special Authority see SA1845 below - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

#### ⇒SA1845 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either
  - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
  - 3.2 The patient has tried but failed to guit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

**Renewal** from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;

### NERVOUS SYSTEM

	Subsidy		Fully	Brand or
(N	Manufacturer's Price)	;	Subsidised	Generic
	\$	Per	•	Manufacturer

continued...

and

- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

# **Chemotherapeutic Agents**

## Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see SA2046 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

### ⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
  - 3.1 Both:
    - 3.1.1 Patient is treatment naive; and
    - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
  - 3.2 All of the following:
    - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
    - 3.2.2 The patient has not received prior bendamustine therapy; and
    - 3.2.3 Fither:
      - 3.2.3.1 Both:
        - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
        - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
      - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Fither:
  - 2.1 Both:

Subsidy (Manufacturer's Price)	Subsi	Fully idised	Brand or Generic
\$	Per	/	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with \* are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

Tob 2 mg	90.05	100	✓ Myleran
Tab 2 mg		100	wiyieran
CARBOPLATIN - PCT only - Specialist			4
Inj 10 mg per ml, 45 ml vial		1	✓ DBL Carboplatin
	45.20		Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
	1,387.00		✓ Bicnu Heritage S29
Inj 100 mg for ECP	710.00	100 mg OP	✓ Baxter
(Bicnu Heritage \$29 Inj 100 mg vial to be delisted 1 Septemb	er 2022)	•	
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	<ul><li>Cisplatin Ebewe</li></ul>
Inj 1 mg per ml, 100 ml vial	21.00	1	<ul><li>Cisplatin Ebewe</li></ul>
	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist		1	✓ Endoxan
, , , , ,	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
Inj 2 g	180.00	1	✓ Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
LOMUSTINE - PCT - Retail pharmacy-Specialist			
Cap 10 mg	132.59	20	✓ CeeNU
Cap 40 mg		20	✓ CeeNU
. •			

(M	Subsidy lanufacturer's Price) \$	Subs Per	Fully sidised	
MELPHALAN				
Tab 2 mg - PCT - Retail pharmacy-Specialist	40.70	25	1	Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	1	Alkeran
			1	Alkeran S29 S29
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	1	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	1	Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	1	Baxter
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	1	Bedford S29
, ,			1	Max Health S29
			1	THIO-TEPA S29
				Tepadina S29
Inj 100 mg vial	CBS	1		Max Health \$29
inj 100 mg vidi		'		Tepadina \$29
			•	i chanilla 222
Antimetabolites				
AZACITIDINE - PCT only - Specialist - Special Authority see SA14	167 below			
Inj 100 mg vial		1	•	Azacitidine Dr Reddy's

AZACITIDINE  – PCT only – Specialist – Special Autl	nority see SA1467 below		
Inj 100 mg vial	75.06	1	Azacitidine Dr
, ,			Reddy's
	605.00		✓ Vidaza
Inj 1 mg for ECP	0.83	1 mg	✓ Baxter

### ⇒SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic
  - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder);
  - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- 4 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
  - 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	e) Per	Subsidised	Generic Manufacturer
CALCUIN FOLINATE	Ψ	rei		iviariuiaciurei
CALCIUM FOLINATE  Tab 15 mg - PCT - Retail pharmacy-Specialist	11// 60	10	1	DBL Leucovorin
Tab 13 mg - FOT - netall pharmacy-Specialist	114.09	10	•	Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	1	Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali		1		Calcium Folinate
				Sandoz
			✓	Calcium Folinate
				Sandoz S29 S29
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	1	Leucovorin
				Pharmacia \$29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓	Calcium Folinate
				Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	/	Calcium Folinate
				Ebewe .
	94.90	10	•	Leucovorin
			_	Pharmacia S29
Inj 300 mg - PCT only - Specialist	22.51	1	•	Calcium Folinate
	05.44		,	Ebewe
	25.14		•	Leucovorin DBL S29
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	1	Calcium Folinate
, 1 3, 1				Sandoz
			✓	Calcium Folinate
				Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	1	Calcium Folinate
				Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓	Calcium Folinate
				Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	•	Baxter
CAPECITABINE - Retail pharmacy-Specialist				
Tab 150 mg		60		Capercit
Tab 500 mg	49.00	120	•	Capercit
CLADRIBINE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml		1		Litak S29
Inj 1 mg per ml, 10 ml		. 1		Leustatin
Inj 10 mg for ECP	749.96	10 mg C	)P	Baxter
CYTARABINE		_	_	
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special	ist400.00	5	•	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail	44.00		,	D#
pharmacy-Specialist		1 10 mg		Pfizer Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Special		00 mg (		Baxter
FLUDARABINE PHOSPHATE		oo nig (	J. •	Durioi
Tab 10 mg - PCT - Retail pharmacy-Specialist	412.00	20	J	Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5		Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist		50 mg C		Baxter
,				

	Subsidy (Manufacturer's Price		Fully bsidised	Brand or Generic
	\$	Per		Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	<b>✓</b>	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	<b>✓</b>	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist		100 mg	<b>✓</b>	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist		ŭ		
Inj 1 g, 26.3 ml vial	62 50	1	1	DBL Gemcitabine
Inj 1 gInj 1 g		1		Gemcitabine Ebewe
Inj 1 mg for ECP		1 mg		Baxter
• •	0.02	inig	• '	Daxici
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist	50.53		,	
Inj 20 mg per ml, 5 ml vial		1	-	Accord
	71.44		<b>√</b>	Irinotecan Actavis 100
	100.00		<b>✓</b>	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	<b>✓</b>	Baxter
MERCAPTOPURINE		ŭ		
	27.00	25	<b>✓</b> I	Puri-nethol
Tab 50 mg – PCT – Retail pharmacy-Specialist		20	•	rui i-iletiitti
Oral suspension 20 mg per ml – Retail pharmacy-Specialis				• ••
Special Authority see SA1725 below	428.00	100 ml OP		Allmercap

■ SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

### **METHOTREXATE**

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ <u>Trexate</u>
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	✓ Methotrexate
			Sandoz
*	Inj 10 mg prefilled syringe14.66	1	✓ Methotrexate
	, 01		Sandoz
*	Inj 15 mg prefilled syringe14.77	1	✓ Methotrexate
	, - 3, , 3-		Sandoz
*	Inj 20 mg prefilled syringe14.88	1	✓ Methotrexate
•	.,	-	Sandoz
*	Inj 25 mg prefilled syringe	1	✓ Methotrexate
	, =0g p. 0	•	Sandoz
*	Inj 30 mg prefilled syringe	1	✓ Methotrexate
-,-	ing of mg promited symigo		Sandoz
*	Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL
-,-	The state of the s	Ü	Onco-Vial
*	Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate
~	Inj 25 mg per mi, 20 mi viai = 1 0 1 = Hetaii phamiacy-opedialist45.00	'	Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
		ı	▼ Methotrexate Enewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	. 1	✓ <u>Methotrexate Ebewe</u>
*	Inj 1 mg for ECP – PCT only – Specialist	_ 1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	/	Baxter

### **⇒SA1679** Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

**Renewal — (mesothelioma)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient has chemotherapy-naïve disease; and
    - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
  - 2.2 All of the following:
    - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
    - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
    - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m<sup>2</sup> every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m<sup>2</sup> every 21 days.

•		
Other Cytotoxic Agents		
AMSACRINE - PCT only - Specialist		
Inj 50 mg per ml, 1.5 ml ampoule1,500.00	6	✓ Amsidine S29
4,736.00		✓ Amsidine S29
Inj 75 mg1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharmacy-Specialist		
Cap 0.5 mg1,175.87	100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist		
Inj 1 mg per ml, 10 ml vial	10	✓ Phenasen
Inj 10 mg for ECP481.70	10 mg OP	✓ Baxter

I anvis

25

	Subsidy (Manufacturer's Price \$	e) Subs Per	Fully sidised	Brand or Generic Manufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist Inj 15,000 iu, vial	185.16	1		BL Bleomycin Gulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ Ba	xter
BORTEZOMIB – PCT only – Specialist – Special Authority see S Inj 2.5 mg vial		1		rtezomib Juno S29 S29
Inj 3.5 mg vial	105.00	1	✓ Bo	rtezomib Dr Reddy's S29 S29 rtezomib Dr-Reddy's rtezomib Juno S29
Inj 1 mg for ECP	31.20	1 mg	✓ Ba	xter
(Bortezomib Juno S29 S29 Inj 2.5 mg vial to be delisted 1 Augus (Bortezomib Dr Reddy's S29 S29 Inj 3.5 mg vial to be delisted 1	st 2022)	•		

(Bortezomib Juno S29 Inj 3.5 mg vial to be delisted 1 August 2022)

## **⇒SA1889** Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Fither:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis \*.

Note: Indications marked with \* are unapproved indications.

DACARBAZINE - PCT only - Specialist		
Inj 200 mg vial62.70	1	✓ DBL Dacarbazine
580.60	10	✓ Dacarbazine  APP \$29
Inj 200 mg for ECP62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist		
Inj 0.5 mg vial255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP255.00	0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		
Inj 2 mg per ml, 10 ml149.50	1	✓ Pfizer
Inj 20 mg vial	10	✓ Daunorubicin
		Zentiva S29
Inj 20 mg for ECP149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist		
Inj 20 mg48.75	1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial46.89	1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial26.95	1	✓ Docetaxel
		Accord S29
Inj 80 mg195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP0.65	1 mg	✓ Baxter

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic  Manufacturer
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial	11.50	1	Doxorubicin Ebewe
	17.00		Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	Arrow-Doxorubicin
	69.99		Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
ETOPOSIDE		Ū	
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	✓ Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	✓ Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specia		1	✓ Rex Medical
Inj 1 mg for ECP - PCT only - Specialist		1 mg	✓ Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist			
Inj 100 mg (of etoposide base)	40.00	1	✓ Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg	✓ Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] - PCT - Retail pha		ŭ	
Cap 500 mg	23.82	100	✓ Devatis
IDARUBICIN HYDROCHLORIDE		100	<u> </u>
Inj 5 mg vial - PCT only - Specialist	100 74	1	✓ Zavedos
Inj 10 mg vial – PCT only – Specialist		1	✓ Zavedos ✓ Zavedos
Inj 1 mg for ECP – PCT only – Specialist		1 mg	
, , ,		ı my	Daxlei
LENALIDOMIDE – Retail pharmacy-Specialist – Special Author Wastage claimable	•		
Cap 5 mg	5,122.76	28	✓ Revlimid
Cap 10 mg	4,655.25	21	✓ Revlimid
	6,207.00	28	✓ Revlimid
Cap 15 mg	5,429.39	21	✓ Revlimid
	7,239.18	28	✓ Revlimid
Cap 25 mg	7,627.00	21	✓ Revlimid

## ⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
  - 3.1 Lenalidomide to be used as third line\* treatment for multiple myeloma; or
  - 3.2 Both:
    - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
    - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with \* is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

### **MESNA**

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ <u>Uromitexan</u>
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	<ul><li>Uromitexan</li></ul>
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	641.70	1	✓ Accord S29
Inj 20 mg vial	3,275.00	1	✓ Omegapharm S29
, ,			✓ Teva
Inj 1 mg for ECP	470.75	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see	e SA1883 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg	3,701.00	56	✓ Lynparza

### ⇒SA1883 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Patient has a high-grade serous\* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: \*Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

## PACLITAXEL - PCT only - Specialist

Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority see SA19	979 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

### ⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN	[DEOXYCOFORMYCIN	<ul><li>I) – PCT only – Specialist</li></ul>
-------------	------------------	--

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
PROCARBAZINE HYDROCHLORIDE - PCT - Retail pharma	cy-Specialist			
Cap 50 mg	980.00	50	✓	Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below - Re	etail pharmacy			
Cap 5 mg	9.13	5	1	Temaccord
Cap 20 mg	16.38	5	1	Temaccord
, •	18.30		1	Apo-Temozolomide
	136.00	14	1	Accord S29
Cap 100 mg	35.98	5	1	Temaccord
•	40.20		1	Apo-Temozolomide
	532.00	14	1	Accord S29
Cap 140 mg	50.12	5	1	Temaccord
	400.00		1	Amneal S29
Cap 180 mg	620.00	14	1	Accord S29
Cap 250 mg		5	1	Temaccord
•	688.00		1	Amneal S29

## ⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
  - 1.2 Patient has newly diagnosed anaplastic astrocytoma\*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour\*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

**Initial application** — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has glioblastoma multiforme; and
  - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
  - 2.1 Patient has anaplastic astrocytoma\*; and
  - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
  - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub: Per	sidised •	Generic Manufacturer	

continued...

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

**Renewal — (ewing's sarcoma)** only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a \* is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Special	I Authority see SA1124 below	
Cap 50 mg	378.00 28	Thalomid
Cap 100 mg	756.00 28	Thalomid

## ⇒SA1124 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

### Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis\*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with \* is an unapproved indication.

### **TRETINOIN**

Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority	see SA1868 belo	)W	
Tab 14 × 10 mg, 7 × 50 mg, 21 × 100 mg	1,771.86	42 OP	✓ Venclexta
Tab 10 mg		14 OP	✓ Venclexta
Tab 50 mg	239.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable	8,209.41	120	✓ Venclexta

### ⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)\* and B-cell prolymphocytic leukaemia (B-PLL)\*. Indications marked with \* are Unapproved indications.

### VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist102.73	5	<ul><li>DBL Vincristine Sulfate</li></ul>
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓ Navelbine
210.00		✓ Vinorelbine Ebewe
328.65		✓ Sagent S29
Inj 1 mg for ECP1.25	1 mg	✓ Baxter
Inj 50 mg for ECP	50 mg OP	✓ Baxter (Sagent)

# **Protein-tyrosine Kinase Inhibitors**

ALECTINIB – Retail pharmacy-Specialist – Special Authority see SA1870 below
Wastage claimable

## **⇒SA1870** Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Al K test: and

3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

## DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable

Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

## ⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
  - 1.2 Maximum dose of 140 mg/day: or
- 2 Both:
  - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
  - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
  - 3.1 The patient has a diagnosis of CML in chronic phase; and
  - 3.2 Maximum dose of 100 mg/day; and
  - 3.3 Any of the following:
    - 3.3.1 Patient has documented treatment failure\* with imatinib; or
    - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
    - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or

3.3.4 Patients is enrolled in the KISS study\*\* and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib\*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: \*treatment failure for CML as defined by Leukaemia Net Guidelines. \*\*Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special Authority se	e SA2115 below		
Tab 100 mg	764.00	30	✓ Tarceva
Tab 150 mg	1,146.00	30	Tarceva

### ⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	) Si	ubsidised	Generic	
\$	Per	/	Manufacturer	

continued

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither:
  - 3.1 Patient is treatment naive: or
  - 3.2 Both:
    - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
    - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see SA2116 below

Tab 250 mg .......1,700.00 30 ✓ Iressa

## ⇒SA2116 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
  - 2.1 Patient is treatment naive; or
  - 2.2 Both:
    - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

### IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg – [Xpharm] – Special Authority see SA1460 on t	he		
	next page	2,400.00	60	✓ Glivec
*	Cap 100 mg	58.23	60	✓ Imatinib-Rex
*	Cap 400 mg	84.79	30	✓ Imatinib-Rex

Subsidy (Manufacturer's Price)

Subsidised Per

Fully

Brand or Generic Manufacturer

## ⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

### Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Tvkerb

## ⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology);
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

### NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6,532.00	120	Tasigna

## ⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither:
  - 2.1 Patient has documented CML treatment failure\* with imatinib; or
  - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

Note: \*treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
PALBOCICLIB – Retail pharmacy-Specialist – Special Authorit Wastage claimable	y see SA1894 below			
Tab 75 mg	4,000.00	21	<b>✓</b>	brance
Tab 100 mg	4,000.00	21	<b>√</b>	brance
Tab 125 mg	4,000.00	21	<b>✓</b> I	brance

### ⇒SA1894 Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
  - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
  - 4.2.2.2 All of the following:
    - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
    - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
    - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

### PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

## ⇒SA1190 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

### All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 Both:
    - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
    - 2.3.2 The cancer did not progress whilst on sunitinib; and

(1	Subsidy Manufacturer's Price)	F Subsid	ully	Brand or Generic
· ·	\$	Per	1	Manufacturer

### continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal; or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

### DOIII.

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage Claimable			
Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg	· ·	56	<ul><li>Jakavi</li></ul>
Tab 15 mg	· ·	56	Jakavi
Tab 20 mg	· ·	56	<ul><li>Jakavi</li></ul>

### ⇒SA1890 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
  - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
  - 2.2 Both:

Wastage claimable

- 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
- 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer
SUNITINIB - Special Authority see SA2117 below - Retail pha	rmacy			
Cap 12.5 mg	208.38	28	<b>√</b> S	Sunitinib Pfizer
, ,	2,315.38		<b>√</b> S	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022	•			
Cap 25 mg	416.77	28	<b>√</b> S	Sunitinib Pfizer
	4.630.77		<b>√</b> S	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022	,			
Cap 50 mg	694.62	28	<b>√</b> S	Sunitinib Pfizer
3	9.261.54		<b>√</b> S	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022	0,=01101		_	
(Sutent Cap 12.5 mg to be delisted 1 July 2022)				

(Sutent Cap 25 mg to be delisted 1 July 2022)

(Sutent Cap 50 mg to be delisted 1 July 2022)

## ⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
  - 2.1 The patient is treatment naive; or
  - 2.2 The patient has only received prior cytokine treatment; or
  - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
  - 2.4 Both:
    - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
  - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
  - 5.2 Haemoglobin level < lower limit of normal: or
  - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
  - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
  - 5.5 Karnofsky performance score of less than or equal to 70; or
  - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
  - 2.1 The patient's disease has progressed following treatment with imatinib; or
  - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

1 No evidence of disease progression; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

continued...

2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria: Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
  - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
  - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
  - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

# Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see \$A2118 below

Wastage claimable

Tab 250 mg .......4,276.19 120 ✓ Zytiga

## ⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
  - 4.1 All of the following:
    - 4.1.1 Patient is symptomatic; and
    - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and

Subsic		Fully	Brand or
(Manufacture	r's Price) Subs	idised	Generic
\$	Per	1	Manufacturer

continued...

- 4.1.3 Patient has ECOG performance score of 0-1; and
- 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
- 4.2 All of the following:
  - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane: and
  - 4.2.2 Patient has ECOG performance score of 0-2; and
  - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression, and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

**Renewal — (pandemic circumstances)** from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

### **BICALUTAMIDE**

Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
· ·	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special	Authority see SA1895 belo	)W	
Inj 50 mg per ml, 5 ml prefilled syringe	1,068.00	2	✓ Faslodex

### ⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

	Subsidy (Manufacturer's Price)	Q.,	Fully bsidised	Brand or Generic
	(Manufacturer's Frice)	Per	JSIUISEU 🗸	Manufacturer
MEGESTROL ACETATE – Subsidy by endorsement Subsidy by endorsement – Subsidised for patients who wer prescription is endorsed accordingly. Pharmacists may ann prior dispensing of megestrol acetate.				
Tab 160 mg	48.80	30	✓ N	Megace S29
(Megace S29 Tab 160 mg to be delisted 1 February 2023)				J
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml vial	30.64	5	1	Octreotide
,				MaxRx S29
	56.87		✓ [	DBL Octreotide
Inj 100 mcg per ml, 1 ml vial	40.00	5	✓ [	DBL Octreotide
Inj 500 mcg per ml, 1 ml vial	145.00	5	✓ [	DBL Octreotide
	222.00		✔ (	Octreotide (Sun) 829
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	<b>✓</b> N	/lax Health
Max Health to be Principal Supply on 1 June 2022	30.64		✓ (	Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32 71	5	✓ N	Max Health
ing roo mag per mi, r mi ampoule		J	-	Octreotide GH \$29
Max Health to be Principal Supply on 1 June 2022			•	Journal and
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	-	Max Health Octreotide GH S29
Max Health to be Principal Supply on 1 June 2022				
(Octreotide MaxRx \$29 Inj 50 mcg per ml, 1 ml vial to be delist (DBL Octreotide Inj 50 mcg per ml, 1 ml vial to be delisted 1 Jun (DBL Octreotide Inj 100 mcg per ml, 1 ml vial to be delisted 1 Jun (DBL Octreotide Inj 500 mcg per ml, 1 ml vial to be delisted 1 Jun (Octreotide (Sun) \$29 Inj 500 mcg per ml, 1 ml vial to be delisted (Octreotide GH \$29 Inj 50 mcg per ml, 1 ml ampoule to be delicoted GH \$29 Inj 100 mcg per ml, 1 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 2 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 2 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 2 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 2 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 3 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 4 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 5 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$29 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 6 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 7 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per ml, 1 ml ampoule to be delicoted GH \$20 Inj 500 mcg per m	ne 2022) Ine 2022) Ine 2022) Ine 2022) Ine 2022) Insted 1 June 2022) Insted 1 June 2022)	acy		
Inj depot 10 mg prefilled syringe	439.97	1	_	Octreotide Depot Teva
Inj depot 20 mg prefilled syringe	647.03	1	<b>√</b> <u>(</u>	Octreotide Depot Teva
Inj depot 30 mg prefilled syringe	718.55	1	✓ (	Octreotide Depot Teva

## **⇒SA2119** Special Authority for Subsidy

**Initial application — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

### All of the following:

- 1 The patient has nausea\* and vomiting\* due to malignant bowel obstruction\*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with \* are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal — (Malignant Bowel Obstruction)** from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
  - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
  - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
  - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

**Renewal — (Acromegaly)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly: and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

**Initial application — (Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
  - 2.1 Gastrinoma; and
  - 2.2 Either:
    - 2.2.1 Patient has failed surgery; or
    - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
  - 3.1 Insulinomas; and
  - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
  - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
  - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

**Renewal — (Other Indications)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

### TAMOXIFEN CITRATE

*	Tab 10 mg	15.00	60	✓ <u>Tamoxifen Sandoz</u>
*	Tab 20 mg	6.65	60	✓ <u>Tamoxifen Sandoz</u>

## **Aromatase Inhibitors**

ANASTROZOLE	30	✓ Anatrole
EXEMESTANE	30	✓ Pfizer Exemestane
LETROZOLE	30	✓ <u>Letrole</u>

## **Immunosuppressants**

## Cytotoxic Immunosuppressants

# AZATHIOPRINE

不	Tab 25 mg	7.30	00	<ul><li>Azamun</li></ul>
*	Tab 50 mg	7.60	100	✓ Azamun
	Inj 50 mg vial199		1	✓ Imuran
	uran Inj 50 mg vial to be delisted 1 January 2023)			
MY	COPHENOLATE MOFETIL			
	Tab 500 mg	5.90	50	<ul><li>Cellcept</li></ul>
	Cap 250 mg	5.90	100	<ul> <li>Cellcept</li> </ul>
	Powder for oral liq 1 g per 5 ml - Subsidy by endorsement187		165 ml OP	✓ Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

7 05

### **Fusion Proteins**

FTANERCEPT - Special Authority see SA2103 below - Retail pharmacy

Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector		4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe	1,050.00	4	✓ Enbrel

### ⇒SA2103 Special Authority for Subsidy

**Initial application — (adult-onset Still's disease)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.1 Either:
  - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
  - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
- 1.2 Fither:
  - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
  - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

**Initial application** — **(ankylosing spondylitis)** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
  - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
  - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

continued...

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

Average normal chiest expansion corrected for a 18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

intolerance: and

- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

### ....

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Per	1	Manufacturer

continued...

- 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Either:
  - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
  - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
  - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
  - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

**Initial application — (Arthritis - rheumatoid )** only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
- 1.2 Fither:
  - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
  - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
  - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
  - 2.5 Either:
    - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
    - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
  - 2.6 Either:
    - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

**Renewal — (Arthritis - rheumatoid)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or

Subsi	.,	
(Manufacture	er's Price) Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Either:
      - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
      - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
  - 2.2 Both:
    - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
    - 2.2.2 Either:
      - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
      - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and

Subsidy (Manufacturer's Pric	e)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

### continued...

- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose): and
- 5 Any of the following:
  - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

### All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

### Immune Modulators

st		
2,774.48	5	✓ ATGAM
Specialist		
149.37	1	✓ OncoTICE
176.90	3	✓ SII-Onco-BCG S29
	2,774.48 Specialist 149.37	2,774.48 5 Specialist 1

## Monoclonal Antibodies

	nacy	A2102 below – Retail pharm	ADALIMUMAB (AMGEVITA) - Special Authority see SA
✓ Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

### ⇒SA2102 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

### Both:

- 1 The patient has severe Behcet's disease\* that is significantly impacting the patient's quality of life; and
- 2 Fither:

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

- 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
- 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with \* are unapproved indications.

**Initial application — (Hidradenitis suppurativa)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

**Renewal** — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
  - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
  - 1.2 Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		ubsidised	Generic	
	Per	✓	Manufacturer	

continued...

- 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
- 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
  - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 2.2 Either:
    - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with \* are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection;
  - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab: or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	

continued...

4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less: or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e); or
  - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < 1/2+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	1	Manufacturer

continued...

**Initial application — (Ocular inflammation - severe)** from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
- 2 All of the following:
  - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
  - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
  - 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
  - 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
  - 2.5 Either:
    - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
    - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
  - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Approvals valid for 6 months for applications meeting the following criteria:

### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA): and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects; or
    - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
  - 2.3 Either:
    - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

### 1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects: or
  - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
  - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 Any of the following:
    - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
  - 12 Fither
    - 1.2.1 The patient has experienced intolerable side effects: or
    - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
    - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

**Initial application** — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
  - 1.2 Either
    - 1.2.1 The patient has experienced intolerable side effects; or
    - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
  - 2 All of the following:
    - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and

S	Subsidy	Fully	Brand or
(Manufa	cturer's Price)	Subsidised	Generic
	\$ Per	r <b>√</b>	Manufacturer

continued...

- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
- 2.5 Fither
  - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
  - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
  - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
  - 12 Fither
    - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
    - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

**Initial application — (ulcerative colitis)** only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
  - 2.1 Patient's SCCAI score is greater than or equal to 4; or
  - 2.2 Patient's PUCAI score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Fither:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis\* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
  - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application: or
  - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with \* are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose: and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
- 5 Any of the following:
  - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
  - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

**Renewal — (inflammatory bowel arthritis – peripheral)** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

#### Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA) - Special Authority see SA2101 below - Retail pharmacy

Inj 20 mg per 0.4 ml prefilled syringe	1,599.96	2	<ul><li>Humira</li></ul>
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	<ul><li>Humira</li></ul>

#### ⇒SA2101 Special Authority for Subsidy

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
  - 1 Either:
    - 1.1 Applicant is a rheumatologist; or
    - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
  - 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
  - 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
  - 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment: or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

(Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
  - 2.1 Either:
    - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
    - 2.1.2 CDAI score is 150 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
    - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Either:
    - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
    - 2.1.2 PCDAI score is 15 or less; or
  - 2.2 Both:
    - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed;
    - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Applicant is a gastroenterologist; or
  - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2 Either:
  - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
  - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

**Renewal — (psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement: and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Applicant is a rheumatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Fither:
  - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
  - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
  - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Applicant is a dermatologist; or
  - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
    - 2.1.2 Either:
      - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
      - 2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or
  - 2.2 Both:

Subsidy	Fully	Brand or
(Manufacturer's		Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2.2 Either:
  - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
  - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value: and
- $3\,$  Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

**Renewal — (severe ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 The patient has had a good clinical response following 3 initial doses; or
  - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
  - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
  - 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

✓ Eylea

#### ⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Wet age-related macular degeneration (wet AMD); or
    - 1.1.2 Polypoidal choroidal vasculopathy; or
    - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
  - 1.2 Either:
    - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
    - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
  - 1.3 There is no structural damage to the central fovea of the treated eye; and
  - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
  - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
  - 2.2 Patient has previously\* (\*before June 2018) received treatment with ranibizumab for wAMD and disease was stable

Subsidy	Ful	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
 \$	Per •	Manufacturer

continued...

while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

**Renewal — (wet age related macular degeneration)** only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

### ⇒SA2096 Special Authority for Subsidy

**Initial application** — (**Treatment of profoundly immunocompromised patients**) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity\*; and
- 3 Patient is profoundly immunocompromised\*\* and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: \* Mild to moderate disease severity as described on the Ministry of Health Website

\*\* Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Erbitux	1	364.00	 Inj 5 mg per ml, 20 ml vial	
✓ Erbitux	1	1,820.00	 Inj 5 mg per ml, 100 ml vial	
✓ Baxter	1 mg	3.82	 Inj 1 mg for ECP	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **⇒SA1697** Special Authority for Subsidy

**Initial application** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB - PCT only - Special Authority see SA2082 below

Inj 100 mg	806.00	1	✓ Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

### ⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
  - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
  - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
  - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
  - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
  - 1.2 CDAI score is 150 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
  - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
  - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and

Subsidy	Fı	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
  - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
  - 1.2 PCDAI score is 15 or less; or
  - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — **(Graft vs host disease)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the qut.

**Initial application — (Pulmonary sarcoidosis)** from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

**Renewal — (ankylosing spondylitis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	(Manufacturer's P
--	-------------------

continued...

- 2 Both:
  - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
  - 2.2 Any of the following:
    - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
    - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
    - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

**Renewal — (chronic ocular inflammation)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Fither:
  - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
  - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
  - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain: and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

- 4 Either:
  - 4.1 IV cyclophosphamide has been tried; or
  - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

**Renewal — (neurosarcoidosis)** only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

- Either:
  - 1 A withdrawal period has been tried and the patient has relapsed; or
  - 2 All of the following:
    - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
    - 2.2 There has been a marked reduction in prednisone dose; and
    - 2.3 Fither:
      - 2.3.1 There has been an improvement in MRI appearances; or
      - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

# 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
- 1.2 Either:
  - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
  - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
  - 2.1 Either:
    - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
    - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
  - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
  - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Roth:

- 1 Fither:
  - 1.1 Both:

Subs (Manufactur		
\$	Per	Manufacturer

continued...

- 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
- 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
- 1.2 Both:
  - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
  - 1.2.2 Fither:
    - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
    - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 Rheumatoid arthritis: or
  - 2.2 Ankylosing spondylitis; or
  - 2.3 Psoriatic arthritis; or
  - 2.4 Severe ocular inflammation: or
  - 2.5 Chronic ocular inflammation: or
  - 2.6 Crohn's disease (adults): or
  - 2.7 Crohn's disease (children); or
  - 2.8 Fistulising Crohn's disease: or
  - 2.9 Severe fulminant ulcerative colitis: or
  - 2.10 Severe ulcerative colitis: or
  - 2.11 Plague psoriasis; or
  - 2.12 Neurosarcoidosis; or
  - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for

- psoriatic arthritis; and 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
  - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

**Renewal — (rheumatoid arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
  - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

**Initial application** — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
  - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
  - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Subsidy	y Full	/ Brand or
(Manufacturer's	s Price) Subsidise	d Generic
\$	Per 🗸	Manufacturer

continued...

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application** — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
    - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
  - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
  - 2.2 Any of the following:
    - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
    - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
    - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

**Initial application** — **(ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Fither:
  - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
  - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

corticosteroids: and

4 Surgery (or further surgery) is considered to be clinically inappropriate.

**Renewal — (ulcerative colitis)** only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Fither:
  - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
  - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

**Initial application — (pyoderma gangrenosum)** only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum\*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with \* are unapproved indications.

**Renewal — (pyoderma gangrenosum)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA1896 below - Retail pharmacy		
Inj 100 mg prefilled pen	) 1	✓ Nucala
Ini 100 mg vial 1,638.00	) 1	✓ Nucala

#### ⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10<sup>9</sup> cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Fither

Subsidy		Fully	Brand or	
(Manufacturer's Pr	ice) S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids: or
- 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
  - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
  - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

## **⇒SA1627** Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts\* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

\* Neutrophil greater than or equal to  $1.5 \times 10^9$ /L and platelets greater than or equal to  $75 \times 10^9$ /L.

#### ⇒SA1744 Special Authority for Subsidy

**Initial application** — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Per 🗸	Manufacturer

continued...

- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
  - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
  - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

**Initial application — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
    - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
  - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
  - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
  - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
  - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
  - 4.1 Treatment to be stopped if inadequate response\* following 4 doses; or
  - 4.2 Complete response\* to 6 doses of omalizumab.

**Renewal — (severe asthma)** only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

**Renewal — (severe chronic spontaneous urticaria)** only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded\* to 6 doses of omalizumab; or
- 2 Both:
  - 2.1 Patient has previously had a complete response\* to 6 doses of omalizumab; and
  - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: \*Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	1	Manufacturer

continued...

Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 1	4 ml vial	1	1	Perjeta
Inj 420 mg for ECF	<sup>2</sup> 3,927.00	420 mg OP	1	Baxter

### ⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
  - 2.1 Patient is chemotherapy treatment naïve; or
  - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

## RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

### ⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
  - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
  - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
  - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
<b>\$</b>	Per 🗸	Manufacturer

continued...

leflunomide alone or in combination with oral or parenteral methotrexate; and

- 6 Either:
  - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
  - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Fither:
  - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
  - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
  - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
  - 1.2 Fither:
    - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
    - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Either:
  - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sub	sidised	Generic
\$	Per	1	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Fither:
  - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
  - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
  - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
  - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2114 below

inj 100 mg per 10 mi viai	2/5.33	2	✓ RIXIMYO
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

### ⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*: and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
  - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
  - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
  - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
  - 3.4 Patient is a female of child-bearing potential; or
  - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with \* are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis\*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> of body-surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection\*.

Note: Indications marked with \* are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
  - 2.1 The patient is rituximab treatment naive; or
  - 2.2 Fither:
    - 2.2.1 The patient is chemotherapy treatment naive: or
    - 2.2.2 Both:
      - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
      - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
  - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Fither:
  - 4.1 The patient does not have chromosome 17p deletion CLL; or
  - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

**Renewal — (Chronic lymphocytic leukaemia)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

#### 1 Either:

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
  - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
  - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
  - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
  - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is

Subsidy	F	ully	Brand or	
(Manufacturer's Pr	rice) Subsid	ised	Generic	
\$	Per	✓	Manufacturer	

continued...

considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
  - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
  - 2.2 All of the following:
    - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
    - 2.2.2 The patient is receiving treatment with mycophenolate; and
    - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

**Initial application — (Post-transplant)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder\*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with \* are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

4 0

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
  - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
  - 2.2 Both:
    - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
    - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
  - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
  - 3.2 Both:
    - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
    - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS\* or FRNS\*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS\* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

**Renewal** — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	•	Manufacturer	

continued...

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome\*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
  - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
  - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
  - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

**Initial application — (haemophilia with inhibitors)** only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:
All of the following:

- 1 Either:
  - 1.1 Patient has immune thrombocytopenic purpura\* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
  - 1.2 Patient has immune thrombocytopenic purpura\* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1 Treatment with steroids and splenectomy have been ineffective; or
- 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
- 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

#### Fither:

- 1 Both:
  - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
  - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia\* requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia\* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. \*Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with \* are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia\* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with \* are unapproved indications.

Subsidy (Manufacturer's Price)	Fu Subsidis	,	
\$	Per	<ul> <li>Manufacturer</li> </ul>	

continued...

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease\*: and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with \* are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks; and
- 2 Either:
  - 2.1 Patient has thrombotic thrombocytopenic purpura\* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
  - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura\* with neurological or cardiovascular pathology.

Note: Indications marked with \* are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura\*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE\*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Note: Indications marked with \* are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE\* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with \* are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia\*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with \* are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
  - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia\*; and
  - 2.2 An initial response lasting at least 12 months was demonstrated; and
  - 2.3 Patient now requires repeat treatment.

Note: Indications marked with \* are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
  - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
  - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

**Renewal — (severe antisynthetase syndrome)** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

**Initial application** — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
    - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
  - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

continued...

- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
  - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Roth
  - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
  - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

All of the follow

- 1 Fither:
  - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy\*; or
  - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks.

**Renewal — (Membranous nephropathy)** only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy\*; and
- 2 Fither:
  - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
  - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Notes:

- a) Indications marked with \* are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma\*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m<sup>2</sup> per dose for a maximum of 18 doses.

Note: Indications marked with \* are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant\*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with \* are unapproved indications.

Initial application — (pemiphiqus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
  - 1.1 Patient has severe rapidly progressive pemphigus; and
  - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
  - 1.3 Any of the following:
    - 1.3.1 Skin involvement is at least 5% body surface area; or
    - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions: or
    - 1.3.3 Involvement of two or more mucosal sites: or
- 2 Both:
  - 2.1 Patient has pemphigus; and
  - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with \* are unapproved indications.

Renewal — (pemiphigus\*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with \* are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy Inj 150 mg per ml, 1 ml prefilled syringe......799.50 ✓ Cosentvx 1.599.00 ✓ Cosentyx

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
  - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
  - 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
  - 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
  - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
  - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course: and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
  - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
  - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
  - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis - second-line biologic) only from a rheumatologist or medical practitioner on the

Subsidy (Manufacturer's Price)	9	Fully ubsidised	Brand or Generic
 (Manufacturer's Frice)	Per	ubsidised ✓	Manufacturer

continued...

recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
  - 1.2 Either:
    - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
    - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
  - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
  - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
  - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
  - 2.4 Either:
    - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
    - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
  - 2.5 Any of the following:
    - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
    - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
    - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

#### 1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the n	next page - Retail pharmacy		
Note: Siltuximab is to be administered at doses no	greater than 11 mg/kg every	3 weeks.	
Inj 100 mg vial	770.57	1	✓ Sylvant
Ini 400 mg vial	3.082.33	1	✓ Svlvant

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	1	Manufacturer	

### ⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

**Renewal** only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB - PCT only - Special Authority see SA2100 below			
Inj 20 mg per ml, 4 ml vial	220.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	880.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial	550.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial	1,100.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	4,400.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

# **⇒SA2100** Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

# 1 All of the following:

- 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
- 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
- 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or

#### 2 All of the following:

- 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
- 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
- 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
  - 2.1 rheumatoid arthritis; or
  - 2.2 systemic juvenile idiopathic arthritis; or
  - 2.3 adult-onset Still's disease; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.4 polyarticular juvenile idiopathic arthritis; or
- 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Fither:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
  - 3.2 Both:
    - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
    - 3.2.2 Either:
      - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
      - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
  - 3.1 Treatment with methotrexate is contraindicated: or
  - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
  - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
  - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
  - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints;
  - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
  - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
  - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
  - 1.1 Fither:
    - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
    - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule: and
  - 1.2 Either:
    - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
    - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
  - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
  - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
  - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 Both:
  - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
  - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
  - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
  - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
  - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
  - 2.4 Any of the following:
    - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
    - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (moderate to severe COVID-19\*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Note: Indications marked with \* are unapproved indications.

**Renewal — (Rheumatoid Arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

2 Either:

Both:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP		1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	/	Manufacturer

continued...

- 2 Either:
  - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 2.2 Both:
    - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 2.2.2 The cancer did not progress whilst on lapatinib: and
- 3 Either:
  - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 3.2 All of the following:
    - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
    - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
  - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
  - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned: or
  - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
  - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

**Renewal** — (early breast cancer\*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

### All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
  - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
  - 3.2 Both:
    - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
    - 3.2.2 The cancer did not progress whilst on lapatinib; or

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	Subs	idised	Generic
	\$	Per	✓	Manufacturer

continued...

- 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
  - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
  - 4.2 All of the following:
    - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
    - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
    - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: \* For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA1871 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

#### ⇒SA1871 Special Authority for Subsidy

**Initial application** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
  - 3.1 The patient has received prior therapy for metastatic disease\*; or
  - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy\*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Fither:
  - 5.1 Patient does not have symptomatic brain metastases; or
  - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

**Renewal** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine;
- 2 Treatment to be discontinued at disease progression.

Note: \*Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

# Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB - PCT only - Specialist - Special Authority	see SA2120 on the next p	oage	
Inj 10 mg per ml, 4 ml vial	1,051.98	1	<ul><li>Opdivo</li></ul>
Inj 10 mg per ml, 10 ml vial	2,629.96	1	<ul><li>Opdivo</li></ul>
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# ⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded pembrolizumab; or
  - 4.2 Roth
    - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

**Renewal** only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
  must have reduction in short axis to < 10 mm.</li>
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
PEMBROLIZUMAB - PCT only - Specialist - Special Authority s	see SA2121 below				
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓	Keytruda	
Inj 1 mg for ECP	49.14	1 mg	·	Baxter	
Tarina in the second		_			

SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV: and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
  - 4.1 Patient has not received funded nivolumab; or
  - 4.2 Both:
    - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
    - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
  - 1.1 Any of the following:
    - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
    - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
    - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
  - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
  - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
  - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
  - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
  - 2.2 Patient has signs of disease progression; and
  - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

#### continued...

- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

# Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg	88.91	50	Neoral
Cap 100 mg	177.81	50	Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail pharr	nacy		
Wastage claimable	0.540.00	00	. Afinitan
Tab 10 mg		30	Afinitor
Tab 5 mg	4,555.76	30	Afinitor

### ⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

#### Both:

- 1 Patient has tuberous sclerosis: and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

**Renewal** only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

#### SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg		100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral lig 1 mg per ml	449.99	60 ml OP	✓ Rapamune

### ⇒SA2005 Special Authority for Subsidy

**Initial application** from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy: or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP: or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid

(1	Subsidy Manufacturer's Price)	F Subsid	ully	Brand or Generic
· ·	\$	Per	1	Manufacturer

continued...

for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation\*; and
- 2 Any of the following:
  - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
  - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
  - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations\*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
  - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
  - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with \* are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex\*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex\*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with \* are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
  - 2.1 Both:
    - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
    - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
  - 2.2 Both:

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	Subsidy	Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price)		
	 \$	Per 🗸	Manufacturer

continued...

- 2.2.1 Vigabatrin is contraindicated; and
- 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex\*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with \* are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

erio zimo o opociai rialiioni, oco orini io zonom i			
Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg	248.20	50	✓ Tacrolimus Sandoz
. •			

#### ⇒SA1745 Special Authority for Subsidy

**Initial application — (organ transplant)** only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications\*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with \* are unapproved indications

#### **JAK** inhibitors

### ⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
  - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
  - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
  - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 3.2 Both:
  - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
  - 3.2.2 Either:
    - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
    - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Antiallergy Preparations**

## **Allergic Emergencies**

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

#### ⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

**Renewal** from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

# Allergy Desensitisation

## **⇒SA1367** Special Authority for Subsidy

**Initial application** only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

**Renewal** only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see S	A1367 above - F	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluer	nt 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see	SA1367 above	- Retail pharr	macy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze	005.00	4.00	411
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	205.00	1 OP	√ Alboy
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml		TOP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent		1 OP	✓ Venomil S29
uneu venom, with unuell		i OF	V CHOIIII 023

227

	Subsidy		Fully	Brand or
	(Manufacturer's Price		idised	
	\$	Per		Manufacturer
Authistonius				
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1 12	100	1	Zista
* Oral liq 1 mg per ml		200 ml	_	Histaclear
CHLORPHENIRAMINE MALEATE				<u></u>
* Oral liq 2 mg per 5 ml	0.07	500 ml	./	Histafen
	9.37	300 1111	•	пізіаіен
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg		40		
	(8.40)			Polaramine
	1.01	20		
4. 0 10 0 - 1	(5.99)			Polaramine
* Oral liq 2 mg per 5 ml		100 ml		<b>.</b> .
	(10.29)			Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
	(8.23)			Telfast
* Tab 120 mg	4.74	10		
	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
LORATADINE				
* Tab 10 mg	1.69	100	1	Lorafix
* Oral liq 1 mg per ml		100 ml	1	Haylor syrup
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg	1.39	50	1	Allersoothe
* Tab 25 mg		50		Allersoothe
* Oral lig 1 mg per 1 ml		100 ml		Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a		5		Hospira
ing to mig por mi, this amposito of the only available of a		•		Поорши
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01 2	00 dose OP	1	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free	8.54 2	00 dose OP		Beclazone 50
Aerosol inhaler, 100 mcg per dose		00 dose OP		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		00 dose OP		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free	22.67 2	00 dose OP	1	Beclazone 250
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	1	Pulmicort
				Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	1	Pulmicort
				Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	1	Pulmicort
				Turbuhaler

	Subsidy		Fully	Brand or
	(Manufacturer's	Price) Subs	idised	Generic
	\$	Per		Manufacturer
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flix	cotide
Powder for inhalation, 50 mcg per dose		60 dose OP		kotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP		cotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP		cotide
Aerosol inhaler, 250 mcg per dose	24.62	120 dose OP	✓ Flix	
Powder for inhalation, 250 mcg per dose		60 dose OP		kotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	ts			
EFORMOTEROL FUMARATE				
EFORMOTEROL FUMARATE	ioo 00.64	CO doos		
Powder for inhalation, 12 mcg per dose, and monodose dev		60 dose	Г	ا:اد د
	(35.80)		FOI	radil
EFORMOTEROL FUMARATE DIHYDRATE				
Powder for inhalation 4.5 mcg per dose, breath activated				
(equivalent to eformoterol fumarate 6 mcg metered dose	e) 10.32	60 dose OP		
	(16.90)		Ox	is Turbuhaler
INDACATEROL				
Powder for inhalation 150 mcg	61.00	30 dose OP	✓ On	brez Breezhaler
Powder for inhalation 300 mcg		30 dose OP	✓ On	brez Breezhaler
SALMETEROL				
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	<b>√</b> Sα	revent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP		revent Accuhaler
1 owder for initial attorit, so may per acce, breath activated		00 0000 01	- 00.	event Addunate
Inhaled Corticosteroids with Long-Acting Beta-	-Adrenocept	tor Agonists		
BUDESONIDE WITH EFORMOTEROL				
Powder for inhalation 160 mcg with 4.5 mcg eformoterol				
fumarate per dose (equivalent to 200 mcg budesonide v	with			
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ Du	oResp Spiromax
,		120 dose OF	• Du	unesp Spirolliax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma				
per dose (equivalent to 400 mcg budesonide with 12 mc	.g			
eformoterol fumarate metered dose) – No more than 2 dose per day	92.50	120 dose OP	✓ D	oResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vai	
· · · · · · · · · · · · · · · · · · ·		120 dose OP		
Powder for inhalation 100 mcg with eformoterol fumarate 6 r	11cg33.74	120 dose OF		mbicort urbuhaler 100/6
Agranal inhalar 200 mag with aformatoral furnariate 6 mag	01.40	120 dose OP	✓ Vai	
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg				
Powder for inhalation 200 mcg with eformoterol fumarate 6 r	110y44.08	120 dose OP		mbicort 'urbuhaler 200/6
Douglas for inholation 400 sees with afarmatary for				ui buildiei 200/0
Powder for inhalation 400 mcg with eformoterol fumarate	44.00	60 46 00	./ 0	mbiaant
12 mcg - No more than 2 dose per day	44.08	60 dose OP		mbicort
			ı	urbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			_	
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Bre	eo Ellipta

Subsidy (Manufacturer's I	Price) Subs	Fully idised	Brand or Generic Manufacturer
•			
	120 dose OP 120 dose OP	_	Seretide Seretide
33.74	60 dose OP	<b>√</b> §	Seretide Accuhaler
44.08	60 dose OP	✓ 9	Seretide Accuhaler
	150 ml	<b>✓</b> \	<u>/entolin</u>
	10 5		/entolin /entolin
3.80	200 dose OP		Respigen SalAir
(6.20)		١	/entolin
8.96	20	<b>√</b> <u>I</u>	<u>Asthalin</u>
9.43	20	✓ <u>I</u>	<u>Asthalin</u>
22.20	120 dose OP	<b>✓</b> E	Bricanyl Turbuhaler
	200 dose OP	<b>√</b> 1	Atrovent
b	20	_	Jnivent
holineraic A	aents	_	
J - 1			
	200 dose OP	<b>√</b> г	Duolin HFA
12.13	200 0036 OF	٠ ١	/uviii i ii A
	(Manufacturer's I \$ \$	(Manufacturer's Price)         Subsequence           \$         120 dose OP	(Manufacturer's Price)         Subsidised Per

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

# **Long-Acting Muscarinic Antagonists**

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

### UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

# Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

#### ⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL − Special Authority see SA1584 above − Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg.....81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see SA1584 above – Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg ......81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg .......77.00 30 dose OP ✓ Anoro Ellipta

### **Antifibrotics**

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

### ⇒SA2012 Special Authority for Subsidy

**Initial application — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with pirfenidone; or
  - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

<ul><li>Esbriet</li></ul>	90	Tab 801 mg3,645.00
✓ Esbriet	90	Tab 267 mg1,215.00

### ⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
  - 5.1 The patient has not previously received treatment with nintedanib; or
  - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
  - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

**Renewal — (idiopathic pulmonary fibrosis)** only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

				_
	Subsidy (Manufacturer's Price)		Fully Subsidised	
	(Manufacturer's Frice)	Per	Jubsidised ✓	Manufacturer
Leukotriene Receptor Antagonists				
MONTELUKAST				
* Tab 4 mg	4.25	28		Montelukast Mylan
* Tab 5 mg		28		Montelukast Mylan
* Tab 10 mg	3.95	28		Montelukast Mylan
Methylxanthines				
AMINOPHYLLINE				
* Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a				
PSO	180.00	5	✓	DBL Aminophylline
THEOPHYLLINE				. ,
* Tab long-acting 250 mg	23.02	100	1	Nuelin-SR
* Oral liq 80 mg per 15 ml		00 m		Nuelin
				<u></u>
Mucolytics				
DORNASE ALFA - Special Authority see SA1978 below - Retail	pharmacy			
Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	1	Pulmozyme
<b>⇒SA1978</b> Special Authority for Subsidy				•
Initial application — (cystic fibrosis) only from a respiratory ph	vsician or paediatrici	an A	nnrovals	valid for 12 months for

**Initial application** — **(cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
  - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
  - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
  - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or</p>
  - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

**Renewal — (cystic fibrosis)** only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

### ⇒SA2017 Special Authority for Subsidy

**Initial application** only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
  - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele: or
  - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N

	Subsidy (Manufacturer's \$	Price) Subsi	Fully dised	Brand or Generic Manufacturer
continued and S549R) in the CFTR gene on at least 1 alle	ele; and			
<ul> <li>3 Patients must have a sweat chloride value of at least 6 sweat collection system; and</li> <li>4 Treatment with ivacaftor must be given concomitantly</li> <li>5 Patient must not have an acute upper or lower respirat (including antibiotics) for pulmonary disease in the last</li> <li>6 The dose of ivacaftor will not exceed one tablet or one</li> <li>7 Applicant has experience and expertise in the manage</li> </ul>	on mmol/L by quanties or infection, pulm 4 weeks prior to cally	apy for this conditionary exacerbations treat	tion; ar on, or	nd changes in therapy
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	<b>✓</b> <u>B</u>	Biomed
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE				
Metered aqueous nasal spray, 50 mcg per dose		200 dose OP	_	<u>iteroClear</u>
Metered aqueous nasal spray, 100 mcg per dose	2.84	200 dose OP	✓ <u>S</u>	<u>iteroClear</u>
FLUTICASONE PROPIONATE  Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	<b>✓</b> <u>F</u>	lixonase Hayfever & Allergy
IPRATROPIUM BROMIDE				
Aqueous nasal spray, 0.03%	5.23	15 ml OP	<b>√</b> <u>U</u>	<u>Inivent</u>
Respiratory Devices				
MASK FOR SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO				
c) Only for children aged six years and under Small	2.20	1	<b>√</b> e	-chamber Mask
PEAK FLOW METER		•	_	
a) Up to 25 dev available on a PSO				
b) Only on a PSO				
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	lini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				

b) Only on a PSO

1

✓ e-chamber Turbo

✓ e-chamber La Grande

✓ Volumatic

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

# **Respiratory Stimulants**

**CAFFEINE CITRATE** 

Oral liq 20 mg per ml (10 mg base per ml).......15.10 25 ml OP ✓ Biomed



	Subsidy		Fully	Brand or
	(Manufacturer's Pri \$	ce) Subs Per	sidised •	Generic Manufacturer
For Propositions				
Ear Preparations				
FLUMETASONE PIVALATE	4.40	7.5 ml OD		annaman Vinform
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	• 1	Locacorten-Viaform ED's
			<b>✓</b> I	Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTATI	N		
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate	F 10	7.5 ml OD		/anaaamh
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	•	Kenacomb
Ear/Eye Preparations				
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN				
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and				
gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP		Sofradex
FRAMYCETIN SULPHATE	(3.27)		,	Jonadex
Ear/Eye drops 0.5%	4.13	8 ml OP		
	(8.65)		5	Soframycin
Eye Preparations				
Eye preparations are only funded for use in the eye, unless expli	icitly stated otherw	ise.		
Anti-Infective Preparations				
ACICLOVIR				
* Eye oint 3%	14.88	4.5 g OP	✓ <u>\</u>	<u>/iruPOS</u>
CHLORAMPHENICOL	4 55	5 - OD		S
Eye oint 1% Eye drops 0.5%		5 g OP 10 ml OP		<u>Devatis</u> Chlorafast
Funded for use in the ear*. Indications marked with * a			-	
CIPROFLOXACIN			_	
Eye drops 0.3% – Subsidy by endorsement		5 ml OP	-	Ciprofloxacin Teva
for the second line treatment of chronic suppurative otiti		,		
Note: Indication marked with a * is an unapproved indication		•	•	3,
GENTAMICIN SULPHATE	44.40	5 LOD		
Eye drops 0.3%	11.40	5 ml OP	•	Genoptic
PROPAMIDINE ISETHIONATE  * Eye drops 0.1%	2.97	10 ml OP		
- Lyo diope 0.1 /0	(14.55)	10 1111 01	E	Brolene
SODIUM FUSIDATE [FUSIDIC ACID]				
Eye drops 1%	5.29	5 g OP	<b>✓</b> I	Fucithalmic
TOBRAMYCIN Eye oint 0.3%	10.45	2 E a OP	./ -	Tobrex
Eye drops 0.3%		3.5 g OP 5 ml OP		l obrex Fobrex

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

# **Corticosteroids and Other Anti-Inflammatory Preparations**

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,44	4.50	1	<ul><li>Ozurdex</li></ul>

#### ⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
  - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
  - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

#### All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

#### DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM Eye drops 0.1%	8.80	5 ml OP	✓ <u>Voltaren Ophtha</u>
FLUOROMETHOLONE  * Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20		✓ Flucon

	Subsidy (Manufacturer's Pric	e) Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
• •	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	<b>✓</b> L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	5.93	10 ml OP	<b>✓</b> P	rednisolone-AFT
•	7.00	5 ml OP	<b>√</b> P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below -	Retail phar	macv	
Eye drops 0.5%, single dose (preservative free)		20 dose	•	linims Prednisolone

## ⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

#### Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE Eye drops 2%	1.79	5 ml OP	✓ <u>Rexacrom</u>
Glaucoma Preparations - Beta Blockers			
BETAXOLOL  * Eye drops 0.25%  * Eye drops 0.5%  TIMOLOL  * Eye drops 0.25%  * Eye drops 0.5%  * Eye drops 0.5%  * Eye drops 0.5%, gel forming	7.50 1.81 2.04	5 ml OP 5 ml OP 5 ml OP 5 ml OP 2.5 ml OP	✓ Betoptic S ✓ Betoptic  ✓ Arrow-Timolol ✓ Arrow-Timolol ✓ Timoptol XE
Glaucoma Preparations - Carbonic Anhydrase Inhil	bitors		
ACETAZOLAMIDE  * Tab 250 mg  BRINZOLAMIDE	17.03	100	✓ Diamox
* Eye drops 1%  DORZOLAMIDE HYDROCHLORIDE	7.30	5 ml OP	✓ <u>Azopt</u>
* Eye drops 2%	9.77 (17.44)	5 ml OP	Trusopt
DORZOLAMIDE WITH TIMOLOL  * Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓ <u>Dortimopt</u>
Glaucoma Preparations - Prostaglandin Analogues			
BIMATOPROST  * Eye drops 0.03%	5.95	3 ml OP	✓ Bimatoprost  Multichem

	Subsidy (Manufacturer's P \$		Fully Brand or dised Generic  Manufacturer
* Eye drops 0.005%	1.82	2.5 ml OP	✓ <u>Teva</u>
* Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Travatan</u>
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE  * Eye drops 0.2%	4.29	5 ml OP	✓ <u>Arrow-Brimonidine</u>
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE  * Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ Arrow - Lattim
PILOCARPINE HYDROCHLORIDE  # Eye drops 1%*  # Eye drops 2%		15 ml OP 15 ml OP	✓ Isopto Carpine ✓ Isopto Carpine
* Eye drops 4%  Subsidised for oral use pursuant to the Standard Formu	7.99	15 ml OP	✓ Isopto Carpine
* Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine

# **⇒SA0895** Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
ATROPINE SULPHATE		
* Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE  * Eye drops 1%8.76	15 ml OP	✓ Cyclogyl
TROPICAMIDE	13 1111 01	• Cyclogyi
* Eye drops 0.5%7.15	15 ml OP	✓ Mydriacyl
* Eye drops 1%	15 ml OP	✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 243		
HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	✓ Methopt
HYPROMELLOSE WITH DEXTRAN		
* Eye drops 0.3% with dextran 0.1%	15 ml OP	✓ Poly-Tears

<sup>▲</sup>Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

### **Preservative Free Ocular Lubricants**

#### ⇒SA1388 Special Authority for Subsidy

**Initial application** from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp of severe secretory dry eye; and
- 2 Either:
  - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
  - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA1388 above - Retail pharma	acy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authority s	see SA1388 al	oove – Retail p	harmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	<ul> <li>Systane Unit Dose</li> </ul>
SODIUM HYALURONATE [HYALURONIC ACID] - Special Authorit	y see SA1388	above - Retai	il pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ <u>Hylo-Fresh</u>
Hylo-Fresh has a 6 month expiry after opening. The Pharm	acy Procedure	s Manual rest	riction allowing one bottle per
month is not relevant and therefore only the prescribed dosa	age to the near	rest OP may b	e claimed.

# **Other Eye Preparations**

NAPHAZOLINE HYDROCHLORIDE  * Eye drops 0.1%	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.20	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT  * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE  Eve oint 138 mcg per g	5 a OP	✓ VitA-POS

✓ BSF Vigisom

Pharma \$29

	Subsidy (Manufacturer's Price) \$	) Per	Fully Subsidised	Brand or Generic Manufacturer	
Various					
PHARMACY SERVICES May only be claimed once per patient.  * Brand switch fee	4.50	1 fee	_	3SF Cinacalcet Devatis 3SF Comtan	

- a) The Pharmacode for BSF Vigisom is 2634112 see also page 136
- b) The Pharmacode for BSF Cinacalcet Devatis is 2634120 see also page 78
- c) The Pharmacode for BSF Comtan is 2634139 see also page 118

(BSF Cinacalcet Devatis Brand switch fee to be delisted 1 July 2022)

(BSF Comtan Brand switch fee to be delisted 1 July 2022)

(BSF Vigisom Brand switch fee to be delisted 1 July 2022)

# Agents Used in the Treatment of Poisonings

#### **Antidotes**

ACETYLCYSTEINE		
Inj 200 mg per ml, 10 ml ampoule58.76	10	✓ DBL Acetylcysteine
		✓ Martindale

#### NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

*	Inj 400 mcg per ml, 1 ml ampoule	22.60	5	DBL Naloxone
				Hydrochloride

#### Removal and Elimination

#### CHARCOAL

*	Oral liq 50 g per 250 m	43.50	250 ml OP	<ul><li>Carbosorb-X</li></ul>
---	-------------------------	-------	-----------	-------------------------------

- a) Up to 250 ml available on a PSO
- b) Only on a PSO

#### DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

# Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

## ⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
  - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2\*; or



Subsidy		Fully	Brand or
(Manufacturer's Price)	_	Subsidised	Generic
\$	Per		Manufacturer

continued...

- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

**Renewal** only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2\* and liver MRI T2\* levels.

DEFERIPRONE - Special Authority see SA1480 below - F	Retail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

## ⇒SA1480 Special Authority for Subsidy

**Initial application** only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

#### Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

* Inj 500 mg vial	151.31	10	✓ DBL  Desferrioxamine  Mesylate for Inj  BP
SODIUM CALCIUM EDETATE			
* Inj 200 mg per ml, 5 ml	53.31	6	
•	(156.71)		Calcium Disodium

Versenate

# **Standard Formulae**

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml)	00		
Codeine phosphate	60 mg	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Glycerol	40 ml	mg per ml)	
Preservative	qs	Phenobarbitone Sodium	400 mg
Water	to 100 ml	Glycerol BP	4 ml
CODEINE LINCTUS (15 mg per 5 ml)		Water	to 40 ml
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol	40 ml	Pilocarpine 4% eye drops	qs
Preservative	qs	Preservative	qs
Water	to 100 ml	Water	to 500 ml
Tracor	10 100 1111	(Preservative should be used if quantity supplied is	
FOLINIC MOUTHWASH		than 5 days.)	ioi more
Calcium folinate 15 mg tab	1 tab	illail 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 500 ml	Methylcellulose	5 g
(Preservative should be used if quantity supplied is	for more	Preservative	qs
than 5 days. Maximum 500 ml per prescription.)		Water	to 500 ml
than o days. Maximum ooo iii por proconption.		(Preservative should be used if quantity supplied is	
METHADONE MIXTURE		than 5 days. Maximum 500 ml per prescription.)	ioi illoic
Methadone powder	qs	than 5 days. Maximum 500 mi per prescription.	
Glycerol	qs	SODIUM CHLORIDE ORAL LIQUID	
Water	to 100 ml	Sodium chloride inj 23.4%, 20 ml	qs
		Water	gs .
METHYL HYDROXYBENZOATE 10% SOLUTION		(Only funded if prescribed for treatment of hyponatr	•
Methyl hydroxybenzoate	10 g	(,,,,,,	,
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqu	uid mixture)	Vancomycin 500 mg injection	10 vials
	,	Glycerol BP	40 ml
OMEPRAZOLE SUSPENSION		Water	to 100 ml
Omeprazole capules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	ım difficile
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	
Water	to 100 ml	3	

## EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy		Fully Brand or
	(Manufacturer's Pr	rice) Sub	sidised Generic  Manufacturer
Extemporaneously Compounded Preparations	and Galenica	ils	
CODEINE PHOSPHATE - Safety medicine; prescriber may de		g frequency	
Powder – Only in combination		25 g	December
Only in extemporaneously compounded codeine linctus	(90.09)		Douglas
COLLODION FLEXIBLE	'•		
Note: This product is no longer being manufactured by the determined.	supplier and will b	e delisted fror	m the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination			
Only in extemporaneously compounded oral mixtures.			
Soln		100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination	1		
Only in combination with Ora-Plus. Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination		1701111	<u> </u>
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prep	arations.		
METHADONE HYDROCHLORIDE  a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing f			
d) Extemporaneously compounded methadone will only be	reimbursed at the	rate of the ch	neapest form available
(methadone powder, not methadone tablets). Powder	7 84	1 g	✓ AFT
METHYL HYDROXYBENZOATE		' 9	· All
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE		-	
Powder		100 g	✓ <u>MidWest</u>
Suspension – Only in combination		473 ml	✓ <u>Ora-Plus</u>
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCI			√ Ove Bland CE
Suspension		473 ml	✓ <u>Ora-Blend SF</u>
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Or Suspension	•	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM		4701111	<u> </u>
Powder – Only in combination	52.50	10 g	✓ MidWest
•	325.00	100 g	✓ MidWest
Only in children up to 12 years			
PROPYLENE GLYCOL	100/	_	
Only in extemporaneously compounded methyl hydroxyben Liq		n. 500 ml	✓ Midwest
SODIUM BICARBONATE		000 1111	· manou
Powder BP – Only in combination	10.05	500 g	✓ Midwest
Only in aytomporangously compounded amonrazale ar		enancion	<del></del>

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

# EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Su Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparation		500 ml	✓ <u>M</u>	idwest	
WATER Tap - Only in combination	0.00	1 ml	<b>✓</b> Ta	ap water	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

# **Nutrient Modules**

## Carbohydrate

#### ⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

**Initial application — (Indications other than cystic fibrosis or renal failure)** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

# Carbohydrate And Fat

# **⇒SA1376** Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
  - 2.1 cancer in children; or
  - 2.2 faltering growth; or
  - 2.3 bronchopulmonary dysplasia; or
  - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

#### Fat

# **⇒SA1523** Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 247

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

## **Protein**

# **⇒SA1524** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	<ul> <li>Special Authority see SA1524 above – Hospital p</li> </ul>	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy
(Manufacturer's Price) Subsider
\$ Per

Fully Subsidised

Brand or Generic Manufacturer

# **Oral and Enteral Feeds**

#### Diabetic Products

#### ⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid	500 ml OP	✓ Glucerna Select
7.50	1,000 ml OP	✓ Diason RTH
		✓ Nutrison Advanced
		Diason

(Diason RTH Liquid to be delisted 1 December 2022)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 mi OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

### **Fat Modified Products**

#### ⇒SA1525 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see SA1525 above – Hospital pharmacy [HP3]

✓ fully subsidised 249

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

# Paediatric Products For Children Awaiting Liver Transplant

#### ⇒SA1098 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

## Paediatric Products For Children With Chronic Renal Failure

# ⇒SA1099 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Powder .......54.00 400 g OP 

✓ Kindergen

#### **Paediatric Products**

# ⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
  - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
  - 2.2 any condition causing malabsorption; or
  - 2.3 faltering growth in an infant/child; or
  - 2.4 increased nutritional requirements; or
  - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

Subsidy		Fully	Brand or	
(Manufacturer's Price	e) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA Liquid		ne previous pag 500 ml OP	e – Hospital pharmacy [HP3]  ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA13 Liquid		previous page 500 ml OP	<ul><li>Hospital pharmacy [HP3]</li><li>✓ Nutrini RTH</li><li>✓ Pediasure RTH</li></ul>
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Aut pharmacy [HP3]	thority see	SA1379 on the	previous page – Hospital
Liquid	.6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED – Special Authority see SA1379 on the previ- Powder (vanilla)		- Hospital phari 850 g OP	macy [HP3]  Pediasure
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see \$A1379 Liquid (strawberry) Liquid (vanilla)	.1.60	revious page – 200 ml OP 200 ml OP	Hospital pharmacy [HP3]  Fortini Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 (Liquid (chocolate)	.1.07 .1.07 .1.07	vious page – Ho 200 ml OP 200 ml OP 200 ml OP 250 ml OP	ospital pharmacy [HP3]  Pediasure Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authori pharmacy [HP3]	ty see SA1	1379 on the pre	vious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	.1.60 .1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		age – Hospital į 400 a OP	oharmacy [HP3]  Peptamen Junior

#### **Renal Products**

## ⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 251

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		ous page – Hos 220 ml OP	<b>✓</b> 1	harmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110		s page – Hospit 237 ml OP	al pha	armacy [HP3]
Liquid, 200 ml bottle	(3.31) 11.52	4 OP	١	NovaSource Renal
Liquid (apricot) 125 mlLiquid (caramel) 125 ml		4 OP 4 OP	✓ [	NovaSource Renal Renilon 7.5 Renilon 7.5
(NovaSource Renal Liquid to be delisted 1 September 2022)				

## **Specialised And Elemental Products**

## **⇒SA1377** Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML - \$	Special Authority sec	e SA1377 above	- Hospital pharmacy [HP3]
Liquid	18.06	1,000 ml OP	✓ Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority	see SA1377 above	- Hospital pharn	nacy [HP3]
Liquid (grapefruit), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority se	e SA1377 above – I	Hospital pharma	cy [HP3]
Powder (unflavoured)	4.50	80 g OP	✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML - Special A	authority see SA137	7 above – Hospi	tal pharmacy [HP3]
Liquid	12.04	1,000 ml OP	✓ Peptisorb

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per 🗸 Brand or Generic Manufacturer

### Paediatric Products For Children With Low Energy Requirements

### ⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

### Standard Supplements

### ⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
  - 2.1 The patient has a condition causing malabsorption; or
  - 2.2 The patient has failure to thrive; or
  - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subs (Manufactur		
\$	Per	Manufacturer

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m<sup>2</sup>; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

**Initial application — (Short-term medical condition)** from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
  - 5.1 Pregnant; and
  - 5.2 Any of the following:
    - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
    - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
    - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm<sup>3</sup>); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

# **SPECIAL FOODS**

	Subsidy (Manufacturer's \$		Fully Brand or dised Generic  Manufacturer
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 Liquid		lospital pharmac 250 ml OP 1,000 ml OP	y [HP3]  ✓ Ensure Plus HN  ✓ Ensure Plus RTH  ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 or Liquid		spital pharmacy 250 ml OP 1,000 ml OP	[HP3]  ✓ Isosource Standard  ✓ Nutrison Standard  RTH  ✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Author Liquid	,	on page 253 – Ho 1,000 ml OP	ospital pharmacy [HP3]  Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority : Liquid		oage 253 - Hosp 1,000 ml OP	ital pharmacy [HP3]  ✓ Jevity RTH  ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority Liquid		page 253 – Hos 1,000 ml OP	pital pharmacy [HP3]  ✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority Liquid		page 253 – Hos 1,000 ml OP	pital pharmacy [HP3]  ✓ Jevity HiCal RTH  ✓ Nutrison Energy  Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on pa Powder (chocolate)	•	al pharmacy [HP 840 g OP	3] ✓ Sustagen Hospital Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	<ul><li>✓ Ensure</li><li>✓ Sustagen Hospital</li><li>Formula Active</li></ul>
	26.00	850 g OP	✓ Ensure

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

### ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 253 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 253 — Hospital pharmacy [HP3]
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.26)	200 0.	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

## **High Calorie Products**

# ⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	<b>✓</b>	

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
  - 1.1 any condition causing malabsorption; or
  - 1.2 faltering growth in an infant/child; or
  - 1.3 increased nutritional requirements; or
  - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
  - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
  - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

### **Food Thickeners**

#### ⇒SA1106 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price	) Sub	Fully sidised	Brand or Generic	
	\$	Per	1	Manufacturer	
FOOD THICKENER – Special Authority see SA1106 on the pre	vious page – Hospita	l pharmac	y [HP3]		
Powder	6.53	00 g OP	✓ N	utilis	
	7.25	80 g OP	✓ F	eed Thickener	
				Karicare Aptamil	

### **Gluten Free Foods**

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

### ⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

pharmacy [HP3]	
1,000 g OP	
•	Healtheries Simple Baking Mix
oharmacy [HP3]	
1,000 g OP	
	NZB Low Gluten Bread Mix
	Horleys Bread Mix
nacy [HP3]	
2,000 g OP	
	Horleys Flour
	1,000 g OP  charmacy [HP3] 1,000 g OP

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	ce) Su Per	ıbsidised •	Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	orevious page – Ho	ospital pha	rmacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)			Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)			Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)			Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)			Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)			Orgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)			Orgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)			Orgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)			Orgran

# Foods And Supplements For Inborn Errors Of Metabolism

### ⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

# **Supplements For Homocystinuria**

## **Supplements For MSUD**

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Por	1	Manufacturer

# **Supplements For PKU**

AMINOACID FORMULA WITHOUT PHENYLALANINE	<ul> <li>Special Authority see SA1</li> </ul>	108 on the pre	evious page	– Hospital
pharmacy [HP3]				
· · · ·		0-	<i>-</i>	

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	<ul> <li>Easiphen Liquid</li> </ul>
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

# Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page	ge – Hospital	pharmacy [HP3]
Powder8.22	500 g OP	<ul><li>Loprofin Mix</li></ul>

OW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

LOW PROTEIN PASTA - Special Authority see SATTOS on the prev	/lous page – i	nospilai priami	acy [nP3]
Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	<ul><li>Loprofin</li></ul>
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 a OP	✓ Loprofin

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

### Infant Formulae

### For Williams Syndrome

### ⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

#### Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

### **Gastrointestinal and Other Malabsorptive Problems**

MINO ACID FORMULA – Special Authority see SA2092 below – H Powder		400 g OP ✓	Alfamino Alfamino Junior
Powder (unflavoured)	53.00	400 g OP 🗸	Elecare Elecare LCP Neocate Gold Neocate Junior Unflavoured
Powder (vanilla)	53.00	400 g OP 🗸	Neocate SYNEO Elecare Neocate Junior Vanilla

#### ⇒SA2092 Special Authority for Subsidy

**Initial application** — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
  - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or alleray or malabsorotion; and
  - 6.2 Either:
    - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
  - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
      - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

#### Either:

- 1 Both:
  - 1.1 Patient has IgE mediated allergy; and
  - 1.2 All of the following:
    - 1.2.1 Patient remains allergic to cow's milk; and
    - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
    - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
    - 1.2.4 Amino acid formula is required for a nutritional deficit; and
    - 1.2.5 It has been more than three months from the previous approval; or

### 2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
  - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
  - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
  - 2.2.3 Amino acid formula is required for a nutritional deficit; and
  - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
  - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
  - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
  - 2.2 Eosinophilic oesophagitis; or
  - 2.3 Ultra-short gut; or
  - 2.4 Severe Immune deficiency; or
  - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
  - 2.6 Both:
    - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
    - 2.6.2 Either:
      - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
      - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

#### SA1953 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
  - 2.1 Severe malabsorption; or
  - 2.2 Short bowel syndrome; or
  - 2.3 Intractable diarrhoea: or
  - 2.4 Biliary atresia; or
  - 2.5 Cholestatic liver diseases causing malabsorption: or
  - 2.6 Cystic fibrosis; or
  - 2.7 Proven fat malabsorption; or
  - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
  - 2.9 Intestinal failure: or
  - 2.10 Both:
    - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

SYNEO 2

(A)	Subsidy //anufacturer's Price)	Subsi	Fully	Brand or Generic
ζ.,	\$	Per	✓	Manufacturer

continued...

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
  - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
  - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

### ⇒SA1557 Special Authority for Subsidy

**Initial application** only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
  - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
  - 1.2 Either:
    - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
    - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption: or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure: or
- 11 All of the following:
  - 11.1 For step down from Amino Acid Formula: and
  - 11.2 The infant is currently receiving funded amino acid formula; and
  - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
  - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IqE mediated allergic reaction.

continued...



Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	Jubsiuiseu	Manufacturer

**Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

### Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML − Special Authority see SA1698 below − Hospital pharmacy [HP3] Liquid.......2.35 125 ml OP ✓ Infatrini

### SA1698 Special Authority for Subsidy

**Initial application** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

**Renewal** only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

# **Ketogenic Diet**

#### ⇒SA1197 Special Authority for Subsidy

**Initial application** only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

**Renewal** only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1
			✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

### **SECTION I: NATIONAL IMMUNISATION SCHEDULE**

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

# **Vaccinations**

#### BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00 10 

✓ BCG Vaccine

#### DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
  Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

### DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE			
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	ND HAEMOPHILUS I	NFLUENZAE TY	PE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of			and a considerable and a second
2) An additional four doses (as appropriate) are funded for			
10 who are patients post haematopoietic stem cell trans post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up p		'	
to complete full primary immunisation. Please refer to the Imi			
programmes.			
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10 <b>✓</b> <u>In</u>	<u>ifanrix-hexa</u>
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)imi			
transplantation, or chemotherapy; functional asplenic; pr			olid organ transplant, pre-
or post cochlear implants, renal dialysis and other sever	, , , , , , , , , , , , , , , , , , , ,	•	val madiaina physisian ar
<ol> <li>For use in testing for primary immunodeficiency disease paediatrician.</li> </ol>	s, on the recommend	allon of an intern	iai medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg	:		
prefilled syringe plus vial 0.5 ml		1 🗸 H	iberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver di	sease; or		
3) One dose of vaccine for close contacts of known hepatit	is A cases.		

Inj 1440 ELISA units in 1 ml syringe......0.00

✓ Havrix

✓ Havrix Junior

1

		NATIONAL	IIVIIVI	UNISATI	ON SCHEDULE
		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B	RECOMBINANT VACCINE - [Xpharm]				
	g per 0.5 ml prefilled syringe		1	✓ E	ngerix-B
	led for patients meeting any of the following criteria:				
	for household or sexual contacts of known acute h				s; or
	for children born to mothers who are hepatitis B su				and the second are an above
3)	for children up to and under the age of 18 years in				achieved a positive
4)	serology and require additional vaccination or requ for HIV positive patients; or	life a primary course of	or vacc	ination, or	
	for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	course: or			
	for patients following immunosuppression; or	,			
8)	for solid organ transplant patients; or				
	for post-haematopoietic stem cell transplant (HSC	T) patients; or			
10)	following needle stick injury.				
Inj 20 mc	g per 1 ml prefilled syringe	0.00	1	<b>√</b> E	ngerix-B
	led for patients meeting any of the following criteria:			_	
1)	for household or sexual contacts of known acute h	epatitis B patients or h	nepatit	is B carriers	s; or
	for children born to mothers who are hepatitis B su				
3)	for children up to and under the age of 18 years in				achieved a positive
4\	serology and require additional vaccination or requ	ire a primary course o	of vacc	cination; or	
	for HIV positive patients; or for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	ourse. or			
	for patients following immunosuppression; or	ouise, or			
,	for solid organ transplant patients; or				
	for post-haematopoietic stem cell transplant (HSC	T) patients; or			
	following needle stick injury; or				
	for dialysis patients; or				
12)	for liver or kidney transplant patients.				
HUMAN PAP	ILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 5	58) VACCINE [HPV] -	- [Xph	arml	
	e following:	,	٠,	•	
1) Max	ximum of two doses for children aged 14 years and	under; or			
	ximum of three doses for patients meeting any of th				
	) People aged 15 to 26 years inclusive; or				
2	) Either:				
	People aged 9 to 26 years inclusive				
	1) Confirmed HIV infection; or				
a\ •-	2) Transplant (including stem cell) patients: or				
3) Max	ximum of four doses for people aged 9 to 26 years i	nclusive post chemoth	nerapy	<i>'</i>	
Inj 270 m	cg in 0.5 ml syringe	0.00	10	<b>√</b> <u>G</u>	ardasil 9

Subsidy	Sı	Fully	Brand or
(Manufacturer's Price)		ubsidised	Generic
\$	Per	•	Manufacturer

#### INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

### A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
  - a) ischaemic heart disease, or
  - b) congestive heart failure, or
  - c) rheumatic heart disease, or
  - d) congenital heart disease, or
  - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
  - a) asthma, if on a regular preventative therapy, or
  - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
  - a) autoimmune disease, or
  - b) immune suppression or immune deficiency, or
  - c) HIV, or
  - d) transplant recipients, or
  - e) neuromuscular and CNS diseases/disorders, or
  - f) haemoglobinopathies, or
  - g) on long term aspirin, or
  - h) have a cochlear implant, or
  - i) errors of metabolism at risk of major metabolic decompensation, or
  - j) pre and post splenectomy, or
  - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)......110.00 10 
✓ Afluria Quad
(2022 formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

c)

### A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
  - i) have any of the following cardiovascular diseases:
    - a) ischaemic heart disease, or
    - b) congestive heart failure, or
    - c) rheumatic heart disease, or
    - d) congenital heart disease, or
    - e) cerebo-vascular disease; or
  - ii) have either of the following chronic respiratory diseases:
    - a) asthma, if on a regular preventative therapy, or
    - b) other chronic respiratory disease with impaired lung function; or
  - iii) have diabetes; or
  - iv) have chronic renal disease; or
  - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
  - vi) have any of the following other conditions:
    - a) autoimmune disease, or
    - b) immune suppression or immune deficiency, or
    - c) HIV, or
    - d) transplant recipients, or
    - e) neuromuscular and CNS diseases/disorders, or
    - f) haemoglobinopathies, or
    - g) are children on long term aspirin, or
    - h) have a cochlear implant, or
    - i) errors of metabolism at risk of major metabolic decompensation, or
    - i) pre and post splenectomy, or
    - k) down syndrome, or
  - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

#### MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

### A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression: or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

# MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Fither:

- A) Any of the following:
  - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
  - 2) One dose for close contacts of meningococcal cases of any group; or
  - 3) One dose for person who has previously had meningococcal disease of any group; or
  - 4) A maximum of two doses for bone marrow transplant patients; or
  - 5) A maximum of two doses for person pre- and post-immunosuppression\*; or
- B) Both:
  - 1) Person is aged between 13 and 25 years, inclusive; and
  - 2) Either
    - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
    - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

\*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

NATIONAL IMMUNISATION SCHEDULE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression\*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients: or v) up to two doses for person pre- and post-immunosuppression\*. \*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression\*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. \*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B. 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10

Synflorix

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
	Per	<b>✓</b>	Manufacturer	

### PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
  - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
  - b) with primary immune deficiencies; or
  - c) with HIV infection; or
  - d) with renal failure, or nephrotic syndrome; or
  - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
  - f) with cochlear implants or intracranial shunts; or
  - g) with cerebrospinal fluid leaks; or
  - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
  - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
  - j) pre term infants, born before 28 weeks gestation; or
  - k) with cardiac disease, with cyanosis or failure; or
  - I) with diabetes; or
  - m) with Down syndrome; or
  - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Ini 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4.

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]		
<ol> <li>Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochl</li> <li>All of the following:</li> </ol>	tional asplenia, pre- or p	post-solid organ t	ransplant, renal dialysis,
<ul> <li>a) Patient is a child under 18 years for (re-)immur</li> <li>b) Treatment is for a maximum of two doses; and</li> <li>c) Any of the following:</li> </ul>	isation; and		
i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or	or	·	
or vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more prednisone of 2 mg/kg per day or greater	than two weeks, and wh		
20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or fail xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	estation; or ure; or	gh-dose corticost	eroid therapy); or
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ng: ndividuals; or	_	neumovax 23
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe ROTAVIRUS ORAL VACCINE – [Xpharm]		tch-up programm 1 ✓ <u>II</u>	
Maximum of two doses for patients meeting the following:  1) first dose to be administered in infants aged under 1- 2) no vaccination being administered to children aged 2			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

	Subsidy (Manufacturer's Price) \$	Subsid Per	Fully lised •	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]	*			
Either:				
1) Maximum of one dose for primary vaccination for eith	er:			
a) Any infant born on or after 1 April 2016; or				
<li>b) For previously unvaccinated children turning 11 varicella infection (chickenpox), or</li>	years old on or after 1	July 2017, \	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
a) Any of the following for non-immune patients:				
<ul> <li>i) with chronic liver disease who may in futur</li> <li>ii) with deteriorating renal function before trans</li> </ul>		ansplantatior	n; or	
iii) prior to solid organ transplant; or				
<ul><li>iv) prior to any elective immunosuppression*,</li><li>v) for post exposure prophylaxis who are imm</li></ul>		onte : or		
b) For patients at least 2 years after bone marrow			snecial	list or
c) For patients at least 6 months after completion of				
d) For HIV positive non immune to varicella with m				
e) For patients with inborn errors of metabolism at varicella, or				
<li>f) For household contacts of paediatric patients where the household continuous where the household continuous.</li>	act has no clinical hist	ory of varicel	lla, or	
<li>g) For household contacts of adult patients who ha immunocompromised, or undergoing a procedul has no clinical history of varicella.</li>				
* immunosuppression due to steroid or other immunosuppr	essive therapy must b	e for a treatn	nent p	eriod of greater than
28 days				
Inj 1350 PFU prefilled syringe	0.00	1		<u>arivax</u>
		10	_	<u>arivax</u>
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUAT Funded for patients meeting the following criteria:	ED VACCINE [SHING	GLES VACCI	NE] -	-[Xpharm]
<ol> <li>One dose for all people aged 65 years</li> </ol>				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1	<b>J</b> 70	ostavax

- Symbols -	Albendazole	89	Anagrelide hydrochloride	
UK Synacthen80	Albey	227	Analgesics	
3TC105	Albustix	77	Anastrozole	168
- A -	Alchemy Oxybutynin		Anatrole	168
A-Scabies67	Aldurazyme	29	Andriol Testocaps	81
Abacavir sulphate105	Alecensa	157	Androderm	80
Abacavir sulphate with	Alectinib	157	Anoro Ellipta	231
lamivudine 105	Alendronate sodium	111	Antabuse	142
Abiraterone acetate164	Alendronate sodium with		Antacids and Antiflatulents	6
Acarbose11	colecalciferol	111	Anthelmintics	89
Accarb11	Alfacalcidol	33	Antiacne Preparations	61
Accuretic 1048	Alfamino	262	Antiallergy Preparations	227
Accuretic 2048	Alfamino Junior	262	Antianaemics	37
Acetazolamide238	Alginic acid	6	Antiandrogen Oral	
Acetec47	Alglucosidase alfa		Contraceptives	75
Acetic acid with hydroxyquinoline and	Alkeran		Antiarrhythmics	
ricinoleic acid75	Alkeran S29	147	Antibacterials	
Acetylcysteine241	Allersoothe	228	Antibacterials Topical	
Aci-Jel75	Allmercap	149	Anticholinergic Agents	
Aciclovir	Allopurinol		Anticholinesterases	
Infection100	Alpha-Adrenoceptor Blockers		Antidepressants	
Sensory236	Alpha-Keri Lotion		Antidiarrhoeals	
Acidex6	Alphamox		Antiepilepsy Drugs	
Acipimox54	Alphamox 125		Antifibrinolytics, Haemostatics and	
Acitretin68	Alphamox 250		Local Sclerosants	
Aclasta114	Alprolix		Antifibrotics	
Actemra215	Alu-Tab		Antifungals	
Actemra S29215	Aluminium hydroxide		Antifungals Topical	
Actinomycin D151	Alvogen		Antihistamines	
Actrapid10	Amantadine hydrochloride		Antihypotensives	
Actrapid Penfill	Ambrisentan		Antimalarials	
Acupan120	Ambrisentan Mylan		Antimigraine Preparations	
Adalimumab (Amgevita)175	Amgevita		Antinausea and Vertigo Agents	
Adalimumab (Humira)184	Amiloride hydrochloride		Antipruritic Preparations	
Adapalene61	Amiloride hydrochloride with		Antipsychotics	
ADR Cartridge 1.823	furosemide	53	Antiretrovirals	
Adrenaline57	Amiloride hydrochloride with		Antirheumatoid Agents	
Advantan64	hydrochlorothiazide	53	Antispasmodics and Other Agents	
Advate41	Aminophylline		Altering Gut Motility	
Adynovate41	Amiodarone hydrochloride		Antithrombotic Agents	
Afinitor223	Amisulpride		Antithymocyte globulin	
Aflibercept188	Amisulpride Mylan		(equine)	179
Afluria Quad	Amitriptyline		Antitrichomonal Agents	
(2022 formulation)	Amlodipine		Antituberculotics and	
Afluria Quad Junior	Amneal		Antileprotics	90
(2022 formulation)	Amorolfine		Antiulcerants	
AFT-Pyrazinamide100	Amoxicillin		Antivirals	
Agents Affecting the	Amoxicillin with clavulanic acid.		Anxiolytics	
Renin-Angiotensin System 47	Amphotericin B		Anzatax	15/
Agents for Parkinsonism and Related	Amsacrine		Apidra	
Disorders118	AmsaLyo		Apidra SoloStar	
Agents Used in the Treatment of	Amsidine		APO-Atomoxetine	
Poisonings241	Amzoate		Apo-Diltiazem CD	
Agrylin	Anaesthetics		Apo-Doxazosin	
<del>g</del> . , 100			. Po Dovaroom	

Apo-Prednisone	80	Aubagio	136	Betaloc CR	50
Apo-Primidone		Augmentin	92	Betamethasone dipropionate	6
Apo-Temozolomide		Aurorix		Betamethasone dipropionate with	
Apomorphine hydrochloride		AutoSoft 30		calcipotriol	
Aprepitant		AutoSoft 90		Betamethasone sodium phosphate	
Apresoline		Avelox		with betamethasone acetate	
Aptamil AllerPro SYNEO 1		Avonex		Betamethasone valerate	
Aptamil AllerPro SYNEO 2		Avonex Pen		Betamethasone valerate with sodi	
Aptamil Gold+ Pepti Junior		Azacitidine		fusidate [fusidic acid]	
Aqueous cream		Azacitidine Dr Reddy's		Betaxolol	
Aratac		Azamun		Betnovate	
Arava		Azathioprine		Betoptic	
Arginine		Azilect		Betoptic S	
		Azithromycin		Bexsero	
Aripiprazole Sandaz		•		Bezafibrate	
Aripiprazole Sandoz		Azopt AZT		Bezalip	
Aristocort		-B-	105–106		
Arrotex-Prazosin S29		<del>-</del>	4-	Bezalip Retard	
Arrow - Lattim		B-D Micro-Fine		Bicalutamide	
Arrow-Amitriptyline		B-D Ultra Fine		Bicillin LA	
Arrow-Bendrofluazide		B-D Ultra Fine II		BiCNU	
Arrow-Brimonidine		Bacillus Calmette-Guerin (BC	,	Bicnu Heritage	
Arrow-Diazepam		vaccine	175	Bile and Liver Therapy	
Arrow-Doxorubicin	152	Bacillus Calmette-Guerin		Biltricide	
Arrow-Losartan &		vaccine		Bimatoprost	
Hydrochlorothiazide		Baclofen		Bimatoprost Multichem	
Arrow-Norfloxacin	109	Bactroban	62	Binarex	
Arrow-Ornidazole		Balance		Binocrit	
Arrow-Quinapril 10	48	Barrier Creams and Emollient	s65	Biodone	
Arrow-Quinapril 20	48	BCG Vaccine		Biodone Extra Forte	
Arrow-Quinapril 5	48	Beclazone 100		Biodone Forte	122
Arrow-Roxithromycin	91	Beclazone 250	<mark>228</mark>	Bisacodyl	
Arrow-Timolol	238	Beclazone 50	228	Bisoprolol fumarate	50
Arrow-Topiramate	128	Beclomethasone dipropionate	228	Bisoprolol Mylan	
Arrow-Tramadol		Bee venom allergy treatment.	227	BK Lotion	
Arsenic trioxide	150	Bendamustine hydrochloride.	145	Bleomycin sulphate	15 <sup>-</sup>
Asacol	7	Bendrofluazide	54	Blood Colony-stimulating	
Ascorbic acid	33	Bendroflumethiazide		Factors	44
Aspen Adrenaline	57	[Bendrofluazide]	54	Blood glucose diagnostic test	
Aspirin		Benzathine benzylpenicillin		meter	14
Blood	41	Benzatropine mesylate		Blood glucose diagnostic test	
Nervous	120	Benzbromaron AL 100		strip	1
Asthalin	230	Benzbromarone	116	Blood glucose test strips (visually	
Atazanavir sulphate		Benztrop		impaired)	1
Atenolol		Benzydamine hydrochloride		Blood Ketone Diagnostic Test	
Atenolol AFT		Benzylpenicillin sodium [Penic		Strip	13
Atenolol AFT S29		G]		Bonjela	
ATGAM		Beta Cream		Boostrix	
Ativan		Beta Ointment		Bortezomib	
Atomoxetine		Beta Scalp		Bortezomib Dr Reddy's S29	
Atorvastatin		Beta-Adrenoceptor Agonists		Bortezomib Dr-Reddy's	
Atropine sulphate		Beta-Adrenoceptor Blockers		Bortezomib Juno	
Cardiovascular	40	Betadine		Bortezomib Juno S29	
Sensory		Betadine Skin Prep		Bosentan	
		Betaferon		Bosentan Dr Reddy's	
AtroptAtrovent	 ევე	Betahistine dihydrochloride			
				Bplex	
AU Synacthen	80	Betaine	21	Breo Ellipta	22

Brevinor 1/28	74	Canesten	62	Chlorothiazide	5
Bricanyl Turbuhaler	230	Capecitabine	148	Chlorpheniramine maleate	22
Brilinta	41	Capercit	148	Chlorpromazine hydrochloride	13
Brimonidine tartrate	239	Capoten	47	Chlortalidone [Chlorthalidone]	5
Brimonidine tartrate with timolol		Capsaicin		Chlorthalidone	5
maleate	239	Musculoskeletal	111	Chlorvescent	40
Brinzolamide	238	Nervous	120	Choice Load 375	
Brolene	236	Captopril	47	Choice TT380 Short	7
Bromocriptine mesylate	118	Carafate		Choice TT380 Standard	
Brufen SR		Carbaccord	146	Choline salicylate with cetalkoniu	um
BSF Comtan	241	Carbamazepine	126	chloride	
BSF Vigisom		Carbimazole		Ciclosporin	
Buccastem	130	Carbomer	240	Cilazapril	4
Budesonide		Carboplatin	146	Cilicaine	
Alimentary	6	Carboplatin Ebewe		Cilicaine VK	
Respiratory22		Carbosorb-X		Cinacalcet	
Budesonide with eformoterol		Cardinol LA		Cipflox	
Bumetanide		Cardizem CD		Ciprofloxacin	
Buprenorphine Naloxone BNM		CareSens Dual		Infection	9:
Buprenorphine with naloxone		CareSens N		Sensory	
Bupropion hydrochloride	142	CareSens N POP		Ciprofloxacin Teva	
Burinex		CareSens N Premier		Cisplatin	
Burinex S29		CareSens PRO		Cisplatin Ebewe	14
Buscopan		Carmellose sodium with gelating		Citalopram hydrobromide	12
Buspirone hydrochloride		pectin		Cladribine	
Buspirone Viatris		Carmustine		Clarithromycin	
Busulfan		Carvedilol		Alimentary	
- C -		Carvedilol Sandoz		Infection	
Cabergoline	88	Casirivimab and imdevimab		Clexane	4:
Cacit		Catapres		Clexane Forte	
Caffeine citrate		CeeNU		Clindamycin	
Calamine		Cefaclor monohydrate		Clinicians	
Calamine-AFT		Cefalexin		Clinicians Renal Vit	
Calci-Tab 500		Cefalexin Sandoz		Clobazam	
Calcipotriol		Cefazolin		Clobetasol propionate	
Calcitonin		Ceftriaxone		Clobetasone butyrate	
Calcitriol		Ceftriaxone-AFT		Clofazimine	
Calcitriol-AFT		Cefuroxime axetil		Clomazol	
Calcium 500 mg Hexal		Celebrex		Dermatological	6
Calcium carbonate		Celecoxib		Genito-Urinary	7
Calcium Channel Blockers	0, 54	Celecoxib Pfizer		Clomifene citrate	8
Calcium Disodium Versenate		Celestone Chronodose		Clomipramine hydrochloride	
Calcium folinate		Cellcept		Clomipramine Teva	
Calcium Folinate Ebewe		Centrally-Acting Agents		Clonazepam1	
Calcium Folinate Sandoz		Cephalexin ABM		Clonidine	
Calcium Folinate Sandoz S29		Cetirizine hydrochloride		Clonidine BNM	
Calcium gluconate		Cetomacrogol		Clonidine hydrochloride	
Calcium Homeostasis		Cetomacrogol with glycerol		Clonidine Teva	
	70				
Calcium lactate gluconate with calcium carbonate	25	Cetuximab		Clopidogrel Clopidogrel Multichem	4
		Charcoal			
Calcium Possarium				Clopine	
Calcium Resonium Calcium-Sandoz Forte		Chemotherapeutic Agents		Clopixol1	132, 134
		Chickenpox vaccine		Clotrimazole	^-
Calogen  Candesartan cilexetil		Chlorafast		Dermatological	
				Genito-Urinary	
Candestar	4Ծ	Chloramphenicol	23b	Clozapine	13

Clozaril	131	Cyclonex	146	DBL Vincristine Sulfate	15
Co-trimoxazole	96	Cyclopentolate hydrochloride	239	Decozol	3
Coal tar	68	Cyclophosphamide	146	Deferasirox	24
Coal tar with allantoin, menthol,		Cyclorin	99	Deferiprone	24
phenol and sulphur		Cycloserine		Denosumab	
Coal tar with salicylic acid and		Cyproterone acetate	80	Deolate	9
sulphur	68	Cyproterone acetate with		Deoxycoformycin	15
Coco-Scalp	68	ethinyloestradiol	75	Depo-Medrol	8
Codeine phosphate		Cystadane		Depo-Provera	
Extemporaneous	244	Cytarabine		Depo-Testosterone	8
Nervous	121	Cytotec	8	Deprim	9
Coenzyme Q10	28	Cytoxan		Dermol	
Colchicine	116	- D -		Desferrioxamine mesilate	24
Colecalciferol	33	D-Penamine		Desmopressin	8
Colestid	54	Dabigatran	44	Desmopressin acetate	
Colestipol hydrochloride		Dacarbazine		Desmopressin-PH&T	8
Colgout		Dacarbazine APP	151	Desuric	
Colifoam		Dactinomycin [Actinomycin D]	151	Detection of Substances in	
Colistin sulphomethate		Daivobet		Urine	<del>7</del>
Colistin-Link		Daivonex		Dexamethasone	
Collodion flexible		Daktarin	63	Hormone	7
Colloidal bismuth subcitrate		Dalacin C	94	Sensory	
Colofac		Dantrium	117	Dexamethasone phosphate	
Coloxyl		Dantrium S29		Dexamethasone Phosphate	
Combigan		Dantrolene	117	Panpharma	7
Compound electrolytes		Daonil	11	Dexamethasone with framycetin	
Compound electrolytes with glu		Dapa-Tabs	54	gramicidin	
[Dextrose]		Dapsone		Dexamethasone with neomycin	
Compound hydroxybenzoate		Daraprim	95	sulphate and polymyxin B	
Comtan		Darunavir		sulphate	23
Concerta	139	Darunavir Mylan		Dexamfetamine sulfate	
Condoms	72	Dasatinib	158	Dexmethsone	<mark>7</mark>
Condyline	70	Daunorubicin	151	Dextrochlorpheniramine	
Contraceptives - Hormonal		Daunorubicin Zentiva		maleate	22
Contraceptives - Non-hormonal		David One Step Cassette Pregn		Dextrose	
Copaxone		Test		DHC Continus	
Corticosteroids and Related Ag		DBL Acetylcysteine		Diabetes	1
for Systemic Use		DBL Adrenaline		Diabetes Management	
Corticosteroids Topical		DBL Aminophylline		Diacomit	
Cortifoam		DBL Bleomycin Sulfate		Diagnostic Agents	
Cosentyx		DBL Carboplatin		Diamide Relief	
Cosmegen		DBL Cisplatin		Diamox	
Coumadin		DBL Dacarbazine		Diasip	
Country Life		DBL Desferrioxamine Mesylate t	for Inj	Diason RTH	
Coversyl		BP		Diazepam1	26, 13
Creon 10000		DBL Docetaxel		Diazoxide	
Creon 25000	24	DBL Ergometrine	75	Dibenzyline	
Creon Micro		DBL Gemcitabine		Diclofenac Sandoz	11
Crotamiton		DBL Gentamicin		Diclofenac sodium	
Crystaderm		DBL Heparin Sodium		Musculoskeletal	11
Curam		DBL Leucovorin Calcium		Sensory	
Curam Duo 500/125		DBL Methotrexate Onco-Vial		Differin	
Cvite		DBL Morphine Sulphate		Difflam	
Cyclizine hydrochloride		DBL Naloxone Hydrochloride	241	Diflucan	
Cyclizine lactate		DBL Octreotide		Digestives Including Enzymes	
Cyclogyl	239	DBL Pethidine Hydrochloride		Digoxin	4
- ,					

Dihydrocodeine tartrate121	Duocal Super Soluble Powder	2/17	Ensure	25
Dilantin	Duolin		Ensure Plus	
Dilantin Infatab	Duolin HFA		Ensure Plus HN	
Diltiazem hydrochloride	DuoResp Spiromax		Ensure Plus RTH	
Dimethicone65–66	Duride		Ensure Two Cal HN RTH	
Dimethyl fumarate	- E -	57	Entacapone	
,	e-chamber La Grande	224	Entecavir	10
Dipentum				
Diphtheria, tetanus and pertussis	e-chamber Mask		Entecavir Sandoz	
vaccine	e-chamber Turbo		Entocort CIR	
Diphtheria, tetanus, pertussis and	E-Mycin		Entresto 24/26	47
polio vaccine	Ear Preparations		Entresto 49/51	
Diphtheria, tetanus, pertussis, polio,	Ear/Eye Preparations		Entresto 97/103	
hepatitis B and haemophilus	Easiphen Liquid		Epilim	120
influenzae type B vaccine 268	Econazole nitrate		Epilim Crushable	120
Diprosone	Efavirenz	105	Epilim IV	
Diprosone OV63	Efavirenz with emtricitabine and	405	Epilim S/F Liquid	
Dipyridamole41	tenofovir disoproxil		Epilim Syrup	
Disopyramide phosphate49	Eformoterol fumarate		Epirubicin Ebewe	
Disulfiram142	Eformoterol fumarate dihydrate	229	Epirubicin hydrochloride	
Diuretics52	Eftrenonacog alfa [Recombinant		Eplerenone	
Docetaxel151	factor IX]		Epoetin alfa	3
Docetaxel Accord151	Efudix		Epoprostenol	6
Docetaxel Sandoz151	Egopsoryl TA		Eptacog alfa [Recombinant factor	
Docusate sodium25	Elaprase		VIIa]	
Docusate sodium with	Eldepryl	118	ERA	
sennosides25	Elecare		Erbitux	
Dolutegravir106	Elecare LCP	262	Ergometrine maleate	
Domperidone 130	Electral		Erlotinib	
Donepezil hydrochloride140	Elelyso	31	Erythrocin IV	
Donepezil-Rex140	Elemental 028 Extra	252	Erythromycin (as lactobionate)	
Dornase alfa233	Elidel	68	Erythromycin ethyl succinate	9
Dortimopt238	Elocon		Erythromycin stearate	9
Dorzolamide hydrochloride238	Elocon Alcohol Free	64	Esbriet	
Dorzolamide with timolol238	Eltrombopag	38	Escitalopram	12
Dostinex88	Eltroxin	83	Escitalopram (Ethics)	12
Dosulepin [Dothiepin]	EMB Fatol	99	Eskazole	
hydrochloride124	Emend Tri-Pack	129	Essential Generics	50
Dosulepin Mylan124	Emicizumab	39	Essential Prednisolone	
Dothiepin124	EMLA	120	Estradot	
Doxazosin47	Empagliflozin	13	Estradot 50 mcg	8
Doxazosin Clinect47	Empagliflozin with metformin		Estrofem	8
Doxine93	hydrochloride	. 13	Etanercept	
Doxorubicin Ebewe152	Emtricitabine		Ethambutol hydrochloride	9
Doxorubicin hydrochloride 152	Emtricitabine with tenofovir		Ethics Aspirin	
Doxycycline93	disoproxil	102	Ethics Aspirin EC	4
DP Lotion65	Emtriva		Ethics Lisinopril	4
DP Lotn HC64	Emulsifying ointment	65	Ethinyloestradiol	
DP-Allopurinol116	Emulsifying Ointment ADE		Ethinyloestradiol with	
Dr Reddy's Omeprazole9	Enalapril maleate		desogestrel	
Drofate51	Enbrel	168	Ethinyloestradiol with	
Drugs Affecting Bone	Endocrine Therapy		levonorgestrel	74
Metabolism 111	Endoxan		Ethinyloestradiol with	
Dual blood glucose and blood ketone	Engerix-B		norethisterone	74
diagnostic test meter 14	Enlafax XR		Ethosuximide	12
Dulaglutide12	Enoxaparin sodium		Etopophos	
Dulcolax SP Drop26	Enstilar		Etoposide	

Etoposide phosphate	152	Flecainide Controlled Release		Frusemide	5
Etravirine	105	Teva	49	Fucicort	6
Eumovate	64	Fleet Phosphate Enema	26	Fucidin	9
Everet	127	Flixonase Hayfever & Allergy	234	Fucithalmic	23
Everolimus		Flixotide		Fulvestrant	
Evista	112	Flixotide Accuhaler		Fungilin	
Exemestane		Florinef		Furosemid-Ratiopharm	
Exjade		Fluanxol		Furosemide [Frusemide]	
Extemporaneously Compounded		Flucil		Furosemide-Baxter	
Preparations and	•	Flucloxacillin		fusidic acid	
Galenicals	244	Flucloxacillin-AFT		Dermatological	62 6
Eye Preparations		Flucloxin		Infection	
Eylea		Flucon		Sensory	
Ezetimibe		Fluconazole		- G -	20
		Fludara Oral			10
Ezetimibe Sandoz				GabapentinGacet	12
Ezetimibe with simvastatin	50	Fludarabine Ebewe			
-F-		Fludarabine phosphate		Galsulfase	
Factor eight inhibitor bypassing		Fludrocortisone acetate		Galvumet	
fraction		Fluids and Electrolytes		Galvus	
Famotidine		Flumetasone pivalate	236	Gardasil 9	
Famotidine Hovid	•	Fluocortolone caproate with		Gastrodenol	
Faslodex		fluocortolone pivalate and		Gaviscon Double Strength	
Fatty Cream AFT	65	cinchocaine	8	Gaviscon Infant	
Febuxostat	116	Fluorometholone	237	Gazyva	19
Febuxostat multichem	116	Fluorouracil	149	Gefitinib	15
Feed Thickener Karicare		Fluorouracil Accord	149	GEM Aqueous Cream	6
Aptamil	259	Fluorouracil sodium	70	Gemcitabine Ebewe	14
FEIBA NF		Fluox	125	Gemcitabine hydrochloride	
Felo 10 ER	51	Fluoxetine hydrochloride	125	Genoptic	
Felo 5 ER	51	Flupenthixol decanoate		Gentamicin sulphate	
Felodipine		Flutamide		Infection	9
Femme-Tab ED		Flutamin		Sensory	
Fenpaed 100 mg per 5 ml		Fluticasone		Gilenya	
Fentanyl		Fluticasone furoate with		Ginet	
Fentanyl Sandoz		vilanterol	220	Glatiramer acetate	
Ferinject		Fluticasone propionate		Glecaprevir with pibrentasvir	
Ferodan		Fluticasone with salmeterol		Glibenclamide	
Ferriprox		FML		Gliclazide	
Ferro-F-Tabs		Foban		Glipizide	
Ferro-tab		Folic acid		Glivec	
Ferrograd		Folic Acid Multichem		Glizide	
Ferrosig		Folic Acid Mylan		Glucagen Hypokit	
Ferrous fumarate		Food Thickeners		Glucagon hydrochloride	
Ferrous fumarate with folic acid.		Foods And Supplements For In		Glucerna Select	
Ferrous sulfate		Errors Of Metabolism		Glucose [Dextrose]	4
Fexofenadine hydrochloride		Foradil		Gluten Free Foods	
Fibro-vein		Forteo	113	Glycerin with sodium saccharin	
Filgrastim	44	Fortini		Glycerin with sucrose	24
Finasteride	76	Fortini Multi Fibre	251	Glycerol	
Fingolimod	135	Fortisip		Alimentary	2
Firazyr	227	Fortisip Multi Fibre	257	Extemporaneous	
Flagyl		Fosamax		Glyceryl trinitrate	
Flagyl-S		Fosamax Plus	111	Alimentary	
Flamazine		Framycetin sulphate		Cardiovascular	5
Flecainide acetate		Frisium		Glycopyrronium	
Flecainide BNM		Frumil		Glycopyrronium bromide	
				,p, bronniae	

Glycopyrronium with		Humulin NPH	10	Indacaterol	229
indacaterol	231	Humulin R		Indapamide	
Go Healthy		Hyaluronic acid		Infanrix IPV	
Gold Knight		Hydralazine		Infanrix-hexa	
Gold Knight XL		Hydralazine hydrochloride		Infant Formulae	
Goserelin		Hydrocortisone	57	Infatrini	
			64		
Gutron		Dermatological		Infliximab	
Gynaecological Anti-infectives	/5	Hormone		Influenza vaccine	
-H-	4.40	Hydrocortisone (PSM)		Inhaled Corticosteroids	228
Habitrol	143	Hydrocortisone acetate	/	Inhaled Long-acting	000
Haemophilus influenzae type B		Hydrocortisone acetate with	_	Beta-adrenoceptor Agonists	
vaccine		pramoxine hydrochloride	7	Inspra	
Haldol		Hydrocortisone and paraffin liquid		Instillagel Lido	
Haldol Concentrate	132	and lanolin		Insulin aspart	11
Haldol Decanoas	132	Hydrocortisone butyrate		Insulin aspart with insulin aspart	
Haloperidol	131	Hydrocortisone with cinchocaine	8	protamine	
Haloperidol decanoate		Hydrocortisone with miconazole	64	Insulin glargine	<b>1</b> 1
Harvoni	102	Hydrocortisone with natamycin and	l	Insulin glulisine	<b>1</b> 1
Havrix	268	neomycin	64	Insulin isophane	
Havrix Junior	268	Hydrogen peroxide	61	Insulin isophane with insulin	
Haylor syrup	228	Hydroxocobalamin		neutral	10
healthE Dimethicone 10%		Hydroxocobalamin Mercury		Insulin lispro	
healthE Dimethicone 4% Lotion		Pharma	33	Insulin lispro with insulin lispro	
healthE Dimethicone 5%		hydroxycarbamide		protamine	11
healthE Glycerol BP		Hydroxychloroquine		Insulin neutral	
healthE Urea Cream		Hydroxyurea		Insulin pen needles	
Healtheries Simple Baking Mix		[hydroxycarbamide]	152	Insulin pump	
Hemastix		Hygroton		Insulin pump cartridge	
Hemlibra		Hylo-Fresh		Insulin pump infusion set (steel	
Heparin sodium		Hymenoptera		cannula)	21
Heparinised saline		Hyoscine butylbromide		Insulin pump infusion set (steel	
Heparon Junior		Hyoscine hydrobromide		cannula, straight insertion)	21
				Insulin pump infusion set (teflon	2
Hepatitis A vaccine	200	Hypam			00
Hepatitis B recombinant	000	Hyperuricaemia and Antigout		cannula)	20
vaccine		Hypromellose		Insulin pump infusion set (teflon	
Herceptin		Hypromellose with dextran	239	cannula, angle insertion with	00
Hiberix		-1-		insertion device)	23
Hiprex		Ibiamox		Insulin pump infusion set (teflon	
Histaclear		Ibrance		cannula, angle insertion)	23
Histafen		Ibuprofen		Insulin pump infusion set (teflon	
Holoxan		Icatibant		cannula, straight insertion with	
Horleys Bread Mix		Idarubicin hydrochloride		insertion device)	23
Horleys Flour	259	Idursulfase		Insulin pump infusion set (teflon	
Hormone Replacement Therapy -		Ifosfamide	146	cannula, straight insertion)	23
Systemic	81	Igroton	54	Insulin pump reservoir	23
HPV	269	lkorel		Insulin syringes, disposable with	
Humalog		lloprost	60	attached needle	16
Humalog Mix 25	11	Imatinib mesilate	159	Intelence	105
Humalog Mix 50		Imatinib-Rex		Interferon beta-1-alpha	
Human papillomavirus (6, 11, 16, 1		Imigran		Interferon beta-1-beta	135
31, 33, 45, 52 and 58) vaccine		Imipramine hydrochloride		Intra-uterine device	73
[HPV]	269	Imiquimod		Invega Sustenna	
Humatin		Immune Modulators		IPCA-Frusemide	53
Humira		Immunosuppressants		IPCA-Metoprolol	
HumiraPen		Imuran		IPCA-Propranolol	
Humulin 30/70		Incruse Ellipta		IPOL	
		300 Empia		🕶	/ <

Ipratropium bromide	.230, 234	Klacid		Levomepromazine	13
Iressa		Alimentary	9	Levomepromazine	
Irinotecan Actavis 100	149	Infection	90	hydrochloride	13
Irinotecan hydrochloride	149	Kliogest	82	Levonorgestrel	
Irinotecan-Rex		Kliovance	82	Genito-Urinary	74–7
Iron (as ferric carboxymaltose)	35	Kogenate FS	41	Hormone	8
Iron polymaltose		Konakion MM	41	Levothyroxine	8
Isentress		Konsyl-D		Lidocaine [Lignocaine]	
Isentress HD	106	Kuvan	30	Lidocaine [Lignocaine]	
Ismo 20	<mark>57</mark>	-L-		hydrochloride	12
Ismo 40 Retard	<mark>57</mark>	Labetalol	50	Lidocaine [Lignocaine] with	
Isoniazid	99	Lacosamide	127	chlorhexidine	12
Isoniazid with rifampicin	99	Lactulose	26	Lidocaine [Lignocaine] with	
Isoptin		Laevolac		prilocaine	12
Isoptin Retard		Lagevrio	103	Lidocaine-Baxter	
Isoptin SR		Lamictal		Lidocaine-Claris	
Isopto Carpine		Lamivudine		Life Extension	3
Isosorbide mononitrate		Lamivudine Alphapharm	,	Lignocaine	
Isosource Standard		Lamotrigine		Lioresal Intrathecal	
Isotretinoin		Lamprene		Lipid-Modifying Agents	
Ispaghula (psyllium) husk		Lanoxin		Liquigen	
Itch-Soothe		Lanoxin Paediatric Elixir		Lisinopril	
Itraconazole		Lanoxin PG		Litak	
Itrazole		Lanoxin S29		Lithium carbonate	
lvacaftor		Lansoprazole		Livostin	
lvermectin		Lantus		LMX4	
- <b>J</b> -	00	Lantus SoloStar		Locacorten-Viaform ED's	
Jadelle	7/	Lanvis		Local preparations for Anal an	
Jakavi		Lanzol Relief		Rectal Disorders	
Jardiamet		Lapatinib ditosylate		Locasol	
Jardiance		Largactil		Locoid	
		Largacii		Locoid Crelo	
Jaydess					
Jevity HiCal RTH		Lasix		Locoid Lipocream	b
Jevity Plus		Latanoprost		Locorten-Vioform	
Jevity RTH		Latanoprost with timolol		Lodoxamide	
Juno Pemetrexed	150	Lax-Suppositories		Logem	
- K -	000	Lax-Tab		Lomide	
Kadcyla		Laxatives		Lomustine	
Kaletra		Laxsol		Loniten	
Kalydeco		Ledipasvir with sofosbuvir		Loperamide hydrochloride	
Kemadrin		Leflunomide		Lopinavir with ritonavir	10
Kenacomb		Lenalidomide		Lopinavir/Ritonavir Mylan	
Kenacort-A 10		Letrole		Loprofin	
Kenacort-A 40		Letrozole		Loprofin Mix	
Kenalog in Orabase	33	Leukeran FC	146	Lorafix	
Ketocal 3:1	266	Leukotriene Receptor		Loratadine	
KetoCal 4:1	266	Antagonists		Lorazepam	
Ketoconazole		Leuprorelin	87	Lorstat	
Dermatological	69	Leustatin		Losartan Actavis	4
Infection	97	Levetiracetam	127	Losartan potassium	4
Ketogenic Diet		Levetiracetam-AFT	127	Losartan potassium with	
Ketoprofen		Levlen ED	74	hydrochlorothiazide	4
KetoSens		Levocabastine	238	Lovir	
Keytruda		Levocarnitine		Loxamine	
Kindergen		Levodopa with benserazide		Lucrin Depot 1-month	
Kivexa	105	Levodopa with carbidopa		Lucrin Depot 3-month	
				- L	

67	Metabolic Disorder Agents27	Midazolam	130
Lynparza153	Metformin hydrochloride11	Midazolam-Baxter	136
Lyrica127	Metformin Mylan11	Midodrine	49
- M -	Methadone hydrochloride	Mifegyne	7
m-Eslon122	Extemporaneous244	Mifepristone	7
Mabthera200	Nervous122	Minerals	3
Macro Organic Psyllium Husk25	Methatabs122	Mini-Wright AFS Low Range	23
Macrobid109	Methenamine (hexamine)	Mini-Wright Standard	23
Macrogol 3350 with potassium	hippurate109	Minidiab	
chloride, sodium bicarbonate and	Methopt239	MiniMed 1.8 Reservoir	
sodium chloride26	Methotrexate149	MMT-326A	2
Macrogol 400 and propylene	Methotrexate DBL Onco-Vial149	MiniMed 3.0 Reservoir	
glycol240	Methotrexate Ebewe149	MMT-332A	2
Madopar 125118	Methotrexate Sandoz149	MiniMed 770G	10
Madopar 250118	Methyl hydroxybenzoate244	MiniMed Mio MMT-921A	2
Madopar 62.5118	Methylcellulose244	MiniMed Mio MMT-923A	2
Madopar HBS118	Methylcellulose with glycerin and	MiniMed Mio MMT-925A	2
Madopar Rapid118	sodium saccharin244	MiniMed Mio MMT-941A	2
Magnesium hydroxide36	Methylcellulose with glycerin and	MiniMed Mio MMT-943A	2
Magnesium sulphate36	sucrose244	MiniMed Mio MMT-945A	2
Mantoux276	Methyldopa52	MiniMed Mio MMT-965A	2
Marevan44	Methyldopa Mylan52	MiniMed Mio MMT-975A	
Marine Blue Lotion SPF 50+70	Methyldopa Mylan S2952	MiniMed Quick-Set MMT-386A	2
Martindale Pharma241	Methylnaltrexone bromide25	MiniMed Quick-Set MMT-387A	
Mask for spacer device234	Methylphenidate ER - Teva138	MiniMed Quick-Set MMT-396A	2
Maviret102	Methylphenidate hydrochloride 138	MiniMed Quick-Set MMT-397A	2
Maxidex237	Methylphenidate hydrochloride	MiniMed Quick-Set MMT-398A	2
Maxitrol237	extended-release139	MiniMed Quick-Set MMT-399A	
MCT oil (Nutricia)248	Methylprednisolone79	MiniMed Silhouette MMT-368A	
Measles, mumps and rubella	Methylprednisolone (as sodium	MiniMed Silhouette MMT-377A	2
vaccine272	succinate) 80	MiniMed Silhouette MMT-378A	2
Mebendazole89	Methylprednisolone aceponate64	MiniMed Silhouette MMT-381A	2
Mebeverine hydrochloride8	Methylprednisolone acetate80	MiniMed Silhouette MMT-382A	2
Medrol79	Methylxanthines233	MiniMed Silhouette MMT-383A	2
Medroxyprogesterone acetate	Metoclopramide Actavis 10130	MiniMed Silhouette MMT-384A	
Genito-Urinary74	Metoclopramide hydrochloride 130	MiniMed Sure-T MMT-864A	2
Hormone81–82	Metolazone53	MiniMed Sure-T MMT-866A	2
Mefenamic acid110	Metopirone88	MiniMed Sure-T MMT-874A	2
Megace166	Metoprolol IV Mylan50	MiniMed Sure-T MMT-876A	2
Megestrol acetate166	Metoprolol succinate50	MiniMed Sure-T MMT-884A	2
Melatonin136	Metoprolol tartrate50	MiniMed Sure-T MMT-886A	2
Melphalan147	Metrogyl98	Minims Pilocarpine	23
Menactra272	Metronidazole98	Minims Prednisolone	23
Meningococcal (groups A, C, Y and	Metyrapone88	Minirin	8
W-135) conjugate vaccine 272	Mexiletine hydrochloride49	Minirin Melt	8
Meningococcal B multicomponent	Miacalcic78	Mino-tabs	9
vaccine273	Micolette26	Minocycline hydrochloride	
Meningococcal C conjugate	Miconazole33	Minomycin	
vaccine273	Miconazole nitrate	Minor Skin Infections	6
Menthol63	Dermatological63	Minoxidil	5
Mepolizumab197	Genito-Urinary75	Mirena	8
Mercaptopurine149	Micreme75	Mirtazapine	
Mercilon 2873	Micreme H64	Misoprostol	
Mesalazine7	Microgynon 20 ED74	Mitomycin C	15
Mesna	Microgynon 3074	Mitozantrone	15
Mestinon110	Microlut74	Mitozantrone Ebewe	

Mixtard 30	10	Naphcon Forte	240	Norethisterone	
MMR II	272	Naprosyn SR 1000		Genito-Urinary	7
Moclobemide	125	Naprosyn SR 750		Hormone	8
Modafinil	140	Naproxen		Norflex	
Modavigil	140	Narcaricin mite		Norfloxacin	10
Moduretic		Nasal Preparations	234	Noriday 28	
Molaxole		Natalizumab	136	Norimin	
Molnupiravir		Natulan		Normacol Plus	
Moments		Nausafix		Normison	13
Mometasone furoate		Nausicalm		Norpress	
Monogen		Navelbine		Nortriptyline hydrochloride	
Montelukast		Nefopam hydrochloride		Norvir	
Montelukast Mylan		Neisvac-C		Noumed	
Moroctocog alfa [Recombinant f		Neo-B12	33	Noumed Paracetamol	12
VIII]		Neo-Mercazole	83	NovaSource Renal	
Morphine hydrochloride		Neocate Gold		Novatretin	
Morphine sulphate		Neocate Junior Unflavoured	262	NovoMix 30 FlexPen	1
Motetis		Neocate Junior Vanilla		NovoRapid	
Mouth and Throat		Neocate SYNEO		NovoRapid FlexPen	
Movapo		Neoral		NovoRapid Penfill	
Moxifloxacin		Neostigmine metilsulfate		NovoSeven RT	
MSUD Maxamum		Nepro HP (strawberry)		Noxafil	
Mucilaginous laxatives with		Nepro HP (vanilla)		Nozinan	
stimulants	25	Nepro HP RTH		Nozinan (Swiss)	
Mucolytics		Neulactil		Nucala	
Mucosoothe		Neulastim		Nuelin	
Multiple Sclerosis Treatments		NeuroTabs		Nuelin-SR	
Multivitamin renal		Nevirapine		Nupentin	
Multivitamins		Nevirapine Alphapharm		Nutilis	25
Mupirocin		Nicorandil		Nutren Diabetes	
Muscle Relaxants		Nicotine		Nutrient Modules	
Mvite		Nifedipine		Nutrini Energy Multi Fibre	
Myambutol		Nifuran		Nutrini Energy RTH	
Mycobutin		Nilotinib		Nutrini Low Energy Multi Fibre	
MycoNail		Nilstat		Nutrini Peptisorb	
Mycophenolate mofetil		Alimentary	33	Nutrini Peptisorb Energy	
Mydriacyl		Genito-Urinary		Nutrini RTH	
Mylan (12 hr release)		Infection		Nutrison 800 Complete Multi	
Mylan (24 hr release)		Nintedanib		Fibre	25
Mylan Atenolol		Nipent		Nutrison Advanced Diason	
Mylan Clomiphen		Nirmatrelvir with ritonavir		Nutrison Concentrated	
Mylan Indapamide		Nitrates		Nutrison Energy	
Mylan Italy (24 hr release)		Nitroderm TTS		Nutrison Energy Multi Fibre	
Myleran		Nitrofurantoin		Nutrison Multi Fibre	
Myometrial and Vaginal Hormor		Nitrolingual Pump Spray		Nutrison Standard RTH	
Preparations		Nivestim		Nyefax Retard	
Myozyme		Nivolumab		Nystatin	
- N -		Nizoral		Alimentary	3
Nadolol	50	Nodia		Genito-Urinary	
Nadolol BNM		Noflam 250		Infection	
Naglazyme		Noflam 500		NZB Low Gluten Bread Mix	
Nalcrom	20 Q	Non-Steroidal Anti-Inflammatory		- 0 -	23
Naloxone hydrochloride	2/1			O/W Fatty Emulsion Cream	G
Naltraccord		Drugs Nonacog gamma, [Recombinan		Obinutuzumab	
Naltrexone hydrochloride		Factor IX]		Obstetric Preparations	
Naphazoline hydrochloride		1 40101 1/1]		Obstatio i reparations	/
TRADITION OF THE TRAINING THE T					

Ocrelizumab	136	Osmolite RTH	256	Paroxetine	12
Ocrevus		Other Endocrine Agents		Paser	
Octocog alfa [Recombinant f		Other Oestrogen Preparations		Paxam	
VIII] (Advate)		Other Progestogen		Paxlovid	
Octocog alfa [Recombinant f		Preparations	82	Pazopanib	
VIII] (Kogenate FS)		Other Skin Preparations		Peak flow meter	
Octreotide		Ovestin		Pedialyte - Bubblegum	
Octreotide (Sun)		Genito-Urinary	75	Pediasure	
Octreotide Depot Teva		Hormone		Pediasure RTH	
Octreotide GH		Oxaliplatin		Pegaspargase	
Octreotide long-acting		Oxaliplatin Accord		Pegasys	
				Pegfilgrastim	
Octreotide MaxRx		Oxaliplatin Actavis 100 Oxaliplatin Ebewe			
Oestradiol		•		Pegylated interferon alfa-2a	10
Oestradiol valerate		Oxis Turbuhaler		Pembrolizumab	
Oestradiol with norethisteror	ie82	Oxpentifylline		Pemetrexed	
Oestriol	75	Oxybutynin		Penicillamine	
Genito-Urinary		Oxycodone hydrochloride		Penicillin G	
Hormone		Oxycodone Sandoz		PenMix 30	
Oestrogens		Oxycodone Sandoz S29		PenMix 40	
Ofev		OxyContin		PenMix 50	
Oil in water emulsion		OxyNorm	123	Pentasa	
Olanzapine		Oxytocin		Pentostatin [Deoxycoformycin]	
Olaparib		Oxytocin BNM	75	Pentoxifylline [Oxpentifylline]	
Olbetam		Oxytocin with ergometrine		Peptamen Junior	
Olbetam S29	54	maleate	75	Peptisorb	25
Olopatadine		Ozurdex	237	Perhexiline maleate	
Olopatadine Teva	240	- P -		Pericyazine	13
Olsalazine		Pacifen	117	Perindopril	4
Omalizumab	198	Pacimol	121	Perjeta	20
Omeprazole	9	Paclitaxel	154	Permethrin	6
Omeprazole actavis 10	9	Paclitaxel Actavis	154	Perrigo	70
Omeprazole actavis 20	9	Paclitaxel Ebewe	154	Pertuzumab	
Omeprazole actavis 40		Paediatric Seravit	34	Peteha	9
Omnitrope		Palbociclib	161	Pethidine hydrochloride	124
Onbrez Breezhaler	229	Paliperidone	133	Pevaryl	
Oncaspar LYO		Pamidronate disodium		Pexsig	5
OncoTiCE		Pamisol		Pfizer Exemestane	
Ondansetron		Pancreatic enzyme		Pharmacy Health Sorbolene with	
Ondansetron ODT-DRLA		Pantoprazole		Glycerin	6
One-Alpha		Panzop Relief		Pharmacy Services	
Onrex		Panzytrat		Pheburane	
Opdivo		Papaverine hydrochloride		Phenasen	
Ora-Blend		Para-amino salicylic acid		Phenobarbitone	
Ora-Blend SF		Paracare		Phenobarbitone sodium	
Ora-Plus		Paracare Double Strength		Extemporaneous	24
Ora-Sweet		Paracetamol		Nervous	
Ora-Sweet SF		Paracetamol + Codeine		Phenoxybenzamine	
Orabase		(Relieve)	124	hydrochloride	4
Oral and Enteral Feeds		Paracetamol with codeine		Phenoxymethylpenicillin (Penicillin	
Oratane		Paraffin		V)	) O
Ordine		Paraffin liquid with wool fat		Phenytoin sodium12	ენ_ 1ე: უ
		·			
Orgran		Parasiticidal Preparations		Phillips Milk of Magnesia	
Ornidazole		Parlodel		Phlexy 10	
Orphenadrine citrate				Phospharus	
Ortho-tolidine		Parnate S29		Phosphorus	
Oruvail SR	I IU	Paromomycin	<del>95</del>	Phytomenadione	4

Pilocarpine hydrochloride	239	Prednisone	80	Quick-Set MMT-392	23
Pimafucort	64	Prednisone Clinect	80	Quick-Set MMT-393	23
Pimecrolimus	68	Pregabalin	127	Quinapril	48
Pine tar with trolamine laurilsuli	fate	Pregabalin Pfizer	127	Quinapril with	
and fluorescein	69	Pregnancy Tests - hCG Urine		hydrochlorothiazide	48
Pinetarsol	69	Premarin		Qvar	
Pioglitazone	11	Prevenar 13		- R -	
Pirfenidone		Priadel		RA-Morph	122
Pizotifen		Primaquine		Raloxifene hydrochloride	
PKU Anamix Infant		Primidone		Raltegravir potassium	
PKU Anamix Junior		Primidone Clinect		Ramipex	
PKU Anamix Junior Chocolate		Primolut N		Ranbaxy-Cefaclor	
PKU Anamix Junior LQ		Priorix		Ranbaxy-Cefaclor S29	
PKU Anamix Junior Orange		Probenecid		Rapamune	
PKU Anamix Junior Vanilla		Probenecid-AFT		Rasagiline	118
PKU Lophlex LQ 10		Procaine penicillin		Reandron 1000	
		Procarbazine hydrochloride		Recombinant factor IX	
PKU Lophlex LQ 20		-			
PKU Lophlex Powder		Prochlorperazine		Recombinant factor VIIa	
PKU Lophlex Sensation 20		Proctofoam		Recombinant factor VIII	
Plaquenil		Proctosedyl		Rectogesic	
Plendil ER		Procyclidine hydrochloride		Redipred	
Pneumococcal (PCV10) conjug		Progesterone		Relieve	
vaccine		Proglicem		Relistor	
Pneumococcal (PCV13) conjug		Proglycem		Remicade	
vaccine	2/4	Progynova		Renilon 7.5	
Pneumococcal (PPV23)		Prolia		Resonium-A	
polysaccharide vaccine		Promethazine hydrochloride		Resource Beneprotein	
Pneumovax 23		Propafenone hydrochloride		Respigen	
Podophyllotoxin		Propamidine isethionate		Respiratory Devices	
Polaramine	228	Propranolol		Respiratory Stimulants	235
Poliomyelitis vaccine	275	Propylene glycol		Retinol palmitate	
Poloxamer	25	Propylthiouracil	83	ReTrieve	61
Poly-Gel	240	Prostacur	165	Retrovir	105
Poly-Tears		Protaphane	10	Revlimid	152
Poly-Visc	240	Protaphane Penfill	10	Revolade	38
Polycal	246	Protifar	248	Rexacrom	238
Ponstan	110	Protionamide	99	Riboflavin	29
Posaconazole	97	Provera	81	Ribomustin	145
Postinor-1	75	Provera HD	82	Ricit	76
Potassium chloride	45-46	PSM Citalopram	125	Rifabutin	100
Potassium citrate	76	Psoriasis and Eczema		Rifadin	100
Potassium iodate	35	Preparations	68	Rifampicin	100
Povidone iodine	66	PTU		Rifaximin	9
Pradaxa	44	Pulmicort Turbuhaler	228	Rifinah	99
Pramipexole hydrochloride		Pulmozyme	233	Rilutek	119
Pravastatin		Puri-nethol		Riluzole	119
Pravastatin Mylan		Puria		RINVOQ	225
Praziquantel		Pyrazinamide	100	Riodine	66
Prazosin		Pyridostigmine bromide		Risedronate Sandoz	
Pred Forte		Pyridoxine hydrochloride		Risedronate sodium	
Prednisolone		Pyridoxine multichem		Risperdal Consta	
Prednisolone acetate		Pyrimethamine		Risperidone	
Prednisolone sodium		Pytazen SR		Risperidone (Teva)	
Prednisolone sodium		- Q -		Risperon	
phosphate	238	Quetapel	132	Ritalin	
Prednisolone-AFT		Quetiapine		Ritalin LA	
i iodinolione Al I	200	addiapino	102	1 III. L/1	103

Ritonavir	106	Senokot	26	Sodium tetradecyl sulphate	4
Rituximab (Mabthera)	200	SensoCard	15	Sodium valproate	12
Rituximab (Riximyo)	202	Serc	130	Sofradex	23
Rivaroxaban	44	Serenace	131	Soframycin	23
Rivastigmine	141	Seretide		Solgar27-	-29, 3°
Rivastigmine Patch BNM 10	141	Seretide Accuhaler	230	Solifenacin Mylan	
Rivastigmine Patch BNM 5	141	Serevent	229	Solifenacin succinate	7
Rivotril	126	Serevent Accuhaler	229	Solu-Cortef	
Riximyo		Sertraline		Solu-Medrol	8
RIXUBIS	40	Setrona	125	Solu-Medrol-Act-O-Vial	
Rizamelt	129	Setrona AU		Somatropin (Omnitrope)	8
Rizatriptan	129	Sevredol	122	Sotalol	5
RoActemra S29	215	Sex Hormones Non		Spacer device	23
Ronapreve	189	Contraceptive	80	Span-K	4
Ropin	118	Shield XL	72	Spiolto Respimat	23
Ropinirole hydrochloride	118	shingles vaccine	276	Spiractin	
Rosuvastatin	55	SII-Onco-BCG	175	Spiriva	23
Rosuvastatin Viatris	55	Sildenafil		Spiriva Respimat	
Rotarix		Silhouette MMT-373		Spironolactone	
Rotavirus oral vaccine		Siltuximab		Sporanox	
Roxane	6	Simvastatin	55	Sprycel	
Roxane-Propranolol		Simvastatin Mylan		Stemetil	
Roxithromycin		Sinemet		SteroClear	
Rubifen		Sinemet CR		Stesolid	
Rubifen SR		Sirolimus	223	Stimulants/ADHD Treatments	
Rugby Capsaicin Topical		Siterone	80	Stiripentol	
Cream	120	Slow-Lopresor		Stocrin	
Rulide D		Smith BioMed Rapid Pregnancy		Stomahesive	
Rurioctocog alfa pegol [Recon	nbinant	Test	76	Strattera	
factor VIII]		Sodibic		Strides Shasun	
Ruxolitinib		Sodium acid phosphate		Stromectol	
Rythmodan		Sodium alginate		Sucralfate	
Rytmonorm		Sodium benzoate		Sulfadiazine Silver	
-S-		Sodium bicarbonate		Sulfadiazine sodium	9
Sabril	128	Blood	45-46	Sulfasalazine	
Sacubitril with valsartan		Extemporaneous		Sulphur	
Sagent		Sodium calcium edetate		Sulprix	
SalAir		Sodium chloride		Sumagran	
Salazopyrin		Blood	45	Sumatriptan	
Salazopyrin EN		Respiratory		Sunitinib	
Salbutamol		Sodium citrate with sodium lauryl	201	Sunitinib Pfizer	
Salbutamol with ipratropium		sulphoacetate	26	Sunscreens	
bromide	230	Sodium citro-tartrate		Sunscreens, proprietary	
Salicylic acid		Sodium cromoglicate		Sure-T MMT-863	
Salmeterol		Alimentary	8	Sure-T MMT-873	
Sandomigran		Sensory		Sustagen Hospital Formula	
Sanofi Primaquine		Sodium fluoride		Sustagen Hospital Formula	20
Sapropterin dihydrochloride		Sodium Fusidate [fusidic acid]		Active	250
Scalp Preparations		Dermatological	62	Sustanon Ampoules	
Scopoderm TTS		Infection		Sutent	
Sebizole		Sensory		Sylvant	
Secukinumab		Sodium hyaluronate [Hyaluronic	200	Symbicort Turbuhaler 100/6	
Sedatives and Hypnotics		acid]	240	Symbicort Turbuhaler 200/6	22
Seebri Breezhaler		Sodium phenylbutyrate	30 30	Symbicort Turbuhaler 400/12	
Selegiline hydrochloride	118	Sodium picosulfate	26	Symmetrel	111
Senna		Sodium polystyrene sulphonate		Sympathomimetics	
OUI II a		oodium polystyrene sulphoriale	40	Oympamommono	

## **INDEX: Generic Chemicals and Brands**

Synacthen	80	Thioguanine	150	Trimethoprim	9
Synacthen Depot		Thiotepa		Trimethoprim with	
Synacthene Retard		Thyroid and Antithyroid Agents		sulphamethoxazole	
Synflorix		Ticagrelor		[Co-trimoxazole]	9
Synthroid		Tilcotil		Trisequens	
Syntometrine		Timolol		Trisul	
Syrup (pharmaceutical grade)		Timoptol XE		Trophic Hormones	
Systane Unit Dose		Tiotropium bromide		Tropicamide	
- T -	2 10	Tiotropium bromide with		Trulicity	1
Tacrolimus		olodaterol	231	Trusopt	23
Dermatological	69	Tivicay		TruSteel	
Oncology		TMP		Tuberculin PPD [Mantoux] test	
Tacrolimus Sandoz		Tobramycin		Tubersol	
Taliglucerase alfa		Infection	05	Two Cal HN	
Tambocor	40	Sensory		Tykerb	
Tamoxifen citrate				Tysabri	
		Tobramycin BNM		- U -	10
Tamoxifen Sandoz		Tobramycin Mylan		-	00
Tamsulosin hydrochloride		Tobrex		Ultibro Breezhaler	
Tamsulosin-Rex		Tocilizumab		Ultraproct	
Tandem Cartridge		Tofranil		Umeclidinium	
Tandem t:slim X2 with Basal-IQ		Tolcapone		Umeclidinium with vilanterol	
Tap water		Topamax	128	Univent2	
Tarceva		Topical Products for Joint and		Upadacitinib	
Tasigna		Muscular Pain		Ural	
Tasmar		Topiramate		Urea	
Taurine		Topiramate Actavis	128	Urex Forte	
Tecfidera		Total parenteral nutrition (TPN)	45	Urinary Agents	
Tegretol		TPN		Urinary Tract Infections	
Tegretol CR		Tramadol hydrochloride		Urinorm	
Telfast		Tramal SR 100		Uromitexan	
Teligent	94	Tramal SR 150	124	Ursodeoxycholic acid	2
Temaccord		Tramal SR 200	124	Ursosan	2
Temazepam	137	Trandate	50	Utrogestan	8
Temozolomide	155	Tranexamic acid	41	- V -	
Tenofovir disoproxil	100	Tranylcypromine sulphate	125	Vaccinations	26
Tenofovir Disoproxil Teva	100	Trastuzumab	218	Vaclovir	10
Tenoxicam	110	Trastuzumab emtansine	220	Valaciclovir	10
Tensipine MR10	51	Travatan	239	Valganciclovir	10
Tepadina		Travoprost	239	Valganciclovir Mylan	
Terbinafine		Treatments for Dementia		Vancomycin	9
Terbutaline sulphate		Treatments for Substance		Vannair	
Teriflunomide		Dependence	141	Varenicline Pfizer	14
Teriparatide		Trental 400		Varenicline tartrate	
Testosterone		Tretinoin		Varicella vaccine [Chickenpox	
Testosterone cipionate		Dermatological	61	vaccine]	27
Testosterone esters		Oncology		Varicella zoster virus (Oka strain	
Testosterone undecanoate		Trexate		attenuated vaccine [shingles	,
Tetrabenazine		Triamcinolone acetonide		vaccine]	27
Tetrabromophenol		Alimentary	33	Various	
Tetracosactrin		Dermatological		Varivax	
Tetracycline		Hormone		Vasodilators	
Thalidomide		Triamcinolone acetonide with		Vasopressin Agonists	
Thalomid		gramicidin, neomycin and nysi	atin	Vasorex	
Theophylline		Dermatological		Vedafil	
Thiamine hydrochloride	200 22	Sensory		Veletri	
THIO-TEPA	147	Triazolam		Venclexta	
11110-1F1 V	147	i iiazviaiii	101	ν οι ΙΟΙ <del>Ο</del> Λια	10

Venetoclax156	XMET Maxamum260
Venlafaxine	Xolair
Venomil227	XP Maxamum261
VENOX227	Xylocaine
Ventavis60	Xylocaine 2% Jelly119
Ventolin230	Xyntha40
Vepesid152	- <b>Z</b> -
Verapamil hydrochloride52	Zapril47
Vergo 16130	Zarontin126
Vermox89	Zaroxolyn53
Versacloz131	Zavedos152
Vesanoid	Zeffix100
Vexazone11	Zematop69
Vfend97	Zetlam100
Viaderm KC64	Ziagen105
Vidaza147	Zidovudine [AZT]105
Vigabatrin128	Zidovudine [AZT] with
Vigisom136	lamivudine 106
Vildagliptin11	Zimybe56
Vildagliptin with metformin	Zinc and castor oil65
hydrochloride12	Zinc sulphate36
Vimpat127	Zincaps36
Vinblastine sulphate157	Zinnat89
Vincristine sulphate157	Ziprasidone132
Vinorelbine157	Zista228
Vinorelbine Ebewe157	Zithromax90
Viramune Suspension105	Zoledronic acid
ViruPOS236	Hormone78
Vit.D333	Musculoskeletal114
Vita-B1233	Zoledronic acid Mylan78
VitA-POS240	Zopiclone
Vitabdeck34	Zopiclone Actavis137
Vital252	Zostavax276
Vitamin B complex33	Zostrix111
Vitamin B6 2533	Zostrix HP120
Vitamins33–34	Zuclopenthixol decanoate134
Vivonex TEN252	Zuclopenthixol hydrochloride132
Voltaren110	Zusdone132
Voltaren D110	Zyban142
Voltaren Ophtha237	Zypine132
Voltaren SR110	Zypine ODT132
Volumatic234	Zyprexa Relprevv133
Voriconazole97	Zytiga164
Votrient161	
Vttack97	
- W -	
Warfarin sodium44	
Wart Preparations	
Wasp venom allergy treatment227	
Water	
Blood	
Extemporaneous	
Wool fat with mineral oil65	
- X - Xarelto44	
Vifayon 0	