

June 2022
Volume 29 Number 1

Editors:

Kaye Wilson, Doris Chong, & Sophie Molloy
email: enquiry@pharmac.govt.nz
Telephone +64 4 460 4990
Facsimile +64 4 460 4995
Level 9, 40 Mercer Street
PO Box 10 254 Wellington

Freephone Information Line

0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at
no cost from the Pharmac website
www.pharmac.govt.nz/schedule.

You can register to have an electronic
version of the Pharmaceutical Schedule (link
to PDF copy) emailed to your nominated
email address each month by subscribing at
pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX.
XML version of the Schedule available from
schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering

email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3686



This work is licensed under the Creative
Commons Attribution 4.0 International licence.
In essence, you are free to copy, distribute
and adapt it, as long as you attribute the work
to Pharmac and abide by the other licence
terms. To view a copy of this licence, visit:
creativecommons.org/licenses/by/4.0/.
Attribution to Pharmac should be in written
form and not by reproduction of the Pharmac
logo. While care has been taken in compiling
this Schedule, Pharmac takes no responsibility
for any errors or omissions, and shall not be
liable for any consequences arising there from.

Introducing Pharmac 2

Section A General Rules **5**

Section B Alimentary Tract & Metabolism **6**

Blood & Blood Forming Organs **37**

Cardiovascular System **47**

Dermatologicals **61**

Genito Urinary System **71**

Hormone Preparations – Systemic **78**

Infections – Agents For Systemic Use **89**

Musculoskeletal System **110**

Nervous System **118**

Oncology Agents & Immunosuppressants **145**

Respiratory System & Allergies **227**

Sensory Organs **236**

Various **241**

Section C Extemporaneous Compounds (ECPs) **243**

Section D Special Foods **246**

Section I National Immunisation Schedule **267**

Index **277**

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list. The funding for pharmaceuticals comes from District Health Boards.

Pharmac's role:

“Secure for eligible people in need of pharmaceuticals, the best health outcomes that can reasonably be achieved, and from within the amount of funding provided.”

New Zealand Public Health and Disability Act 2000

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at <https://www.pharmac.govt.nz/about>.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in DHB Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in DHB Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to DHB Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in DHB Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in DHB hospitals. Section H lists the Pharmaceuticals that can be used in DHB hospitals and is a separate publication.

The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example

ANATOMICAL HEADING			
THERAPEUTIC HEADING			
CHEMICAL			
Presentation, form and strength		Subsidy (Manufacturer's Price) \$ Per	Fully Brand or Subsidised Generic ✓ Manufacturer
Presentation - Available on a PSO		10.00 100	Brand A ✓ Brand B
Presentation - Retail pharmacy-specialist		15.00 50	✓ Brand C
Presentation - Retail pharmacy-specialist		18.00 250 ml OP	✓ Brand D
a) Prescriptions must be written by a paediatrician or paediatric cardiologist; or b) on the recommendation of a paediatrician or a paediatric cardiologist			
CHEMICAL			
Presentation, form and strength		26.53 100	Brand E
		(35.27)	
Sole Supply/Principal Supply			
✓ Fully Subsidised			
▲ Three months supply may be dispensed at one time if endorsed 'certified exemption' by the prescriber or pharmacist.			

Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Practitioner's Supply Order

Conditions of and restrictions on prescribing (including Special Authority where it applies)

Three months or six months, as applicable, dispensed all-at-once

Brand or manufacturer's name

Sole Subsidised Supply product or Principal Supply Status

Fully subsidised product

Original Pack - Subsidy is rounded up to a multiple of whole packs

Quantity the Subsidy applies to

Subsidy paid on a product before mark-ups and GST

Manufacturer's Price if different from Subsidy

Glossary

Units of Measure

gram	g	microgram.....	mcg	millimole.....	mmol
kilogram.....	kg	milligram	mg	unit.....	u
international unit.....	iu	millilitre.....	ml		

Abbreviations

Ampoule	Amp	Gelatinous	Gel	Solution.....	Soln
Capsule	Cap	Granules	Gran	Suppository.....	Supp
Cream.....	Crm	Infusion	Inf	Tablet.....	Tab
Device.....	Dev	Injection	Inj	Tincture.....	Tinc
Dispersible.....	Disp	Liquid.....	Liq	Trans Dermal Delivery	
Effervescent.....	Eff	Long Acting.....	LA	System.....	TDDS
Emulsion.....	Emul	Ointment.....	Oint		
Enteric Coated.....	EC	Sachet	Sach		

Read the [General Rules](https://www.pharmac.govt.nz/section-a) : <https://www.pharmac.govt.nz/section-a>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Antacids and Antiflatulents

Antacids and Reflux Barrier Agents

ALGINIC ACID

Sodium alginate 225 mg and magnesium alginate 87.5 mg per sachet.....	5.31	30	✓	Gaviscon Infant
---	------	----	---	-----------------

SODIUM ALGINATE

* Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour.....	1.80 (8.60)	60		Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml.....	1.50 (5.50)	500 ml		Acidex

Phosphate Binding Agents

ALUMINIUM HYDROXIDE

* Tab 600 mg	12.56	100	✓	Alu-Tab
--------------------	-------	-----	---	---------

CALCIUM CARBONATE

Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml) – Subsidy by endorsement	39.00	500 ml	✓	Roxane
Only when prescribed for patients unable to swallow calcium carbonate tablets or where calcium carbonate tablets are inappropriate and the prescription is endorsed accordingly.				

Antidiarrhoeals

Agents Which Reduce Motility

LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on a PSO

* Tab 2 mg	10.75	400	✓	Nodia
* Cap 2 mg	6.25	400	✓	Diamide Relief

Rectal and Colonic Anti-inflammatories

BUDESONIDE

Cap 3 mg – Special Authority see SA1886 below – Retail pharmacy.....	166.50	90	✓	Entocort CIR
--	--------	----	---	--------------

► [SA1886](#) Special Authority for Subsidy

Initial application — (Crohn's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications).....	26.55	15 g OP 21.1 g OP	✓ Cortifoam ^{\$29} ✓ Colifoam
--	-------	----------------------	---

HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE

Topical aerosol foam, 1% with pramoxine hydrochloride 1%.....	26.55	10 g OP	✓ Proctofoam ^{\$29}
---	-------	---------	------------------------------

MESALAZINE

Tab 400 mg	49.50	100	✓ Asacol
Tab long-acting 500 mg.....	56.10	100	✓ Pentasa
Tab 800 mg	85.50	90	✓ Asacol
Modified release granules, 1 g	118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml	41.30	7	✓ Pentasa
Suppos 500 mg	22.80	20	✓ Asacol
Suppos 1 g	50.96	28	✓ Pentasa

OLSALAZINE

Tab 500 mg	93.37	100	✓ Dipentum
Cap 250 mg	53.00	100	✓ Dipentum

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications).....	74.10	1 OP	✓	Essential Prednisolone ^{S29}
SODIUM CROMOGLICATE				
Cap 100 mg.....	92.91	100	✓	Nalcrom
SULFASALAZINE				
* Tab 500 mg.....	14.00	100	✓	Salazopyrin
Tab EC 500 mg.....	15.53	100	✓	Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CINCHOCAINE				
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g.....	11.06	30 g OP	✓	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg.....	7.30	12	✓	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE				
Oint 5 mg with cinchocaine hydrochloride 5 mg per g.....	15.00	30 g OP	✓	Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g.....	9.90	12	✓	Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE – Special Authority see SA1329 below – Retail pharmacy				
* Oint 0.2%.....	22.00	30 g OP	✓	Rectogesic

► [SA1329](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE				
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO.....	65.45	10	✓	Max Health
HYOSCINE BUTYLBROMIDE				
* Tab 10 mg.....	6.35	100	✓	Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO.....	6.35	5	✓	Buscopan
MEBEVERINE HYDROCHLORIDE				
* Tab 135 mg.....	9.20	90	✓	Colofac

Antiulcerants

Antisecretory and Cytoprotective

MISOPROSTOL				
* Tab 200 mcg – Up to 120 tab available on a PSO.....	41.50	120	✓	Cytotec

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

Helicobacter Pylori Eradication

CLARITHROMYCIN

Tab 500 mg – Subsidy by endorsement.....	14.58	14	✓ Klacid
a) Maximum of 28 tab per prescription			
b) Subsidised only if prescribed for helicobacter pylori eradication and prescription is endorsed accordingly.			
Note: the prescription is considered endorsed if clarithromycin is prescribed in conjunction with a proton pump inhibitor and either amoxicillin or metronidazole.			

H2 Antagonists

FAMOTIDINE – Only on a prescription

* Tab 20 mg	4.91	100	✓ Famotidine Hovid <small>\$29</small>
* Tab 40 mg	8.48	100	✓ Famotidine Hovid <small>\$29</small>
* Inj 10 mg per ml, 4 ml – Subsidy by endorsement	57.02	10	✓ Mylan <small>\$29</small>
Subsidy by endorsement – Subsidised for patients receiving treatment as part of palliative care.			

Proton Pump Inhibitors

LANSOPRAZOLE

* Cap 15 mg	4.20	100	✓ Lanzol Relief
* Cap 30 mg	5.26	100	✓ Lanzol Relief

OMEPRAZOLE

For omeprazole suspension refer Standard Formulae, [page 243](#)

* Cap 10 mg	1.94	90	✓ Omeprazole actavis 10
* Cap 20 mg	1.86	90	✓ Omeprazole actavis 20
* Cap 40 mg	3.11	90	✓ Omeprazole actavis 40
* Powder – Only in combination.....	42.50	5 g	✓ Midwest
Only in extemporaneously compounded omeprazole suspension.			
* Inj 40 mg ampoule with diluent.....	33.98	5	✓ Dr Reddy's Omeprazole

PANTOPRAZOLE

* Tab EC 20 mg	2.02	100	✓ Panzop Relief
* Tab EC 40 mg	2.85	100	✓ Panzop Relief

Site Protective Agents

COLLOIDAL BISMUTH SUBCITRATE

Tab 120 mg	14.51	50	✓ Gastrodenol <small>\$29</small>
------------------	-------	----	--

SUCRALFATE

Tab 1 g	35.50 (48.28)	120	Carafate
---------------	------------------	-----	----------

Bile and Liver Therapy

RIFAXIMIN – Special Authority see [SA1461 on the next page](#) – Retail pharmacy

Tab 550 mg	625.00	56	✓ Xifaxan
------------------	--------	----	------------------

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

►SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE – Special Authority see SA1320 below – Retail pharmacy

Cap 25 mg	110.00	100	✓ Proglycem ^{\$29}
Cap 100 mg	280.00	100	✓ Proglycem ^{\$29}
Oral liq 50 mg per ml	620.00	30 ml OP	✓ Proglycem ^{\$29}

►SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit – Up to 5 kit available on a PSO.....	32.00	1	✓ Glucagen Hypokit
--	-------	---	--------------------

Insulin - Short-acting Preparations

INSULIN NEUTRAL

▲ Inj human 100 u per ml.....	25.26	10 ml OP	✓ Actrapid ✓ Humulin R
▲ Inj human 100 u per ml, 3 ml.....	42.66	5	✓ Actrapid Penfill ✓ Humulin R

Insulin - Intermediate-acting Preparations

INSULIN ASPART WITH INSULIN ASPART PROTAMINE

▲ Inj 100 iu per ml, 3 ml prefilled pen.....	52.15	5	✓ NovoMix 30 FlexPen
--	-------	---	----------------------

INSULIN ISOPHANE

▲ Inj human 100 u per ml.....	17.68	10 ml OP	✓ Humulin NPH ✓ Protaphane
▲ Inj human 100 u per ml, 3 ml.....	29.86	5	✓ Humulin NPH ✓ Protaphane Penfill

INSULIN ISOPHANE WITH INSULIN NEUTRAL

▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70 ✓ Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 40 ✓ PenMix 50

(PenMix 40 Inj human with neutral insulin 100 u per ml, 3 ml to be delisted 1 December 2022)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
▲ Inj lispro 25% with insulin lispro protamine 75% 100 u per ml, 3 ml.....	42.66	5	✓	Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per ml, 3 ml.....	42.66	5	✓	Humalog Mix 50
Insulin - Long-acting Preparations				
INSULIN GLARGINE				
▲ Inj 100 u per ml, 10 ml.....	63.00	1	✓	Lantus
▲ Inj 100 u per ml, 3 ml.....	94.50	5	✓	Lantus
▲ Inj 100 u per ml, 3 ml disposable pen.....	94.50	5	✓	Lantus SoloStar
Insulin - Rapid Acting Preparations				
INSULIN ASPART				
▲ Inj 100 u per ml, 10 ml.....	30.03	1	✓	NovoRapid
▲ Inj 100 u per ml, 3 ml.....	51.19	5	✓	NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓	NovoRapid FlexPen
INSULIN GLULISINE				
▲ Inj 100 u per ml, 10 ml.....	27.03	1	✓	Apidra
▲ Inj 100 u per ml, 3 ml.....	46.07	5	✓	Apidra
▲ Inj 100 u per ml, 3 ml disposable pen.....	46.07	5	✓	Apidra SoloStar
INSULIN LISPRO				
▲ Inj 100 u per ml, 10 ml.....	34.92	10 ml OP	✓	Humalog
▲ Inj 100 u per ml, 3 ml.....	59.52	5	✓	Humalog
Alpha Glucosidase Inhibitors				
ACARBOSE				
* Tab 50 mg	8.95	90	✓	Accarb
* Tab 100 mg	15.29	90	✓	Accarb
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
* Tab 5 mg	7.50	100	✓	Daonil
GLICLAZIDE				
* Tab 80 mg	15.18	500	✓	Glizide
GLIPIZIDE				
* Tab 5 mg	4.58	100	✓	Minidiab
METFORMIN HYDROCHLORIDE				
* Tab immediate-release 500 mg.....	14.74	1,000	✓	Metformin Mylan
* Tab immediate-release 850 mg.....	11.28	500	✓	Metformin Mylan
PIOGLITAZONE				
* Tab 15 mg	6.80	90	✓	Vexazone
* Tab 30 mg	7.30	90	✓	Vexazone
* Tab 45 mg	12.25	90	✓	Vexazone
VILDAGLIPTIN				
Tab 50 mg	35.00	60	✓	Galvus

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE				
Tab 50 mg with 1,000 mg metformin hydrochloride	35.00	60	✓	Galvumet
Tab 50 mg with 850 mg metformin hydrochloride	35.00	60	✓	Galvumet

GLP-1 Agonists

►SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

DULAGLUTIDE – Special Authority see [SA2065 above](#) – Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

* Inj 1.5mg per 0.5 ml prefilled pen 115.23 4 ✓ **Trulicity**

SGLT2 Inhibitors

►SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

- 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
- 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
- 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m² in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN – Special Authority see [SA2068 on the previous page](#) – Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

* Tab 10 mg	58.56	30	✓ Jardiance
* Tab 25 mg	58.56	30	✓ Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see [SA2068 on the previous page](#) – Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

* Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ Jardiamet
* Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	✓ Jardiamet
* Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	✓ Jardiamet
* Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	✓ Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP – Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
- 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Test strips	15.50	10 strip OP	✓ KetoSens
-------------------	-------	-------------	-------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER – Subsidy by endorsement

- Maximum of 1 pack per prescription
- Up to 1 pack available on a PSO
- A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - type 1 diabetes; or
 - permanent neonatal diabetes; or
 - undergone a pancreatectomy; or
 - cystic fibrosis-related diabetes; or
 - metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips.....	20.00	1 OP	✓ <u>CareSens Dual</u>
-----------------------------	-------	------	------------------------

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER – Subsidy by endorsement

- Maximum of 1 pack per prescription
- Up to 1 pack available on a PSO
- A diagnostic blood glucose test meter is subsidised for a patient who:
 - is receiving insulin or sulphonylurea therapy; or
 - is pregnant with diabetes; or
 - is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- type 1 diabetes; or
- permanent neonatal diabetes; or
- undergone a pancreatectomy; or
- cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

strips.....	10.00	1 OP	✓ <u>CareSens N</u>
			✓ <u>CareSens N POP</u>
	20.00		✓ <u>CareSens N Premier</u>

Note: Only 1 meter available per PSO

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP – Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	10.56	50 test OP	✓ CareSens N ✓ CareSens PRO
-------------------	-------	------------	--

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- 2) Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips.....	26.20	50 test OP	✓ SensoCard
--------------------------------	-------	------------	--------------------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES – Maximum of 200 dev per prescription

* 29 g x 12.7 mm	10.95	100	✓ B-D Micro-Fine
* 31 g x 5 mm	12.26	100	✓ B-D Micro-Fine
* 31 g x 6 mm	9.50	100	✓ Berpu
* 31 g x 8 mm	10.95	100	✓ B-D Micro-Fine
* 32 g x 4 mm	10.95	100	✓ B-D Micro-Fine

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE – Maximum of 200 dev per prescription				
* Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
	1.36	10		
	(1.99)			B-D Ultra Fine
* Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓	B-D Ultra Fine II
	1.30	10		
	(1.99)			B-D Ultra Fine II
* Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
	1.36	10		
	(1.99)			B-D Ultra Fine
* Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	✓	B-D Ultra Fine II
	1.36	10		
	(1.99)			B-D Ultra Fine II
* Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine
	1.36	10		
	(1.99)			B-D Ultra Fine
* Syringe 1 ml with 31 g x 8 mm needle	13.56	100	✓	B-D Ultra Fine II
	1.36	10		
	(1.99)			B-D Ultra Fine II

Insulin Pumps

INSULIN PUMP – Special Authority see [SA1603 below](#) – Retail pharmacy

- Maximum of 1 dev per prescription
- Only on a prescription
- Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓	MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓	Tandem t:slim X2 with Basal-IQ

►SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Patient has permanent neonatal diabetes; and
- A MDI regimen trial is inappropriate; and
- Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- Either:
 - Applicant is a relevant specialist; or
 - Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

4 Either:

- 4.1 Applicant is a relevant specialist; or
- 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatotomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatotomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Insulin Pump Consumables

►SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1c; and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

- than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from initial application; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

INSULIN PUMP CARTRIDGE – Special Authority see [SA1985 on page 19](#) – Retail pharmacy

- a) Maximum of 3 sets per prescription
 - b) Only on a prescription
 - c) Maximum of 13 packs of cartridge sets will be funded per year.
- | | | | |
|-----------------------------------|-------|------|--------------------|
| Cartridge 300 U, t:lock x 10..... | 50.00 | 1 OP | ✓ Tandem Cartridge |
|-----------------------------------|-------|------|--------------------|

INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special Authority see [SA1985 on page 19](#) – Retail pharmacy

- a) Maximum of 3 sets per prescription
 - b) Only on a prescription
 - c) Maximum of 13 infusion sets will be funded per year.
- | | | | |
|--|--------|------|------------------------------|
| 10 mm steel needle; 60 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-884A |
| 10 mm steel needle; 80 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-886A |
| 6 mm steel needle; 60 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-864A |
| 6 mm steel needle; 80 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-866A |
| 8 mm steel needle; 60 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-874A |
| 8 mm steel needle; 80 cm tubing x 10 | 130.00 | 1 OP | ✓ MiniMed Sure-T
MMT-876A |
| 6 mm steel needle; 29 G; manual insertion; 60 cm tubing x
10 with 10 needles; luer lock | 130.00 | 1 OP | ✓ Sure-T MMT-863 |
| 8 mm steel needle; 29 G; manual insertion; 60 cm tubing x
10 with 10 needles; luer lock | 130.00 | 1 OP | ✓ Sure-T MMT-873 |

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) – Special Authority see [SA1985 on page 19](#) –
Retail pharmacy

- a) Maximum of 3 sets per prescription
 - b) Only on a prescription
 - c) Maximum of 13 infusion sets will be funded per year.
- | | | | |
|--|--------|------|------------|
| 6 mm steel cannula; straight insertion; 80 cm line x 10 with
10 needles | 130.00 | 1 OP | ✓ TruSteel |
| 8 mm steel cannula; straight insertion; 80 cm line x 10 with
10 needles | 130.00 | 1 OP | ✓ TruSteel |
| 6 mm steel cannula; straight insertion; 60 cm line x 10 with
10 needles | 130.00 | 1 OP | ✓ TruSteel |
| 8 mm steel cannula; straight insertion; 60 cm line x 10 with
10 needles | 130.00 | 1 OP | ✓ TruSteel |

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA) – Special Authority see SA1985 on page 19 – Retail pharmacy				
a) Maximum of 3 set per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓	MiniMed Silhouette MMT-384A
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-398A
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	✓	MiniMed Mio MMT-941A
6 mm teflon needle, 45 cm pink tubing × 10.....	130.00	1 OP	✓	MiniMed Mio MMT-921A
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	✓	MiniMed Mio MMT-943A
6 mm teflon needle, 60 cm pink tubing × 10.....	130.00	1 OP	✓	MiniMed Mio MMT-923A
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-399A
6 mm teflon needle, 80 cm blue tubing.....	130.00	1 OP	✓	MiniMed Mio MMT-945A
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓	MiniMed Mio MMT-965A
6 mm teflon needle, 80 cm pink tubing × 10.....	130.00	1 OP	✓	MiniMed Mio MMT-925A
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-387A
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-396A
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-397A
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	✓	MiniMed Mio MMT-975A
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓	MiniMed Quick-Set MMT-386A

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION WITH INSERTION DEVICE) – Special Authority see SA1985 on page 19 – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon cannula; angle insertion; insertion device; 110 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
17 mm teflon cannula; angle insertion; 60 cm line x 10 with 10 needles; luer lock	130.00	1 OP	✓ Silhouette MMT-373
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION WITH INSERTION DEVICE) – Special Authority see SA1985 on page 19 – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
6 mm teflon cannula; straight insertion; insertion device; 110 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 110 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cm line x 10 with 10 needles	140.00	1 OP	✓ AutoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGHT INSERTION) – Special Authority see SA1985 on page 19 – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 infusion sets will be funded per year.			
6 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with 10 needles; luer lock	130.00	1 OP	✓ Quick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 on page 19 – Retail pharmacy			
a) Maximum of 3 sets per prescription			
b) Only on a prescription			
c) Maximum of 13 packs of reservoir sets will be funded per year.			
10 x luer lock conversion cartridges 1.8 ml for Paradigm pumps	50.00	1 OP	✓ ADR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml x 10	50.00	1 OP	✓ MiniMed 1.8 Reservoir MMT-326A
Cartridge for 7 series pump; 3.0 ml x 10	50.00	1 OP	✓ MiniMed 3.0 Reservoir MMT-332A

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Digestives Including Enzymes

PANCREATIC ENZYME

Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓	Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓	Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓	Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph Eur U)	34.93	20 g OP	✓	Creon Micro
<i>(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease)) to be delisted 1 June 2023)</i>				

URSODEOXYCHOLIC ACID – Special Authority see [SA1739 below](#) – Retail pharmacy

Cap 250 mg	32.95	100	✓	Ursosan
------------------	-------	-----	---	----------------

► [SA1739](#) Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults; and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogeneic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Agents

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription

* Powder for oral soln.....	6.00	250 g OP	✓ Macro Organic Psyllium Husk
	12.20	500 g OP	✓ Konsyl-D

MUCILAGINOUS LAXATIVES WITH STIMULANTS

* Dry.....	6.02 (17.32)	500 g OP	Normacol Plus
	2.41 (8.72)	200 g OP	Normacol Plus

Faecal Softeners

DOCUSATE SODIUM – Only on a prescription

* Tab 50 mg	2.31	100	✓ Coloxyl
* Tab 120 mg	3.13	100	✓ Coloxyl

DOCUSATE SODIUM WITH SENNOSIDES

* Tab 50 mg with sennosides 8 mg.....	3.50	200	✓ Laxsol
---------------------------------------	------	-----	-----------------

POLOXAMER – Only on a prescription

Not funded for use in the ear.

* Oral drops 10%.....	3.98	30 ml OP	✓ Coloxyl
-----------------------	------	----------	------------------

Opioid Receptor Antagonists - Peripheral

METHYLNALTREXONE BROMIDE – Special Authority see [SA1691 below](#) – Retail pharmacy

Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246.00	7	✓ Relistor

► **SA1691** Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL				
* Suppos 3.6 g – Only on a prescription	9.25	20	✓ PSM	
LACTULOSE – Only on a prescription				
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ <u>Laevolac</u>	
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM BICARBONATE AND SODIUM CHLORIDE				
Powder for oral soln 13.125 g with potassium chloride 46.6 mg, sodium bicarbonate 178.5 mg and sodium chloride 350.7 mg	6.70	30	✓ <u>Molaxole</u>	
SODIUM ACID PHOSPHATE – Only on a prescription				
Enema 16% with sodium phosphate 8%	2.50	1	✓ <u>Fleet Phosphate Enema</u>	
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATE – Only on a prescription				
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per ml, 5 ml	29.98	50	✓ <u>Micolette</u>	

Stimulant Laxatives

BISACODYL – Only on a prescription				
* Tab 5 mg	5.80	200	✓ <u>Pharmacy Health</u>	
* Suppos 10 mg	3.69	10	✓ <u>Lax-Suppositories</u>	
SENNA – Only on a prescription				
* Tab, standardised	2.17	100		
	(8.21)			Senokot
	0.43	20		
	(2.06)			Senokot
SODIUM PICOSULFATE – Special Authority see SA2053 below – Retail pharmacy				
Oral soln 7.5 mg per ml	7.40	30 ml OP	✓ <u>Dulcolax SP Drop</u>	

►SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – Special Authority see SA1986 on the next page – Retail pharmacy				
Inj 50 mg vial	1,142.60	1	✓ <u>Myozyme</u>	

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authority see [SA2042 below](#) – Retail pharmacy

Tab 1,000 mg	CBS	90	✓ Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder	CBS	400 g	✓ Biomed

►SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE – Special Authority see [SA1987 on the next page](#) – Retail pharmacy

Powder for oral soln.....	575.00	180 g OP	✓ Cystadane
---------------------------	--------	----------	-------------

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

►SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 – Special Authority see [SA2039 below](#) – Retail pharmacy

Cap 120 mg.....	CBS	30	✓ Solgar
Cap 160 mg.....	CBS	60	✓ Go Healthy

►SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see [SA1988 below](#) – Retail pharmacy

Inj 1 mg per ml, 5 ml vial.....	2,234.00	1	✓ Naglazyme
---------------------------------	----------	---	-------------

►SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:
Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

IDURSULFASE – Special Authority see [SA1623 on the next page](#) – Retail pharmacy

Inj 2 mg per ml, 3 ml vial.....	4,608.30	1	✓ Elaprase
---------------------------------	----------	---	------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria:
All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – Special Authority see [SA1695 below](#) – Retail pharmacy

Inj 100 U per ml, 5 ml vial	1,335.16	1	✓ Aldurazyme
-----------------------------------	----------	---	--------------

►SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria:
All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysaccharidosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE – Special Authority see [SA2040 below](#) – Retail pharmacy

Tab 500 mg	CBS	30	✓ Solgar
Cap 250 mg	CBS	30	✓ Solgar
Cap 500 mg	CBS	60	✓ Balance
Oral liq 500 mg per 10 ml	CBS	300 ml	✓ Balance

►SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

- Both:
- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
 - 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN – Special Authority see [SA2041 on the next page](#) – Retail pharmacy

Tab 100 mg	CBS	100	✓ Country Life
Cap 100 mg	CBS	100	✓ Solgar

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

►SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE – Special Authority see [SA1989 below](#) – Retail pharmacy

Tab soluble 100 mg1,452.70 30 OP ✓ **Kuvan**

►SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE – Special Authority see [SA1599 below](#) – Retail pharmacy

Soln 100 mg per mlCBS 100 ml ✓ **Amzoate** S29

►SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see [SA1990 on the next page](#) – Retail pharmacy

Grans 483 mg per g2,016.00 174 g OP ✓ **Pheburane**

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE – Special Authority see [SA2043 below](#) – Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	✓ Life Extension

►SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA – Special Authority see [SA1880 below](#) – Retail pharmacy

Inj 200 unit vial	1,072.00	1	✓ Elelyso
-------------------------	----------	---	-----------

►SA1880 Special Authority for Subsidy

Special Authority approved by the Gaucher Treatment Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Co-ordinator, Gaucher Treatment Panel Phone: 04 460 4990
Pharmac PO Box 10 254 Facsimile: 04 916 7571
Wellington Email: gaucherpanel@pharmac.govt.nz

Completed application forms must be sent to the coordinator for the Gaucher Treatment Panel and will be considered by the Gaucher Treatment Panel at the next practicable opportunity.

Notification of the Gaucher Treatment Panel's decision will be sent to the patient, the applying clinician and the patient's GP (if specified).

Access Criteria

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1) The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2) Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by taliglucerase alfa or might be reasonably expected to compromise a response to therapy with taliglucerase alfa; and
- 3) Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 4) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations, are submitted to the Gaucher Panel for assessment; and
- 5) Any of the following:
- 6) 1) Patient has haematological complications such as haemoglobin less than 95 g/l, symptomatic anaemia, thrombocytopenia; at least two episodes of severely symptomatic splenic infarcts confirmed with imagery; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- massive symptomatic splenomegaly; or
- 2) Patient has skeletal complications such as acute bone crisis requiring hospitalisation or major pain management strategies; radiological MRI Evidence of incipient destruction of any major joint (e.g. hips or shoulder); spontaneous fractures or vertebral collapse; chronic bone pain not controlled by other pharmaceuticals; or
- 3) Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
- 4) Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
- 5) Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period.

*Unapproved indication

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1) Patient has demonstrated a symptomatic improvement or no deterioration in the main symptom for which therapy was initiated; and
- 2) Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and
- 3) Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and three yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4) Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 5) Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 6) Patient is compliant with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units), unless otherwise agreed by Pharmac; and
- 7) Supporting clinical information including test reports, MRI whole body STIR, haematological data, and other relevant investigations are submitted to the Gaucher Panel for assessment as required.

Mouth and Throat

Agents Used in Mouth Ulceration

BENZYDAMINE HYDROCHLORIDE

Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with

Endorsement	9.00	500 ml	
	(20.31)		Difflam

Additional subsidy by endorsement for a patient who has oral mucositis as a result of treatment for cancer, and the prescription is endorsed accordingly.

CARMELLOSE SODIUM WITH GELATIN AND PECTIN

Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder	8.48	28 g OP	
	(10.95)		Stomahesive

CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE

* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
	(6.00)		Bonjela

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
TRIAMCINOLONE ACETONIDE Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
Oropharyngeal Anti-infectives			
AMPHOTERICIN B Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE Oral gel 20 mg per g	4.74	40 g OP	✓ Decozol
NYSTATIN Oral liq 100,000 u per ml	1.76	24 ml OP	✓ Nilstat
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a PSO	1.89 2.46 2.84 3.15	3 5	✓ Vita-B12 ✓ Hydroxocobalamin Panpharma ✓ Neo-B12 ✓ Hydroxocobalamin Mercury Pharma
<i>(Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 November 2022)</i> <i>(Neo-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 November 2022)</i> <i>(Hydroxocobalamin Mercury Pharma Inj 1 mg per ml, 1 ml ampoule to be delisted 1 November 2022)</i>			
PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription * Tab 25 mg – No patient co-payment payable..... Tab 50 mg	2.70 23.45	90 500	✓ Vitamin B6 25 ✓ Pyridoxine multichem
THIAMINE HYDROCHLORIDE – Only on a prescription * Tab 50 mg	7.09	100	✓ Max Health
VITAMIN B COMPLEX * Tab, strong, BPC	7.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription * Tab 100 mg	9.90	500	✓ Cvite
Vitamin D			
ALFACALCIDOL * Cap 0.25 mcg	26.32	100	✓ One-Alpha
* Cap 1 mcg	87.98	100	✓ One-Alpha
* Oral drops 2 mcg per ml	60.68	20 ml OP	✓ One-Alpha

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
CALCITRIOL				
* Cap 0.25 mcg	7.95	100	✓	Calcitriol-AFT
* Cap 0.5 mcg	13.75	100	✓	Calcitriol-AFT
COLECALCIFEROL				
* Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription.....	2.95	12	✓	Vit.D3
* Oral liq 188 mcg per ml (7,500 iu per ml)	9.00	4.8 ml OP	✓	Puria

Multivitamin Preparations

MULTIVITAMIN RENAL – Special Authority see [SA1546 below](#) – Retail pharmacy

* Cap.....	6.49	30	✓	Clinicians Renal Vit
------------	------	----	---	-----------------------------

►SA1546 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or
- 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA).

MULTIVITAMINS – Special Authority see [SA1036 below](#) – Retail pharmacy

* Powder	72.00	200 g OP	✓	Paediatric Seravit
----------------	-------	----------	---	---------------------------

►SA1036 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins.

VITAMINS

* Tab (BPC cap strength).....	11.45	1,000	✓	Mvite
* Cap (fat soluble vitamins A, D, E, K) – Special Authority see SA1720 below – Retail pharmacy	23.40	60	✓	Vitabdeck

►SA1720 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has cystic fibrosis with pancreatic insufficiency; or
- 2 Patient is an infant or child with liver disease or short gut syndrome; or
- 3 Patient has severe malabsorption syndrome.

Minerals

Calcium

CALCIUM CARBONATE

* Tab 1.25 g (500 mg elemental)	6.69	250	✓	Calci-Tab 500
* Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsement.....	54.60	76	✓	Cacit ^{S29}
	260.00	100	✓	Calcium 500 mg Hexal ^{S29}

Subsidy by endorsement – Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable.

(Cacit ^{S29} Tab eff 1.25 g (500 mg elemental) to be delisted 1 July 2022)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
CALCIUM GLUCONATE				
* Inj 10%, 10 ml ampoule	32.00	10	✓	Max Health - Hameln ^{\$29}
	64.00	20	✓	Max Health ^{\$29}
CALCIUM LACTATE GLUCONATE WITH CALCIUM CARBONATE				
Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg elemental)				
– Subsidy by endorsement	52.00	20	✓	Calcium-Sandoz Forte ^{\$29}
	260.00	100	✓	Calcium-Sandoz Forte ^{\$29}
Subsidy by endorsement – Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable.				
<i>(Calcium-Sandoz Forte ^{\$29} Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg elemental) to be delisted 1 July 2022)</i>				
<i>(Calcium-Sandoz Forte ^{\$29} Tab eff 2.94 g with calcium carbonate 0.3 g (500 mg elemental) to be delisted 1 July 2022)</i>				

Fluoride

SODIUM FLUORIDE				
* Tab 1.1 mg (0.5 mg elemental)	8.64	100	✓	PSM

Iodine

POTASSIUM IODATE				
* Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓	NeuroTabs

Iron

FERROUS FUMARATE				
* Tab 200 mg (65 mg elemental)	3.04	100	✓	Ferro-tab
FERROUS FUMARATE WITH FOLIC ACID				
* Tab 310 mg (100 mg elemental) with folic acid 350 mcg	5.98	100	✓	Ferro-F-Tabs
Ferro-F-Tabs to be Principal Supply on 1 August 2022				
FERROUS SULFATE				
* Tab long-acting 325 mg (105 mg elemental)	3.09	30	✓	Ferrograd
* Oral liq 30 mg (6 mg elemental) per 1 ml	12.08	500 ml	✓	Ferodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority see SA1840 below – Retail pharmacy				
Inj 50 mg per ml, 10 ml vial	150.00	1	✓	Ferinject

►SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

* Inj 50 mg per ml, 2 ml ampoule	34.50	5	✓ Ferrosig
--	-------	---	------------

Magnesium

MAGNESIUM HYDROXIDE

Suspension 8%.....	33.60	355 ml	✓ Phillips Milk of Magnesia ^{\$29}
--------------------	-------	--------	--

MAGNESIUM SULPHATE

* Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ <u>Martindale</u>
---	-------	----	---------------------

Zinc

ZINC SULPHATE

* Cap 137.4 mg (50 mg elemental)	11.00	100	✓ <u>Zincaps</u>
--	-------	-----	------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Antianaemics

Hypoplastic and Haemolytic

►SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure; and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
EPOETIN ALFA – Special Authority see SA1775 on the previous page – Retail pharmacy				
Wastage claimable				
Inj 1,000 iu in 0.5 ml, syringe.....	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe.....	100.00	6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe.....	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe.....	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe.....	125.00	6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe.....	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe.....	175.00	6	✓	Binocrit
Inj 10,000 iu in 1 ml, syringe.....	197.50	6	✓	Binocrit
Inj 40,000 iu in 1 ml, syringe.....	250.00	1	✓	Binocrit

Megaloblastic

FOLIC ACID				
Tab 0.8 mg	26.60	1,000	✓	Folic Acid multichem
* Tab 5 mg	5.82	100	✓	Folic Acid Mylan
Oral liq 50 mcg per ml	27.82	25 ml OP	✓	Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] – [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial.....	612.50	1	✓	Alprolix
Inj 500 iu vial.....	1,225.00	1	✓	Alprolix
Inj 1,000 iu vial.....	2,450.00	1	✓	Alprolix
Inj 2,000 iu vial.....	4,900.00	1	✓	Alprolix
Inj 3,000 iu vial.....	7,350.00	1	✓	Alprolix
Inj 4,000 iu vial.....	9,800.00	1	✓	Alprolix

ELTROMBOPAG – Special Authority see [SA1743 below](#) – Retail pharmacy

Wastage claimable				
Tab 25 mg	1,550.00	28	✓	Revolade
Tab 50 mg	3,100.00	28	✓	Revolade

►SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB – [Xpharm] – Special Authority see [SA1969 below](#)

Inj 30 mg in 1 ml vial.....	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial.....	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial.....	12,492.00	1	✓ Hemlibra
Inj 150 mg in 1 ml vial.....	17,846.00	1	✓ Hemlibra

►SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

2 Either:

- 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
- 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and

3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and

4 There is no immediate plan for major surgery within the next 12 months; and

5 Either:

- 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and

6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] – [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	2,356.60	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION – [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FEIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] – [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu prefilled syringe	287.50	1	✓ Xyntha
Inj 500 iu prefilled syringe	575.00	1	✓ Xyntha
Inj 1,000 iu prefilled syringe	1,150.00	1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	✓ Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	✓ Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] – [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	435.00	1	✓ RIXUBIS
Inj 1,000 iu vial	870.00	1	✓ RIXUBIS
Inj 2,000 iu vial	1,740.00	1	✓ RIXUBIS
Inj 3,000 iu vial	2,610.00	1	✓ RIXUBIS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) – [Xpharm]				
For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.				
Inj 250 iu vial.....	210.00	1	✓	Advate
Inj 500 iu vial.....	420.00	1	✓	Advate
Inj 1,000 iu vial.....	840.00	1	✓	Advate
Inj 1,500 iu vial.....	1,260.00	1	✓	Advate
Inj 2,000 iu vial.....	1,680.00	1	✓	Advate
Inj 3,000 iu vial.....	2,520.00	1	✓	Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE FS) – [Xpharm]				
For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.				
Inj 250 iu vial.....	237.50	1	✓	Kogenate FS
Inj 500 iu vial.....	475.00	1	✓	Kogenate FS
Inj 1,000 iu vial.....	950.00	1	✓	Kogenate FS
Inj 2,000 iu vial.....	1,900.00	1	✓	Kogenate FS
Inj 3,000 iu vial.....	2,850.00	1	✓	Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII] – [Xpharm]				
For patients with haemophilia A receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.				
Inj 250 iu vial.....	300.00	1	✓	Adynovate
Inj 500 iu vial.....	600.00	1	✓	Adynovate
Inj 1,000 iu vial.....	1,200.00	1	✓	Adynovate
Inj 2,000 iu vial.....	2,400.00	1	✓	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml.....	28.50 (73.00)	5		Fibro-vein
TRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓	Mercury Pharma

Vitamin K

PHYTOMENADIONE				
Inj 2 mg per 0.2 ml – Up to 5 inj available on a PSO.....	8.00	5	✓	Konakion MM
Inj 10 mg per ml, 1 ml – Up to 5 inj available on a PSO.....	9.21	5	✓	Konakion MM

Antithrombotic Agents

Antiplatelet Agents

ASPIRIN				
* Tab 100 mg	10.80	990	✓	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	✓	Clopidogrel Multichem
DIPYRIDAMOLE				
* Tab long-acting 150 mg.....	10.90	60	✓	Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next page – Retail pharmacy				
* Tab 90 mg	90.00	56	✓	Brilinta

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM – Special Authority see [SA1646 below](#) – Retail pharmacy

Inj 20 mg in 0.2 ml syringe.....	31.28	10	✓	Clexane
Inj 40 mg in 0.4 ml syringe.....	42.49	10	✓	Clexane
Inj 60 mg in 0.6 ml syringe.....	60.67	10	✓	Clexane
Inj 80 mg in 0.8 ml syringe.....	80.89	10	✓	Clexane
Inj 100 mg in 1 ml syringe.....	101.30	10	✓	Clexane
Inj 120 mg in 0.8 ml syringe.....	125.87	10	✓	Clexane Forte
Inj 150 mg in 1 ml syringe.....	143.86	10	✓	Clexane Forte

►SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner.

Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule	72.84	50	✓	Pfizer
Inj 5,000 iu per ml, 1 ml	32.66	5	✓	DBL Heparin Sodium ^{S29}
	70.33		✓	Hospira
Inj 5,000 iu per ml, 5 ml ampoule	289.05	50	✓	Pfizer
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓	Hospira
	42.40		✓	Heparin DBL ^{S29}
	482.20	50	✓	Heparin DBL ^{S29}

HEPARINISED SALINE

Inj 10 iu per ml, 5 ml	65.48	50	✓	Pfizer
------------------------------	-------	----	---	--------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg – No more than 2 cap per day	76.36	60	✓	Pradaxa
Cap 110 mg	76.36	60	✓	Pradaxa
Cap 150 mg	76.36	60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg – No more than 1 tab per day	83.10	30	✓	Xarelto
Tab 15 mg – Up to 14 tab available on a PSO	77.56	28	✓	Xarelto
Tab 20 mg	77.56	28	✓	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	✓	Coumadin
	6.46	100	✓	Marevan
* Tab 2 mg	4.31	50	✓	Coumadin
* Tab 3 mg	10.03	100	✓	Marevan
* Tab 5 mg	5.93	50	✓	Coumadin
	11.48	100	✓	Marevan

Blood Colony-stimulating Factors

FILGRASTIM – Special Authority see SA1259 below – Retail pharmacy				
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	✓	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓	Nivestim

►SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia ($ANC < 0.5 \times 10^9/L$); or
- 5 Treatment of drug-induced prolonged neutropenia ($ANC < 0.5 \times 10^9/L$).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM – Special Authority see SA1912 below – Retail pharmacy				
Inj 6 mg per 0.6 ml syringe	1,080.00	1	✓	Neulastim

►SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*).

Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE]				
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO	30.65	5	✓	Biomed
* Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO	15.00	1	✓	Biomed
POTASSIUM CHLORIDE				
* Inj 75 mg per ml, 10 ml	65.00	50	✓	Juno
SODIUM BICARBONATE				
Inj 8.4%, 50 ml	21.40	1	✓	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination				
Inj 8.4%, 100 ml	21.95	1	✓	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.				
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.23	500 ml	✓	Baxter
	1.26	1,000 ml	✓	Baxter
Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule	35.50	5	✓	Biomed
For Sodium chloride oral liquid formulation refer Standard Formulae, page 243				
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO	2.80	20	✓	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO	5.40	50	✓	Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	✓	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)				
Infusion	CBS	1 OP	✓	TPN
WATER				
1) On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or				
2) On a bulk supply order; or				
3) When used in the extemporaneous compounding of eye drops; or				
4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.				
Inj 10 ml ampoule – Up to 5 inj available on a PSO	7.19	50	✓	Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO	5.00	20	✓	Fresenius Kabi
			✓	Multichem
Oral Administration				
CALCIUM POLYSTYRENE SULPHONATE				
Powder	169.85	300 g OP	✓	Calcium Resonium
COMPOUND ELECTROLYTES				
Powder for oral soln – Up to 5 sach available on a PSO	9.77	50	✓	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]				
Soln with electrolytes (2 x 500 ml)	6.55	1,000 ml OP	✓	Pedialyte - Bubblégum

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

BLOOD AND BLOOD FORMING ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓	Phosphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq).....	5.26 (17.10)	60		Chlorvescent
* Tab long-acting 600 mg (8 mmol).....	8.90	200	✓	Span-K
SODIUM BICARBONATE				
Cap 840 mg.....	8.52	100	✓	Sodibic
			✓	Sodibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84.65	454 g OP	✓	Resonium-A

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN

* Tab 2 mg	17.35	500	✓ Apo-Doxazosin ✓ Doxazosin Clinect
* Tab 4 mg	20.94	500	✓ Apo-Doxazosin ✓ Doxazosin Clinect

(Apo-Doxazosin Tab 2 mg to be delisted 1 September 2022)

(Apo-Doxazosin Tab 4 mg to be delisted 1 September 2022)

PHENOXYBENZAMINE HYDROCHLORIDE

* Cap 10 mg	65.00	30	✓ BNM ^{\$29}
	216.67	100	✓ Dibenzyline ^{\$29}

PAZOSIN

* Tab 1 mg	5.53	100	✓ Arrotex-Prazosin S29 ^{\$29}
* Tab 2 mg	7.00	100	✓ Arrotex-Prazosin S29 ^{\$29}
* Tab 5 mg	11.70	100	✓ Arrotex-Prazosin S29 ^{\$29}

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAPTOPRIL

* Oral liq 5 mg per ml	94.99	95 ml OP	✓ Capoten
------------------------------	-------	----------	-----------

Oral liquid restricted to children under 12 years of age.

CILAZAPRIL – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

* Tab 0.5 mg	2.09	90	✓ <u>Zapril</u>
* Tab 2.5 mg	4.80	90	✓ <u>Zapril</u>
* Tab 5 mg	8.35	90	✓ <u>Zapril</u>

ENALAPRIL MALEATE

* Tab 5 mg	1.82	100	✓ <u>Acetec</u>
* Tab 10 mg	2.02	100	✓ <u>Acetec</u>
* Tab 20 mg	2.42	100	✓ <u>Acetec</u>

LISINOPRIL

* Tab 5 mg	11.07	90	✓ <u>Ethics Lisinopril</u>
* Tab 10 mg	11.67	90	✓ <u>Ethics Lisinopril</u>
* Tab 20 mg	14.69	90	✓ <u>Ethics Lisinopril</u>

PERINDOPRIL

Tab 2 mg	1.58	30	✓ <u>Coversyl</u>
Tab 4 mg	2.95	30	✓ <u>Coversyl</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
QUINAPRIL				
Tab 5 mg	5.97	90	✓	Arrow-Quinapril 5
Tab 10 mg	5.18	90	✓	Arrow-Quinapril 10
Tab 20 mg	7.95	90	✓	Arrow-Quinapril 20

ACE Inhibitors with Diuretics

QUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide.

Tab 10 mg with hydrochlorothiazide 12.5 mg	4.10	30	✓	Accuretic 10
Tab 20 mg with hydrochlorothiazide 12.5 mg	5.25	30	✓	Accuretic 20

Angiotensin II Antagonists

CANDESARTAN CILEXETIL

* Tab 4 mg	2.00	90	✓	Candestar
* Tab 8 mg	2.28	90	✓	Candestar
* Tab 16 mg	3.31	90	✓	Candestar
* Tab 32 mg	5.26	90	✓	Candestar

LOSARTAN POTASSIUM

Tab 12.5 mg	1.56	84	✓	Losartan Actavis
Tab 25 mg	1.84	84	✓	Losartan Actavis
Tab 50 mg	2.25	84	✓	Losartan Actavis
Tab 100 mg	3.50	84	✓	Losartan Actavis

Angiotensin II Antagonists with Diuretics

LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE

Tab 50 mg with hydrochlorothiazide 12.5 mg	15.25	30	✓	Arrow-Losartan & Hydrochlorothiazide
--	-------	----	---	---

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN – Special Authority see [SA1905 below](#) – Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓	Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓	Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓	Entresto 97/103

➔ [SA1905](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

3 Either:

- 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
- 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and

4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Local, [page 118](#)

AMIODARONE HYDROCHLORIDE

▲ Tab 100 mg	3.80	30	✓ Aratac
▲ Tab 200 mg	5.25	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a PSO	16.37	10	✓ Max Health

ATROPINE SULPHATE

* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	15.09	10	✓ Martindale
--	-------	----	---------------------

DIGOXIN

* Tab 62.5 mcg – Up to 30 tab available on a PSO	7.00	240	✓ Lanoxin PG
* Tab 250 mcg – Up to 30 tab available on a PSO	15.20	240	✓ Lanoxin
* Oral liq 50 mcg per ml	16.60	60 ml	✓ Lanoxin
			✓ Lanoxin Paediatric
			Elixir ^{\$29}
			✓ Lanoxin S29 ^{\$29}

DISOPYRAMIDE PHOSPHATE

▲ Cap 100 mg	23.87	100	✓ Rythmodan
--------------------	-------	-----	--------------------

FLECAINIDE ACETATE

▲ Tab 50 mg	19.95	60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	39.51	90	✓ Flecainide
			Controlled
			Release Teva
▲ Cap long-acting 200 mg	61.06	90	✓ Flecainide
			Controlled
			Release Teva
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambacor

MEXILETINE HYDROCHLORIDE

▲ Cap 150 mg	162.00	100	✓ Teva ^{\$29}
▲ Cap 250 mg	202.00	100	✓ Teva ^{\$29}

PROPAFENONE HYDROCHLORIDE

▲ Tab 150 mg	40.90	50	✓ Rytmonorm
--------------------	-------	----	--------------------

Antihypotensives

MIDODRINE – Special Authority see [SA1474 on the next page](#) – Retail pharmacy

Tab 2.5 mg	53.00	100	✓ Gutron
Tab 5 mg	79.00	100	✓ Gutron

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

►SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL

* Tab 50 mg	9.33	500	✓ <u>Mylan Atenolol</u>
* Tab 100 mg	14.20	500	✓ <u>Mylan Atenolol</u>
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ <u>Atenolol AFT</u>
			S29 S29
	38.20		✓ <u>Essential</u>
			Generics S29
	49.85		✓ <u>Atenolol AFT</u>

Restricted to children under 12 years of age.

BISOPROLOL FUMARATE

* Tab 2.5 mg	1.84	90	✓ <u>Bisoprolol Mylan</u>
* Tab 5 mg	2.55	90	✓ <u>Bisoprolol Mylan</u>
* Tab 10 mg	3.62	90	✓ <u>Bisoprolol Mylan</u>

CARVEDILOL

* Tab 6.25 mg	2.24	60	✓ <u>Carvedilol Sandoz</u>
* Tab 12.5 mg	2.30	60	✓ <u>Carvedilol Sandoz</u>
* Tab 25 mg	2.95	60	✓ <u>Carvedilol Sandoz</u>

LABETALOL

* Tab 100 mg	14.50	100	✓ <u>Trandate</u>
* Tab 200 mg	27.00	100	✓ <u>Trandate</u>
* Inj 5 mg per ml, 20 ml ampoule	59.06	5	
	(88.60)		Trandate
* inj 5 mg per ml, 20 ml vial	42.29	1	
	(48.20)		Alvogen S29

METOPROLOL SUCCINATE

* Tab long-acting 23.75 mg	1.45	30	✓ <u>Betaloc CR</u>
* Tab long-acting 47.5 mg	1.43	30	✓ <u>Betaloc CR</u>
* Tab long-acting 95 mg	2.15	30	✓ <u>Betaloc CR</u>
* Tab long-acting 190 mg	4.27	30	✓ <u>Betaloc CR</u>

METOPROLOL TARTRATE

Tab 50 mg	5.66	100	✓ <u>IPCA-Metoprolol</u>
Tab 100 mg	7.55	60	✓ <u>IPCA-Metoprolol</u>
* Tab long-acting 200 mg	23.40	28	✓ <u>Slow-Lopresor</u>
* Inj 1 mg per ml, 5 ml vial	26.50	5	✓ <u>Metoprolol IV Mylan</u>

NADOLOL

Tab 40 mg	19.19	100	✓ <u>Nadolol BNM</u> S29
Tab 80 mg	30.39	100	✓ <u>Nadolol BNM</u> S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PROPRANOLOL				
Tab 10 mg	7.04	100	✓	Drofate
Tab 40 mg	8.75	100	✓	IPCA-Propranolol
* Cap long-acting 160 mg	18.17	100	✓	Cardinol LA
* Oral liq 4 mg per ml – Special Authority see SA1327 below –				
Retail pharmacy.....	CBS	500 ml	✓	Roxane-Propranolol <small>\$29</small>

➤ **SA1327 Special Authority for Subsidy**

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrhythmias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrhythmias or congenital cardiac abnormalities.

SOTALOL

* Tab 80 mg	32.58	500	✓	Mylan
* Tab 160 mg	10.98	100	✓	Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

AMLODIPINE

* Tab 2.5 mg	1.08	90	✓	Vasorex
* Tab 5 mg	0.96	90	✓	Vasorex
* Tab 10 mg	1.19	90	✓	Vasorex

FELODIPINE

* Tab long-acting 2.5 mg	1.45	30	✓	Plendil ER
* Tab long-acting 5 mg	4.07	90	✓	Felo 5 ER
* Tab long-acting 10 mg	4.32	90	✓	Felo 10 ER

NIFEDIPINE

* Tab long-acting 10 mg	18.80	56	✓	Tensipine MR10 <small>\$29</small>
* Tab long-acting 20 mg	9.12	50	✓	Mylan (12 hr release) <small>\$29</small>
	17.72	100	✓	Nyefax Retard
* Tab long-acting 30 mg	4.78	14	✓	Mylan Italy (24 hr release) <small>\$29</small>
	34.10	100	✓	Mylan (24 hr release) <small>\$29</small>
* Tab long-acting 60 mg	52.81	100	✓	Mylan (24 hr release) <small>\$29</small>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	✓	Accord ^{S29}
* Cap long-acting 120 mg	33.42	500	✓	Apo-Diltiazem CD
* Cap long-acting 180 mg	7.00	30	✓	Cardizem CD
* Cap long-acting 240 mg	9.30	30	✓	Cardizem CD
PERHEXILINE MALEATE				
* Tab 100 mg	62.90	100	✓	Pexsig
VERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	✓	Isoptin
* Tab 80 mg	11.74	100	✓	Isoptin
* Tab long-acting 120 mg	36.02	100	✓	Isoptin Retard ^{S29}
			✓	Isoptin SR
			✓	Isoptin SR
* Tab long-acting 240 mg	15.12	30		
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	25.00	5	✓	Isoptin
Centrally-Acting Agents				
CLONIDINE				
* Patch 2.5 mg, 100 mcg per day – Only on a prescription	10.34	4	✓	Mylan
* Patch 5 mg, 200 mcg per day – Only on a prescription	13.18	4	✓	Mylan
* Patch 7.5 mg, 300 mcg per day – Only on a prescription	16.93	4	✓	Mylan
CLONIDINE HYDROCHLORIDE				
* Tab 25 mcg	8.75	112	✓	Clonidine BNM
	29.32		✓	Clonidine Teva
* Tab 150 mcg	37.07	100	✓	Catapres
* Inj 150 mcg per ml, 1 ml ampoule	29.68	10	✓	Medsurge
<i>(Clonidine BNM Tab 25 mcg to be delisted 1 November 2022)</i>				
METHYLDOPA				
* Tab 250 mg	15.10	100	✓	Methyldopa Mylan
	52.85	500	✓	Methyldopa Mylan
				S29 ^{S29}

Diuretics

Loop Diuretics

BUMETANIDE				
* Tab 1 mg	4.91	30	✓	Burinex S29 ^{S29}
	16.36	100	✓	Burinex
* Inj 500 mcg per ml, 4 ml vial	7.95	5	✓	Burinex

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
FUROSEMIDE [FRUSEMIDE]				
Tab 40 mg – Up to 30 tab available on a PSO	8.00	1,000	✓	IPCA-Frusemide
* Tab 500 mg	25.00	50	✓	Urex Forte
	89.48		✓	Furosemid- Ratiopharm <small>\$29</small>
	169.96	100	✓	Furosemid- Ratiopharm <small>\$29</small>
* Oral liq 10 mg per ml	11.20	30 ml OP	✓	Lasix
* Inj 10 mg per ml, 25 ml ampoule	60.65	6	✓	Lasix
* Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	1.15	5	✓	Furosemide-Baxter

Potassium Sparing Diuretics

AMILORIDE HYDROCHLORIDE				
Oral liq 1 mg per ml	32.10	25 ml OP	✓	Biomed
EPLERENONE – Special Authority see SA1728 below – Retail pharmacy				
Tab 25 mg	18.50	30	✓	Inspira
Tab 50 mg	25.00	30	✓	Inspira

►SA1728 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has heart failure with ejection fraction less than 40%; and
- 2 Either:
 - 2.1 Patient is intolerant to optimal dosing of spironolactone; or
 - 2.2 Patient has experienced a clinically significant adverse effect while on optimal dosing of spironolactone.

METOLAZONE				
Tab 5 mg	CBS	1	✓	Metolazone <small>\$29</small>
		50	✓	Zaroxolyn <small>\$29</small>
SPIRONOLACTONE				
* Tab 25 mg	3.68	100	✓	Spiractin
Spiractin to be Principal Supply on 1 September 2022				
* Tab 100 mg	10.65	100	✓	Spiractin
Spiractin to be Principal Supply on 1 September 2022				
Oral liq 5 mg per ml	30.60	25 ml OP	✓	Biomed

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE				
* Tab 5 mg with furosemide 40 mg	8.63	28	✓	Frumil
AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE				
* Tab 5 mg with hydrochlorothiazide 50 mg	5.00	50	✓	Moduretic

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
* Tab 2.5 mg – Up to 150 tab available on a PSO	20.00	500	✓	<u>Arrow-Bendrofluazide</u>
May be supplied on a PSO for reasons other than emergency.				
* Tab 5 mg	34.55	500	✓	<u>Arrow-Bendrofluazide</u>
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	27.82	25 ml OP	✓	<u>Biomed</u>
CHLORTALIDONE [CHLORTHALIDONE]				
Tab 25 mg	3.90	30	✓	<u>Igroton</u> ^{\$29}
	6.50	50	✓	<u>Hygroton</u>
INDAPAMIDE				
* Tab 2.5 mg	10.45	90	✓	<u>Dapa-Tabs</u>
	11.61	100	✓	<u>Mylan Indapamide</u> ^{\$29}

Lipid-Modifying Agents

Fibrates

BEZAFIBRATE				
* Tab 200 mg	19.46	90	✓	<u>Bezalip</u>
* Tab long-acting 400 mg	21.21	30	✓	<u>Bezalip Retard</u>

Other Lipid-Modifying Agents

ACIPIMOX				
* Cap 250 mg	21.56	30	✓	<u>Olbetam</u>
			✓	<u>Olbetam</u> ^{\$29}

Resins

COLESTIPOL HYDROCHLORIDE				
Grans for oral liq 5 g	32.89	30	✓	<u>Colestid</u>

HMG CoA Reductase Inhibitors (Statins)

ATORVASTATIN				
* Tab 10 mg	6.16	500	✓	<u>Lorstat</u>
* Tab 20 mg	9.24	500	✓	<u>Lorstat</u>
* Tab 40 mg	14.92	500	✓	<u>Lorstat</u>
* Tab 80 mg	26.54	500	✓	<u>Lorstat</u>
PRAVASTATIN				
* Tab 20 mg	2.11	28	✓	<u>Pravastatin Mylan</u>
* Tab 40 mg	3.61	28	✓	<u>Pravastatin Mylan</u>

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ROSUVASTATIN – Special Authority see SA2093 below – Retail pharmacy				
* Tab 5 mg	1.70	30	✓	Rosuvastatin Viatris
* Tab 10 mg	2.42	30	✓	Rosuvastatin Viatris
* Tab 20 mg	3.92	30	✓	Rosuvastatin Viatris
* Tab 40 mg	5.28	30	✓	Rosuvastatin Viatris

► **SA2093 Special Authority for Subsidy**

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient is considered to be at risk of cardiovascular disease; and
- 1.2 Patient is Māori or any Pacific ethnicity; or

2 Both:

- 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
- 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIMVASTATIN

* Tab 10 mg	1.23	90	✓	Simvastatin Mylan
* Tab 20 mg	2.03	90	✓	Simvastatin Mylan
* Tab 40 mg	3.58	90	✓	Simvastatin Mylan
* Tab 80 mg	7.12	90	✓	Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZETIMIBE – Special Authority see [SA1045 on the next page](#) – Retail pharmacy

* Tab 10 mg	1.95	30	✓	Ezetimibe Sandoz
-------------------	------	----	---	-------------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN – Special Authority see SA1046 below – Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	✓ Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	✓ Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

►SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:
All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

* Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Spray
* Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
* Patch 50 mg, 10 mg per day	18.62	30	✓ Nitroderm TTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ISOSORBIDE MONONITRATE				
* Tab 20 mg	19.55	100	✓	Ismo 20
* Tab long-acting 40 mg.....	8.20	30	✓	Ismo 40 Retard
* Tab long-acting 60 mg.....	9.25	90	✓	Duride

Sympathomimetics

ADRENALINE

Inj 1 in 1,000, 1 ml ampoule – Up to 5 inj available on a PSO	4.98	5	✓	Aspen Adrenaline
	10.76		✓	DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PSO	27.00	5	✓	Hospira
	49.00	10	✓	Aspen Adrenaline

Vasodilators

HYDRALAZINE HYDROCHLORIDE

* Tab 25 mg – Special Authority see SA1321 below – Retail pharmacy.....	CBS	1	✓	Hydralazine
		56	✓	Onelink ^{S29}
		84	✓	AMDIPHARM ^{S29}
		100	✓	Onelink ^{S29}
* Inj 20 mg ampoule.....	25.90	5	✓	Apresoline

►SA1321 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 For the treatment of refractory hypertension; or
- 2 For the treatment of heart failure in combination with a nitrate, in patients who are intolerant or have not responded to ACE inhibitors and/or angiotensin receptor blockers.

MINOXIDIL

▲ Tab 10 mg	70.00	100	✓	Loniten
-------------------	-------	-----	---	----------------

NICORANDIL

▲ Tab 10 mg	25.57	60	✓	Ikorel
▲ Tab 20 mg	32.28	60	✓	Ikorel

PAPAVERINE HYDROCHLORIDE

* Inj 12 mg per ml, 10 ml ampoule.....	257.12	5	✓	Hospira
--	--------	---	---	----------------

PENTOXIFYLLINE [OXPENTIFYLLINE]

Tab 400 mg	42.26	50	✓	Trental 400
------------------	-------	----	---	--------------------

Endothelin Receptor Antagonists

AMBRISENTAN – Special Authority see [SA1702 below](#) – Retail pharmacy

Tab 5 mg	1,550.00	30	✓	Ambrisentan Mylan
Tab 10 mg	1,550.00	30	✓	Ambrisentan Mylan
			✓	Mylan

►SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
BOSENTAN – Special Authority see SA1991 below – Retail pharmacy				
Tab 62.5 mg	119.85	60	✓	Bosentan Dr Reddy's
Tab 125 mg	119.85	60	✓	Bosentan Dr Reddy's

➔SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

- 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
- 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail pharmacy

Tab 25 mg	0.85	4	✓ Vedafil
Tab 50 mg	1.70	4	✓ Vedafil
Tab 100 mg	10.20	12	✓ Vedafil

➡SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm⁻⁵); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see [SA1696 below](#) – Retail pharmacy

Inj 500 mcg vial.....	36.61	1	✓ Veletri
Inj 1.5 mg vial	73.21	1	✓ Veletri

►► **SA1696** Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST – Special Authority see [SA1705 below](#) – Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml	740.10	30	✓ Ventavis
--	--------	----	-------------------

►► **SA1705** Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, [page 89](#)

ADAPALENE

- Maximum of 30 g per prescription
- Only on a prescription

Crm 0.1%.....	22.89	30 g OP	✓ Differin
Gel 0.1%.....	22.89	30 g OP	✓ Differin

ISOTRETINOIN – Special Authority see [SA2023 below](#) – Retail pharmacy

Cap 5 mg.....	11.26	60	✓ Oratane
Cap 10 mg.....	18.75	120	✓ Oratane
Cap 20 mg.....	26.73	120	✓ Oratane

►SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- Either:
 - Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRETINOIN

Crm 0.5 mg per g – Maximum of 50 g per prescription	15.57	50 g OP	✓ ReTrieve
---	-------	---------	------------

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, [page 89](#)

HYDROGEN PEROXIDE

* Crm 1%.....	8.56	10 g OP	✓ Crystaderm
		15 g OP	✓ Crystaderm

DERMATOLOGICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
MUPIROCI				
Oint 2%.....	6.60 (11.50)	15 g OP		Bactroban
a) Only on a prescription				
b) Not in combination				
SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 2%.....	1.59	5 g OP	✓ Foban	
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
Oint 2%.....	1.59	5 g OP	✓ Foban	
a) Maximum of 5 g per prescription				
b) Only on a prescription				
c) Not in combination				
SULFADIAZINE SILVER				
Crm 1%.....	10.80	50 g OP	✓ Flamazine	
a) Up to 250 g available on a PSO				
b) Not in combination				

Antifungals Topical

For systemic antifungals, refer to INFECTIONS, Antifungals, [page 96](#)

AMOROLFINE				
a) Only on a prescription				
b) Not in combination				
Nail soln 5%.....	14.93	5 ml OP	✓ MycoNail	
CLOTRIMAZOLE				
* Crm 1%.....	0.77	20 g OP	✓ Clomazol	
a) Only on a prescription				
b) Not in combination				
* Soln 1%.....	4.36 (7.55)	20 ml OP		Canesten
a) Only on a prescription				
b) Not in combination				
ECONAZOLE NITRATE				
Crm 1%.....	1.00 (7.48)	20 g OP		Pevaryl
a) Only on a prescription				
b) Not in combination				
Foaming soln 1%, 10 ml sachets.....	9.89 (17.23)	3		Pevaryl
a) Only on a prescription				
b) Not in combination				

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
MICONAZOLE NITRATE				
* Crm 2%.....	0.81	15 g OP	✓	Multichem
a) Only on a prescription				
b) Not in combination				
* Lotn 2%	4.36 (10.03)	30 ml OP		Daktarin
a) Only on a prescription				
b) Not in combination				
* Tinct 2%.....	4.36 (12.10)	30 ml OP		Daktarin
a) Only on a prescription				
b) Not in combination				

Antipruritic Preparations

CALAMINE

- a) Only on a prescription
- b) Not in combination

Crm, aqueous, BP 1.08 100 g ✓ **Calamine-AFT**

CROTAMITON

- a) Only on a prescription
- b) Not in combination

Crm 10%..... 3.29 20 g OP ✓ **Itch-Soothe**

MENTHOL – Only in combination

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain
- 2) With or without other dermatological galenicals.

Crystals..... 6.92 25 g ✓ **MidWest**
 29.60 100 g ✓ **MidWest**

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, [page 79](#)

Corticosteroids - Plain

BETAMETHASONE DIPROPIONATE

Crm 0.05%..... 2.96 15 g OP ✓ **Diprosone**
 36.00 50 g OP ✓ **Diprosone**
 Oint 0.05%..... 2.96 15 g OP ✓ **Diprosone**
 36.00 50 g OP ✓ **Diprosone**
 Oint 0.05% in propylene glycol base 4.33 30 g OP ✓ **Diprosone OV**

BETAMETHASONE VALERATE

* Crm 0.1%..... 4.53 50 g OP ✓ **Beta Cream**
 * Oint 0.1%..... 5.84 50 g OP ✓ **Beta Ointment**
 * Lotn 0.1% 25.00 50 ml OP ✓ **Betnovate**

CLOBETASOL PROPIONATE

* Crm 0.05%..... 2.18 30 g OP ✓ **Dermol**
 * Oint 0.05%..... 2.12 30 g OP ✓ **Dermol**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
CLOBETASONE BUTYRATE				
Crm 0.05%.....	5.38 (10.00)	30 g OP		Eumovate
HYDROCORTISONE				
* Crm 1% – Only on a prescription.....	3.70	100 g OP	✓	<u>Hydrocortisone (PSM)</u>
	17.15	500 g	✓	<u>Hydrocortisone (PSM)</u>
* Powder – Only in combination.....	49.95	25 g	✓	<u>ABM</u>
Up to 5% in a dermatological base (not proprietary Topical Corticosteroid – Plain) with or without other dermatological galenicals				
HYDROCORTISONE AND PARAFFIN LIQUID AND LANOLIN				
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% – Only on a prescription	10.57	250 ml	✓	<u>DP Lotn HC</u>
HYDROCORTISONE BUTYRATE				
Lipocream 0.1%.....	4.85	100 g OP	✓	<u>Locoid Lipocream</u>
Oint 0.1%.....	10.28	100 g OP	✓	<u>Locoid</u>
Milky emul 0.1%	12.33	100 ml OP	✓	<u>Locoid Crelo</u>
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%.....	4.46	15 g OP	✓	<u>Advantan</u>
Oint 0.1%.....	4.46	15 g OP	✓	<u>Advantan</u>
MOMETASONE FUROATE				
Crm 0.1%.....	1.95	15 g OP	✓	<u>Elocon Alcohol Free</u>
	3.10	50 g OP	✓	<u>Elocon Alcohol Free</u>
Oint 0.1%.....	1.95	15 g OP	✓	<u>Elocon</u>
	2.90	50 g OP	✓	<u>Elocon</u>
Lotn 0.1%	4.50	30 ml OP	✓	<u>Elocon</u>
TRIAMCINOLONE ACETONIDE				
Crm 0.02%.....	6.30	100 g OP	✓	<u>Aristocort</u>
Oint 0.02%.....	6.35	100 g OP	✓	<u>Aristocort</u>

Corticosteroids - Combination

BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUSIDIC ACID]				
Crm 0.1% with sodium fusidate (fusidic acid) 2%.....	3.49 (10.45)	15 g OP		Fucicort
a) Maximum of 15 g per prescription				
b) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE – Only on a prescription				
* Crm 1% with miconazole nitrate 2%.....	1.89	15 g OP	✓	<u>Micreme H</u>
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN – Only on a prescription				
Crm 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	15 g OP	✓	<u>Pimafucort</u>
Oint 1% with natamycin 1% and neomycin sulphate 0.5%.....	3.35	15 g OP	✓	<u>Pimafucort</u>
<i>(Pimafucort Crm 1% with natamycin 1% and neomycin sulphate 0.5% to be delisted 1 April 2023)</i>				
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN				
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g – Only on a prescription	3.49 (9.28)	15 g OP		Viaderm KC

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE			
* Crm 5% pump bottle.....	4.48	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 5%</u>
* Crm 10% pump bottle.....	4.52	500 ml OP	✓ <u>healthE</u> <u>Dimethicone 10%</u>
ZINC AND CASTOR OIL			
* Oint.....	4.65	500 g	✓ <u>Boucher</u>
Emollients			
AQUEOUS CREAM			
* Crm.....	1.73	500 g	✓ <u>Boucher</u> ✓ <u>GEM Aqueous Cream</u>
GEM Aqueous Cream to be Principal Supply on 1 July 2022 (Boucher Crm to be delisted 1 August 2022)			
CETOMACROGOL			
* Crm BP.....	1.99	500 g	✓ <u>Cetomacrogol-AFT</u>
CETOMACROGOL WITH GLYCEROL			
Crm 90% with glycerol 10%.....	2.35	500 ml OP	✓ <u>Boucher</u> ✓ <u>Pharmacy Health Sorbolene with Glycerin</u>
	3.10	1,000 ml OP	✓ <u>Boucher</u>
EMULSIFYING OINTMENT			
* Oint BP.....	3.40	500 g	✓ <u>Emulsifying Ointment ADE</u>
OIL IN WATER EMULSION			
* Crm.....	2.04	500 g	✓ <u>Fatty Cream AFT</u> ✓ <u>O/W Fatty Emulsion Cream</u>
	2.19		
Fatty Cream AFT to be Principal Supply on 1 September 2022 (O/W Fatty Emulsion Cream Crm to be delisted 1 September 2022)			
PARAFFIN			
Oint liquid paraffin 50% with white soft paraffin 50%.....	5.35	500 ml OP	✓ <u>healthE</u>
UREA			
* Crm 10%.....	1.37	100 g OP	✓ <u>healthE Urea Cream</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
WOOL FAT WITH MINERAL OIL – Only on a prescription				
* Lotn hydrous 3% with mineral oil.....	5.60 (11.95)	1,000 ml		DP Lotion
	1.40 (4.53)	250 ml OP		DP Lotion
	5.60 (20.53)	1,000 ml		Alpha-Keri Lotion
	(23.91)			BK Lotion
	1.40 (7.73)	250 ml OP		BK Lotion

Other Dermatological Bases

PARAFFIN

White soft – Only in combination.....	4.99	450 g	✓ healthE
	19.99	2,500 g	✓ healthE

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid – Plain.

Minor Skin Infections

POVIDONE IODINE

Oint 10%.....	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%.....	4.15	100 ml	✓ Riodine
Antiseptic soln 10%.....	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol.....	1.63 (3.48)	100 ml	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol.....	1.63 (7.78)	100 ml	Pfizer

Parasiticial Preparations

DIMETHICONE

* Lotn 4%	4.98	200 ml OP	✓ healthE Dimethicone 4% Lotion
-----------------	------	-----------	--

IVERMECTIN – Special Authority see [SA1225 below](#) – Retail pharmacy

Tab 3 mg – Up to 100 tab available on a PSO	17.20	4	✓ Stromectol
---	-------	---	---------------------

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- 3) For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

► [SA1225](#) Special Authority for Subsidy

Initial application — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist.

Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides; or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%.....	5.75	30 g OP	✓ Lyderm
Lotn 5%	3.99	30 ml OP	✓ A-Scabies

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see [SA2024 below](#) – Retail pharmacy

Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

►SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment; or
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL

Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	✓ Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP	✓ Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	✓ Daivobet

CALCIPOTRIOL

Oint 50 mcg per g	40.00	120 g OP	✓ Daivonex
-------------------------	-------	----------	-------------------

COAL TAR

Soln BP – Only in combination.....	36.25	200 ml	✓ Midwest
------------------------------------	-------	--------	------------------

- 1) Up to 10% only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain
- 2) With or without other dermatological galenicals.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR				
Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and allantoin crm 2.5%.....	6.59 (8.00) 3.43 (4.35)	75 g OP 30 g OP		Egopsoryl TA Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR				
Soln 12% with salicylic acid 2% and sulphur 4% oint.....	4.97 7.95	25 g OP 40 g OP	✓ ✓	Coco-Scalp Coco-Scalp
PIMECROLIMUS – Special Authority see SA1970 below – Retail pharmacy				
a) Maximum of 15 g per prescription				
b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.				
Cream 1%.....	28.50	15 g OP	✓	Elidel
►SA1970 Special Authority for Subsidy				
Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Both:				
1 Patient has atopic dermatitis on the eyelid; and				
2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.				
PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN – Only on a prescription				
* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.....	4.44	500 ml	✓	Pinetarsol
SALICYLIC ACID				
Powder – Only in combination.....	18.88	250 g	✓	Midwest
1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain or collodion flexible				
2) With or without other dermatological galenicals.				
SULPHUR				
Precipitated – Only in combination.....	6.35	100 g	✓	Midwest
1) Only in combination with a dermatological base or proprietary Topical Corticosteroid – Plain				
2) With or without other dermatological galenicals.				
TACROLIMUS				
Oint 0.1% – Special Authority see SA2074 below – Retail pharmacy.....	33.00	30 g OP	✓	Zematop
a) Maximum of 30 g per prescription				
b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.				
►SA2074 Special Authority for Subsidy				
Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Both:				
1 Patient has atopic dermatitis on the face; and				
2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.				

Scalp Preparations

BETAMETHASONE VALERATE		
* Scalp app 0.1%	9.84	100 ml OP ✓ <u>Beta Scalp</u>
CLOBETASOL PROPIONATE		
* Scalp app 0.05%	5.69	30 ml OP ✓ <u>Dermol</u>
HYDROCORTISONE BUTYRATE		
Scalp lotn 0.1%	6.57	100 ml OP ✓ <u>Locoid</u>
KETOCONAZOLE		
Shampoo 2%	3.23	100 ml OP ✓ <u>Sebizole</u> ✓ <u>Sebizole</u>
a) Maximum of 100 ml per prescription		
b) Only on a prescription		

Sunscreens

SUNSCREENS, PROPRIETARY – Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Lotn,	5.10	200 g OP ✓ <u>Marine Blue Lotion</u> <u>SPF 50+</u>
-------------	------	--

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, [page 68](#)

IMIQUIMOD		
Crm 5%, 250 mg sachet	21.72	24 ✓ <u>Perrigo</u>
PODOPHYLLOTOXIN		
Soln 0.5%	33.60	3.5 ml OP ✓ <u>Condyline</u>
a) Maximum of 3.5 ml per prescription		
b) Only on a prescription		

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM		
Crm 5%	6.95	20 g OP ✓ <u>Efudix</u>

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Contraceptives - Non-hormonal				
Condoms				
CONDOMS				
* 49 mm – Up to 144 dev available on a PSO	11.42	144	✓	<u>Moments</u>
* 53 mm.....	0.95	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				
* 53 mm, 0.05 mm thickness.....	0.95	10	✓	<u>Moments</u>
	11.42	144	✓	<u>Moments</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 53 mm, chocolate, brown	0.95	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 53 mm, strawberry, red.....	0.95	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 56 mm.....	0.97	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				
* 56 mm, 0.05 mm thickness.....	1.30	12	✓	<u>Gold Knight</u>
	15.57	144	✓	<u>Gold Knight</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓	<u>Gold Knight</u>
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				
* 56 mm, 0.08 mm thickness.....	0.97	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 56 mm, 0.08 mm thickness, red	0.97	10	✓	<u>Moments</u>
	11.64	144	✓	<u>Moments</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 56 mm, chocolate	1.30	12	✓	<u>Gold Knight</u>
	15.57	144	✓	<u>Gold Knight</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 56 mm, strawberry.....	1.30	12	✓	<u>Gold Knight</u>
	15.57	144	✓	<u>Gold Knight</u>
a) Up to 60 dev available on a PSO				
b) Maximum of 60 dev per prescription				
* 60 mm.....	1.42	12	✓	<u>Gold Knight XL</u>
	14.87	144	✓	<u>Shield XL</u>
	17.02		✓	<u>Gold Knight XL</u>

- a) Maximum of 60 dev per prescription
 b) Up to 60 dev available on a PSO

\$29 Unapproved medicine supplied under Section 29

Sole Subsidised Supply

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
* 60 mm (bulk pack)	14.87	144	✓	<u>Gold Knight XL</u>
a) Maximum of 60 dev per prescription				
b) Up to 60 dev available on a PSO				

Contraceptive Devices

INTRA-UTERINE DEVICE

a) Up to 40 dev available on a PSO				
b) Only on a PSO				
* IUD 29.1 mm length × 23.2 mm width	18.45	1	✓	<u>Choice TT380 Short</u>
* IUD 33.6 mm length × 29.9 mm width	18.45	1	✓	<u>Choice</u> <u>TT380 Standard</u>
* IUD 35.5 mm length × 19.6 mm width	15.50	1	✓	<u>Choice Load 375</u>

Contraceptives - Hormonal

Combined Oral Contraceptives

►SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab – Up to 84 tab available on a PSO	10.00	84	✓	<u>Mercilon 28</u>
--	-------	----	---	--------------------

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets – Up to 112 tab available on a PSO	2.18 6.45	84 112	✓	Microgynon 20 ED Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg..... (16.50)	6.62	63		Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Authority see SA0500 on the previous page b) Up to 63 tab available on a PSO				
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets – Up to 112 tab available on a PSO	1.77 6.45	84 112	✓	Levlen ED Femme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab – Up to 84 tab available on a PSO.....	6.95	84	✓	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab – Up to 84 tab available on a PSO.....	21.99 29.32	84 112	✓	Norimin Norimin

Progestogen-only Contraceptives

► **SA0500** Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL

* Tab 30 mcg – Up to 84 tab available on a PSO	16.50	84	✓	Microlut
	22.00	112	✓	Microlut
* Subdermal implant (2 × 75 mg rods) – Up to 3 pack available on a PSO.....	106.92	1	✓	Jadelle

MEDROXYPROGESTERONE ACETATE

Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO	7.98	1	✓	Depo-Provera
--	------	---	---	---------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
NORETHISTERONE				
Tab 350 mcg – Up to 84 tab available on a PSO	12.25	84	✓	Noriday 28

Emergency Contraceptives

LEVONORGESTREL				
* Tab 1.5 mg	4.95	1	✓	Postinor-1
a) Maximum of 2 tab per prescription				
b) Up to 5 tab available on a PSO				
c) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.				

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions “contraceptive” (code “O”) when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

* Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO	4.98	168	✓	Ginet
--	------	-----	---	--------------

Gynaecological Anti-infectives

ACETIC ACID WITH HYDROXYQUINOLINE AND RICINOLEIC ACID

Jelly with glacial acetic acid 0.94%, hydroxyquinoline sulphate 0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator	8.43 (24.00)	100 g OP		Aci-Jel
---	-----------------	----------	--	---------

CLOTRIMAZOLE

* Vaginal crm 1% with applicators	2.50	35 g OP	✓	Clomazol
* Vaginal crm 2% with applicators	3.00	20 g OP	✓	Clomazol

MICONAZOLE NITRATE

* Vaginal crm 2% with applicator	6.89	40 g OP	✓	Micreme
--	------	---------	---	----------------

NYSTATIN

Vaginal crm 100,000 u per 5 g with applicator(s)	4.00	75 g OP	✓	Nilstat
--	------	---------	---	----------------

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE

Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	160.00	5	✓	DBL Ergometrine
---	--------	---	---	------------------------

OESTRIOL

* Crm 1 mg per g with applicator	6.62	15 g OP	✓	Ovestin
* Pessaries 500 mcg	6.86	15	✓	Ovestin

OXYTOCIN – Up to 5 inj available on a PSO

Inj 5 iu per ml, 1 ml ampoule	3.98	5	✓	Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	✓	Oxytocin BNM

OXYTOCIN WITH ERGOMETRINE MALEATE – Up to 5 inj available on a PSO

Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml	30.00	5	✓	Syntometrine
--	-------	---	---	---------------------

▲Three months supply may be dispensed at one time if endorsed “certified exemption” by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Cassette	12.00	40 test OP	✓ David One Step Cassette Pregnancy Test
			✓ Smith BioMed Rapid Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, [page 106](#)

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see [SA0928 below](#) – Retail pharmacy

* Tab 5 mg	4.81	100	✓ <u>Ricit</u>
------------------	------	-----	----------------

➔ [SA0928](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE – Special Authority see [SA1032 below](#) – Retail pharmacy

* Cap 400 mcg	17.73	100	✓ <u>Tamsulosin-Rex</u>
---------------------	-------	-----	-------------------------

➔ [SA1032](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

* Tab 5 mg	5.42	100	✓ Alchemy Oxybutynin ^{\$29}
------------------	------	-----	---

POTASSIUM CITRATE

Oral liq 3 mmol per ml – Special Authority see [SA1083 on the](#)

next page – Retail pharmacy	31.80	200 ml OP	✓ Biomed
---	-------	-----------	----------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:
Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE

* Grans eff 4 g sachets2.22 28 ✓ **Ural**

SOLIFENACIN SUCCINATE

Tab 5 mg2.05 30 ✓ **Solifenacin Mylan**
Tab 10 mg3.72 30 ✓ **Solifenacin Mylan**

Detection of Substances in Urine

ORTHO-TOLIDINE

* Compound diagnostic sticks7.50 50 test OP
(8.25) Hemastix

TETRABROMOPHENOL

* Blue diagnostic strips7.02 100 test OP
(13.92) Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE

Tab 200 mg60.00 1 ✓ **Mifegyne**
180.00 3 ✓ **Mifegyne**
a) Up to 15 tab available on a PSO
b) Only on a PSO

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Calcium Homeostasis

CALCITONIN

* Inj 100 iu per ml, 1 ml ampoule 121.00 5 ✓ **Miacalcic**

CINACALCET – Special Authority see [SA1618 below](#) – Retail pharmacy

Tab 30 mg 42.06 28 ✓ **Cinacalcet Devatis**

a) Brand switch fee payable (Pharmacode 2634120) - see [page 241](#) for details

b) Wastage claimable

Tab 60 mg – Wastage claimable 84.12 28 ✓ **Cinacalcet Devatis**

► **SA1618** Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial – Special Authority see [SA2109 below](#) –

Retail pharmacy 18.00 1 ✓ **Zoledronic acid
Mylan**

► **SA2109** Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting

continued...

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia .

Note: Indications marked with * are unapproved indications.

Corticosteroids and Related Agents for Systemic Use

BETAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE

* Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	19.20 (36.96)	5	Celestone Chronodose
---	------------------	---	-------------------------

DEXAMETHASONE

* Tab 0.5 mg – Up to 60 tab available on a PSO	1.50	30	✓ <u>Dexamethsone</u>
* Tab 4 mg – Up to 30 tab available on a PSO	2.65	30	✓ <u>Dexamethsone</u>
Oral liq 1 mg per ml	48.15	25 ml OP	✓ <u>Biomed</u>

DEXAMETHASONE PHOSPHATE

Dexamethasone phosphate injection will not be funded for oral use.

* Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	9.25	10	✓ <u>Dexamethasone Phosphate Panpharma</u>
* Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	16.37	10	✓ <u>Dexamethasone Phosphate Panpharma</u>

FLUDROCORTISONE ACETATE

* Tab 100 mcg	14.32	100	✓ <u>Florinef</u>
---------------------	-------	-----	-------------------

HYDROCORTISONE

* Tab 5 mg	8.10	100	✓ <u>Douglas</u>
* Tab 20 mg	20.32	100	✓ <u>Douglas</u>
* Inj 100 mg vial	4.38	1	✓ <u>Solu-Cortef</u>
a) Up to 5 inj available on a PSO			
b) Only on a PSO			

METHYLPREDNISOLONE

* Tab 4 mg	112.00	100	✓ <u>Medrol</u>
* Tab 100 mg	223.10	20	✓ <u>Medrol</u>

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
METHYLPREDNISOLONE (AS SODIUM SUCCINATE)				
Inj 40 mg vial	22.30	1	✓	Solu-Medrol-Act-O-Vial
Inj 125 mg vial	34.10	1	✓	Solu-Medrol-Act-O-Vial
Inj 500 mg vial	26.88	1	✓	Solu-Medrol-Act-O-Vial
Inj 1 g vial	32.84	1	✓	Solu-Medrol
METHYLPREDNISOLONE ACETATE				
Inj 40 mg per ml, 1 ml vial	47.06	5	✓	Depo-Medrol
PREDNISOLONE				
* Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓	Redipred
PREDNISONE				
* Tab 1 mg	18.58	500	✓	Apo-Prednisone
			✓	Prednisone Clinect
* Tab 2.5 mg	21.04	500	✓	Apo-Prednisone
			✓	Prednisone Clinect
* Tab 5 mg – Up to 30 tab available on a PSO	19.30	500	✓	Apo-Prednisone
			✓	Prednisone Clinect
* Tab 20 mg – Up to 30 tab available on a PSO	50.51	500	✓	Apo-Prednisone
			✓	Prednisone Clinect
<i>(Apo-Prednisone Tab 1 mg to be delisted 1 November 2022)</i>				
<i>(Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022)</i>				
<i>(Apo-Prednisone Tab 5 mg to be delisted 1 November 2022)</i>				
<i>(Apo-Prednisone Tab 20 mg to be delisted 1 November 2022)</i>				
TETRACOSACTRIN				
* Inj 250 mcg per ml, 1 ml ampoule	75.00	1	✓	UK Synacthen ^{\$29}
			✓	AU Synacthen
			✓	Synacthen
* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓	Synacthen Depot
			✓	Synacthene
				Retard ^{\$29}
TRIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule	51.10	5	✓	Kenacort-A 40

Sex Hormones Non Contraceptive

Androgen Agonists and Antagonists

CYPROTERONE ACETATE				
Tab 50 mg	14.37	50	✓	Siterone
Tab 100 mg	28.03	50	✓	Siterone
TESTOSTERONE				
Patch 5 mg per day	90.00	30	✓	Androderm
TESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	✓	Depo-Testosterone

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
TESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12.98	1	✓	Sustanon Ampoules
TESTOSTERONE UNDECANOATE				
Cap 40 mg – Subsidy by endorsement	21.00	60	✓	Andriol Testocaps
Subsidy by endorsement – subsidised for patients who were taking testosterone undecanoate cap 40mg prior to 1 November 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of testosterone undecanoate cap 40 mg in the preceding 12 months.				
Inj 250 mg per ml, 4 ml vial.....	86.00	1	✓	Reandron 1000

Hormone Replacement Therapy - Systemic

Prescribing Guideline

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG “Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004”.

Oestrogens

OESTRADIOL – See prescribing guideline [above](#)

* Tab 1 mg	4.12 (11.10)	28 OP		
* Tab 2 mg	4.12 (11.10)	28 OP		Estrofem
Patch 25 mcg per day.....	6.12	8	✓	Estradot
a) No more than 2 patch per week				
b) Only on a prescription				
Patch 50 mcg per day.....	7.04	8	✓	Estradot 50 mcg
a) No more than 2 patch per week				
b) Only on a prescription				
Patch 75 mcg per day.....	7.91	8	✓	Estradot
a) No more than 2 patch per week				
b) Only on a prescription				
Patch 100 mcg per day.....	7.91	8	✓	Estradot
a) No more than 2 patch per week				
b) Only on a prescription				

OESTRADIOL VALERATE – See prescribing guideline [above](#)

* Tab 1 mg	12.36	84	✓	Progynova
* Tab 2 mg	12.36	84	✓	Progynova
OESTROGENS – See prescribing guideline above				
* Conjugated, equine tab 300 mcg.....	3.01 (17.50)	28		Premarin
* Conjugated, equine tab 625 mcg.....	4.12 (17.50)	28		Premarin

Progestogens

MEDROXYPROGESTERONE ACETATE – See prescribing guideline [above](#)

* Tab 2.5 mg	4.69	30	✓	Provera
* Tab 5 mg	17.50	100	✓	Provera
* Tab 10 mg	8.94	30	✓	Provera

▲Three months supply may be dispensed at one time if endorsed “certified exemption” by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Progestogen and Oestrogen Combined Preparations				
OESTRADIOL WITH NORETHISTERONE – See prescribing guideline on the previous page				
* Tab 1 mg with 0.5 mg norethisterone acetate	5.40 (18.10)	28 OP		Kliovance
* Tab 2 mg with 1 mg norethisterone acetate	5.40 (18.10)	28 OP		Kliogest
* Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg oestradiol tab (12) and 1 mg oestradiol tab (6).....	5.40 (18.10)	28 OP		Trisequens

Other Oestrogen Preparations

ETHINYLOESTRADIOL – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking ethinyloestradiol prior to 1 March 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ethinyloestradiol.

Tab 10 mcg.....	17.60	100	✓ NZ Medical and Scientific
-----------------	-------	-----	-----------------------------

(NZ Medical and Scientific Tab 10 mcg to be delisted 1 February 2023)

OESTRIOL

* Tab 2 mg	7.00	30	✓ <u>Ovestin</u>
------------------	------	----	------------------

Other Progestogen Preparations

LEVONORGESTREL

* Intra-uterine device 52 mg.....	269.50	1	✓ <u>Mirena</u>
* Intra-uterine device 13.5 mg.....	215.60	1	✓ <u>Jaydess</u>

MEDROXYPROGESTERONE ACETATE

Tab 100 mg	116.15	100	✓ <u>Provera HD</u>
------------------	--------	-----	---------------------

NORETHISTERONE

* Tab 5 mg – Up to 30 tab available on a PSO	5.49	30	✓ <u>Primolut N</u>
--	------	----	---------------------

PROGESTERONE

Cap 100 mg – Special Authority see SA1609 below – Retail pharmacy.....	16.50	30	✓ <u>Utrogestan</u>
---	-------	----	---------------------

►SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Either:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and

continued...

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

3 Either:

- 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
- 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

CARBIMAZOLE

* Tab 5 mg	7.56	100	✓ Neo-Mercazole
------------------	------	-----	-----------------

Neo-Mercazole to be Principal Supply on 1 September 2022

LEVOTHYROXINE

* Tab 25 mcg.....	5.55	90	✓ Synthroid
* Tab 50 mcg.....	1.71	28	✓ Mercury Pharma
	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tab 100 mcg.....	1.78	28	✓ Mercury Pharma
	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin

PROPYLTHIOURACIL – Special Authority see [SA1199 below](#) – Retail pharmacy

Propylthiouracil is not recommended for patients under the age of 18 years unless the patient is pregnant and other treatments are contraindicated.

Tab 50 mg	35.00	100	✓ PTU <small>\$29</small>
-----------------	-------	-----	---------------------------

► **SA1199** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SOMATROPIN (OMNITROPE) – Special Authority see [SA2032 below](#) – Retail pharmacy

* Inj 5 mg cartridge.....	69.75	1	✓ Omnitrope
* Inj 10 mg cartridge.....	69.75	1	✓ Omnitrope
* Inj 15 mg cartridge.....	139.50	1	✓ Omnitrope

► **SA2032** Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist.

Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

2 All of the following:

- 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
- 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
- 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
- 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and

5 Either:

5.1 Both:

5.1.1 The patient is aged two years or older; and

5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or

5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:

continued...

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.1 The patient has been treated with somatropin for < 12 months; and
- 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
- 1.3 Serum IGF-I levels have been increased within $\pm 1SD$ of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within $\pm 1SD$ of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

GOSERELIN

Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamber syringe – Higher subsidy of \$221.60 per 1 inj with Endorsement	66.48 (221.60)	1	Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe – Higher subsidy of \$591.68 per 1 inj with Endorsement	177.50 (591.68)	1	Lucrin Depot 3-month

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

HORMONE PREPARATIONS - SYSTEMIC EXCLUDING CONTRACEPTIVE HORMONES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Vasopressin Agonists				
DESMOPRESSIN				
Wafer 120 mcg	47.00	30	✓	Minirin Melt
DESMOPRESSIN ACETATE				
Tab 100 mcg.....	25.00	30	✓	Minirin
Tab 200 mcg.....	54.45	30	✓	Minirin
▲ Nasal spray 10 mcg per dose.....	27.95	6 ml OP	✓	Desmopressin- PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	✓	Minirin
Other Endocrine Agents				
CABERGOLINE				
Tab 0.5 mg – Maximum of 2 tab per prescription; can be waived by Special Authority see SA2070 below.....	3.75 15.20	2 8	✓ ✓	Dostinex Dostinex
»SA2070 Special Authority for Waiver of Rule				
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Any of the following:				
1 Hyperprolactinemia; or				
2 Acromegaly*; or				
3 Inhibition of lactation.				
Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.				
Note: Indication marked with * is an unapproved indication.				
CLOMIFENE CITRATE				
Tab 50 mg	29.84	10	✓	Mylan Clomiphen ^{\$29}
METYRAPONE				
Cap 250 mg.....	558.00	50	✓	Metopirone

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

Anthelmintics

ALBENDAZOLE – Special Authority see [SA1318 below](#) – Retail pharmacy

Tab 400 mg	469.20	60	✓ Eskazole ^{S29}
------------------	--------	----	----------------------------------

►SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE – Only on a prescription

Tab 100 mg	7.97	6	✓ Vermox
Oral liq 100 mg per 5 ml	2.18	15 ml	Vermox
	(7.53)		

PRAZIQUANTEL

Tab 600 mg	68.00	8	✓ Biltricide
------------------	-------	---	---------------------

Antibacterials

- a) For topical antibacterials, refer to DERMATOLOGICALS, [page 61](#)
b) For anti-infective eye preparations, refer to SENSORY ORGANS, [page 236](#)

Cephalosporins and Cephamycins

CEFACLOR MONOHYDRATE

Cap 250 mg	24.70	100	✓ Ranbaxy-Cefaclor ✓ Ranbaxy-Cefaclor ^{S29} ^{S29}
Grans for oral liq 125 mg per 5 ml – Wastage claimable.....	3.53	100 ml	✓ Ranbaxy-Cefaclor ✓ Ranbaxy-Cefaclor ^{S29} ^{S29}

CEFALEXIN

Cap 250 mg	3.33	20	✓ Cephalexin ABM
Cap 500 mg	3.95	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml – Wastage claimable.....	8.75	100 ml	✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml – Wastage claimable.....	11.75	100 ml	✓ Cefalexin Sandoz

CEFAZOLIN – Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a DHB approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	3.39	5	✓ AFT
Inj 1 g vial	3.49	5	✓ AFT

CEFTRIAXONE – Subsidy by endorsement

- a) Up to 10 inj available on a PSO
b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly.

Inj 500 mg vial	0.89	1	✓ Ceftriaxone-AFT
Inj 1 g vial	3.99	5	✓ Ceftriaxone-AFT

CEFUROXIME AXETIL – Subsidy by endorsement

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly.

Tab 250 mg	45.93	50	✓ Zinnat
------------------	-------	----	-----------------

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see [SA1683 below](#)

A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓	Apo-Azithromycin
Tab 500 mg – Up to 8 tab available on a PSO	2.57	2	✓	Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) – Wastage claimable	16.97	15 ml	✓	Zithromax

►SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with *Pseudomonas aeruginosa* or *Pseudomonas*-related gram negative organisms*; or
- 4 Patient has an atypical *Mycobacterium* infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN – Maximum of 500 mg per prescription; can be waived by Special Authority see [SA1857 below](#)

Tab 250 mg	8.53	14	✓	Klacid
Grans for oral liq 250 mg per 5 ml – Wastage claimable.....	192.00	50 ml	✓	Klacid

►SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
continued...			
1 Atypical mycobacterial infection; or			
2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.			
Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:			
Both:			
1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and			
2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.			
Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated.			
Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.			
ERYTHROMYCIN (AS LACTOBIONATE)			
Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	✓ E-Mycin
a) Up to 300 ml available on a PSO			
b) Up to 2 x the maximum PSO quantity for RFPP			
c) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	✓ E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 500 mg	29.90 (44.58)	100	ERA
<i>(ERA Tab 500 mg to be delisted 1 September 2022)</i>			
ROXITHROMYCIN			
Tab disp 50 mg.....	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow-Roxithromycin
Tab 300 mg	16.33	50	✓ Arrow-Roxithromycin
<i>(Rulide D Tab disp 50 mg to be delisted 1 September 2022)</i>			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg.....	22.50	500	✓	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg.....	36.98	500	✓	Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml.....	1.40	100 ml	✓	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml.....	1.73	100 ml	✓	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
Inj 250 mg vial	15.97	10	✓	Ibiamox
Inj 500 mg vial	17.43	10	✓	Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO	21.64	10	✓	Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID				
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab available on a PSO	0.89	10	✓	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml.....	5.00	100 ml	✓	Augmentin
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO	2.20	100 ml OP	✓	Curam
BENZATHINE BENZYL PENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj available on a PSO	375.97	10	✓	Bicillin LA
BENZYL PENICILLIN SODIUM [PENICILLIN G]				
Inj 600 mg (1 million units) vial – Up to 5 inj available on a PSO	11.09	10	✓	Sandoz
FLUCLOXACILLIN				
Cap 250 mg – Up to 30 cap available on a PSO.....	15.79	250	✓	Flucloxacillin-AFT
Cap 500 mg – Up to 30 cap available on a PSO.....	52.99	500	✓	Flucloxacillin-AFT
Grans for oral liq 25 mg per ml.....	3.29	100 ml	✓	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 50 mg per ml.....	3.68	100 ml	✓	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial	17.56	10	✓	Flucloxin
Inj 500 mg vial	18.87	10	✓	Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	✓	Flucil

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg – Up to 30 cap available on a PSO.....	3.84	50	✓	Cilicaine VK
Cap 500 mg.....	6.86	50	✓	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml.....	2.99	100 ml	✓	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml.....	3.99	100 ml	✓	AFT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe – Up to 5 inj available on a PSO.....	123.50	5	✓	Cilicaine

Tetracyclines

DOXYCYCLINE				
* Tab 100 mg – Up to 30 tab available on a PSO.....	64.43	500	✓	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg – Additional subsidy by Special Authority see SA1355 below – Retail pharmacy	5.79 (12.05)	60		Mino-tabs
* Cap 100 mg.....	19.32 (52.04)	100		Minomycin

►SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE – Special Authority see SA1332 below – Retail pharmacy

Tab 250 mg	21.42	28	✓	Accord ^{S29}
------------------	-------	----	---	------------------------------

►SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg – Up to 5 tab available on a PSO.....	2.42	28	✓	Ciptflox
Tab 500 mg – Up to 5 tab available on a PSO.....	3.40	28	✓	Ciptflox
Tab 750 mg	5.95	28	✓	Ciptflox

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
CLINDAMYCIN				
Cap hydrochloride 150 mg	4.61	24	✓	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓	Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – Subsidy by endorsement				
Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly.				
Inj 150 mg	65.00	1	✓	Colistin-Link
GENTAMICIN SULPHATE				
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement.....	95.00	5	✓	DBL Gentamicin
Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.				
Inj 10 mg per ml, 2 ml ampoule – Subsidy by endorsement.....	91.00	5	✓	Wockhardt <small>\$29</small>
	182.00	10	✓	Teligent <small>\$29</small>
Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.				
Inj 40 mg per ml, 2 ml ampoule – Subsidy by endorsement.....	17.50	10	✓	Pfizer
	87.50	50	✓	Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient or complicated urinary tract infection and the prescription is endorsed accordingly.				
MOXIFLOXACIN – Special Authority see SA1740 below – Retail pharmacy				
No patient co-payment payable				
Tab 400 mg	42.00	5	✓	Avelox

➔SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

1 Both:

1.1 Active tuberculosis*; and

1.2 Any of the following:

1.2.1 Documented resistance to one or more first-line medications; or

1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or

1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or

1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or

1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or

2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or

3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and

2 Either:

2.1 Has tried and failed to clear infection using azithromycin; or

2.2 Has laboratory confirmed azithromycin resistance; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
continued...				
3 Treatment is only for 7 days.				
Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eye injury and treatment is for 5 days only.				
Note: Indications marked with * are unapproved indications.				
PAROMOMYCIN – Special Authority see SA1689 below – Retail pharmacy				
Cap 250 mg	126.00	16	✓	Humatin ^{S29}
➔ SA1689 Special Authority for Subsidy				
Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:				
Either:				
1 Patient has confirmed cryptosporidium infection; or				
2 For the eradication of Entamoeba histolytica carriage.				
Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria:				
Either:				
1 Patient has confirmed cryptosporidium infection; or				
2 For the eradication of Entamoeba histolytica carriage.				
PYRIMETHAMINE – Special Authority see SA1328 below – Retail pharmacy				
Tab 25 mg	48.00	30	✓	Daraprim ^{S29}
➔ SA1328 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Any of the following:				
1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or				
2 For pregnant patients for the term of the pregnancy; or				
3 For infants with congenital toxoplasmosis until 12 months of age.				
SODIUM FUSIDATE [FUSIDIC ACID]				
Tab 250 mg	67.85	36	✓	Fucidin
SULFADIAZINE SODIUM – Special Authority see SA1331 below – Retail pharmacy				
Tab 500 mg	543.20	56	✓	Wockhardt ^{S29}
➔ SA1331 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Any of the following:				
1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or				
2 For pregnant patients for the term of the pregnancy; or				
3 For infants with congenital toxoplasmosis until 12 months of age.				
TOBRAMYCIN				
Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement	18.50	5	✓	Tobramycin Mylan
Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly.				
Solution for inhalation 60 mg per ml, 5 ml – Subsidy by endorsement.....	395.00	56 dose	✓	Tobramycin BNM
a) Wastage claimable				
b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly.				
TRIMETHOPRIM				
* Tab 300 mg – Up to 30 tab available on a PSO	18.55	50	✓	TMP

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOXAZOLE]				
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – Up to 30 tab available on a PSO.....	64.80	500	✓	Trisul
* Oral liq 8 mg sulphamethoxazole 40 mg per ml – Up to 200 ml available on a PSO.....	2.97	100 ml	✓	Deprim
VANCOMYCIN – Subsidy by endorsement				
Only if prescribed for a dialysis or cystic fibrosis patient or for prophylaxis of endocarditis or for treatment of Clostridium difficile following metronidazole failure and the prescription is endorsed accordingly.				
Inj 500 mg vial	2.35	1	✓	Mylan

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, [page 62](#)
b) For topical antifungals refer to GENITO URINARY, [page 75](#)

FLUCONAZOLE				
Cap 50 mg	2.75	28	✓	Mylan
Cap 150 mg	0.65	1	✓	Mylan
Cap 200 mg	12.89	28	✓	Mylan
Powder for oral suspension 10 mg per ml – Special Authority				
see SA1359 below – Retail pharmacy.....	109.34	35 ml	✓	Diflucan
Wastage claimable				

►SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE				
Cap 100 mg	4.27	15	✓	Itrazole
Oral liq 10 mg per ml – Special Authority see SA1322 on the next page – Retail pharmacy.....	141.80	150 ml OP	✓	Sporanox

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

►SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg – PCT	CBS	30	✓ Link Healthcare S29
		100	✓ Nizoral S29
			✓ Strides Shasun S29

NYSTATIN

Tab 500,000 u	14.16 (17.09)	50	Nilstat
Cap 500,000 u	12.81 (15.47)	50	Nilstat

POSACONAZOLE – Special Authority see SA1285 below – Retail pharmacy

Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml	761.13	105 ml OP	✓ Noxafil

►SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERBINAFINE

* Tab 250 mg	8.15	84	✓ Deolate
--------------------	------	----	-----------

VORICONAZOLE – Special Authority see SA1273 on the next page – Retail pharmacy

Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml – Wastage claimable	1,523.22	70 ml	✓ Vfend

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

►SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as *Fusarium* spp. and *Scedosporium* spp.

Antimalarials

PRIMAQUINE – Special Authority see [SA1684 below](#) – Retail pharmacy

Tab 15 mg	400.00	100	✓ Sanofi Primaquine ^{S29}
-----------------	--------	-----	---------------------------------------

►SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE

Tab 200 mg – Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg – Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl

ORNIDAZOLE

Tab 500 mg	36.16	10	✓ Arrow-Ornidazole
------------------	-------	----	--------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

Antituberculosics and Antileptotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculosics and Antileptotics group regardless of immigration status.

CLOFAZIMINE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

* Cap 50 mg 442.00 100 ✓ **Lamprene** \$29

CYCLOSERINE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.

Cap 250 mg 344.00 60 ✓ **Cyclorin** \$29

DAPSONE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Tab 25 mg 268.50 100 ✓ **Dapsone**

Tab 100 mg 329.50 100 ✓ **Dapsone**

ETHAMBUTOL HYDROCHLORIDE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician

Tab 100 mg 85.73 100 ✓ **EMB Fatol** \$29

Tab 400 mg 49.34 56 ✓ **Myambutol** \$29

ISONIAZID – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician

* Tab 100 mg 23.00 100 ✓ **PSM**

ISONIAZID WITH RIFAMPICIN – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician

* Tab 100 mg with rifampicin 150 mg 89.82 100 ✓ **Rifinah**

* Tab 150 mg with rifampicin 300 mg 179.13 100 ✓ **Rifinah**

PARA-AMINO SALICYLIC ACID – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

Grans for oral liq 4 g sachet 280.00 30 ✓ **Paser** \$29

PROTONAMIDE – Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

Tab 250 mg 305.00 100 ✓ **Peteha** \$29

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PYRAZINAMIDE – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician				
* Tab 500 mg	64.95	100	✓	AFT-Pyrazinamide
RIFABUTIN – Retail pharmacy-Specialist				
a) No patient co-payment payable				
b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, respiratory physician or gastroenterologist				
* Cap 150 mg	299.75	30	✓	Mycobutin
RIFAMPICIN – Subsidy by endorsement				
a) No patient co-payment payable				
b) For confirmed recurrent <i>Staphylococcus aureus</i> infection in combination with other effective anti-staphylococcal antimicrobial based on susceptibilities and the prescription is endorsed accordingly; can be waived by endorsement - Retail pharmacy - Specialist. Specialist must be an internal medicine physician, clinical microbiologist, dermatologist, paediatrician, or public health physician.				
* Cap 150 mg	58.54	100	✓	Rifadin
* Cap 300 mg	122.06	100	✓	Rifadin
* Oral liq 100 mg per 5 ml	12.60	60 ml	✓	Rifadin

Antivirals

For eye preparations refer to Eye Preparations, Anti-Infective Preparations, [page 236](#)

Hepatitis B Treatment

ENTECAVIR				
* Tab 0.5 mg	52.00	30	✓	Entecavir Sandoz
LAMIVUDINE – Special Authority see SA1685 below – Retail pharmacy				
Tab 100 mg	6.95	28	✓	Zetlam
Oral liq 5 mg per ml	270.00	240 ml OP	✓	Zeffix

➔SA1685 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 1 year where used for the treatment or prevention of hepatitis B.

Renewal from any relevant practitioner. Approvals valid for 2 years where used for the treatment or prevention of hepatitis B.

TENOFOVIR DISOPROXIL

Tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals for the purposes of Special Authority SA1651., [page 104](#)

* Tab 245 mg (300.6 mg as a succinate)	38.10	30	✓	Tenofovir Disoproxil Teva
--	-------	----	---	----------------------------------

Herpesvirus Treatments

ACICLOVIR				
* Tab dispersible 200 mg	1.60	25	✓	Lovir
* Tab dispersible 400 mg	5.38	56	✓	Lovir
* Tab dispersible 800 mg	5.98	35	✓	Lovir
VALACICLOVIR				
Tab 500 mg	6.50	30	✓	Vaclovir
Tab 1,000 mg	13.76	30	✓	Vaclovir

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
VALGANCICLOVIR – Special Authority see SA1993 below – Retail pharmacy				
Tab 450 mg	132.00	60	✓	Valganciclovir Mylan

► **SA1993** Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
- 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR – [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website <https://pharmac.govt.nz/maviret>

Tab 100 mg with pibrentasvir 40 mg	24,750.00	84 OP	✓ Maviret
--	-----------	-------	-----------

LEDIPASVIR WITH SOFOSBUVIR – [Xpharm] – Special Authority see [SA1605 below](#)

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg.....	24,363.46	28	✓ Harvoni
---------------------------------------	-----------	----	-----------

► [SA1605](#) Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website <http://www.pharmac.govt.nz/maviret> or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: heccpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL – Subsidy by endorsement; can be waived by Special Authority see [SA1994 below](#)

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA1651 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA1651, [page 104](#) There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate)	61.15	30	✓ Teva
---	-------	----	--------

► [SA1994](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and
- 2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and
- 3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 3 months and is not contraindicated for treatment; and
- 4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and

6 Either:

6.1 All of the following:

6.1.1 Patient is male or transgender; and

6.1.2 Patient has sex with men; and

6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and

6.1.4 Any of the following:

6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or

6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or

6.1.4.3 Patient has used methamphetamine in the last three months; or

6.2 All of the following:

6.2.1 Patient has a regular partner who has HIV infection; and

6.2.2 Partner is either not on treatment or has a detectable viral load; and

6.2.3 Condoms have not been consistently used.

Renewal from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Applicant has an up to date knowledge of the safety issues and is competent to prescribe pre-exposure prophylaxis (refer to local health pathways or <https://ashm.org.au/HIV/PrEP/> for training materials); and

2 Patient has undergone testing for HIV, syphilis and Hep B if not immune and a full STI screen in the previous two weeks; and

3 Patient has had renal function testing (creatinine, phosphate and urine protein/creatinine ratio) within the last 12 months and is not contraindicated for treatment; and

4 Patient has received advice regarding the reduction of risk of HIV and sexually transmitted infections and how to reduce those risks; and

5 Patient has tested HIV negative and is not at risk of HIV seroconversion; and

6 Either:

6.1 All of the following:

6.1.1 Patient is male or transgender; and

6.1.2 Patient has sex with men; and

6.1.3 Patient is likely to have multiple episodes of condomless anal intercourse in the next 3 months; and

6.1.4 Any of the following:

6.1.4.1 Patient has had at least one episode of condomless receptive anal intercourse with one or more casual male partners in the last 3 months; or

6.1.4.2 A diagnosis of rectal chlamydia, rectal gonorrhoea, or infectious syphilis within the last 3 months; or

6.1.4.3 Patient has used methamphetamine in the last three months; or

6.2 All of the following:

6.2.1 Patient has a regular partner who has HIV infection; and

6.2.2 Partner is either not on treatment or has a detectable viral load; and

6.2.3 Condoms have not been consistently used.

COVID-19 Treatments

MOLNUPIRAVIR – [Xpharm] – Subsidy by endorsement

a) No patient co-payment payable

b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on [Pharmac's website](#)) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Cap 200 mg 0.00 40 ✓ **Lagevrio**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
NIRMATRELVIR WITH RITONAVIR – [Xpharm] – Subsidy by endorsement				
a) No patient co-payment payable				
b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on Pharmac's website) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.				
Tab 150 mg with ritonavir 100 mg	0.00	30	✓	Paxlovid

Antiretrovirals

►SA1651 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following non-occupational exposure to HIV) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (second or subsequent post-exposure prophylaxis) only from a named specialist. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 2.1 Patient has had unprotected receptive anal intercourse with a known HIV positive person; or
- 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
- 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA1651 on the previous page – Retail pharmacy			
Tab 200 mg	190.15	90	✓ Stocrin
Tab 600 mg	63.38	30	✓ Stocrin
ETRAVIRINE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Tab 200 mg	770.00	60	✓ Intencele
NEVIRAPINE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Tab 200 mg	84.00	60	✓ Nevirapine
			Alphapharm
Oral suspension 10 mg per ml.....	203.55	240 ml	✓ Viramune Suspension

Nucleosides Reverse Transcriptase Inhibitors

ABACAVIR SULPHATE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Tab 300 mg	180.00	60	✓ Ziagen
Oral liq 20 mg per ml	256.31	240 ml OP	✓ Ziagen
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Note: abacavir with lamivudine (combination tablets) counts as two anti-retroviral medications for the purposes of the anti-retroviral Special Authority.			
Tab 600 mg with lamivudine 300 mg.....	63.00	30	✓ Kivexa
EFAVIRENZ WITH ETRICITABINE AND TENOFOVIR DISOPROXIL – Special Authority see SA1651 on the previous page – Retail pharmacy			
Note: Efavirenz with emtricitabine and tenofovir disoproxil counts as three anti-retroviral medications for the purposes of the anti-retroviral Special Authority			
Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil 245 mg (300 mg as a maleate).....	106.88	30	✓ Mylan
EMTRICITABINE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Cap 200 mg	307.20	30	✓ Emtriva
LAMIVUDINE – Special Authority see SA1651 on the previous page – Retail pharmacy			
Tab 150 mg	84.50	60	✓ Lamivudine
			Alphapharm
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC
ZIDOVUDINE [AZT] – Special Authority see SA1651 on the previous page – Retail pharmacy			
Cap 100 mg	152.25	100	✓ Retrovir
Oral liq 10 mg per ml	30.45	200 ml OP	✓ Retrovir

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority see SA1651 on page 104 – Retail pharmacy				
Note: zidovudine [AZT] with lamivudine (combination tablets) counts as two anti-retroviral medications for the purposes of the anti-retroviral Special Authority.				
Tab 300 mg with lamivudine 150 mg.....	33.00	60	✓	Alphapharm

Protease Inhibitors

ATAZANAVIR SULPHATE – Special Authority see SA1651 on page 104 – Retail pharmacy				
Cap 150 mg.....	141.68	60	✓	Teva
Cap 200 mg.....	188.91	60	✓	Teva
DARUNAVIR – Special Authority see SA1651 on page 104 – Retail pharmacy				
Tab 400 mg.....	132.00	60	✓	Darunavir Mylan
Tab 600 mg.....	196.65	60	✓	Darunavir Mylan
LOPINAVIR WITH RITONAVIR – Special Authority see SA1651 on page 104 – Retail pharmacy				
Tab 100 mg with ritonavir 25 mg.....	150.00	60	✓	Lopinavir/Ritonavir Mylan
Tab 200 mg with ritonavir 50 mg.....	295.00	120	✓	Lopinavir/Ritonavir Mylan
Oral liq 80 mg with ritonavir 20 mg per ml.....	735.00	300 ml OP	✓	Kaletra
RITONAVIR – Special Authority see SA1651 on page 104 – Retail pharmacy				
Tab 100 mg.....	43.31	30	✓	Norvir

Strand Transfer Inhibitors

DOLUTEGRAVIR – Special Authority see SA1651 on page 104 – Retail pharmacy				
Tab 50 mg.....	1,090.00	30	✓	Tivicay
RALTEGRAVIR POTASSIUM – Special Authority see SA1651 on page 104 – Retail pharmacy				
Tab 400 mg.....	1,090.00	60	✓	Isentress
Tab 600 mg.....	1,090.00	60	✓	Isentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

1) Diagnosis

- Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
- PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
- Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- 1) Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2) Pregnancy.
- 3) Neutropenia ($< 2.0 \times 10^9$) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A – Special Authority see [SA2034 below](#) – Retail pharmacy

- a) See prescribing guideline [on the previous page](#)
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

Inj 180 mcg prefilled syringe.....500.00 4 ✓ **Pegasys**

►SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 3.2 Patient was a partial responder; or
- 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naïve; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

continued...

3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or

3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE

* Tab 1 g	40.01	100	✓ Hiprex
-----------------	-------	-----	----------

NITROFURANTOIN

* Tab 50 mg – Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
* Tab 100 mg	37.50	100	✓ Nifuran
* Cap modified-release 100 mg – Wastage claimable	86.40	100	✓ <u>Macrobid</u>

NORFLOXACIN

Tab 400 mg – Subsidy by endorsement.....	245.00	100	✓ Arrow-Norfloxacin
--	--------	-----	---------------------

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

MUSCULOSKELETAL SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Anticholinesterases

NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	✓	<u>Max Health</u>
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.79	100	✓	<u>Mestinon</u>

Non-Steroidal Anti-Inflammatory Drugs

DICLOFENAC SODIUM				
* Tab EC 25 mg	1.99	50	✓	<u>Diclofenac Sandoz</u>
* Tab 50 mg dispersible	1.50	20	✓	<u>Voltaren D</u>
* Tab EC 50 mg	1.99	50	✓	<u>Diclofenac Sandoz</u>
* Tab long-acting 75 mg	19.60	100	✓	<u>Voltaren SR</u>
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a PSO	13.20	5	✓	<u>Voltaren</u>
* Suppos 12.5 mg	2.04	10	✓	<u>Voltaren</u>
* Suppos 25 mg	2.44	10	✓	<u>Voltaren</u>
* Suppos 50 mg – Up to 10 supp available on a PSO	4.22	10	✓	<u>Voltaren</u>
* Suppos 100 mg	7.00	10	✓	<u>Voltaren</u>
IBUPROFEN				
* Tab 200 mg	21.40	1,000	✓	<u>Relieve</u>
* Tab long-acting 800 mg	3.05	30	✓	<u>Brufen SR</u>
* Oral liq 20 mg per ml	2.25	200 ml	✓	<u>Ethics</u>
	11.29		✓	<u>Fenpaed 100 mg per 5 ml</u>
KETOPROFEN				
* Cap long-acting 200 mg	12.07	28	✓	<u>Oruvail SR</u>
MEFENAMIC ACID				
* Cap 250 mg	1.25	50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
NAPROXEN				
* Tab 250 mg	32.69	500	✓	<u>Noflam 250</u>
* Tab 500 mg	28.71	250	✓	<u>Noflam 500</u>
* Tab long-acting 750 mg	6.47	28	✓	<u>Naprosyn SR 750</u>
* Tab long-acting 1 g	8.62	28	✓	<u>Naprosyn SR 1000</u>
TENOXICAM				
* Tab 20 mg	9.15	100	✓	<u>Tilcotil</u>
* Inj 20 mg vial	9.95	1	✓	<u>AFT</u>

NSAIDs Other

CELECOXIB				
Cap 100 mg	3.45	60	✓	<u>Celecoxib Pfizer</u>
Cap 200 mg	2.30	30	✓	<u>Celebrex</u>
	3.20		✓	<u>Celecoxib Pfizer</u>

(Celebrex Cap 200 mg to be delisted 1 November 2022)

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% – Special Authority see [SA1289 below](#) – Retail

pharmacy..... 9.75 45 g OP ✓ **Zostrix**

►SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE – Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly.

Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg 8.78 100 ✓ **Plaquenil**

LEFLUNOMIDE

Tab 10 mg 6.00 30 ✓ **Arava**

Tab 20 mg 6.00 30 ✓ **Arava**

PENICILLAMINE

Tab 125 mg 67.23 100 ✓ **D-Penamine**

Tab 250 mg 110.12 100 ✓ **D-Penamine**

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM

* Tab 70 mg 2.44 4 ✓ **Fosamax**

ALENDRONATE SODIUM WITH COLECALCIFEROL

* Tab 70 mg with colecalciferol 5,600 iu 1.51 4 ✓ **Fosamax Plus**

Other Treatments

DENOSUMAB – Special Authority see [SA1777 below](#) – Retail pharmacy

Inj 60 mg prefilled syringe..... 326.00 1 ✓ **Prolia**

►SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Either:
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
- 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

Inj 3 mg per ml, 10 ml vial.....	27.53	1	✓ Pamisol
Inj 6 mg per ml, 10 ml vial.....	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial.....	79.95	1	✓ Pamisol

RALOXIFENE HYDROCHLORIDE – Special Authority see [SA1779 below](#) – Retail pharmacy

* Tab 60 mg	53.76	28	✓ Evista
-------------------	-------	----	----------

►SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- History of two significant osteoporotic fractures demonstrated radiologically; or
- Documented T-Score less than or equal to -3.0 (see Notes); or
- A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- Patient has had a Special Authority approval for zoledronic acid (Underlying cause – Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

RISEDRONATE SODIUM

Tab 35 mg 3.10 4 ✓ **Risedronate Sandoz**

TERIPARATIDE – Special Authority see [SA1139 below](#) – Retail pharmacy

Inj 250 mcg per ml, 2.4 ml 490.00 1 ✓ **Forteo**

► [SA1139](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- The patient has severe, established osteoporosis; and
- The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- The patient has had two or more fractures due to minimal trauma; and
- The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- A maximum of 18 months of treatment (18 cartridges) will be subsidised.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

ZOLEDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial – Special Authority see

SA2110 below – Retail pharmacy 60.00 100 ml OP ✓ **Aclasta**

►SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -1.5) (see Note); or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause - glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Any of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and

2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause - Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL

* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg	28.57	500	✓ DP-Allopurinol

BENZBROMARONE – Special Authority see [SA1963 below](#) – Retail pharmacy

Tab 50 mg	22.50	100	✓ Narcaricin mite ^{\$29}
Tab 100 mg	13.50	30	✓ Desuric ^{\$29}
	45.00	100	✓ Urinorm ^{\$29}
			✓ Benzbromaron AL
			100 ^{\$29}

➔ [SA1963](#) Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	6.00	100	✓ Colgout
Colgout to be Principal Supply on 1 September 2022			

FEBUXOSTAT – Special Authority see [SA2054 below](#) – Retail pharmacy

Tab 80 mg	20.00	28	✓ Febuxostat multichem
Tab 120 mg	20.00	28	✓ Febuxostat multichem

➔ [SA2054](#) Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout.

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

* Tab 500 mg	66.95	100	✓ Probenecid-AFT
--------------------	-------	-----	------------------

Muscle Relaxants

BACLOFEN

* Tab 10 mg	4.20	100	✓ Pacifen
Inj 0.05 mg per ml, 1 ml ampoule – Subsidy by endorsement.....	11.55	1	✓ Lioresal Intrathecal
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.			
Inj 2 mg per ml, 5 ml ampoule – Subsidy by endorsement.....	306.82	5	✓ Medsurge
Subsidised only for use in a programmable pump in patients where oral antispastic agents have been ineffective or have caused intolerable side effects and the prescription is endorsed accordingly.			

DANTROLENE

Cap 25 mg	97.50	100	✓ Dantrium
Cap 50 mg	77.00	100	✓ Dantrium S29 ^{S29}
			✓ Dantrium

ORPHENADRINE CITRATE

Tab 100 mg	20.76	100	✓ Norflex
------------------	-------	-----	-----------

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE

▲ Cap 100 mg 38.24 60 ✓ **Symmetrel**

APOMORPHINE HYDROCHLORIDE

▲ Inj 10 mg per ml, 2 ml ampoule 59.50 5 ✓ **Movapo**

▲ Inj 10 mg per ml, 5 ml ampoule 121.84 5 ✓ **Movapo**

BROMOCRIPTINE MESYLATE – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of bromocriptine mesylate.

* Tab 2.5 mg 11.70 30 ✓ **Parlodel** ^{\$29}

(Parlodel ^{\$29} Tab 2.5 mg to be delisted 1 September 2022)

ENTACAPONE – Brand switch fee payable (Pharmacode 2634139) - see page 241 for details

▲ Tab 200 mg 18.04 100 ✓ **Comtan**

LEVODOPA WITH BENSERAZIDE

* Tab dispersible 50 mg with benserazide 12.5 mg 13.25 100 ✓ **Madopar Rapid**

* Cap 50 mg with benserazide 12.5 mg 13.75 100 ✓ **Madopar 62.5**

* Cap 100 mg with benserazide 25 mg 15.80 100 ✓ **Madopar 125**

* Cap long-acting 100 mg with benserazide 25 mg 22.85 100 ✓ **Madopar HBS**

* Cap 200 mg with benserazide 50 mg 26.25 100 ✓ **Madopar 250**

LEVODOPA WITH CARBIDOPA

* Tab 100 mg with carbidopa 25 mg 21.11 100 ✓ **Sinemet**

* Tab long-acting 200 mg with carbidopa 50 mg 43.65 100 ✓ **Sinemet CR**

* Tab 250 mg with carbidopa 25 mg 38.39 100 ✓ **Sinemet**

PRAMIPEXOLE HYDROCHLORIDE

▲ Tab 0.25 mg 6.12 100 ✓ **Ramipex**

▲ Tab 1 mg 20.73 100 ✓ **Ramipex**

RASAGILINE

* Tab 1 mg 53.50 30 ✓ **Azilect** ^{\$29}

ROPINIROLE HYDROCHLORIDE

▲ Tab 0.25 mg 2.85 84 ✓ **Ropin**

..... 3.39 100 ✓ **Mylan** ^{\$29}

▲ Tab 1 mg 3.95 84 ✓ **Ropin**

..... 4.70 100 ✓ **Mylan** ^{\$29}

▲ Tab 2 mg 5.48 84 ✓ **Ropin**

▲ Tab 5 mg 12.50 84 ✓ **Ropin**

(Mylan ^{\$29} Tab 1 mg to be delisted 1 December 2022)

SELEGILINE HYDROCHLORIDE – Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

* Tab 5 mg 48.00 100 ✓ **Eldepryl** ^{\$29}

TOLCAPONE

▲ Tab 100 mg 152.38 100 ✓ **Tasmar**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Anticholinergics				
BENZATROPINE MESYLATE				
Tab 2 mg	9.59	60	✓	Benztrop
Inj 1 mg per ml, 2 ml	95.00	5	✓	Phebra
a) Up to 10 inj available on a PSO				
b) Only on a PSO				
PROCYCLIDINE HYDROCHLORIDE				
Tab 5 mg	7.40	100	✓	Kemadrin

Agents for Essential Tremor, Chorea and Related Disorders

RILUZOLE – Special Authority see [SA1403 below](#) – Retail pharmacy

Wastage claimable

Tab 50 mg	130.00	56	✓	Rilutek
-----------------	--------	----	---	----------------

»SA1403 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has amyotrophic lateral sclerosis with disease duration of 5 years or less; and
- 2 The patient has at least 60 percent of predicted forced vital capacity within 2 months prior to the initial application; and
- 3 The patient has not undergone a tracheostomy; and
- 4 The patient has not experienced respiratory failure; and
- 5 Any of the following:
 - 5.1 The patient is ambulatory; or
 - 5.2 The patient is able to use upper limbs; or
 - 5.3 The patient is able to swallow.

Renewal from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 The patient has not undergone a tracheostomy; and
- 2 The patient has not experienced respiratory failure; and
- 3 Any of the following:
 - 3.1 The patient is ambulatory; or
 - 3.2 The patient is able to use upper limbs; or
 - 3.3 The patient is able to swallow.

TETRABENAZINE				
Tab 25 mg	91.10	112	✓	Motetis

Anaesthetics

Local

LIDOCAINE [LIGNOCAINE]

Gel 2%, tube – Subsidy by endorsement	14.50	30 ml	✓	Xylocaine 2% Jelly
a) Up to 150 ml available on a PSO				
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.				
Gel 2%, 11 ml urethral syringe – Subsidy by endorsement	42.00	10	✓	Instillagel Lido
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral, cervical or rectal administration and the prescription is endorsed accordingly.				

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%.....	38.00	200 ml	✓	Mucosoothe
Inj 1%, 5 ml ampoule – Up to 25 inj available on a PSO.....	8.75	25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule – Up to 5 inj available on a PSO.....	8.25	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule – Up to 5 inj available on a PSO.....	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial – Up to 5 inj available on a PSO.....	6.20	5	✓	Lidocaine-Baxter
			✓	Lidocaine-Clarix
			✓	Lidocaine-Baxter
Inj 2%, 20 ml vial – Up to 5 inj available on a PSO.....	6.45	5		
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes –				
Subsidy by endorsement.....	103.32	10	✓	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical administration and the prescription is endorsed accordingly.				

Topical Local Anaesthetics

►SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see SA0906 above – Retail pharmacy

Crm 4%.....	5.40	5 g OP	✓	LMX4
	27.00	30 g OP	✓	LMX4

LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE – Special Authority see SA0906 above – Retail pharmacy

Crm 2.5% with prilocaine 2.5%.....	45.00	30 g OP	✓	EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	✓	EMLA

Analgesics

For Anti-inflammatory NSAIDs refer to MUSCULOSKELETAL, [page 110](#)

Non-opioid Analgesics

ASPIRIN

* Tab dispersible 300 mg – Up to 30 tab available on a PSO.....4.50 100 ✓ **Ethics Aspirin**

CAPSAICIN – Subsidy by endorsement

Subsidised only if prescribed for post-herpetic neuralgia or diabetic peripheral neuropathy and the prescription is endorsed accordingly.

Crm 0.075%.....	11.95	45 g OP	✓	Zostrix HP
	15.83	57 g OP	✓	Rugby Capsaicin
				Topical
				Cream \$29

NEFOPAM HYDROCHLORIDE

Tab 30 mg23.40 90 ✓ **Acupan**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PARACETAMOL				
Tab 500 mg - blister pack	19.75	1,000	✓	<u>Pacimol</u>
a) Maximum of 300 tab per prescription; can be waived by endorsement b) Up to 30 tab available on a PSO c) <ol style="list-style-type: none"> 1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition. 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing. 				
Tab 500 mg - bottle pack – Maximum of 300 tab per prescription; can be waived by endorsement	17.92	1,000	✓	<u>Noumed Paracetamol</u>
1) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater, and the prescription is annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition. 2) Maximum of 100 tab per dispensing for non-endorsed patients. If quantities prescribed for more than 100 tabs (for non-endorsed patients), then dispense in repeat dispensings not exceeding 100 tab per dispensing.				
Oral liq 120 mg per 5 ml	5.45	1,000 ml	✓	<u>Paracare</u>
a) Maximum of 600 ml per prescription; can be waived by endorsement b) Up to 200 ml available on a PSO c) Not in combination d) <ol style="list-style-type: none"> 1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing. 2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition. 				
Oral liq 250 mg per 5 ml	6.25	1,000 ml	✓	<u>Paracare Double Strength</u>
a) Maximum of 600 ml per prescription; can be waived by endorsement b) Up to 100 ml available on a PSO c) Not in combination d) <ol style="list-style-type: none"> 1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing. 2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition. 				
* Suppos 125 mg	3.59	10	✓	<u>Gacet</u>
* Suppos 250 mg	4.18	10	✓	<u>Gacet</u>
* Suppos 500 mg	12.40	50	✓	<u>Gacet</u>

Opioid Analgesics

CODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency

Tab 15 mg	6.25	100	✓	<u>PSM</u>
Tab 30 mg	7.45	100	✓	<u>PSM</u>
Tab 60 mg	14.25	100	✓	<u>PSM</u>

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
DIHYDROCODEINE TARTRATE				
Tab long-acting 60 mg.....	8.60	60	✓	DHC Continus
FENTANYL				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Inj 50 mcg per ml, 2 ml ampoule	3.75	10	✓	Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	9.41	10	✓	Boucher and Muir
Patch 12.5 mcg per hour	6.99	5	✓	Fentanyl Sandoz
Patch 25 mcg per hour	7.99	5	✓	Fentanyl Sandoz
Patch 50 mcg per hour	9.49	5	✓	Fentanyl Sandoz
Patch 75 mcg per hour	17.99	5	✓	Fentanyl Sandoz
Patch 100 mcg per hour	18.59	5	✓	Fentanyl Sandoz
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).				
e) For methadone hydrochloride oral liquid refer Standard Formulae, page 243				
Tab 5 mg	1.40	10	✓	Methatabs
Oral liq 2 mg per ml	6.40	200 ml	✓	Biodone
Oral liq 5 mg per ml	6.40	200 ml	✓	Biodone Forte
Oral liq 10 mg per ml	7.50	200 ml	✓	Biodone Extra Forte
Inj 10 mg per ml, 1 ml	61.00	10	✓	AFT
MORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Oral liq 1 mg per ml	9.28	200 ml	✓	RA-Morph
Oral liq 2 mg per ml	16.24	200 ml	✓	RA-Morph
Oral liq 5 mg per ml	19.44	200 ml	✓	Ordine ^{S29}
			✓	RA-Morph
Oral liq 10 mg per ml	27.74	200 ml	✓	Ordine ^{S29}
			✓	RA-Morph

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab immediate-release 10 mg.....	2.80	10	✓	<u>Sevredol</u>
Tab immediate-release 20 mg.....	5.52	10	✓	<u>Sevredol</u>
Cap long-acting 10 mg.....	2.05	10	✓	<u>m-Eslon</u>
Cap long-acting 30 mg.....	3.00	10	✓	<u>m-Eslon</u>
Cap long-acting 60 mg.....	6.12	10	✓	<u>m-Eslon</u>
Cap long-acting 100 mg.....	7.13	10	✓	<u>m-Eslon</u>
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	6.99	5	✓	<u>DBL Morphine Sulphate</u>
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	5.61	5	✓	<u>DBL Morphine Sulphate</u>
Inj 15 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	7.08	5	✓	<u>DBL Morphine Sulphate</u>
Inj 30 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	7.28	5	✓	<u>DBL Morphine Sulphate</u>
OXYCODONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab controlled-release 5 mg.....	2.69	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 10 mg.....	2.69	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 20 mg.....	3.49	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 40 mg.....	5.49	20	✓	<u>Oxycodone Sandoz</u>
Tab controlled-release 80 mg.....	12.99	20	✓	<u>Oxycodone Sandoz</u>
Cap immediate-release 5 mg.....	1.88	20	✓	<u>OxyNorm</u>
Cap immediate-release 10 mg.....	3.32	20	✓	<u>OxyNorm</u>
Cap immediate-release 20 mg.....	5.23	20	✓	<u>OxyNorm</u>
Oral liq 5 mg per 5 ml.....	11.20	250 ml	✓	<u>OxyNorm</u>
Inj 10 mg per ml, 1 ml ampoule.....	5.82	5	✓	<u>Hameln</u>
	7.28		✓	<u>OxyNorm</u>
Hameln to be Principal Supply on 1 July 2022				
Inj 10 mg per ml, 2 ml ampoule.....	11.49	5	✓	<u>Hameln</u>
	14.36		✓	<u>OxyNorm</u>
Hameln to be Principal Supply on 1 July 2022				
Inj 50 mg per ml, 1 ml ampoule.....	22.92	5	✓	<u>Hameln</u>
	30.60		✓	<u>OxyNorm</u>
Hameln to be Principal Supply on 1 July 2022				
<i>(OxyNorm Inj 10 mg per ml, 1 ml ampoule to be delisted 1 July 2022)</i>				
<i>(OxyNorm Inj 10 mg per ml, 2 ml ampoule to be delisted 1 July 2022)</i>				
<i>(OxyNorm Inj 50 mg per ml, 1 ml ampoule to be delisted 1 July 2022)</i>				
PARACETAMOL WITH CODEINE – Safety medicine; prescriber may determine dispensing frequency				
* Tab paracetamol 500 mg with codeine phosphate 8 mg.....	26.51	1,000	✓	<u>Paracetamol + Codeine (Relieve)</u>

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PETHIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
Tab 50 mg	4.70	10	✓ PSM	
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	29.88	5	✓ DBL Pethidine Hydrochloride	
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	30.72	5	✓ DBL Pethidine Hydrochloride	
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg	1.52	20	✓ Tramal SR 100	
Tab sustained-release 150 mg	2.10	20	✓ Tramal SR 150	
Tab sustained-release 200 mg	2.75	20	✓ Tramal SR 200	
Cap 50 mg	2.80	100	✓ Arrow-Tramadol	

Antidepressants

Cyclic and Related Agents

AMITRIPTYLINE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg	2.49	100	✓ Arrow-Amitriptyline	
Tab 25 mg	1.51	100	✓ Arrow-Amitriptyline	
Tab 50 mg	2.51	100	✓ Arrow-Amitriptyline	
CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg	10.17	30	✓ Clomipramine Teva	
Tab 25 mg	11.99	30	✓ Clomipramine Teva	
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by endorsement				
a) Safety medicine; prescriber may determine dispensing frequency				
b) Subsidy by endorsement – Subsidised for patients who were taking dosulepin [dothiepin] hydrochloride prior to 1 June 2019 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of dosulepin [dothiepin] hydrochloride.				
Tab 75 mg	3.85	30	✓ Dosulepin Mylan	
Cap 25 mg	7.83	50	✓ Dosulepin Mylan ^{S29}	
IMIPRAMINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg	5.48	50	✓ Tofranil	
Tab 25 mg	10.96	100	✓ Tofranil	
Tab 25 mg	8.80	50	✓ Tofranil	
NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg	2.44	100	✓ Norpress	
Tab 25 mg	5.98	180	✓ Norpress	

Monoamine-Oxidase Inhibitors (MAOIs) - Non Selective

TRANLYCYPROMINE SULPHATE				
Tab 10 mg	12.85	28	✓ Parnate ^{S29} ^{S29}	
	22.94	50	✓ Parnate	
	45.88	100	✓ Parnate ^{S29} ^{S29}	
	96.00		✓ Parnate ^{S29} ^{S29}	

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

Monoamine-Oxidase Type A Inhibitors

MOCLOBEMIDE

* Tab 150 mg	11.80	60	✓ <u>Aurorix</u>
* Tab 300 mg	19.25	60	✓ <u>Aurorix</u>

Selective Serotonin Reuptake Inhibitors

CITALOPRAM HYDROBROMIDE

* Tab 20 mg	1.91	84	✓ <u>PSM Citalopram</u>
-------------------	------	----	-------------------------

ESCITALOPRAM

* Tab 10 mg	1.07	28	✓ <u>Escitalopram</u> (Ethics)
* Tab 20 mg	1.92	28	✓ <u>Escitalopram</u> (Ethics)

FLUOXETINE HYDROCHLORIDE

* Tab dispersible 20 mg, scored – Subsidy by endorsement	1.84	28	✓ <u>Fluox</u>
	1.98	30	✓ <u>Fluox</u>

Subsidised by endorsement

- 1) When prescribed for a patient who cannot swallow whole tablets or capsules and the prescription is endorsed accordingly; or
- 2) When prescribed in a daily dose that is not a multiple of 20 mg in which case the prescription is deemed to be endorsed. Note: Tablets should be combined with capsules to facilitate incremental 10 mg doses.

Cap 20 mg	2.91	84	✓ <u>Fluox</u>
-----------------	------	----	----------------

PAROXETINE

* Tab 20 mg	3.61	90	✓ <u>Loxamine</u>
-------------------	------	----	-------------------

SERTRALINE

* Tab 50 mg	0.92	30	✓ <u>Setrona</u>
			✓ <u>Setrona AU</u>
* Tab 100 mg	1.61	30	✓ <u>Setrona</u>
			✓ <u>Setrona AU</u>

Other Antidepressants

MIRTAZAPINE

Tab 30 mg	2.60	28	✓ <u>Noumed</u>
Tab 45 mg	3.45	28	✓ <u>Noumed</u>

VENLAFAXINE

* Cap 37.5 mg	6.38	84	✓ <u>Enlafax XR</u>
* Cap 75 mg	8.11	84	✓ <u>Enlafax XR</u>
* Cap 150 mg	11.16	84	✓ <u>Enlafax XR</u>

Antiepilepsy Drugs

Agents for Control of Status Epilepticus

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency			
Inj 5 mg per ml, 2 ml ampoule – Subsidy by endorsement.....	23.66	5	✓ Hospira
a) Up to 5 inj available on a PSO			
b) Only on a PSO			
c) PSO must be endorsed “not for anaesthetic procedures”.			
Rectal tubes 5 mg – Up to 5 tube available on a PSO	43.50	5	✓ Stesolid
PHENYTOIN SODIUM			
* Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	104.58	5	✓ Hospira
* Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a PSO	154.01	5	✓ Hospira

Control of Epilepsy

CARBAMAZEPINE			
* Tab 200 mg	14.53	100	✓ Tegretol
* Tab long-acting 200 mg	16.98	100	✓ Tegretol CR
* Tab 400 mg	34.58	100	✓ Tegretol
* Tab long-acting 400 mg	39.17	100	✓ Tegretol CR
* Oral liq 20 mg per ml	26.37	250 ml	✓ Tegretol
CLOBAZAM – Safety medicine; prescriber may determine dispensing frequency			
Tab 10 mg	9.12	50	✓ Frisium
CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency			
Oral drops 2.5 mg per ml	7.38	10 ml OP	✓ Rivotril
ETHOSUXIMIDE			
Cap 250 mg	140.88	100	✓ Zarontin
Oral liq 250 mg per 5 ml	56.35	200 ml	✓ Zarontin
GABAPENTIN			
Note: Not subsidised in combination with subsidised pregabalin			
* Cap 100 mg	6.45	100	✓ Nupentin
* Cap 300 mg	8.45	100	✓ Nupentin
* Cap 400 mg	10.26	100	✓ Nupentin
LACOSAMIDE – Special Authority see SA1125 below – Retail pharmacy			
▲ Tab 50 mg	25.04	14	✓ Vimpat
▲ Tab 100 mg	50.06	14	✓ Vimpat
	200.24	56	✓ Vimpat
▲ Tab 150 mg	75.10	14	✓ Vimpat
	300.40	56	✓ Vimpat
▲ Tab 200 mg	400.55	56	✓ Vimpat

►SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:
Both:

- 1 Patient has partial-onset epilepsy; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note).

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

LAMOTRIGINE

▲ Tab dispersible 2 mg	55.00	30	✓ Lamictal
▲ Tab dispersible 5 mg	50.00	30	✓ Lamictal
* Tab dispersible 25 mg	2.76	56	✓ Logem
* Tab dispersible 50 mg	3.31	56	✓ Logem
* Tab dispersible 100 mg	4.40	56	✓ Logem

LEVETIRACETAM

Tab 250 mg	4.99	60	✓ Everet
Tab 500 mg	8.79	60	✓ Everet
Tab 750 mg	14.39	60	✓ Everet
Tab 1,000 mg	18.59	60	✓ Everet
Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT

PHENOBARBITONE

For phenobarbitone oral liquid refer Standard Formulae, [page 243](#)

* Tab 15 mg	40.00	500	✓ PSM
* Tab 30 mg	40.00	500	✓ PSM

PHENYTOIN SODIUM

* Tab 50 mg	75.00	200	✓ Dilantin Infatab
Cap 30 mg	74.00	200	✓ Dilantin
Cap 100 mg	37.00	200	✓ Dilantin
* Oral liq 30 mg per 5 ml	22.03	500 ml	✓ Dilantin

PREGABALIN

Note: Not subsidised in combination with subsidised gabapentin

Cap 25 mg	2.25	56	✓ Pregabalin Pfizer
* Cap 75 mg	2.65	56	✓ Pregabalin Pfizer
Cap 150 mg	4.01	56	✓ Lyrica
			✓ Pregabalin Pfizer
Cap 300 mg	7.38	56	✓ Pregabalin Pfizer

PRIMIDONE

* Tab 250 mg	37.35	100	✓ Apo-Primidone
			✓ Primidone Clinect

(Apo-Primidone Tab 250 mg to be delisted 1 January 2023)

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC	52.24	100	✓	Epilim
* Oral liq 200 mg per 5 ml	20.48	300 ml	✓	Epilim S/F Liquid
			✓	Epilim Syrup
			✓	Epilim IV
STIRIPENTOL – Special Authority see SA1330 below – Retail pharmacy				
Cap 250 mg	509.29	60	✓	Diacomit \$29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit \$29

►SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

TOPIRAMATE

▲ Tab 25 mg	11.07	60	✓	Arrow-Topiramate
	26.04		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 50 mg	18.81	60	✓	Arrow-Topiramate
	44.26		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 100 mg	31.99	60	✓	Arrow-Topiramate
	75.25		✓	Topiramate Actavis
			✓	Topamax
▲ Tab 200 mg	55.19	60	✓	Arrow-Topiramate
	129.85		✓	Topiramate Actavis
			✓	Topamax
▲ Sprinkle cap 15 mg	20.84	60	✓	Topamax
▲ Sprinkle cap 25 mg	26.04	60	✓	Topamax

VIGABATRIN – Special Authority see SA2088 below – Retail pharmacy

▲ Tab 500 mg	119.30	100	✓	Sabril
--------------------	--------	-----	---	---------------

►SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

1.3 Patient has tuberous sclerosis complex; and

2 Either:

2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or

2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Notes: "Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and

2 Either:

2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or

2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields..

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, [page 110](#)

Acute Migraine Treatment

RIZATRIPTAN

Tab orodispersible 10 mg	3.65	30	✓ Rizamelt
--------------------------------	------	----	-------------------

SUMATRIPTAN

Tab 50 mg	14.41	90	✓ Sumagran
-----------------	-------	----	-------------------

Tab 100 mg	22.68	90	✓ Sumagran
------------------	-------	----	-------------------

Inj 12 mg per ml, 0.5 ml prefilled pen – Maximum of 10 inj per prescription	34.00	2 OP	✓ Imigran
---	-------	------	------------------

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, [page 50](#)

PIZOTIFEN

* Tab 500 mcg	23.21	100	✓ Sandomigran
---------------------	-------	-----	----------------------

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, [page 8](#)

APREPITANT – Special Authority see [SA0987 on the next page](#) – Retail pharmacy

Cap 2 x 80 mg and 1 x 125 mg	30.00	3 OP	✓ Emend Tri-Pack
------------------------------------	-------	------	-------------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
►SA0987 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.				
Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.				
BETAHISTINE DIHYDROCHLORIDE				
* Tab 16 mg	3.88	84	✓	Vergo 16
	4.62	100	✓	Serc
<i>(Vergo 16 Tab 16 mg to be delisted 1 July 2022)</i>				
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.49	10	✓	Nausicalm
CYCLIZINE LACTATE				
Inj 50 mg per ml, 1 ml	21.53	10	✓	Hameln
DOMPERIDONE				
* Tab 10 mg	2.85	100	✓	Pharmacy Health
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓	Martindale <small>\$29</small>
Patch 1.5 mg – Special Authority see SA1998 below – Retail pharmacy.....	14.11	2	✓	Scopoderm TTS
►SA1998 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:				
Either:				
1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or				
2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.				
Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.				
METOCLOPRAMIDE HYDROCHLORIDE				
* Tab 10 mg – Up to 30 tab available on a PSO	1.30	100	✓	Metoclopramide Actavis 10
* Inj 5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	9.50	10	✓	Pfizer
ONDANSETRON				
* Tab 4 mg	2.68	50	✓	Onrex
* Tab disp 4 mg – Up to 10 tab available on a PSO	0.76	10	✓	 Ondansetron ODT-DRLA
* Tab 8 mg	4.57	50	✓	Onrex
* Tab disp 8 mg – Up to 10 tab available on a PSO	1.13	10	✓	 Ondansetron ODT-DRLA
PROCHLORPERAZINE				
* Tab 3 mg buccal.....	5.97	50		Buccastem
	(30.00)			
* Tab 5 mg – Up to 30 tab available on a PSO	8.00	250	✓	Nausafix
* Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO	25.81	10	✓	Stemetil

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Antipsychotics				
General				
AMISULPRIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 100 mg	5.15	30	✓	Sulprix
	17.16	100	✓	Amisulpride Mylan <small>§29</small>
Tab 200 mg	14.96	60	✓	Sulprix
Tab 400 mg	29.78	60	✓	Sulprix
<i>(Amisulpride Mylan <small>§29</small> Tab 100 mg to be delisted 1 August 2022)</i>				
ARIPIPRAZOLE – Safety medicine; prescriber may determine dispensing frequency				
Tab 5 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 10 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 15 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 20 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 30 mg	10.50	30	✓	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg – Up to 30 tab available on a PSO	14.83	100	✓	Largactil
Tab 25 mg – Up to 30 tab available on a PSO	15.62	100	✓	Largactil
Tab 100 mg – Up to 30 tab available on a PSO	36.73	100	✓	Largactil
Inj 25 mg per ml, 2 ml – Up to 5 inj available on a PSO	30.79	10	✓	Largactil
CLOZAPINE – Hospital pharmacy [HP4] Safety medicine; prescriber may determine dispensing frequency				
Tab 25 mg	6.69	50	✓	Clopine
	13.37	100	✓	Clozaril
			✓	Clopine
			✓	Clozaril
Tab 50 mg	8.67	50	✓	Clopine
	17.33	100	✓	Clopine
Tab 100 mg	17.33	50	✓	Clopine
			✓	Clozaril
	34.65	100	✓	Clopine
			✓	Clozaril
Tab 200 mg	34.65	50	✓	Clopine
	69.30	100	✓	Clopine
Suspension 50 mg per ml	67.62	100 ml	✓	Versacloz
HALOPERIDOL – Safety medicine; prescriber may determine dispensing frequency				
Tab 500 mcg – Up to 30 tab available on a PSO	6.23	100	✓	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO	9.43	100	✓	Serenace
Tab 5 mg – Up to 30 tab available on a PSO	14.86	50	✓	Serenace
	29.72	100	✓	Serenace
Oral liq 2 mg per ml – Up to 200 ml available on a PSO	23.84	100 ml	✓	Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	21.55	10	✓	Serenace
LEVOMEPRMAZINE – Safety medicine; prescriber may determine dispensing frequency				
Tab 25 mg (33.8 mg as a maleate)	16.10	100	✓	Nozinan (Swiss)
Tab 25 mg as a maleate	16.10	100	✓	Nozinan
Tab 100 mg (135 mg as a maleate)	41.75	100	✓	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	✓	Nozinan

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

NERVOUS SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LEVOMEPROMAZINE HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Inj 25 mg per ml, 1 ml ampoule	33.50	10	✓	Nozinan
LITHIUM CARBONATE – Safety medicine; prescriber may determine dispensing frequency				
Tab long-acting 400 mg	72.00	100	✓	Priadel
Cap 250 mg	9.42	100	✓	Douglas
OLANZAPINE – Safety medicine; prescriber may determine dispensing frequency				
Tab 2.5 mg	1.35	28	✓	Zypine
Tab 5 mg	1.58	28	✓	Zypine
Tab orodispersible 5 mg	1.81	28	✓	Zypine ODT
Tab 10 mg	2.01	28	✓	Zypine
Tab orodispersible 10 mg	2.38	28	✓	Zypine ODT
PERICAZINE – Safety medicine; prescriber may determine dispensing frequency				
Tab 2.5 mg	10.49	84	✓	Neulactil
	12.49	100	✓	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
	44.45	100	✓	Neulactil
QUETIAPINE – Safety medicine; prescriber may determine dispensing frequency				
Tab 25 mg	2.15	90	✓	Quetapel
Tab 100 mg	5.06	90	✓	Quetapel
Tab 200 mg	8.90	90	✓	Quetapel
Tab 300 mg	12.86	90	✓	Quetapel
RISPERIDONE – Safety medicine; prescriber may determine dispensing frequency				
Tab 0.5 mg	1.86	60	✓	Risperidone (Teva)
Tab 1 mg	2.06	60	✓	Risperidone (Teva)
Tab 2 mg	2.29	60	✓	Risperidone (Teva)
Tab 3 mg	2.50	60	✓	Risperidone (Teva)
Tab 4 mg	3.42	60	✓	Risperidone (Teva)
Oral liq 1 mg per ml	8.90	30 ml	✓	Risperon
ZIPRASIDONE – Safety medicine; prescriber may determine dispensing frequency				
Cap 20 mg	17.90	60	✓	Zusdone
Cap 40 mg	27.41	60	✓	Zusdone
Cap 60 mg	38.39	60	✓	Zusdone
Cap 80 mg	46.55	60	✓	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; prescriber may determine dispensing frequency				
Tab 10 mg	31.45	100	✓	Clopixol

Depot Injections

FLUPENTHIXOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency				
Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO	13.14	5	✓	Fluanxol
Inj 20 mg per ml, 2 ml – Up to 5 inj available on a PSO	20.90	5	✓	Fluanxol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO	40.87	5	✓	Fluanxol
HALOPERIDOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency				
Inj 50 mg per ml, 1 ml – Up to 5 inj available on a PSO	28.39	5	✓	Haldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO	55.90	5	✓	Haldol Concentrate
			✓	Haldol
				Decanoas \$29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
OLANZAPINE – Special Authority see SA1428 below – Retail pharmacy				
Safety medicine; prescriber may determine dispensing frequency				
Inj 210 mg vial	252.00	1	✓	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓	Zyprexa Relprevv
Inj 405 mg vial	504.00	1	✓	Zyprexa Relprevv

► **SA1428** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE – Special Authority see [SA1429 below](#) – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg syringe	194.25	1	✓	Invega Sustenna
Inj 50 mg syringe	271.95	1	✓	Invega Sustenna
Inj 75 mg syringe	357.42	1	✓	Invega Sustenna
Inj 100 mg syringe	435.12	1	✓	Invega Sustenna
Inj 150 mg syringe	435.12	1	✓	Invega Sustenna

► **SA1429** Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE – Special Authority see [SA1427 on the next page](#) – Retail pharmacy

Safety medicine; prescriber may determine dispensing frequency

Inj 25 mg vial	135.98	1	✓	Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓	Risperdal Consta
Inj 50 mg vial	217.56	1	✓	Risperdal Consta

►SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE – Safety medicine; prescriber may determine dispensing frequency
 Inj 200 mg per ml, 1 ml – Up to 5 inj available on a PSO 19.80 5 ✓ **Clopixol**

Anxiolytics

BUSPIRONE HYDROCHLORIDE

* Tab 5 mg	18.50	100	✓ Buspirone Viatris
* Tab 10 mg	12.50	100	✓ Buspirone Viatris

CLONAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam

DIAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam

LORAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ Ativan

Multiple Sclerosis Treatments

►SA2051 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) must be confirmed by a neurologist. Diagnosis must include MRI confirmation; and
- 2 Patients must have Clinically Definite Relapsing multiple sclerosis with or without underlying progression; and
- 3 Patients must have an EDSS score between 0 – 6.0; and
- 4 Patient has had at least 1 significant relapse of multiple sclerosis in the previous 12 months or 2 significant relapses in the past 24 months; and
- 5 All of the following:
 - 5.1 Each significant relapse must be confirmed by the applying neurologist or general physician (the patient may not

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- necessarily have been seen by them during the relapse, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
- 5.2 Each significant relapse is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 5.3 Each significant relapse has lasted at least one week and has started at least one month after the onset of a previous relapse; and
 - 5.4 Each significant relapse can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 5.5 Either:
 - 5.5.1 Each significant relapse is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 5.5.2 Each significant relapse is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
 - 6 Evidence of new inflammatory activity on an MR scan within the past 24 months; and
 - 7 Any of the following:
 - 7.1 A sign of that new inflammatory activity is a gadolinium enhancing lesion; or
 - 7.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 7.3 A sign of that new inflammatory activity is a T2 lesion with associated local swelling; or
 - 7.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent relapse that occurred within the last 2 years; or
 - 7.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MR scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE – Special Authority see [SA2051 on the previous page](#) – Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg.....	520.00	14	✓ Tecfidera
Cap 240 mg.....	2,000.00	56	✓ Tecfidera

FINGOLIMOD – Special Authority see [SA2051 on the previous page](#) – Retail pharmacy

a) Wastage claimable

b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 0.5 mg.....	2,200.00	28	✓ Gilenya
-----------------	----------	----	------------------

GLATIRAMER ACETATE – Special Authority see [SA2051 on the previous page](#) – Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Inj 40 mg prefilled syringe.....	1,137.48	12	✓ Copaxone
----------------------------------	----------	----	-------------------

INTERFERON BETA-1-ALPHA – Special Authority see [SA2051 on the previous page](#) – Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Inj 6 million iu prefilled syringe.....	1,170.00	4	✓ Avonex
Injection 6 million iu per 0.5 ml pen injector.....	1,170.00	4	✓ Avonex Pen

INTERFERON BETA-1-BETA – Special Authority see [SA2051 on the previous page](#) – Retail pharmacy

Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Inj 8 million iu per 1 ml.....	1,322.89	15	✓ Betaferon
--------------------------------	----------	----	--------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
NATALIZUMAB – Special Authority see SA2051 on page 134 – Retail pharmacy				
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.				
Inj 20 mg per ml, 15 ml vial.....	1,750.00	1	✓	Tysabri
OCRELIZUMAB – Special Authority see SA2051 on page 134 – Retail pharmacy				
Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.				
Inj 30 mg per ml, 10 ml vial.....	9,346.00	1	✓	Ocrevus
TERIFLUNOMIDE – Special Authority see SA2051 on page 134 – Retail pharmacy				
a) Wastage claimable				
b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.				
Tab 14 mg	659.90	28	✓	Aubagio

Sedatives and Hypnotics

MELATONIN – Special Authority see SA1666 below – Retail pharmacy				
Brand switch fee payable (Pharmacode 2634112) - see page 241 for details				
Tab modified-release 2 mg – No more than 5 tab per day	11.50	30	✓	Vigisom

►SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispensing frequency				
Inj 1 mg per ml, 5 ml ampoule	5.50	10	✓	Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule – Up to 10 inj available on a PSO	17.28	10	✓	Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only.				
Inj 5 mg per ml, 3 ml ampoule	4.50	5	✓	Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule – Up to 5 inj available on a PSO	13.09	5	✓	Pfizer
On a PSO for status epilepticus use only. PSO must be endorsed for status epilepticus use only.				
PHENOBARBITONE SODIUM – Special Authority see SA1386 on the next page – Retail pharmacy				
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓	Max Health <small>S29</small>

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

►SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine dispensing frequency

Tab 10 mg	1.33	25	✓ <u>Normison</u>
-----------------	------	----	-------------------

TRIAZOLAM – Safety medicine; prescriber may determine dispensing frequency

Tab 125 mcg.....	5.10	100	
	(9.85)		Hypam
Tab 250 mcg.....	4.10	100	
	(11.20)		Hypam

ZOPICLONE – Safety medicine; prescriber may determine dispensing frequency

Tab 7.5 mg	10.80	500	✓ <u>Zopiclone Actavis</u>
------------------	-------	-----	----------------------------

Stimulants/ADHD Treatments

ATOMOXETINE

Cap 10 mg	18.41	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29 ✓ Generic Partners
	107.03		✓ Strattera
Cap 18 mg	27.06	28	✓ APO-Atomoxetine ✓ Generic Partners
	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ APO-Atomoxetine ✓ Generic Partners
Cap 40 mg	29.22	28	✓ APO-Atomoxetine ✓ Generic Partners
	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29 ✓ Generic Partners
Cap 80 mg	56.45	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29 ✓ Generic Partners
Cap 100 mg	58.48	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine S29 S29 ✓ Generic Partners

DEXAMFETAMINE SULFATE – Special Authority see SA1149 on the next page – Retail pharmacy

a) Only on a controlled drug form

b) Safety medicine; prescriber may determine dispensing frequency

Tab 5 mg	21.00	100	✓ <u>PSM</u>
----------------	-------	-----	--------------

►SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE – Special Authority see [SA1964 on the next page](#) – Retail pharmacy

a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing frequency			
Tab immediate-release 5 mg.....	3.20	30	✓ Rubifen
Tab immediate-release 10 mg.....	3.00	30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg.....	7.75	30	✓ Methylphenidate ER - Teva
Tab immediate-release 20 mg.....	7.85	30	✓ Rubifen
Tab sustained-release 20 mg.....	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg.....	11.45	30	✓ Methylphenidate ER - Teva
Tab extended-release 36 mg.....	15.50	30	✓ Methylphenidate ER - Teva
Tab extended-release 54 mg.....	22.25	30	✓ Methylphenidate ER - Teva

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE – Special Authority see SA1965 below – Retail pharmacy

a) Only on a controlled drug form			
b) Safety medicine; prescriber may determine dispensing frequency			
Tab extended-release 18 mg.....	58.96	30	✓ Concerta
Tab extended-release 27 mg.....	65.44	30	✓ Concerta
Tab extended-release 36 mg.....	71.93	30	✓ Concerta
Tab extended-release 54 mg.....	86.24	30	✓ Concerta
Cap modified-release 10 mg.....	15.60	30	✓ Ritalin LA
Cap modified-release 20 mg.....	20.40	30	✓ Ritalin LA
Cap modified-release 30 mg.....	25.52	30	✓ Ritalin LA
Cap modified-release 40 mg.....	30.60	30	✓ Ritalin LA

►SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL – Special Authority see [SA1999 below](#) – Retail pharmacy

Tab 100 mg	29.13	60	✓ Modavigil
------------------	-------	----	--------------------

►SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE

* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
RIVASTIGMINE – Special Authority see SA1488 below – Retail pharmacy				
Patch 4.6 mg per 24 hour	38.00	30	✓	Rivastigmine Patch BNM 5
Patch 9.5 mg per 24 hour	38.00	30	✓	Rivastigmine Patch BNM 10

► **SA1488** **Special Authority for Subsidy**

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE – Special Authority see SA1203 below – Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg	18.37	28	✓	Buprenorphine Naloxone BNM
Tab sublingual 8 mg with naloxone 2 mg	53.12	28	✓	Buprenorphine Naloxone BNM

► **SA1203** **Special Authority for Subsidy**

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health..

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg 11.00 30 ✓ **Zyban**

DISULFIRAM

Tab 200 mg 236.40 100 ✓ **Antabuse** S29

NALTREXONE HYDROCHLORIDE – Special Authority see [SA1408 below](#) – Retail pharmacy

Tab 50 mg 133.33 30 ✓ **Naltrexone**

► [SA1408](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to one of the District Health Boards or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
NICOTINE				
a) Nicotine will not be funded in amounts less than 4 weeks of treatment.				
b) Note: Direct Provision by a pharmacist permitted under the provisions in Part I of Section A.				
Patch 7 mg – Up to 28 patch available on a PSO	18.14	28	✓	Habitrol
Patch 7 mg for direct distribution only – [Xpharm]	3.94	7	✓	Habitrol
Patch 14 mg – Up to 28 patch available on a PSO	19.95	28	✓	Habitrol
Patch 14 mg for direct distribution only – [Xpharm]	4.52	7	✓	Habitrol
Patch 21 mg – Up to 28 patch available on a PSO	22.86	28	✓	Habitrol
Patch 21 mg for direct distribution only – [Xpharm]	5.18	7	✓	Habitrol
Lozenge 1 mg – Up to 216 loz available on a PSO	19.18	216	✓	Habitrol
Lozenge 1 mg for direct distribution only – [Xpharm]	3.20	36	✓	Habitrol
Lozenge 2 mg – Up to 216 loz available on a PSO	21.02	216	✓	Habitrol
Lozenge 2 mg for direct distribution only – [Xpharm]	3.24	36	✓	Habitrol
Gum 2 mg (Fruit) – Up to 384 piece available on a PSO	38.21	384	✓	Habitrol
Gum 2 mg (Fruit) for direct distribution only – [Xpharm]	8.64	96	✓	Habitrol
Gum 2 mg (Mint) – Up to 384 piece available on a PSO	38.21	384	✓	Habitrol
Gum 2 mg (Mint) for direct distribution only – [Xpharm]	8.64	96	✓	Habitrol
Gum 4 mg (Fruit) – Up to 384 piece available on a PSO	44.17	384	✓	Habitrol
Gum 4 mg (Fruit) for direct distribution only – [Xpharm]	10.01	96	✓	Habitrol
Gum 4 mg (Mint) – Up to 384 piece available on a PSO	44.17	384	✓	Habitrol
Gum 4 mg (Mint) for direct distribution only – [Xpharm]	10.01	96	✓	Habitrol

VARENICLINE TARTRATE – Special Authority see [SA1845 below](#) – Retail pharmacy

- A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- Varenicline will not be funded in amounts less than 4 weeks of treatment.
- The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg x 11 and 1 mg x 42	16.67	53 OP	✓	Varenicline Pfizer
Tab 1 mg	17.62	56	✓	Varenicline Pfizer

➔ **SA1845** **Special Authority for Subsidy**

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- Either:
 - The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- The patient is not pregnant; and
- The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria:

All of the following:

- Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking;

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.
Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.
This includes the 4-week 'starter' pack.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE – PCT only – Specialist – Special Authority see [SA2046 below](#)

Inj 25 mg vial	77.00	1	✓ Ribomustin
Inj 100 mg vial	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	3.23	1 mg	✓ Baxter

►SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Either:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

continued...

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
continued...				
2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and				
2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or				
2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.				
Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.				
Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:				
All of the following:				
1 Patient has Hodgkin's lymphoma requiring treatment; and				
2 Patient has a ECOG performance status of 0-2; and				
3 Patient has received one prior line of chemotherapy; and				
4 Patient's disease relapsed or was refractory following prior chemotherapy; and				
5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m ² twice per cycle, for a maximum of four cycles.				
Note: Indications marked with * are unapproved indications.				
BUSULFAN – PCT – Retail pharmacy-Specialist				
Tab 2 mg	89.25	100	✓	Myleran
CARBOPLATIN – PCT only – Specialist				
Inj 10 mg per ml, 45 ml vial.....	32.59	1	✓	DBL Carboplatin
	45.20		✓	Carboplatin Ebewe
	48.50		✓	Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓	Baxter
CARMUSTINE – PCT only – Specialist				
Inj 100 mg vial	710.00	1	✓	BiCNU
	1,387.00		✓	Bicnu Heritage ^{\$29}
BiCNU to be Principal Supply on 1 September 2022				
Inj 100 mg for ECP	710.00	100 mg OP	✓	Baxter
<i>(Bicnu Heritage ^{\$29} Inj 100 mg vial to be delisted 1 September 2022)</i>				
CHLORAMBUCIL – PCT – Retail pharmacy-Specialist				
Tab 2 mg	29.06	25	✓	Leukeran FC
CISPLATIN – PCT only – Specialist				
Inj 1 mg per ml, 50 ml vial.....	15.00	1	✓	Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial.....	21.00	1	✓	Cisplatin Ebewe
	29.66		✓	DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓	Baxter
CYCLOPHOSPHAMIDE				
Tab 50 mg – PCT – Retail pharmacy-Specialist.....	145.00	50	✓	Cyclonex
Inj 1 g vial – PCT – Retail pharmacy-Specialist.....	35.65	1	✓	Endoxan
	127.80	6	✓	Cytosan
Inj 2 g vial – PCT only – Specialist.....	71.25	1	✓	Endoxan
Inj 1 mg for ECP – PCT only – Specialist.....	0.04	1 mg	✓	Baxter
IFOSFAMIDE – PCT only – Specialist				
Inj 1 g.....	96.00	1	✓	Holoxan
Inj 2 g.....	180.00	1	✓	Holoxan
Inj 1 mg for ECP	0.10	1 mg	✓	Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LOMUSTINE – PCT – Retail pharmacy-Specialist				
Cap 10 mg.....	132.59	20	✓	CeeNU
Cap 40 mg.....	399.15	20	✓	CeeNU
MELPHALAN				
Tab 2 mg – PCT – Retail pharmacy-Specialist.....	40.70	25	✓	Alkeran
Inj 50 mg – PCT only – Specialist.....	67.80	1	✓	Alkeran
			✓	Alkeran S29
OXALIPLATIN – PCT only – Specialist				
Inj 100 mg vial	25.01	1	✓	Oxaliplatin Actavis 100
	110.00		✓	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial.....	46.32	1	✓	Oxaliplatin Accord
Inj 1 mg for ECP	0.48	1 mg	✓	Baxter
THIOTEPA – PCT only – Specialist				
Inj 15 mg vial	CBS	1	✓	Bedford S29
			✓	Max Health S29
			✓	THIO-TEPA S29
			✓	Tepadina S29
Inj 100 mg vial	CBS	1	✓	Max Health S29
			✓	Tepadina S29

Antimetabolites

AZACITIDINE – PCT only – Specialist – Special Authority see SA1467 below				
Inj 100 mg vial	75.06	1	✓	Azacitidine Dr Reddy's
	605.00		✓	Vidaza
Inj 1 mg for ECP	0.83	1 mg	✓	Baxter

➡SA1467 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- Any of the following:
 - The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- The patient has performance status (WHO/ECOG) grade 0-2; and
- The patient does not have secondary myelodysplastic syndrome resulting from chemical injury or prior treatment with chemotherapy and/or radiation for other diseases; and
- The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- No evidence of disease progression; and
- The treatment remains appropriate and patient is benefitting from treatment.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
CALCIUM FOLINATE				
Tab 15 mg – PCT – Retail pharmacy-Specialist.....	114.69	10	✓	DBL Leucovorin Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist	17.10	5	✓	Hospira
Inj 10 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.....	7.28	1	✓	Calcium Folate Sandoz
			✓	Calcium Folate Sandoz S29 <small>S29</small>
Inj 50 mg – PCT – Retail pharmacy-Specialist.....	72.80	10	✓	Leucovorin Pharmacia <small>S29</small>
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist	9.49	1	✓	Calcium Folate Sandoz
Inj 100 mg – PCT only – Specialist.....	7.33	1	✓	Calcium Folate Ebewe
	94.90	10	✓	Leucovorin Pharmacia <small>S29</small>
Inj 300 mg – PCT only – Specialist.....	22.51	1	✓	Calcium Folate Ebewe
	25.14		✓	Leucovorin DBL <small>S29</small>
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	✓	Calcium Folate Sandoz
			✓	Calcium Folate Sandoz S29 <small>S29</small>
Inj 1 g – PCT only – Specialist.....	67.51	1	✓	Calcium Folate Ebewe
Inj 10 mg per ml, 100 ml vial – PCT only – Specialist	72.00	1	✓	Calcium Folate Sandoz
Inj 1 mg for ECP – PCT only – Specialist.....	0.06	1 mg	✓	Baxter
CAPECITABINE – Retail pharmacy-Specialist				
Tab 150 mg	10.00	60	✓	Capercit
Tab 500 mg	49.00	120	✓	Capercit
CLADRIBINE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml.....	749.96	1	✓	Litak <small>S29</small>
Inj 1 mg per ml, 10 ml.....	749.96	1	✓	Leustatin
Inj 10 mg for ECP	749.96	10 mg OP	✓	Baxter
CYTARABINE				
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist.....	400.00	5	✓	Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist	41.36	1	✓	Pfizer
Inj 1 mg for ECP – PCT only – Specialist.....	0.25	10 mg	✓	Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Specialist.....	80.00	100 mg OP	✓	Baxter
FLUDARABINE PHOSPHATE				
Tab 10 mg – PCT – Retail pharmacy-Specialist.....	412.00	20	✓	Fludara Oral
Inj 50 mg vial – PCT only – Specialist	576.45	5	✓	Fludarabine Ebewe
Inj 50 mg for ECP – PCT only – Specialist.....	115.29	50 mg OP	✓	Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial – PCT only – Specialist	10.51	1	✓	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial – PCT only – Specialist	29.44	1	✓	Fluorouracil Accord
Inj 1 mg for ECP – PCT only – Specialist	0.62	100 mg	✓	Baxter
GEMCITABINE HYDROCHLORIDE – PCT only – Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	✓	DBL Gemcitabine
Inj 1 g	15.89	1	✓	Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	✓	Baxter
IRINOTECAN HYDROCHLORIDE – PCT only – Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	✓	Accord
	71.44		✓	Irinotecan Actavis 100
	100.00		✓	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	✓	Baxter
MERCAPTOPURINE				
Tab 50 mg – PCT – Retail pharmacy-Specialist	37.00	25	✓	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist – Special Authority see SA1725 below	428.00	100 ml OP	✓	Allmercap
►SA1725 Special Authority for Subsidy				
Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.				
Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.				
METHOTREXATE				
* Tab 2.5 mg – PCT – Retail pharmacy-Specialist	9.98	90	✓	Trexate
* Tab 10 mg – PCT – Retail pharmacy-Specialist	33.71	90	✓	Trexate
* Inj 2.5 mg per ml, 2 ml – PCT – Retail pharmacy-Specialist	47.50	5	✓	Methotrexate DBL
* Inj 7.5 mg prefilled syringe	14.61	1	✓	Methotrexate Sandoz
* Inj 10 mg prefilled syringe	14.66	1	✓	Methotrexate Sandoz
* Inj 15 mg prefilled syringe	14.77	1	✓	Methotrexate Sandoz
* Inj 20 mg prefilled syringe	14.88	1	✓	Methotrexate Sandoz
* Inj 25 mg prefilled syringe	14.99	1	✓	Methotrexate Sandoz
* Inj 30 mg prefilled syringe	15.09	1	✓	Methotrexate Sandoz
* Inj 25 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist	30.00	5	✓	Methotrexate DBL Onco-Vial
* Inj 25 mg per ml, 20 ml vial – PCT – Retail pharmacy-Specialist	45.00	1	✓	DBL Methotrexate Onco-Vial
* Inj 100 mg per ml, 10 ml – PCT – Retail pharmacy-Specialist	25.00	1	✓	Methotrexate Ebewe
* Inj 100 mg per ml, 50 ml vial – PCT – Retail pharmacy-Specialist	79.99	1	✓	Methotrexate Ebewe
* Inj 1 mg for ECP – PCT only – Specialist	0.06	1 mg	✓	Baxter
* Inj 5 mg intrathecal syringe for ECP – PCT only – Specialist	4.73	5 mg OP	✓	Baxter

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PEMETREXED – PCT only – Specialist – Special Authority see SA1679 below				
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter
► SA1679 Special Authority for Subsidy				
Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
Both:				
1 Patient has been diagnosed with mesothelioma; and				
2 Pemetrexed to be administered at a dose of 500 mg/m ² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.				
Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
All of the following:				
1 No evidence of disease progression; and				
2 The treatment remains appropriate and the patient is benefitting from treatment; and				
3 Pemetrexed to be administered at a dose of 500mg/m ² every 21 days for a maximum of 6 cycles.				
Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
Both:				
1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and				
2 Either:				
2.1 Both:				
2.1.1 Patient has chemotherapy-naïve disease; and				
2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m ² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or				
2.2 All of the following:				
2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and				
2.2.2 Patient has not received prior funded treatment with pemetrexed; and				
2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m ² every 21 days for a maximum of 6 cycles.				
Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:				
All of the following:				
1 No evidence of disease progression; and				
2 The treatment remains appropriate and the patient is benefitting from treatment; and				
3 Pemetrexed is to be administered at a dose of 500mg/m ² every 21 days.				
THIOGUANINE – PCT – Retail pharmacy-Specialist				
Tab 40 mg	126.31	25	✓	Lanvis
Other Cytotoxic Agents				
AMSACRINE – PCT only – Specialist				
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓	Amsidine ^{\$29}
	4,736.00		✓	Amsidine ^{\$29}
Inj 75 mg	1,250.00	5	✓	AmsaLyo ^{\$29}
ANAGRELIDE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist				
Cap 0.5 mg	1,175.87	100	✓	Agrylin
ARSENIC TRIOXIDE – PCT only – Specialist				
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓	Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓	Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
BLEOMYCIN SULPHATE – PCT only – Specialist				
Inj 15,000 iu, vial.....	185.16	1	✓	DBL Bleomycin Sulfate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓	Baxter
BORTEZOMIB – PCT only – Specialist – Special Authority see SA1889 below				
Inj 2.5 mg vial	75.00	1	✓	Bortezomib Juno S29
Inj 3.5 mg vial	105.00	1	✓	Bortezomib Dr Reddy's S29
			✓	Bortezomib Dr-Reddy's
			✓	Bortezomib Juno S29
Inj 1 mg for ECP	31.20	1 mg	✓	Baxter
<i>(Bortezomib Juno S29 Inj 2.5 mg vial to be delisted 1 August 2022)</i>				
<i>(Bortezomib Dr Reddy's S29 Inj 3.5 mg vial to be delisted 1 August 2022)</i>				
<i>(Bortezomib Juno S29 Inj 3.5 mg vial to be delisted 1 August 2022)</i>				
►SA1889 Special Authority for Subsidy				
Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:				
Either:				
1 The patient has symptomatic multiple myeloma; or				
2 The patient has symptomatic systemic AL amyloidosis *.				
Note: Indications marked with * are unapproved indications.				
DACARBAZINE – PCT only – Specialist				
Inj 200 mg vial	62.70	1	✓	DBL dacarbazine
	580.60	10	✓	Dacarbazine APP S29
Inj 200 mg for ECP	62.70	200 mg OP	✓	Baxter
DACTINOMYCIN [ACTINOMYCIN D] – PCT only – Specialist				
Inj 0.5 mg vial	255.00	1	✓	Cosmegen
Inj 0.5 mg for ECP	255.00	0.5 mg OP	✓	Baxter
DAUNORUBICIN – PCT only – Specialist				
Inj 2 mg per ml, 10 ml	149.50	1	✓	Pfizer
Inj 20 mg vial	1,495.00	10	✓	Daunorubicin Zentiva S29
Inj 20 mg for ECP	149.50	20 mg OP	✓	Baxter
DOCETAXEL – PCT only – Specialist				
Inj 20 mg.....	48.75	1	✓	Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial.....	46.89	1	✓	DBL Docetaxel
Inj 20 mg per ml, 4 ml vial.....	26.95	1	✓	Docetaxel Accord S29
Inj 80 mg.....	195.00	1	✓	Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓	Baxter

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
DOXORUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial.....	10.00	1	✓	Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial.....	11.50	1	✓	Doxorubicin Ebewe
	17.00		✓	Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial.....	23.00	1	✓	Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial	65.00	1	✓	Arrow-Doxorubicin
	69.99		✓	Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓	Baxter
EPIRUBICIN HYDROCHLORIDE – PCT only – Specialist				
Inj 2 mg per ml, 5 ml vial.....	25.00	1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial.....	30.00	1	✓	Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial.....	99.99	1	✓	Epirubicin Ebewe
Inj 1 mg for ECP	0.50	1 mg	✓	Baxter
ETOPOSIDE				
Cap 50 mg – PCT – Retail pharmacy-Specialist	340.73	20	✓	Vepesid
Cap 100 mg – PCT – Retail pharmacy-Specialist	340.73	10	✓	Vepesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Specialist	7.90	1	✓	Rex Medical
Inj 1 mg for ECP – PCT only – Specialist.....	0.09	1 mg	✓	Baxter
ETOPOSIDE PHOSPHATE – PCT only – Specialist				
Inj 100 mg (of etoposide base).....	40.00	1	✓	Etopophos
Inj 1 mg (of etoposide base) for ECP	0.47	1 mg	✓	Baxter
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pharmacy-Specialist				
Cap 500 mg.....	23.82	100	✓	Devatis
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial – PCT only – Specialist	109.74	1	✓	Zavedos
Inj 10 mg vial – PCT only – Specialist	233.64	1	✓	Zavedos
Inj 1 mg for ECP – PCT only – Specialist.....	25.77	1 mg	✓	Baxter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authority see SA2047 below				
Wastage claimable				
Cap 5 mg.....	5,122.76	28	✓	Revlimid
Cap 10 mg.....	4,655.25	21	✓	Revlimid
	6,207.00	28	✓	Revlimid
Cap 15 mg.....	5,429.39	21	✓	Revlimid
	7,239.18	28	✓	Revlimid
Cap 25 mg.....	7,627.00	21	✓	Revlimid

►SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg – PCT – Retail pharmacy-Specialist.....	314.00	50	✓ Uromitexan
Tab 600 mg – PCT – Retail pharmacy-Specialist.....	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule – PCT only – Specialist.....	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule – PCT only – Specialist.....	407.40	15	✓ Uromitexan
Inj 1 mg for ECP – PCT only – Specialist.....	2.96	100 mg	✓ Baxter

MITOMYCIN C – PCT only – Specialist

Inj 5 mg vial	641.70	1	✓ Accord ^{S29}
Inj 20 mg vial	3,275.00	1	✓ Omegapharm ^{S29}
			✓ Teva
Inj 1 mg for ECP	470.75	1 mg	✓ Baxter

MITOZANTRONE – PCT only – Specialist

Inj 2 mg per ml, 10 ml vial.....	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter

OLAPARIB – Retail pharmacy-Specialist – Special Authority see [SA1883 below](#)

Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg	3,701.00	56	✓ Lynparza

➡ **SA1883** Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment; and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL – PCT only – Specialist

Inj 30 mg.....	47.30	5	✓	Paclitaxel Ebewe
Inj 100 mg.....	24.00	1	✓	Paclitaxel Ebewe
	91.67		✓	Paclitaxel Actavis
Inj 150 mg.....	26.69	1	✓	Paclitaxel Ebewe
	137.50		✓	Anzatax
			✓	Paclitaxel Actavis
Inj 300 mg.....	44.00	1	✓	Paclitaxel Ebewe
	275.00		✓	Anzatax
			✓	Paclitaxel Actavis
Inj 1 mg for ECP.....	0.20	1 mg	✓	Baxter

PEGASPARGASE – PCT only – Special Authority see SA1979 below

Inj 750 iu per ml, 5 ml vial.....	3,455.00	1	✓	Oncaspar LYO ^{\$29}
-----------------------------------	----------	---	---	------------------------------

► **SA1979 Special Authority for Subsidy**

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] – PCT only – Specialist

Inj 10 mg.....	CBS	1	✓	Nipent ^{\$29}
----------------	-----	---	---	------------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-Specialist				
Cap 50 mg	980.00	50	✓	Natulan ^{\$29}
TEMOZOLOMIDE – Special Authority see SA1741 below – Retail pharmacy				
Cap 5 mg	9.13	5	✓	Temaccord
Cap 20 mg	16.38	5	✓	Temaccord
	18.30		✓	Apo-Temozolomide
	136.00	14	✓	Accord ^{\$29}
Cap 100 mg	35.98	5	✓	Temaccord
	40.20		✓	Apo-Temozolomide
	532.00	14	✓	Accord ^{\$29}
Cap 140 mg	50.12	5	✓	Temaccord
	400.00		✓	Amneal ^{\$29}
Cap 180 mg	620.00	14	✓	Accord ^{\$29}
Cap 250 mg	86.34	5	✓	Temaccord
	688.00		✓	Amneal ^{\$29}

►SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE – Retail pharmacy-Specialist – Special Authority see [SA1124 below](#)

Cap 50 mg.....	378.00	28	✓	Thalomid
Cap 100 mg.....	756.00	28	✓	Thalomid

►SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg – PCT – Retail pharmacy-Specialist.....	479.50	100	✓	Vesanoid
---	--------	-----	---	----------

VENETOCLAX – Retail pharmacy-Specialist – Special Authority see [SA1868 below](#)

Tab 14 x 10 mg, 7 x 50 mg, 21 x 100 mg.....	1,771.86	42 OP	✓	Venclexta
Tab 10 mg.....	95.78	14 OP	✓	Venclexta
Tab 50 mg.....	239.44	7 OP	✓	Venclexta
Tab 100 mg – Wastage claimable.....	8,209.41	120	✓	Venclexta

►SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist.....	270.37	5	✓ Hospira
Inj 1 mg for ECP – PCT only – Specialist.....	6.00	1 mg	✓ Baxter

VINCISTINE SULPHATE

Inj 1 mg per ml, 1 ml vial – PCT – Retail pharmacy-Specialist.....	74.52	5	✓ DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist.....	102.73	5	✓ DBL Vincristine Sulfate
Inj 1 mg for ECP – PCT only – Specialist.....	12.60	1 mg	✓ Baxter

VINORELBINE – PCT only – Specialist

Inj 10 mg per ml, 1 ml vial.....	12.00	1	✓ Navelbine
	42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial.....	56.00	1	✓ Navelbine
	210.00		✓ Vinorelbine Ebewe
	328.65		✓ Sagent ^{S29}
Inj 1 mg for ECP.....	1.25	1 mg	✓ Baxter
Inj 50 mg for ECP.....	328.65	50 mg OP	✓ Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

ALECTINIB – Retail pharmacy-Specialist – Special Authority see [SA1870 below](#)

Wastage claimable			
Cap 150 mg.....	7,935.00	224	✓ Alecensa

➡SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

ALK test; and

3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB – Special Authority see [SA1805 below](#) – Retail pharmacy

Wastage claimable

Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

► **SA1805** Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 Both:

- 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
- 1.2 Maximum dose of 140 mg/day; or

2 Both:

- 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
- 2.2 Maximum dose of 140 mg/day; or

3 All of the following:

- 3.1 The patient has a diagnosis of CML in chronic phase; and
- 3.2 Maximum dose of 100 mg/day; and
- 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
 - 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up
[https://www.cancertrialsnz.ac.nz/kiss/](https://www.cancertrials.nz.ac.nz/kiss/)

ERLOTINIB – Retail pharmacy-Specialist – Special Authority see [SA2115 below](#)

Tab 100 mg	764.00	30	✓ Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

► **SA2115** Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 4 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Either:
 - 3.1 Patient is treatment naive; or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB – Retail pharmacy-Specialist – Special Authority see [SA2116 below](#)

Tab 250 mg	1,700.00	30	✓ Iressa
------------------	----------	----	----------

➔SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

Tab 100 mg – [Xpharm] – Special Authority see [SA1460 on the next page](#).....

	2,400.00	60	✓ Glivec
* Cap 100 mg	58.23	60	✓ Imatinib-Rex
* Cap 400 mg	84.79	30	✓ Imatinib-Rex

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990
 Pharmac Facsimile: (04) 916 7571
 PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz
 Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- Maximum dose of 400 mg/day.
- Applications to be made and subsequent prescriptions can be written by an oncologist.
- Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE – Special Authority see [SA2035 below](#) – Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

Tab 250 mg	1,899.00	70	✓ Tykerb
------------------	----------	----	----------

►SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- Lapatinib not to be given in combination with trastuzumab; and
- Lapatinib to be discontinued at disease progression.

NILOTINIB – Special Authority see [SA1489 below](#) – Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	✓ Tasigna
Cap 200 mg	6,532.00	120	✓ Tasigna

►SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- Either:
 - Patient has documented CML treatment failure* with imatinib; or
 - Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- Maximum nilotinib dose of 800 mg/day; and
- Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- Maximum nilotinib dose of 800 mg/day; and
- Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
PALBOCICLIB – Retail pharmacy-Specialist – Special Authority see SA1894 below				
Wastage claimable				
Tab 75 mg	4,000.00	21	✓	Ibrance
Tab 100 mg	4,000.00	21	✓	Ibrance
Tab 125 mg	4,000.00	21	✓	Ibrance

►SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:
 - second or subsequent line setting
 - 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
 - 4.2 Both:
 - first line setting
 - 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB – Special Authority see [SA1190 below](#) – Retail pharmacy

Tab 200 mg	1,334.70	30	✓	Votrient
Tab 400 mg	2,669.40	30	✓	Votrient

►SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB – Special Authority see [SA1890 below](#) – Retail pharmacy

Wastage claimable

Tab 5 mg	2,500.00	56	✓ Jakavi
Tab 10mg	5,000.00	56	✓ Jakavi
Tab 15 mg	5,000.00	56	✓ Jakavi
Tab 20 mg	5,000.00	56	✓ Jakavi

► **SA1890** **Special Authority for Subsidy**

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
SUNITINIB – Special Authority see SA2117 below – Retail pharmacy				
Cap 12.5 mg.....	208.38	28	✓	Sunitinib Pfizer
	2,315.38		✓	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022				
Cap 25 mg.....	416.77	28	✓	Sunitinib Pfizer
	4,630.77		✓	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022				
Cap 50 mg.....	694.62	28	✓	Sunitinib Pfizer
	9,261.54		✓	Sutent
Sunitinib Pfizer to be Principal Supply on 1 July 2022				
<i>(Sutent Cap 12.5 mg to be delisted 1 July 2022)</i>				
<i>(Sutent Cap 25 mg to be delisted 1 July 2022)</i>				
<i>(Sutent Cap 50 mg to be delisted 1 July 2022)</i>				

► **SA2117** Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naïve; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

1 Any of the following:

- 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
- 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and

2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES – refer to HORMONE PREPARATIONS, Tropic Hormones, [page 83](#)

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see [SA2118 below](#)

Wastage claimable

Tab 250 mg4,276.19 120 ✓ Zytiga

➔ [SA2118](#) Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:

4.1 All of the following:

- 4.1.1 Patient is symptomatic; and
- 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and

continued...

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
MEGESTROL ACETATE – Subsidy by endorsement				
Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.				
Tab 160 mg	48.80	30	✓	Megace S29
<i>(Megace S29 Tab 160 mg to be delisted 1 February 2023)</i>				
OCTREOTIDE				
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓	Max Health
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓	Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓	Max Health
OCTREOTIDE LONG-ACTING – Special Authority see SA2119 below – Retail pharmacy				
Inj depot 10 mg prefilled syringe.....	439.97	1	✓	Octreotide Depot <u>Teva</u>
Inj depot 20 mg prefilled syringe.....	647.03	1	✓	Octreotide Depot <u>Teva</u>
Inj depot 30 mg prefilled syringe.....	718.55	1	✓	Octreotide Depot <u>Teva</u>

►SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*;
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed; or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas - for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

* Tab 10 mg	15.00	60	✓ Tamoxifen Sandoz
* Tab 20 mg	6.65	60	✓ Tamoxifen Sandoz

Aromatase Inhibitors

ANASTROZOLE

* Tab 1 mg	4.55	30	✓ Anatrole
------------------	------	----	-------------------

EXEMESTANE

* Tab 25 mg	14.50	30	✓ Pfizer Exemestane
-------------------	-------	----	----------------------------

LETROZOLE

* Tab 2.5 mg	5.84	30	✓ Letrole
--------------------	------	----	------------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE

* Tab 25 mg	7.35	60	✓	Azamun
* Tab 50 mg	7.60	100	✓	Azamun
* Inj 50 mg vial	199.00	1	✓	Imuran

(Imuran Inj 50 mg vial to be delisted 1 January 2023)

MYCOPHENOLATE MOFETIL

Tab 500 mg	35.90	50	✓	Cellcept
Cap 250 mg	35.90	100	✓	Cellcept
Powder for oral liq 1 g per 5 ml – Subsidy by endorsement	187.25	165 ml OP	✓	Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT – Special Authority see SA2103 below – Retail pharmacy

Inj 25 mg	690.00	4	✓	Enbrel
Inj 25 mg autoinjector	690.00	4	✓	Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓	Enbrel
Inj 50 mg prefilled syringe	1,050.00	4	✓	Enbrel

➔SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the HML rules; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
 - 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Both:
- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

the patient continues with etanercept treatment; and

- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and

- 1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or

- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or

- 2 All of the following:

- 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and

- 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and

- 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and

- 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and

- 2.5 Either:

- 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

- 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

- 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm

25-34 years - Male: 7.5 cm; Female: 5.5 cm

35-44 years - Male: 6.5 cm; Female: 4.5 cm

45-54 years - Male: 6.0 cm; Female: 5.0 cm

55-64 years - Male: 5.5 cm; Female: 4.0 cm

65-74 years - Male: 4.0 cm; Female: 4.0 cm

75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:

- 1.1 Applicant is a rheumatologist; or

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and

- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and

- 1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or

- 2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

- 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and

- 2.3 Any of the following:

- 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

- 2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and

- 1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
- 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or

- 2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and

- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and

- 2.3 Any of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or

2 All of the following:

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a dermatologist; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and

3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) – PCT only – Specialist

Inj 50 mg per ml, 5 ml.....	2,774.48	5	✓ ATGAM
-----------------------------	----------	---	---------

BACILLUS CALMETTE-GUERIN (BCG) VACCINE – PCT only – Specialist

Subsidised only for bladder cancer.

Inj 2-8 × 100 million CFU.....	149.37	1	✓ OncoTICE
--------------------------------	--------	---	------------

Inj 40 mg per ml, vial.....	176.90	3	✓ SI-Onco-BCG <small>\$29</small>
-----------------------------	--------	---	-----------------------------------

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) – Special Authority see [SA2102 below](#) – Retail pharmacy

Inj 20 mg per 0.4 ml prefilled syringe.....	190.00	1	✓ Amgevita
---	--------	---	------------

Inj 40 mg per 0.8 ml prefilled pen.....	375.00	2	✓ Amgevita
---	--------	---	------------

Inj 40 mg per 0.8 ml prefilled syringe.....	375.00	2	✓ Amgevita
---	--------	---	------------

► [SA2102](#) Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
- 2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
- 3 Patient has 3 or more active lesions; and
- 4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

- Either:
- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
 - 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
- 1 Patient has pyoderma gangrenosum*; and
 - 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.3 Patient has complex peri-anal fistula; and
- 3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
- 1.2 Either:

- 1.2.1 The patient has experienced intolerable side effects; or
- 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or

2 All of the following:

- 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
- 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
- 2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and
- 2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
- 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
- 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or

2 All of the following:

- 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
- 2.3 Either:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects; or
 - 1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
 - 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- Either:
- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis; or
 - 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2 Patient's PUCAI score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
- 3 Any of the following:
 - 3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has axial inflammatory pain for six months or more; and
- 3 Patient is unable to take NSAIDs; and
- 4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
- 5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
- 6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
- 2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
- 3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
- 5 Any of the following:
 - 5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA) – Special Authority see [SA2101 on the next page](#) – Retail pharmacy

Inj 20 mg per 0.4 ml prefilled syringe.....	1,599.96	2	✓ Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe.....	1,599.96	2	✓ Humira

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

►SA2101 Special Authority for Subsidy

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Either:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 CDAI score is 150 or less; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

2.2 Both:

- 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

2.1 Either:

- 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2.1.2 PCDAI score is 15 or less; or

2.2 Both:

- 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

- 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and

4 Either:

- 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
- 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Either:

- 1.1 Applicant is a dermatologist; or
- 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and

2 Either:

2.1 Both:

2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

2.1.2 Either:

2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or

2.2 Both:

2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and

2.2.2 Either:

2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or

2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and

3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

- 1.1 The patient has had a good clinical response following 3 initial doses; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and

- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

ALIBERCEPT – Special Authority see [SA1772 below](#) – Retail pharmacy

Inj 40 mg per ml, 0.1 ml vial.....	1,250.00	1	✓ Eylea
------------------------------------	----------	---	---------

► [SA1772](#) Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months; or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 – 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
--	---	----------------------------	-------------------------------------

continued...

- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CASIRIVIMAB AND IMDEVIMAB – [Xpharm] – Special Authority see [SA2096 below](#)

Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 120 mg per ml imdevimab, 11.1 ml vial (1).....	0.00	1 OP	✓ Ronapreve
--	------	------	--------------------

►SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the [Ministry of Health Website](#)

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB – PCT only – Specialist – Special Authority see [SA1697 below](#)

Inj 5 mg per ml, 20 ml vial.....	364.00	1	✓ Erbitux
Inj 5 mg per ml, 100 ml vial.....	1,820.00	1	✓ Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

►SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

INFLIXIMAB – PCT only – Special Authority see [SA2082 below](#)

Inj 100 mg.....	806.00	1	✓ Remicade
Inj 1 mg for ECP	8.29	1 mg	✓ Baxter

►SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
- 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist.

Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoidosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Either:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist.

Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
- 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or

2 All of the following:

- 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.2 Ankylosing spondylitis; or
- 2.3 Psoriatic arthritis; or
- 2.4 Severe ocular inflammation; or
- 2.5 Chronic ocular inflammation; or
- 2.6 Crohn's disease (adults); or
- 2.7 Crohn's disease (children); or
- 2.8 Fistulising Crohn's disease; or
- 2.9 Severe fulminant ulcerative colitis; or
- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis; or
- 2.12 Neurosarcoidosis; or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

2.2 Any of the following:

- 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
- 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
- 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB – Special Authority see [SA1896 below](#) – Retail pharmacy

Inj 100 mg prefilled pen	1,638.00	1	✓ Nucala
Inj 100 mg vial	1,638.00	1	✓ Nucala

➔SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded; and
- 4 Patient has a blood eosinophil count of greater than 0.5×10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB – PCT only – Specialist – Special Authority see [SA1627 on the next page](#)

Inj 25 mg per ml, 40 ml vial.....	5,910.00	1	✓ Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
		✓	

►SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL; and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to $1.5 \times 10^9/L$ and platelets greater than or equal to $75 \times 10^9/L$.

OMALIZUMAB – Special Authority see SA1744 below – Retail pharmacy

Inj 150 mg prefilled syringe.....	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

►SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or formoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
- 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
- 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB – PCT only – Specialist – Special Authority see [SA2128 below](#)

Inj 100 mg per ml, 1 ml vial..... 1,700.00 1 ✓ Synagis

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

► **SA2128** Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - 2.2.2.2 Either:
 - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2.2 Patient is Māori or any Pacific ethnicity; or

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

2.2.3 Both:

2.2.3.1 Patient has haemodynamically significant heart disease; and

2.2.3.2 Any of the following:

2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a); or

2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or

2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or

2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

- Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient will require surgical palliation/definitive repair within the next 3 months.
- Mean pulmonary artery pressure more than 45 mmHg.
- LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician.

Approvals valid for 6 months where patient still meets initial criteria.

PERTUZUMAB – PCT only – Specialist – Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial.....	3,927.00	1	✓ Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	✓ Baxter

►SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- Either:
 - Patient is chemotherapy treatment naïve; or
 - Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- The patient has good performance status (ECOG grade 0-1); and
- Pertuzumab to be administered in combination with trastuzumab; and
- Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) – PCT only – Specialist – Special Authority see SA1976 below

Inj 100 mg per 10 ml vial	1,075.50	2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

►SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Either:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) – PCT only – Specialist – Special Authority see [SA2114 below](#)

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

► **SA2114** **Special Authority for Subsidy**

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

3.4 Patient is a female of child-bearing potential; or

3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

1 Patient has been diagnosed with ANCA associated vasculitis*; and

2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and

3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and

2 Any of the following:

2.1 The patient is rituximab treatment naive; or

2.2 Either:

2.2.1 The patient is chemotherapy treatment naive; or

2.2.2 Both:

2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and

2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or

2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and

3 The patient has good performance status; and

4 Either:

4.1 The patient does not have chromosome 17p deletion CLL; or

4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and

5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and

6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Either:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
- 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m² administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m² administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Either:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*;
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks; and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

3 Either:

- 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not been effective at controlling active disease; or
- 3.2 Rapid treatment is required due to life threatening complications; and

4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not been effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis; and
- 2 Either:
 - 2.1 Both:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
- 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
- 2.2 Rapid treatment is required due to life threatening complications; and

- 3 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m² of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m² every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m²; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note); and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m² of body surface area per week for a total of 4 weeks.

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
- 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*;
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensitisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogeneic stem cell transplant*;
- 2 Patient would receive no more than two doses at 375 mg/m² of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

skin ulceration and reduction in corticosteroid requirement; and

2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB – Special Authority see [SA2084 below](#) – Retail pharmacy

Inj 150 mg per ml, 1 ml prefilled syringe.....	799.50	1	✓ Cosentyx
	1,599.00	2	✓ Cosentyx

►[SA2084](#) Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule, for severe chronic plaque psoriasis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or

continued...

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and

- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
--	---	----------------------------	-------------------------------------

continued...

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and

2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB – Special Authority see [SA1596 below](#) – Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	✓ Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

►SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TOCILIZUMAB – PCT only – Special Authority see [SA2100 below](#)

Inj 20 mg per ml, 4 ml vial.....	220.00	1	✓ Actemra
			✓ Actemra S29 S29
	880.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial.....	550.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial.....	1,100.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
	4,400.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	2.85	1 mg	✓ Baxter

►SA2100 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

1 All of the following:

- 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
- 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
- 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or

2 All of the following:

- 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
- 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
- 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a DHB hospital in accordance with the General Rules of the Pharmaceutical Schedule; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
- 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with baricitinib.

Note: Indications marked with * are unapproved indications.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

continued...

recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB – PCT only – Specialist – Special Authority see [SA1632 below](#)

Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	9.36	1 mg	✓ Baxter

► **SA1632** Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Either:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE – PCT only – Specialist – Special Authority see [SA1871 below](#)

Inj 100 mg vial	2,320.00	1	✓ Kadcyła
Inj 160 mg vial	3,712.00	1	✓ Kadcyła
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

►SA1871 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Treatment to be discontinued at disease progression.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

NIVOLUMAB – PCT only – Specialist – Special Authority see [SA2120 below](#)

Inj 10 mg per ml, 4 ml vial.....	1,051.98	1	✓ Opdivo
Inj 10 mg per ml, 10 ml vial.....	2,629.96	1	✓ Opdivo
Inj 1 mg for ECP	27.62	1 mg	✓ Baxter

➡SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks.

Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB – PCT only – Specialist – Special Authority see [SA2121](#) below

Inj 25 mg per ml, 4 ml vial.....	4,680.00	1	✓	Keytruda
Inj 1 mg for ECP.....	49.14	1 mg	✓	Baxter

►SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks.

Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target) must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN

Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral

EVEROLIMUS – Special Authority see [SA2008 below](#) – Retail pharmacy

Wastage claimable			
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

►SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
SIROLIMUS – Special Authority see SA2005 below – Retail pharmacy				
Tab 1 mg	749.99	100	✓	Rapamune
Tab 2 mg	1,499.99	100	✓	Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓	Rapamune

►SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR < 30 ml/min; or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- HUS or TTP; or
- Leukoencephalopathy; or
- Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomylipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS – Special Authority see [SA1745 below](#) – Retail pharmacy

Cap 0.5 mg.....	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg.....	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg.....	84.30	100	✓ Tacrolimus Sandoz
Cap 5 mg.....	248.20	50	✓ Tacrolimus Sandoz

► **SA1745** Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

- Both:
- 1 Patient requires long-term systemic immunosuppression; and
 - 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see [SA2079 on the next page](#) – Retail pharmacy

Tab 15 mg	1,271.00	28	✓ RINVOQ
-----------------	----------	----	----------

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

➤SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a DHB hospital in accordance with the Section H rules; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist.

Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see [SA1558 below](#) – Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe.....	2,668.00	1	✓ Firazyr
---	----------	---	------------------

► **SA1558** **Special Authority for Subsidy**

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

► **SA1367** **Special Authority for Subsidy**

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT – Special Authority see [SA1367 above](#) – Retail pharmacy

Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX <small>\$29</small>
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX <small>\$29</small>
Maintenance kit - 6 vials 120 mcg freeze dried venom, with diluent	285.00	1 OP	✓ Venomil <small>\$29</small>
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent 9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera <small>\$29</small>

WASP VENOM ALLERGY TREATMENT – Special Authority see [SA1367 above](#) – Retail pharmacy

Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera <small>\$29</small>
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Venomil <small>\$29</small>
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera <small>\$29</small>
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Venomil <small>\$29</small>

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1.12	100	✓ Zista	
* Oral liq 1 mg per ml	2.84	200 ml	✓ Histaclear	
CHLORPHENIRAMINE MALEATE				
* Oral liq 2 mg per 5 ml	9.37	500 ml	✓ Histafen	
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg	2.02	40		
	(8.40)			Polaramine
	1.01	20		
	(5.99)			Polaramine
* Oral liq 2 mg per 5 ml	1.77	100 ml		
	(10.29)			Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
	(8.23)			Telfast
* Tab 120 mg	4.74	10		
	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
LORATADINE				
* Tab 10 mg	1.69	100	✓ Lorafix	
* Oral liq 1 mg per ml	1.43	100 ml	✓ Haylor syrup	
PROMETHAZINE HYDROCHLORIDE				
* Tab 10 mg	1.39	50	✓ Allersoothe	
Allersoothe to be Principal Supply on 1 September 2022				
* Tab 25 mg	1.58	50	✓ Allersoothe	
Allersoothe to be Principal Supply on 1 September 2022				
* Oral liq 1 mg per 1 ml	3.39	100 ml	✓ Allersoothe	
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO	17.87	5	✓ Hospira	
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01	200 dose OP	✓ Qvar	
Aerosol inhaler, 50 mcg per dose CFC-free	8.54	200 dose OP	✓ Beclazone 50	
Aerosol inhaler, 100 mcg per dose	17.52	200 dose OP	✓ Qvar	
Aerosol inhaler, 100 mcg per dose CFC-free	12.50	200 dose OP	✓ Beclazone 100	
Aerosol inhaler, 250 mcg per dose CFC-free	22.67	200 dose OP	✓ Beclazone 250	
BUDESONIDE				
Powder for inhalation, 100 mcg per dose	17.00	200 dose OP	✓ Pulmicort Turbuhaler	
Powder for inhalation, 200 mcg per dose	19.00	200 dose OP	✓ Pulmicort Turbuhaler	
Powder for inhalation, 400 mcg per dose	32.00	200 dose OP	✓ Pulmicort Turbuhaler	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
FLUTICASONE				
Aerosol inhaler, 50 mcg per dose.....	7.19	120 dose OP	✓	Flixotide
Powder for inhalation, 50 mcg per dose.....	7.50	60 dose OP	✓	Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose.....	7.50	60 dose OP	✓	Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose.....	13.60	120 dose OP	✓	Flixotide
Aerosol inhaler, 250 mcg per dose.....	24.62	120 dose OP	✓	Flixotide
Powder for inhalation, 250 mcg per dose.....	13.60	60 dose OP	✓	Flixotide Accuhaler

Inhaled Long-acting Beta-adrenoceptor Agonists

EFORMOTEROL FUMARATE

Powder for inhalation, 12 mcg per dose, and monodose device.....	20.64 (35.80)	60 dose		Foradil
--	------------------	---------	--	---------

EFORMOTEROL FUMARATE DIHYDRATE

Powder for inhalation 4.5 mcg per dose, breath activated (equivalent to eformoterol fumarate 6 mcg metered dose).....	10.32 (16.90)	60 dose OP		Oxis Turbuhaler
--	------------------	------------	--	-----------------

INDACATEROL

Powder for inhalation 150 mcg.....	61.00	30 dose OP	✓	Onbrez Breezhaler
Powder for inhalation 300 mcg.....	61.00	30 dose OP	✓	Onbrez Breezhaler

SALMETEROL

Aerosol inhaler CFC-free, 25 mcg per dose.....	26.25	120 dose OP	✓	Serevent
Powder for inhalation, 50 mcg per dose, breath activated	26.25	60 dose OP	✓	Serevent Accuhaler

Inhaled Corticosteroids with Long-Acting Beta-Adrenoceptor Agonists

BUDESONIDE WITH EFORMOTEROL

Powder for inhalation 160 mcg with 4.5 mcg eformoterol fumarate per dose (equivalent to 200 mcg budesonide with 6 mcg eformoterol fumarate metered dose)	41.50	120 dose OP	✓	DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumarate per dose (equivalent to 400 mcg budesonide with 12 mcg eformoterol fumarate metered dose) – No more than 2 dose per day.....	82.50	120 dose OP	✓	DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg.....	18.23	120 dose OP	✓	Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 mcg	33.74	120 dose OP	✓	Symbicort Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg.....	21.40	120 dose OP	✓	Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 mcg	44.08	120 dose OP	✓	Symbicort Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day	44.08	60 dose OP	✓	Symbicort Turbuhaler 400/12

FLUTICASONE FUROATE WITH VILANTEROL

Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓	Breo Ellipta
--	-------	------------	---	---------------------

RESPIRATORY SYSTEM AND ALLERGIES

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
FLUTICASONE WITH SALMETEROL				
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose OP	✓	<u>Seretide</u>
Aerosol inhaler 125 mcg with salmeterol 25 mcg	32.60	120 dose OP	✓	<u>Seretide</u>
Powder for inhalation 100 mcg with salmeterol 50 mcg – No more than 2 dose per day	33.74	60 dose OP	✓	<u>Seretide Accuhaler</u>
Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day	44.08	60 dose OP	✓	<u>Seretide Accuhaler</u>

Beta-Adrenoceptor Agonists

SALBUTAMOL				
Oral liq 400 mcg per ml	40.00	150 ml	✓	<u>Ventolin</u>
Infusion 1 mg per ml, 5 ml	118.38	10	✓	<u>Ventolin</u>
Inj 500 mcg per ml, 1 ml – Up to 5 inj available on a PSO	53.00	5	✓	<u>Ventolin</u>

Inhaled Beta-Adrenoceptor Agonists

SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free – Up to 1000 dose available on a PSO	3.80	200 dose OP	✓	<u>Respigen</u>
	(6.20)		✓	<u>SalAir</u> Ventolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	8.96	20	✓	<u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO	9.43	20	✓	<u>Asthalin</u>
TERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	✓	<u>Bricanyl Turbuhaler</u>

Anticholinergic Agents

IPRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose available on a PSO	16.20	200 dose OP	✓	<u>Atrovent</u>
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 neb available on a PSO	11.73	20	✓	<u>Univent</u>

Inhaled Beta-Adrenoceptor Agonists with Anticholinergic Agents

SALBUTAMOL WITH IPRATROPIUM BROMIDE				
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg per dose CFC-free	12.19	200 dose OP	✓	<u>Duolin HFA</u>
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	11.04	20	✓	<u>Duolin</u>

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
	✓	

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM – Subsidy by endorsement

- Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium.
- Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Powder for inhalation 50 mcg per dose 61.00 30 dose OP ✓ **Seebri Breezhaler**

TIOTROPIUM BROMIDE – Subsidy by endorsement

- Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

Powder for inhalation, 18 mcg per dose 50.37 30 dose ✓ **Spiriva**
Soln for inhalation 2.5 mcg per dose 50.37 60 dose OP ✓ **Spiriva Respimat**

UMECLIDINIUM – Subsidy by endorsement

- Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Powder for inhalation 62.5 mcg per dose 61.50 30 dose OP ✓ **Incruse Ellipta**

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

►SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg 81.00 30 dose OP ✓ **Ultibro Breezhaler**

TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg 81.00 60 dose OP ✓ **Spiolto Respimat**

UMECLIDINIUM WITH VILANTEROL – Special Authority see [SA1584 above](#) – Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg 77.00 30 dose OP ✓ **Anoro Ellipta**

Antifibrotics

NINTEDANIB – Special Authority see [SA2012 on the next page](#) – Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

Cap 100 mg 2,554.00 60 OP ✓ **Ofev**
Cap 150 mg 3,870.00 60 OP ✓ **Ofev**

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

➤SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Tab 801 mg	3,645.00	90	✓ Esbriet
Tab 267 mg	1,215.00	90	✓ Esbriet

➤SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:
All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

Leukotriene Receptor Antagonists

MONTELUKAST

* Tab 4 mg	4.25	28	✓ Montelukast Mylan
* Tab 5 mg	4.25	28	✓ Montelukast Mylan
* Tab 10 mg	3.95	28	✓ Montelukast Mylan

Methylxanthines

AMINOPHYLLINE

* Inj 25 mg per ml, 10 ml ampoule – Up to 5 inj available on a PSO	180.00	5	✓ DBL Aminophylline
---	--------	---	----------------------------

THEOPHYLLINE

* Tab long-acting 250 mg.....	23.02	100	✓ Nuelin-SR
* Oral liq 80 mg per 15 ml	16.60	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA – Special Authority see [SA1978 below](#) – Retail pharmacy

Nebuliser soln, 2.5 mg per 2.5 ml ampoule	250.00	6	✓ Pulmozyme
---	--------	---	--------------------

► **SA1978** Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period; or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR – PCT only – Specialist – Special Authority see [SA2017 below](#)

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

► **SA2017** Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N)

continued...

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

and S549R) in the CFTR gene on at least 1 allele; and

- 3 Patients must have a sweat chloride value of at least 60 mmol/L by quantitative pilocarpine iontophoresis or by Macroduct sweat collection system; and
- 4 Treatment with ivacaftor must be given concomitantly with standard therapy for this condition; and
- 5 Patient must not have an acute upper or lower respiratory infection, pulmonary exacerbation, or changes in therapy (including antibiotics) for pulmonary disease in the last 4 weeks prior to commencing treatment with ivacaftor; and
- 6 The dose of ivacaftor will not exceed one tablet or one sachet twice daily; and
- 7 Applicant has experience and expertise in the management of cystic fibrosis.

SODIUM CHLORIDE

Not funded for use as a nasal drop.

Soln 7%24.50 90 ml OP ✓ **Biomed**

Nasal Preparations

Allergy Prophylactics

BUDESONIDE

Metered aqueous nasal spray, 50 mcg per dose2.54 200 dose OP ✓ **SteroClear**

Metered aqueous nasal spray, 100 mcg per dose2.84 200 dose OP ✓ **SteroClear**

FLUTICASONE PROPIONATE

Metered aqueous nasal spray, 50 mcg per dose1.98 120 dose OP ✓ **Flixonase Hayfever
& Allergy**

IPRATROPIUM BROMIDE

Aqueous nasal spray, 0.03%.....5.23 15 ml OP ✓ **Univent**

Respiratory Devices

MASK FOR SPACER DEVICE

a) Up to 50 dev available on a PSO

b) Only on a PSO

c) Only for children aged six years and under

Small.....2.20 1 ✓ **e-chamber Mask**

PEAK FLOW METER

a) Up to 25 dev available on a PSO

b) Only on a PSO

Low range.....9.54 1 ✓ **Mini-Wright AFS
Low Range**

Normal range.....9.54 1 ✓ **Mini-Wright
Standard**

SPACER DEVICE

a) Up to 50 dev available on a PSO

b) Only on a PSO

220 ml (single patient)2.95 1 ✓ **e-chamber Turbo**

510 ml (single patient)5.12 1 ✓ **e-chamber La
Grande**

800 ml.....6.50 1 ✓ **Volumatic**

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

Respiratory Stimulants

CAFFEINE CITRATE
Oral liq 20 mg per ml (10 mg base per ml)..... 15.10 25 ml OP ✓ **Biomed**

Ear Preparations

FLUMETASONE PIVALATE

Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's ✓ Locorten-Vioform
--	------	-----------	---

TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCIN AND NYSTATIN

Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
--	------	-----------	-------------------

Ear/Eye Preparations

DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN

Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and gramicidin 50 mcg per ml	4.50 (9.27)	8 ml OP	Sofradex
---	----------------	---------	----------

FRAMYCETIN SULPHATE

Ear/Eye drops 0.5%	4.13 (8.65)	8 ml OP	Soframycin
--------------------------	----------------	---------	------------

Eye Preparations

Eye preparations are only funded for use in the eye, unless explicitly stated otherwise.

Anti-Infective Preparations

ACICLOVIR

* Eye oint 3%	14.88	4.5 g OP	✓ ViruPOS
---------------------	-------	----------	------------------

CHLORAMPHENICOL

Eye oint 1%	1.55	5 g OP	✓ Devatis
Eye drops 0.5%	1.54	10 ml OP	✓ Chlorafast

Funded for use in the ear*. Indications marked with * are unapproved indications.

CIPROFLOXACIN

Eye drops 0.3% – Subsidy by endorsement	9.73	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis or severe bacterial conjunctivitis resistant to chloramphenicol; or for the second line treatment of chronic suppurative otitis media (CSOM)*; and the prescription is endorsed accordingly.			
Note: Indication marked with a * is an unapproved indication.			

GENTAMICIN SULPHATE

Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
----------------------	-------	---------	-------------------

PROPAMIDINE ISETHIONATE

* Eye drops 0.1%	2.97 (14.55)	10 ml OP	Brolene
------------------------	-----------------	----------	---------

SODIUM FUSIDATE [FUSIDIC ACID]

Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
--------------------	------	--------	----------------------

TOBRAMYCIN

Eye oint 0.3%	10.45	3.5 g OP	✓ Tobrex
Eye drops 0.3%	11.48	5 ml OP	✓ Tobrex

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
Corticosteroids and Other Anti-Inflammatory Preparations			
DEXAMETHASONE			
* Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
* Eye drops 0.1%	4.50	5 ml OP	✓ Maxidex
Ocular implant 700 mcg – Special Authority see SA1680 below			
– Retail pharmacy.....	1,444.50	1	✓ Ozurdex
» SA1680 Special Authority for Subsidy			
Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:			
All of the following:			
1 Patient has diabetic macular oedema with pseudophakic lens; and			
2 Patient has reduced visual acuity of between 6/9 - 6/48 with functional awareness of reduction in vision; and			
3 Either:			
3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or			
3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and			
4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.			
Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:			
Both:			
1 Patient's vision is stable or has improved (prescriber determined); and			
2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.			
Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:			
All of the following:			
1 Patient has diabetic macular oedema; and			
2 Patient has reduced visual acuity of between 6/9 - 6/48 with functional awareness of reduction in vision; and			
3 Patient is of child bearing potential and has not yet completed a family; and			
4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.			
Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:			
All of the following:			
1 Patient's vision is stable or has improved (prescriber determined); and			
2 Patient is of child bearing potential and has not yet completed a family; and			
3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.			
DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE			
* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g.....	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM			
Eye drops 0.1%	8.80	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE			
* Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20		✓ Flucon

SENSORY ORGANS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71 (10.34)	4 ml OP		Livostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓	Lomide
PREDNISOLONE ACETATE				
Eye drops 1%	5.93 7.00	10 ml OP 5 ml OP	✓	Prednisolone-AFT Pred Forte
PREDNISOLONE SODIUM PHOSPHATE – Special Authority see SA1715 below – Retail pharmacy				
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	✓	Minims Prednisolone

► [SA1715](#) Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE

Eye drops 2%	1.79	5 ml OP	✓	Rexacrom
--------------------	------	---------	---	-----------------

Glaucoma Preparations - Beta Blockers

BETAXOLOL

* Eye drops 0.25%	11.80	5 ml OP	✓	Betoptic S
* Eye drops 0.5%	7.50	5 ml OP	✓	Betoptic

TIMOLOL

* Eye drops 0.25%	1.81	5 ml OP	✓	Arrow-Timolol
* Eye drops 0.5%	2.04	5 ml OP	✓	Arrow-Timolol
* Eye drops 0.5%, gel forming	3.78	2.5 ml OP	✓	Timoptol XE

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

ACETAZOLAMIDE

* Tab 250 mg	17.03	100	✓	Diamox
--------------------	-------	-----	---	---------------

BRINZOLAMIDE

* Eye drops 1%	7.30	5 ml OP	✓	Azopt
----------------------	------	---------	---	--------------

DORZOLAMIDE HYDROCHLORIDE

* Eye drops 2%	9.77 (17.44)	5 ml OP		Trusopt
----------------------	-----------------	---------	--	---------

DORZOLAMIDE WITH TIMOLOL

* Eye drops 2% with timolol 0.5%	2.73	5 ml OP	✓	Dortimopt
--	------	---------	---	------------------

Glaucoma Preparations - Prostaglandin Analogues

BIMATOPROST

* Eye drops 0.03%	5.95	3 ml OP	✓	Bimatoprost Multichem
-------------------------	------	---------	---	--

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
LATANOPROST				
* Eye drops 0.005%	1.82	2.5 ml OP	✓	Teva
TRAVOPROST				
* Eye drops 0.004%	9.75	2.5 ml OP	✓	Travatan

Glaucoma Preparations - Other

BRIMONIDINE TARTRATE				
* Eye drops 0.2%	4.29	5 ml OP	✓	Arrow-Brimonidine
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE				
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓	Combigan
LATANOPROST WITH TIMOLOL				
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓	Arrow - Lattim
PILOCARPINE HYDROCHLORIDE				
* Eye drops 1%	4.26	15 ml OP	✓	Isopto Carpine
* Eye drops 2%	5.35	15 ml OP	✓	Isopto Carpine
* Eye drops 4%	7.99	15 ml OP	✓	Isopto Carpine
Subsidised for oral use pursuant to the Standard Formulae.				
* Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓	Minims Pilocarpine

► [SA0895](#) Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:
Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be “tools of trade” and are not approved as special authority items.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics

ATROPINE SULPHATE				
* Eye drops 1%	17.36	15 ml OP	✓	Atrop
CYCLOPENTOLATE HYDROCHLORIDE				
* Eye drops 1%	8.76	15 ml OP	✓	Cyclogyl
TROPICAMIDE				
* Eye drops 0.5%	7.15	15 ml OP	✓	Mydriacyl
* Eye drops 1%	8.66	15 ml OP	✓	Mydriacyl

Preparations for Tear Deficiency

For acetylcysteine eye drops refer Standard Formulae, [page 243](#)

HYPROMELLOSE				
* Eye drops 0.5%	19.50	15 ml OP	✓	Methopt
HYPROMELLOSE WITH DEXTRAN				
* Eye drops 0.3% with dextran 0.1%	2.30	15 ml OP	✓	Poly-Tears

Preservative Free Ocular Lubricants

►SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see [SA2134 above](#) – Retail pharmacy

Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
----------------------------------	------	----	-------------------

MACROGOL 400 AND PROPYLENE GLYCOL – Special Authority see [SA2134 above](#) – Retail pharmacy

Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	✓ Systane Unit Dose
--	------	----	----------------------------

SODIUM HYALURONATE [HYALURONIC ACID] – Special Authority see [SA2134 above](#) – Retail pharmacy

Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hilo-Fresh
-----------------------------	-------	----------	---------------------

Hilo-Fresh has a 6 month expiry after opening. The Pharmacy Procedures Manual restriction allowing one bottle per month is not relevant and therefore only the prescribed dosage to the nearest OP may be claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE

* Eye drops 0.1%	4.15	15 ml OP	✓ Naphcon Forte
------------------------	------	----------	------------------------

OLOPATADINE

Eye drops 0.1%	2.20	5 ml OP	✓ Olopatadine Teva
----------------------	------	---------	---------------------------

PARAFFIN LIQUID WITH WOOL FAT

* Eye oint 3% with wool fat 3%	3.63	3.5 g OP	✓ Poly-Visc
--------------------------------------	------	----------	--------------------

RETINOL PALMITATE

Eye oint 138 mcg per g	3.80	5 g OP	✓ Vita-POS
------------------------------	------	--------	-------------------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

Various

PHARMACY SERVICES

May only be claimed once per patient.

* Brand switch fee	4.50	1 fee	✓ BSF Cinacalcet Devatis
			✓ BSF Comtan
			✓ BSF Vigisom

a) The Pharmacode for BSF Vigisom is 2634112 - see also [page 136](#)

b) The Pharmacode for BSF Cinacalcet Devatis is 2634120 - see also [page 78](#)

c) The Pharmacode for BSF Comtan is 2634139 - see also [page 118](#)

(BSF Cinacalcet Devatis Brand switch fee to be delisted 1 July 2022)

(BSF Comtan Brand switch fee to be delisted 1 July 2022)

(BSF Vigisom Brand switch fee to be delisted 1 July 2022)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE

Inj 200 mg per ml, 10 ml ampoule	58.76	10	✓ DBL Acetylcysteine Martindale Pharma ^{S29}
--	-------	----	---

NALOXONE HYDROCHLORIDE

a) Up to 5 inj available on a PSO

b) Only on a PSO

* Inj 400 mcg per ml, 1 ml ampoule	22.60	5	✓ DBL Naloxone Hydrochloride
--	-------	---	---------------------------------

Removal and Elimination

CHARCOAL

* Oral liq 50 g per 250 ml	43.50	250 ml OP	✓ Carbosorb-X
a) Up to 250 ml available on a PSO			
b) Only on a PSO			

DEFERASIROX – Special Authority see [SA1492 below](#) – Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	✓ Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	✓ Exjade

► [SA1492](#) Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or

continued...

▲ Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

* Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

continued...

- 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
- 3.3 Treatment with deferiprone has resulted in arthritis; or
- 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 - 1.0 cells per μL).

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE – Special Authority see [SA1480 below](#) – Retail pharmacy

Tab 500 mg	533.17	100	✓ Feriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Feriprox

➔ **SA1480** Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DEFERIOXAMINE MESILATE

* Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP
-------------------------	--------	----	--

SODIUM CALCIUM EDETATE

* Inj 200 mg per ml, 5 ml	53.31 (156.71)	6	Calcium Disodium Versenate
---------------------------------	-------------------	---	-------------------------------

Standard Formulae

ACETYLCYSTEINE EYE DROPS

Acetylcysteine inj 200 mg per ml, 10 ml	qs
Suitable eye drop base	qs

PHENOBARBITONE ORAL LIQUID

Phenobarbitone Sodium	1 g
Glycerol BP	70 ml
Water	to 100 ml

CODEINE LINCTUS (3 mg per 5 ml)

Codeine phosphate	60 mg
Glycerol	40 ml
Preservative	qs
Water	to 100 ml

PHENOBARBITONE SODIUM PAEDIATRIC ORAL LIQUID (10 mg per ml)

Phenobarbitone Sodium	400 mg
Glycerol BP	4 ml
Water	to 40 ml

CODEINE LINCTUS (15 mg per 5 ml)

Codeine phosphate	300 mg
Glycerol	40 ml
Preservative	qs
Water	to 100 ml

PILOCARPINE ORAL LIQUID

Pilocarpine 4% eye drops	qs
Preservative	qs
Water	to 500 ml

(Preservative should be used if quantity supplied is for more than 5 days.)

FOLINIC MOUTHWASH

Calcium folinate 15 mg tab	1 tab
Preservative	qs
Water	to 500 ml

(Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.)

SALIVA SUBSTITUTE FORMULA

Methylcellulose	5 g
Preservative	qs
Water	to 500 ml

(Preservative should be used if quantity supplied is for more than 5 days. Maximum 500 ml per prescription.)

METHADONE MIXTURE

Methadone powder	qs
Glycerol	qs
Water	to 100 ml

SODIUM CHLORIDE ORAL LIQUID

Sodium chloride inj 23.4%, 20 ml	qs
Water	qs

METHYL HYDROXYBENZOATE 10% SOLUTION

Methyl hydroxybenzoate	10 g
Propylene glycol	to 100 ml

(Use 1 ml of the 10% solution per 100 ml of oral liquid mixture)

(Only funded if prescribed for treatment of hyponatraemia)

VANCOMYCIN ORAL SOLUTION (50 mg per ml)

Vancomycin 500 mg injection	10 vials
Glycerol BP	40 ml
Water	to 100 ml

OMEPRAZOLE SUSPENSION

Omeprazole capules or powder	qs
Sodium bicarbonate powder BP	8.4 g
Water	to 100 ml

(Only funded if prescribed for treatment of Clostridium difficile following metronidazole failure)

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
Extemporaneously Compounded Preparations and Galenicals				
CODEINE PHOSPHATE – Safety medicine; prescriber may determine dispensing frequency				
Powder – Only in combination.....	63.09 (90.09)	25 g		Douglas
Only in extemporaneously compounded codeine linctus.				
COLLODION FLEXIBLE				
Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.				
Collodion flexible	19.30	100 ml	✓	PSM
COMPOUND HYDROXYBENZOATE – Only in combination				
Only in extemporaneously compounded oral mixtures.				
Soln	30.00	100 ml	✓	Midwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination				
Only in combination with Ora-Plus.				
Suspension.....	30.95	473 ml	✓	Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination				
Only in combination with Ora-Plus.				
Suspension.....	30.95	473 ml	✓	Ora-Sweet
GLYCEROL				
* Liquid – Only in combination	3.23	500 ml	✓	healthE Glycerol BP
Only in extemporaneously compounded oral liquid preparations.				
METHADONE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequency				
d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).				
Powder	7.84	1 g	✓	AFT
METHYL HYDROXYBENZOATE				
Powder	8.98	25 g	✓	Midwest
METHYLCELLULOSE				
Powder	36.95	100 g	✓	MidWest
Suspension – Only in combination	30.95	473 ml	✓	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHARIN – Only in combination				
Suspension.....	30.95	473 ml	✓	Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE – Only in combination				
Suspension.....	30.95	473 ml	✓	Ora-Blend
PHENOBARBITONE SODIUM				
Powder – Only in combination.....	52.50 325.00	10 g 100 g	✓ ✓	MidWest MidWest
Only in children up to 12 years				
PROPYLENE GLYCOL				
Only in extemporaneously compounded methyl hydroxybenzoate 10% solution.				
Liq.....	11.25	500 ml	✓	Midwest
SODIUM BICARBONATE				
Powder BP – Only in combination	10.05	500 g	✓	Midwest
Only in extemporaneously compounded omeprazole and lansoprazole suspension.				

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$	Fully Subsidised ✓	Brand or Generic Manufacturer
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparations.			
Liq.....	14.95	500 ml	✓ Midwest
WATER			
Tap – Only in combination.....	0.00	1 ml	✓ Tap water

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

*Three months or six months, as applicable, dispensed all-at-once

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Nutrient Modules

Carbohydrate

►SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Either:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 cancer in children; or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism.

Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT – Special Authority see [SA1930 above](#) – Hospital pharmacy [HP3]

Powder5.29 400 g OP ✓ Polycal

Carbohydrate And Fat

►SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE AND FAT SUPPLEMENT – Special Authority see [SA1376 on the previous page](#) – Hospital pharmacy [HP3]

Powder (neutral)	60.31	400 g OP	✓ Duocal Super Soluble Powder
------------------------	-------	----------	--------------------------------------

Fat

»SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome; or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 10 ascites; or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Renewal — (Inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 3 years for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see [SA1523 on the previous page](#) – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry).....	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

Protein

►SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT – Special Authority see [SA1524 above](#) – Hospital pharmacy [HP3]

Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource Beneprotein

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Oral and Enteral Feeds

Diabetic Products

►SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML – Special Authority see [SA1095 above](#) – Hospital pharmacy [HP3]

Liquid	3.75	500 ml OP	✓ Glucerna Select
	7.50	1,000 ml OP	✓ Diason RTH
			✓ Nutrison Advanced Diason

(Diason RTH Liquid to be delisted 1 December 2022)

DIABETIC ORAL FEED 1KCAL/ML – Special Authority see [SA1095 above](#) – Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 ml OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

Fat Modified Products

►SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see [SA1525 above](#) – Hospital pharmacy [HP3]

Powder	60.48	400 g OP	✓ Monogen
--------------	-------	----------	-----------

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Paediatric Products For Children Awaiting Liver Transplant

►SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML – Special Authority see [SA1098 above](#) – Hospital pharmacy [HP3]

Powder (unflavoured) 78.97 400 g OP ✓ **Heparon Junior**

Paediatric Products For Children With Chronic Renal Failure

►SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML – Special Authority see [SA1099 above](#) – Hospital pharmacy [HP3]

Powder 54.00 400 g OP ✓ **Kindergen**

Paediatric Products

►SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
continued...			
applications meeting the following criteria:			
Both:			
1 The treatment remains appropriate and the patient is benefiting from treatment; and			
2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid.....	6.00	500 ml OP	✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid.....	2.68	500 ml OP	✓ Nutrini RTH ✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid.....	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Powder (vanilla).....	21.07	850 g OP	✓ Pediasure
<i>(Pediasure Powder (vanilla) to be delisted 1 September 2022)</i>			
PAEDIATRIC ORAL FEED 1.5KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid (strawberry).....	1.60	200 ml OP	✓ Fortini
Liquid (vanilla).....	1.60	200 ml OP	✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid (chocolate).....	1.07	200 ml OP	✓ Pediasure
Liquid (strawberry).....	1.07	200 ml OP	✓ Pediasure
Liquid (vanilla).....	1.07	200 ml OP	✓ Pediasure
	1.34	250 ml OP	✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Liquid (unflavoured).....	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (chocolate).....	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (strawberry).....	1.60	200 ml OP	✓ Fortini Multi Fibre
Liquid (vanilla).....	1.60	200 ml OP	✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the previous page – Hospital pharmacy [HP3]			
Powder.....	43.60	400 g OP	✓ Peptamen Junior

Renal Products

►SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML – Special Authority see [SA1101 above](#) – Hospital pharmacy [HP3]

Liquid..... 6.08 500 ml OP ✓ **Nepro HP RTH**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1101 on the previous page – Hospital pharmacy [HP3]				
Liquid.....	2.67	220 ml OP	✓	Nepro HP (strawberry) ✓ Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA1101 on the previous page – Hospital pharmacy [HP3]				
Liquid.....	2.88 (3.31)	237 ml OP		NovaSource Renal
Liquid, 200 ml bottle	11.52 (13.24)	4 OP		NovaSource Renal
Liquid (apricot) 125 ml.....	11.52	4 OP	✓	Renilon 7.5
Liquid (caramel) 125 ml.....	11.52	4 OP	✓	Renilon 7.5
<i>(NovaSource Renal Liquid to be delisted 1 September 2022)</i>				

Specialised And Elemental Products

➔ [SA1377](#) Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Special Authority see SA1377 above – Hospital pharmacy [HP3]				
Liquid.....	18.06	1,000 ml OP	✓	Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML – Special Authority see SA1377 above – Hospital pharmacy [HP3]				
Liquid (grapefruit), 250 ml carton.....	171.00	18 OP	✓	Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton.....	171.00	18 OP	✓	Elemental 028 Extra
Liquid (summer fruits), 250 ml carton	171.00	18 OP	✓	Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML – Special Authority see SA1377 above – Hospital pharmacy [HP3]				
Powder (unflavoured)	4.50	80 g OP	✓	Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority see SA1377 above – Hospital pharmacy [HP3]				
Liquid.....	12.04	1,000 ml OP	✓	Nutrison Advanced Peptisorb ✓ Peptisorb

(Peptisorb Liquid to be delisted 1 June 2023)

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Paediatric Products For Children With Low Energy Requirements

►SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML – Special Authority see [SA1196 above](#) – Hospital pharmacy [HP3]

Liquid	4.00	500 ml OP	✓ Nutrini Low Energy Multi Fibre
--------------	------	-----------	---

Standard Supplements

►SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - Patient is Malnourished
 - 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:
 - Patient has not responded to first-line dietary measures over a 4 week period by:
 - 2.1 Increasing their food intake frequency (eg snacks between meals); or
 - 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
 - 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:
 - Patient is Malnourished
 - 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
 - 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
 - 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions; or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube - refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	1.75	250 ml OP	✓	Ensure Plus HN
	7.00	1,000 ml OP	✓	Ensure Plus RTH
			✓	Nutrison Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	1.24	250 ml OP	✓	Isosource Standard
	5.29	1,000 ml OP	✓	Nutrison Standard RTH
			✓	Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	5.29	1,000 ml OP	✓	Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	5.29	1,000 ml OP	✓	Jevity RTH
			✓	Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	6.35	1,000 ml OP	✓	Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Liquid.....	7.00	1,000 ml OP	✓	Jevity HiCal RTH
			✓	Nutrison Energy Multi Fibre
ORAL FEED (POWDER) – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]				
Powder (chocolate).....	14.00	840 g OP	✓	Sustagen Hospital Formula
	26.00	850 g OP	✓	Ensure
Powder (vanilla).....	14.00	840 g OP	✓	Sustagen Hospital Formula Active
	26.00	850 g OP	✓	Ensure

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
ORAL FEED 1.5KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]			
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO ₂ value exceeding 55mmHg. The prescription must be endorsed accordingly.			
Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26) (1.26)	200 ml OP	Ensure Plus Fortisip
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26) (1.26)	200 ml OP	Ensure Plus Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26)	200 ml OP	Ensure Plus
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26)	200 ml OP	Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85 (1.33) 0.72 (1.26) (1.26)	237 ml OP 200 ml OP	Ensure Plus Ensure Plus Fortisip
ORAL FEED WITH FIBRE 1.5 KCAL/ML – Special Authority see SA1859 on page 253 – Hospital pharmacy [HP3]			
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.			
Liquid (chocolate) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26)	200 ml OP	Fortisip Multi Fibre
Liquid (strawberry) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26)	200 ml OP	Fortisip Multi Fibre
Liquid (vanilla) – Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72 (1.26)	200 ml OP	Fortisip Multi Fibre

High Calorie Products

►SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML – Special Authority see [SA1195 on the previous page](#) – Hospital pharmacy [HP3]

Liquid.....	5.50	500 ml OP	✓ Nutrison Concentrated
	11.00	1,000 ml OP	✓ Ensure Two Cal HN RTH

ORAL FEED 2 KCAL/ML – Special Authority see [SA1195 on the previous page](#) – Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) – Higher subsidy of \$1.90 per 200 ml with

Endorsement	0.96 (1.90)	200 ml OP	Two Cal HN
-------------------	----------------	-----------	------------

Food Thickeners

►SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
FOOD THICKENER – Special Authority see SA1106 on the previous page – Hospital pharmacy [HP3]			
Powder	6.53	300 g OP	✓ Nutilis
	7.25	380 g OP	✓ Feed Thickener Karicare Aptamil

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

►SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner.

Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist.

Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see [SA1729 above](#) – Hospital pharmacy [HP3]

Powder	2.81	1,000 g OP	
	(5.15)		Healthies Simple Baking Mix

GLUTEN FREE BREAD MIX – Special Authority see [SA1729 above](#) – Hospital pharmacy [HP3]

Powder	3.93	1,000 g OP	
	(7.32)		NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix

GLUTEN FREE FLOUR – Special Authority see [SA1729 above](#) – Hospital pharmacy [HP3]

Powder	5.62	2,000 g OP	
	(18.10)		Horleys Flour

SPECIAL FOODS

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
GLUTEN FREE PASTA – Special Authority see SA1729 on the previous page – Hospital pharmacy [HP3]				
Buckwheat Spirals.....	2.00 (3.11)	250 g OP		Orgran
Corn and Vegetable Shells.....	2.00 (2.92)	250 g OP		Orgran
Corn and Vegetable Spirals.....	2.00 (2.92)	250 g OP		Orgran
Rice and Corn Lasagne Sheets.....	1.60 (3.82)	200 g OP		Orgran
Rice and Corn Macaroni.....	2.00 (2.92)	250 g OP		Orgran
Rice and Corn Penne	2.00 (2.92)	250 g OP		Orgran
Rice and Maize Pasta Spirals.....	2.00 (2.92)	250 g OP		Orgran
Rice and Millet Spirals.....	2.00 (3.11)	250 g OP		Orgran
Rice and corn spaghetti noodles	2.00 (2.92)	375 g OP		Orgran
Vegetable and Rice Spirals	2.00 (2.92)	250 g OP		Orgran
Italian long style spaghetti	2.00 (3.11)	220 g OP		Orgran

Foods And Supplements For Inborn Errors Of Metabolism

► [SA1108](#) Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

AMINOACID FORMULA WITHOUT METHIONINE – Special Authority see [SA1108 above](#) – Hospital pharmacy [HP3]

Powder461.94 500 g OP ✓ **XMET Maxamum**

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE – Special Authority see [SA1108 above](#) – Hospital pharmacy [HP3]

Powder437.22 500 g OP ✓ **MSUD Maxamum**

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
---	----------------------------	-------------------------------------

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Tabs.....	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 28 g sachets.....	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets.....	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula.....	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange).....	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured).....	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton.....	540.00	18 OP	✓ Easiphen Liquid
Liquid (juicy tropical) 125 ml.....	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml.....	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml.....	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml.....	936.00	30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Powder	8.22	500 g OP	✓ Loprofin Mix
--------------	------	----------	----------------

LOW PROTEIN PASTA – Special Authority see [SA1108 on the previous page](#) – Hospital pharmacy [HP3]

Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne.....	5.95	250 g OP	✓ Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni.....	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals.....	11.91	500 g OP	✓ Loprofin

Infant Formulae

For Williams Syndrome

►SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA – Special Authority see [SA1110 above](#) – Hospital pharmacy [HP3]

Powder	44.40	400 g OP	✓ Locasol
--------------	-------	----------	-----------

Gastrointestinal and Other Malabsorptive Problems

AMINO ACID FORMULA – Special Authority see [SA2092 below](#) – Hospital pharmacy [HP3]

Powder	43.60	400 g OP	✓ Alfamino
			✓ Alfamino Junior
Powder (unflavoured)	53.00	400 g OP	✓ Elecare
			✓ Elecare LCP
			✓ Neocate Gold
			✓ Neocate Junior
			Unflavoured
			✓ Neocate SYNEO
Powder (vanilla)	53.00	400 g OP	✓ Elecare
			✓ Neocate Junior
			Vanilla

►SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:

6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and

6.2 Either:

6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or

6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist.

Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
---	-----	--------------------------	-------------------------------------

continued...

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA – Special Authority see [SA1953 below](#) – Hospital pharmacy [HP3]

Liquid 1 kcal/ml.....	10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml.....	15.68	500 ml OP	✓ Nutrini Peptisorb Energy

►SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea; or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption; or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure; or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

continued...

2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and

3 Either:

- 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
- 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA – Special Authority see [SA1557 below](#) – Hospital pharmacy [HP3]

Powder	15.21	450 g OP	✓ Aptamil Gold+ Pepti Junior
	30.42	900 g OP	✓ Aptamil AllerPro SYNEO 1
			✓ Aptamil AllerPro SYNEO 2

»SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

1 Both:

- 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
- 1.2 Either:

- 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
- 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or

2 Severe malabsorption; or

3 Short bowel syndrome; or

4 Intractable diarrhoea; or

5 Biliary atresia; or

6 Cholestatic liver diseases causing malsorption; or

7 Cystic fibrosis; or

8 Proven fat malabsorption; or

9 Severe intestinal motility disorders causing significant malabsorption; or

10 Intestinal failure; or

11 All of the following:

11.1 For step down from Amino Acid Formula; and

11.2 The infant is currently receiving funded amino acid formula; and

11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and

11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IgE mediated allergic reaction.

continued...

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

continued...

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML – Special Authority see [SA1698 below](#) – Hospital pharmacy [HP3]

Liquid.....	2.35	125 ml OP	✓ Infatrini
-------------	------	-----------	--------------------

►SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

►SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA – Special Authority see [SA1197 above](#) – Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1
			✓ Ketocal 3:1
Powder (vanilla).....	35.50	300 g OP	✓ KetoCal 4:1

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE – [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent..... 0.00 10 ✓ **BCG Vaccine**

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE – [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- 3) A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

haemagglutinin and 2.5 mcg pertactin in 0.5 ml syringe 0.00 10 ✓ **Boostrix**

1

✓ **Boostrix**

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- 2) A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

poliomyelitis virus in 0.5ml syringe 0.00 10 ✓ **Infanrix IPV**

NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AND HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Up to four doses for children up to and under the age of 10 for primary immunisation; or				
2) An additional four doses (as appropriate) are funded for (re-)immunisation for children up to and under the age of 10 who are patients post haematopoietic stem cell transplantation, or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or				
3) Up to five doses for children up to and under the age of 10 receiving solid organ transplantation.				
Note: A course of up-to four vaccines is funded for catch up programmes for children (up to and under the age of 10 years) to complete full primary immunisation. Please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes.				
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus, 10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	✓	Infanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE – [Xpharm]				
One dose for patients meeting any of the following:				
1) For primary vaccination in children; or				
2) An additional dose (as appropriate) is funded for (re-)immunisation for patients post haematopoietic stem cell transplantation, or chemotherapy; functional asplenic; pre or post splenectomy; pre- or post solid organ transplant, pre- or post cochlear implants, renal dialysis and other severely immunosuppressive regimens; or				
3) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.				
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg; prefilled syringe plus vial 0.5 ml.....	0.00	1	✓	Hiberix
HEPATITIS A VACCINE – [Xpharm]				
Funded for patients meeting any of the following criteria:				
1) Two vaccinations for use in transplant patients; or				
2) Two vaccinations for use in children with chronic liver disease; or				
3) One dose of vaccine for close contacts of known hepatitis A cases.				
Inj 1440 ELISA units in 1 ml syringe.....	0.00	1	✓	Havrix
Inj 720 ELISA units in 0.5 ml syringe.....	0.00	1	✓	Havrix Junior

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm]			
Inj 10 mcg per 0.5 ml prefilled syringe.....	0.00	1	✓ Engerix-B
Funded for patients meeting any of the following criteria:			
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or			
2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or			
4) for HIV positive patients; or			
5) for hepatitis C positive patients; or			
6) for patients following non-consensual sexual intercourse; or			
7) for patients following immunosuppression; or			
8) for solid organ transplant patients; or			
9) for post-haematopoietic stem cell transplant (HSCT) patients; or			
10) following needle stick injury.			
Inj 20 mcg per 1 ml prefilled syringe.....	0.00	1	✓ Engerix-B
Funded for patients meeting any of the following criteria:			
1) for household or sexual contacts of known acute hepatitis B patients or hepatitis B carriers; or			
2) for children born to mothers who are hepatitis B surface antigen (HBsAg) positive; or			
3) for children up to and under the age of 18 years inclusive who are considered not to have achieved a positive serology and require additional vaccination or require a primary course of vaccination; or			
4) for HIV positive patients; or			
5) for hepatitis C positive patients; or			
6) for patients following non-consensual sexual intercourse; or			
7) for patients following immunosuppression; or			
8) for solid organ transplant patients; or			
9) for post-haematopoietic stem cell transplant (HSCT) patients; or			
10) following needle stick injury; or			
11) for dialysis patients; or			
12) for liver or kidney transplant patients.			
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 58) VACCINE [HPV] – [Xpharm]			
Any of the following:			
1) Maximum of two doses for children aged 14 years and under; or			
2) Maximum of three doses for patients meeting any of the following criteria:			
1) People aged 15 to 26 years inclusive; or			
2) Either:			
People aged 9 to 26 years inclusive			
1) Confirmed HIV infection; or			
2) Transplant (including stem cell) patients: or			
3) Maximum of four doses for people aged 9 to 26 years inclusive post chemotherapy			
Inj 270 mcg in 0.5 ml syringe.....	0.00	10	✓ Gardasil 9

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

– [Xpharm] 11.00 1 ✓ **Afluria Quad Junior**
(2022 formulation)

A) INFLUENZA VACCINE – child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebro-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.

- B) Doctors are the only Contractors entitled to claim payment from the Funder for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine) 110.00

10 ✓ **Afluria Quad**
(2022 formulation)

Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
---	------------------------------	-------------------------------------

- a) Only on a prescription
- b) No patient co-payment payable
- c)

A) INFLUENZA VACCINE – people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebro-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
 - d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
 - b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment from the Funder for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

MEASLES, MUMPS AND RUBELLA VACCINE

- Only on a prescription
- No patient co-payment payable
-

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- For primary vaccination in children; or
- For revaccination following immunosuppression; or
- For any individual susceptible to measles, mumps or rubella; or
- A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- Contractors will be entitled to claim payment from the Funder for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with their DHB for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.

- Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50, Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml	112.50	5	✓ MMR II
	250.00	10	✓ <u>Priorix</u>

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE – [Xpharm]

Either:

A) Any of the following:

- Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
- One dose for close contacts of meningococcal cases of any group; or
- One dose for person who has previously had meningococcal disease of any group; or
- A maximum of two doses for bone marrow transplant patients; or
- A maximum of two doses for person pre- and post-immunosuppression*; or

B) Both:

- Person is aged between 13 and 25 years, inclusive; and
- Either:
 - One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to a total of approximately 48 mcg of diphtheria toxoid carrier per 0.5 ml vial	0.00	1	✓ <u>Menactra</u>
---	------	---	-------------------

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
--	---	------------------------------	-------------------------------------

MENINGOCOCCAL B MULTICOMPONENT VACCINE – [Xpharm]

Either:

A) Both:

- 1) Child is under one year of age; and
- 2) Any of the following:
 - i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to three doses for close contacts of meningococcal cases of any group; or
 - iii) up to three doses for child who has previously had meningococcal disease of any group; or
 - iv) up to three doses for bone marrow transplant patients; or
 - v) up to three doses for child pre- and post-immunosuppression*; or

B) Both:

- 1) Person is one year of age or over; and
- 2) Any of the following:
 - i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or
 - ii) up to two doses for close contacts of meningococcal cases of any group; or
 - iii) up to two doses for person who has previously had meningococcal disease of any group; or
 - iv) up to two doses for bone marrow transplant patients; or
 - v) up to two doses for person pre- and post-immunosuppression*.

*Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 175 mcg per 0.5 ml prefilled syringe.....0.00 1 ✓ **Bexsero**

MENINGOCOCCAL C CONJUGATE VACCINE – [Xpharm]

Both:

- 1) The child is under 9 months of age; and
- 2) Any of the following:
 - 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant; or
 - 2) Two doses for close contacts of meningococcal cases of any group; or
 - 3) Two doses for child who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for child pre- and post-immunosuppression*.

Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 10 mcg in 0.5 ml syringe.....0.00 1 ✓ **Neisvac-C**

PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE – [Xpharm]

- 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B,

7F, 9V, 14 and 23F; 3 mcg of pneumococcal

polysaccharide serotypes 4, 18C and 19F in 0.5 ml

prefilled syringe0.00 10 ✓ **Synflorix**

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
--	---	-----	--------------------------	-------------------------------------

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE – [Xpharm]

Any of the following:

- 1) Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - l) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Inj 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

5, 6A, 6B, 7F, 9V, 14, 18C, 19A, 19F and 23F in 0.5ml

syringe	0.00	10	✓ Prevenar 13
		1	✓ Prevenar 13

	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per ✓	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE – [Xpharm]			
Either:			
1) Up to three doses (as appropriate) for patients with HIV, for patients post haematopoietic stem cell transplant, or chemotherapy; pre- or post-splenectomy or with functional asplenia, pre- or post-solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, or primary immunodeficiency; or			
2) All of the following:			
a) Patient is a child under 18 years for (re-)immunisation; and			
b) Treatment is for a maximum of two doses; and			
c) Any of the following:			
i) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or			
ii) with primary immune deficiencies; or			
iii) with HIV infection; or			
iv) with renal failure, or nephrotic syndrome; or			
v) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or			
vi) with cochlear implants or intracranial shunts; or			
vii) with cerebrospinal fluid leaks; or			
viii) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or			
ix) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or			
x) pre term infants, born before 28 weeks gestation; or			
xi) with cardiac disease, with cyanosis or failure; or			
xii) with diabetes; or			
xiii) with Down syndrome; or			
xiv) who are pre- or post-splenectomy, or with functional asplenia.			
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1	✓ <u>Pneumovax 23</u>
POLIOMYELITIS VACCINE – [Xpharm]			
Up to three doses for patients meeting either of the following:			
1) For partially vaccinated or previously unvaccinated individuals; or			
2) For revaccination following immunosuppression.			
Note: Please refer to the Immunisation Handbook for appropriate schedule for catch-up programmes.			
Inj 80D antigen units in 0.5 ml syringe.....	0.00	1	✓ <u>IPOL</u>
ROTAVIRUS ORAL VACCINE – [Xpharm]			
Maximum of two doses for patients meeting the following:			
1) first dose to be administered in infants aged under 14 weeks of age; and			
2) no vaccination being administered to children aged 24 weeks or over.			
Oral susp live attenuated human rotavirus			
1,000,000 CCID50 per dose, prefilled oral applicator.....	0.00	10	✓ <u>Rotarix</u>

NATIONAL IMMUNISATION SCHEDULE

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised ✓	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm]				
Either:				
1) Maximum of one dose for primary vaccination for either:				
a) Any infant born on or after 1 April 2016; or				
b) For previously unvaccinated children turning 11 years old on or after 1 July 2017, who have not previously had a varicella infection (chickenpox), or				
2) Maximum of two doses for any of the following:				
a) Any of the following for non-immune patients:				
i) with chronic liver disease who may in future be candidates for transplantation; or				
ii) with deteriorating renal function before transplantation; or				
iii) prior to solid organ transplant; or				
iv) prior to any elective immunosuppression*, or				
v) for post exposure prophylaxis who are immune competent inpatients.; or				
b) For patients at least 2 years after bone marrow transplantation, on advice of their specialist, or				
c) For patients at least 6 months after completion of chemotherapy, on advice of their specialist, or				
d) For HIV positive non immune to varicella with mild or moderate immunosuppression on advice of HIV specialist, or				
e) For patients with inborn errors of metabolism at risk of major metabolic decompensation, with no clinical history of varicella, or				
f) For household contacts of paediatric patients who are immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella, or				
g) For household contacts of adult patients who have no clinical history of varicella and who are severely immunocompromised, or undergoing a procedure leading to immune compromise where the household contact has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosuppressive therapy must be for a treatment period of greater than 28 days				
Inj 1350 PFU prefilled syringe	0.00	1	✓	Varivax
		10	✓	Varivax
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATED VACCINE [SHINGLES VACCINE] – [Xpharm]				
Funded for patients meeting the following criteria:				
1) One dose for all people aged 65 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1	✓	Zostavax
		10	✓	Zostavax

Diagnostic Agents

TUBERCULIN PPD [MANTOUX] TEST – [Xpharm]				
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	✓	Tubersol

- Symbols -		
UK Synacthen.....	80	
3TC.....	105	
- A -		
A-Scabies.....	68	
Abacavir sulphate.....	105	
Abacavir sulphate with lamivudine.....	105	
Abiraterone acetate.....	164	
Acarbose.....	11	
Accarb.....	11	
Accuretic 10.....	48	
Accuretic 20.....	48	
Acetazolamide.....	238	
Acetec.....	47	
Acetic acid with hydroxyquinoline and ricinoleic acid.....	75	
Acetylcysteine.....	241	
Aci-Jel.....	75	
Aciclovir.....		
Infection.....	100	
Sensory.....	236	
Acidex.....	6	
Acipimox.....	54	
Acitretin.....	68	
Aclasta.....	114	
Actemra.....	215	
Actemra S29.....	215	
Actinomycin D.....	151	
Actrapid.....	10	
Actrapid Penfill.....	10	
Acupan.....	120	
Adalimumab (Amgevita).....	175	
Adalimumab (Humira).....	183	
Adapalene.....	61	
ADR Cartridge 1.8.....	23	
Adrenaline.....	57	
Advantan.....	64	
Advate.....	41	
Adynovate.....	41	
Afinitor.....	223	
Aflibercept.....	188	
Afluria Quad (2022 formulation).....	270	
Afluria Quad Junior (2022 formulation).....	270	
AFT-Pyrazinamide.....	100	
Agents Affecting the Renin-Angiotensin System.....	47	
Agents for Parkinsonism and Related Disorders.....	118	
Agents Used in the Treatment of Poisonings.....	241	
Agrylin.....	150	
Albendazole.....	89	
Albey.....	227	
Albustix.....	77	
Alchemy Oxybutynin.....	76	
Aldurazyme.....	29	
Alecensa.....	157	
Alectinib.....	157	
Alendronate sodium.....	111	
Alendronate sodium with colecalciferol.....	111	
Alfacalcidol.....	33	
Alfamino.....	262	
Alfamino Junior.....	262	
Alginic acid.....	6	
Alglucosidase alfa.....	26	
Alkeran.....	147	
Alkeran S29.....	147	
Allersoothe.....	228	
Allmercap.....	149	
Allopurinol.....	116	
Alpha-Adrenoceptor Blockers.....	47	
Alpha-Keri Lotion.....	66	
Alphamox.....	92	
Alphamox 125.....	92	
Alphamox 250.....	92	
Alprolix.....	38	
Alu-Tab.....	6	
Aluminium hydroxide.....	6	
Alvogen.....	50	
Amantadine hydrochloride.....	118	
Ambrisentan.....	57	
Ambrisentan Mylan.....	57	
Amgevita.....	175	
Amiloride hydrochloride.....	53	
Amiloride hydrochloride with furosemide.....	53	
Amiloride hydrochloride with hydrochlorothiazide.....	53	
Aminophylline.....	233	
Amiodarone hydrochloride.....	49	
Amisulpride.....	131	
Amisulpride Mylan.....	131	
Amitriptyline.....	124	
Amlodipine.....	51	
Amneal.....	155	
Amorolfine.....	62	
Amoxicillin.....	92	
Amoxicillin with clavulanic acid.....	92	
Amphotericin B.....	33	
Amsacrine.....	150	
AmsaLyo.....	150	
Amsidine.....	150	
Amzoate.....	30	
Anaesthetics.....	119	
Anagrelide hydrochloride.....	150	
Analgesics.....	120	
Anastrozole.....	167	
Anatrole.....	167	
Andriol Testocaps.....	81	
Androderm.....	80	
Anoro Ellipta.....	231	
Antabuse.....	142	
Antacids and Antiflatulents.....	6	
Anthelmintics.....	89	
Antiacne Preparations.....	61	
Antiallergy Preparations.....	227	
Antianaeemics.....	37	
Antiandrogen Oral Contraceptives.....	75	
Antiarrhythmics.....	49	
Antibacterials.....	89	
Antibacterials Topical.....	61	
Anticholinergic Agents.....	230	
Anticholinesterases.....	110	
Antidepressants.....	124	
Antidiarrhoeals.....	6	
Antiepilepsy Drugs.....	126	
Antifibrinolytics, Haemostatics and Local Sclerosants.....	38	
Antifibrotics.....	231	
Antifungals.....	96	
Antifungals Topical.....	62	
Antihistamines.....	228	
Antihypotensives.....	49	
Antimalarials.....	98	
Antimigraine Preparations.....	129	
Antinausea and Vertigo Agents.....	129	
Antipruritic Preparations.....	63	
Antipsychotics.....	131	
Antiretrovirals.....	104	
Antirheumatoid Agents.....	111	
Antispasmodics and Other Agents Altering Gut Motility.....	8	
Antithrombotic Agents.....	41	
Antithymocyte globulin (equine).....	175	
Antitrichomonal Agents.....	98	
Antituberculosics and Antileprotics.....	99	
Antulcerants.....	8	
Antivirals.....	100	
Anxiolytics.....	134	
Anzatax.....	154	
Apidra.....	11	
Apidra SoloStar.....	11	
APO-Atomoxetine.....	137	
APO-Atomoxetine S29.....	137	
Apo-Azithromycin.....	90	

Apo-Diltiazem CD	52	Atrovent	230	Betahistine dihydrochloride	130
Apo-Doxazosin	47	AU Synacthen	80	Betaine	27
Apo-Prednisone	80	Aubagio	136	Betaloc CR	50
Apo-Primidone	127	Augmentin	92	Betamethasone dipropionate	63
Apo-Temozolomide	155	Aurorix	125	Betamethasone dipropionate with calcipotriol	68
Apomorphine hydrochloride	118	AutoSoft 30	23	Betamethasone sodium phosphate with betamethasone acetate	79
Appetitant	129	AutoSoft 90	23	Betamethasone valerate	63, 70
Apresoline	57	Avonex	135	Betamethasone valerate with sodium fusidate [fusidic acid]	64
Aptamil AllerPro SYNEO 1	265	Avonex Pen	135	Betaxolol	238
Aptamil AllerPro SYNEO 2	265	Azaciditidine	147	Betnovate	63
Aptamil Gold+ Pepti Junior	265	Azaciditidine Dr Reddy's	147	Betoptic	238
Aqueous cream	65	Azithromycin	90	Betoptic S	238
Aratac	49	Azopt	238	Bexsero	273
Arava	111	AZT	105-106	Bezafibrate	54
Arginine	27	- B -			54
Aripiprazole	131	B-D Micro-Fine	15	Bezalip	54
Aripiprazole Sandoz	131	B-D Ultra Fine	16	Bezalip Retard	54
Aristocort	64	B-D Ultra Fine II	16	Bicalutamide	165
Artotex-Prazosin S29	47	Bacillus Calmette-Guerin (BCG) vaccine	175	Bicillin LA	92
Arrow - Lattim	239	Bacillus Calmette-Guerin vaccine	267	BiCNU	146
Arrow-Amitriptyline	124	Baclofen	117	Bicu Heritage	146
Arrow-Bendrofluazide	54	Bactroban	62	Bile and Liver Therapy	9
Arrow-Brimonidine	239	Balance	29	Biltricide	89
Arrow-Diazepam	134	Barrier Creams and Emollients	65	Bimatoprost	238
Arrow-Doxorubicin	152	BCG Vaccine	267	Bimatoprost Multichem	238
Arrow-Losartan & Hydrochlorothiazide	48	Beclazone 100	228	Binarex	165
Arrow-Norfloxacin	109	Beclazone 250	228	Binocrit	38
Arrow-Ornidazole	98	Beclazone 50	228	Biodone	122
Arrow-Quinapril 10	48	Beclomethasone dipropionate	228	Biodone Extra Forte	122
Arrow-Quinapril 20	48	Bee venom allergy treatment	227	Biodone Forte	122
Arrow-Quinapril 5	48	Bendamustine hydrochloride	145	Bisacodyl	26
Arrow-Roxithromycin	91	Bendrofluazide	54	Bisoprolol fumarate	50
Arrow-Timolol	238	Bendroflumethiazide [Bendrofluazide]	54	Bisoprolol Mylan	50
Arrow-Topiramate	128	Benzathine benzylpenicillin	92	BK Lotion	66
Arrow-Tramadol	124	Benzatropine mesylate	119	Bleomycin sulphate	151
Arsenic trioxide	150	Benzbromaron AL 100	116	Blood Colony-stimulating Factors	44
Asacol	7	Benzbromarone	116	Blood glucose diagnostic test meter	14
Ascorbic acid	33	Benztrop	119	Blood glucose diagnostic test strip	15
Aspen Adrenaline	57	Benzylamine hydrochloride	32	Blood glucose test strips (visually impaired)	15
Aspirin		Benzylpenicillin sodium [Penicillin G]	92	Blood Ketone Diagnostic Test Strip	13
Blood	41	Beta Cream	63	Bonjela	32
Nervous	120	Beta Ointment	63	Boostrix	267
Asthalin	230	Beta Scalp	70	Bortezomib	151
Atazanavir sulphate	106	Beta-Adrenoceptor Agonists	230	Bortezomib Dr Reddy's S29	151
Atenolol	50	Beta-Adrenoceptor Blockers	50	Bortezomib Dr-Reddy's	151
Atenolol AFT	50	Betadine	66	Bortezomib Juno	151
Atenolol AFT S29	50	Betadine Skin Prep	66	Bortezomib Juno S29	151
ATGAM	175	Betaferon	135	Bosentan	58
Ativan	134			Bosentan Dr Reddy's	58
Atomoxetine	137				
Atorvastatin	54				
Atropine sulphate					
Cardiovascular	49				
Sensory	239				
Atropt	239				

Bplex.....	33	Candesartan cilexetil	48	Chlorambucil.....	146
Breo Ellipta	229	Candestar	48	Chloramphenicol.....	236
Brevinor 1/28	74	Canesten	62	Chlorothiazide.....	54
Bricanyl Turbuhaler	230	Capecitabine.....	148	Chlorpheniramine maleate	228
Brilinta.....	41	Capercit.....	148	Chlorpromazine hydrochloride.....	131
Brimonidine tartrate	239	Capoten	47	Chlortalidone [Chlorthalidone]	54
Brimonidine tartrate with timolol maleate.....	239	Capsaicin		Chlorthalidone	54
Brinzolamide.....	238	Musculoskeletal	111	Chlorvescent.....	46
Brolene	236	Nervous.....	120	Choice Load 375	73
Bromocriptine mesylate	118	Captopril	47	Choice TT380 Short	73
Brufen SR	110	Carafate.....	9	Choice TT380 Standard	73
BSF Comtan	241	Carbaccord	146	Choline salicylate with cetalkonium chloride	32
BSF Vigisom.....	241	Carbamazepine	126	Ciclosporin.....	223
Buccastem	130	Carbimazole	83	Cilazapril.....	47
Budesonide		Carbomer	240	Cilicaine.....	93
Alimentary	6	Carboplatin	146	Cilicaine VK	93
Respiratory.....	228, 234	Carboplatin Ebewe	146	Cinacalcet.....	78
Budesonide with eformoterol	229	Carbosorb-X	241	Cipflox	93
Bumetanide	52	Cardinol LA	51	Ciprofloxacin	
Buprenorphine Naloxone BNM.....	141	Cardizem CD	52	Infection	93
Buprenorphine with naloxone	141	CareSens Dual	14	Sensory	236
Bupropion hydrochloride.....	142	CareSens N	14-15	Ciprofloxacin Teva.....	236
Burinex	52	CareSens N POP	14	Cisplatin	146
Burinex S29	52	CareSens N Premier	14	Cisplatin Ebewe.....	146
Buscopan.....	8	CareSens PRO.....	15	Citalopram hydrobromide	125
Buspirone hydrochloride.....	134	Carmellose sodium with gelatin and pectin.....	32	Cladribine	148
Buspirone Viatrix	134	Carmustine	146	Clarithromycin	
Busulfan.....	146	Carvedilol	50	Alimentary.....	9
- C -		Carvedilol Sandoz	50	Infection	90
Cabergoline	88	Casirivimab and imdevimab	189	Clexane	43
Cacit	34	Catapres	52	Clexane Forte	43
Caffeine citrate	235	CeeNU	147	Clindamycin	94
Calamine	63	Cefaclor monohydrate	89	Clinicians	27
Calamine-AFT	63	Cefalexin.....	89	Clinicians Renal Vit.....	34
Calci-Tab 500	34	Cefalexin Sandoz	89	Clobazam	126
Calcipotriol.....	68	Cefazolin.....	89	Clobetasol propionate.....	63, 70
Calcitonin	78	Ceftriaxone	89	Clobetasone butyrate	64
Calcitriol.....	34	Ceftriaxone-AFT	89	Clofazimine.....	99
Calcitriol-AFT	34	Cefuroxime axetil.....	89	Clomazol	
Calcium 500 mg Hexal	34	Celebrex	110	Dermatological	62
Calcium carbonate.....	6, 34	Celecoxib.....	110	Genito-Urinary.....	75
Calcium Channel Blockers	51	Celecoxib Pfizer.....	110	Clomifene citrate.....	88
Calcium Disodium Versenate	242	Celestone Chronodose.....	79	Clomipramine hydrochloride.....	124
Calcium folinate.....	148	Cellcept.....	168	Clomipramine Teva	124
Calcium Folate Ebewe.....	148	Centrally-Acting Agents	52	Clonazepam	126, 134
Calcium Folate Sandoz.....	148	Cephalexin ABM.....	89	Clonidine	52
Calcium Folate Sandoz S29	148	Cetirizine hydrochloride	228	Clonidine BNM.....	52
Calcium gluconate.....	35	Cetomacrogol	65	Clonidine hydrochloride.....	52
Calcium Homeostasis.....	78	Cetomacrogol with glycerol	65	Clonidine Teva.....	52
Calcium lactate gluconate with calcium carbonate	35	Cetomacrogol-AFT	65	Clopidogrel	41
Calcium polystyrene sulphonate.....	45	Cetuximab	189	Clopidogrel Multichem.....	41
Calcium Resonium	45	Charcoal	241	Clopine	131
Calcium-Sandoz Forte.....	35	Chemotherapeutic Agents	145	Clopixol.....	132, 134
Calogen	248	Chickenpox vaccine.....	276	Clotrimazole	
		Chlorafast	236	Dermatological	62

Genito-Urinary.....	75	Cyclizine lactate.....	130	DBL Pethidine Hydrochloride	124
Clozapine.....	131	Cyclogyl.....	239	DBL Vincristine Sulfate.....	157
Clozaril.....	131	Cyclonex.....	146	Decozol.....	33
Co-trimoxazole	96	Cyclopentolate hydrochloride	239	Deferasirox.....	241
Coal tar.....	68	Cyclophosphamide.....	146	Deferiprone.....	242
Coal tar with allantoin, menthol, phenol and sulphur.....	69	Cyclorin.....	99	Denosumab.....	111
Coal tar with salicylic acid and sulphur.....	69	Cycloserine.....	99	Deolate.....	97
Coco-Scalp.....	69	Cyproterone acetate.....	80	Deoxycoformycin.....	154
Codeine phosphate		Cyproterone acetate with ethinyloestradiol.....	75	Depo-Medrol.....	80
Extemporaneous.....	244	Cystadane.....	27	Depo-Provera.....	74
Nervous.....	121	Cytarabine.....	148	Depo-Testosterone.....	80
Coenzyme Q10.....	28	Cytotec.....	8	Deprim.....	96
Colchicine.....	116	Cytoxan.....	146	Dermol.....	63, 70
Colecalciferol.....	34	- D -		Desferrioxamine mesilate.....	242
Colestid.....	54	D-Penamine.....	111	Desmopressin.....	88
Colestipol hydrochloride	54	Dabigatran.....	44	Desmopressin acetate.....	88
Colgout.....	116	Dacarbazine.....	151	Desmopressin-PH&T.....	88
Colifoam.....	7	Dacarbazine APP.....	151	Desuric.....	116
Colistin sulphomethate.....	94	Dactinomycin [Actinomycin D].....	151	Detection of Substances in Urine.....	77
Colistin-Link.....	94	Daivobet.....	68	Dexamethasone	
Collodion flexible.....	244	Daivonex.....	68	Hormone.....	79
Colloidal bismuth subcitrate.....	9	Daktarin.....	63	Sensory.....	237
Colofac.....	8	Dalacin C.....	94	Dexamethasone phosphate.....	79
Coloxyl.....	25	Dantrium.....	117	Dexamethasone Phosphate	
Combigan.....	239	Dantrium S29.....	117	Panpharma.....	79
Compound electrolytes.....	45	Dantrolene.....	117	Dexamethasone with framycetin and gramicidin.....	236
Compound electrolytes with glucose [Dextrose].....	45	Daonil.....	11	Dexamethasone with neomycin sulphate and polymyxin B sulphate.....	237
Compound hydroxybenzoate.....	244	Dapa-Tabs.....	54	Dexamfetamine sulfate.....	137
Comtan.....	118	Dapsone.....	99	Dexmethsone.....	79
Concerta.....	139	Daraprim.....	95	Dextrochlorpheniramine maleate.....	228
Condoms.....	72	Darunavir.....	106	Dextrose.....	45
Condylane.....	70	Dasatinib.....	158	DHC Continus.....	122
Contraceptives - Hormonal.....	73	Daunorubicin.....	151	Diabetes.....	10
Contraceptives - Non-hormonal.....	71	Daunorubicin Zentiva.....	151	Diabetes Management	13
Copaxone.....	135	David One Step Cassette Pregnancy Test.....	76	Diacomit.....	128
Corticosteroids and Related Agents for Systemic Use	79	DBL Acetylcysteine.....	241	Diagnostic Agents.....	276
Corticosteroids Topical.....	63	DBL Adrenaline.....	57	Diamide Relief.....	6
Cortifoam.....	7	DBL Aminophylline.....	233	Diamox.....	238
Cosentyx.....	213	DBL Bleomycin Sulfate.....	151	Diasip.....	249
Cosmegen.....	151	DBL Carboplatin.....	146	Diason RTH.....	249
Coumadin.....	44	DBL Cisplatin.....	146	Diazepam.....	126, 134
Country Life.....	29	DBL Dacarbazine.....	151	Diazoxide.....	10
Coversyl.....	47	DBL Desferrioxamine Mesylate for Inj BP.....	242	Dibenzyline.....	47
Creon 10000.....	24	DBL Docetaxel.....	151	Diclofenac Sandoz.....	110
Creon 25000.....	24	DBL Ergometrine.....	75	Diclofenac sodium	
Creon Micro.....	24	DBL Gemcitabine.....	149	Musculoskeletal.....	110
Crotamiton.....	63	DBL Gentamicin.....	94	Sensory.....	237
Crystaderm.....	61	DBL Heparin Sodium.....	43	Differin.....	61
Curam.....	92	DBL Leucovorin Calcium.....	148	Diffiam.....	32
Curam Duo 500/125.....	92	DBL Methotrexate Onco-Vial.....	149	Diffucan.....	96
Cvite.....	33	DBL Morphine Sulphate.....	123	Digestives Including Enzymes.....	24
Cyclizine hydrochloride.....	130	DBL Naloxone Hydrochloride	241		

Digoxin.....	49	Dulcolax SP Drop	26	Enstilar.....	68
Dihydrocodeine tartrate	122	Duocal Super Soluble Powder.....	247	Ensure	256
Dilantin.....	127	Duolin	230	Ensure Plus	257
Dilantin Infatab.....	127	Duolin HFA	230	Ensure Plus HN.....	256
Diltiazem hydrochloride	52	DuoResp Spiromax	229	Ensure Plus RTH.....	256
Dimethicone.....	65-66	Duride.....	57	Ensure Two Cal HN RTH	258
Dimethyl fumarate	135	- E -			
Dipentum.....	7	e-chamber La Grande	234	Entacapone	118
Diphtheria, tetanus and pertussis vaccine	267	e-chamber Mask.....	234	Entecavir.....	100
Diphtheria, tetanus, pertussis and polio vaccine.....	267	e-chamber Turbo.....	234	Entecavir Sandoz	100
Diphtheria, tetanus, pertussis, polio, hepatitis B and haemophilus influenzae type B vaccine.....	268	E-Mycin.....	91	Entocort CIR.....	6
Diprosone	63	Ear Preparations.....	236	Entresto 24/26	48
Diprosone OV.....	63	Ear/Eye Preparations	236	Entresto 49/51	48
Dipyridamole.....	41	Easiphen Liquid	261	Entresto 97/103.....	48
Disopyramide phosphate.....	49	Econazole nitrate.....	62	Epilim	128
Disulfiram.....	142	Efavirenz.....	105	Epilim Crushable	128
Diuretics.....	52	Efavirenz with emtricitabine and tenofovir disoproxil.....	105	Epilim IV	128
Docetaxel.....	151	Eformoterol fumarate	229	Epilim S/F Liquid.....	128
Docetaxel Accord	151	Eformoterol fumarate dihydrate	229	Epilim Syrup	128
Docetaxel Sandoz	151	Eftrenonacog alfa [Recombinant factor IX]	38	Epirubicin Ebewe.....	152
Docusate sodium.....	25	Efudix.....	70	Epirubicin hydrochloride	152
Docusate sodium with sennosides	25	Egopsoryl TA.....	69	Eplerenone	53
Dolutegravir	106	Elaprase	28	Epoetin alfa.....	38
Domperidone.....	130	Eldepryl.....	118	Epoprostenol	60
Donepezil hydrochloride.....	140	Elecare	262	Eptacog alfa [Recombinant factor VIIa]	40
Donepezil-Rex.....	140	Elecare LCP	262	ERA	91
Dornase alfa	233	Electral.....	45	Eribitux	189
Dortimopt.....	238	Elelyso	31	Ergometrine maleate	75
Dorzolamide hydrochloride.....	238	Elemental 028 Extra	252	Erlotinib.....	158
Dorzolamide with timolol.....	238	Elidel.....	69	Erythrocin IV	91
Dostinex.....	88	Elocon.....	64	Erythromycin (as lactobionate).....	91
Dosulepin [Dothiepin] hydrochloride	124	Elocon Alcohol Free	64	Erythromycin ethyl succinate.....	91
Dosulepin Mylan.....	124	Eltrombopag	38	Erythromycin stearate.....	91
Dothiepin	124	Eltroxin.....	83	Esbriet	232
Doxazosin.....	47	EMB Fatol.....	99	Escitalopram.....	125
Doxazosin Clinect.....	47	Emend Tri-Pack.....	129	Escitalopram (Ethics).....	125
Doxine	93	Emicizumab	39	Eskazole	89
Doxorubicin Ebewe.....	152	EMLA.....	120	Essential Generics.....	50
Doxorubicin hydrochloride.....	152	Empagliflozin	13	Essential Prednisolone.....	8
Doxycycline	93	Empagliflozin with metformin hydrochloride.....	13	Estradot	81
DP Lotion.....	66	Emtricitabine.....	105	Estradot 50 mcg	81
DP Lotn HC	64	Emtricitabine with tenofovir disoproxil	102	Estrofem	81
DP-Allopurinol.....	116	Emtriva	105	Etanercept	168
Dr Reddy's Omeprazole	9	Emulsifying ointment	65	Ethambutol hydrochloride.....	99
Drofade.....	51	Emulsifying Ointment ADE	65	Ethics Aspirin.....	120
Drugs Affecting Bone Metabolism	111	Enalapril maleate.....	47	Ethics Aspirin EC.....	41
Dual blood glucose and blood ketone diagnostic test meter	14	Enbrel.....	168	Ethics Lisinopril.....	47
Dulaglutide.....	12	Endocrine Therapy	164	Ethinylestradiol	82
		Endoxan	146	Ethinylestradiol with desogestrel	73
		Engerix-B.....	269	Ethinylestradiol with levonorgestrel	74
		Enlaxaf XR.....	125	Ethinylestradiol with norethisterone.....	74
		Enoxaparin sodium.....	43	Ethosuximide.....	126
				Etopophos	152

Etoposide.....	152	Flecainide BNM	49	Frumil.....	53
Etoposide phosphate.....	152	Flecainide Controlled Release Teva.....	49	Frusemide.....	53
Etravirine	105	Fleet Phosphate Enema.....	26	Fucicort.....	64
Eumovate	64	Flixonase Hayfever & Allergy	234	Fucidin.....	95
Everet	127	Flixotide	229	Fucithalmic	236
Everolimus	223	Flixotide Accuhaler	229	Fulvestrant.....	165
Evista.....	112	Florinef.....	79	Fungilin.....	33
Exemestane.....	167	Fluanxol	132	Furosemid-Ratiopharm.....	53
Exjade.....	241	Flucil.....	92	Furosemide [Frusemide].....	53
Extemporaneously Compounded Preparations and Galenicals.....	244	Flucloxacillin	92	Furosemide-Baxter	53
Eye Preparations.....	236	Flucloxacillin-AFT	92	fusidic acid Dermatological.....	62, 64
Eylea.....	188	Flucloxin	92	Infection	95
Ezetimibe.....	55	Flucon.....	237	Sensory.....	236
Ezetimibe Sandoz.....	55	Fluconazole	96		
Ezetimibe with simvastatin	56	Fludara Oral.....	148	- G -	
		Fludarabine Ebewe.....	148	Gabapentin.....	126
- F -		Fludarabine phosphate.....	148	Gacet.....	121
Factor eight inhibitor bypassing fraction.....	40	Fludrocortisone acetate.....	79	Galsulfase.....	28
Famotidine.....	9	Fluids and Electrolytes	45	Galvumet	12
Famotidine Hovid.....	9	Flumetasone pivalate	236	Galvus	11
Faslodex.....	165	Fluocortolone caproate with fluocortolone pivalate and cinchocaine.....	8	Gardasil 9	269
Fatty Cream AFT	65	Fluorometholone.....	237	Gastrodenol.....	9
Febuxostat.....	116	Fluorouracil.....	149	Gaviscon Double Strength.....	6
Febuxostat multichem	116	Fluorouracil Accord.....	149	Gaviscon Infant.....	6
Feed Thickener Karicare Aptamil.....	259	Fluorouracil sodium	70	Gazyva	197
FEIBA NF.....	40	Fluox.....	125	Gefitinib	159
Felo 10 ER.....	51	Fluoxetine hydrochloride	125	GEM Aqueous Cream	65
Felo 5 ER.....	51	Flupenthixol decanoate	132	Gemcitabine Ebewe	149
Felodipine.....	51	Flutamide	165	Gemcitabine hydrochloride.....	149
Femme-Tab ED.....	74	Flutamin.....	165	Genoptic	236
Fenpaed 100 mg per 5 ml	110	Fluticasone	229	Gentamicin sulphate Infection	94
Fentanyl.....	122	Fluticasone furoate with vilanterol	229	Sensory.....	236
Fentanyl Sandoz.....	122	Fluticasone propionate	234	Gilenya	135
Ferinject.....	35	Fluticasone with salmeterol	230	Ginet.....	75
Ferodan	35	FML.....	237	Glatiramer acetate.....	135
Ferriprox	242	Foban	62	Glecaprevir with pibrentasvir	102
Ferro-F-Tabs	35	Folic acid	38	Glibenclamide.....	11
Ferro-tab.....	35	Folic Acid multichem.....	38	Gliclazide	11
Ferrograd.....	35	Folic Acid Mylan	38	Glipizide.....	11
Ferrosig	36	Food Thickeners.....	258	Glivec.....	159
Ferrous fumarate.....	35	Foods And Supplements For Inborn Errors Of Metabolism.....	260	Glizide.....	11
Ferrous fumarate with folic acid.....	35	Foradil.....	229	Glucagen Hypokit	10
Ferrous sulfate.....	35	Forteo	113	Glucagon hydrochloride.....	10
Fexofenadine hydrochloride	228	Fortini.....	251	Glucerna Select.....	249
Fibro-vein.....	41	Fortini Multi Fibre.....	251	Glucose [Dextrose].....	45
Filgrastim.....	44	Fortisip.....	257	Gluten Free Foods.....	259
Finasteride.....	76	Fortisip Multi Fibre	257	Glycerin with sodium saccharin	244
Fingolimod	135	Fosamax.....	111	Glycerin with sucrose	244
Firazyr.....	227	Fosamax Plus.....	111	Glycerol	
Flagyl.....	98	Framycetin sulphate	236	Alimentary	26
Flagyl-S.....	98	Frسيوم.....	126	Extemporaneous.....	244
Flamazine.....	62			Glyceryl trinitrate	
Flecainide acetate	49			Alimentary.....	8

Glycopyrronium bromide	8	Humulin 30/70	10	Imuran	168
Glycopyrronium with indacaterol	231	Humulin NPH	10	Incruse Ellipta	231
Go Healthy	28	Humulin R	10	Indacaterol	229
Gold Knight	72	Hyaluronic acid	240	Indapamide	54
Gold Knight XL	72-73	Hydralazine	57	Infanrix IPV	267
Goserelin	87	Hydralazine hydrochloride	57	Infanrix-hexa	268
Gutron	49	Hydrocortisone		Infant Formulae	262
Gynaecological Anti-infectives	75	Dermatological	64	Infatrin	266
- H -		Hormone	79	Infliximab	189
Habitrol	143	Hydrocortisone (PSM)	64	Influenza vaccine	270
Haemophilus influenzae type B vaccine	268	Hydrocortisone acetate	7	Inhaled Corticosteroids	228
Haldol	132	Hydrocortisone acetate with pramoxine hydrochloride	7	Inhaled Long-acting Beta-adrenoceptor Agonists	229
Haldol Concentrate	132	Hydrocortisone and paraffin liquid and lanolin	64	Inspira	53
Haldol Decanoas	132	Hydrocortisone butyrate	64, 70	Instillagel Lido	119
Haloperidol	131	Hydrocortisone with cinchocaine	8	Insulin aspart	11
Haloperidol decanoate	132	Hydrocortisone with miconazole	64	Insulin aspart with insulin aspart protamine	10
Harvoni	102	Hydrocortisone with natamycin and neomycin	64	Insulin glargine	11
Havrix	268	Hydrogen peroxide	61	Insulin glulisine	11
Havrix Junior	268	Hydroxocobalamin	33	Insulin isophane	10
Haylor syrup	228	Hydroxocobalamin Mercury Pharma	33	Insulin isophane with insulin neutral	10
healthE Dimethicone 10%	65	Hydroxocobalamin Panpharma	33	Insulin lispro	11
healthE Dimethicone 4% Lotion	66	hydroxycarbamide	152	Insulin lispro with insulin lispro protamine	11
healthE Dimethicone 5%	65	Hydroxychloroquine	111	Insulin neutral	10
healthE Glycerol BP	244	Hydroxyurea		Insulin pen needles	15
healthE Urea Cream	65	[hydroxycarbamide]	152	Insulin pump	16
Healtheries Simple Baking Mix	259	Hygroton	54	Insulin pump cartridge	21
Hemastix	77	Hylo-Fresh	240	Insulin pump infusion set (steel cannula)	21
Hemilibra	39	Hymenoptera	227	Insulin pump infusion set (steel cannula, straight insertion)	21
Heparin sodium	43	Hyoscine butylbromide	8	Insulin pump infusion set (teflon cannula)	22
Heparinised saline	43	Hyoscine hydrobromide	130	Insulin pump infusion set (teflon cannula, angle insertion with insertion device)	23
Heparon Junior	250	Hypam	137	Insulin pump infusion set (teflon cannula, angle insertion)	23
Hepatitis A vaccine	268	Hyperuricaemia and Antigout	116	Insulin pump infusion set (teflon cannula, straight insertion)	23
Hepatitis B recombinant vaccine	269	Hypromellose	239	Insulin pump reservoir	23
Herceptin	219	Hypromellose with dextran	239	Insulin syringes, disposable with attached needle	16
Hiberix	268	- I -		Intelence	105
Hiprex	109	Ibiamox	92	Interferon beta-1-alpha	135
Histaclear	228	Ibrance	161	Interferon beta-1-beta	135
Histafen	228	Ibuprofen	110	Intra-uterine device	73
Holoxan	146	Icatibant	227	Invega Sustenna	133
Horleys Bread Mix	259	Idarubicin hydrochloride	152	IPCA-Frusamide	53
Horleys Flour	259	Idursulfase	28	IPCA-Metoprolol	50
Hormone Replacement Therapy - Systemic	81	Ifosfamide	146		
HPV	269	Igroton	54		
Humalog	11	Ikorel	57		
Humalog Mix 25	11	Iloprost	60		
Humalog Mix 50	11	Imatinib mesilate	159		
Human papillomavirus (6, 11, 16, 18, 31, 33, 45, 52 and 58) vaccine [HPV]	269	Imatinib-Rex	159		
Humatin	95	Imigran	129		
Humira	183	Imipramine hydrochloride	124		
HumiraPen	183	Imiquimod	70		
		Immune Modulators	106		
		Immunosuppressants	168		

IPCA-Propranolol.....	51	Kindergen.....	250	Levodopa with carbidopa.....	118
IPOL.....	275	Kivexa.....	105	Levomepromazine.....	131
Ipratropium bromide.....	230, 234	Klacid.....		Levomepromazine hydrochloride.....	132
Iressa.....	159	Alimentary.....	9	Levonorgestrel Genito-Urinary.....	74–75
Irinotecan Actavis 100.....	149	Infection.....	90	Hormone.....	82
Irinotecan hydrochloride.....	149	Kliogest.....	82	Levothyroxine.....	83
Irinotecan-Rex.....	149	Kliovance.....	82	Lidocaine [Lignocaine].....	119–120
Iron (as ferric carboxymaltose).....	35	Kogenate FS.....	41	Lidocaine [Lignocaine] hydrochloride.....	120
Iron polymaltose.....	36	Konakion MM.....	41	Lidocaine [Lignocaine] with chlorhexidine.....	120
Isentress.....	106	Konsyl-D.....	25	Lidocaine [Lignocaine] with prilocaine.....	120
Isentress HD.....	106	Kuvan.....	30	Lidocaine-Baxter.....	120
Ismo 20.....	57			Lidocaine-Claris.....	120
Ismo 40 Retard.....	57	- L -		Life Extension.....	31
Isoniazid.....	99	Labetalol.....	50	Lignocaine.....	119–120
Isoniazid with rifampicin.....	99	Lacosamide.....	126	Lioresal Intrathecal.....	117
Isoptin.....	52	Lactulose.....	26	Lipid-Modifying Agents.....	54
Isoptin Retard.....	52	Laevolac.....	26	Liquigen.....	248
Isoptin SR.....	52	Lagevrio.....	103	Lisinopril.....	47
Isopto Carpine.....	239	Lamictal.....	127	Litak.....	148
Isosorbide mononitrate.....	57	Lamivudine.....	100, 105	Lithium carbonate.....	132
Isosource Standard.....	256	Lamivudine Alphapharm.....	105	Livostin.....	238
Isotretinoin.....	61	Lamotrigine.....	127	LMX4.....	120
Ispaghula (psyllium) husk.....	25	Lamprene.....	99	Locacorten-Viaform ED's.....	236
Itch-Soothe.....	63	Lanoxin.....	49	Local preparations for Anal and Rectal Disorders.....	8
Itraconazole.....	96	Lanoxin Paediatric Elixir.....	49	Locasol.....	262
Itrazole.....	96	Lanoxin PG.....	49	Locoid.....	64, 70
Ivacafter.....	233	Lanoxin S29.....	49	Locoid Crelo.....	64
Ivermectin.....	66	Lansoprazole.....	9	Locoid Lipocream.....	64
		Lantus.....	11	Locorten-Vioform.....	236
- J -		Lantus SoloStar.....	11	Lodoxamide.....	238
Jadelle.....	74	Lanvis.....	150	Logem.....	127
Jakavi.....	162	Lanzol Relief.....	9	Lomide.....	238
Jardiamet.....	13	Lapatinib ditosylate.....	160	Lomustine.....	147
Jardiance.....	13	Largactil.....	131	Loniten.....	57
Jaydess.....	82	Laronidase.....	29	Loperamide hydrochloride.....	6
Jevity HiCal RTH.....	256	Lasix.....	53	Lopinavir with ritonavir.....	106
Jevity Plus.....	256	Latanoprost.....	239	Lopinavir/Ritonavir Mylan.....	106
Jevity RTH.....	256	Latanoprost with timolol.....	239	Loprofin.....	261
Juno Pemetrexed.....	150	Lax-Suppositories.....	26	Loprofin Mix.....	261
		Laxatives.....	25	Lorafix.....	228
- K -		Laxsol.....	25	Loratadine.....	228
Kadcyla.....	220	Ledipasvir with sofosbuvir.....	102	Lorazepam.....	134
Kaletra.....	106	Leflunomide.....	111	Lorstat.....	54
Kalydeco.....	233	Lenalidomide.....	152	Losartan Actavis.....	48
Kemadrin.....	119	Letrole.....	167	Losartan potassium.....	48
Kenacomb.....	236	Letrozole.....	167	Losartan potassium with hydrochlorothiazide.....	48
Kenacort-A 10.....	80	Leukeran FC.....	146	Lovir.....	100
Kenacort-A 40.....	80	Leukotriene Receptor Antagonists.....	233	Loxamine.....	125
Kenalog in Orabase.....	33	Leuporelin.....	87	Lucrin Depot 1-month.....	87
Ketocal 3:1.....	266	Leustatin.....	148		
KetoCal 4:1.....	266	Levetiracetam.....	127		
Ketoconazole.....		Levetiracetam-AFT.....	127		
Dermatological.....	70	Levlen ED.....	74		
Infection.....	97	Levocabastine.....	238		
Ketogenic Diet.....	266	Levocarnitine.....	29		
Ketoprofen.....	110	Levodopa with benserazide.....	118		
KetoSens.....	13				
Keytruda.....	222				

Lucrin Depot 3-month.....	87	Mestinon.....	110	Microlut.....	74
Lyderm.....	68	Metabolic Disorder Agents.....	26	Midazolam.....	136
Lynparza.....	153	Metformin hydrochloride.....	11	Midazolam-Baxter.....	136
Lyrica.....	127	Metformin Mylan.....	11	Midodrine.....	49
- M -					
m-Eslon.....	123	Methadone hydrochloride		Mifegyne.....	77
Mabthera.....	200	Extemporaneous.....	244	Mifepristone.....	77
Macro Organic Psyllium Husk.....	25	Nervous.....	122	Minerals.....	34
Macrobid.....	109	Methatabs.....	122	Mini-Wright AFS Low Range.....	234
Macrogol 3350 with potassium		Methenamine (hexamine)		Mini-Wright Standard.....	234
chloride, sodium bicarbonate and		hippurate.....	109	Minidiab.....	11
sodium chloride.....	26	Methopt.....	239	MiniMed 1.8 Reservoir	
Macrogol 400 and propylene		Methotrexate.....	149	MMT-326A.....	23
glycol.....	240	Methotrexate DBL Onco-Vial.....	149	MiniMed 3.0 Reservoir	
Madopar 125.....	118	Methotrexate Ebewe.....	149	MMT-332A.....	23
Madopar 250.....	118	Methotrexate Sandoz.....	149	MiniMed 770G.....	16
Madopar 62.5.....	118	Methyl hydroxybenzoate.....	244	MiniMed Mio MMT-921A.....	22
Madopar HBS.....	118	Methylcellulose.....	244	MiniMed Mio MMT-923A.....	22
Madopar Rapid.....	118	Methylcellulose with glycerin and		MiniMed Mio MMT-925A.....	22
Magnesium hydroxide.....	36	sodium saccharin.....	244	MiniMed Mio MMT-941A.....	22
Magnesium sulphate.....	36	Methylcellulose with glycerin and		MiniMed Mio MMT-943A.....	22
Mantoux.....	276	sucrose.....	244	MiniMed Mio MMT-945A.....	22
Marevan.....	44	Methylidopa.....	52	MiniMed Mio MMT-965A.....	22
Marine Blue Lotion SPF 50+.....	70	Methylidopa Mylan.....	52	MiniMed Mio MMT-975A.....	22
Martindale Pharma.....	241	Methylidopa Mylan S29.....	52	MiniMed Quick-Set MMT-386A.....	22
Mask for spacer device.....	234	Methylnaltrexone bromide.....	25	MiniMed Quick-Set MMT-387A.....	22
Maviret.....	102	Methylphenidate ER - Teva.....	138	MiniMed Quick-Set MMT-396A.....	22
Maxidex.....	237	Methylphenidate hydrochloride.....	138	MiniMed Quick-Set MMT-397A.....	22
Maxitrol.....	237	Methylphenidate hydrochloride		MiniMed Quick-Set MMT-398A.....	22
MCT oil (Nutricia).....	248	extended-release.....	139	MiniMed Quick-Set MMT-399A.....	22
Measles, mumps and rubella		Methylprednisolone.....	79	MiniMed Silhouette MMT-368A.....	22
vaccine.....	272	Methylprednisolone (as sodium		MiniMed Silhouette MMT-377A.....	22
Meclizine.....	89	succinate).....	80	MiniMed Silhouette MMT-378A.....	22
Meclizine.....	89	Methylprednisolone aceponate.....	64	MiniMed Silhouette MMT-381A.....	22
Mebeverine hydrochloride.....	8	Methylprednisolone acetate.....	80	MiniMed Silhouette MMT-382A.....	22
Medrol.....	79	Methylxanthines.....	233	MiniMed Silhouette MMT-383A.....	22
Medroxyprogesterone acetate		Metoclopramide Actavis 10.....	130	MiniMed Silhouette MMT-384A.....	22
Genito-Urinary.....	74	Metoclopramide hydrochloride.....	130	MiniMed Sure-T MMT-864A.....	21
Hormone.....	81-82	Metolazone.....	53	MiniMed Sure-T MMT-866A.....	21
Mefenamic acid.....	110	Metopirone.....	88	MiniMed Sure-T MMT-874A.....	21
Megace.....	166	Metoprolol IV Mylan.....	50	MiniMed Sure-T MMT-876A.....	21
Megestrol acetate.....	166	Metoprolol succinate.....	50	MiniMed Sure-T MMT-884A.....	21
Melatonin.....	136	Metoprolol tartrate.....	50	MiniMed Sure-T MMT-886A.....	21
Melphalan.....	147	Metrogyl.....	98	Minims Pilocarpine.....	239
Menactra.....	272	Metronidazole.....	98	Minims Prednisolone.....	238
Meningococcal (groups A, C, Y and		Metyrapone.....	88	Minirin.....	88
W-135) conjugate vaccine.....	272	Mexiletine hydrochloride.....	49	Minirin Melt.....	88
Meningococcal B multicomponent		Miacalcin.....	78	Mino-tabs.....	93
vaccine.....	273	Micolette.....	26	Minocycline hydrochloride.....	93
Meningococcal C conjugate		Miconazole.....	33	Minomycin.....	93
vaccine.....	273	Miconazole nitrate		Minor Skin Infections.....	66
Menthol.....	63	Dermatological.....	63	Minoxidil.....	57
Mepolizumab.....	197	Genito-Urinary.....	75	Mirena.....	82
Mercaptopurine.....	149	Micreme.....	75	Mirtazapine.....	125
Mercilon 28.....	73	Micreme H.....	64	Misoprostol.....	8
Mesalazine.....	7	Microgynon 20 ED.....	74	Mitomycin C.....	153
Mesna.....	153	Microgynon 30.....	74	Mitozantrone.....	153

Mitozantrone Ebewe.....	153	Naphazoline hydrochloride.....	240	Norethisterone	
Mixtard 30.....	10	Naphcon Forte.....	240	Genito-Urinary.....	75
MMR II.....	272	Naprosyn SR 1000.....	110	Hormone.....	82
Moclobemide.....	125	Naprosyn SR 750.....	110	Norflex.....	117
Modafinil.....	140	Naproxen.....	110	Norflexacin.....	109
Modavigil.....	140	Narcarcin mite.....	116	Noriday 28.....	75
Moduretic.....	53	Nasal Preparations.....	234	Norimin.....	74
Molaxole.....	26	Natalizumab.....	136	Normacol Plus.....	25
Molnupiravir.....	103	Natulan.....	155	Normison.....	137
Moments.....	72	Nausafix.....	130	Norpress.....	124
Mometasone furoate.....	64	Nausicalm.....	130	Nortriptyline hydrochloride.....	124
Monogen.....	249	Navelbine.....	157	Norvir.....	106
Montelukast.....	233	Nefopam hydrochloride.....	120	Noured.....	125
Montelukast Mylan.....	233	Neisvac-C.....	273	Noured Paracetamol.....	121
Morotocog alfa [Recombinant factor VIII].....	40	Neo-B12.....	33	NovaSource Renal.....	252
Morphine hydrochloride.....	122	Neo-Mercazole.....	83	Novatretin.....	68
Morphine sulphate.....	123	Neocate Gold.....	262	NovoMix 30 FlexPen.....	10
Motetis.....	119	Neocate Junior Unflavoured.....	262	NovoRapid.....	11
Mouth and Throat.....	32	Neocate Junior Vanilla.....	262	NovoRapid FlexPen.....	11
Movapo.....	118	Neocate SYNEO.....	262	NovoRapid Penfill.....	11
Moxifloxacin.....	94	Neoral.....	223	NovoSeven RT.....	40
MSUD Maxamum.....	260	Neostigmine metilsulfate.....	110	Noxafil.....	97
Mucilaginous laxatives with stimulants.....	25	Nepro HP (strawberry).....	252	Nozinan.....	131-132
Mucolytics.....	233	Nepro HP (vanilla).....	252	Nozinan (Swiss).....	131
Mucosoothe.....	120	Nepro HP RTH.....	251	Nucala.....	197
Multiple Sclerosis Treatments.....	134	Neulactil.....	132	Nuelin.....	233
Multivitamin renal.....	34	Neulastim.....	44	Nuelin-SR.....	233
Multivitamins.....	34	NeuroTabs.....	35	Nupentin.....	126
Mupirocin.....	62	Nevirapine.....	105	Nutilis.....	259
Muscle Relaxants.....	117	Nevirapine Alphapharm.....	105	Nutren Diabetes.....	249
Mvite.....	34	Nicorandil.....	57	Nutrient Modules.....	246
Myambutol.....	99	Nicotine.....	143	Nutrini Energy Multi Fibre.....	251
Mycobutin.....	100	Nifedipine.....	51	Nutrini Energy RTH.....	251
MycoNail.....	62	Nifuran.....	109	Nutrini Low Energy Multi Fibre.....	253
Mycophenolate mofetil.....	168	Nilotinib.....	160	Nutrini Peptisorb.....	264
Mydriacyl.....	239	Nilstat.....		Nutrini Peptisorb Energy.....	264
Mylan (12 hr release).....	51	Alimentary.....	33	Nutrini RTH.....	251
Mylan (24 hr release).....	51	Genito-Urinary.....	75	Nutrison 800 Complete Multi Fibre.....	256
Mylan Atenolol.....	50	Infection.....	97	Nutrison Advanced Diason.....	249
Mylan Clomiphen.....	88	Nintedanib.....	231	Nutrison Advanced Peptisorb.....	252
Mylan Indapamide.....	54	Nipent.....	154	Nutrison Concentrated.....	258
Mylan Italy (24 hr release).....	51	Nirmatrelvir with ritonavir.....	104	Nutrison Energy.....	256
Myleran.....	146	Nitrates.....	56	Nutrison Energy Multi Fibre.....	256
Myometrial and Vaginal Hormone Preparations.....	75	Nitroderm TTS.....	56	Nutrison Multi Fibre.....	256
Myozyme.....	26	Nitrofurantoin.....	109	Nutrison Standard RTH.....	256
- N -		Nitrolingual Pump Spray.....	56	Nyefax Retard.....	51
Nadolol.....	50	Nivestim.....	44	Nystatin	
Nadolol BNM.....	50	Nivolumab.....	221	Alimentary.....	33
Naglazyme.....	28	Nizoral.....	97	Genito-Urinary.....	75
Nalcrom.....	8	Nodia.....	6	Infection.....	97
Naloxone hydrochloride.....	241	Noflam 250.....	110	NZB Low Gluten Bread Mix.....	259
Naltraccord.....	142	Noflam 500.....	110	- O -	
Naltrexone hydrochloride.....	142	Non-Steroidal Anti-Inflammatory Drugs.....	110	O/W Fatty Emulsion Cream.....	65
		Nonacog gamma, [Recombinant Factor IX].....	40	Obinutuzumab.....	197
				Obstetric Preparations.....	77

Ocrelizumab	136	Other Progestogen	Pazopanib.....	161	
Ocrevus	136	Preparations	82	Peak flow meter.....	234
Octocog alfa [Recombinant factor		Other Skin Preparations	70	Pedialyte - Bubblegum	45
VIII] (Advate).....	41	Ovestin		Pediasure	251
Octocog alfa [Recombinant factor		Genito-Urinary.....	75	Pediasure RTH.....	251
VIII] (Kogenate FS).....	41	Hormone	82	Pegaspargase	154
Octreotide	166	Oxaliplatin.....	147	Pegasy.....	107
Octreotide Depot Teva	166	Oxaliplatin Accord.....	147	Pegfilgrastim.....	44
Octreotide long-acting	166	Oxaliplatin Actavis 100	147	Pegylated interferon alfa-2a	107
Oestradiol	81	Oxaliplatin Ebewe.....	147	Pembrolizumab.....	222
Oestradiol valerate	81	Oxis Turbuhaler	229	Pemetrexed.....	150
Oestradiol with norethisterone.....	82	Oxpentifylline	57	Penicillamine	111
Oestriol		Oxybutynin.....	76	Penicillin G.....	92
Genito-Urinary.....	75	Oxycodone hydrochloride.....	123	PenMix 30.....	10
Hormone	82	Oxycodone Sandoz	123	PenMix 40.....	10
Oestrogens	81	OxyNorm	123	PenMix 50.....	10
Ofev	231	Oxytocin.....	75	Pentasa	7
Oil in water emulsion	65	Oxytocin BNM.....	75	Pentostatin [Deoxycorymycin].....	154
Olanzapine	132-133	Oxytocin with ergometrine		Pentoxifylline [Oxpentifylline].....	57
Olaparib	153	maleate.....	75	Peptamen Junior	251
Olbetam	54	Ozurdex.....	237	Peptisorb	252
Olbetam S29.....	54			Perhexiline maleate.....	52
Olopatadine	240	- P -		Pericyazine.....	132
Olopatadine Teva	240	Pacifen.....	117	Perindopril	47
Olisalazine	7	Pacimol.....	121	Perjeta	200
Omalizumab	198	Paclitaxel	154	Permethrin	68
Omeprazole	9	Paclitaxel Actavis.....	154	Perrigo	70
Omeprazole actavis 10.....	9	Paclitaxel Ebewe.....	154	Pertuzumab	200
Omeprazole actavis 20	9	Paediatric Seravit	34	Peteha	99
Omeprazole actavis 40	9	Palbociclib	161	Pethidine hydrochloride	124
Omnitrope	83	Paliperidone.....	133	Pevaryl.....	62
Onbrez Breezhaler	229	Palivizumab	199	Pexsig.....	52
Oncaspar LYO	154	Pamidronate disodium	112	Pfizer Exemestane	167
OncoTICE	175	Pamisol	112	Pharmacy Health Sorbolene with	
Ondansetron.....	130	Pancreatic enzyme	24	Glycerin	65
Ondansetron ODT-DRLA	130	Pantoprazole	9	Pharmacy Services.....	241
One-Alpha	33	Panzop Relief	9	Pheburane	30
Onrex	130	Panzyltrat	24	Phenaxen	150
Opdivo	221	Papaverine hydrochloride.....	57	Phenobarbitone	127
Ora-Blend	244	Para-amino salicylic acid	99	Phenobarbitone sodium	
Ora-Blend SF.....	244	Paracare	121	Extemporaneous.....	244
Ora-Plus	244	Paracare Double Strength	121	Nervous.....	136
Ora-Sweet	244	Paracetamol	121	Phenoxybenzamine	
Ora-Sweet SF.....	244	Paracetamol + Codeine		hydrochloride	47
Orabase	32	(Relieve)	123	Phenoxyethylpenicillin (Penicillin	
Oral and Enteral Feeds	249	Paracetamol with codeine	123	V).....	93
Oratane	61	Paraffin	65-66	Phenytoln sodium.....	126-127
Ordine	122	Paraffin liquid with wool fat	240	Phillips Milk of Magnesia	36
Orgran	260	Parasiticial Preparations.....	66	Phlexy 10.....	261
Ornidazole	98	Parlodol	118	Phosphate Phebra.....	46
Orphenadrine citrate.....	117	Parnate	124	Phosphorus	46
Ortho-tolidine	77	Parnate S29.....	124	Phytomenadione.....	41
Oruvail SR	110	Paromomycin.....	95	Pilocarpine hydrochloride	239
Osmolite RTH.....	256	Paroxetine	125	Pimafucort	64
Other Endocrine Agents	88	Paser	99	Pimecolimus.....	69
Other Oestrogen Preparations	82	Paxam	134	Pine tar with trolamine laurilsulfate	
		Paxlovid.....	104		

and fluorescein	69	Pregnancy Tests - hCG Urine	76	hydrochlorothiazide	48
Pinetarsol.....	69	Premarin.....	81	Qvar.....	228
Pioglitazone.....	11	Prevenar 13.....	274	- R -	
Pirfenidone.....	232	Priadel.....	132	RA-Morph.....	122
Pizotifen.....	129	Primaquine.....	98	Raloxifene hydrochloride.....	112
PKU Anamix Infant.....	261	Primidone.....	127	Raltegravir potassium.....	106
PKU Anamix Junior.....	261	Primidone Clinect.....	127	Ramipex.....	118
PKU Anamix Junior Chocolate.....	261	Primolut N.....	82	Ranbaxy-Cefaclor.....	89
PKU Anamix Junior LQ.....	261	Priorix.....	272	Ranbaxy-Cefaclor S29.....	89
PKU Anamix Junior Orange.....	261	Probenecid.....	117	Rapamune.....	224
PKU Anamix Junior Vanilla.....	261	Probenecid-AFT.....	117	Rasagiline.....	118
PKU Lophlex LQ 10.....	261	Procaine penicillin.....	93	Reandron 1000.....	81
PKU Lophlex LQ 20.....	261	Procabazine hydrochloride.....	155	Recombinant factor IX.....	38, 40
PKU Lophlex Powder.....	261	Prochlorperazine.....	130	Recombinant factor VIIa.....	40
PKU Lophlex Sensation 20.....	261	Proctofoam.....	7	Recombinant factor VIII.....	40–41
Plaquenil.....	111	Proctosedyl.....	8	Rectogesic.....	8
Plendil ER.....	51	Procyclidine hydrochloride.....	119	Redipred.....	80
Pneumococcal (PCV10) conjugate vaccine.....	273	Progesterone.....	82	Relieve.....	110
Pneumococcal (PCV13) conjugate vaccine.....	274	Proglicem.....	10	Relistor.....	25
Pneumococcal (PPV23) polysaccharide vaccine.....	275	Proglycem.....	10	Remicade.....	189
Pneumovax 23.....	275	Prodynova.....	81	Renilon 7.5.....	252
Podophyllotoxin.....	70	Prolia.....	111	Resonium-A.....	46
Polaramine.....	228	Promethazine hydrochloride.....	228	Resource Beneprotein.....	248
Poliomyelitis vaccine.....	275	Propafenone hydrochloride.....	49	Respigen.....	230
Poloxamer.....	25	Propamide isethionate.....	236	Respiratory Devices.....	234
Poly-Gel.....	240	Propranolol.....	51	Respiratory Stimulants.....	235
Poly-Tears.....	239	Propylene glycol.....	244	Retinol palmitate.....	240
Poly-Visc.....	240	Propylthiouracil.....	83	ReTrieve.....	61
Polycal.....	246	Prostacur.....	165	Retrovir.....	105
Ponstan.....	110	Protaphane.....	10	Revlimid.....	152
Posaconazole.....	97	Protaphane Penfill.....	10	Revolade.....	38
Postinor-1.....	75	Protifar.....	248	Rexacrom.....	238
Potassium chloride.....	45–46	Protionamide.....	99	Riboflavin.....	29
Potassium citrate.....	76	Provera.....	81	Ribomustin.....	145
Potassium iodate.....	35	Provera HD.....	82	Ricit.....	76
Povidone iodine.....	66	PSM Citalopram.....	125	Rifabutin.....	100
Pradaxa.....	44	Psoriasis and Eczema Preparations.....	68	Rifadin.....	100
Pramipexole hydrochloride.....	118	PTU.....	83	Rifampicin.....	100
Pravastatin.....	54	Pulmicort Turbuhaler.....	228	Rifaximin.....	9
Pravastatin Mylan.....	54	Pulmozyme.....	233	Rifinah.....	99
Praziquantel.....	89	Puri-nethol.....	149	Rilutek.....	119
Prazosin.....	47	Puria.....	34	Riluzole.....	119
Pred Forte.....	238	Pyrazinamide.....	100	RINVOQ.....	225
Prednisolone.....	80	Pyridostigmine bromide.....	110	Riodine.....	66
Prednisolone acetate.....	238	Pyridoxine hydrochloride.....	33	Risedronate Sandoz.....	113
Prednisolone sodium.....	8	Pyridoxine multichem.....	33	Risedronate sodium.....	113
Prednisolone sodium phosphate.....	238	Pyrimethamine.....	95	Risperdal Consta.....	133
Prednisolone-AFT.....	238	Pytazen SR.....	41	Risperidone.....	132–133
Prednisone.....	80	- Q -		Risperidone (Teva).....	132
Prednisone Clinect.....	80	Quetapel.....	132	Risperon.....	132
Pregabalin.....	127	Quetiapine.....	132	Ritalin.....	138
Pregabalin Pfizer.....	127	Quick-Set MMT-392.....	23	Ritalin LA.....	139
		Quick-Set MMT-393.....	23	Ritonavir.....	106
		Quinapril.....	48	Rituximab (Mabthera).....	200
		Quinapril with		Rituximab (Riximyo).....	202
				Rivaroxaban.....	44

Rivastigmine	141	Seretide	230	Solgar	27–29, 31
Rivastigmine Patch BNM 10	141	Seretide Accuhaler	230	Solifenacin Mylan	77
Rivastigmine Patch BNM 5	141	Serevent	229	Solifenacin succinate	77
Rivotril	126	Serevent Accuhaler	229	Solu-Cortef	79
Riximyo	202	Sertraline	125	Solu-Medrol	80
RIXUBIS	40	Setrona	125	Solu-Medrol-Act-O-Vial	80
Rizamelt	129	Setrona AU	125	Somatropin (Omnitrope)	83
Rizatriptan	129	Sevredol	123	Sotalol	51
RoActemra S29	215	Sex Hormones Non Contraceptive	80	Spacer device	234
Ronapreve	189	Shield XL	72	Span-K	46
Ropin	118	shingles vaccine	276	Spolto Respimat	231
Ropinirole hydrochloride	118	SII-Onco-BCG	175	Spiractin	53
Rosuvastatin	55	Sildenafil	59	Spiriva	231
Rosuvastatin Viatris	55	Silhouette MMT-373	23	Spiriva Respimat	231
Rotarix	275	Siltuximab	215	Spironolactone	53
Rotavirus oral vaccine	275	Simvastatin	55	Sporanox	96
Roxane	6	Simvastatin Mylan	55	Sprycel	158
Roxane-Propranolol	51	Sinemet	118	Stemetil	130
Roxithromycin	91	Sinemet CR	118	SteroClear	234
Rubifen	138	Sirolimus	224	Stesolid	126
Rubifen SR	138	Siterone	80	Stimulants/ADHD Treatments	137
Rugby Capsaicin Topical Cream	120	Slow-Lopresor	50	Stiripentol	128
Rulide D	91	Smith BioMed Rapid Pregnancy Test	76	Stocrin	105
Rurioctocog alfa pegol [Recombinant factor VIII]	41	Sodibic	46	Stomahesive	32
Ruxolitinib	162	Sodium acid phosphate	26	Strattera	137
Rythmodan	49	Sodium alginate	6	Strides Shasun	97
Rytmonorm	49	Sodium benzoate	30	Stromectol	66
- S -		Sodium bicarbonate Blood	45–46	Sucrafate	9
Sabril	128	Extemporaneous	244	Sulfadiazine Silver	62
Sacubitril with valsartan	48	Sodium calcium edetate	242	Sulfadiazine sodium	95
Sagent	157	Sodium chloride Blood	45	Sulfasalazine	8
SalAir	230	Respiratory	234	Sulphur	69
Salazopyrin	8	Sodium citrate with sodium lauryl sulphoacetate	26	Sulprix	131
Salazopyrin EN	8	Sodium citro-tartrate	77	Sumagran	129
Salbutamol	230	Sodium cromoglicate	8	Sumatriptan	129
Salbutamol with ipratropium bromide	230	Alimentary	238	Sunitinib	163
Salicylic acid	69	Sensory	236	Sunitinib Pfizer	163
Salmeterol	229	Sodium fluoride	35	Sunscreens	70
Sandomigran	129	Sodium Fusidate [fusidic acid] Dermatological	62	Sunscreens, proprietary	70
Sanofi Primaquine	98	Infection	95	Sure-T MMT-863	21
Sapropterin dihydrochloride	30	Sensory	236	Sure-T MMT-873	21
Scalp Preparations	70	Sodium hyaluronate [Hyaluronic acid]	240	Sustagen Hospital Formula	256
Scopoderm TTS	130	Sodium phenylbutyrate	30	Sustagen Hospital Formula Active	256
Sebizole	70	Sodium picosulfate	26	Sustanon Ampoules	81
Secukinumab	213	Sodium polystyrene sulphonate	46	Sutant	163
Sedatives and Hypnotics	136	Sodium tetradecyl sulphate	41	Sylvant	215
Seebri Breezhaler	231	Sodium valproate	128	Symbicort Turbuhaler 100/6	229
Selegiline hydrochloride	118	Sofradex	236	Symbicort Turbuhaler 200/6	229
Senna	26	Soframycin	236	Symbicort Turbuhaler 400/12	229
Senokot	26			Symmetrel	118
SensoCard	15			Sympathomimetics	57
Serc	130			Synacthen	80
Serenace	131			Synacthen Depot	80
				Synacthene Retard	80
				Synagis	199

Synflorix.....	273	Ticagrelor.....	41	[Co-trimoxazole].....	96
Synthroid.....	83	Tilcotil.....	110	Trisequens.....	82
Syntometrine.....	75	Timolol.....	238	Trisol.....	96
Syrup (pharmaceutical grade).....	245	Timoptol XE.....	238	Trophic Hormones.....	83
Systane Unit Dose.....	240	Tiotropium bromide.....	231	Tropicamide.....	239
- T -					
Tacrolimus		Tiotropium bromide with olodaterol.....	231	Trulicity.....	12
Dermatological.....	69	Tivicay.....	106	Trusopt.....	238
Oncology.....	225	TMP.....	95	TruSteel.....	21
Tacrolimus Sandoz.....	225	Tobramycin		Tuberculin PPD [Mantoux] test.....	276
Taliglucerase alfa.....	31	Infection.....	95	Tubersol.....	276
Tambacor.....	49	Sensory.....	236	Two Cal HN.....	258
Tamoxifen citrate.....	167	Tobramycin BNM.....	95	Tykerb.....	160
Tamoxifen Sandoz.....	167	Tobramycin Mylan.....	95	Tysabri.....	136
Tamsulosin hydrochloride.....	76	Tobrex.....	236	- U -	
Tamsulosin-Flex.....	76	Tocilizumab.....	215	Ultibro Breezhaler.....	231
Tandem Cartridge.....	21	Tofranil.....	124	Ultraproct.....	8
Tandem t:slim X2 with Basal-IQ.....	16	Tolcapone.....	118	Umeclidinium.....	231
Tap water.....	245	Topamax.....	128	Umeclidinium with vilanterol.....	231
Tarceva.....	158	Topical Products for Joint and Muscular Pain.....	111	Univent.....	230, 234
Tasigna.....	160	Topiramate.....	128	Upadacitinib.....	225
Tasmar.....	118	Topiramate Actavis.....	128	Ural.....	77
Taurine.....	31	Total parenteral nutrition (TPN).....	45	Urea.....	65
Tecfidera.....	135	TPN.....	45	Urex Forte.....	53
Tegretol.....	126	Tramadol hydrochloride.....	124	Urinary Agents.....	76
Tegretol CR.....	126	Tramal SR 100.....	124	Urinary Tract Infections.....	109
Telfast.....	228	Tramal SR 150.....	124	Urinorm.....	116
Teligent.....	94	Tramal SR 200.....	124	Uromitexan.....	153
Temaccord.....	155	Trandate.....	50	Ursodeoxycholic acid.....	24
Temazepam.....	137	Tranexamic acid.....	41	Ursosan.....	24
Temozolomide.....	155	Tranlycypromine sulphate.....	124	Utrogestan.....	82
Tenofovir disoproxil.....	100	Trastuzumab.....	219	- V -	
Tenofovir Disoproxil Teva.....	100	Trastuzumab emtansine.....	220	Vaccinations.....	267
Tenoxicam.....	110	Travatan.....	239	Vaclovir.....	100
Tensipine MR10.....	51	Travoprost.....	239	Valaciclovir.....	100
Tepadina.....	147	Treatments for Dementia.....	140	Valganciclovir.....	101
Terbinafine.....	97	Treatments for Substance Dependence.....	141	Valganciclovir Mylan.....	101
Terbutaline sulphate.....	230	Trental 400.....	57	Vancomycin.....	96
Teriflunomide.....	136	Tretinoin		Vannair.....	229
Teriparatide.....	113	Dermatological.....	61	Varenicline Pfizer.....	143
Testosterone.....	80	Oncology.....	156	Varenicline tartrate.....	143
Testosterone cipionate.....	80	Trexate.....	149	Varicella vaccine [Chickenpox vaccine].....	276
Testosterone esters.....	81	Triamcinolone acetonide		Varicella zoster virus (Oka strain) live attenuated vaccine [shingles vaccine].....	276
Testosterone undecanoate.....	81	Alimentary.....	33	Various.....	241
Tetrabenazine.....	119	Dermatological.....	64	Varivax.....	276
Tetrabromophenol.....	77	Hormone.....	80	Vasodilators.....	57
Tetracosactrin.....	80	Triamcinolone acetonide with gramicidin, neomycin and nystatin		Vasopressin Agonists.....	88
Tetracycline.....	93	Dermatological.....	64	Vasorex.....	51
Thalidomide.....	156	Sensory.....	236	Vedafil.....	59
Thalomid.....	156	Triazolam.....	137	Veletri.....	60
Theophylline.....	233	Trimethoprim.....	95	Venclexta.....	156
Thiamine hydrochloride.....	33	Trimethoprim with sulphamethoxazole		Venetoclax.....	156
THIO-TEPA.....	147			Venlafaxine.....	125
Thioguanine.....	150			Venomil.....	227
Thiotepa.....	147				
Thyroid and Antithyroid Agents.....	83				

VENOX.....	227	Xylocaine.....	120
Ventavis.....	60	Xylocaine 2% Jelly.....	119
Ventolin.....	230	Xyntha.....	40
Vepesid.....	152	- Z -	
Verapamil hydrochloride.....	52	Zapril.....	47
Vergo 16.....	130	Zarontin.....	126
Vermox.....	89	Zaroxolyn.....	53
Versacloz.....	131	Zavedos.....	152
Vesanoid.....	156	Zeffix.....	100
Vexazone.....	11	Zematop.....	69
Vfend.....	97	Zetlam.....	100
Viaderm KC.....	64	Ziagen.....	105
Vidaza.....	147	Zidovudine [AZT].....	105
Vigabatrin.....	128	Zidovudine [AZT] with	
Vigisom.....	136	lamivudine.....	106
Vildagliptin.....	11	Zimybe.....	56
Vildagliptin with metformin		Zinc and castor oil.....	65
hydrochloride.....	12	Zinc sulphate.....	36
Vimpat.....	126	Zincaps.....	36
Vinblastine sulphate.....	157	Zinnat.....	89
Vincristine sulphate.....	157	Ziprasidone.....	132
Vinorelbine.....	157	Zista.....	228
Vinorelbine Ebewe.....	157	Zithromax.....	90
Viramune Suspension.....	105	Zoledronic acid	
ViruPOS.....	236	Hormone.....	78
Vit.D3.....	34	Musculoskeletal.....	114
Vita-B12.....	33	Zoledronic acid Mylan.....	78
Vita-POS.....	240	Zopiclone.....	137
Vitabdeck.....	34	Zopiclone Actavis.....	137
Vital.....	252	Zostavax.....	276
Vitamin B complex.....	33	Zostrix.....	111
Vitamin B6 25.....	33	Zostrix HP.....	120
Vitamins.....	33-34	Zuclopenthixol decanoate.....	134
Vivonex TEN.....	252	Zuclopenthixol hydrochloride.....	132
Voltaren.....	110	Zusdone.....	132
Voltaren D.....	110	Zyban.....	142
Voltaren Ophtha.....	237	Zypine.....	132
Voltaren SR.....	110	Zypine ODT.....	132
Volumatic.....	234	Zyprexa Relprevv.....	133
Voriconazole.....	97	Zytiga.....	164
Votrient.....	161		
Vttack.....	97		
- W -			
Warfarin sodium.....	44		
Wart Preparations.....	70		
Wasp venom allergy treatment.....	227		
Water			
Blood.....	45		
Extemporaneous.....	245		
Wool fat with mineral oil.....	66		
- X -			
Xarelto.....	44		
Xifaxan.....	9		
XMET Maxamum.....	260		
Xolair.....	198		
XP Maxamum.....	261		



















