July 2022 Volume 29 Number 1

Editors:

Kaye Wilson, & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Facsimile +64 4 460 4995 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

Accessible in an electronic format at no cost from the Pharmac website www.pharmac.govt.nz/schedule.

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering
email: texschedule@pharmac.govt.nz
©Pharmaceutical Management Agency
ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

	introducing Fharmac	
Section A	General Rules	5
Section B	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	37
	Cardiovascular System	47
	Dermatologicals	61
	Genito Urinary System	71
	Hormone Preparations – Systemic	78
	Infections – Agents For Systemic Use	89
	Musculoskeletal System	110
	Nervous System	118
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	227
	Sensory Organs	236
	Various	241
Section C	Extemporaneous Compounds (ECPs)	243
Section D	Special Foods	246
Section I	National Immunisation Schedule	267
	Index	277

Introducing Pharmac

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION R. ALIMENTARY TRACT AND METAROLISM

SECTION B: ALIMENTARY TRACT AND ME	TABOLISM			
	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
ALGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg p sachet		30	√ (Gaviscon Infant
** Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour	1.80 (8.60)	60	(Gaviscon Double Strength
* Oral liq 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg per 10 ml		500 ml	ļ	Acidex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE * Tab 600 mg	39.00	100 500 ml	√ F	Alu-Tab Roxane
inappropriate and the prescription is endorsed according		.5 OI WI	iere calciu	iii carbonale labiels are
Antidiarrhoeals				
Agents Which Reduce Motility				
LOPERAMIDE HYDROCHLORIDE – Up to 30 cap available on * Tab 2 mg* * Cap 2 mg	10.75	400 400		Nodia Diamide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy SA1886 Special Authority for Subsidy Initial application — (Crohn's disease) from any relevant prac		90 alid for		Entocort CIR
the following criteria:				1,1

Both:

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*: and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam S29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa
OLSALAZINE		
Tab 500 mg93.37	100	Dipentum
Cap 250 mg53.00	100	✓ Dipentum

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PREDNISOLONE SODIUM				
Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone \$29
SODIUM CROMOGLICATE Cap 100 mgSULFASALAZINE	92.91	100	•	Nalcrom
* Tab 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND CIT	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and		
cinchocaine hydrochloride 5 mg per g11.06	30 g OP	Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and		
cinchocaine hydrochloride 1 mg	12	Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

GLYCERYL TRINITRATE - Special Authority see SA1329 below - Retail pharmacy

* Oint 0.2%......22.00 30 g OP

* Rectogesic

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule – Up to 10 inj available on a PSO	65.45	10	✓ Max Health
HYOSCINE BUTYLBROMIDE * Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml – Up to 5 inj available on a PSO		5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE * Tab 135 mg	9.20	90	✓ Colofac

Antiulcerants

Antisecretory and Cytoprotective

MIS	SOPROSTOL		
*	Tab 200 mcg - Up to 120 tab available on a PSO41.50	120	Cytotec

		ALIMENTARY	TR	ACT AN	D METABOLISM
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Н	elicobacter Pylori Eradication				
CL	ARITHROMYCIN Tab 500 mg — Subsidy by endorsement	eradication and prescri		is endorse	
Н	2 Antagonists				
	MOTIDINE – Only on a prescription Tab 20 mg	4.91	100	1	Famotidine Hovid S29
*	Tab 40 mg	8.48	100	✓	Famotidine Hovid \$29
*	Inj 10 mg per ml, 4 ml - Subsidy by endorsement		10 of pa		Mylan §29 re.
P	roton Pump Inhibitors				
*	NSOPRAZOLE Cap 15 mg Cap 30 mg IEPRAZOLE For omeprazole suspension refer Standard Formulae, page	5.26	100 100		Lanzol Relief Lanzol Relief
*	Cap 10 mg		90	•	Omeprazole actavis
*	Cap 20 mg	1.86	90	✓	Omeprazole actavis 20
*	Cap 40 mg	3.11	90	/	Omeprazole actavis 40
*	Powder – Only in combination Only in extemporaneously compounded omeprazole su		5 g	✓	Midwest
*	Inj 40 mg ampoule with diluent		5	✓	Dr Reddy's Omeprazole
	NTOPRAZOLE	0.00	100	,	Dannan Daliaf
	Tab EC 20 mg		100 100		Panzop Relief Panzop Relief
S	ite Protective Agents				
	LLOIDAL BISMUTH SUBCITRATE Tab 120 mg	14.51	50	/	Gastrodenol S29
SU	CRALFATE Tab 1 g	35.50	120		Countries

Bile and Liver Therapy

(48.28)

Carafate

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	sidised	Generic	
\$	Per	1	Manufacturer	

⇒SA1461 Special Authority for Subsidy

Initial application only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid for 6 months where the patient has hepatic encephalopathy despite an adequate trial of maximum tolerated doses of lactulose.

Renewal only from a gastroenterologist, hepatologist or Practitioner on the recommendation of a gastroenterologist or hepatologist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Diabetes

Hyperglycaemic Agents

DIAZOXIDE - Special Authority see SA1320 below - Retail pha	armacy		
Cap 25 mg	110.00	100	✓ Proglicem S29
Cap 100 mg	280.00	100	✓ Proglicem S29
Oral liq 50 mg per ml	620.00	30 ml OP	✓ Proglycem S29
			✓ e5 Pharma S29

⇒SA1320 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where used for the treatment of confirmed hypoglycaemia caused by hyperinsulinism.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

GLUCAGON HYDROCHLORIDE

Inj 1 mg syringe kit	- Up to 5 kit available on a PSO	32.00 1	•	Glucagen Hypokit
----------------------	----------------------------------	---------	---	------------------

Insulin - Short-acting Preparations

INS	SULIN NEUTRAL		
lack	Inj human 100 u per ml25.26	10 ml OP	✓ Actrapid
			✓ Humulin R
\blacktriangle	Inj human 100 u per ml, 3 ml42.66	5	Actrapid Penfill
			✓ Humulin R

Insulin - Intermediate-acting Preparations

INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	✓ NovoMix 30 FlexPen
INSULIN ISOPHANE			•
▲ Inj human 100 u per ml	17.68	10 ml OP	✓ Humulin NPH✓ Protaphane
▲ Inj human 100 u per ml, 3 ml	29.86	5	✓ Humulin NPH
·			Protaphane Penfill
INSULIN ISOPHANE WITH INSULIN NEUTRAL			
▲ Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70
			Mixtard 30
▲ Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70
			✓ PenMix 30
			✓ PenMix 40
			✓ PenMix 50

(PenMix 40 Inj human with neutral insulin 100 u per ml, 3 ml to be delisted 1 December 2022)

	Subsidy (Manufacturer's Price	e) Subs	Fully idised	Brand or Generic
	\$	Per	1	Manufacturer
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE				
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml,	,			
3 ml		5	✓	Humalog Mix 25
Inj lispro 50% with insulin lispro protamine 50% 100 u per ml,	ı			
3 ml	42.66	5	✓	Humalog Mix 50
Insulin - Long-acting Preparations				
NSULIN GLARGINE				
Inj 100 u per ml, 10 ml	63.00	1	✓	Lantus
Inj 100 u per ml, 3 ml		5	✓	Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓	Lantus SoloStar
Insulin - Rapid Acting Preparations				
NSULIN ASPART				
Inj 100 u per ml, 10 ml	30.03	1	✓	NovoRapid
Inj 100 u per ml, 3 ml		5		NovoRapid Penfill
Inj 100 u per ml, 3 ml syringe		5	✓	NovoRapid FlexPen
NSULIN GLULISINE				
Inj 100 u per ml, 10 ml	27.03	1	1	Apidra
Inj 100 u per ml, 3 ml	46.07	5	1	Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5		Apidra SoloStar
NSULIN LISPRO				
Inj 100 u per ml, 10 ml		10 ml OP	✓	Humalog
Inj 100 u per ml, 3 ml	59.52	5	√	Humalog
Alpha Glucosidase Inhibitors				
CARBOSE				
★ Tab 50 mg	8.95	90	1	Accarb
₭ Tab 100 mg	15.29	90	1	<u>Accarb</u>
Oral Hypoglycaemic Agents				
GLIBENCLAMIDE				
₹ Tab 5 mg	7.50	100	✓	Daonil
GLICLAZIDE			-	
₹ Tab 80 mg	15.18	500	1	Glizide
GLIPIZIDE	- · · · ·	*	-	
₭ Tab 5 mg	4.58	100	✓ I	Minidiab
METFORMIN HYDROCHLORIDE				
F Tab immediate-release 500 mg	14 74	1.000	√ 1	Metformin Mylan
★ Tab immediate-release 500 mg		500		Metformin Mylan
PIOGLITAZONE				
Tab 15 mg	6.80	90	1	Vexazone
· · · · · · · · · · · · · · · · · · ·		90	1	Vexazone Vexazone
€ Tab 30 mg				
≮ Tab 30 mg ≮ Tab 45 mg	12.25	90		Vexazone
· · · · · · · · · · · · · · · · · · ·	12.25	90		Vexazone

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VILDAGLIPTIN WITH METFORMIN HYDROCHLORIDE Tab 50 mg with 1,000 mg metformin hydrochloride Tab 50 mg with 850 mg metformin hydrochloride		60 60	-	Galvumet Galvumet

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

DULAGLUTIDE - Special Authority see SA2065 above - Retail pharmacy

Note: Not to be given in combination with a funded SGLT-2 inhibitor.

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	

continued...

- 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
- 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
- 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN – Special Authority see SA2068 on the previous page – Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 10 mg58	3.56	30	Jardiance
*	Tab 25 mg	3.56	30	Jardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE - Special Authority see SA2068 on the previous page - Retail pharmacy

Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

✓ CareSens Dual 1 OP

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes; or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes; or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 diagnostic test

1 OP ✓ CareSens N ✓ CareSens N POP 20.00 ✓ CareSens N Premier

Note: Only 1 meter available per PSO

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
, , , , , , , , , , , , , , , , , , , ,	Per 🗸	Manufacturer

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- 1) Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips	26.20	50 test OP	SensoCard
---------------------------	-------	------------	-----------------------------

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES - Maximum of 200 dev per prescription

*	29 g × 12.7 mm10	0.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm	2.26	100	✓ B-D Micro-Fine
	31 g × 6 mm		100	✓ Berpu
	31 g × 8 mm10		100	✓ B-D Micro-Fine
	32 g × 4 mm		100	✓ B-D Micro-Fine

		Subsidy		Fully	Brand or					
		(Manufacturer's Price)		Subsidised	Generic					
		\$	Per		Manufacturer					
INS	INSULIN SYRINGES, DISPOSABLE WITH ATTACHED NEEDLE – Maximum of 200 dev per prescription									
*	Syringe 0.3 ml with 29 g × 12.7 mm needle	13.56	100	1	B-D Ultra Fine					
		1.36	10							
		(1.99)			B-D Ultra Fine					
*	Syringe 0.3 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II					
		1.30	10							
		(1.99)			B-D Ultra Fine II					
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine					
		1.36	10							
		(1.99)			B-D Ultra Fine					
*	Syringe 0.5 ml with 31 g × 8 mm needle	13.56	100	✓	B-D Ultra Fine II					
		1.36	10							
		(1.99)			B-D Ultra Fine II					
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓	B-D Ultra Fine					
	, ,	1.36	10							
		(1.99)			B-D Ultra Fine					
*	Syringe 1 ml with 31 g x 8 mm needle	13.56	100	✓	B-D Ultra Fine II					
		1.36	10							
		(1.99)			B-D Ultra Fine II					

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription
- c) Maximum of 1 insulin pump per patient each four year period.

Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi		Generic	
	Per 🗸		Manufacturer	

continued...

- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Fither:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Fither:
 - 8.1 Applicant is a relevant specialist: or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — **(severe unexplained hypoglycaemia)** only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 Applicant is a relevant specialist; or

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (**Previous use before 1 September 2012**) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less

1 OP

✓ TruSteel

	Subsidy (Manufacturer's Price) \$	Subsid Per	Fully ised ✓	Brand or Generic Manufacturer
continued than 80 mmol/mol; and The patient's HbA1c has not deteriorated more than 5 mm The patient has not had an increase in severe unexplained Either: 4.1 Applicant is a relevant specialist; or 4.2 Applicant is a nurse practitioner working within their	d hypoglycaemic epis			ne; and
INSULIN PUMP CARTRIDGE — Special Authority see SA1985 or a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of cartridge sets will be funded per Cartridge 300 U. t:lock × 10	year.	narmacy	✓ T	andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) – Special A a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	Authority see SA1985		9 – Re	etail pharmacy
10 mm steel needle; 60 cm tubing × 10		1 OP 1 OP		liniMed Sure-T MMT-884A liniMed Sure-T
6 mm steel needle; 60 cm tubing × 10		1 OP	✓ M	MMT-886A liniMed Sure-T MMT-864A
6 mm steel needle; 80 cm tubing × 10		1 OP 1 OP		liniMed Sure-T MMT-866A liniMed Sure-T
8 mm steel needle; 80 cm tubing × 10		1 OP		MMT-874A liniMed Sure-T MMT-876A
6 mm steel needle; 29 G; manual insertion; 60 cm tubing × 10 with 10 needles; luer lock	130.00	1 OP	√ S	ure-T MMT-863
8 mm steel needle; 29 G; manual insertion; 60 cm tubing x 10 with 10 needles; luer lock		1 OP	_	ure-T MMT-873
INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles		cial Authorii 1 OP		SA1985 on page 19 –
8 mm steel cannula; straight insertion; 80 cm line × 10 with 10 needles	130.00	1 OP	✓ T	ruSteel
10 needles	130.00	1 OP	✓ T	ruSteel

8 mm steel cannula; straight insertion; 60 cm line × 10 with

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

\$ Per

Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year

c) Maximum of 13 infusion sets will be funded per year.			
13 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-382A
13 mm teflon needle, 45 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-368A
13 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-381A
13 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-383A
17 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-377A
17 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette MMT-378A
17 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	✓ MiniMed Silhouette
6 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	MMT-384A ✓ MiniMed Quick-Set
6 mm teflon needle, 45 cm blue tubing × 10	130.00	1 OP	MMT-398A ✓ MiniMed Mio
6 mm teflon needle, 45 cm pink tubing × 10	130.00	1 OP	MMT-941A ✓ MiniMed Mio
6 mm teflon needle, 60 cm blue tubing × 10	130.00	1 OP	MMT-921A ✓ MiniMed Mio
6 mm teflon needle, 60 cm pink tubing × 10	130.00	1 OP	MMT-943A ✓ MiniMed Mio
6 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	MMT-923A ✓ MiniMed Quick-Set
6 mm teflon needle, 80 cm blue tubing	130.00	1 OP	MMT-399A ✓ MiniMed Mio
6 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	MMT-945A ✓ MiniMed Mio
6 mm teflon needle, 80 cm pink tubing × 10	130.00	1 OP	MMT-965A ✓ MiniMed Mio
6 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	MMT-925A ✓ MiniMed Quick-Set
9 mm teflon needle, 110 cm tubing × 10	130.00	1 OP	MMT-387A ✓ MiniMed Quick-Set
9 mm teflon needle, 60 cm tubing × 10	130.00	1 OP	MMT-396A ✓ MiniMed Quick-Set
9 mm teflon needle, 80 cm clear tubing × 10	130.00	1 OP	MMT-397A ✓ MiniMed Mio
9 mm teflon needle, 80 cm tubing × 10	130.00	1 OP	MMT-975A ✓ MiniMed Quick-Set
			MMT-386A

	Subsidy (Manufacturer's Price) \$) Sub Per	Fully osidised	Brand or Generic Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	ISERTION WITH IN	SERTION	DEVICE	E) - Special Authority see
SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
13 mm teflon cannula; angle insertion; insertion device; 110 c	:m			
line x 10 with 10 needles		1 OP	✓ Aı	utoSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cn line x 10 with 10 needles		1 OP	✓ A	utoSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE IN	SERTION) - Spec	cial Author	ity see S/	A1985 on page 19 –
Retail pharmacy	, ,		•	
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
17 mm teflon cannula; angle insertion; 60 cm line × 10 with 10 needles; luer lock	120.00	1 OP	√ Si	ilhouette MMT-373
•		-		
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH see SA1985 on page 19 – Retail pharmacy	I INSERTION WITH	H INSEKI	ION DEV	/ICE) - Special Authority
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; insertion device;				
110 cm line × 10 with 10 needles	140.00	1 OP	✓ Aı	utoSoft 90
6 mm teflon cannula; straight insertion; insertion device; 60 cr				
line × 10 with 10 needles	140.00	1 OP	✓ Ai	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device;	140.00	4 OD		
110 cm line × 10 with 10 needles		1 OP	♥ Al	utoSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 cr		1 OP	✓ A	utoSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH				
Retail pharmacy	I INSERTION) - 3	ppeciai Au	Honly Se	e 3A 1905 on page 19 -
a) Maximum of 3 sets per prescription				
b) Only on a prescription				
c) Maximum of 13 infusion sets will be funded per year.				
6 mm teflon cannula; straight insertion; 60 cm tubing \times 10 with	h			
10 needles; luer lock		1 OP	✓ Q	uick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing x 10 with		4.00		
10 needles; luer lock		1 OP	₽ Q	uick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 or	n page 19 – Retail p	harmacy		
A) Maximum of 3 sets per prescription				
b) Only on a prescriptionc) Maximum of 13 packs of reservoir sets will be funded per	voor			
10 x luer lock conversion cartridges 1.8 ml for Paradigm pum		1 OP	✓ Δ1	DR Cartridge 1.8
Cartridge for 5 and 7 series pump; 1.8 ml × 10		1 OP		iniMed
1 17				1.8 Reservoir
				MMT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		iniMed
				3.0 Reservoir
				MMT-332A

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
· · · · ·	Dor 🗸	Manufacturor

Digestives Including Enzymes

Р	Δ	١	J	CI	R	F	Δ	Т	IC.	F	N	7	1	И	F

1 / WOTE/WIO ENZIME			
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	✓ Creon 10000
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase, 1,250 U protease))	94.40	100	✓ Panzytrat
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	✓ Creon 25000
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph			
Eur U)	34.93	20 g OP	✓ Creon Micro
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amy	lase, 1,250 U p	rotease)) to	be delisted 1 June 2023)
URSODEOXYCHOLIC ACID – Special Authority see SA1739 below		nacy 100	✓ <u>Ursosan</u>

⇒SA1739 Special Authority for Subsidy

Initial application — (Alaquille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has chronic severe drug induced cholestatic liver injury; and
 - 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
 - 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (Pregnancy) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN): and
- 2 Liver function has not improved with modifying the TPN composition.

Subsidy		ully	Brand or	_
(Manufacturer's Price)	Subsid	ised	Generic	
\$	Per	•	Manufacturer	

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Note: Ursodeoxycholic acid is not an appropriate therapy for patients requiring a liver transplant (bilirubin > 100 micromol/l; decompensated cirrhosis). These patients should be referred to an appropriate transplant centre. Treatment failure -- doubling of serum bilirubin levels, absence of a significant decrease in ALP or ALT and AST, development of varices, ascites or encephalopathy, marked worsening of pruritus or fatigue, histological progression by two stages, or to cirrhosis, need for transplantation.

Laxatives

Bulk-forming Ager	nts
--------------------------	-----

ISPAGHULA (PSYLLIUM) HUSK – Only on a prescription * Powder for oral soln	6.00	250 g OP	✓ Macro Organic
	12.20	500 g OP	Psyllium Husk ✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS * Dry	6.02	500 g OP	
The state of the s	(17.32)	300 g Oi	Normacol Plus
	2.41 (8.72)	200 g OP	Normacol Plus
Faecal Softeners			
DOCUSATE SODIUM - Only on a prescription			
* Tab 50 mg * Tab 120 mg		100 100	✓ <u>Coloxyl</u> ✓ Coloxyl
DOCUSATE SODIUM WITH SENNOSIDES			
Tab 50 mg with sennosides 8 mg	3.50	200	✓ Laxsol
POLOXAMER – Only on a prescription Not funded for use in the ear.			
* Oral drops 10%	3.98	30 ml OP	✓ Coloxyl
Opioid Receptor Antagonists - Peripheral			
METHYLNALTREXONE BROMIDE – Special Authority see SA1 Inj 12 mg per 0.6 ml vial		ail pharmacy 1	✓ Relistor

⇒SA1691 Special Authority for Subsidy

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

246.00

continued...

✓ Relistor

Subsidy	Fu	ılly	Brand or
(Manufacturer's Price)	Subsidis	ed	Generic
 \$	Per	✓	Manufacturer

continued...

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Osmotic Laxatives

GLYCEROL * Suppos 3.6 g - Only on a prescription	9.25	20	✓ PSM
LACTULOSE – Only on a prescription			
* Oral liq 10 g per 15 ml	3.33	500 ml	✓ Laevolac
MACROGOL 3350 WITH POTASSIUM CHLORIDE, SODIUM Powder for oral soln 13.125 g with potassium chloride 46		D SODIUM (CHLORIDE
sodium bicarbonate 178.5 mg and sodium chloride 3	350.7 mg 6.70	30	✓ Molaxole
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACET	ATE - Only on a pres	scription	
Enema 90 mg with sodium lauryl sulphoacetate 9 mg per	r ml,		
5 ml	29.98	50	✓ Micolette
Ottom death and the co			

Stimulant Laxatives

BISACODYL - Only on a prescription

* Tab 5 mg	5.80	200	✓ Pharmacy Health
* Suppos 10 mg		10	✓ Lax-Suppositories
SENNA - Only on a prescription			
* Tab, standardised	2.17	100	
	(8.21)		Senokot
	0.43	20	
	(2.06)		Senokot
SODIUM PICOSULFATE - Special Authority see \$A2053 belo	w – Retail pharma	icv	
Oral soln 7.5 mg per ml	7.40	30 ml OP	✓ Dulcolax SP Drop

⇒SA2053 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient is a child with problematic constipation despite an adequate trial of other oral pharmacotherapies including macrogol where practicable; and
- 2 The patient would otherwise require a high-volume bowel cleansing preparation or hospital admission.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Metabolic Disorder Agents

ALGLUCOSIDASE ALFA – Special Authority see SA1986	on the next page – Retail	pharmacy	
Inj 50 mg vial	1,142.60	1	✓ Myozyme

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT: and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ADCININE	 Special Authority see 	CA2042 bolow	Potail pharmacy
ABUILINE	- Special Aumoniv See	SAZU4Z DEIOW -	- Belali Dharmacv

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg	CBS	50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE - Special Authority see SA19	87 on the next page – Retail pharmacy	
_ , , , ,		

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsi	dised	Generic	
\$	Per	•	Manufacturer	

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5,10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - S	pecial Authority see SA2039 below – Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assav in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene, and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

LARONIDASE – Special Authority see SA1695 below – Retail pharmacy
Inj 100 U per ml, 5 ml vial......1,335.16

✓ Aldurazyme

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT): and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the next page - Retail pharmacy
Tab 100 mgCBS 100
Cap 100 mgCBS 100
✓ Country Life
✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

SAPROPTERIN DIHYDROCHLORIDE - Special Authority see SA1989 below - Retail pharmacy

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

SODIUM BENZOATE - Special Authority see SA1599 below - Retail pharmacy Soln 100 mg per mlCBS

DECOMING pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE - Special Authority see SA1990 on the next page - Retail pharmacy

Grans 483 mg per g......2,016.00 174 g OP ✓ **Pheburane**

100 ml

✓ Amzoate S29

Subsidy		Fully	Brand or	
(Manufacturer's Price)	,	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg		90	✓ Life Extension
Powder	000	300 g	✓ Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy		
Inj 200 unit vial1,072.00	1	✓ Elelyso

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease; or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size; and

Subsidy		Fully	Brand or
(Manufacturer's Pri	ice)	Subsidised	Generic
\$	Pe	r 🗸	Manufacturer

continued...

- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration			
BENZYDAMINE HYDROCHLORIDE Soln 0.15% - Higher subsidy of \$20.31 per 500 ml with			
Endorsement	9.00	500 ml	
	(20.31)		Difflam
Additional subsidy by endorsement for a patient who hap rescription is endorsed accordingly.	is oral mucositis a	as a result of tre	eatment for cancer, and the
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	
	(3.60)		Orabase
Powder	8.48	28 g OP	
	(10.95)		Stomahesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE			
* Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 a OP	
•	(6.00)	J	Bonjela
TRIAMCINOLONE ACETONIDE	, ,		·
Paste 0.1%	5.33	5 a OP	✓ Kenalog in Orabase
1 4545 51176		0 y 0.	- Nonarog III Grazaco
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 a OP	✓ Decozol
NYSTATIN	•	3	
Oral liq 100,000 u per ml	1 76	24 ml OP	✓ Nilstat
Oral ing 100,000 u per IIII	1.70	24 IIII OF	· Mistat

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
Vitamins				
/itamin B				
YDROXOCOBALAMIN				
Inj 1 mg per ml, 1 ml ampoule - Up to 6 inj available on a PS	O 1.89	3		Vita-B12
	2.46		•	Hydroxocobalamin Panpharma
	2.84	_		Neo-B12
	3.15	5	•	Hydroxocobalamin Mercury Pharma
iita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 Novembe leo-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 Novembe lydroxocobalamin Mercury Pharma Inj 1 mg per ml, 1 ml ampoul ('RIDOXINE HYDROCHLORIDE	er 2022)	Novem.	ber 2022)	·
a) No more than 100 mg per dose				
b) Only on a prescription				
Tab 25 mg - No patient co-payment payable		90	✓	Vitamin B6 25
Tab 50 mg	23.45	500	•	Pyridoxine
				multichem
HIAMINE HYDROCHLORIDE – Only on a prescription				
Tab 50 mg	7.09	100	•	Max Health
TAMIN B COMPLEX			_	
Tab, strong, BPC	7.15	500		Bplex
/itamin C				
SCORBIC ACID				
a) No more than 100 mg per dose				
b) Only on a prescription			_	
Tab 100 mg	9.90	500	✓	Cvite
/itamin D				
LFACALCIDOL				
Cap 0.25 mcg	26.32	100	✓	One-Alpha
Cap 1 mcg		100		One-Alpha
Oral drops 2 mcg per ml	60.68 2	20 ml C)P 🗸	One-Alpha
ALCITRIOL			_	
Cap 0.25 mcg		100		Calcitriol-AFT
Cap 0.5 mcg	13.68	100	•	Calcitriol-AFT
OLECALCIFEROL (50 000) M ; (40	0.5-			W. Do
Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescription		12		Vit.D3
Oral liq 188 mcg per ml (7,500 iu per ml)	9.00 4	l.8 ml ()P •	Puria
Multivitamin Preparations				
LILTIVITAMINI DENIAL Consist Authority and CA4E4C on the m	ovt page - Detail p	harma	CV	
ULTIVITAMIN RENAL – Special Authority see SA1546 on the n	ext page – netali p	ınanına	O y	
Cap		30 30	•	Clinicians Renal Vit

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Pri \$	ce) Subs	Fully sidised	Brand or Generic Manufacturer
➤ SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals va the following criteria: Either:				
 The patient has chronic kidney disease and is receiving The patient has chronic kidney disease grade 5, defined ml/min/1.73 m² body surface area (BSA). 				
MULTIVITAMINS - Special Authority see SA1036 below - Ret * Powder		200 g OP	✓ Pa	aediatric Seravit
Initial application from any relevant practitioner. Approvals vainborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid withou approval for multivitamins.				·
VITAMINS * Tab (BPC cap strength)* * Cap (fat soluble vitamins A, D, E, K) – Special Authority se		1,000	✓ M	vite
SA1720 below – Retail pharmacy		60	✓ V	itabdeck
Initial application from any relevant practitioner. Approvals va the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; 2 Patient is an infant or child with liver disease or short gu 3 Patient has severe malabsorption syndrome.	or	niewai unies.	s nounec	To applications meeting
Minerals				
Calcium				
CALCIUM CARBONATE * Tab 1.25 g (500 mg elemental) * Tab eff 1.25 g (500 mg elemental) – Subsidy by endorsem		250 100	_	alci-Tab 500 alcium 500 mg Hexal ^{©29}
Subsidy by endorsement – Only when prescribed for p considered unsuitable.	aediatric patients (<	5 years) whe	ere calci	
CALCIUM GLUCONATE * Inj 10%, 10 ml ampoule	32.00	10	✓ M	ax Health -
	64.00	20	✓ M	ax Health \$29
Fluoride				
Fluoride SODIUM FLUORIDE * Tab 1.1 mg (0.5 mg elemental)		100	√ P:	

POTASSIUM IODATE

✓ NeuroTabs

90

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully sidised	Brand or Generic Manufacturer	
Iron					
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.04	100	√ <u>F</u>	erro-tab	
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg Ferro-F-Tabs to be Principal Supply on 1 August 2022	5.98	100	√ F	erro-F-Tabs	
FERROUS SULFATE * Tab long-acting 325 mg (105 mg elemental) * Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml		errograd erodan	
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority se Inj 50 mg per ml, 10 ml vial		letail phar 1	,	erinject	

⇒SA1840 Special Authority for Subsidy

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective; or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

*	Inj 50 mg per ml, 2 ml ampoule	34.50	5	✓ Ferrosig
---	--------------------------------	-------	---	------------

	Subsidy (Manufacturer's Price \$) Su Per	Fully bsidised	Brand or Generic Manufacturer	
Magnesium					
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml		hillips Milk of Magnesia §29	
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ <u>M</u>	artindale	
Zinc					
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ Z	incaps	

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus: and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus: and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*: and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Epoetin alfa is indicated in the treatment of anaemia associated with chronic renal failure (CRF) where no cause for anaemia other than CRF is detected and there is adequate monitoring of iron stores and iron replacement therapy.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
EPOETIN ALFA - Special Authority see SA1775 on the previous	page – Retail pharm	асу		
Wastage claimable		•		
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	✓	Binocrit
Inj 2,000 iu in 1 ml, syringe	100.00	6	✓	Binocrit
Inj 3,000 iu in 0.3 ml, syringe	150.00	6	✓	Binocrit
Inj 4,000 iu in 0.4 ml, syringe	96.50	6	✓	Binocrit
Inj 5,000 iu in 0.5 ml, syringe	125.00	6	✓	Binocrit
Inj 6,000 iu in 0.6 ml, syringe	145.00	6	✓	Binocrit
Inj 8,000 iu in 0.8 ml, syringe	175.00	6	1	Binocrit
Inj 10,000 iu in 1 ml, syringe	197.50	6	1	Binocrit
Inj 40,000 iu in 1 ml, syringe	250.00	1	✓	Binocrit
Megaloblastic				

FC	LIC ACID			
	Tab 0.8 mg	26.60	1,000	✓ Folic Acid multichem
*	Tab 5 mg	5.82	100	✓ Folic Acid Mylan
	Oral liq 50 mcg per ml	27.82	25 ml OP	✓ Biomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial		1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	✓ Alprolix
ELTROMBOPAG – Special Authority see SA1743 belo	w – Retail pharmacy		·

Е

Wastage olaimable			
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	✓ Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding: or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist.

continued...

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic
 \$	Per	•	Manufacturer

continued...

Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy.

Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab);
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- Fither:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

FMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

✓ Hemlibra	1	3,570.00	Inj 30 mg in 1 ml vial
✓ Hemlibra	1	7,138.00	Inj 60 mg in 0.4 ml vial
✓ Hemlibra	1	12,492.00	Inj 105 mg in 0.7 ml vial
✓ Hemlibra	1	17.846.00	Ini 150 mg in 1 ml vial

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more: and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or
 - 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
- 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe		1	✓ NovoSeven RT
Inj 5 mg syringe		1	✓ NovoSeven RT
Inj 8 mg syringe		1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	5.00 1	FEIBA NF
Inj 1,000 U2,630	0.00 1	✓ FEIBA NF
Inj 2,500 U	5.00 1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria.

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	✓ Xyntha
Inj 1,000 iu prefilled syringe		1	✓ Xyntha
Inj 2,000 iu prefilled syringe	2,300.00	1	Xyntha
Inj 3,000 iu prefilled syringe	3,450.00	1	Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	0 1	✓ RIXUBIS
Inj 1,000 iu vial870.0	0 1	✓ RIXUBIS
Inj 2,000 iu vial	0 1	✓ RIXUBIS
Inj 3,000 iu vial2,610.0	0 1	✓ RIXUBIS

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) -	- [Xpharm]		
For patients with haemophilia. Preferred Brand of short hal			
managed by the Haemophilia Treaters Group in conjunction			
Inj 250 iu vial		1	✓ Advate
Inj 500 iu vial		1	✓ Advate
Inj 1,000 iu vial		1	✓ Advate
Inj 1,500 iu vial	·	1	Advate
Inj 2,000 iu vial	,	1	Advate
Inj 3,000 iu vial	•	1	✓ Advate
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE For patients with haemophilia. Rare Clinical Circumstances treatment is managed by the Haemophilia Treaters Group in subject to criteria.	Brand of short half-life		
Inj 250 iu vial	237.50	1	Kogenate FS
Inj 500 iu vial	475.00	1	✓ Kogenate FS
Inj 1,000 iu vial	950.00	1	Kogenate FS
Inj 2,000 iu vial		1	✓ Kogenate FS
Inj 3,000 iu vial		1	✓ Kogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII For patients with haemophilia A receiving prophylaxis treatn Treaters Group in conjunction with the National Haemophilia Inj 250 iu vial	nent. Access to funder a Management group. 300.00 600.00 1,200.00	1 1 1 1	Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate Adynovate
· · · · · · · · · · · · · · · · · · ·	2,400.00	'	Adynovate
SODIUM TETRADECYL SULPHATE * Inj 3% 2 ml	28.50 (73.00)	5	Fibro-vein
TRANEXAMIC ACID	(/		
Tab 500 mg	9.45	60	✓ Mercury Pharma
- 1 ab 500 mg		-	- moroury i marma
Vitamin K			
PHYTOMENADIONE			
Inj 2 mg per 0.2 ml - Up to 5 inj available on a PSO	8.00	5	✓ Konakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓ Konakion MM
Antithrombotic Agents			
Antiplatelet Agents			
ASPIRIN * Tab 100 mg	10.80	990	✓ Ethics Aspirin EC
· ·		550	- Lanco Aspirii Lo
CLOPIDOGREL * Tab 75 mg	4.60	84	✓ Clopidogrel Multichem
DIPYRIDAMOLE	10.00	60	√ Dutomor: CD
* Tab long-acting 150 mg		60	✓ Pytazen SR
TICAGRELOR – Special Authority see SA1955 on the next pag * Tab 90 mg		56	✓ Brilinta

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	

⇒SA1955 Special Authority for Subsidy

Initial application — (acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Either:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Either:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment: and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	(Manufacturer's Price)	Subsi Per	dised	Generic Manufacturer	
Hamanin and Antononiat Dromonations					

Heparin and Antagonist Preparations

•			
ENOXAPARIN SODIUM - Special Authority see SA1646	below – Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	✓ Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe		10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe		10	 Clexane Forte
Inj 150 mg in 1 ml syringe		10	 Clexane Forte

⇒SA1646 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy: or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

HEPARIN SODIUM

Inj 1,000 iu per ml, 5 ml ampoule	72.84	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml	32.66	5	DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	289.05	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml	22.42	5	✓ Hospira
	42.40		✓ Heparin DBL S29
	482.20	50	✓ Heparin DBL S29
HEPARINISED SALINE			
Inj 10 iu per ml, 5 ml	65.48	50	✓ Pfizer

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
Oral Anticoagulants				
DABIGATRAN				
Cap 75 mg - No more than 2 cap per day	76.36	60	1	Pradaxa
Cap 110 mg	76.36	60	1	Pradaxa
Cap 150 mg	76.36	60	✓	Pradaxa
RIVAROXABAN				
Tab 10 mg - No more than 1 tab per day	83.10	30	1	Xarelto
Tab 15 mg - Up to 14 tab available on a PSO	77.56	28	1	Xarelto
Tab 20 mg	77.56	28	1	Xarelto
WARFARIN SODIUM				
Note: Marevan and Coumadin are not interchangeable.				
* Tab 1 mg	3.46	50	1	Coumadin
	6.46	100		Marevan
* Tab 2 mg		50	_	Coumadin
* Tab 3 mg		100		Marevan
* Tab 5 mg		50		Coumadin
	11.48	100	•	Marevan
Blood Colony-stimulating Factors				
FILGRASTIM - Special Authority see SA1259 below - Retail p	harmacy			
Inj 300 mcg per 0.5 ml prefilled syringe	96.22	10	1	Nivestim
Inj 480 mcg per 0.5 ml prefilled syringe	148.58	10	✓	Nivestim

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC $< 0.5 \times 10^9$ /L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 below - Retail pharmacy

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

	Subsidy		Fully	Brand or
	(Manufacturer's Pric		Subsidised	Generic
	\$	Per		Manufacturer
Fluids and Electrolytes				
Intravenous Administration				
GLUCOSE [DEXTROSE]				
* Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO		5 1		Biomed Biomed
* Inj 50%, 90 ml bottle - Up to 5 inj available on a PSO	15.00	ı	•	Biomed
* Inj 75 mg per ml, 10 ml	65.00	50	/	Juno
SODIUM BICARBONATE		00		ouno .
Inj 8.4%, 50 ml	21.40	1	1	Biomed
a) Up to 5 inj available on a PSO				
b) Not in combination	04.05			Diamed
Inj 8.4%, 100 ml	21.95	1	•	Biomed
b) Not in combination				
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Not funded for nebuliser	use except when the	used in o	conjunctio	n with an antibiotic intended
for nebuliser use.	1.00	F00 l		Daviday
Inj 0.9%, bag – Up to 2000 ml available on a PSO	1.26	500 ml		Baxter Baxter
Only if prescribed on a prescription for renal dialysis, ma				
for emergency use. (500 ml and 1,000 ml packs)				
Inj 23.4% (4 mmol/ml), 20 ml ampoule For Sodium chloride oral liquid formulation refer Standare		5	•	Biomed
Inj 0.9%, 5 ml ampoule – Up to 5 inj available on a PSO		20	1	Fresenius Kabi
Inj 0.9%, 10 ml ampoule – Up to 5 inj available on a PSO	5.40	50		Fresenius Kabi
Inj 0.9%, 20 ml ampoule	5.00	20	1	Fresenius Kabi
TOTAL PARENTERAL NUTRITION (TPN)			_	
Infusion	CBS	1 OP	•	TPN
WATER			. tata atau	Pakadia dha Dhamaa ay dad
 On a prescription or Practitioner's Supply Order only wh Schedule requiring a solvent or diluent; or 	nen on the same to	rm as ar	1 injection	listed in the Pharmaceutical
2) On a bulk supply order; or				
3) When used in the extemporaneous compounding of eye				
4) When used for the dilution of sodium chloride soln 7% f	or cystic fibrosis pa	atients o	nly.	
Inj 10 ml ampoule - Up to 5 inj available on a PSO	7 19	50	1	Pfizer
Inj 20 ml ampoule – Up to 5 inj available on a PSO		20		Fresenius Kabi
			1	Multichem
Oral Administration				
Oral Auministration				
CALCIUM POLYSTYRENE SULPHONATE	100.05	000 - 0	n .	Oalaium Dagerier
Powder	169.85	300 g O	۲ 🗸	Calcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO	9.53	50	/	Electral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE]		50	•	Livellai
Soln with electrolytes (2 × 500 ml)	6.55 1	,000 ml	OP 🗸	Pedialyte -
, ,				Bubblegum

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	Brand or Generic Manufacturer
PHOSPHORUS				
Tab eff 500 mg (16 mmol)	82.50	100	✓ F	Phosphate Phebra
POTASSIUM CHLORIDE				
* Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26	60		
	(17.10)		(Chlorvescent
* Tab long-acting 600 mg (8 mmol)	8.90	200	✓ 9	Span-K
SODIUM BICARBONATE				
Cap 840 mg	8.52	100	√ 9	Sodibic
			√ 9	Sodibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84 65	154 a C)P 🗸 F	Resonium-A

S29 S29

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	1	Manufacturer	

Alpha-Adrenoceptor Blockers

Alpha Adrenoceptor Blockers

DOXAZOSIN		
* Tab 2 mg	500	✓ Apo-Doxazosin✓ Doxazosin Clinect
* Tab 4 mg	500	✓ Apo-Doxazosin✓ Doxazosin Clinect
(Apo-Doxazosin Tab 2 mg to be delisted 1 September 2022)		
(Apo-Doxazosin Tab 4 mg to be delisted 1 September 2022)		
PHENOXYBENZAMINE HYDROCHLORIDE		
* Cap 10 mg65.00	30	✓ BNM S29
216.67	100	✓ Dibenzyline S29
PRAZOSIN		
* Tab 1 mg	100	 ✓ Arrotex-Prazosin S29 S29
* Tab 2 mg7.00	100	✓ Arrotex-Prazosin S29 S29
* Tab 5 mg 11.70	100	✓ Arrotex-Prazosin

Agents Affecting the Renin-Angiotensin System

ACE Inhibitors

CAP	

*	Oral liq 5 mg per ml94.99	95 ml OP	Capoten
	Oral liquid restricted to children under 12 years of age.		

CILAZAPRIL - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking cilazapril prior to 1 May 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of cilazapril.

* T	ab 0.5 mg2.09	90	✓ Zapril
	ab 2.5 mg4.80	90	✓ Zapril
T	ab 5 mg8.35	90	✓ Zapril
ENAL	APRIL MALEATE		
* T	ab 5 mg1.82	100	✓ Acetec
	ab 10 mg2.02	100	✓ Acetec
* T	ab 20 mg2.42	100	✓ Acetec
LISIN	OPRIL		
* T	ab 5 mg11.07	90	 Ethics Lisinopril
	ab 10 mg11.67	90	✓ Ethics Lisinopril
* T	ab 20 mg14.69	90	 Ethics Lisinopril
PERI	NDOPRIL		
Т	ab 2 mg1.58	30	✓ Coversyl
T	ab 4 mg2.95	30	✓ Coversyl

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
QUINAPRIL				
Tab 5 mg	5.97	90	✓ .	Arrow-Quinapril 5
Tab 10 mg	5.18	90	✓ .	Arrow-Quinapril 10
Tab 20 mg	7.95	90	✓	Arrow-Quinapril 20

ACE Inhibitors with Diuretics

QUINAPRIL WITH HYDROCHLOROTHIAZIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking quinapril with hydrochlorothiazide prior to 1 May 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of quinapril with hydrochlorothiazide.

✓ Accuretic 10	30	g4.10	Tab 10 mg with hydrochlorothiazide 12.5 mg
✓ Accuretic 20	30	g5.25	Tab 20 mg with hydrochlorothiazide 12.5 mg

Angiotensin II Antagonists

* Tab 4 mg	2.00	90	Candestar
* Tab 8 mg		90	✓ Candestar
* Tab 16 mg		90	✓ Candestar
* Tab 32 mg	5.26	90	✓ Candestar
LOSARTAN POTASSIUM			
Tab 12.5 mg	1.56	84	✓ Losartan Actavis
Tab 25 mg	1.84	84	✓ Losartan Actavis
Tab 50 mg	2.25	84	✓ Losartan Actavis
Tab 100 mg	3.50	84	✓ Losartan Actavis

Angiotensin II Antagonists with Diuretics

Tab 50 mg with hydrochlorothiazide 12.5 mg	15.25	30	✓ Arrow-Losartan &
			Hydrochlorothiazide

Angiotensin II Antagonists with Neprilysin Inhibitors

SACUBITRIL WITH VALSARTAN - Special Authority see SA1905 below - Retail pharmacy

Note: Due to the angiotensin II receptor blocking activity of sacubitril with valsartan it should not be co-administered with an ACE inhibitor or another ARB.

Tab 24.3 mg with valsartan 25.7 mg	190.00	56	✓ Entresto 24/26
Tab 48.6 mg with valsartan 51.4 mg	190.00	56	✓ Entresto 49/51
Tab 97.2 mg with valsartan 102.8 mg	190.00	56	✓ Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III: or
 - 2.3 Patient is in NYHA/WHO functional class IV; and

continued...

Subsidy (Manufacturer's Pri	ce)	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

- 3 Either:
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or
 - 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics			
For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesth	netics, Local, p	age 118	
AMIODARONE HYDROCHLORIDE			
▲ Tab 100 mg		30	✓ Aratac
▲ Tab 200 mg	4.49	30	✓ Aratac
Inj 50 mg per ml, 3 ml ampoule – Up to 10 inj available on a	45.00	10	/ May Haalib
PSO	15.22	10	✓ Max Health
ATROPINE SULPHATE			
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	15.00	10	✓ Martindale
DIGOXIN	13.03	10	<u> inartiridate</u>
* Tab 62.5 mcg - Up to 30 tab available on a PSO	7 00	240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO		240	✓ Lanoxin
* Oral lig 50 mcg per ml		60 ml	✓ Lanoxin
			 Lanoxin Paediatric
			Elixir S29
			✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE			
▲ Cap 100 mg	23.87	100	Rythmodan
FLECAINIDE ACETATE			
▲ Tab 50 mg		60	✓ Flecainide BNM
▲ Cap long-acting 100 mg	39.51	90	✓ Flecainide
			Controlled
A Can long acting 200 mg	61.06	90	Release Teva ✓ Flecainide
▲ Cap long-acting 200 mg	01.00	90	Controlled
			Release Teva
Inj 10 mg per ml, 15 ml ampoule	100.00	5	✓ Tambocor
MEXILETINE HYDROCHLORIDE			
▲ Cap 150 mg	162.00	100	✓ Teva S29
▲ Cap 250 mg		100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE			
▲ Tab 150 mg	40.90	50	✓ Rytmonorm
-			·
Antihypotensives			
MIDODRINE - Special Authority see SA1474 on the next page - F	Retail pharmac	/	

IIDODRINE - Special Authority see SA1474 on the ne	xt page – Retail pharmacy		
Tab 2.5 mg	53.00	00	Gutron
Tab 5 mg	79.00 1	00	Gutron

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Note: Treatment should be started with small doses and titrated upwards as necessary. Hypertension should be avoided, and the usual target is a standing systolic blood pressure of 90 mm Hg.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOI OI

ATENOLOL			
* Tab 50 mg	9.33	500	✓ Mylan Atenolol
* Tab 100 mg	14.20	500	✓ Mylan Atenolol
* Oral liq 25 mg per 5 ml	21.25	300 ml OP	✓ Atenolol AFT
			S29 S29
	38.20		✓ Essential
	00.20		Generics \$29
	49.85		✓ Atenolol AFT
Restricted to children under 12 years of age.	49.00		▼ Ateriolol AFT
, ,			
BISOPROLOL FUMARATE			
* Tab 2.5 mg		90	✓ Bisoprolol Mylan
* Tab 5 mg		90	✓ Bisoprolol Mylan
* Tab 10 mg	3.62	90	✓ Bisoprolol Mylan
CARVEDILOL			
* Tab 6.25 mg	2.24	60	Carvedilol Sandoz
* Tab 12.5 mg	2.30	60	Carvedilol Sandoz
* Tab 25 mg	2.95	60	Carvedilol Sandoz
LABETALOL			
* Tab 100 mg	14.50	100	✓ Trandate
* Tab 200 mg		100	✓ Trandate
* Inj 5 mg per ml, 20 ml ampoule		5	
, 01	(88.60)		Trandate
* inj 5 mg per ml, 20 ml vial	42.29 [′]	1	
, 01	(48.20)		Alvogen S29
METOPROLOL SUCCINATE	(10.00)		9
* Tab long-acting 23.75 mg	1 //5	30	✓ Betaloc CR
* Tab long-acting 25.75 mg		30	✓ Betaloc CR
* Tab long-acting 47.5 mg		30	✓ Betaloc CR
* Tab long-acting 95 mg		30	✓ Betaloc CR
	4.21	30	• Detailor Cit
METOPROLOL TARTRATE	F 00	100	/ IDOA Matarasalal
Tab 50 mg		100	✓ IPCA-Metoprolol
Tab 100 mg		60	✓ IPCA-Metoprolol
* Tab long-acting 200 mg		28	✓ Slow-Lopresor
* Inj 1 mg per ml, 5 ml vial	26.50	5	Metoprolol IV Mylan
NADOLOL			
Tab 40 mg	19.19	100	✓ Nadolol BNM S29
Tab 80 mg	30.39	100	✓ Nadolol BNM S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPRANOLOL				
Tab 10 mg	7.04	100	✓ D	rofate
Tab 40 mg	8.75	100	✓ II	PCA-Propranolol
* Cap long-acting 160 mg	18.17	100	√ 0	ardinol LA
★ Oral lig 4 mg per ml - Special Authority see SA1327 be	elow –			
Retail pharmacy		500 m	l √ R	loxane- Propranolol S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

*	Tab 80 mg32.58	500	Mylan
	Tab 160 mg10.98		✓ Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑN	ILODIPINE		
*	Tab 2.5 mg1.08	90	✓ Vasorex
*	Tab 5 mg0.96	90	✓ Vasorex
*	Tab 10 mg1.19	90	✓ Vasorex
FE	LODIPINE		
*	Tab long-acting 2.5 mg1.45	30	✓ Plendil ER
*	Tab long-acting 5 mg4.07	90	✓ Felo 5 ER
*	Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
NIF	EDIPINE		
*	Tab long-acting 10 mg	56	✓ Tensipine MR10 S29
*	Tab long-acting 20 mg9.12	50	✓ Mylan (12 hr release) S29
	17.72	100	✓ Nyefax Retard
*	Tab long-acting 30 mg4.78	14	✓ Mylan Italy (24 hr
			release) \$29
	34.10	100	✓ Mylan (24 hr
			release) \$29
*	Tab long-acting 60 mg52.81	100	✓ Mylan (24 hr
			release) \$29
			- /

	Subsidy (Manufacturer's Price) \$	Per	Subsidised G	rand or eneric anufacturer
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	✓ Acco	ord S29
₭ Cap long-acting 120 mg		500	✓ Apo-	Diltiazem CD
₭ Cap long-acting 180 mg		30		izem CD
★ Cap long-acting 240 mg		30	✓ Card	izem CD
PERHEXILINE MALEATE				
★ Tab 100 mg	62 90	100	✓ Pexs	in
/ERAPAMIL HYDROCHLORIDE	02.00	100	- I CAC	"9
	7.01	100	√ loon	lin.
★ Tab 40 mg Tab 80 mg		100	✓ Isop	
<u> </u>			•	
* Tab long-acting 120 mg	36.02	100		tin Retard \$29
★ Tab long-acting 240 mg	15 10	30	✓ Isop	
		30	• Isop	un ən
Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO		5	./ loon	u
P3U	25.00	Э	✓ Isop	un
Centrally-Acting Agents CLONIDINE				
▶ Patch 2.5 mg, 100 mcg per day – Only on a prescription	10.34	4	✓ Myla	n
F Patch 5 mg, 200 mcg per day − Only on a prescription		4	✓ Myla	_
★ Patch 7.5 mg, 300 mcg per day – Only on a prescription		4	✓ Myla	_
CLONIDINE HYDROCHLORIDE			_	
★ Tab 25 mcg	8 75	112	✓ Clon	idine BNM
- 145 25 116g	29.32			idine Teva
★ Tab 150 mcg		100	✓ Cata	
lt Inj 150 mcg per ml, 1 ml ampoule		10	✓ Med	
Clonidine BNM Tab 25 mcg to be delisted 1 November 2022)				
METHYLDOPA				
∦ Tab 250 mg	15 10	100	✓ Math	yldopa Mylan
r 140 200 mg	52.85	500		yldopa Mylan
	02.00	000		9 S29
			- JE	3 023
Diuretics				
Loop Diuretics				
BUMETANIDE				
₭ Tab 1 mg	4 91	30	✓ Rurii	nex S29 S29
r 100 r mg	16.36	100	✓ Buri	
₭ Inj 500 mcg per ml, 4 ml vial		, 00	· Duiii	

_		Subsidy		Fully Brai	nd or
		(Manufacturer's F	Price) S Per	ubsidised Ger ✓ Mar	eric jufacturer
=		φ	rei	V IVIAI	lulacturei
FU	ROSEMIDE [FRUSEMIDE]	0.00	1 000	∠ IDCA I	
*	Tab 40 mg — Up to 30 tab available on a PSO Tab 500 mg		1,000 50	✓ IPCA-I	Frusemide Corte
~	Tab 300 mg	89.48	30	✓ Furose	
		00.10			opharm S29
		169.96	100	✓ Furose	emid-
				Ratio	opharm S29
*	Oral liq 10 mg per ml	11.20	30 ml OP	✓ Lasix	
	Inj 10 mg per ml, 25 ml ampoule		6	Lasix	
*	Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj available on a P	SO 1.15	5	✓ Furose	emide-Baxter
P	otassium Sparing Diuretics				
ΑM	ILORIDE HYDROCHLORIDE				
	Oral liq 1 mg per ml	32.10	25 ml OP	Biome	d
ΕP	LERENONE – Special Authority see SA1728 below – Retail pl	harmacy			
	Tab 25 mg		30	✓ Inspra	
_	Tab 50 mg	25.00	30	✓ Inspra	
>	SA1728 Special Authority for Subsidy				
	ial application from any relevant practitioner. Approvals valid following criteria:	I without further	renewal unl	ess notified for a	applications meeting
Bot	•				
Do	 Patient has heart failure with ejection fraction less than 40° 	%· and			
	2 Either:	70, and			
	2.1 Patient is intolerant to optimal dosing of spironolact	one; or			
	2.2 Patient has experienced a clinically significant adve	erse effect while	on optimal of	dosing of spiron	olactone.
ME	TOLAZONE				
	Tab 5 mg	CBS	1	✓ Metola	zone S29
	-		50	✓ Zaroxo	olyn S29
SP	RONOLACTONE				

100

100

25 ml OP

28

50

✓ Spiractin

✓ Spiractin

✓ Biomed

✓ Frumil

✓ Moduretic

▲Three months supply may be dispensed at one time if endorsed	"cartified examption" by the prescriber or pharmacist
Trifee months supply may be dispensed at one time if endorsed	i certified exemption by the prescriber of pharmacist.

* Tab 5 mg with hydrochlorothiazide 50 mg......5.00

AMILORIDE HYDROCHLORIDE WITH HYDROCHLOROTHIAZIDE

Spiractin to be Principal Supply on 1 September 2022

Potassium Sparing Combination Diuretics

AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully sidised	Brand or Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE] * Tab 2.5 mg - Up to 150 tab available on a PSO	20.00	500	✓ <u>A</u>	rrow- Bendrofluazide
May be supplied on a PSO for reasons other than em * Tab 5 mg	• .	500	✓ <u>A</u>	rrow- Bendrofluazide
CHLOROTHIAZIDE Oral liq 50 mg per ml	27.82	25 ml OP	✓ B	iomed
CHLORTALIDONE [CHLORTHALIDONE] Tab 25 mg	3.90 6.50	30 50		roton §29 ygroton
NDAPAMIDE * Tab 2.5 mg	10.45 11.61	90 100	_	apa-Tabs ylan Indapamide §29
Lipid-Modifying Agents				
Fibrates				
BEZAFIBRATE * Tab 200 mg * Tab long-acting 400 mg		90 30	_	ezalip ezalip Retard
Other Lipid-Modifying Agents				
ACIPIMOX * Cap 250 mg	21.56	30	-	Ibetam Ibetam S29 S29
Resins				
COLESTIPOL HYDROCHLORIDE Grans for oral liq 5 g	32.89	30	✓ C	olestid
HMG CoA Reductase Inhibitors (Statins)				
ATORVASTATIN * Tab 10 mg * Tab 20 mg * Tab 40 mg * Tab 80 mg	9.24 14.92	500 500 500 500	✓ <u>L</u>	orstat orstat orstat orstat
* Tab 20 mg	2.11	28	√ <u>P</u>	ravastatin Mylan

	Subsidy (Manufacturer's Price) \$		Subsidised	Brand or Generic Manufacturer
ROSUVASTATIN - Special Authority see SA2093 below - Retail	pharmacy			
* Tab 5 mg	1.70	30	✓	Rosuvastatin Viatris
* Tab 10 mg	2.42	30	✓	Rosuvastatin Viatris
* Tab 20 mg	3.92	30	✓	Rosuvastatin Viatris
* Tab 40 mg	5.28	30	•	Rosuvastatin Viatris

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

. . .

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity: or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atoryastatin and/or simyastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	//VASTATIN			
*	Tab 10 mg	1.23	90	 Simvastatin Mylan
*	Tab 20 mg	2.03	90	✓ Simvastatin Mylan
*	Tab 40 mg	3.58	90	✓ Simvastatin Mylan
	Tab 80 mg		90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

ΕZI	ETIMIBE - Special Authority see SA1045 on the next page - Retail pharmacy		
*	Tab 10 mg1.95	30	✓ Ezetimibe Sandoz

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	1	Manufacturer

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than 10 × normal) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to < 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy. If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

EZETIMIBE WITH SIMVASTATIN - Special Authority see SA1046 below - Retail pharmacy

Tab 10 mg with simvastatin 10 mg	5.15	30	Zimybe
Tab 10 mg with simvastatin 20 mg	6.15	30	Zimybe
Tab 10 mg with simvastatin 40 mg	7.15	30	✓ Zimybe
Tab 10 mg with simvastatin 80 mg	8.15	30	✓ Zimybe

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Notes: A patient who has failed to reduce their LDL cholesterol to less than or equal to 2.0 mmol/litre with the use of a less potent statin should use a more potent statin prior to consideration being given to the use of non-statin therapies.

Other treatment options including fibrates, resins and nicotinic acid should be considered after failure of statin therapy.

If a patient's LDL cholesterol cannot be calculated because the triglyceride level is too high then a repeat test should be performed and if the LDL cholesterol again cannot be calculated then it can be considered that the LDL cholesterol is greater than 2.0 mmol/litre.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose – Up to 250 dose available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Sprav
	Patch 25 mg, 5 mg per day Patch 50 mg, 10 mg per day		30 30	✓ Nitroderm TTS ✓ Nitroderm TTS

	C	ARI	DIOVAS	CULAR SYSTEM
	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
ISOSORBIDE MONONITRATE				
* Tab 20 mg	19.55	100		Ismo 20
* Tab long-acting 40 mg		30		Ismo 40 Retard
* Tab long-acting 60 mg	9.25	90	•	<u>Duride</u>
Sympathomimetics				
ADRENALINE				
Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO		5		Aspen Adrenaline
lai 1 in 10 000 10 ml annaula . Lla ta 5 ini aunilable en a Bú	10.76	_		DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule – Up to 5 inj available on a PS	49.00	5 10		Hospira Aspen Adrenaline
	49.00	10		Aspen Aurenanne
Vasodilators				
HYDRALAZINE HYDROCHLORIDE				
* Tab 25 mg - Special Authority see SA1321 below - Retail				
pharmacy	CBS	1	✓	Hydralazine
		56	✓	Onelink \$29
		84	✓	AMDIPHARM \$29
		100	✓	Onelink S29
* Inj 20 mg ampoule	25.90	5	✓	Apresoline
SA1321 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Either:	without further rene	wal u	nless notif	fied for applications meeting
 For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nitrainhibitors and/or angiotensin receptor blockers. 	ate, in patients who a	are in	tolerant or	have not responded to AC
MINOXIDIL			,	
▲ Tab 10 mg	70.00	100	•	Loniten
NICORANDIL			_	
▲ Tab 10 mg		60		Ikorel
▲ Tab 20 mg	32.28	60	•	Ikorel
PAPAVERINE HYDROCHLORIDE				
* Inj 12 mg per ml, 10 ml ampoule	257.12	5	•	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	•	Trental 400
Endothelin Receptor Antagonists				
AMBRISENTAN - Special Authority see SA1702 below - Retail p	•			
Tab 5 mg	1,550.00	30	/	Ambrisentan Mylan

⇒SA1702 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

30

✓ Ambrisentan Mylan

✓ Mylan

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic	
BOSENTAN - Special Authority see SA1991 below - Retail phar	macy				
Tab 62.5 mg	119.85	60	•	Bosentan Dr Reddy's	
Tab 125 mg	119.85	60	•	Bosentan Dr Reddy's	

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II. III. or IV: and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.1.2 Either
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil: or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy; or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan; or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or

continued...

	Subsidy		Fully	Brand or
(Ma	anufacturer's Price)	Subs	idised	Generic
	\$	Per	1	Manufacturer

continued...

- 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
- 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL - Special Authority see SA1992 below - Retail pharmacy		
Tab 25 mg	4	✓ Vedafil
Tab 50 mg1.70	4	✓ Vedafil
Tab 100 mg10.20	12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs): and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II: or
 - 3.2 PAH is in NYHA/WHO functional class III: or
 - 3.3 PAH is in NYHA/WHO functional class IV: and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

continued...

Subsid (Manufacturer		ılly Brand or ed Generic	
	Per	✓ Manufacturer	

continued...

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Prostacyclin Analogues

EPOPROSTENOL – Special Authority see SA1696 below – Retail pharmacy		
Inj 500 mcg vial36.61	1	✓ Veletri
Inj 1.5 mg vial73.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

ILOPROST - Special Authority see SA1705 below - Retail pharmacy

Nebuliser soln 10 mcg per ml, 2 ml740.10 30 ✓ Ventavis

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

ADAPALENE

- a) Maximum of 30 g per prescription
- b) Only on a prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - R	letail pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice: and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
 - 3.2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment; or
- 2 Patient is not of child bearing potential.

Note: Applicants are recommended to either have used or be familiar with using a decision support tool accredited by their professional body.

TRFTINOIN

Crm 0.5 mg per q − Maximum of 50 g per prescription......15.57 50 g OP ✓ ReTrieve

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 89

HYDROGEN PEROXIDE

DERMATOLOGICALS

	Subsidy (Manufacturer's F \$	Price) Subs Per	Fully Brand or sidised Generic Manufacturer
MUPIROCIN			
Oint 2%		15 g OP	
a). Oak an a marandation	(11.50)		Bactroban
a) Only on a prescriptionb) Not in combination			
ODIUM FUSIDATE [FUSIDIC ACID]			
Crm 2%	1.59	5 g OP	✓ Foban
 a) Maximum of 5 g per prescription 			
b) Only on a prescription			
c) Not in combination	1.50	E ~ OD	√ Cabo∷
Oint 2%	1.59	5 g OP	✓ <u>Foban</u>
a) Maximum of 5 g per prescriptionb) Only on a prescription			
c) Not in combination			
SULFADIAZINE SILVER			
Crm 1%	10.80	50 g OP	✓ Flamazine
a) Up to 250 g available on a PSO	10.00	30 g Oi	• I lalliazille
b) Not in combination			
Antitungals Tonical			
Antifungals Topical			
	ale page 96		
or systemic antifungals, refer to INFECTIONS, Antifunga	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga MOROLFINE a) Only on a prescription	als, page 96		
For systemic antifungals, refer to INFECTIONS, Antifunga MOROLFINE a) Only on a prescription b) Not in combination		5 ml OP	✓ MycoNail
for systemic antifungals, refer to INFECTIONS, Antifungal MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		5 ml OP	✓ <u>MycoNail</u>
For systemic antifungals, refer to INFECTIONS, Antifunga AMOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93	5 ml OP 20 g OP	✓ <u>MycoNail</u> ✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93		
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%	14.93	20 g OP	
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
ior systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP	✓ Clomazol
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP	✓ Clomazol Canesten
for systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP 20 g OP	✓ Clomazol Canesten
For systemic antifungals, refer to INFECTIONS, Antifungals, MOROLFINE a) Only on a prescription b) Not in combination Nail soln 5%		20 g OP 20 ml OP 20 g OP	✓ Clomazol Canesten Pevaryl

	Subsidy (Manufacturer's F \$	Price) Sub	Fully sidised	Brand or Generic Manufacturer
ICONAZOLE NITRATE	·			
Crm 2%	0.81	15 g OP	✓ M	lultichem
a) Only on a prescription		J		
b) Not in combination				
: Lotn 2%	4.36	30 ml OP		
	(10.03)		D	aktarin
a) Only on a prescription				
b) Not in combination				
Finct 2%	4.36	30 ml OP		
	(12.10)		D	aktarin
a) Only on a prescription				
b) Not in combination				

a) Only on a prescription

b) Not in combination

100 a ✓ Calamine-AFT

CROTAMITON

a) Only on a prescription

b) Not in combination

Crm 10%......3.29 20 g OP ✓ Itch-Soothe

MENTHOL - Only in combination

1) Only in combination with a dermatological base or proprietary Topical Corticosteriod - Plain

2) With or without other dermatological galenicals.

Crystals.......6.92 25 g 29.60 100 g ✓ MidWest ✓ MidWest

Corticosteroids Topical

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 79

Corticosteroids - Plain

BET	AMETHASONE DIPROPIONATE			
	Crm 0.05%	2.96	15 g OP	✓ Diprosone
	3	6.00	50 g OP	✓ Diprosone
	Oint 0.05%	2.96	15 g OP	✓ Diprosone
	3	6.00	50 g OP	✓ Diprosone
	Oint 0.05% in propylene glycol base	4.33	30 g OP	✓ Diprosone OV
BET	AMETHASONE VALERATE			
*	Crm 0.1%	4.53	50 g OP	✓ Beta Cream
*	Oint 0.1%	5.84	50 g OP	✓ Beta Ointment
*	Lotn 0.1%	25.00	50 ml OP	✓ Betnovate
CLC	DBETASOL PROPIONATE			
*	Crm 0.05%	2.18	30 g OP	✓ Dermol
*	Oint 0.05%	2.12	30 g OP	✓ Dermol

	Subsidy F (Manufacturer's Price) Subsidi			Brand or
	(Manufacturer's F \$	Price) Subs Per	idised	Generic Manufacturer
LODETACONE BUTYBATE	Ψ	1 01		Warranacturer
LOBETASONE BUTYRATE	5.00	00 00		
Crm 0.05%		30 g OP		
	(10.00)		t	Eumovate
YDROCORTISONE				
Crm 1% - Only on a prescription	3.70	100 g OP	✓	Hydrocortisone
, , ,		Ü		(PSM)
	17.15	500 g	✓	Hydrocortisone
	17.10	000 g		(PSM)
Powder – Only in combination	40.05	25 g	1	ABM
Up to 5% in a dermatological base (not proprietary To				
	opicai Conticosterio	a – Piain) with c	or with c	out other dermatologic
galenicals				
YDROCORTISONE AND PARAFFIN LIQUID AND LANOLII	N			
Lotn 1% with paraffin liquid 15.9% and lanolin 0.6% - On	nly on			
a prescription		250 ml	✓ [OP Lotn HC
YDROCORTISONE BUTYRATE			-	
Lipocream 0.1%	4.05	100 ~ OD	./ 1	
		100 g OP		Locoid Lipocream
Oint 0.1%		100 g OP	_	_ocoid
Milky emul 0.1%	12.33	100 ml OP	✓ [_ocoid Crelo
ETHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	1	<u>Advantan</u>
Oint 0.1%	4.46	15 g OP		Advantan
OMETASONE FUROATE		3 -	-	
Crm 0.1%	1.05	45 × OD		Tinnam Alaahal Fuan
GIII 0.1%		15 g OP	_	Elocon Alcohol Free
0: 10.40	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP	_	Elocon
	2.90	50 g OP	_	Elocon
Lotn 0.1%	4.50	30 ml OP	✓	<u>Elocon</u>
RIAMCINOLONE ACETONIDE				
Crm 0.02%	6.30	100 g OP	1	Aristocort
Oint 0.02%		100 g OP		Aristocort
5			-	<u></u>
Corticosteroids - Combination				
ETAMETHASONE VALERATE WITH SODIUM FUSIDATE [(FUSIDIC ACID)			
Crm 0.1% with sodium fusidate (fusidic acid) 2%		15 g OP		
om o. 170 with obdiant labidate (labidae acid) 270	(10.45)	10 9 01	F	ucicort
a) Maximum of 15 a per properintion	(10.43)		'	dolooit
a) Maximum of 15 g per prescription				
b) Only on a prescription				
YDROCORTISONE WITH MICONAZOLE - Only on a presi	cription			
	1.89	15 g OP	✓ <u>I</u>	Micreme H
Crm 1% with miconazole nitrate 2%	_ Only on a procesi	ntion		
Crm 1% with miconazole nitrate 2%		Juon		Name of the same
Crm 1% with miconazole nitrate 2% YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN		15 a OP	√ [rimatucori
Crm 1% with miconazole nitrate 2%YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5%	3.35	15 g OP		Pimafucort
Crm 1% with miconazole nitrate 2% YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35 3.35	15 g OP	√ F	Pimafucort
Crm 1% with miconazole nitrate 2% YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% Pimafucort Crm 1% with natamycin 1% and neomycin sulphate	3.35 3.35 ate 0.5% to be delist	15 g OP ted 1 April 2023	√ F	
Crm 1% with miconazole nitrate 2% YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5%	3.35 3.35 ate 0.5% to be delist	15 g OP ted 1 April 2023	√ F	
Crm 1% with miconazole nitrate 2% YDROCORTISONE WITH NATAMYCIN AND NEOMYCIN Crm 1% with natamycin 1% and neomycin sulphate 0.5% Oint 1% with natamycin 1% and neomycin sulphate 0.5% Pimafucort Crm 1% with natamycin 1% and neomycin sulphate	o3.35 o3.35 ate 0.5% to be delist YCIN AND NYSTAT	15 g OP ted 1 April 2023	√ F	
Crm 1% with miconazole nitrate 2%	3.35 3.35 3.35 3.46 3.77 3.35 3.77 3.78 3.78 3.78 3.78 3.78 3.78 3.78	15 g OP ted 1 April 2023 TIN	√ F	
Crm 1% with miconazole nitrate 2%	3.35 3.35 3.35 3.46 3.77 3.35 3.77 3.78 3.78 3.78 3.78 3.78 3.78 3.78	15 g OP ted 1 April 2023	✓ F	

Subsidy (Manufacturer's Price) \$

Fully Subsidised Per

Brand or Generic Manufacturer

Barrier Creams and Emollients			
Barrier Creams			
DIMETHICONE	4.00	500 100	
* Crm 5% pump bottle	4.30	500 ml OP	✓ healthE Dimethicone 5%
* Crm 10% pump bottle	4.52	500 ml OP	✓ healthE Dimethicone 10%
ZINC AND CASTOR OIL * Oint	4.65	500 g	✓ Boucher
Emollients			
AQUEOUS CREAM			
* Crm	1.73	500 g	✓ Boucher ✓ GEM Aqueous Cream
(Boucher Crm to be delisted 1 August 2022)			
CETOMACROGOL	1.00	F00 =	/ Ostomosyanal AFT
* Crm BP	1.99	500 g	✓ Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL Crm 90% with glycerol 10%	2.35	500 ml OP	✓ Boucher
Citi 60 / A milit gijodi di Toloni		000 1111 01	✓ Pharmacy Health Sorbolene with Glycerin
	3.10	1,000 ml OP	✓ Boucher
EMULSIFYING OINTMENT	0.40	F00 =	/ Faculaifuina
* Oint BP	3.40	500 g	Emulsifying Ointment ADE
OIL IN WATER EMULSION			
* Crm		500 g	✓ Fatty Cream AFT
	2.19		✓ O/W Fatty Emulsion Cream
Fatty Cream AFT to be Principal Supply on 1 September (O/W Fatty Emulsion Cream Crm to be delisted 1 September 202)			
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50%	5.35	500 ml OP	✓ healthE
URFA		000 1111 01	- Housting
* Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
WOOL FAT WITH MINERAL OIL - Only on a prescription		-	
* Lotn hydrous 3% with mineral oil		1,000 ml	
	(11.95) 1.40	050 ml OD	DP Lotion
	(4.53)	250 ml OP	DP Lotion
	5.60	1,000 ml	
	(20.53)		Alpha-Keri Lotion
	(23.91) 1.40	250 ml OP	BK Lotion
	(7.73)	200 IIII OF	BK Lotion
	. ,		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Other Dermatological Bases

PARAFFIN

Only in combination with a dermatological galenical or as a diluent for a proprietary Topical Corticosteroid - Plain.

Minor Skin Infections

POVIDONE IODINE			
Oint 10%	7.40	65 g OP	✓ Betadine
a) Maximum of 130 g per prescription			
b) Only on a prescription			
Antiseptic Solution 10%	4.15	100 ml	✓ Riodine
Antiseptic soln 10%	3.83	15 ml	✓ Riodine
	5.40	500 ml	✓ Riodine
Skin preparation, povidone iodine 10% with 30% alcohol	1.63	100 ml	
	(3.48)		Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol	1.63	100 ml	
	(7.78)		Pfizer

Parasiticidal Preparations

DIMETHICONE

 ★ Lotn 4%
 4.25
 200 ml OP
 ✓ healthE

 Dimethicone 4%

 Lotion

IVERMECTIN - Special Authority see SA1225 below - Retail pharmacy

Tab 3 mg − Up to 100 tab available on a PSO......17.20 4 Stromectol

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or

continued...

Subsidy (Manufacturer's Price)	Sı	Fully ibsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scables hyperinfestation (Crusted/ Norwegian scables); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria: Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Note: Ivermectin is no more effective than topical therapy for treatment of standard scabies infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- Strongyloidiasis.

PE	RM	١FT	ΉF	RIN

Crm 5%5.75	30 g OP	Lyderm
Lotn 5%	30 ml OP	✓ A-Scables

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Psoriasis and Eczema Preparations

ACITRETIN – Special Authority see SA2024 below – Retail pharmacy		
Cap 10 mg17.86	60	✓ Novatretin
Cap 25 mg41.36	60	✓ Novatretin

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
 - 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment;
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL	

Foam spray 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP 60 g OP 30 g OP	✓ Enstilar✓ <u>Daivobet</u>✓ <u>Daivobet</u>
CALCIPOTRIOL Oint 50 mcg per g		120 g OP	✓ Daivonex
COAL TAR Soln BP - Only in combination	36.25	200 ml	✓ Midwest
Up to 10% only in combination with a dermatological		ietary Topical C	Corticosteriod – Plai

ain

2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and

allantoin crm 2.5%	6.59	75 a OP	
	(8.00)	- 3 -	Egopsoryl TA
	3.43	30 g OP	0, ,
	(4.35)	-	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 on the next page - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks.

Cream 1%	15 g OP	✓ Elidel
----------	---------	----------

DERMATOLOGICALS

Subsidy	Full	y Brand or
(Manufacturer's Price)	Subsidise	d Generic
\$	Per 🗸	Manufacturer

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

★ Soln 2.3% with trolamine laurilsulfate and fluorescein sodium.......4.44 500 ml ✓ Pinetarsol

SALICYLIC ACID

Powder − Only in combination.......18.88 250 g ✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
- 2) With or without other dermatological galenicals.

SULPHUR

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

TACROLIMUS

Oint 0.1% − Special Authority see SA2074 below − Retail pharmacy.......33.00 30 g OP ✓ Zematop

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician, . Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

BETAMETHASONE VALERATE * Scalp app 0.1%	9.84	100 ml OP	✓ Beta Scalp
CLOBETASOL PROPIONATE * Scalp app 0.05%		30 ml OP	✓ Dermol
HYDROCORTISONE BUTYRATE Scalp lotn 0.1%		100 ml OP	✓ Locoid
KETOCONAZOLE Shampoo 2%		100 ml OP	✓ Sebizole
Grampoo 275	0.20	100 1111 01	✓ <u>Sebizole</u>

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	•	Manufacturer	

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

Contraceptives - Non-hormonal

Condoms

_	a) Maximum of 60 dev per prescription			supplied under Section 20
		17.02		✓ Gold Knight XL
		14.87	144	✓ Shield XL
K	60 mm		12	✓ Gold Knight XL
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		15.57	144	✓ Gold Knight
+	56 mm, strawberry		12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		15.57	144	✓ Gold Knight
	56 mm, chocolate		12	✓ Gold Knight
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
		11.64	144	✓ Moments
	56 mm, 0.08 mm thickness, red		10	✓ <u>Moments</u>
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO			
	\	11.64	144	✓ Moments
	56 mm, 0.08 mm thickness		10	✓ Moments
	b) Up to 60 dev available on a PSO	2.27	40	/ H
	a) Maximum of 60 dev per prescription			
	56 mm, 0.05mm thickness (bulk pack)	14.61	144	✓ Gold Knight
	b) Maximum of 60 dev per prescription		4	40.11
	a) Up to 60 dev available on a PSO			
)	15.57	144	✓ Gold Knight
	56 mm, 0.05 mm thickness		12	✓ Gold Knight
	b) Up to 60 dev available on a PSO	1.00	10	Cold Value
	a) Maximum of 60 dev per prescription			
	a) Maniana of CO day now prescription	11.04	144	✓ <u>Moments</u>
	ווווו סכ	0.97	10 144	
. ,	b) Maximum of 60 dev per prescription 56 mm	0.07	10	✓ Moments
	a) Up to 60 dev available on a PSO			
	a). Un to 60 day available on a BCO	11.04	144	• WOMENTS
		11.64	144	✓ Moments
. ;	53 mm, strawberry, red	0.95	10	✓ Moments
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	11.07	דדי	- momento
	or min, vilocolato, blown	11.64	144	✓ Moments
. ,	53 mm, chocolate, brown	0.95	10	✓ Moments
	b) Maximum of 60 dev per prescription			
	a) Up to 60 dev available on a PSO	11.72	1 1 1 1	· <u>momento</u>
	50 mm, 0.00 mm unounou	11.42	144	✓ Moments
. ;	53 mm, 0.05 mm thickness	0.95	10	✓ Moments
	b) Up to 60 dev available on a PSO			
	a) Maximum of 60 dev per prescription	11.04	144	<u>wionients</u>
	33 11111	11.64	144	✓ Moments
	49 mm – Up to 144 dev available on a PSO53 mm.		10	✓ <u>Moments</u> ✓ Moments
			144	

a) Maximumosidisedev per prescription b) Hindopassdayiqvailable on a PSO

GENITO-URINARY SYSTEM

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
60 mm (bulk pack) a) Maximum of 60 dev per prescription b) Up to 60 dev available on a PSO	14.87	144	•	Gold Knight XL

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- h) Only on a PSO

	b) Only on a PSO			
*	IUD 29.1 mm length × 23.2 mm width	18.45	1	✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	18.45	1	✓ Choice
				TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	15.50	1	Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- Either:
 - 1 Patient is on a Social Welfare benefit: or
 - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit: or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

*	Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up	to		
	84 tab available on a PSO	10.00	84	Mercilon 28

GENITO-URINARY SYSTEM

	Subsidy		Fully	Brand or
	(Manufacturer's Price)	S	ubsidised	Generic
	\$	Per	•	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	2.18	84	✓ N	licrogynon 20 ED
'	6.45	112		emme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)		N	Microgynon 30
 a) Higher subsidy of \$15.00 per 63 tab with Special Auth b) Up to 63 tab available on a PSO Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets 	•	the pr	evious paç	ge
		84	./ 1	evlen ED
Up to 112 tab available on a PSO				
	6.45	112	• -	emme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to)			
84 tab available on a PSO	6.95	84	✓ E	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U	D			
to 84 tab available on a PSO	•	84	✓ N	lorimin
	29.32	112	✓ N	lorimin
	20.02		- 1	

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

LEVONORGESTREL			
* Tab 30 mcg - Up to 84 tab available on a PSO	16.50	84	✓ Microlut
	22.00	112	✓ Microlut
* Subdermal implant (2 x 75 mg rods) - Up to 3 pack available)		
on a PSO	106.92	1	✓ <u>Jadelle</u>
MEDROXYPROGESTERONE ACETATE			
Inj 150 mg per ml, 1 ml syringe - Up to 5 inj available on a PS	SO7.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Si Per	Fully ubsidised	Brand or Generic Manufacturer	
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	✓ <u>N</u>	oriday 28	
Emergency Contraceptives					
LEVONORGESTREL * Tab 1.5 mg		1 Part I of	•	ostinor-1	

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate			
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43 (24.00)	100 g OP	Aci-Jel	
CLOTRIMAZOLE			
* Vaginal crm 1% with applicators	35 g OP	Clomazol	
* Vaginal crm 2% with applicators	20 g OP	Clomazol	
MICONAZOLE NITRATE			
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme	
NYSTATIN	Ü		
Vaginal crm 100,000 u per 5 g with applicator(s)4.00	75 g OP	✓ Nilstat	

Myometrial and Vaginal Hormone Preparations

160.00	5	DBL Ergometrine
6.62	15 g OP	✓ Ovestin
6.86	15	✓ Ovestin
3.98	5	Oxytocin BNM
4.98	5	Oxytocin BNM
ole on a PSO		
e32.40	5	✓ Syntometrine
	160.00 6.62 6.86 4.98 ble on a PSO e32.40	6.62 15 g OP6.86 153.98 54.98 5 ble on a PSO

GENITO-URINARY SYSTEM

	Subsidy		ully	Brand or
(Ma	nufacturer's Price)	Subsid	ised	Generic
	\$	Per	✓	Manufacturer

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

Pregnancy Test

✓ Smith BioMed Rapid

Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 106

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

* Tab 5 mg4.81

.81 100 **✓ Ricit**

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Note: Patients with enlarged prostates are the appropriate candidates for therapy with finasteride.

Alpha-1A Adrenoreceptor Blockers

TAMSULOSIN HYDROCHLORIDE - Special Authority see SA1032 below - Retail pharmacy

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

POTASSIUM CITRATE

Oral liq 3 mmol per ml - Special Authority see SA1083 on the

next page − Retail pharmacy.......31.80 200 ml OP ✓ Biomed

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	5	Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has recurrent calcium oxalate urolithiasis; and
- 2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE			
* Grans eff 4 g sachets	2.22	28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE			
Tab 5 mg	2.05	30	Solifenacin Mylan
Tab 10 mg	3.72	30	✓ Solifenacin Mylan

Detection of Substances in Urine

ORTHO-TOLIDINE	<u>. </u>	_	_
* Compound diagnostic sticks	7.50	50 test OP	
, , , , , , ,	(8.25)		Hemastix
TETRABROMOPHENOL			
* Blue diagnostic strips	7.02	100 test OP	
	(13.92)		Albustix

Obstetric Preparations

Antiprogesterones

MIFEPRISTONE			
Tab 200 mg	60.00	1	✓ Mifegyne
-	180.00	3	✓ Mifegyne

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

Subsidy (Manufacturer's Price)	Cub	Fully	Brand or Generic
(Manufacturer's Price)		osiaisea	
\$	Per	/	Manufacturer

Calcium Homeostasis

CALCITONIN			
* Inj 100 iu per ml, 1 ml ampoule	121.00	5	Miacalcic
CINACALCET - Special Authority see SA1618 below - Retail	pharmacy		
Tab 30 mg - Wastage claimable	42.06	28	✓ Cinacalet Devatis
Tab 60 mg - Wastage claimable	84.12	28	✓ Cinacalet Devatis

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

Inj 4 mg per 5 ml, vial − Special Authority see SA2109 below −

Retail pharmacy.......18.00 1

✓ Zoledronic acid

Mvlan

⇒SA2109 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement: and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia .

Note: Indications marked with * are unapproved indications.

C	Corticosteroids and Related Agents for Systemic Use			
	TAMETHASONE SODIUM PHOSPHATE WITH BETAMETHASONE ACETATE			
*	Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml19.20	5		
	(36.96)		Celestone	
			Chronodose	
DE	XAMETHASONE			
*	Tab 0.5 mg - Up to 60 tab available on a PSO1.50	30	✓ <u>Dexmethsone</u>	
*	Tab 4 mg - Up to 30 tab available on a PSO2.65	30	✓ <u>Dexmethsone</u>	
	Oral liq 1 mg per ml48.15	25 ml OP	✓ Biomed	
DE	XAMETHASONE PHOSPHATE			
	Dexamethasone phosphate injection will not be funded for oral use.	40	4.5	
*	Inj 4 mg per ml, 1 ml ampoule – Up to 5 inj available on a PSO9.25	10	✓ Dexamethasone Phosphate Panpharma	
*	Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 16.37	10	✓ Dexamethasone Phosphate Panpharma	
	UDROCORTISONE ACETATE		i anphama	
*	Tab 100 mcg11.46	100	✓ Florinef	
	3	100	• Homici	
# *	'DROCORTISONE Tab 5 mg8.10	100	✓ Douglas	
	Tab 20 mg	100	✓ Douglas ✓ Douglas	
	Inj 100 mg vial4.38	1	✓ Solu-Cortef	
	a) Up to 5 inj available on a PSO b) Only on a PSO	·	<u> </u>	
ME	THYLPREDNISOLONE			
*	Tab 4 mg112.00	100	✓ Medrol	
*	Tab 100 mg223.10	20	✓ Medrol	
ME	THYLPREDNISOLONE (AS SODIUM SUCCINATE)			
	Inj 40 mg vial	1	✓ Solu-Medrol-Act- O-Vial	•
	Inj 125 mg vial34.10	1	✓ Solu-Medrol-Act- O-Vial	-
	Inj 500 mg vial	1	✓ Solu-Medrol-Act O-Vial	-
	Inj 1 g vial32.84	1	✓ Solu-Medrol	

	Subsidy	,	Fully	Brand or
	(Manufacturer's Pric	e) Sub: Per	sidised •	Generic Manufacturer
ETHYLPREDNISOLONE ACETATE	· ·			
Inj 40 mg per ml, 1 ml vial	47.06	5	✓	Depo-Medrol
REDNISOLONE		·		- opoou.o.
Oral liq 5 mg per ml – Up to 30 ml available on a PSO Restricted to children under 12 years of age.	6.00	30 ml OP	✓	<u>Redipred</u>
REDNISONE				
: Tab 1 mg	18.58	500		Apo-Prednisone
	24.24	500	_	Prednisone Clinect
F Tab 2.5 mg	21.04	500		Apo-Prednisone Prednisone Clinect
Tab 5 mg - Up to 30 tab available on a PSO	10.30	500		Apo-Prednisone
Tab 3 mg - op to 30 tab available on a 1 30	19.00	300		Prednisone Clinect
F Tab 20 mg - Up to 30 tab available on a PSO	50.51	500	1	Apo-Prednisone Prednisone Clinect
Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 20 mg to be delisted 1 November 2022)				
ETRACOSACTRIN				
Inj 250 mcg per ml, 1 ml ampoule	75.00	1		UK Synacthen S29
				AU Synacthen
for Inj 1 mg per ml, 1 ml ampoule	690.00	1	1	Synacthen Synacthen Depot Synacthene Retard S29
RIAMCINOLONE ACETONIDE				
Inj 10 mg per ml, 1 ml ampoule	20.80	5	✓	Kenacort-A 10
Inj 40 mg per ml, 1 ml ampoule		5		Kenacort-A 40
Sex Hormones Non Contraceptive				
Androgen Agonists and Antagonists				
YPROTERONE ACETATE				
Tab 50 mg	14.37	50	1	Siterone
Tab 100 mg	28.03	50	1	Siterone
ESTOSTERONE				
Patch 5 mg per day	90.00	30	1	Androderm
ESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	✓	Depo-Testosterone
ESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12.98	1	1	Sustanon Ampoules
ESTOSTERONE UNDECANOATE				•
Cap 40 mg - Subsidy by endorsement	21.00	60	1	Andriol Testocaps
Subsidy by endorsement - subsidised for patients who was	were taking testoste	erone undec	anoate	cap 40mg prior to
1 November 2021 and the prescription is endorsed acco	ordingly. Pharmacis	sts may ann	otate th	ne prescription as endo
where there exists a record of prior dispensing of testost	terone undecanoate		in the	preceding 12 months.
Inj 250 mg per ml, 4 ml vial	86.00	1	✓	Reandron 1000

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Hormone Replacement Therapy - Systemic

Prescribing Guideline

Oestrogens

HRT should be taken at the lowest dose for the shortest period of time necessary to control symptoms. Patients should be reviewed 6 monthly in line with the updated NZGG "Evidence-based Best Practice Guideline on Hormone Replacement Therapy March 2004".

OESTRADIOL - See prescribing guideline above			
* Tab 1 mg	4.12	28 OP	
•	(11.10)		Estrofem
* Tab 2 mg	4.12	28 OP	
	(11.10)		Estrofem
Patch 50 mcg per 24 hours	7.04 [′]	4	✓ Climara
a) No more than 1 patch per week			
b) Only on a prescription			
Patch 25 mcg per day	6.12	8	✓ Estradot
a) No more than 2 patch per week			
b) Only on a prescription			
Patch 50 mcg per day	7.04	8	✓ Estradot 50 mcg
			•

a) No more than 2 patch per week b) Only on a prescription			
Patch 75 mcg per day	7.91	8	✓ Estradot
a) No more than 2 patch per week			
b) Only on a prescription			

Patch 100 mcg per day	7.91	8	✓ Estradot
a) No more than 2 patch per week			

OESTRADIOL VALERATE – See prescribing guideline above			
* Tab 1 mg	12.36	84	✓ Progynova
* Tab 2 mg	12.36	84	✓ Progynova
OESTROGENS - See prescribing guideline above			
* Conjugated equipe tab 300 mcg	3.01	28	

~~	Conjugated, equilie tab 500 meg		20	
		(17.50)		Premarin
*	Conjugated, equine tab 625 mcg	4.12	28	
		(17.50)		Premarin

Progestogens

b) Only on a prescription

ME	DROXYPROGESTERONE ACETATE - See prescribing guideline abo	ove		
*	Tab 2.5 mg4	.69	30	✓ Provera
	Tab 5 mg17		100	✓ Provera
	Tab 10 mg		30	✓ Provera

Subsidy

(Manufacturer's Price)

(18.10)

(18.10)

Fully

Subsidised

28 OP

Brand or

Generic

Kliogest

Trisequens

	\$	Per	1	Manufacturer	
Progestogen and Oestrogen Combined Prepar	ations				
OESTRADIOL WITH NORETHISTERONE - See prescribing gr	uideline on the pre	vious page			
* Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP			
	(18.10)		ŀ	Kliovance	
* Tab 2 mg with 1 mg norethisterone acetate	5.40	28 OP			

Other Oestrogen Preparations

ETHINYLOESTRADIOL - Subsidy by endorsement

Tab 2 mg with 1 mg norethisterone acetate (10), and 2 mg

oestradiol tab (12) and 1 mg oestradiol tab (6)...............................5.40

Subsidy by endorsement – Subsidised for patients who were taking ethinyloestradiol prior to 1 March 2022 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of ethinyloestradiol.

OFSTRIOL

Other Progestogen Preparations

LEVONORGESTREL			
* Intra-uterine device 52 mg	269.50	1	✓ Mirena
* Intra-uterine device 13.5 mg	215.60	1	 Jaydess
MEDROXYPROGESTERONE ACETATE Tab 100 mg	116 15	100	✓ Provera HD
3	110.15	100	▼ Flovela IID
NORETHISTERONE			_
* Tab 5 mg – Up to 30 tab available on a PSO	5.49	30	✓ Primolut N
PROGESTERONE			
Cap 100 mg - Special Authority see SA1609 below - Retail			
pharmacy	16.50	30	Utrogestan

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Fither:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and

Subsidy	Subsidy F		Brand or
(Manufacturer's Price)	(Manufacturer's Price) Subsid		Generic
\$	Per	1	Manufacturer

continued...

- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

3-11-3			
CARBIMAZOLE			
* Tab 5 mg	7.56	100	✓ Neo-Mercazole
Neo-Mercazole to be Principal Supply on 1 September 2022			
LEVOTHYROXINE			
* Tab 25 mcg	5.55	90	✓ Synthroid
* Tab 50 mcg	1.71	28	✓ Mercury Pharma
· ·	5.79	90	✓ Synthroid
	64.28	1,000	✓ Eltroxin
* Tab 100 mcg	1.78	28	✓ Mercury Pharma
·	6.01	90	✓ Synthroid
	66.78	1,000	✓ Eltroxin
PROPYLTHIOURACIL - Special Authority see SA1199 below - Reta	il pharmacy	,	
Propylthiouracil is not recommended for patients under the age of treatments are contraindicated.	18 years u	nless the patie	ent is pregnant and other
Tab 50 mg	35.00	100	✓ PTU S29

SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SC	MATROPIN (OMNITROPE) - Special Authority see SA203	2 below – Retail pha	rmacy	
*	Inj 5 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
*	Inj 15 mg cartridge	139.50	1	✓ Omnitrope

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>

continued...

- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred: and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist.

Subsidy (Manufacturer's Price)	Full Subsidise		
(Wandacties S Frice)	Per •	Manufacturer	

continued...

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or

bsidy	Fully	Brand or
turer's Price) Subsid	dised	Generic
 \$ Per	✓	

continued...

if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and

- 5 Either:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

1 All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	1	Manufacturer

continued...

- 1.1 The patient has been treated with somatropin for < 12 months; and
- 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
- 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
- 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or

2 All of the following:

- 2.1 The patient has been treated with somatropin for more than 12 months; and
- 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
- 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
- 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or

3 All of the following:

- 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
- 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3.3 The patient has severe growth hormone deficiency (see notes); and
- 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

G)S	FR	FΙ	IN

Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe	122.37	1	✓ Teva

LEUPRORELIN

Additional subsidy by endorsement where the patient is a child or adolescent and is unable to tolerate administration of goserelin and the prescription is endorsed accordingly.

Inj 3.75 mg prefilled dual chamb	er syringe - Higher subsidy of
400400 4:: ::	

\$221.60 per 1 inj with Endorsement	66.48	1	
,	(221.60)		Lucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy			
of \$591.68 per 1 inj with Endorsement	177.50	1	
, ,	(591.68)		Lucrin Depot 3-month

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
Vasopressin Agonists				
DESMOPRESSIN Wafer 120 mcg DESMOPRESSIN ACETATE	47.00	30	✓ N	Minirin Melt
Tab 100 mcg		30 30		Ainirin Ainirin
▲ Nasal spray 10 mcg per dose		6 ml OP	✓ [Desmopressin- PH&T
Inj 4 mcg per ml, 1 ml	67.18	10	✓ N	<i>l</i> inirin
Other Endocrine Agents				
CABERGOLINE Tab 0.5 mg - Maximum of 2 tab per prescription: can be				

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Hyperprolactinemia; or
- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

CLOMIFENE CITRATE Tab 50 mg	29.84	10	✓ Mylan Clomiphen \$29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Anthelmintics** ALBENDAZOLE - Special Authority see SA1318 below - Retail pharmacy 60 Fskazole S29 ⇒SA1318 Special Authority for Subsidy Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids. Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment. MEBENDAZOLE - Only on a prescription Tab 100 mg7.97 6 Vermox 15 ml (7.53)Vermox PRAZIQUANTFI 8 Biltricide **Antibacterials** a) For topical antibacterials, refer to DERMATOLOGICALS, page 61 b) For anti-infective eve preparations, refer to SENSORY ORGANS, page 236 Cephalosporins and Cephamycins CEFACLOR MONOHYDRATE Cap 250 mg......24.70 ✓ Ranbaxy-Cefaclor 100 ✓ Ranbaxy-Cefaclor S29 S29 ✓ Ranbaxy-Cefaclor 100 ml ✓ Ranbaxy-Cefaclor S29 S29 **CEFALEXIN** 20 ✓ Cephalexin ABM ✓ Cephalexin ABM 20 Grans for oral lig 25 mg per ml - Wastage claimable.....8.75 100 ml ✓ Cefalexin Sandoz ✓ Cefalexin Sandoz Grans for oral liq 50 mg per ml - Wastage claimable......11.75 100 ml CEFAZOLIN - Subsidy by endorsement Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly. CEFTRIAXONE - Subsidy by endorsement a) Up to 10 ini available on a PSO b) Subsidised only if prescribed for a dialysis or cystic fibrosis patient, or the treatment of gonorrhoea, or the treatment of pelvic inflammatory disease, or the treatment of suspected meningococcal disease, and the prescription or PSO is endorsed accordingly. ✓ Ceftriaxone-AFT ✓ Ceftriaxone-AFT

Only if prescribed for prophylaxis of endocarditis and the prescription is endorsed accordingly

CEFUROXIME AXETIL - Subsidy by endorsement

✓ Zinnat

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	1	Manufacturer	

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg	8.19	30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO		2	✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage			
claimable	16.97	15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Either:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and
- 3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

1 Atypical mycobacterial infection; or

FRYTHROMYCIN (AS LACTORIONATE)

2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial	10.00	1	✓ Erythrocin IV
ERYTHROMYCIN ETHYL SUCCINATE			
Tab 400 mg	16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSOb) Up to 2 x the maximum PSO quantity for RFPP			
Grans for oral liq 200 mg per 5 ml	5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSOb) Up to 2 x the maximum PSO quantity for RFPPc) Wastage claimable			
Grans for oral liq 400 mg per 5 ml	6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO			
b) Wastage claimable			
ERYTHROMYCIN STEARATE			
Tab 500 mg	29.90	100	
•	(44.58)		ERA
(ERA Tab 500 mg to be delisted 1 September 2022)			
ROXITHROMYCIN			
Tab disp 50 mg	8.29	10	✓ Rulide D
Restricted to children under 12 years of age.			
Tab 150 mg	8.28	50	✓ Arrow- Roxithromycin
Tab 300 mg	16.33	50	✓ Arrow- Roxithromycin

(Rulide D Tab disp 50 mg to be delisted 1 September 2022)

	Subsidy (Manufacturer's Price		Fully ubsidised	
	\$	Per		Manufacturer
Penicillins				
AMOXICILLIN				
Cap 250 mg	22.50	500	✓	Alphamox
 a) Up to 30 cap available on a PSO 				
b) Up to 10 x the maximum PSO quantity for RFPP				
Cap 500 mg	36.98	500		Alphamox
a) Up to 30 cap available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPP			_	
Grans for oral liq 125 mg per 5 ml	1.40	100 ml	•	Alphamox 125
a) Up to 200 ml available on a PSO				
b) Wastage claimable	4.70	400	,	Alb050
Grans for oral liq 250 mg per 5 ml	1./3	100 ml	•	Alphamox 250
a) Up to 300 ml available on a PSO				
b) Up to 10 x the maximum PSO quantity for RFPPc) Wastage claimable				
Inj 250 mg vial	15 97	10	1	Ibiamox
Inj 500 mg vial		10		Ibiamox
Inj 1 g vial – Up to 5 inj available on a PSO		10		Ibiamox
AMOXICILLIN WITH CLAVULANIC ACID		. •		
Tab 500 mg with clavulanic acid 125 mg – Up to 30 tab				
available on a PSO	0.80	10	1	Curam Duo 500/125
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 r		10	•	Curain Duo 300/123
per ml	•	100 ml	1	Augmentin
a) Up to 200 ml available on a PSO		100 1111	_	, taginonim
b) Wastage claimable				
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 r	ma			
per ml – Up to 200 ml available on a PSO		00 ml OF	•	Curam
BENZATHINE BENZYLPENICILLIN				
Inj 900 mg (1.2 million units) in 2.3 ml syringe – Up to 5 inj				
available on a PSO	375 97	10	1	Bicillin LA
		10		Diomini Ert
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial – Up to 5 inj available on a PS	SO 11.00	10	1	Sandoz
	30 11.09	10	•	Sandoz
FLUCLOXACILLIN Cap 250 mg – Up to 30 cap available on a PSO	15.70	250	./	Flucloxacillin-AFT
Cap 500 mg - Up to 30 cap available on a PSO		250 500		Flucioxacillin-AFT
Grans for oral lig 25 mg per ml		100 ml		AFT
a) Up to 200 ml available on a PSO		100 1111	-	<u>a</u>
b) Wastage claimable				
Grans for oral liq 50 mg per ml	3.68	100 ml	1	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Inj 250 mg vial	17.56	10	✓	Flucloxin
Inj 500 mg vial		10	✓	Flucloxin
Inj 1 g vial – Up to 5 inj available on a PSO	5.70	5	✓	Flucil

	Subsidy Manufacturer's Price)		Fully dised	
<u> </u>	\$	Per	1	Manufacturer
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				-
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK
Cap 500 mg		50	✓	Cilicaine VK
a) Up to 20 cap available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
Grans for oral liq 125 mg per 5 ml	2.99	100 ml	1	AFT
a) Up to 200 ml available on a PSO				
b) Wastage claimable				
Grans for oral liq 250 mg per 5 ml	3.99	100 ml	✓	AFT
a) Up to 300 ml available on a PSO				
b) Up to 2 x the maximum PSO quantity for RFPP				
c) Wastage claimable				
PROCAINE PENICILLIN				
Inj 1.5 g in 3.4 ml syringe - Up to 5 inj available on a PSO	123.50	5	1	Cilicaine
Takes avalines				
Tetracyclines				
DOXYCYCLINE				
* Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	1	Doxine
MINOCYCLINE HYDROCHLORIDE				
* Tab 50 mg - Additional subsidy by Special Authority see				
SA1355 below – Retail pharmacy	5.79	60		
o	(12.05)	••		Mino-tabs
* Cap 100 mg		100		- /
	(52.04)			Minomycin
OA4055 On a dal Authority for Manufacturery Bullet	` '			•

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

 $\label{temperature} \textbf{TETRACYCLINE} \ - \ \textbf{Special Authority see} \ \frac{\textbf{SA1332 below}}{\textbf{SA1332 below}} - \ \textbf{Retail pharmacy}$

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 61

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis: or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	
CLINDAMYCIN	·		
	4.61	24	✓ Dalacin C
Cap hydrochloride 150 mg			
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	✓ Dalacin C
COLISTIN SULPHOMETHATE - Retail pharmacy-Specialist - S	Subsidy by endorseme	ent	
Only if prescribed for dialysis or cystic fibrosis patient and the	e prescription is endo	rsed	accordingly.
Inj 150 mg	65.00	1	✓ Colistin-Link
GENTAMICIN SULPHATE			
	05.00	5	✓ DBL Gentamicin
Inj 10 mg per ml, 1 ml ampoule – Subsidy by endorsement.			
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	ct intection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	ct infection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓ Pfizer
, , ,	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient endorsed accordingly.	or complicated urinary	trac	ct infection and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retai No patient co-payment payable	I pharmacy		
Tab 400 mg	42.00	5	✓ <u>Avelox</u>

⇒SA1740 Special Authority for Subsidy

Initial application — (**Tuberculosis**) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications: or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications; or
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and

	Subsidy (Manufacturer's Price) \$	Subsi Per	Fully dised	Brand or Generic Manufacturer	
continued 3 Treatment is only for 7 days.					
Initial application — (Penetrating eye injury) only from an opht requires prophylaxis following a penetrating eye injury and treatment of the indications marked with * are unapproved indications.			or 1 mon	nth where the patient	
PAROMOMYCIN - Special Authority see SA1689 below - Retail	pharmacy				
Cap 250 mg	126.00	16	✓ Hu	umatin \$29	
■ SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinic month for applications meeting the following criteria: Either:	al microbiologist or q	gastroente	rologist.	Approvals valid for 1	
1 Patient has confirmed cryptosporidium infection; or2 For the eradication of Entamoeba histolyica carriage.					
Renewal only from an infectious disease specialist, clinical microbapplications meeting the following criteria: Either:	iologist or gastroent	erologist.	Approva	als valid for 1 month fo	r
 Patient has confirmed cryptosporidium infection; or For the eradication of Entamoeba histolyica carriage. 					
PYRIMETHAMINE – Special Authority see SA1328 below – Retai Tab 25 mg		30	✓ Da	araprim \$29	
Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of	a period of 3 months		notified	for applications meeti	ng
SODIUM FUSIDATE [FUSIDIC ACID] Tab 250 mg	67.85	36	√ Fu	ıcidin	
SULFADIAZINE SODIUM – Special Authority see SA1331 below Tab 500 mg	– Retail pharmacy	56	✓ W	ockhardt \$29	
➤ SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid the following criteria: Any of the following:					ng
For the treatment of toxoplasmosis in patients with HIV for For pregnant patients for the term of the pregnancy; or For infants with congenital toxoplasmosis until 12 months or	•	s; or			
TOBRAMYCIN Inj 40 mg per ml, 2 ml vial – Subsidy by endorsement Only if prescribed for dialysis or cystic fibrosis patient and				obramycin Mylan gly.	
Solution for inhalation 60 mg per ml, 5 ml — Subsidy by endorsement		6 dose		bramycin BNM	
b) Only if prescribed for a cystic fibrosis patient and the p	prescription is endors	sed accord	ingly.		
TRIMETHOPRIM * Tab 300 mg – Up to 30 tab available on a PSO	18.55	50	✓ <u>TN</u>	<u>NP</u>	

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [C	O-TRIMOXAZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole to 30 tab available on a PSO		500	✓ <u>T</u>	<u>risul</u>
* Oral liq 8 mg sulphamethoxazole 40 mg per ml - available on a PSO	•	00 m	√ D	eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis p difficile following metronidazole failure and the pr	escription is endorsed accordingly		s or for treat	tment of Clostridium
Inj 500 mg vial	2.35	1	✓ <u>M</u>	<u>lylan</u>

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 62
- b) For topical antifungals refer to GENITO URINARY, page 75

FLUCONAZOLE

Cap 50 mg2.75	28	✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg	28	✓ Mylan
Powder for oral suspension 10 mg per ml – Special Authority		
see SA1359 below – Retail pharmacy109.34	35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg4.27	15	✓ Itrazole
Oral liq 10 mg per ml - Special Authority see SA1322 on the		
next page – Retail pharmacy	150 ml OP	✓ Sporanox

Subsidy		Fully	Brand or	
(Manufacturer's Price)		bsidised	Generic	
	Per	✓	Manufacturer	

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOLE

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 below - Ret	ail pharmacy		
Tab modified-release 100 mg		24	✓ Noxafil
Oral liq 40 mg per ml		105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Fither:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

* Tab 250 mg	8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the next page	ge – Retail phar	macy	
Tab 50 mg	91.00	56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,523.22	70 ml	✓ Vfend

Subsidy (Manufacturer:		Fully Subsidised	Brand or Generic	
\$	Pe	er 🗸	Manufacturer	

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised: and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA16	84 below – Retail pharmacy			
Tab 15 mg	400.00	100	✓ Sanofi	
-			Primaguine \$2	9

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg	24.48	10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ <u>Arrow-Ornidazole</u>

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antituberculotics and Antileprotics

Note: There is no co-payment charge for all pharmaceuticals listed in the Antituberculotics and Antileprotics group regardless of immigration status.

CLOFAZIMINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist.

CYCLOSERINE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician.
- Cap 250 mg.......344.00 60 **✓ Cyclorin** 529

DAPSONE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or dermatologist

Dapsone	100	Tab 25 mg268.50	Т
Dapsone	100	Tab 100 mg329.50	Т

ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an infectious disease physician, clinical microbiologist or respiratory physician

Tab 100 mg	85.73	100	✓ EMB Fatol S29
Tab 400 mg	49.34	56	✓ Myambutol S29

ISONIAZID - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician
- ***** Tab 100 mg23.00 100 ✓ <u>PSM</u>

ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist

- a) No patient co-payment payable
- b) Prescriptions must be written by, or on the recommendation of, an internal medicine physician, paediatrician, clinical microbiologist, dermatologist or public health physician

*	Tab 100 mg with rifampicin 150 mg	89.82	100	✓ Rifinah
*	Tab 150 mg with rifampicin 300 mg	179.13	100	Rifinah

PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

PROTIONAMIDE - Retail pharmacy-Specialist

- a) No patient co-payment payable
- Prescriptions must be written by, or on the recommendation of, an infectious disease specialist, clinical microbiologist or respiratory physician

		Subsidy (Manufacturer's Price \$	e) Per	Full Subsidise	ed Generic
PΥ	RAZINAMIDE – Retail pharmacy-Specialist				
	 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendarespiratory physician 	ation of, an infectious	disease	e physicia	an, clinical microbiologist
*	Tab 500 mg	64.95	100	•	AFT-Pyrazinamide
	ABUTIN - Retail pharmacy-Specialist				•
	a) No patient co-payment payable b) Prescriptions must be written by, or on the recommendation and transfer payable.	ation of, an infectious	disease	e physicia	an, respiratory physician c
k	gastroenterologist Cap 150 mg	299.75	30		Mycobutin
	AMPICIN – Subsidy by endorsement				,
	a) No patient co-payment payable				
	 For confirmed recurrent Staphylococcus aureus infectio antimicrobial based on susceptibilities and the prescripti Retail pharmacy - Specialist. Specialist must be an inte paediatrician, or public health physician. 	ion is endorsed accor ernal medicine physic	rdingly; ian, clin	can be w	vaived by endorsement - obiologist, dermatologist,
K	Cap 150 mg		100		Rifadin
	Cap 300 mg Oral lig 100 mg per 5 ml		100 60 ml		∕ <u>Rifadin</u> ∕ Rifadin
	epatitis B Treatment				
K	TECAVIR Tab 0.5 mg		30	•	Entecavir Sandoz
Αl	MIVUDINE - Special Authority see SA1685 below - Retail p	•	28		Zetlam
	Tab 100 mg Oral liq 5 mg per ml		20 240 ml (Zeffix
-	SA1685 Special Authority for Subsidy				
it	ial application only from a relevant specialist or medical pra		mmend	ation of a	a relevant specialist.
eı	provals valid for 1 year where used for the treatment or prevenewal from any relevant practitioner. Approvals valid for 2 y		the trea	itment or	prevention of hepatitis B.
	Tenofovir disoproxil prescribed under endorsement for the talking antiretrovirals for the purposes of Special Authority SA2139		cluded	in the co	unt of up to 4 subsidised
+	Tab 245 mg (300.6 mg as a maleate)		30	•	Tenofovir Disoproxil Mylan
÷	Tab 245 mg (300.6 mg as a succinate)	38.10	30	•	Tenofovir Disoproxil Teva
	nofovir Disoproxil Teva Tab 245 mg (300.6 mg as a succina	te) to be delisted 1 D	ecembe	er 2022)	
H	erpesvirus Treatments				
.C	CLOVIR Tab dispersible 200 mg	1.60	25	,	/ Lovir
K	Tab dispersible 400 mg		56		Lovir
÷	Tab dispersible 800 mg	5.98	35	•	Lovir

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VALACICLOVIR				
Tab 500 mg	6.50	30	_	<u>aclovir</u>
Tab 1,000 mg	13.76	30	✓ <u>v</u>	/aclovir
VALGANCICLOVIR – Special Authority see SA1993 below – Ret Tab 450 mg		60	√ ∨	/alganciclovir
1 ab 450 mg	132.00	00	• <u>•</u>	Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — **(transplant cytomegalovirus prophylaxis)** only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or

2 Both:

- 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
- 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months); and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive; and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Roth:

Subsidy (Manufacturer's Price)	Full Subsidise	' · · · · · ·	
 \$	Per •	Manufacturer	

continued...

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 84 OP ✓ Maviret

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990,

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 below

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 103 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

* Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a

(Teva Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a succinate) to be delisted 1 December 2022)

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

continued...

✓ Teva

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (https://ashm.org.au/HIV/PrEP/)

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Note: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines (https://ashm.org.au/HIV/PrEP/)

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
	Per	✓	Manufacturer

continued...

ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Non-nucleosides Reverse Transcriptase Inhibitors

EFAVIRENZ – Special Authority see SA2139 on the previous page – Retail pharmacy						
Tab 200 mg	190.15	90	✓ Stocrin			
Tab 600 mg	63.38	30	✓ Stocrin			
ETRAVIRINE - Special Authority see SA2139 on the previous page - Retail pharmacy						
Tab 200 mg	770.00	60	✓ Intelence			

INFECTIONS - AGENTS FOR SYSTEMIC USE						
	Subsidy (Manufacturer's P \$		Fully Brand or dised Generic Manufacturer			
NEVIRAPINE - Special Authority see SA2139 on page 103 - F Tab 200 mg		60	✓ <u>Nevirapine</u> Alphapharm			
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Viramune Suspension			
Nucleosides Reverse Transcriptase Inhibitors						
ABACAVIR SULPHATE - Special Authority see SA2139 on particle 300 mg	180.00	armacy 60 240 ml OP	✓ Ziagen ✓ Ziagen			
ABACAVIR SULPHATE WITH LAMIVUDINE – Special Authorit Note: abacavir with lamivudine (combination tablets) count anti-retroviral Special Authority.						
Tab 600 mg with lamivudine 300 mg	63.00	30	✓ Kivexa			
EFAVIRENZ WITH EMTRICITABINE AND TENOFOVIR DISOF pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxil of anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxil of the control	counts as three an	ti-retroviral med	dications for the purposes of the			
245 mg (300 mg as a maleate)		30	✓ Mylan			
EMTRICITABINE – Special Authority see SA2139 on page 103 Cap 200 mg	307.20	y 30	✓ Emtriva			
LAMIVUDINE - Special Authority see SA2139 on page 103 - F Tab 150 mg		60	✓ <u>Lamivudine</u> Alphapharm			
Oral liq 10 mg per ml	102.50	240 ml OP	✓ 3TC			
ZIDOVUDINE [AZT] - Special Authority see SA2139 on page 1		•				
Cap 100 mgOral liq 10 mg per ml		100 200 ml OP	✓ Retrovir✓ Retrovir			
ZIDOVUDINE [AZT] WITH LAMIVUDINE — Special Authority se Note: zidovudine [AZT] with lamivudine (combination tablet the anti-retroviral Special Authority. Tab 300 mg with lamivudine 150 mg	ee SA2139 on pag ts) counts as two a	e 103 – Retail p	harmacy			
Protease Inhibitors						
ATAZANAVIR SULPHATE – Special Authority see SA2139 on Cap 150 mg Cap 200 mg	141.68 188.91	pharmacy 60 60	✓ Teva ✓ Teva			
DARUNAVIR - Special Authority see SA2139 on page 103 - R Tab 400 mg Tab 600 mg	132.00	60 60	✓ <u>Darunavir Mylan</u> ✓ <u>Darunavir Mylan</u>			
LOPINAVIR WITH RITONAVIR – Special Authority see SA2138 Tab 100 mg with ritonavir 25 mg	150.00	etail pharmacy 60	✓ <u>Lopinavir/Ritonavir</u> <u>Mylan</u>			
Tab 200 mg with ritonavir 50 mg	295.00	120	✓ <u>Lopinavir/Ritonavir</u> <u>Mylan</u>			
Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml OP	✓ Kaletra			

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	(Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
RITONAVIR – Special Authority see SA2139 on page 103 – Reta Tab 100 mg		30	✓ N	lorvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR – Special Authority see SA2139 on page 103- Tab 50 mg	, ,	30	✓ T	ïvicay
RALTEGRAVIR POTASSIUM – Special Authority see SA2139 of Tab 400 mg	1,090.00	harm 60 60	✓ ls	sentress sentress HD

Immune Modulators

Guidelines for the use of interferon in the treatment of hepatitis C:

Physicians considering treatment of patients with hepatitis C should discuss cases with a gastroenterologist or an infectious disease physician. All subjects undergoing treatment require careful monitoring for side effects.

Patients should be otherwise fit.

Hepatocellular carcinoma should be excluded by ultrasound examination and alpha-fetoprotein level.

Criteria for Treatment

- 1) Diagnosis
 - Anti-HCV positive on at least two occasions with a positive PCR for HCV-RNA and preferably confirmed by a supplementary RIBA test; or
 - PCR-RNA positive for HCV on at least 2 occasions if antibody negative; or
 - Anti-HCV positive on at least two occasions with a positive supplementary RIBA test with a negative PCR for HCV RNA but with a liver biopsy consistent with 2(b) following.

Exclusion Criteria

- Autoimmune liver disease. (Interferon may exacerbate autoimmune liver disease as well as other autoimmune diseases such as thyroid disease).
- 2) Pregnancy.
- 3) Neutropenia (< 2.0 × 10⁹) and/or thrombocytopenia.
- 4) Continuing alcohol abuse and/or continuing intravenous drug users.

Dosage

The current recommended dosage is 3 million units of interferon alfa-2a or interferon alfa-2b administered subcutaneously 3 times a week for 52 weeks (twelve months)

Exit Criteria

The patient's response to interferon treatment should be reviewed at either three or four months. Interferon treatment should be discontinued in patients who do not show a substantial reduction (50%) in their mean pre-treatment ALT level at this stage.

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

- a) See prescribing guideline above
- b) Note: Pharmac will consider funding ribavirin for the small group of patients who have a clinical need for ribavirin and meet Special Authority criteria. Please contact the Hepatitis C Coordinator at Pharmac on 0800-023-588 option 4.

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	•	Manufacturer

continued...

- 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
- 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
- 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Notes:

- Consider stopping treatment if there is absence of a virological response (defined as at least a 2-log reduction in viral load) following 12 weeks of treatment since this is predictive of treatment failure.
- Consider reducing treatment to 24 weeks if serum HCV RNA level at Week 4 is undetectable by sensitive PCR assay (less than 50IU/ml) AND Baseline serum HCV RNA is less than 400,000IU/ml

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Fither:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
- 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal; and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Either:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2,000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis): and

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer	
---	-----	---------------------	-------------------------------------	--

continued...

- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with anagrelide and busulfan is not clinically appropriate; or
 - Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:
 - 3.2.1 Patient has a myeloproliferative disorder*; and
 - 3.2.2 Fither:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Notes: Indications marked with * are unapproved indications.

- Approved dose is 180 mcg once weekly.
- The recommended dose of Pegylated Interferon alfa-2a is 180 mcg once weekly.
- In patients with renal insufficiency (calculated creatinine clearance less than 50ml/min), Pegylated Interferon-alfa 2a dose should be reduced to 135 mcg once weekly.
- In patients with neutropaenia and thrombocytopaenia, dose should be reduced in accordance with the datasheet guidelines.
- Pegylated Interferon-alfa 2a is not approved for use in children.

Initial application — **(post-allogenic bone marrow transplant)** from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

	-	,	4.	0.04	400	
*	Tab 1 g		4	0.01	100	Hiprex

METHENAMINE (HEXAMINE) HIPPURATE

INFECTIONS - AGENTS FOR SYSTEMIC USE

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	
NITROFURANTOIN				
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓	Nifuran
* Tab 100 mg	37.50	100	✓	Nifuran
* Cap modified-release 100 mg - Up to 15 cap available on a PSO	86.40	100	/	Macrobid
NORFLOXACIN Tab 400 mg - Subsidy by endorsement	245.00	100	/	Arrow-Norfloxacin
Only if prescribed for a patient with an uncomplicated uri with proven resistance to first line agents and the prescri	nary tract infection th			ive to a first line agent or

	Subsidy		Fully	
	(Manufacturer's Price)		Subsidised	
	\$	Per		Manufacturer
Anticholinesterases				
Antichonnesterases				
NEOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	1	Max Health
PYRIDOSTIGMINE BROMIDE				
▲ Tab 60 mg	45.79	100	1	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
DICLOFENAC SODIUM	4.00		,	Distriction of Courts
* Tab EC 25 mg		50		Diclofenac Sandoz
* Tab 50 mg dispersible		20	_	Voltaren D
* Tab EC 50 mg		50		Diclofenac Sandoz
* Tab long-acting 75 mg		100		Voltaren SR
* Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a		5 10		Voltaren Voltaren
* Suppos 25 mg		10		Voltaren
* Suppos 50 mg - Up to 10 supp available on a PSO		10		Voltaren
* Suppos 100 mg*		10		Voltaren
	7.00	10	•	Voltaren
IBUPROFEN	04.40	4 000	,	. "
* Tab 200 mg		1,000		Relieve
* Tab long-acting 800 mg		30		Brufen SR
* Oral liq 20 mg per ml		200 ml		Ethics
	11.29		•	Fenpaed 100 mg per 5 ml
KETODDOFFN				J IIII
KETOPROFEN * Cap long-acting 200 mg	10.07	28	./	Oruvail SR
	12.07	20	•	Oruvali Sh
MEFENAMIC ACID				
* Cap 250 mg		50		
	(9.16)			Ponstan
	0.50	20		
	(5.60)			Ponstan
NAPROXEN				
* Tab 250 mg	32.69	500		Noflam 250
* Tab 500 mg		250	_	Noflam 500
* Tab long-acting 750 mg		28		Naprosyn SR 750
* Tab long-acting 1 g	8.62	28	/	Naprosyn SR 1000
TENOXICAM				
* Tab 20 mg	9.15	100	1	Tilcotil
* Inj 20 mg vial	9.95	1	1	AFT
NSAIDs Other				
CELECOXIB				
Cap 100 mg	3.45	60	1	Celecoxib Pfizer
Cap 200 mg		30		Celebrex
- σαρ 200 mg	3.20	00		Celecoxib Pfizer
(Celebrex Cap 200 mg to be delisted 1 November 2022)	0.20		•	COLOUNID I HEOL
(1000000 1000000 1000000 10000000 1000000				

Subsidy
(Manufacturer's Price) Subsidy

\$ Per

Fully Subsidised Brand or Generic Manufacturer

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% - Special Authority see SA1289 below - Retail

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly.

Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg	8.78	100	Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg		30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	✓ D-Penamine
Tab 250 mg	110.12	100	✓ D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM			
* Tab 70 mg	2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECAL CIFEROL			

Other Treatments

AL ENDOONATE CODULA

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
- 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

inj 3 mg per mi, 10 mi viai	27.53	1	Pamisoi
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	79.95	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA177	79 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or

Subsidy (Manufacturer's P	rice)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

70I FDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

SA2110 below – Retail pharmacy60.00 100 ml OP 🗸 Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in voung adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note): or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg		500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see S	SA1963 below – Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	✓ Benzbromaron AL
			100 S29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 m	cg	6.00	100	Colgout
Colgo	ut to be Principal Supply on 1 September 2022			•
FEBUXOSTAT	- Special Authority see SA2054 below - Retail pharma	су		
Tab 80 mg		.20.00	28	✓ Febuxostat
				multichem
Tab 120 m	g	.20.00	28	✓ Febuxostat
				multichem

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose: or
- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

Note: In chronic renal insufficiency, particularly when the glomerular filtration rate is 30 ml/minute or less, probenecid may not be effective. The efficacy and safety of febuxostat have not been fully evaluated in patients with severe renal impairment (creatinine clearance less than 30 ml/minute). No dosage adjustment of febuxostat is necessary in patients with mild or moderate renal impairment. Optimal treatment with allopurinol in patients with renal impairment is defined as treatment to the creatinine clearance-adjusted dose of allopurinol then, if serum urate remains greater than 0.36 mmol/l, a gradual increase of the dose of allopurinol to 600 mg or the maximum tolerated dose.

PROBENECID

*	Tab 500 mg	66.95	100	✓	Probenecid-AFT
---	------------	-------	-----	---	----------------

Muscle Relaxants

BA	CLOFEN				
*	Tab 10 mg	4.20	100	✓ Pacifen	
	Inj 0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal	
	Subsidised only for use in a programmable pump in patien caused intolerable side effects and the prescription is endo		, ,	ents have been ineffective or	have
	Inj 2 mg per ml, 5 ml ampoule - Subsidy by endorsement	306.82	5	✓ Medsurge	
	Subsidised only for use in a programmable pump in patien caused intolerable side effects and the prescription is endo		, ,	ents have been ineffective or	have

100

Norflex

DANTROLENE Con 05 mg	07.50	100	✓ Dantrium
Cap 25 mg	97.50	100	✓ Dantrium S29®
Cap 50 mg	77.00	100	✓ Dantrium
ORPHENADRINE CITRATE			

117

Subsidy Fully Brand or (Manufacturer's Price) Generic Subsidised Per Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists and Related Agents

AMANTADINE HYDROCHLORIDE ▲ Cap 100 mg	38.24	60	✓ Symmetrel
APOMORPHINE HYDROCHLORIDE			•
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule1	21.84	5	✓ Movapo

BROMOCRIPTINE MESYLATE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking bromocriptine mesylate prior to 1 March 2021 and the sts a record of

prescription is endorsed accordingly. Pharmacists may anno prior dispensing of bromocriptine mesylate.	tate the prescript	ion as endor	sed where there exists a
* Tab 2.5 mg	11 70	30	✓ Parlodel S29
(Parlodel \$29 Tab 2.5 mg to be delisted 1 September 2022)		00	- Turioudi
, ,			
ENTACAPONE A Table 2002 are a	10.04	100	. Comton
▲ Tab 200 mg	18.04	100	✓ <u>Comtan</u>
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg		100	Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	Madopar 62.5
* Cap 100 mg with benserazide 25 mg		100	Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg		100	Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg		100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg		100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	5 51	100	✓ Ramipex
▲ Tab 1 mg		100	✓ Ramipex
RASAGILINE		100	- Hampox
	50.50		4 A B B
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	2.85	84	✓ Ropin
	3.39	100	✓ Mylan S29
▲ Tab 1 mg	3.95	84	✓ Ropin
- 	4.70	100	✓ Mylan S29
▲ Tab 2 mg	*****	84	✓ Ropin
▲ Tab 5 mg		84	✓ Ropin
(Mulan con Tab 1 mg to be delicted 1 December 2022)			- -

(Mylan S29) Tab 1 mg to be delisted 1 December 2022)

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement - Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

*	Tab 5 mg	48.00	100	✓ Eldepryl S29
TO	LCAPONE			
\blacktriangle	Tab 100 mg	152.38	100	✓ Tasmar

	Subsidy		Fully	Brand or	
	(Manufacturer's Price)	S Per	Subsidised	Generic Manufacturer	
Anticholinergics					
BENZATROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 10 inj available on a PSO b) Only on a PSO		60 5		Benztrop Phebra	
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ F	Kemadrin	
Agents for Essential Tremor, Chorea and Rela	ted Disorders				
RILUZOLE – Special Authority see SA1403 below – Retail pha Wastage claimable Tab 50 mg	•	56	√ <u>F</u>	Rilutek	
■ SA1403 Special Authority for Subsidy	se duration of 5 years of the distribution of 5 years of 5 ye	or less; nonths p	and orior to the	initial application; a	
TETRABENAZINE Tab 25 mg	91.10	112	✓ N	Motetis	
Anaesthetics					
Local					
LIDOCAINE [LIGNOCAINE] Gel 2%, tube — Subsidy by endorsement	l administration and the	30 ml e presci 10	ription is e	(ylocaine 2% Jelly ndorsed accordingl nstillagel Lido	

accordingly.

	Subsidy (Manufacturer's Price		Fully	
	(Manufacturer's Price	Per	JISEU	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	1	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	8.75	25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓	Lidocaine-Baxter
			✓	Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓	Lidocaine-Baxter
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -				
Subsidy by endorsement	103.32	10	✓	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical a	administration and th	e prescriptio	n is	endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] — Special Authority see	e SA0906 above – Retail pharn	nacy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE -	Special Authority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

ASPIRIN * Tab dispersible 300 mg – Up to 30 tab available on a PSO	4.50	100	✓ Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diabet accordingly.	ic periphera	ıl neuropathy aı	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream \$29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

	c · · ·			
	Subsidy (Manufacturer's Price) \$) Su Per	Fully bsidised	Brand or Generic Manufacturer
ARACETAMOL				
Tab 500 mg - blister pack		1,000	✓ <u>F</u>	Pacimol
 a) Maximum of 300 tab per prescription; can be wai b) Up to 30 tab available on a PSO c) 	ved by endorsement			
1) Subsidy by endorsement for higher quantiti regular daily dosing for one month or greate annotate the prescription as endorsed when 2) Maximum of 100 tab per dispensing for nor (for non-endorsed patients), then dispense Tab 500 mg - bottle pack — Maximum of 300 tab per	er, and the prescription is re dispensing history sup n-endorsed patients. If q	s annotate ports a lo uantities	ed accord ong-term prescribe	dingly. Pharmacists may condition. ed for more than 100 tabs
prescription; can be waived by endorsement	17.92	1,000	√ <u> </u> <u> </u>	loumed Paracetamol
 Subsidy by endorsement for higher quantities i daily dosing for one month or greater, and the prescription as endorsed where dispensing his Maximum of 100 tab per dispensing for non-en non-endorsed patients), then dispense in repeated 	prescription is annotated tory supports a long-tern dorsed patients. If quar	l accordin n conditio ntities pres	gly. Pha n. scribed fo	armacists may annotate the or more than 100 tabs (for
Oral liq 120 mg per 5 ml		1,000 ml	✓ <u>F</u>	Paracare
1) Maximum of 200 ml per dispensing for non- non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantiti regular daily dosing for one month or greate Pharmacists may annotate the prescription condition.	epeat dispensing not exc es is available for patien er and the prescription is	eeding 20 ts with lor endorse	00 ml per ng term o d or anno	dispensing. conditions who require stated accordingly.
Oral liq 250 mg per 5 ml	6.25	1,000 ml	√ <u>F</u>	Paracare Double Strength
 a) Maximum of 600 ml per prescription; can be wain b) Up to 100 ml available on a PSO c) Not in combination d) 	ved by endorsement			
1) Maximum of 200 ml per dispensing for non- non-endorsed patients), then dispense in re 2) Subsidy by endorsement for higher quantiti regular daily dosing for one month or greate Pharmacists may annotate the prescription condition.	epeat dispensing not exc es is available for patien er and the prescription is	eeding 20 ts with lor endorse	00 ml per ng term o d or anno	dispensing. conditions who require stated accordingly.
Suppos 125 mg Suppos 250 mg Suppos 500 mg	4.18	10 10 50	✓ (Gacet Gacet Gacet
Outstill Associated				
Opioid Analgesics				

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy		Fully Brand or
	(Manufacturer's Price)		Subsidised Generic
	\$	Per	✓ Manufacturer
DIHYDROCODEINE TARTRATE			
Tab long-acting 60 mg	8.60	60	DHC Continus
FENTANYL			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	equency		
Inj 50 mcg per ml, 2 ml ampoule	3.75	10	 Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule	9.41	10	 Boucher and Muir
Patch 12.5 mcg per hour	6.99	5	✓ Fentanyl Sandoz
Patch 25 mcg per hour		5	✓ Fentanyl Sandoz
Patch 50 mcg per hour		5	✓ Fentanyl Sandoz
Patch 75 mcg per hour		5	✓ Fentanyl Sandoz
Patch 100 mcg per hour	18.59	5	✓ Fentanyl Sandoz
METHADONE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	equency		
d) Extemporaneously compounded methadone will only be	reimbursed at the rat	e of th	ne cheapest form available
(methadone powder, not methadone tablets).			
e) For methadone hydrochloride oral liquid refer Standard F			
Tab 5 mg		10	✓ Methatabs
Oral liq 2 mg per ml		200 m	<u></u>
Oral liq 5 mg per ml		200 m	
Oral liq 10 mg per ml		200 m	
Inj 10 mg per ml, 1 ml	61.00	10	✓ AFT
MORPHINE HYDROCHLORIDE			
a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing from	equency		
Oral liq 1 mg per ml	11.98	200 m	
Oral liq 2 mg per ml	16.24	200 m	RA-Morph
Oral liq 5 mg per ml	19.44	200 m	
			✓ RA-Morph

Oral liq 10 mg per ml27.74

✓ Ordine S29

✓ RA-Morph

200 ml

	Subsidy		Fully	
	(Manufacturer's Price)	Per	Subsidised <	Generic Manufacturer
NORTH SUPPLIATE	Ψ	1 61		Ivianulaciurei
MORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre		4.0	,	
Tab immediate-release 10 mg		10		Sevredol Sevredol
Tab immediate-release 20 mg		10		Sevredol
Cap long-acting 10 mg		10		m-Eslon
Cap long-acting 30 mg		10		m-Eslon
Cap long-acting 60 mg		10		m-Eslon
Cap long-acting 100 mg		10		m-Esion
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a PS	506.99	5	•	DBL Morphine
			_	Sulphate
Inj 10 mg per ml, 1 ml ampoule – Up to 5 inj available on a F	PSO5.61	5	/	DBL Morphine
				Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	PSO7.08	5	✓	DBL Morphine
				Sulphate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on a F	PSO7.28	5	1	DBL Morphine
, 31 , 1 ,				Sulphate
OXYCODONE HYDROCHLORIDE				•
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	auanav			
, , , , , , , , , , , , , , , , , , , ,	, ,	00	./	Ovvendene Candan
Tab controlled-release 5 mg		20		Oxycodone Sandoz
Tab controlled-release 10 mg		20		Oxycodone Sandoz
Tab controlled-release 20 mg		20		Oxycodone Sandoz
Tab controlled-release 40 mg		20		Oxycodone Sandoz
Tab controlled-release 80 mg		20		Oxycodone Sandoz
Cap immediate-release 5 mg		20		OxyNorm .
Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 20 mg		20		<u>OxyNorm</u>
Oral liq 5 mg per 5 ml		250 m		<u>OxyNorm</u>
Inj 10 mg per ml, 1 ml ampoule		5		Hameln
Inj 10 mg per ml, 2 ml ampoule		5		<u>Hameln</u>
Inj 50 mg per ml, 1 ml ampoule	22.92	5	/	<u>Hameln</u>
PARACETAMOL WITH CODEINE - Safety medicine; prescriber	may determine dispe	ensing	frequenc	V
* Tab paracetamol 500 mg with codeine phosphate 8 mg	26.51	1,000		Paracetamol +
				Codeine (Relieve)
PETHIDINE HYDROCHLORIDE				, ,
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing fre	an an an			
Tab 50 mg	' '	10	./	PSM
Inj 50 mg per ml, 1 ml ampoule – Up to 5 inj available on a F		_		DBL Pethidine
ing 50 mg per mi, i mi ampoule – Op to 5 mg available on a F	25029.88	5	•	
late of the second of the seco	200 00 70	_		Hydrochloride
Inj 50 mg per ml, 2 ml ampoule – Up to 5 inj available on a F	25U30.72	5	•	DBL Pethidine
				Hydrochloride
TRAMADOL HYDROCHLORIDE				
Tab sustained-release 100 mg		20		Tramal SR 100
Tab sustained-release 150 mg		20		Tramal SR 150
Tab sustained-release 200 mg		20	1	Tramal SR 200
Cap 50 mg	2.80	100	1	Arrow-Tramadol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy	Full	
	(Manufacturer's Price) \$	Subsidised Per 🗸	
	Ψ	101	Marialactarer
Antidepressants			
Cyclic and Related Agents			
AMITRIPTYLINE - Safety medicine; prescriber may determine of	dispensing frequency		
Tab 10 mg	2.49		Arrow-Amitriptyline
Tab 25 mg			Arrow-Amitriptyline
Tab 50 mg			Arrow-Amitriptyline
CLOMIPRAMINE HYDROCHLORIDE – Safety medicine; presci			
Tab 10 mg			Clomipramine Teva
Tab 25 mg		30	Clomipramine Teva
DOSULEPIN [DOTHIEPIN] HYDROCHLORIDE – Subsidy by er			
Safety medicine; prescriber may determine dispensing from the Subsidiary for patients who are a subsidiary for patients. Comparison of the comparison of t		[dathianin] hud	rachlarida prior to 1 luna
 Subsidy by endorsement – Subsidised for patients who v 2019 and the prescription is endorsed accordingly. Phar 			
exists a record of prior dispensing of dosulepin [dothiepir		and prosoripul	on as chasissa where there
Tab 75 mg		30	Dosulepin Mylan
Cap 25 mg	7.83	50	Dosulepin
			Mylan S29
IMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine dispe	nsing frequenc	y
Tab 10 mg	5.48	50	Tofranil
	10.96		Tofranil
Tab 25 mg			Tofranil
NORTRIPTYLINE HYDROCHLORIDE – Safety medicine; presc			
Tab 10 mg			Norpress
Tab 25 mg	5.98	180	Norpress
Monoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective		
TRANYLCYPROMINE SULPHATE			
Tab 10 mg	12.85	28	Parnate S29 S29
	22.94	50	Parnate
	45.88		Parnate S29 S29
	96.00	•	Parnate S29 S29
Monoamine-Oxidase Type A Inhibitors			
MOCLOBEMIDE			
* Tab 150 mg			<u>Aurorix</u>
* Tab 300 mg	19.25	60	<u>Aurorix</u>
Selective Serotonin Reuptake Inhibitors			
CITALOPRAM HYDROBROMIDE			
* Tab 20 mg	1.91	84	PSM Citalopram
ESCITALOPRAM			
* Tab 10 mg	1.07	28	Escitalopram
· ·		-	(Ethics)
* Tab 20 mg	1.92	28	Escitalopram
			(Ethics)

		Subsidy		illy Brand or
		(Manufacturer's Price)	Subsidis Per	
		\$	rei	✓ Manufacturer
	E HYDROCHLORIDE	4.04	00	/ Fluor
₭ Tab dispe	ersible 20 mg, scored - Subsidy by endorsement	1.98		✓ Fluox ✓ Fluox
Subs	sidised by endorsement	1.50	30	FIUUX
	When prescribed for a patient who cannot swallow	whole tablets or can	sules and the	nrescription is endorsed
',	accordingly; or	wildle lablete of dap	ouloo una mo	prosonption is chasised
2)	When prescribed in a daily dose that is not a multi	ple of 20 mg in which	case the pres	scription is deemed to be
	endorsed. Note: Tablets should be combined with			
Cap 20 m	ng	2.91	84	Fluox
PAROXETINE	Ē			
★ Tab 20 m	ng	3.61	90	Loxamine
SERTRALINE				
★ Tab 50 m	ng	0.92		✓ Setrona
k T-1-100		4.04		Setrona AU
₭ Tab 100 i	mg	1.61		✓ Setrona ✓ Setrona AU
				Selfolia AU
Other An	tidepressants			
	·			
/IRTAZAPIN		0.60	00	./ Naumad
	ng			✓ <u>Noumed</u> ✓ Noumed
	•		20	Noullieu
/ENLAFAXIN	N⊑ ⊢mg	6 20	84	✓ Enlafax XR
	ng			✓ Enlafax XR
	mg			✓ Enlafax XR
	ů .	•		
Antiepile	psy Drugs			
Agents fo	or Control of Status Epilepticus			
NAZEPAM -	- Safety medicine; prescriber may determine disper	nsing frequency		
	per ml, 2 ml ampoule - Subsidy by endorsement		5	✓ Hospira
	lp to 5 inj available on a PSO			r
,	Only on a PSO			
c) P	SO must be endorsed "not for anaesthetic procedu	ıres".		
Rectal tul	bes 5 mg - Up to 5 tube available on a PSO	43.50	5	✓ Stesolid
PHENYTOIN	SODIUM			
₭ Inj 50 mg	per ml, 2 ml ampoule - Up to 5 inj available on a			
PSO		104.58	5	✓ Hospira
★ Inj 50 mg	per ml, 5 ml ampoule - Up to 5 inj available on a			
PSO		154.01	5	✓ Hospira
Control o	f Epilepsy			
CARBAMAZE	PINE			
	mg	14.53		✓ Tegretol
	-acting 200 mg	16.98		✓ Tegretol CR
* Tab long-		04.50	100	/ Taggatal
* Tab 400 i	mg			✓ Tegretol
* Tab 400 i * Tab long-	mg -acting 400 mg 0 mg per ml	39.17	100	✓ Tegretol CR ✓ Tegretol

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pri		Fully Brand or sidised Generic	
	<u></u>	Per	✓ Manufacturer	
CLOBAZAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Tab 10 mg	9.12	50	✓ Frisium	
CLONAZEPAM - Safety medicine; prescriber may determine di	spensing frequenc	V		
Oral drops 2.5 mg per ml		10 ml OP	✓ Rivotril	
ETHOSUXIMIDE				
Cap 250 mg	140 88	100	✓ Zarontin	
Oral liq 250 mg per 5 ml		200 ml	✓ Zarontin	
GABAPENTIN				
Note: Not subsidised in combination with subsidised pregat	alin			
* Cap 100 mg		100	✓ Nupentin	
* Cap 300 mg		100	✓ Nupentin	
* Cap 400 mg		100	✓ Nupentin	
LACOSAMIDE – Special Authority see SA1125 below – Retail p Tab 50 mg		14	✓ Vimpat	
▲ Tab 100 mg		14	✓ Vimpat ✓ Vimpat	
Tab 100 flig	200.24	56	✓ Vimpat ✓ Vimpat	
▲ Tab 150 mg		14	✓ Vimpat ✓ Vimpat	
Tab 150 flig	300.40	56	✓ Vimpat ✓ Vimpat	
▲ Tab 200 mg		56	✓ Vimpat ✓ Vimpat	
<u> </u>		50	• viiiipat	

⇒SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see Note).

Note: "Optimal treatment" is defined as treatment which is indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight and other features affecting the pharmacokinetics of the drug with good evidence of compliance. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment (see Note)

Note: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

\blacktriangle	Tab dispersible 2 mg	55.00	30	✓ Lamictal
\blacktriangle	Tab dispersible 5 mg	50.00	30	✓ Lamictal
*	Tab dispersible 25 mg		56	✓ Logem
*	Tab dispersible 50 mg		56	✓ Logem
*	Tab dispersible 100 mg	4.40	56	✓ Logem
LE	VETIRACETAM			-
	Tab 250 mg	4.99	60	✓ Everet
	Tab 500 mg		60	✓ Everet
	Tab 750 mg		60	✓ Everet
	Tab 1,000 mg		60	✓ Everet
	Oral liq 100 mg per ml	44.78	300 ml OP	✓ Levetiracetam-AFT
РΗ	ENOBARBITONE			
	For phenobarbitone oral liquid refer Standard Formulae,	page 243		
*	Tab 15 mg	40.00	500	✓ PSM
*	Tab 30 mg	40.00	500	✓ PSM

	Subsidy		Fully	
	(Manufacturer's Pric		Subsidised	
	\$	Per		Manufacturer
PHENYTOIN SODIUM				
* Tab 50 mg	75.00	200	1	Dilantin Infatab
Cap 30 mg	74.00	200	✓	Dilantin
Cap 100 mg	37.00	200	1	Dilantin
* Oral liq 30 mg per 5 ml	22.03	500 ml	1	Dilantin
PREGABALIN				
Note: Not subsidised in combination with subsidised gabage	entin			
Cap 25 mg	2.25	56	1	Pregabalin Pfizer
* Cap 75 mg		56		Pregabalin Pfizer
Cap 150 mg		56	1	Lyrica
			1	Pregabalin Pfizer
Cap 300 mg	7.38	56		Pregabalin Pfizer
PRIMIDONE				· ·
* Tab 250 mg	37.35	100	1	Apo-Primidone
1 22 250 mg				Primidone Clinect
(Apo-Primidone Tab 250 mg to be delisted 1 January 2023)				
SODIUM VALPROATE				
Tab 100 mg	13.65	100	1	Epilim Crushable
Tab 200 mg EC		100		Epilim
Tab 500 mg EC		100		Epilim
* Oral lig 200 mg per 5 ml		300 ml	1	Epilim S/F Liquid
			1	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	✓	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail p				•
Cap 250 mg	•	60	1	Diacomit \$29
Powder for oral lig 250 mg sachet		60		Diacomit S29
T OWNOUT TOT OTAL THY 200 THY SAUTIEL		00	•	Diacollilit

⇒SA1330 Special Authority for Subsidy

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

IOPIRAMATE A Tob 25 mg 11.07	60	✓ Arrow-Topiramate
▲ Tab 25 mg11.07	00	✓ Topiramate Actavis
26.04		✓ Topamax
▲ Tab 50 mg18.81	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
44.26		✓ Topamax
▲ Tab 100 mg31.99	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
75.25		✓ Topamax
▲ Tab 200 mg55.19	60	✓ Arrow-Topiramate
·		✓ Topiramate Actavis
129.85		✓ Topamax
▲ Sprinkle cap 15 mg	60	✓ Topamax
▲ Sprinkle cap 25 mg	60	✓ Topamax

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
	\$	Per	✓	Manufacturer	
VIGABATRIN – Special Authority see SA2088 below – Retail ph.	•	100	✓ Si	السمام	
▲ Tab 500 mg	119.30	100	♥ 58	aDrii	

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and
 - 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
 - 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter); or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: ``Optimal treatment with other antiepilepsy agents" is defined as treatment with other antiepilepsy agents which are indicated and clinically appropriate for the patient, given in adequate doses for the patient's age, weight, and other features affecting the pharmacokinetics of the drug with good evidence of compliance.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Either:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Notes: As a guideline, clinical trials have referred to a notional 50% reduction in seizure frequency as an indicator of success with anticonvulsant therapy and have assessed quality of life from the patient's perspective.

Vigabatrin is associated with a risk of irreversible visual field defects, which may be asymptomatic in the early stages.

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Acute Migraine	Treatment
-----------------------	-----------

RIZATRIPTAN			
Tab orodispersible 10 mg	3.65	30	✓ <u>Rizamelt</u>
SUMATRIPTAN			
Tab 50 mg	14.41	90	✓ Sumagran
Tab 100 mg	22.68	90	✓ Sumagran
Inj 12 mg per ml, 0.5 ml prefilled pen - Maximum of 10 inj per			
prescription	34.00	2 OP	Imigran
			_

Sub	bsidy	Fully	Brand or
(Manufactu	urer's Price) Subsic	lised	Generic
•	\$ Per	1	Manufacturer

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 50

PIZOTIFEN

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

* Tab 16 mg	4.62	100	✓ <u>Serc</u>
CYCLIZINE HYDROCHLORIDE Tab 50 mg	0.49	10	✓ Nausicalm
CYCLIZINE LACTATE			
Inj 50 mg per ml, 1 ml ampoule	16.36	10	✓ Hameln
DOMPERIDONE			_
* Tab 10 mg	2.85	100	✓ Pharmacy Health
HYOSCINE HYDROBROMIDE			
* Inj 400 mcg per ml, 1 ml ampoule		10	✓ Martindale S29
Patch 1.5 mg - Special Authority see SA1998 below		_	
pharmacy	14.11	2	✓ Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

METOCLOPRAMIDE HYDROCHLORIDE

*	Tab 10 mg - Up to 30 tab available on a PSO	1.30	100	✓ <u>Metoclopramide</u>
				Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSO	7.00	10	✓ Baxter
	, , ,	9.50		✓ Pfizer
(Pt	izer Ini 5 ma per ml. 2 ml ampoule to be delisted 1 December 2022)			

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
NDANSETRON	<u> </u>			
Tab 4 mg	2.68	50	1	Onrex
Tab disp 4 mg - Up to 10 tab available on a PSO		10		Ondansetron
Tab disp Ting - Sp to To tab available on a T SS				ODT-DRLA
Tab 8 mg	4 57	50	1	Onrex
Tab disp 8 mg - Up to 10 tab available on a PSO		10		Ondansetron
Tab disp of fig — op to 10 tab available off a 1 30	1.10	10	•	ODT-DRLA
OCHLORPERAZINE				
Tab 3 mg buccal	5.97	50		
. as 0g 2000a	(30.00)			Buccastem
Tab 5 mg - Up to 30 tab available on a PSO	(/	250	1	Nausafix
Inj 12.5 mg per ml, 1 ml – Up to 5 inj available on a PSO		10		Stemetil
and it is a second point of the configuration of th		. •		••••
ntipsychotics				
General				
MISULPRIDE - Safety medicine; prescriber may determine of	lienancina fraguency			
Tab 100 mg		30	J	Sulprix
rab roomly	17.16	100	_	Amisulpride
	17.10	100	•	Mylan S29
Tab 200 mg	14.06	60	./	Sulprix
Tab 200 mg Tab 400 mg		60		Sulprix
IIPIPRAZOLE – Safety medicine; prescriber may determine Tab 5 mg		30	1	Aripiprazole Sandoz
Tab 10 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 15 mg	10.50	30	1	Aripiprazole Sandoz
Tab 20 mg	10.50	30	✓	Aripiprazole Sandoz
Tab 30 mg	10.50	30	1	Aripiprazole Sandoz
ILORPROMAZINE HYDROCHLORIDE - Safety medicine; p	orescriber may determi	ne di	spensing fr	requency
Tab 10 mg - Up to 30 tab available on a PSO	14.83	100		Largactil
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	1	Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	1	Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO	30.79	10	1	Largactil
OZAPINE - Hospital pharmacy [HP4]				
Safety medicine; prescriber may determine dispensing frequency	quency			
Tab 25 mg	. ,	50	/	Clopine
•			1	Clozaril
	13.37	100	1	Clopine
				Clozaril
Tab 50 mg	8.67	50	/	Clopine
•	17.33	100		Clopine
Tab 100 mg	17.33	50		Clopine
•				Clozaril
	34.65	100		Clopine
				Clozaril
Tab 200 mg	34.65	50		Clopine
Ÿ	69.30	100		Clopine

Suspension 50 mg per ml......67.62

✓ Versacloz

100 ml

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Per	Subsidised <	Generic Manufacturer
ALOPERIDOL – Safety medicine; prescriber may determine di	isnensing frequency			
Tab 500 mcg – Up to 30 tab available on a PSO		100	1	Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100		Serenace
Tab 5 mg - Up to 30 tab available on a PSO		50		Serenace
Tab 3 fing Op to 30 tab available off a 1 00	29.72	100		Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 m		Serenace
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		10		Serenace
EVOMEPROMAZINE - Safety medicine; prescriber may deter		allanev		
Tab 25 mg (33.8 mg as a maleate)	, ,	100	_	Nozinan (Swiss)
Tab 25 mg as a maleate		100		Nozinan (Swiss)
Tab 100 mg (135 mg as a maleate)		100		Nozinan (Swiss)
Tab 100 mg as a maleate		100		Nozinan (Owiss)
EVOMEPROMAZINE HYDROCHLORIDE – Safety medicine;	-			
Inj 25 mg per ml, 1 ml ampoule		5		Neuraxpharm S29
	33.50	10	•	Nozinan
ITHIUM CARBONATE - Safety medicine; prescriber may dete	rmine dispensing fre	quency		
Tab long-acting 400 mg	72.00	100		<u>Priadel</u>
Cap 250 mg	9.42	100	/	Douglas
LANZAPINE - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 2.5 mg	1.35	28	1	Zypine
Tab 5 mg	1.58	28	_	Zypine
Tab orodispersible 5 mg	1.81	28	1	Zypine ODT
Tab 10 mg	2.01	28	_	Zypine
Tab orodispersible 10 mg		28		Zypine ODT
ERICYAZINE - Safety medicine; prescriber may determine dis	spensing frequency			
Tab 2.5 mg		84	/	Neulactil
· · · · · · · · · · · · · · · · · · ·	12.49	100		Neulactil
Tab 10 mg		84		Neulactil
	44.45	100		Neulactil
NUETIAPINE - Safety medicine; prescriber may determine disp	nensina frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90		Quetapel
ISPERIDONE - Safety medicine; prescriber may determine di				
Tab 0.5 mg		60	1	Risperidone (Teva)
Tab 1 mg		60	_	Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral liq 1 mg per ml		30 ml		Risperon
, ,,		00 1111	•	эрстоп
IPRASIDONE – Safety medicine; prescriber may determine dis		60		Zuadana
Cap 20 mg		60		Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
	4h 55	60	•	Zusdone
Cap 80 mgUCLOPENTHIXOL HYDROCHLORIDE - Safety medicine; pre				



	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
Depot Injections			

FLUPENTHIXOL DECANOATE – Safety medicine; prescriber may	determine disp	pensing frequ	iency
Inj 20 mg per ml, 1 ml - Up to 5 inj available on a PSO	13.14	5	✓ Fluanxol
Inj 20 mg per ml, 2 ml - Up to 5 inj available on a PSO	20.90	5	✓ Fluanxol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may	determine dispe	ensing freque	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO	28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml - Up to 5 inj available on a PSO	55.90	5	✓ Haldol Concentrate
			✓ Haldol
			Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail phar	rmacy		
Safety medicine; prescriber may determine dispensing frequen	су		
Inj 210 mg vial	252.00	1	Zyprexa Relprevv
Inj 300 mg vial	414.00	1	✓ Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and
 - 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: The patient should be monitored for post-injection syndrome for at least two hours after each injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Ini 405 mg vial504.00

Safety medicine; prescriber may determine dispens	sing frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe		1	✓ Invega Sustenna
Inj 75 mg syringe	357.42	1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Fither:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the

continued...

✓ Zyprexa Relprevy

Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

initiation of an atypical antipsychotic depot injection.

Note: Paliperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialling paliperidone depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Safety medicine; prescriber may determine dispensin	ng frequency		
Inj 25 mg vial	135.98	1	Risperdal Consta
Inj 37.5 mg vial	178.71	1	✓ Risperdal Consta
Inj 50 mg vial	217.56	1	 Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

Note: Risperidone depot injection should ideally be used as monotherapy (i.e. without concurrent use of any other antipsychotic medication). In some cases, it may be clinically appropriate to attempt to treat a patient with typical antipsychotic agents in depot injectable form before trialing risperidone depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine: prescriber may determine dispensing frequency

Inj 200 mg per ml, 1 ml − Up to 5 inj available on a PSO.......19.80 5 ✓ Clopixol

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	18.50	100	 Buspirone Viatris
* Tab 10 mg	12.50	100	 Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may de	termine dispensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determ	nine dispensing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may dete	ermine dispensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ Ativan

Subsidy (Manufacturer's Price) Fully Subsidised

Per

Brand or Generic Manufacturer

Multiple Sclerosis Treatments

⇒SA2140 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist; and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Either:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia, Lhermitte's symptom); and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion; or
 - 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or
 - 6.4 A sign of that new inflammatory activity is a prominent T2 lesion that clearly is responsible for the clinical features of a recent attack that occurred within the last 2 years; or
 - 6.5 A sign of that new inflammatory activity is new T2 lesions compared with a previous MRI scan.

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Renewal — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months where patient has had an EDSS course of 0 to 6.0 (inclusive) with or without the use of unilstoned or billstoned and the court time in the lost six months.

had an EDSS score of 0 to 6.0 (inclusive) with or without the use of unilateral or bilateral aids at any time in the last six months (i.e. the patient has walked 100 metres or more with or without aids in the last six months).

Note: Natalizumab can only be dispensed from a pharmacy registered in the Tysabri Australasian Prescribing Programme operated by the supplier. Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

DIMETHYL FUMARATE - Special Authority see SA2140 above - Retail pharmacy

- a) Wastage claimable
- b) Note: Treatment on two or more funded multiple sclerosis treatments simultaneously is not permitted.

Cap 120 mg) 14	Tecfidera
Cap 240 mg2,000.00	56	Tecfidera

Brand or

Fully

	Subsidy		rully	Diana or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	✓	Manufacturer
FINGOLIMOD - Special Authority see SA2140 on the previous	page – Retail pharma	су		
a) Wastage claimable				
 Note: Treatment on two or more funded multiple scleros 	sis treatments simultar	eousl	y is not per	mitted.
Cap 0.5 mg	2,200.00	28	✓ (Silenya
GLATIRAMER ACETATE - Special Authority see SA2140 on the	ne previous page – Re	tail ph	armacv	
Note: Treatment on two or more funded multiple sclerosis to				ed.
Inj 40 mg prefilled syringe		12		Copaxone
INTERFERON BETA-1-ALPHA - Special Authority see SA2140				
Note: Treatment on two or more funded multiple sclerosis to		•		
Inj 6 million iu prefilled syringe		4	-	vonex
Injection 6 million iu per 0.5 ml pen injector	1,170.00	4	✓	vonex Pen
INTERFERON BETA-1-BETA - Special Authority see SA2140	on the previous page -	- Reta	il pharmacy	/
Note: Treatment on two or more funded multiple sclerosis to				
Inj 8 million iu per 1 ml		15		Betaferon
·	*		_	
NATALIZUMAB – Special Authority see SA2140 on the previous		•		1
Note: Treatment on two or more funded multiple sclerosis to		isiy is		
Inj 20 mg per ml, 15 ml vial	1,750.00	1	✓ 1	ysabri
OCRELIZUMAB - Special Authority see SA2140 on the previous	is page – Retail pharm	ıacy		
Note: Treatment on two or more funded multiple sclerosis to	reatments simultaneou	ısly is	not permitt	ed.
Inj 30 mg per ml, 10 ml vial		í		Ocrevus
TERIFLUNOMIDE – Special Authority see SA2140 on the previ		rmacı	,	
	ous page – Hetali pila	illacy	!	
a) Wastage claimable				
b) Note: Treatment on two or more funded multiple scleros			• •	
Tab 14 mg	659.90	28	✓ <u>F</u>	<u>lubagio</u>

Subeidy

Sedatives and Hypnotics

MELATONIN - Special Authority see SA1666 below - Retail pharmacy
Tab modified-release 2 mg - No more than 5 tab per day11.50 30 ✓ Vigisom

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

Note: Indications marked with * are unapproved indications.

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	Generic
	\$	Per	· •	Manufacturer
MIDAZOLAM - Safety medicine; prescriber may determine dispe	ensing frequency			
Inj 1 mg per ml, 5 ml ampoule	5.50	10	✓	Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available)			
on a PSO	17.28	10	✓	Pfizer
On a PSO for status epilepticus use only. PSO must be	endorsed for status e	epiler	oticus use	only.
Inj 5 mg per ml, 3 ml ampoule		5		Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available				
a PSO		5	✓	Pfizer
On a PSO for status epilepticus use only. PSO must be		epilep	oticus use	only.
PHENOBARBITONE SODIUM - Special Authority see SA1386	pelow - Retail pharm	acv .		·
Inj 200 mg per ml, 1 ml ampoule	•	10	1	Max Health S29
⇒SA1386 Special Authority for Subsidy				
Initial application from any relevant practitioner. Approvals valid	d without further rene	wali	ınless notif	ied for applications meeting
the following criteria:	a without further forte	ware	1111000 110111	ica for applications meeting
Both:				
1 For the treatment of terminal agitation that is unresponsive	to other agents: and	4		
2 The applicant is part of a multidisciplinary team working in	0 /	,		
	•			
TEMAZEPAM – Safety medicine; prescriber may determine disp	. ,			
Tab 10 mg	1.33	25	•	<u>Normison</u>
TRIAZOLAM - Safety medicine; prescriber may determine dispe	nsing frequency			
Tab 125 mcg	5.10	100		
	(9.85)			Hypam

Tab 250 mcg......4.10

ZOPICLONE - Safety medicine; prescriber may determine dispensing frequency

Tab 7.5 mg10.80

100

500

(11.20)

Hypam

✓ Zopiclone Actavis

	Subsidy		Fully Brand or
	(Manufacturer's Price) \$	Per	Subsidised Generic Manufacturer
Stimulants/ADHD Treatments			
ATOMOXETINE			
Cap 10 mg	18.41	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			S29 S29
			✓ Generic Partners
	107.03		✓ Strattera
Cap 18 mg	27.06	28	✓ APO-Atomoxetine
			✓ Generic Partners
	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ APO-Atomoxetine
			✓ Generic Partners
Cap 40 mg	29.22	28	✓ APO-Atomoxetine
			✓ Generic Partners
000	107.03		✓ Strattera
Cap 60 mg	46.51	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			S29 S29
			Generic Partners
Cap 80 mg	56.45	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine S29 S29
			Generic Partners
Cap 100 mg	58.48	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			S29 S29
			✓ Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see SA1149 b	nelow – Retail nharma	CV	
a) Only on a controlled drug form	Tiolaii pilaiilia	Оу	
b) Safety medicine; prescriber may determine dispensing free	earrency		
Tab 5 mg	' '	100	✓ PSM
1 ab o mg		100	* <u>I OM</u>

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and



Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

a) Only on a controlled drug form

 b) Safety medicine; prescriber may determine dispensi 	ng frequency		
Tab immediate-release 5 mg	3.20	30	✓ Rubifen
Tab immediate-release 10 mg		30	✓ Ritalin
			✓ Rubifen
Tab extended-release 18 mg	7.75	30	 Methylphenidate ER
			- Teva
Tab immediate-release 20 mg		30	✓ Rubifen
Tab sustained-release 20 mg	10.95	30	✓ Rubifen SR
Tab extended-release 27 mg		30	Methylphenidate ERTeva
Tab extended-release 36 mg	15.50	30	Methylphenidate ERTeva
Tab extended-release 54 mg	22.25	30	Methylphenidate ERTeva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria: and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
	Per	1	Manufacturer	

continued...

2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg		30	Concerta
Tab extended-release 36 mg		30	Concerta
Tab extended-release 54 mg	86.24	30	Concerta
Cap modified-release 10 mg	15.60	30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Fither:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months

Subsidy		ully Brand or	
(Manufacturer's \$	Price) Subsidis Per	sed Generic Manufacturer	

continued...

for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Fither:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below	- Retail pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	 Rivastigmine Patch
			<u>BNM 5</u>
Patch 9.5 mg per 24 hour	38.00	30	 Rivastigmine Patch
			BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

28

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 8 mg with naloxone 2 mg34.00

✓ Buprenorphine Naloxone BNM

✓ Buprenorphine
Naloxone BNM

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

BUPROPION HYDROCHLORIDE

Tab modified-release 150 mg......11.00 30 ✓ Zyban

	Subsidy (Manufacturer's Price)		Fully Subsidised		
	\$	Per	✓	Manufacturer	
DISULFIRAM					
Tab 200 mg	236.40	100	✓	Antabuse S29	
NALTREXONE HYDROCHLORIDE - Special	I Authority see SA1408 below – Retail p	harm	асу		
Tab 50 mg	133.33	30	•	Naltraccord	

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the provisions in P	art I of Sect	ion A.
Patch 7 mg - Up to 28 patch available on a PSO18.14	28	1
Patch 7 mg for direct distribution only – [Xpharm] 3.94	7	1

Tator 7 mg for direct distribution only [Aprilaning	,	• Habition
Patch 14 mg - Up to 28 patch available on a PSO19.95	28	Habitrol
Patch 14 mg for direct distribution only – [Xpharm]4.52	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO22.86	28	Habitrol
Patch 21 mg for direct distribution only – [Xpharm]5.18	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]3.20	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO21.02	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]3.24	36	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]8.64	96	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO44.17	384	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.

Gum 4 mg (Mint) - Up to 384 piece available on a PSO......44.17

Gum 4 mg (Mint) for direct distribution only - [Xpharm]......10.01

c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

96

Habitrol

Habitrol

✓ Habitrol✓ Habitrol

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHLORIDE - PCT only - Specialist - Special Authority see \$A2046 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2046 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6; and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Either:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 All of the following:
 - 3.2.1 Patient has relapsed refractory disease following prior chemotherapy; and
 - 3.2.2 The patient has not received prior bendamustine therapy; and
 - 3.2.3 Fither:
 - 3.2.3.1 Both:
 - 3.2.3.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.2.3.1.2 Patient has had a rituximab treatment-free interval of 12 months or more: or
 - 3.2.3.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

Both:

- 1 Patients have not received a bendamustine regimen within the last 12 months; and
- 2 Either:
 - 2.1 Both:

(Ma	Subsidy anufacturer's Price)	Sub	Fully	Brand or Generic
	\$	Per	•	Manufacturer

continued...

- 2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
- 2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
- 2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has Hodgkin's lymphoma requiring treatment; and
- 2 Patient has a ECOG performance status of 0-2; and
- 3 Patient has received one prior line of chemotherapy; and
- 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
- 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m² twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

BUSULFAN - PCT - Retail pharmacy-Specialist

BUSULFAN - PUT - Retail pharmacy-Specialist			
Tab 2 mg	89.25	100	✓ Myleran
CARBOPLATIN - PCT only - Specialist			
Inj 10 mg per ml, 45 ml vial	32.59	1	 DBL Carboplatin
, , ,	45.20		✓ Carboplatin Ebewe
	48.50		✓ Carbaccord
Inj 1 mg for ECP	0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist			
Inj 100 mg vial	710.00	1	✓ BiCNU
	1,387.00		✓ Bicnu Heritage S29
BiCNU to be Principal Supply on 1 September 2022			· ·
Inj 100 mg for ECP	710.00	100 mg OP	✓ Baxter
(Bicnu Heritage S29 Inj 100 mg vial to be delisted 1 September	2022)		
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist			
Tab 2 mg	29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist			
Inj 1 mg per ml, 50 ml vial	15.00	1	 Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial		1	✓ Cisplatin Ebewe
, .,	29.66		✓ DBL Cisplatin
Inj 1 mg for ECP	0.31	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE			
Tab 50 mg - PCT - Retail pharmacy-Specialist	145.00	50	✓ Cyclonex
Inj 1 g vial - PCT - Retail pharmacy-Specialist	35.65	1	✓ Endoxan
	127.80	6	✓ Cytoxan
Inj 2 g vial - PCT only - Specialist	71.25	1	✓ Endoxan
Inj 1 mg for ECP - PCT only - Specialist	0.04	1 mg	✓ Baxter
IFOSFAMIDE - PCT only - Specialist			
Inj 1 g	96.00	1	✓ Holoxan
lnj 2 g		1	✓ Holoxan
Inj 1 mg for ECP		1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OMLISTINE DCT Patail pharmacy Specialist	<u> </u>	1 01		Managadio
LOMUSTINE – PCT – Retail pharmacy-Specialist	122.50	20	1	CeeNU
Cap 40 mg		20		CeeNU
Cap 40 mg		20	•	CECINO
MELPHALAN			_	
Tab 2 mg - PCT - Retail pharmacy-Specialist		25		Alkeran
Inj 50 mg - PCT only - Specialist	67.80	1	✓	Alkeran
			✓	Alkeran S29 S29
DXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	•	Oxaliplatin Actavis
	110.00		1	Oxaliplatin Ebewe
Ini 5 ma nor ml. 20 ml vial		1		Oxaliplatin Accord
Inj 5 mg per ml, 20 ml vial Inj 1 mg for ECP		1 mg	_	Baxter
	0.40	1 1119	,	Daxiei
THIOTEPA - PCT only - Specialist				
Inj 15 mg vial	CBS	1	✓	Bedford S29
			✓	Max Health S29
			1	THIO-TEPA \$29
			1	Tepadina S29
Inj 100 mg vial	CBS	1		Max Health S29
iiij 100 iiig viai		'	_	
			•	Tepadina S29

Antimetabolites

AZACITIDINE - PCT only - Specialist - Special Authority see SA214	41 below		
Inj 100 mg vial	75.06	1	✓ Azacitidine Dr
			Reddy's
Inj 1 mg for ECP	0.83	1 mg	✓ Baxter

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or
	(Manufacturer's Price		ubsidised Generic
	\$	Per	✓ Manufacturer
CALCIUM FOLINATE			
Tab 15 mg - PCT - Retail pharmacy-Specialist	114.69	10	✓ DBL Leucovorin
		_	Calcium
Inj 3 mg per ml, 1 ml – PCT – Retail pharmacy-Specialist		5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali	st 7.28	1	✓ Calcium Folinate
			Sandoz
			✓ Calcium Folinate
Life DOT Division On the	70.00	40	Sandoz S29 S29
Inj 50 mg - PCT - Retail pharmacy-Specialist	/2.80	10	Leucovorin
			Pharmacia S29
Inj 10 mg per ml, 10 ml vial – PCT only – Specialist	9.49	1	✓ Calcium Folinate
Let 400 mm - BOT code - Operate list	7.00		Sandoz
Inj 100 mg - PCT only - Specialist	7.33	1	✓ Calcium Folinate Ebewe
	94.90	10	✓ Leucovorin
	94.90	10	
Let 000 mm BOT color Operatellist	00.54		Pharmacia S29
Inj 300 mg - PCT only - Specialist	22.51	1	✓ Calcium Folinate Ebewe
	25.14		✓ Leucovorin DBL ©29
	23.14		Leucovoriii DBL 329
Inj 10 mg per ml, 35 ml vial - PCT only - Specialist	25.14	1	✓ Calcium Folinate
			Sandoz
			Calcium Folinate
			Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67.51	1	Calcium Folinate
			Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	Calcium Folinate
			Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter
CAPECITABINE - Retail pharmacy-Specialist			
Tab 150 mg		60	✓ Capercit
Tab 500 mg	49.00	120	Capercit
CLADRIBINE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml		1	✓ Litak S29
Inj 1 mg per ml, 10 ml		1	Leustatin
Inj 10 mg for ECP	749.96	10 mg OF	○ ✓ Baxter
CYTARABINE			
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali	st400.00	5	✓ Pfizer
Inj 100 mg per ml, 20 ml vial – PCT – Retail			_
pharmacy-Specialist		. 1	✓ Pfizer
Inj 1 mg for ECP – PCT only – Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP – PCT only – Speciali	si80.00	100 mg O	P ✓ Baxter
FLUDARABINE PHOSPHATE			4 - 1 - 1
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5 50 mg 05	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	115.29	50 mg OF	○ ✓ Baxter

	Subsidy (Manufacturer's Pri	ice) Subs	Fully	
	\$	Per	√	Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	1	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
lnj 1 g	15.89	1	1	Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	1	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
, , ,	71.44		•	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE		•		
Tab 50 mg - PCT - Retail pharmacy-Specialist	25.90	25	1	Puri-nethol
Oral suspension 20 mg per ml – Retail pharmacy-Specialist		-		
Special Authority see SA1725 below		100 ml OP	✓	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ <u>Trexate</u>
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ <u>Trexate</u>
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*	Inj 7.5 mg prefilled syringe14.61	1	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe	1	✓ Methotrexate Sandoz
*	Inj 15 mg prefilled syringe	1	✓ Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	1	✓ Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	1	✓ Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	1	✓ Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	✓ Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg OP	✓ Baxter

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	•	Manufacturer
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharm.	acy-Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy Manufacturer's Price \$	e) Subs	sidised Ger	nd or neric nufacturer
BLEOMYCIN SULPHATE - PCT only - Specialist Inj 15,000 iu, vial	185.16	1	✓ DBL E	Bleomycin ate
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ Baxte	r
BORTEZOMIB – PCT only – Specialist – Special Authority see SA Inj 2.5 mg vial		1	✓ Borte:	zomib Juno
Inj 3.5 mg vial	105.00	1	✓ Borte: Dr-F ✓ Borte:	dy's S29 S29 zomib leddy's
Inj 1 mg for ECP	2022) August 2022)	1 mg	✓ Baxte	r

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

Note: Indications marked with are unapproved indications.		
DACARBAZINE - PCT only - Specialist		
Inj 200 mg vial62.7	0 1	✓ DBL Dacarbazine
580.6		✓ Dacarbazine
		APP S29
Inj 200 mg for ECP62.7	0 200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist		
Inj 0.5 mg vial255.0	0 1	✓ Cosmegen
Inj 0.5 mg for ECP255.0	0 0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		
Inj 2 mg per ml, 10 ml149.5	0 1	✓ Pfizer
Inj 20 mg vial1,495.0		✓ Daunorubicin
		Zentiva S29
Inj 20 mg for ECP149.5	0 20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist		
Inj 20 mg48.7	5 1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial46.8		✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial26.9	5 1	✓ Docetaxel
		Accord \$29
Inj 80 mg195.0	0 1	✓ Docetaxel Sandoz
Inj 1 mg for ECP		✓ Baxter
	-	

	Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist	Ψ		-	
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Do	oxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		oxorubicin Ebewe
· , - · · g p · · · · , - · · · · · · · · · · · · · ·	17.00			rrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1		oxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Ar	rrow-Doxorubicin
	69.99		✓ Do	oxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Ba	axter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist				
Inj 2 mg per ml, 5 ml vial	25.00	1	√ Fr	oirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1		pirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Er	pirubicin Ebewe
Inj 1 mg for ECP		1 mg		
ETOPOSIDE		9		
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	✓ Va	epesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10		epesid
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Special		1		ex Medical
Inj 1 mg for ECP – PCT only – Specialist		1 mg		
ETOPOSIDE PHOSPHATE – PCT only – Specialist		9		27.01
, ,	40.00	1	./ E+	opophos
Inj 100 mg (of etoposide base)		ı 1 mg		
, , ,		ı ıııy	▼ Do	axici
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail phar		400		
Cap 500 mg	23.82	100	✓ <u>De</u>	<u>evatis</u>
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial - PCT only - Specialist	109.74	1	✓ Za	rvedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓ Za	rvedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	✓ Ba	axter
LENALIDOMIDE – Retail pharmacy-Specialist – Special Authorit Wastage claimable	y see SA2047 below			
Cap 5 mg	5,122.76	28	✓ Re	evlimid
Cap 10 mg	4,655.25	21	✓ Re	evlimid
. •	6,207.00	28	✓ Re	evlimid
Cap 15 mg	5,429.39	21	✓ Re	evlimid
· · · ·	7,239.18	28	✓ Re	evlimid
Cap 25 mg	7,627.00	21	✓ Re	evlimid

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist314.00	50	Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist407.40	15	Uromitexan
Inj 1 mg for ECP - PCT only - Specialist2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist		
Inj 5 mg vial641.70	1	✓ Accord S29
Inj 20 mg vial	1	✓ Omegapharm S29
		✓ Teva
Inj 1 mg for ECP470.75	1 mg	✓ Baxter
MITOZANTRONE - PCT only - Specialist		
Inj 2 mg per ml, 10 ml vial97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see SA1883 below		
Tab 100 mg3,701.00	56	✓ Lynparza
Tab 150 mg3,701.00	56	✓ Lynparza

SA1883 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

	Subsidy		Fully	Brand or
(Mar	nufacturer's Price)	Subs	sidised	Generic
	\$	Per	1	Manufacturer

continued...

- 1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
- 2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
- 3 Patient has received at least two lines of previous treatment with platinum-based chemotherapy; and
- 4 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line of platinum-based chemotherapy; and
- 5 Patient's disease must have achieved partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 6 Patient's disease has not progressed following prior treatment with olaparib; and
- 7 Treatment will be commenced within 8 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 8 Treatment to be administered as maintenance treatment: and
- 9 Treatment not to be administered in combination with other chemotherapy.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 No evidence of progressive disease: and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy.

Note: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.

PACLITAXEL - PCT only - Specialist

Inj 30 mg	47.30	5	Paclitaxel Ebewe
Inj 100 mg		1	✓ Paclitaxel Ebewe
	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority s	ee SA1979 below		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

PENTOSTATIN [DEOXYCOFORMYCIN] - PCT only - Specialist

Inj 10 mg	CBS	1	✓ Nipent S2:
-----------	-----	---	--------------

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PROCARBAZINE HYDROCHLORIDE – PCT – Retail pharmacy-	Specialist			
Cap 50 mg	980.00	50	✓	Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 below - Reta	il pharmacy			
Cap 5 mg	9.13	5	✓	Temaccord
Cap 20 mg	16.38	5	1	Temaccord
	18.30		1	Apo-Temozolomide
	136.00	14	✓	Accord S29
Cap 100 mg	35.98	5	✓	Temaccord
•	40.20		1	Apo-Temozolomide
	532.00	14	1	Accord \$29
Cap 140 mg	50.12	5	✓	Temaccord
	400.00		✓	Amneal S29
Cap 180 mg	620.00	14	1	Accord S29
Cap 250 mg		5	✓	Temaccord
	688.00		1	Amneal S29

⇒SA1741 Special Authority for Subsidy

Initial application — **(high grade gliomas)** only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*: and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Su	ubsidy	Fully	Brand or
(Manufac	turer's Price) Subsid	lised	Generic
	\$ Per	✓	Manufacturer

continued...

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE - Retail pharmacy-Specialist - Speci	ial Authority see SA1124 below		
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

TRETINOIN

Cap 10 mg - PCT - Retail pharmacy-Specialist4	79.50	100	Vesanoid
VENETOCLAX - Retail pharmacy-Specialist - Special Authority see SA	A1868 below		
Tab 14 \times 10 mg, 7 \times 50 mg, 21 \times 100 mg	71.86	42 OP	✓ Venclexta
Tab 10 mg	95.78	14 OP	✓ Venclexta
Tab 50 mg2	39.44	7 OP	✓ Venclexta
Tab 100 mg - Wastage claimable8,2	09.41	120	✓ Venclexta

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued

recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist102.73	5	DBL Vincristine Sulfate
Inj 1 mg for ECP - PCT only - Specialist12.60	1 mg	✓ Baxter
VINORELBINE - PCT only - Specialist		
Inj 10 mg per ml, 1 ml vial12.00	1	✓ Navelbine
42.00		✓ Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial56.00	1	✓ Navelbine
210.00		✓ Vinorelbine Ebewe
328.65		✓ Sagent S29
Inj 1 mg for ECP	1 mg	✓ Baxter
Inj 50 mg for ECP328.65	50 mg OP	✓ Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

ALECTINIB - Retail pharmacy-Specialist - Special Authority see SA1870 below Wastage claimable

224 ✓ Alecensa

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
\$	Per	✓	Manufacturer	

continued...

Al K test: and

3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Tab 20 mg	3,774.06	60	Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day: or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or
 - 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
 - 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or

3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*: and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

Tab 100 mg	764.00	30	Tarceva
Tab 150 mg	1,146.00	30	✓ Tarceva

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fither
 - 3.1 Patient is treatment naive: or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either:
 - 2.1 Patient is treatment naive; or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460 on the		
	next page2,400.00	60	✓ Glivec
*	Cap 100 mg58.23	60	✓ Imatinib-Rex
*	Cap 400 mg84.79	30	✓ Imatinib-Rex

Subsidy (Manufacturer's Price)

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST - access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 below - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	Tasigna
Cap 200 mg	6.532.00	120	✓ Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Either:
 - 2.1 Patient has documented CML treatment failure* with imatinib; or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
 - 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
 - 3 Maximum nilotinib dose of 800 mg/day; and
 - 4 Subsidised for use as monotherapy only.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PALBOCICLIB – Retail pharmacy-Specialist – Special Authority Wastage claimable	see SA1894 below			
Tab 75 mg	4,000.00	21	✓	Ibrance
Tab 100 mg	4,000.00	21	✓	Ibrance
Tab 125 mg	4,000.00	21	✓	Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Fither:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB -	Special Aut	hority see	SA1190	below -	Retail p	harmacy
Tab 000 m	~					4.0

✓ Votrient	30	1,334.70	······································	Tab 200 mg
✓ Votrient	30	2.669.40		Tab 400 mg

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
 - 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and

Subsidy (Manufacturer's Price	e)	Fully Subsidised	Brand or Generic	
<u> </u>	Per	1	Manufacturer	

continued...

- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

RUXOLITINIB - Special Authority see SA1890 below - Retail pharmacy

Wastage claimable

Tab 5 mg	2,500.00	56	Jakavi
Tab 10mg		56	Jakavi
Tab 15 mg		56	Jakavi
Tab 20 mg	5,000.00	56	Jakavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 on the next page - Retail pharmacy

Cap 12.5 mg	208.38	28	 Sunitinib Pfizer
Cap 25 mg	416.77	28	✓ Sunitinib Pfizer
Cap 50 mg		28	✓ Sunitinib Pfizer

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and
 - The patient has intermediate or poor prognosis defined as:
- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib: or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable

\$ Per ✔ Manufacturer		Subsidy (Manufacturer's Price) \$		Fully Subsidised	Brand or Generic Manufacturer
-----------------------	--	---	--	---------------------	-------------------------------------

continued...

disease): or

- 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 83

ABIRATERONE ACETATE - Retail pharmacy-Specialist - Special Authority see SA2118 below

Wastage claimable

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Fither:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer	
--	--

continued...

4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

BICAL LITAMIDE

Tab 50 mg4.21	28	✓ Binarex
FLUTAMIDE		
Tab 250 mg107.55	90	✓ Prostacur S29
119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Authority see SA1895 below		
Inj 50 mg per ml, 5 ml prefilled syringe1,068.00	2	✓ Faslodex

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

Tab 160 mg	48.80	30	✓ Megace S29
(Megace S29 Tab 160 mg to be delisted 1 February 2023)			
OCTREOTIDE			
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
OCTREOTIDE LONG-ACTING - Special Authority see SA21	19 on the next page -	- Retail pha	rmacy
Inj depot 10 mg prefilled syringe	439.97	1	 Octreotide Depot
			Teva
Inj depot 20 mg prefilled syringe	647.03	1	 Octreotide Depot
			<u>Teva</u>
Inj depot 30 mg prefilled syringe	718.55	1	✓ Octreotide Depot
			Teva

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:
Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Either:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas: and
 - 3.2 Surgery is contraindicated or has failed; or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or

continued...

- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 Patient has a large pituitary tumour, greater than 10 mm at its widest; and
- 3 Patient is scheduled to undergo pituitary surgery in the next six months.

TAMOXIFEN CITRATE

*	Tab 10 mg15.0	0 60	/	Tamoxifen Sandoz
	Tab 20 mg		/	Tamoxifen Sandoz

Aromatase Inhibitors

ANASTROZOLE

*	Tab 1 mg4.55	30	✓ Anatrole
EX	EMESTANE		
*	Tab 25 mg14.50	30	✓ Pfizer Exemestane
LE ⁻	TROZOLE		
*	Tab 2.5 mg	30	✓ Letrole

Immunosuppressants

Cytotoxic Immunosuppressants

AZATHIOPRINE

*	Tab 25 mg7.35	60	Azamun
*	Tab 50 mg	100	Azamun
*	Inj 50 mg vial199.00	1	Imuran

(Imuran Inj 50 mg vial to be delisted 1 January 2023)

MYCOPHENOLATE MOFETIL

Tab 500 mg	35.90	50	 Cellcept
Cap 250 mg	35.90	100	✓ Cellcept
Powder for oral liq 1 g per 5 ml - Subsidy by endorsement	187.25	165 ml OP	 Cellcept

Mycophenolate powder for oral liquid is subsidised only for patients unable to swallow tablets and capsules, and when the prescription is endorsed accordingly.

Fusion Proteins

ETANERCEPT	- Special Authority	y see SA2103 on the ne	ext page - Retail pharmacy
-------------------	---------------------	------------------------	----------------------------

in a contract of the contract			
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector		4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Ini 50 mg prefilled syringe	1.050.00	4	✓ Enbrel

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

⇒SA2103 Special Authority for Subsidy

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
- 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis; or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroiliitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or

Subsidy (Manufacturer's Price)	Fully Subsidised		,		Brand or Generic
 \$	Per	✓	Manufacturer		

continued...

2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and

2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm 45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA): and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance: and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 12 Fither
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or

Subsidy	Ful	ly Brand or	
(Manufacturer's Price)	Subsidise	ed Generic	
\$	Per •	 Manufacture 	r

continued...

1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or

2 All of the following:

- 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
- 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
- 2.4 Fithou
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

1 Either:

- 1.1 Applicant is a rheumatologist; or
- 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and

2 Either:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic
\$ Per ✔ Manufacturer

continued...

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:
 - 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
 - 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses unless contraindicated); and
 - 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate: and
 - 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

- 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin. or acitretin: and
- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a

- All of the following:

 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
 - 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Either:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
 - 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

((Subsidy Manufacturer's Price)	Fully Subsidised		. ,		
<u></u>	\$	Per	✓	Manufacturer		

continued...

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:
 - 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBULIN (EQUINE) - PCT only - Specialist			
Inj 50 mg per ml, 5 ml	.2,774.48	5	✓ ATGAM
BACILLUS CALMETTE-GUERIN (BCG) VACCINE - PCT only - Sp	ecialist		
Subsidised only for bladder cancer.			
Inj 2-8 × 100 million CFU	149.37	1	✓ OncoTICE
Ini 40 mg per ml. vial	176.90	3	✓ SII-Onco-BCG S29

Monoclonal Antibodies

	ıarmacy	 Retail pha 	ee SA2142 on the next page	ADALIMUMAB (AMGEVITA) – Special Authority se
✓ Amgevita	1	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	✓	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	/	2	375.00	Ini 40 mg per 0.8 ml prefilled syringe.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2142 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira: or
- 2 Both:
 - 2.1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
 - 2.2 Either:
 - 2.2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or
 - 2.2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
 - 2.2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
 - 2.3 Patient has 3 or more active lesions; and
 - 2.4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and
 - 2.1.2 Either:
 - 2.1.2.1 Patient has experienced intolerable side effects; or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis: or
 - 2.2 All of the following:
 - 2.2.1 Either:
 - 2.2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and

Subsidy (Manufacturer's Price)	Sul	Fully bsidised	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2.2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Either:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Both:
 - 2.1 Patient has pyoderma gangrenosum*; and
 - 2.2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has severe active Crohn's disease; and
 - 2.2 Any of the following:
 - 2.2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Subsidy (Manufacturer's Price)	Su	Fully ibsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Paediatric patient has active Crohn's disease: and
 - 2.2 Either:
 - 2.2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2.2 Patient has extensive small intestine disease; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids: and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has confirmed Crohn's disease; and
 - 2.2 Any of the following:
 - 2.2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.2.3 Patient has complex peri-anal fistula; and
 - 2.3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
- 2.2 Both:
 - 2.2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
 - 2.2 Both:
 - 2.2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses: or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient has previously had an approval for Humira; or
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 212 Fither
 - 2.1.2.1 The patient has experienced intolerable side effects; or
 - 2.1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
 - 2.2 All of the following:
 - 2.2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.2.3 Patient has bilateral sacroillitis demonstrated by radiology imaging; and
 - 2.2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.2.5 Either:
 - 2.2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
 - 2.2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:
Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 2.1.2 Fither:
 - 2.1.2.1 Patient has experienced intolerable side effects: or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
 - 2.2 All of the following:
 - 2.2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.2.3 Either:
 - 2.2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Si	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 2.1.2 Fither:
 - 2.1.2.1 Patient has experienced intolerable side effects; or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA; or
 - 2.2 All of the following:
 - 2.2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 2.1.2 Either:
 - 2.1.2.1 The patient has experienced intolerable side effects; or
 - 2.1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
- 2.2.4 Either:
 - 2.2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.2.5 Any of the following:
 - 2.2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.2.5.2 Patient has an ESR greater than 25 mm per hour; or
 - 2.2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 2.1.2 Either:
 - 2.1.2.1 The patient has experienced intolerable side effects; or
 - 2.1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
 - 2.2 All of the following:
 - 2.2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.2.5 Either:
 - 2.2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin: or
 - 2.2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.2.6 Fither:

Subsidy	Fully	Brand or	Т
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 2.2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
- 2.2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 2.1.2 Either:
 - 2.1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 2.1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
 - 2.2 All of the following:
 - 2.2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has histologically confirmed ulcerative colitis; and
 - 2.2 Fither:
 - 2.2.1 Patient's SCCAI score is greater than or equal to 4; or
 - 2.2.2 Patient's PUCAI score is greater than or equal to 65; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids; and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

Subsidy (Manufacturer's Price)	Ful Subsidise	,	
\$	Per •	Manufacturer	

continued...

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
 - 2.3 Any of the following:
 - 2.3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 2.3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
 - I of the following:
 - 2 All of the following:
 - 2.1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
 - 2.2 Patient has axial inflammatory pain for six months or more; and
 - 2.3 Patient is unable to take NSAIDs; and
 - 2.4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
 - 2.5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
 - 2.2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose; and

Subsidy (Manufacturer's Price)	Sub	Fully	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Following initi
 - 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA) - Special Authority see SA2101 below - Retail pharmacy

(Humira Inj 20 mg per 0.4 ml prefilled syringe to be delisted 1 December 2022)

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	Humira
Inj 20 mg per 0.4 ml prefilled syringe	•	2	Humira
Inj 40 mg per 0.8 ml prefilled pen	1,599.96	2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	Humira

⇒SA2101 Special Authority for Subsidy

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy	F	ılly	Brand or
(Manufacturer's	s Price) Subsidis	sed	Generic
\$	Per	✓	Manufacturer

continued...

- 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
- 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

Renewal — (Crohn's disease - adults) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Fither:
 - 2.1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab;
 - 2.1.2 CDAI score is 150 or less; or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that CDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a gastroenterologist; or
 - 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Fither:
 - 2.1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 2.1.2 PCDAI score is 15 or less: or
 - 2.2 Both:
 - 2.2.1 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
 - 2.2.2 Applicant to indicate the reason that PCDAI score cannot be assessed; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Fither:

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	1	Manufacturer

continued...

- 1.1 Applicant is a gastroenterologist; or
- 1.2 Applicant is a Practitioner and confirms that a gastroenterologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 2.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Renewal — (hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or

Subsidy (Manufacturer's Price)	Full Subsidise		
\$	Per 🗸	Manufacturer	

continued...

- 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment: and
- 3 A maximum of 8 doses.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 3 Either:
 - 3.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 3.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved guality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (severe chronic plague psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with adalimumab treatment; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Either:
 - 2.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value: or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline valuee; or

2.2 Both:

- 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
- 2.2.2 Fither:
 - 2.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A treatment course is defined as a minimum of 12 weeks adalimumab treatment

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if adalimumab is withdrawn.

AFLIBERCEPT – Special Authority see SA1772 below – Retail pharmacy

Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy; or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Either:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab; or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
- 2 Either:

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
- 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid): and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Authority see SA2096 below

⇒SA2096 Special Authority for Subsidy

Initial application — (Treatment of profoundly immunocompromised patients) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 The patient is in the community with mild to moderate disease severity*; and
- 3 Patient is profoundly immunocompromised** and is at risk of not having mounted an adequate response to vaccination against COVID-19 or is unvaccinated; and
- 4 Patient's symptoms started within the last 10 days; and
- 5 Patient is not receiving high flow oxygen or assisted/mechanical ventilation; and
- 6 Casirivimab and imdevimab is to be administered at a maximum dose of no greater than 2,400 mg.

Notes: * Mild to moderate disease severity as described on the Ministry of Health Website

** Examples include B-cell depletive illnesses or patients receiving treatment that is B-Cell depleting.

CETUXIMAB - PCT only - Specialist - Special Authority see SA1697 on the next page

Inj 5 mg per ml, 20 ml vial	364.00	1	Erbitux
Inj 5 mg per ml, 100 ml vial	1,820.00	1	✓ Erbitux
Inj 1 mg for ECP	3.82	1 mg	✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$
Per ✓ Manufacturer

⇒SA1697 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, non-metastatic, squamous cell cancer of the head and neck; and
- 2 Patient is contraindicated to, or is intolerant of, cisplatin; and
- 3 Patient has good performance status; and
- 4 To be administered in combination with radiation therapy.

GEMTUZUMAB OZOGAMICIN − PCT only − Specialist − Special Authority see SA2136 below Inj 5 mg vial12,973.00 1 ✓ Mylotarg

⇒SA2136 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with daunorubicin and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only; and
- 9 Either:
 - 9.1 Gemtuzumab ozogamicin to be administered as one dose at 3 mg per m² body surface area; or
 - 9.2 Up to 10 mg of gemtuzumab ozogamicin to be administered.

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2082 below

 Inj 100 mg
 806.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 8.29
 1 mg
 ✓ Baxter

⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease: and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Paediatric patient has severe active Crohn's disease; and
- 2 Either:
 - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (**Graft vs host disease**) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Fither

Subsidy	F	ully	Brand or	_
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

continued...

- 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
- 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or
- 2 Both:
 - 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease; and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or

	Subsidy		ully	Brand or
(I	Manufacturer's Price)	Subsid	ised	Generic
	\$	Per	1	Manufacturer

continued...

2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement; and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Fither:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsid	dised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Fither:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis: or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis: or

continued...

- 2.10 Severe ulcerative colitis; or
- 2.11 Plaque psoriasis; or
- 2.12 Neurosarcoidosis: or
- 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept: and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Either:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved guality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fullminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.</p>

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
 - 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids: and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
MEPOLIZUMAB - Special Authority see SA1896 below - Retail	pharmacy			
Inj 100 mg prefilled pen	1,638.00	1	✓	Nucala
Inj 100 mg vial	1,638.00	1	✓	Nucala

⇒SA1896 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eq. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
 - 2 Fither:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Author	rity see SA1627 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	✓ Gazyva
Ini 1 ma for ECP	6.21	1 ma	✓ Baxter

⇒SA1627 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive: and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	✓ Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids; and
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Either:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- Eilliel.
 - 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
 - 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and
 - 2.2.2.2 Either:
 - 2.2.2.2.1 Patient has chronic lung disease; or
 - 2.2.2.2.2 Patient is Māori or any Pacific ethnicity; or
 - 2.2.3 Both:
 - 2.2.3.1 Patient has haemodynamically significant heart disease; and
 - 2.2.3.2 Any of the following:
 - 2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see note a): or
 - 2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or
 - 2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or
 - 2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

 a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient will require surgical palliation/definitive repair within the next 3 months.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Ejection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 below

	colai riainoni, coo crittoro zonon	
Inj 100 mg per 10 ml vial	1,075.50 2	✓ Mabthera
Inj 500 mg per 50 ml vial	2,688.30 1	✓ Mabthera
Inj 1 mg for ECP	5.64 1 mg	✓ Baxter (Mabthera)

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated; and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroquine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

intramuscular gold: or

- 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Fither:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and

Subsidy	:)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
<u> </u>	Per	✓	Manufacturer

continued...

4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Fither:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special Authority see SA2114 below

Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:
 - 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
 - 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
 - 3.3 Cyclophosphamide and methotrexate are contraindicated; or
 - 3.4 Patient is a female of child-bearing potential; or
 - 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment; and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Fither:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia; and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL; and
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

continued...

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks: and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*: and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective: or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Fither:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microlitre; or

Subsidy	Full	/ Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	Manufacturer Manufacturer	

continued...

- 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia*

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Either:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*: and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment: and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and

Subsidy	Full	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per 🗸	 Manufacturer 	

continued...

- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*; and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin: and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Either:
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1.000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	9	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 × 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
 - 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
 - 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and
- 3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

experienced intolerable side effects.

d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and
 - 2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	ıbsidised	Generic	
\$	Per	1	Manufacturer	

continued...

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab; or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or

	Subsidy	Ful	ly Brand or	
((Manufacturer's Price)	Subsidise	ed Generic	
	\$	Per •	 Manufacturer 	

continued...

2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 on the next page - Retail pharmacy

Note	: Siltuximab is to be administered at doses no grea	ter than 11 mg/kg every	3 weeks.	
Inj 1	00 mg vial	770.57	1	Sylvant
Inj 4	00 mg vial	3,082.33	1	✓ Sylvant

Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per ml, 1.5 ml vial with cilgavimab 100 mg per ml,1.5 ml vial	1	✓ Evusheld
TOCILIZUMAB - PCT only - Special Authority see SA2100 below		
Inj 20 mg per ml, 4 ml vial220.00	1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
880.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial550.00	1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial1,100.00	1	✓ Actemra
		✓ Actemra S29 S29
		✓ RoActemra S29 S29
4,400.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP2.85	1 mg	✓ Baxter

⇒SA2100 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the

Subsidy	· F	ully Bra	nd or
(Manufacturer's	s Price) Subsidi	sed Ger	neric
\$	Per	✓ Mar	nufacturer

continued...

following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis; or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Roth
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Fither:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2 Tocilizumab is to be used as monotherapy; and
- 3 Either:
 - 3.1 Treatment with methotrexate is contraindicated: or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Either:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Fither:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Fither:

Subsidy (Manufacturer's Pric	ce)	Fully Subsidised	Brand or Generic	
	Per	1	Manufacturer	

continued...

- 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Roth:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Subsidy		Fully	Brand or	
(Manufacturer's Price)) 5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19*) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Note: Indications marked with * are unapproved indications.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:

Both:

- 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

 TRASTUZUMAB – PCT only – Specialist – Special Authority see SA1632 on the next page

 Inj 150 mg vial
 1,350.00
 1
 ✓ Herceptin

 Inj 440 mg vial
 3,875.00
 1
 ✓ Herceptin

 Inj 1 mg for ECP
 9.36
 1 mg
 ✓ Baxter

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib: and
- 3 Fither:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria: All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology):
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsid	dised	Generic
\$	Per	✓	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	2,320.00	1	Kadcyla
Inj 160 mg vial	3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	24.52	1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and
- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Fither:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:

ibsidy	Fully	Brand or
turer's Price) Sub	osidised	Generic
 \$ Per	/	

continued...

- 5.1 Patient does not have symptomatic brain metastases; or
- 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

		NIVOLUMAB – PCT only – Specialist – Special Authority see SA2120 below
Opdivo	1	Inj 10 mg per ml, 4 ml vial
✓ Opdivo	1	Inj 10 mg per ml, 10 ml vial2,629.96
✓ Baxter	1 mg	Inj 1 mg for ECP27.62

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
 - 2 All of the following:

Subsidy (Manufacturer's Pri	ice)	Fu Subsidis	,	Brand or Generic
\$	P	er	/	Manufacturer

continued...

- 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
- 2.2 Patient has signs of disease progression; and
- 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or

Subsidy		Fully	Brand or
(Manufacturer's Price)	5	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
- 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg		50	✓ Neoral
Cap 100 mg		50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	Neoral
EVEROLIMUS – Special Authority see SA2008 below Wastage claimable	- Retail pharmacy		
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has tuberous sclerosis; and
- 2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

Note: MRI should be performed at minimum once every 12 months, more frequent scanning should be performed with new onset of symptoms such as headaches, visual complaints, nausea or vomiting, or increase in seizure activity.

SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- · Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy: or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	✓	Manufacturer

continued...

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
 - 3 Seizures have a significant impact on quality of life; and
 - 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: "Optimal treatment" is defined as treatment, which is indicated and clinically appropriate for the patient, given in adequate doses for the patients age, weight and other features affecting the pharmacokinetics of the drug, with good evidence of adherence. Women of childbearing age are not required to have a trial of sodium valproate.

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

to to bottom the transfer of the tottom the tottom	an priarriacy		
Cap 0.5 mg	49.60	100	✓ Tacrolimus Sandoz
Cap 0.75 mg	99.30	100	✓ Tacrolimus Sandoz
Cap 1 mg		100	✓ Tacrolimus Sandoz
Cap 5 mg		50	✓ Tacrolimus Sandoz

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT – Special Authority see SA1558 below – Retail pharmacy
Inj 10 mg per ml, 3 ml prefilled syringe.......2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

BEE VENOM ALLERGY TREATMENT - Special Authority see S	A1367 above - F	Retail pharma	су
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluer	nt 305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see	SA1367 above	- Retail pharr	macy
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze	005.00	4.00	411
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze	205.00	1 OP	√ Alboy
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml		TOP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze dried venom, with diluent		1 OP	✓ Venomil S29
uneu venom, with unuell		i OF	V CHOIIII 023

227

	Subsidy		Fully	
	(Manufacturer's Price		Subsidised	
	\$	Per		Manufacturer
Antihistamines				
CETIRIZINE HYDROCHLORIDE				
* Tab 10 mg	1 12	100	1	Zista
* Oral liq 1 mg per ml		200 ml		Histaclear
CHLORPHENIRAMINE MALEATE				
* Oral liq 2 mg per 5 ml	0.27	500 ml	_	Histafen
	9.37	300 1111	•	riistaleli
DEXTROCHLORPHENIRAMINE MALEATE				
* Tab 2 mg		40		
	(8.40)			Polaramine
	1.01	20		
	(5.99)			Polaramine
* Oral liq 2 mg per 5 ml		100 ml		-
	(10.29)			Polaramine
FEXOFENADINE HYDROCHLORIDE				
* Tab 60 mg	4.34	20		
-	(8.23)			Telfast
* Tab 120 mg	4.74	10		
	(8.23)			Telfast
	14.22	30		
	(26.44)			Telfast
LORATADINE				
* Tab 10 mg	1.69	100	/	Lorafix
* Oral lig 1 mg per ml		100 ml		Haylor syrup
PROMETHAZINE HYDROCHLORIDE				,
	1 20	50		Allersoothe
* Tab 10 mg		30	•	AllerSouthe
* Tab 25 mg		50		Allersoothe
Allersoothe to be Principal Supply on 1 September 2022		30	•	AllerSouthe
* Oral lig 1 mg per 1 ml		100 ml	_	Allersoothe
* Inj 25 mg per ml, 2 ml ampoule – Up to 5 inj available on a F		5		Hospira
* III 25 IIIg per IIII, 2 IIII ampodie – op to 5 IIIj avallable off a f	30 17.07	J		поэрпа
Inhaled Corticosteroids				
BECLOMETHASONE DIPROPIONATE				
Aerosol inhaler, 50 mcg per dose	14.01 2	200 dose	OP 🗸	Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose		Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose		Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose		Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose	OP 🗸	Beclazone 250
BUDESONIDE				
	17.00	200 dose	∩D ./	Pulmicort
Powder for inhalation, 100 mcg per dose	17.00 2	200 dose	OP •	
Decides for inhelation, 000 many and as	40.00	000 de e	OD 4	Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	200 dose	OP 🗸	Pulmicort
5				Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	200 dose	OP 🗸	Pulmicort
				Turbuhaler

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic Manufacturer
	\$	Per	✓ Manufacturer
FLUTICASONE			4
Aerosol inhaler, 50 mcg per dose		120 dose OP	
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	
Aerosol inhaler, 250 mcg per dose		120 dose OP	
Powder for inhalation, 250 mcg per dose	13.60	60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agonis	ts		
EFORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose devi		60 dose	
	(35.80)		Foradil
(Foradil Powder for inhalation, 12 mcg per dose, and monodose	aevice to be deli	sted 1 July 202	23)
EFORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dose	e) 10.32	60 dose OP	
	(16.90)		Oxis Turbuhaler
INDACATEROL			
Powder for inhalation 150 mcg	61.00	30 dose OP	Onbrez Breezhaler
Powder for inhalation 300 mcg	61.00	30 dose OP	 Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	✓ Serevent
Powder for inhalation, 50 mcg per dose, breath activated		60 dose OP	Serevent Accuhaler
Inhaled Corticosteroids with Long-Acting Beta-	Adrenocepto	or Agonists	
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide w	vith		
6 mcg eformoterol fumarate metered dose)		120 dose OP	✓ DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fumar			
per dose (equivalent to 400 mcg budesonide with 12 mc			
eformoterol fumarate metered dose) - No more than 2	3		
dose per day	82.50	120 dose OP	✓ DuoResp Spiromax
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg	18.23	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 n	ncg33.74	120 dose OP	✓ Symbicort
-	-		Turbuhaler 100/6
Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Vannair
Powder for inhalation 200 mcg with eformoterol fumarate 6 n	ncg 44.08	120 dose OP	✓ Symbicort
-			Turbuhaler 200/6
Powder for inhalation 400 mcg with eformoterol fumarate			
12 mcg - No more than 2 dose per day	44.08	60 dose OP	✓ Symbicort
			Turbuhaler 400/12
FLUTICASONE FUROATE WITH VILANTEROL			
Powder for inhalation 100 mcg with vilanterol 25 mcg	44.08	30 dose OP	✓ Breo Ellipta

	Subsidy (Manufacturer's \$	Price) Subs	Fully sidised	Brand or Generic Manufacturer
FLUTICASONE WITH SALMETEROL	•			
Aerosol inhaler 50 mcg with salmeterol 25 mcg		120 dose OP 120 dose OP	_	<u>Seretide</u> Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No more than 2 dose per day		60 dose OP	_	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day		60 dose OP	√ 9	Seretide Accuhaler
Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Oral lig 400 mcg per ml	40.00	150 ml	/ \	/entolin
Infusion 1 mg per ml, 5 ml		10	_	/entolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO		5	✓ \	/entolin
Inhaled Beta-Adrenoceptor Agonists				
SALBUTAMOL				
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000				
dose available on a PSO	3.80	200 dose OP		Respigen SalAir
	(6.20)		١	/entolin
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ <u>I</u>	<u>Asthalin</u>
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb available on a PSO		20	✓ <u>/</u>	<u>Asthalin</u>
ERBUTALINE SULPHATE				
Powder for inhalation, 200 mcg per dose (equivalent to 250 mcg metered dose), breath activated	22.20	120 dose OP	✓ E	Bricanyl Turbuhaler
,				
Anticholinergic Agents				
PRATROPIUM BROMIDE				
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dos available on a PSO		200 dose OP	1	Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule – Up to 40 ne available on a PSO		20	√ (Jnivent
Inhaled Beta-Adrenoceptor Agonists with Antic	holinergic <i>F</i>	Agents		
SALBUTAMOL WITH IPRATROPIUM BROMIDE Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p dose CFC-free		200 doss OD	./ 1	Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per	12.19	200 dose OP	• [Judiiii HFA
vial, 2.5 ml ampoule – Up to 20 neb available on a PSO	11.04	20	√ [<u>Duolin</u>

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL − Special Authority see SA1584 above − Retail pharmacy
Powder for Inhalation 50 mcg with indacaterol 110 mcg.....81.00 30 dose OP ✓ Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL – Special Authority see SA1584 above – Retail pharmacy

Soln for inhalation 2.5 mcg with olodaterol 2.5 mcg81.00 60 dose OP ✓ Spiolto Respimat

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for inhalation 62.5 mcg with vilanterol 25 mcg77.00 30 dose OP ✓ Anoro Ellipta

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE - Retail pharmacy-Specialist - Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib.

Esbriet	90	Tab 801 mg3,645.00
✓ Esbriet	90	Tab 267 mg1,215.00

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price \$) Sub Per	Fully osidised	Brand or Generic Manufacturer
eukotriene Receptor Antagonists				
ONTELUKAST				
Tab 4 mg	3.10	28	✓ N	lontelukast Mylan
Tab 5 mg	3.10	28		lontelukast Mylan
Tab 10 mg	2.90	28	✓ N	lontelukast Mylan
Methylxanthines				
MINOPHYLLINE				
Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available on a				
PSO	180.00	5	✓ D	BL Aminophylline
HEOPHYLLINE				
Tab long-acting 250 mg	23.02	100	✓ N	uelin-SR
Oral liq 80 mg per 15 ml		500 ml	✓ N	uelin
Mucolytics				
DRIACE ALEA — Chaoial Authority and CA1079 halour — Batai	Inharmany			
DRNASE ALFA - Special Authority see SA1978 below - Retai Nebuliser soln, 2.5 mg per 2.5 ml ampoule		6	√ D	ulmozyme
nebuliser soin, 2.5 mg per 2.5 mi ampoule	200.00	U	• -	uiiiiozyiile

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month period: or
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25; or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

Tab 150 mg	29,386.00	56	✓ Kalydeco
Oral granules 50 mg, sachet	29,386.00	56	✓ Kalydeco
Oral granules 75 mg, sachet	29,386.00	56	✓ Kalydeco

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Either:
 - 2.1 Patient must have G551D mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene on at least 1 allele; or
 - 2.2 Patient must have other gating (class III) mutation (G1244E, G1349D, G178R, G551S, S1251N, S1255P, S549N

RESPIRATORY SYSTEM AND ALLERGIES				
	Subsidy (Manufacturer's \$	Price) Subs	Fully idised	Brand or Generic Manufacturer
continued and S549R) in the CFTR gene on at least 1 allele	; and			
3 Patients must have a sweat chloride value of at least 60 sweat collection system; and	mmol/L by quant	itative pilocarpir	ne ionto	phoresis or by Macroduct
4 Treatment with ivacaftor must be given concomitantly wi 5 Patient must not have an acute upper or lower respirator (including antibiotics) for pulmonary disease in the last 4 6 The dose of ivacaftor will not exceed one tablet or one so 7 Applicant has experience and expertise in the managem	y infection, pulmo weeks prior to co achet twice daily;	onary exacerbat ommencing trea and	ion, or o	changes in therapy
SODIUM CHLORIDE				
Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	✓ B	iomed
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP		teroClear teroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	✓ <u>F</u>	lixonase Hayfever & Allergy
IPRATROPIUM BROMIDE Aqueous nasal spray, 0.03%	5.23	15 ml OP	√ <u>U</u>	nivent
Respiratory Devices				
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO c) Only for children aged six years and under	2 20	1		ohambar Mask

1	✓ e-chamber Mask
1	Mini-Wright AFS Low Range
1	Mini-Wright Standard
1	✓ e-chamber Turbo
1	e-chamber La Grande
1	✓ Volumatic
	1 1 1 1 1

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml).......15.10 25 ml OP **✓ Biomed**



	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
Ear Preparations			
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYC	IN AND NYSTAT	ΓIN	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5.16	7.5 ml OP	✓ Kenacomb
Ear/Eye Preparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4 50	8 ml OP	
gramoran oo mog por mi	(9.27)	0 1111 01	Sofradex
FRAMYCETIN SULPHATE	(- /		
Ear/Eye drops 0.5%	4.13	8 ml OP	
	(8.65)		Soframycin
	, ,		•
Eye Preparations			
Eye preparations are only funded for use in the eye, unless expl	icitly stated other	wise.	
Anti-Infective Preparations			
And intective repairations			
ACICLOVIR			
* Eye oint 3%	14.88	4.5 g OP	✓ <u>ViruPOS</u>
CHLORAMPHENICOL			
Eye oint 1%		5 g OP	✓ Devatis
Eye drops 0.5%		10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * a	re unapproved in	uications.	
CIPROFLOXACIN	0.70	E ml OD	Cinvellevesin Tave
Eye drops 0.3% – Subsidy by endorsement		5 ml OP	✓ Ciprofloxacin Teva
for the second line treatment of chronic suppurative otiti		•	•
Note: Indication marked with a * is an unapproved indic		, and the pres	onphornic chached accordingly.
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)			
PROPAMIDINE ISETHIONATE			
* Eye drops 0.1%	2.97	10 ml OP	
, , , , , , , , , , , , , , , , , , , ,	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
TOBRAMYCIN		-	
Eye oint 0.3%	10.45	3.5 g OP	✓ Tobrex
Eye drops 0.3%		5 ml OP	✓ Tobrex
•			

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Corticosteroids and Other Anti-Inflammatory Preparations

DE	XAMETHASONE			
*	Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
*	Eye drops 0.1%	4.50	5 ml OP	Maxidex
	Ocular implant 700 mcg - Special Authority see SA1680 below			
	- Retail pharmacy1,44	4.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per g	5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin b sulphate 6,000 u per ml	4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM Eye drops 0.1%	8.80	5 ml OP	✓ <u>Voltaren Ophtha</u>
FLUOROMETHOLONE * Eye drops 0.1%	3.09	5 ml OP	✓ FML
	5.20		✓ Flucon

	Subsidy (Manufacturer's Pric	e) Sub	Fully sidised	Brand or Generic
	\$	Per	1	Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
• •	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	5.93	10 ml OP	✓ P	rednisolone-AFT
•	7.00	5 ml OP	√ P	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Authority	see SA1715 below -	Retail phar	macv	
Eye drops 0.5%, single dose (preservative free)		20 dose	•	linims Prednisolone

⇒SA1715 Special Authority for Subsidy

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE Eye drops 2%1.79	5 ml OP	✓ Rexacrom
Glaucoma Preparations - Beta Blockers		
BETAXOLOL * Eye drops 0.25%	5 ml OP 5 ml OP	✓ Betoptic S ✓ Betoptic
** Eye drops 0.25% 1.81 ** Eye drops 0.5% 2.04 ** Eye drops 0.5%, gel forming 3.78	5 ml OP 5 ml OP 2.5 ml OP	✓ <u>Arrow-Timolol</u> ✓ <u>Arrow-Timolol</u> ✓ Timoptol XE
Glaucoma Preparations - Carbonic Anhydrase Inhibitors		
ACETAZOLAMIDE * Tab 250 mg17.03 BRINZOLAMIDE	100	✓ Diamox
* Eye drops 1%	5 ml OP	✓ <u>Azopt</u>
DORZOLAMIDE HYDROCHLORIDE * Eye drops 2%	5 ml OP	Trusopt
DORZOLAMIDE WITH TIMOLOL * Eye drops 2% with timolol 0.5%	5 ml OP	✓ <u>Dortimopt</u>
Glaucoma Preparations - Prostaglandin Analogues		
BIMATOPROST * Eye drops 0.03%	3 ml OP	✓ <u>Bimatoprost</u> <u>Multichem</u>

	Subsidy (Manufacturer's P \$		Fully Brand or dised Generic Manufacturer
* Eye drops 0.005%	1.82	2.5 ml OP	✓ <u>Teva</u>
* Eye drops 0.004%	9.75	2.5 ml OP	✓ <u>Travatan</u>
Glaucoma Preparations - Other			
BRIMONIDINE TARTRATE * Eye drops 0.2%	4.29	5 ml OP	✓ <u>Arrow-Brimonidine</u>
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE * Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ Arrow - Lattim
PILOCARPINE HYDROCHLORIDE # Eye drops 1%* # Eye drops 2%		15 ml OP 15 ml OP	✓ Isopto Carpine ✓ Isopto Carpine
* Eye drops 4% Subsidised for oral use pursuant to the Standard Formu	7.99	15 ml OP	✓ Isopto Carpine
* Eye drops 2% single dose – Special Authority see SA0895 below – Retail pharmacy	31.95	20 dose	✓ Minims Pilocarpine

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
ATROPINE SULPHATE		
* Eye drops 1%	15 ml OP	✓ <u>Atropt</u>
CYCLOPENTOLATE HYDROCHLORIDE * Eye drops 1%8.76	15 ml OP	✓ Cyclogyl
TROPICAMIDE	13 1111 01	• Cyclogyi
* Eye drops 0.5%7.15	15 ml OP	✓ Mydriacyl
* Eye drops 1%	15 ml OP	✓ Mydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 243		
HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	✓ Methopt
HYPROMELLOSE WITH DEXTRAN		
* Eye drops 0.3% with dextran 0.1%	15 ml OP	✓ Poly-Tears

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER - Special Authority see SA2134 above - Retail pha	armacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Author	ity see SA2134 at	ove – Retail į	pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	Systane Unit Dose
SODIUM HYALURONATE [HYALURONIC ACID] - Special Auth	ority see SA2134	above – Reta	il pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pha	armacy Procedure	s Manual rest	riction allowing one bottle per
month is not relevant and therefore only the prescribed d	losage to the near	est OP may b	e claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%4.15	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%2.17	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g	5 g OP	✓ VitA-POS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Agents Used in the Treatment of Poisonings

Antidotes

ACETYL CYSTEINE

S29 S29

(DBL Acetylcysteine Inj 200 mg per ml, 10 ml ampoule to be delisted 1 December 2022)

(Martindale Pharma S29 S29 Inj 200 mg per ml, 10 ml ampoule to be delisted 1 December 2022)

NALOXONE HYDROCHLORIDE

- a) Up to 5 ini available on a PSO
- b) Only on a PSO

Removal and Elimination

CHARCOAL

D

- - a) Up to 250 ml available on a PSO
 - b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

Tab 125 mg dispersible	276.00	28	Exjade
Tab 250 mg dispersible	552.00	28	✓ Exjade
Tab 500 mg dispersible	1,105.00	28	Exjade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and
- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels, liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis; or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE	 Special Authori 	ty see SA1480	on the next pa	age – Retail pharmacy	
Tab EOO ma				E00 17	

Tab 500 mg	533.17	100	Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox



Subsidy		Fully	Brand or	
(Manufacturer's Pric	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1480 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

DESFERRIOXAMINE MESILATE * Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	53.31	6	
* III 200 IIIg per IIII, 3 III	(156.71)	O	Calcium Disodium Versenate

Standard Formulae

ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP	1 g 70 ml
Culture Cyc Grop Buse	40	Water	to 100 ml
CODEINE LINCTUS (3 mg per 5 ml)			
Codeine phosphate	60 mg	PHENOBARBITONE SODIUM PAEDIATRIC ORAL	LIQUID (10
Glycerol Preservative	40 ml	mg per ml)	400
Water	qs to 100 ml	Phenobarbitone Sodium	400 mg 4 ml
vvaler	10 100 1111	Glycerol BP Water	to 40 ml
CODEINE LINCTUS (15 mg per 5 ml)		Water	10 40 1111
Codeine phosphate	300 mg	PILOCARPINE ORAL LIQUID	
Glycerol	40 ml	Pilocarpine 4% eye drops	qs
Preservative	qs	Preservative	qs
Water	to 100 ml	Water	to 500 ml
FOLINIC MOUTHWASH		(Preservative should be used if quantity supplied is	for more
Calcium folinate 15 mg tab	1 tab	than 5 days.)	
Preservative	qs	SALIVA SUBSTITUTE FORMULA	
Water	to 500 ml	Methylcellulose	5 g
(Preservative should be used if quantity supplied is f	or more	Preservative	qs
than 5 days. Maximum 500 ml per prescription.)		Water	to 500 ml
METUADONE MIXTUDE		(Preservative should be used if quantity supplied is	for more
METHADONE MIXTURE Methadone powder	90	than 5 days. Maximum 500 ml per prescription.)	
Glycerol	qs qs	SODIUM CHLORIDE ORAL LIQUID	
Water	to 100 ml	Sodium chloride inj 23.4%, 20 ml	qs
Water	10 100 1111	Water	qs
METHYL HYDROXYBENZOATE 10% SOLUTION		(Only funded if prescribed for treatment of hyponatra	•
Methyl hydroxybenzoate	10 g		,
Propylene glycol	to 100 ml	VANCOMYCIN ORAL SOLUTION (50 mg per ml)	
(Use 1 ml of the 10% solution per 100 ml of oral liqui	id mixture)	Vancomycin 500 mg injection	10 vials
OMEPRAZOLE SUSPENSION		Glycerol BP Water	40 ml to 100 ml
Omeprazole capules or powder	qs	(Only funded if prescribed for treatment of Clostridiu	
Sodium bicarbonate powder BP	8.4 g	following metronidazole failure)	iii uiiiicii c
Water	to 100 ml	ionowing menoridazore iandre)	

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy	-i\ 0h	Fully Brand or
	(Manufacturer's Pi \$	rice) Subs Per	sidised Generic Manufacturer
-	101		
Extemporaneously Compounded Preparations	and Galenica	IIS	
CODEINE PHOSPHATE - Safety medicine; prescriber may de		g frequency	
Powder – Only in combination		25 g	Douglas
Only in extemporaneously compounded codeine linctus	(90.09)		Douglas
COLLODION FLEXIBLE			
Note: This product is no longer being manufactured by the determined.	supplier and will b	e delisted fror	n the Schedule at a date to be
Collodion flexible	19.30	100 ml	✓ PSM
COMPOUND HYDROXYBENZOATE - Only in combination			
Only in extemporaneously compounded oral mixtures.			
Soln		100 ml	✓ Midwest
GLYCERIN WITH SODIUM SACCHARIN – Only in combination	1		
Only in combination with Ora-Plus. Suspension	30.95	473 ml	✓ Ora-Sweet SF
GLYCERIN WITH SUCROSE – Only in combination	00.00	4701111	• Old-Oweel Ol
Only in combination with Ora-Plus.			
Suspension	30.95	473 ml	✓ Ora-Sweet
GLYCEROL			
* Liquid – Only in combination		500 ml	✓ healthE Glycerol BP
Only in extemporaneously compounded oral liquid prep	arations.		
METHADONE HYDROCHLORIDE a) Only on a controlled drug form			
b) No patient co-payment payable			
c) Safety medicine; prescriber may determine dispensing f	requency		
d) Extemporaneously compounded methadone will only be	reimbursed at the	rate of the ch	eapest form available
(methadone powder, not methadone tablets). Powder	7 84	1 g	✓ AFT
METHYL HYDROXYBENZOATE	7.04	1 9	· All
Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE		ŭ	
Powder		100 g	✓ MidWest
Suspension – Only in combination		473 ml	✓ Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCI			A Ove Bland OF
Suspension		473 ml	✓ Ora-Blend SF
METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Or Suspension	•	473 ml	✓ Ora-Blend
PHENOBARBITONE SODIUM		4701111	o o o o o o o o o o o o o o o o o o o
Powder – Only in combination	52.50	10 g	✓ MidWest
	325.00	100 g	✓ MidWest
Only in children up to 12 years			
PROPYLENE GLYCOL	700to 100/ 001::::-		
Only in extemporaneously compounded methyl hydroxyben Liq		n. 500 ml	✓ Midwest
SODIUM BICARBONATE		333 1111	
Powder BP - Only in combination		500 g	✓ Midwest
Only in aytomographously compounded amongszala ar	id lanconrazola cui	enoneion	

Only in extemporaneously compounded omeprazole and lansoprazole suspension.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price) \$) Su Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio		500 ml	✓ M	lidwest	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Pri	ce)	Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia; or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

continued...

✓ fully subsidised 247

Subsidy		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	1	Manufacturer

continued...

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT – Special Authority see SA1523 on the previous page – Hospital pharmacy [HP3]

Emulsion (neutral)	12.30 200	ml OP	Calogen
	30.75 500	ml OP 🗸	Calogen
Emulsion (strawberry)	12.30 200	ml OP 🗸	Calogen
Oil	30.00 500	ml OP 🗸	MCT oil (Nutricia)
Oil, 250 ml1	14.92 4	OP 🗸	Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PROTEIN SUPPLEMENT	 Special Authority see SA1524 above – Hospital p 	narmacy [HP3]	
Powder	7.90	225 g OP	✓ Protifar
	8.95	227 g OP	✓ Resource
		•	Beneprotein

Subsidy
(Manufacturer's Price)
\$ Per

Fully Subsidised

Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid3.75	500 ml OP	✓ Glucerna Select
7.50	1,000 ml OP	Diason RTH
		✓ Nutrison Advanced
		Diason

(Diason RTH Liquid to be delisted 1 December 2022)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)	1.50	200 mi OP	✓ Diasip
Liquid (vanilla)	1.50	200 ml OP	✓ Diasip
	2.10		✓ Nutren Diabetes

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED – Special Authority see SA1525 above – Hospital pharmacy [HP3]

✓ fully subsidised 249

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Powder54.00 400 g OP

✓ Kindergen

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

Subsidy		Fully	Brand or	
(Manufacturer's Price	e) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML - Special Authority see SA Liquid		ne previous pag 500 ml OP	e – Hospital pharmacy [HP3] ✓ Nutrini Energy RTH
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see SA13 Liquid		previous page 500 ml OP	Hospital pharmacy [HP3]✓ Nutrini RTH✓ Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Aut pharmacy [HP3]	thority see	SA1379 on the	previous page – Hospital
Liquid	.6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED – Special Authority see SA1379 on the previ- Powder (vanilla)		- Hospital phari 850 g OP	macy [HP3] Pediasure
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see \$A1379 Liquid (strawberry) Liquid (vanilla)	.1.60	revious page – 200 ml OP 200 ml OP	Hospital pharmacy [HP3] ✓ Fortini ✓ Fortini
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1379 (Liquid (chocolate)	.1.07 .1.07 .1.07	vious page – Ho 200 ml OP 200 ml OP 200 ml OP 250 ml OP	ospital pharmacy [HP3] Pediasure Pediasure Pediasure Pediasure Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Authori pharmacy [HP3]	ty see SA1	1379 on the pre	vious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	.1.60 .1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	 ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 on the Powder		age – Hospital į 400 a OP	oharmacy [HP3] Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

✓ fully subsidised 251

	Subsidy (Manufacturer's F \$	Price) Subsi Per	Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		ous page – Hos 220 ml OP	✓ 1	harmacy [HP3] Nepro HP (strawberry) Nepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML – Special Authority see SA110		s page – Hospit 237 ml OP	al pha	armacy [HP3]
Liquid, 200 ml bottle	(3.31) 11.52	4 OP	١	NovaSource Renal
Liquid (apricot) 125 mlLiquid (caramel) 125 ml		4 OP 4 OP	✓ [NovaSource Renal Renilon 7.5 Renilon 7.5
(NovaSource Renal Liquid to be delisted 1 September 2022)				

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption; or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas; or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease; or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML — Special Au Liquid	•	.1377 above – H 00 ml OP ✓	
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA133 Liquid (grapefruit), 250 ml carton	71.00 71.00	18 OP 18 OP	(HP3) Elemental 028 Extra Elemental 028 Extra Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 Powder (unflavoured)			HP3] Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority se Liquid		00 ml OP	harmacy [HP3] Nutrison Advanced Peptisorb Peptisorb

(Peptisorb Liquid to be delisted 1 June 2023)

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per 🗸 Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age: and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subs (Manufactur		
\$	Per	Manufacturer

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per ✓	Manufacturer

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis: or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease: or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula: or
- 9 Severe chronic neurological conditions.

SPECIAL FOODS

	Subsidy (Manufacturer's \$		Fully Brand or dised Generic Manufacturer
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 Liquid		lospital pharmac 250 ml OP 1,000 ml OP	y [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML – Special Authority see SA1859 or Liquid		spital pharmacy 250 ml OP 1,000 ml OP	[HP3] ✓ Isosource Standard ✓ Nutrison Standard RTH ✓ Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Author Liquid	,	on page 253 – Ho 1,000 ml OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority : Liquid		oage 253 - Hosp 1,000 ml OP	ital pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority Liquid		page 253 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority Liquid		page 253 – Hos 1,000 ml OP	pital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on pa Powder (chocolate)	•	al pharmacy [HP 840 g OP	3] ✓ Sustagen Hospital Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	✓ Ensure✓ Sustagen HospitalFormula Active
	26.00	850 g OP	✓ Ensure

	Subsidy	Fully	Brand or
(1)	Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 253 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip
Liquid (fruit of the forest) – Higher subsidy of \$1.26 per 200 ml			
with Endorsement		200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) - Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement	0.85	237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML — Special Authority see SA1859 on page 253 — Hospital pharmacy [HP3]
Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Endorsement	0.72	200 ml OP	
	(1.26)	200 0.	Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with

continued...



Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
 \$	Per	✓	

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements; or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

(1.90) Two Cal HN

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	Subsidy (Manufacturer's Price) Sub	Fully sidised	Brand or Generic	
	\$	Per	1	Manufacturer	
FOOD THICKENER – Special Authority see SA1106 on the pre	vious page – Hospita	l pharmac	y [HP3]		
Powder	6.53	00 g OP	✓ N	utilis	
	7.25	80 g OP	✓ F	eed Thickener	
				Karicare Aptamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

pharmacy [HP3]	
1,000 g OP	
•	Healtheries Simple Baking Mix
oharmacy [HP3]	
1,000 g OP	
	NZB Low Gluten Bread Mix
	Horleys Bread Mix
nacy [HP3]	
2,000 g OP	
	Horleys Flour
	1,000 g OP charmacy [HP3] 1,000 g OP

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	ce) Su Per	ıbsidised •	Generic Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	orevious page – Ho	ospital pha	rmacy [H	P3]
Buckwheat Spirals	2.00	250 g OP		
	(3.11)			Orgran
Corn and Vegetable Shells	2.00	250 g OP		
	(2.92)			Orgran
Corn and Vegetable Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Lasagne Sheets	1.60	200 g OP		
	(3.82)			Orgran
Rice and Corn Macaroni	2.00	250 g OP		
	(2.92)			Orgran
Rice and Corn Penne	2.00	250 g OP		
	(2.92)			Orgran
Rice and Maize Pasta Spirals	2.00	250 g OP		
	(2.92)			Orgran
Rice and Millet Spirals	2.00	250 g OP		
	(3.11)			Orgran
Rice and corn spaghetti noodles	2.00	375 g OP		
	(2.92)			Orgran
Vegetable and Rice Spirals	2.00	250 g OP		
	(2.92)			Orgran
Italian long style spaghetti	2.00	220 g OP		
	(3.11)			Orgran

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	idised	Generic
\$	Por	1	Manufacturer

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE	 Special Authority see SA1 	108 on the pre	evious page	– Hospital
pharmacy [HP3]				
· · · ·		0-	<i>-</i>	

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (berry) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (orange) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)	320.00	500 g OP	✓ XP Maxamum
Liquid (berry)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g	1,123.20	36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml		60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

Foods

LOW PROTEIN BAKING MIX – Special Authority see SA1108 on the previous page	ge – Hospital	pharmacy [HP3]
Powder8.22	500 g OP	Loprofin Mix

OW PROTEIN PASTA - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

LOW PROTEIN PASTA - Special Authority see SATTOS on the prev	/lous page – i	nospilai priami	acy [nP3]
Animal shapes	11.91	500 g OP	✓ Loprofin
Lasagne	5.95	250 g OP	Loprofin
Low protein rice pasta	11.91	500 g OP	✓ Loprofin
Macaroni	5.95	250 g OP	✓ Loprofin
Penne	11.91	500 g OP	✓ Loprofin
Spaghetti	11.91	500 g OP	✓ Loprofin
Spirals	11.91	500 a OP	✓ Loprofin

Subsidy (Manufacturer's Price)

Fully Subsidised Brand or Generic Manufacturer

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FORMULA – Special Authority see SA2092 below – H Powder		400 g OP ✓	Alfamino Alfamino Junior
Powder (unflavoured)	53.00	400 g OP 🗸	Elecare Elecare LCP Neocate Gold Neocate Junior Unflavoured
Powder (vanilla)	53.00	400 g OP 🗸	Neocate SYNEO Elecare Neocate Junior Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis: or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency: or
- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or alleray or malabsorotion; and
 - 6.2 Either:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or

2 Both:

- 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
- 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and
 - 2.2.3 Amino acid formula is required for a nutritional deficit; and
 - 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	✓	Manufacturer

- 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time; and
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable; and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome; or
 - 2.3 Intractable diarrhoea: or
 - 2.4 Biliary atresia; or
 - 2.5 Cholestatic liver diseases causing malabsorption: or
 - 2.6 Cystic fibrosis; or
 - 2.7 Proven fat malabsorption; or
 - 2.8 Severe intestinal motility disorders causing significant malabsorption; or
 - 2.9 Intestinal failure: or
 - 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and

continued...

SYNEO 2

(A)	Subsidy //anufacturer's Price)	Subsi	Fully	Brand or Generic
ζ.,	\$	Per	✓	Manufacturer

continued...

- 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms; or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or
- 3 Short bowel syndrome; or
- 4 Intractable diarrhoea; or
- 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption: or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure: or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula: and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IqE mediated allergic reaction.

continued...



Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	Jubsiuiseu	Manufacturer

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

PAEDIATRIC ORAL/ENTERAL FEED 1 KCAL/ML − Special Authority see SA1698 below − Hospital pharmacy [HP3] Liquid.......2.35 125 ml OP ✓ Infatrini

SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: "Volume intolerant" patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)	35.50	300 g OP	✓ KetoCal 4:1
			✓ Ketocal 3:1
Powder (vanilla)	35.50	300 g OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Generic Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000

Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcqatlas.org/index.php.

Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent......0.00 10

✓ BCG Vaccine

DIPHTHERIA, TETANUS AND PERTUSSIS VACCINE - [Xpharm]

Funded for any of the following criteria:

- 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
- A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care
 Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
- A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
- 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 5) A single dose for vaccination of patients aged from 65 years old; or
- 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
- 7) For vaccination of previously unimmunised or partially immunised patients; or
- 8) For revaccination following immunosuppression; or
- 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid, 8 mcg pertussis filamentous

DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE - [Xpharm]

Funded for any of the following:

- 1) A single dose for children up to the age of 7 who have completed primary immunisation; or
- A course of four vaccines is funded for catch up programmes for children (to the age of 10 years) to complete full primary immunisation; or
- 3) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post HSCT, or chemotherapy; pre- or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
- 4) Five doses will be funded for children requiring solid organ transplantation.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

Ini 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg

pertussis toxoid, 25 mcg pertussis filamentous

haemagglutinin, 8 mcg pertactin and 80 D-antigen units

NATIONAL IMMUNISATION SCHEDULE			
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN	ND HAEMOPHILUS I	NFLUENZAE TY	PE B VACCINE -
[Xpharm]			
Funded for patients meeting any of the following criteria:			
Up to four doses for children up to and under the age of			and a considerable and a second
2) An additional four doses (as appropriate) are funded for			
10 who are patients post haematopoietic stem cell trans post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up p		'	
to complete full primary immunisation. Please refer to the Imi			
programmes.			
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10 ✓ <u>In</u>	<u>ifanrix-hexa</u>
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)imi			
transplantation, or chemotherapy; functional asplenic; pr			olid organ transplant, pre-
or post cochlear implants, renal dialysis and other sever	, , , , , , , , , , , , , , , , , , , ,	•	val madiaina physisian ar
 For use in testing for primary immunodeficiency disease paediatrician. 	s, on the recommend	allon of an intern	iai medicine physician or
paediatrician.			
Haemophilus Influenzae type B polysaccharide 10 mcg			
conjugated to tetanus toxoid as carrier protein 20-40 mcg	:		
prefilled syringe plus vial 0.5 ml		1 🗸 H	iberix
HEPATITIS A VACCINE - [Xpharm]			
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
2) Two vaccinations for use in children with chronic liver di	sease; or		
3) One dose of vaccine for close contacts of known hepatit	is A cases.		

Inj 1440 ELISA units in 1 ml syringe......0.00

✓ Havrix

✓ Havrix Junior

1

		NATIONAL	IIVIIVI	UNISATI	ON SCHEDULE
		Subsidy (Manufacturer's Price)	Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B	RECOMBINANT VACCINE - [Xpharm]				
	g per 0.5 ml prefilled syringe		1	✓ E	ngerix-B
	led for patients meeting any of the following criteria:				
	for household or sexual contacts of known acute h				s; or
	for children born to mothers who are hepatitis B su				and the second are second to the
3)	for children up to and under the age of 18 years in				achieved a positive
4)	serology and require additional vaccination or requ for HIV positive patients; or	life a primary course of	or vacc	ination, or	
	for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	course: or			
	for patients following immunosuppression; or	,			
8)	for solid organ transplant patients; or				
	for post-haematopoietic stem cell transplant (HSC	T) patients; or			
10)	following needle stick injury.				
Inj 20 mc	g per 1 ml prefilled syringe	0.00	1	√ E	ngerix-B
	led for patients meeting any of the following criteria:			_	<u> </u>
1)	for household or sexual contacts of known acute h	epatitis B patients or h	nepatit	is B carriers	s; or
	for children born to mothers who are hepatitis B su				
3)	for children up to and under the age of 18 years in				achieved a positive
4\	serology and require additional vaccination or requ	ire a primary course o	of vacc	cination; or	
	for HIV positive patients; or for hepatitis C positive patients; or				
	for patients following non-consensual sexual interc	ourse. or			
	for patients following immunosuppression; or	ouise, or			
,	for solid organ transplant patients; or				
	for post-haematopoietic stem cell transplant (HSC	T) patients; or			
	following needle stick injury; or				
	for dialysis patients; or				
12)	for liver or kidney transplant patients.				
HUMAN PAP	ILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, 52 AND 5	58) VACCINE [HPV] -	- [Xph	arml	
	e following:	,	٠,	•	
1) Max	ximum of two doses for children aged 14 years and	under; or			
	ximum of three doses for patients meeting any of th				
) People aged 15 to 26 years inclusive; or				
2) Either:				
	People aged 9 to 26 years inclusive				
	1) Confirmed HIV infection; or				
a\ •-	2) Transplant (including stem cell) patients: or				
3) Max	ximum of four doses for people aged 9 to 26 years i	nclusive post chemoth	nerapy	<i>'</i>	
Inj 270 m	cg in 0.5 ml syringe	0.00	10	√ <u>G</u>	ardasil 9

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

✓ Afluria Quad Junior - [Xpharm].......11.00 (2022 formulation)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes: or
- iv) have chronic renal disease: or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV. or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness:

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	110.00	10	1	Afluria Quad
				(2022 formulation)

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

C)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or

vii) are pregnant; or

- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 July 2022 to 31 December 2022;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

(Manu	Subsidy	Fully	Brand or
	facturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of diluent 0.5 ml112.50

....112.50 5 **MMR II**250.00 10 **Priorix**

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

NATIONAL IMMUNISATION SCHEDULE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients: or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B. 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10

Synflorix

Subsidy			Brand or	
(Manufacturer's Price)			Generic	
	Per	✓	Manufacturer	

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Ini 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4.

	NATIONAL	IMMUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE - Either:	- [Xpharm]		
 Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with func complement deficiency (acquired or inherited), cochl All of the following: 	tional asplenia, pre- or p	post-solid organ t	ransplant, renal dialysis,
 a) Patient is a child under 18 years for (re-)immur b) Treatment is for a maximum of two doses; and c) Any of the following: 	isation; and		
i) on immunosuppressive therapy or radiation immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or	or	·	
or vi) with cochlear implants or intracranial shur vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more prednisone of 2 mg/kg per day or greater	than two weeks, and wh		
20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks ge xi) with cardiac disease, with cyanosis or fail xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	estation; or ure; or	gh-dose corticost	eroid therapy); or
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	ng: ndividuals; or	_	neumovax 23
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe ROTAVIRUS ORAL VACCINE – [Xpharm]		tch-up programm 1 ✓ <u>II</u>	
Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 1- 2) no vaccination being administered to children aged 2			
Oral susp live attenuated human rotavirus			

10

✓ Rotarix

1,000,000 CCID50 per dose, prefilled oral applicator......0.00

	Subsidy (Manufacturer's Price) \$	Subsid Per	Fully lised •	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] - [Xpharm]	*			
Either:				
1) Maximum of one dose for primary vaccination for eith	er:			
a) Any infant born on or after 1 April 2016; or				
b) For previously unvaccinated children turning 11 varicella infection (chickenpox), or	years old on or after 1	July 2017, \	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
a) Any of the following for non-immune patients:				
 i) with chronic liver disease who may in futur ii) with deteriorating renal function before trans 		ansplantatior	n; or	
iii) prior to solid organ transplant; or				
iv) prior to any elective immunosuppression*,v) for post exposure prophylaxis who are imm		onte : or		
b) For patients at least 2 years after bone marrow			snecial	list or
c) For patients at least 6 months after completion of				
d) For HIV positive non immune to varicella with m				
e) For patients with inborn errors of metabolism at varicella, or				
f) For household contacts of paediatric patients where the household continuous where the household continuous.	act has no clinical hist	ory of varicel	lla, or	
g) For household contacts of adult patients who ha immunocompromised, or undergoing a procedul has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosuppr	essive therapy must b	e for a treatn	nent p	eriod of greater than
28 days				
Inj 1350 PFU prefilled syringe	0.00	1		<u>arivax</u>
		10	_	<u>arivax</u>
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUAT Funded for patients meeting the following criteria:	ED VACCINE [SHING	GLES VACCI	NE] -	-[Xpharm]
 One dose for all people aged 65 years 				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1	J 70	ostavax

- Symbols -	Albendazole	89	Anagrelide hydrochloride	149
UK Synacthen80	Albey	227	Analgesics	120
3TC105	Albustix	77	Anastrozole	166
- A -	Alchemy Oxybutynin	76	Anatrole	166
A-Scabies67	Aldurazyme	29	Andriol Testocaps	80
Abacavir sulphate105	Alecensa	156	Androderm	80
Abacavir sulphate with	Alectinib	156	Anoro Ellipta	231
lamivudine 105	Alendronate sodium	111	Antabuse	142
Abiraterone acetate163	Alendronate sodium with		Antacids and Antiflatulents	6
Acarbose11	colecalciferol	111	Anthelmintics	89
Accarb11	Alfacalcidol	33	Antiacne Preparations	61
Accuretic 1048	Alfamino	262	Antiallergy Preparations	227
Accuretic 2048	Alfamino Junior	262	Antianaemics	37
Acetazolamide238	Alginic acid	6	Antiandrogen Oral	
Acetec47	Alglucosidase alfa		Contraceptives	75
Acetic acid with hydroxyquinoline and	Alkeran		Antiarrhythmics	
ricinoleic acid75	Alkeran S29	146	Antibacterials	
Acetylcysteine241	Allersoothe		Antibacterials Topical	
Aci-Jel75	Allmercap	148	Anticholinergic Agents	
Aciclovir	Allopurinol		Anticholinesterases	
Infection 100	Alpha-Adrenoceptor Blockers		Antidepressants	
Sensory236	Alpha-Keri Lotion		Antidiarrhoeals	
Acidex6	Alphamox		Antiepilepsy Drugs	
Acipimox54	Alphamox 125		Antifibrinolytics, Haemostatics and	
Acitretin68	Alphamox 250		Local Sclerosants	
Aclasta114	Alprolix		Antifibrotics	
Actemra215	Alu-Tab		Antifungals	
Actemra S29215	Aluminium hydroxide		Antifungals Topical	
Actinomycin D	Alvogen		Antihistamines	
Actrapid10	Amantadine hydrochloride		Antihypotensives	
Actrapid Penfill	Ambrisentan		Antimalarials	
Acupan120	Ambrisentan Mylan		Antimigraine Preparations	
Adalimumab (Amgevita)	Amgevita		Antinausea and Vertigo Agents	
Adalimumab (Humira)	Amiloride hydrochloride		Antipruritic Preparations	
Adapalene	Amiloride hydrochloride with		Antipsychotics	
ADR Cartridge 1.823	furosemide	53	Antiretrovirals	
Adrenaline57	Amiloride hydrochloride with		Antirheumatoid Agents	
Advantan	hydrochlorothiazide	53	Antispasmodics and Other Agents	
Advate41	Aminophylline		Altering Gut Motility	
Adynovate41	Amiodarone hydrochloride		Antithrombotic Agents	
Afinitor223	Amisulpride		Antithymocyte globulin	
Aflibercept187	Amisulpride Mylan		(equine)	173
Afluria Quad	Amitriptyline		Antitrichomonal Agents	
(2022 formulation)	Amlodipine		Antituberculotics and	
Afluria Quad Junior	Amneal		Antileprotics	gc
(2022 formulation)	Amorolfine		Antiulcerants	
AFT-Pyrazinamide100	Amoxicillin		Antivirals	
Agents Affecting the	Amoxicillin with clavulanic acid		Anxiolytics	
Renin-Angiotensin System 47	Amphotericin B		Anzatax	159
Agents for Parkinsonism and Related	Amsacrine		Apidra	
Disorders 118	AmsaLyo		Apidra SoloStar	
Agents Used in the Treatment of	Amsidine		APO-Atomoxetine	
Poisonings241	Amzoate		APO-Atomoxetine S29	
Agrylin	Anaesthetics		Apo-Azithromycin	
, w. y	/ u/acou/ouco	113	THO TIEILIIOITIYOIII	50

Apo-Diltiazem CD	52	Atrovent	230	Betahistine dihydrochloride	12
Apo-Doxazosin	47	AU Synacthen	80	Betaine	2
Apo-Prednisone	80	Aubagio	135	Betaloc CR	5
Apo-Primidone	127	Augmentin	92	Betamethasone dipropionate	6
Apo-Temozolomide		Aurorix	124	Betamethasone dipropionate with	
Apomorphine hydrochloride	118	AutoSoft 30	23	calcipotriol	6
Aprepitant		AutoSoft 90	23	Betamethasone sodium phosphate	е
Apresoline		Avelox	94	with betamethasone acetate	7
Aptamil AllerPro SYNEO 1		Avonex		Betamethasone valerate	63, 6
Aptamil AllerPro SYNEO 2		Avonex Pen	135	Betamethasone valerate with sodi	um
Aptamil Gold+ Pepti Junior	265	Azacitidine	146	fusidate [fusidic acid]	6
Aqueous cream	65	Azacitidine Dr Reddy's	146	Betaxolol	
Aratac	49	Azamun	166	Betnovate	6
Arava	111	Azathioprine	166	Betoptic	23
Arginine	27	Azilect	118	Betoptic S	23
Aripiprazole		Azithromycin	90	Bexsero	27
Aripiprazole Sandoz		Azopt	238	Bezafibrate	
Aristocort		AZT		Bezalip	5
Arrotex-Prazosin S29		- B -		Bezalip Retard	5
Arrow - Lattim	239	B-D Micro-Fine	15	Bicalutamide	
Arrow-Amitriptyline		B-D Ultra Fine	16	Bicillin LA	
Arrow-Bendrofluazide	54	B-D Ultra Fine II	16	BiCNU	14
Arrow-Brimonidine		Bacillus Calmette-Guerin (BCG		Bicnu Heritage	14
Arrow-Diazepam		vaccine		Bile and Liver Therapy	
Arrow-Doxorubicin		Bacillus Calmette-Guerin		Biltricide	
Arrow-Losartan &		vaccine	267	Bimatoprost	
Hydrochlorothiazide	48	Baclofen		Bimatoprost Multichem	
Arrow-Norfloxacin		Bactroban		Binarex	
Arrow-Ornidazole	98	Balance	29	Binocrit	3
Arrow-Quinapril 10		Barrier Creams and Emollients.	65	Biodone	12
Arrow-Quinapril 20		BCG Vaccine	267	Biodone Extra Forte	12
Arrow-Quinapril 5		Beclazone 100	228	Biodone Forte	12
Arrow-Roxithromycin		Beclazone 250	228	Bisacodyl	
Arrow-Timolol		Beclazone 50		Bisoprolol fumarate	5
Arrow-Topiramate		Beclomethasone dipropionate		Bisoprolol Mylan	
Arrow-Tramadol		Bee venom allergy treatment		BK Lotion	
Arsenic trioxide		Bendamustine hydrochloride		Bleomycin sulphate	
Asacol		Bendrofluazide		Blood Colony-stimulating	
Ascorbic acid		Bendroflumethiazide		Factors	4
Aspen Adrenaline		[Bendrofluazide]	54	Blood glucose diagnostic test	
Aspirin		Benzathine benzylpenicillin		meter	1
Blood	41	Benzatropine mesylate		Blood glucose diagnostic test	
Nervous	120	Benzbromaron AL 100		strip	1
Asthalin	230	Benzbromarone	116	Blood glucose test strips (visually	
Atazanavir sulphate	105	Benztrop	119	impaired)	1
Atenolol	50	Benzydamine hydrochloride	32	Blood Ketone Diagnostic Test	
Atenolol AFT		Benzylpenicillin sodium [Penicil		Strip	1
Atenolol AFT S29		G]		Bonjela	3
ATGAM	173	Beta Cream		Boostrix	
Ativan		Beta Ointment		Bortezomib	
Atomoxetine		Beta Scalp		Bortezomib Dr Reddy's S29	
Atorvastatin	54	Beta-Adrenoceptor Agonists		Bortezomib Dr-Reddy's	
Atropine sulphate	-	Beta-Adrenoceptor Blockers		Bortezomib Juno	
Cardiovascular	49	Betadine		Bortezomib Juno S29	
Sensory		Betadine Skin Prep		Bosentan	
Atropt		Betaferon		Bosentan Dr Reddy's	
•				•	

Bplex33	Capsaicin	Chlorthalidone5
Breo Ellipta229	Musculoskeletal111	Chlorvescent4
Brevinor 1/2874	Nervous120	Choice Load 3757
Bricanyl Turbuhaler230	Captopril47	Choice TT380 Short7
Brilinta41	Carafate9	Choice TT380 Standard7
Brimonidine tartrate239	Carbaccord145	Choline salicylate with cetalkonium
Brimonidine tartrate with timolol	Carbamazepine125	chloride3
maleate239	Carbimazole83	Ciclosporin22
Brinzolamide238	Carbomer240	Cilazapril4
Brolene236	Carboplatin145	Cilicaine9
Bromocriptine mesylate118	Carboplatin Ebewe145	Cilicaine VK9
Brufen SR110	Carbosorb-X241	Cinacalcet7
Buccastem130	Cardinol LA51	Cipflox9
Budesonide	Cardizem CD52	Ciprofloxacin
Alimentary6	CareSens Dual14	Infection9
Respiratory228, 234	CareSens N14–15	Sensory23
Budesonide with eformoterol229	CareSens N POP14	Ciprofloxacin Teva23
Burnetanide52	CareSens N Premier14	Cisplatin14
Buprenorphine Naloxone BNM141	CareSens PRO15	Cisplatin Ebewe14
Buprenorphine with naloxone141	Carmellose sodium with gelatin and	Citalopram hydrobromide12
Bupropion hydrochloride141	pectin32	Cladribine14
Burinex52	Carmustine 145	Clarithromycin
Burinex S2952	Carvedilol50	Alimentary
Buscopan8	Carvedilol Sandoz50	Infection9
Buspirone hydrochloride133	Casirivimab and imdevimab188	Clexane4
Buspirone Viatris133	Catapres52	Clexane Forte4
Busulfan145	CeeNU146	Climara8
- C -	Cefaclor monohydrate89	Clindamycin9
Cabergoline88	Cefalexin89	Clinicians2
Caffeine citrate235	Cefalexin Sandoz89	Clinicians Renal Vit3
Calamine63	Cefazolin89	Clobazam12
Calamine-AFT63	Ceftriaxone89	Clobetasol propionate63, 6
Calci-Tab 50034	Ceftriaxone-AFT89	Clobetasone butyrate6
Calcipotriol68	Cefuroxime axetil89	Clofazimine9
Calcitonin78	Celebrex110	Clomazol
Calcitriol33	Celecoxib110	Dermatological6
Calcitriol-AFT33	Celecoxib Pfizer110	Genito-Urinary7
Calcium 500 mg Hexal34	Celestone Chronodose79	Clomifene citrate8
Calcium carbonate	Cellcept166	Clomipramine hydrochloride12
Calcium Channel Blockers51	Centrally-Acting Agents52	Clomipramine Teva12
Calcium Disodium Versenate242	Cephalexin ABM89	Clonazepam 126, 13
Calcium folinate147	Cetirizine hydrochloride228	Clonidine5
Calcium Folinate Ebewe147	Cetomacrogol65	Clonidine BNM5
Calcium Folinate Sandoz147	Cetomacrogol with glycerol65	Clonidine hydrochloride5
Calcium Folinate Sandoz S29 147	Cetomacrogol-AFT65	Clonidine Teva5
Calcium gluconate34	Cetuximab188	Clopidogrel4
Calcium Homeostasis78	Charcoal241	Clopidogrel Multichem4
Calcium polystyrene sulphonate45	Chemotherapeutic Agents144	Clopine13
Calcium Resonium45	Chickenpox vaccine276	Clopixol131, 13
Calogen	Chlorafast236	Clotrimazole
Candesartan cilexetil48	Chlorambucil145	Dermatological6
Candestar48	Chloramphenicol236	Genito-Urinary7
Canesten	Chlorothiazide54	Clozapine13
Capecitabine147	Chlorpheniramine maleate228	Clozaril13
Capercit	Chlorpromazine hydrochloride130	Co-trimoxazole9
Capoten47	Chlortalidone [Chlorthalidone]54	Coal tar6

Coal tar with allantoin, menthol,	Cyclorin	99	Denosumab	11
phenol and sulphur6	8 Cycloserine	99	Deolate	97
Coal tar with salicylic acid and	Cyproterone acetate	80	Deoxycoformycin	153
sulphur6	8 Cyproterone acetate with		Depo-Medrol	80
Coco-Scalp6	8 ethinyloestradiol	75	Depo-Provera	74
Codeine phosphate	Cystadane		Depo-Testosterone	80
Extemporaneous24	4 Cytarabine	147	Deprim	96
Nervous12			Dermol	
Coenzyme Q102	8 Cytoxan	145	Desferrioxamine mesilate	
Colchicine11			Desmopressin	8
Colecalciferol3	3 D-Penamine	111	Desmopressin acetate	
Colestid5	4 Dabigatran	44	Desmopressin-PH&T	
Colestipol hydrochloride5			Desuric	116
Colgout11	6 Dacarbazine APP	150	Detection of Substances in	
Colifoam			Urine	7
Colistin sulphomethate9		68	Dexamethasone	
Colistin-Link9			Hormone	79
Collodion flexible24		63	Sensory	
Colloidal bismuth subcitrate	9 Dalacin C	94	Dexamethasone phosphate	79
Colofac		117	Dexamethasone Phosphate	
Coloxyl2			Panpharma	
Combigan23			Dexamethasone with framycetin	
Compound electrolytes4	5 Daonil		gramicidin	236
Compound electrolytes with glucose	Dapa-Tabs		Dexamethasone with neomycin	
[Dextrose]4			sulphate and polymyxin B	
Compound hydroxybenzoate24	4 Daraprim	95	sulphate	
Comtan11			Dexamfetamine sulfate	
Concerta13	,		Dexmethsone	79
Condoms7			Dextrochlorpheniramine	
Condyline7			maleate	
Contraceptives - Hormonal7			Dextrose	
Contraceptives - Non-hormonal7		egnancy	DHC Continus	122
Copaxone13			Diabetes	
Corticosteroids and Related Agents	DBL Acetylcysteine		Diabetes Management	
for Systemic Use7			Diacomit	
Corticosteroids Topical6			Diagnostic Agents	
Cortifoam			Diamide Relief	
Cosentyx21			Diamox	
Cosmegen 15			Diasip	
Coumadin4			Diason RTH	
Country Life2			Diazepam1	
Coversyl4			Diazoxide	
Creon 100002			Dibenzyline	
Creon 250002		75	Diclofenac Sandoz	110
Creon Micro2			Diclofenac sodium	
Crotamiton6			Musculoskeletal	
Crystaderm6			Sensory	
Curam9			Differin	
Curam Duo 500/1259			Difflam	32
Cvite			Diflucan	
Cyclizine hydrochloride12			Digestives Including Enzymes	
Cyclizine lactate12	9 DBL Pethidine Hydrochloride		Digoxin	49
Cyclogyl23	9 DBL Vincristine Sulfate		Dihydrocodeine tartrate	
Cyclonex14			Dilantin	
Cyclopentolate hydrochloride23			Dilantin Infatab	
Cyclophosphamide14	5 Deferiprone	241	Diltiazem hydrochloride	52

Dimethicone65–66	Duride	Ensure Plus RTH25
Dimethyl fumarate134	-E-	Ensure Two Cal HN RTH25
Dipentum7	e-chamber La Grande234	Entacapone11
Diphtheria, tetanus and pertussis	e-chamber Mask234	Entecavir10
vaccine267	e-chamber Turbo234	Entecavir Sandoz10
Diphtheria, tetanus, pertussis and	E-Mycin91	Entocort CIR
polio vaccine267	e5 Pharma10	Entresto 24/264
Diphtheria, tetanus, pertussis, polio,	Ear Preparations236	Entresto 49/514
hepatitis B and haemophilus	Ear/Eye Preparations236	Entresto 97/1034
influenzae type B vaccine 268	Easiphen Liquid261	Epilim12
Diprosone63	Econazole nitrate62	Epilim Crushable12
Diprosone OV63	Efavirenz104	Epilim IV12
Dipyridamole41	Efavirenz with emtricitabine and	Epilim S/F Liquid12
Disopyramide phosphate49	tenofovir disoproxil105	Epilim Syrup 12
Disulfiram142	Eformoterol fumarate229	Epirubicin Ebewe15
Diuretics52	Eformoterol fumarate dihydrate 229	Epirubicin hydrochloride15
Docetaxel150	Eftrenonacog alfa [Recombinant	Eplerenone5
Docetaxel Accord150	factor IX]	Epoetin alfa3
Docetaxel Sandoz150	Efudix70	Epoprostenol6
Docusate sodium25	Egopsoryl TA68	Eptacog alfa [Recombinant factor
Docusate sodium with	Elaprase28	VIIa]4
sennosides25	Eldepryl118	ERA9
Dolutegravir106	Elecare262	Erbitux
Domperidone 129	Elecare LCP262	Ergometrine maleate7
Donepezil hydrochloride140	Electral45	Erlotinib15
Donepezil-Rex140	Elelyso31	Erythrocin IV9
Dornase alfa233	Elemental 028 Extra252	Erythromycin (as lactobionate)9
Dortimopt238	Elidel68	Erythromycin ethyl succinate9
Dorzolamide hydrochloride238	Elocon64	Erythromycin stearate9
Dorzolamide with timolol238	Elocon Alcohol Free64	Esbriet23
Dostinex88	Eltrombopag38	Escitalopram12
Dosulepin [Dothiepin]	Eltroxin83	Escitalopram (Ethics)12
hydrochloride124	EMB Fatol99	Eskazole8
Dosulepin Mylan124	Emend Tri-Pack129	Essential Generics5
Dothiepin	Emicizumab39	Essential Prednisolone
Doxazosin47	EMLA120	Estradot8
Doxazosin Clinect47	Empagliflozin13	Estradot 50 mcg8
Doxine93	Empagliflozin with metformin	Estrofem8
Doxorubicin Ebewe151	hydrochloride13	Etanercept16
Doxorubicin hydrochloride 151	Emtricitabine105	Ethambutol hydrochloride99
Doxycycline93	Emtricitabine with tenofovir	Ethics Aspirin12
DP Lotion65	disoproxil 102	Ethics Aspirin EC4
DP Lotn HC64	Emtriva105	Ethics Lisinopril4
DP-Allopurinol116	Emulsifying ointment65	Ethinyloestradiol8
Dr Reddy's Omeprazole9	Emulsifying Ointment ADE65	Ethinyloestradiol with
Drofate51	Enalapril maleate47	desogestrel
Drugs Affecting Bone	Enbrel166	Ethinyloestradiol with
Metabolism 111	Endocrine Therapy163	levonorgestrel7
Dual blood glucose and blood ketone	Endoxan145	Ethinyloestradiol with
diagnostic test meter 14	Engerix-B269	norethisterone74
Dulaglutide12	Enlafax XR125	Ethosuximide12
Dulcolax SP Drop26	Enoxaparin sodium43	Etopophos15
Duocal Super Soluble Powder247	Enstilar68	Etoposide15
Duolin230	Ensure	Etoposide phosphate15
Duolin HFA230	Ensure Plus257	Etravirine10
DuoResp Spiromax229	Ensure Plus HN256	Eumovate6

Everet	126	Fleet Phosphate Enema	26	Fucidin	9
Everolimus	223	Flixonase Hayfever & Allergy .	234	Fucithalmic	23
Evista	112	Flixotide	229	Fulvestrant	16
Evusheld	215	Flixotide Accuhaler	229	Fungilin	3
Exemestane		Florinef		Furosemid-Ratiopharm	
Exjade	241	Fluanxol	132	Furosemide [Frusemide]	
Extemporaneously Compounded		Flucil	92	Furosemide-Baxter	
Preparations and		Flucloxacillin	92	fusidic acid	
Galenicals	244	Flucloxacillin-AFT	92	Dermatological	62, 6
Eye Preparations	236	Flucloxin	92	Infection	
Eylea	187	Flucon	237	Sensory	23
Ezetimibe		Fluconazole		- G -	
Ezetimibe Sandoz	55	Fludara Oral	147	Gabapentin	12
Ezetimibe with simvastatin	56	Fludarabine Ebewe	147	Gacet	
-F-		Fludarabine phosphate	147	Galsulfase	2
Factor eight inhibitor bypassing		Fludrocortisone acetate		Galvumet	1
fraction	40	Fluids and Electrolytes		Galvus	1
Famotidine	9	Flumetasone pivalate		Gardasil 9	26
Famotidine Hovid	9	Fluocortolone caproate with		Gastrodenol	
Faslodex		fluocortolone pivalate and		Gaviscon Double Strength	
Fatty Cream AFT		cinchocaine	8	Gaviscon Infant	
Febuxostat		Fluorometholone		Gazyva	
Febuxostat multichem		Fluorouracil	148	Gefitinib	
Feed Thickener Karicare		Fluorouracil Accord		GEM Aqueous Cream	6
Aptamil	259	Fluorouracil sodium		Gemcitabine Ebewe	
FEIBA NF		Fluox	125	Gemcitabine hydrochloride	14
Felo 10 ER		Fluoxetine hydrochloride		Gemtuzumab ozogamicin	
Felo 5 ER		Flupenthixol decanoate		Genoptic	
Felodipine	51	Flutamide		Gentamicin sulphate	
Femme-Tab ED		Flutamin	164	Infection	9
Fenpaed 100 mg per 5 ml		Fluticasone	229	Sensory	
Fentanyl		Fluticasone furoate with		Gilenya	13
Fentanyl Sandoz		vilanterol	229	Ginet	
Ferinject		Fluticasone propionate		Glatiramer acetate	
Ferodan		Fluticasone with salmeterol		Glecaprevir with pibrentasvir	
Ferriprox		FML	237	Glibenclamide	
Ferro-F-Tabs		Foban	62	Gliclazide	
Ferro-tab		Folic acid		Glipizide	1
Ferrograd	35	Folic Acid multichem		Glivec	
Ferrosig		Folic Acid Mylan		Glizide	1
Ferrous fumarate		Food Thickeners		Glucagen Hypokit	1
Ferrous fumarate with folic acid	35	Foods And Supplements For I	nborn	Glucagon hydrochloride	
Ferrous sulfate	35	Errors Of Metabolism		Glucerna Select	
Fexofenadine hydrochloride	228	Foradil	229	Glucose [Dextrose]	
Fibro-vein	41	Forteo	113	Gluten Free Foods	
Filgrastim		Fortini	251	Glycerin with sodium saccharin.	
Finasteride		Fortini Multi Fibre		Glycerin with sucrose	
Fingolimod	135	Fortisip	257	Glycerol	
Firazyr		Fortisip Multi Fibre		Alimentary	2
Flagyl		Fosamax		Extemporaneous	
Flagyl-S		Fosamax Plus		Glyceryl trinitrate	
Flamazine		Framycetin sulphate		Alimentary	
Flecainide acetate		Frisium		Cardiovascular	
Flecainide BNM		Frumil		Glycopyrronium	
Flecainide Controlled Release	-	Frusemide		Glycopyrronium bromide	
Teva	49	Fucicort		Glycopyrronium with	

indacaterol2	31	Humulin R	10	Indacaterol	229
Go Healthy		Hyaluronic acid		Indapamide	
Gold Knight		Hydralazine		Infanrix IPV	
Gold Knight XL72–		Hydralazine hydrochloride		Infanrix-hexa	
Goserelin		Hydrocortisone		Infant Formulae	
Gutron		Dermatological	64	Infatrini	
Gynaecological Anti-infectives		Hormone		Infliximab	
- H -		Hydrocortisone (PSM)		Influenza vaccine	
Habitrol1	42	Hydrocortisone acetate		Inhaled Corticosteroids	
Haemophilus influenzae type B	_	Hydrocortisone acetate with		Inhaled Long-acting	
vaccine2	68	pramoxine hydrochloride	7	Beta-adrenoceptor Agonists	22
Haldol1		Hydrocortisone and paraffin liquid		Inspra	
Haldol Concentrate1		and lanolin	. 64	Instillagel Lido	
Haldol Decanoas1		Hydrocortisone butyrate64		Insulin aspart	
Haloperidol1		Hydrocortisone with cinchocaine		Insulin aspart with insulin aspart	
Haloperidol decanoate1		Hydrocortisone with miconazole		protamine	10
Harvoni1		Hydrocortisone with natamycin and		Insulin glargine	
Havrix2		neomycin	64	Insulin glulisine	
Havrix Junior2		Hydrogen peroxide		Insulin isophane	
Haylor syrup2		Hydroxocobalamin		Insulin isophane with insulin	! \
healthE Dimethicone 10%		Hydroxocobalamin Mercury	00	neutral	10
healthE Dimethicone 4% Lotion		Pharma	33	Insulin lispro	
healthE Dimethicone 5%		Hydroxocobalamin Panpharma		Insulin lispro with insulin lispro	
healthE Glycerol BP2		hydroxycarbamide		protamine	41
healthE Urea Cream		Hydroxychloroquine		Insulin neutral	
Healtheries Simple Baking Mix2		Hydroxyurea		Insulin pen needles	
Hemastix		[hydroxycarbamide]	151	Insulin pump	
Hemlibra		Hygroton		Insulin pump cartridge	۱۰۰۰۰۰
Heparin sodium		Hylo-Fresh		Insulin pump infusion set (steel	2
Heparinised saline		Hymenoptera		cannula)	9
Heparon Junior2		Hyoscine butylbromide		Insulin pump infusion set (steel	2
Hepatitis A vaccine2		Hyoscine hydrobromide		cannula, straight insertion)	9
Hepatitis B recombinant	00	Hypam		Insulin pump infusion set (teflon	
vaccine2	60	Hyperuricaemia and Antigout		cannula)	2
Herceptin2		Hypromellose		Insulin pump infusion set (teflon	21
Hiberix2		Hypromellose with dextran		cannula, angle insertion with	
Hiprex1		- -	200	insertion device)	2
Histaclear2		lbiamox	92	Insulin pump infusion set (teflon	2
Histafen2		Ibrance		cannula, angle insertion)	2
Holoxan1		Ibuprofen		Insulin pump infusion set (teflon	2
Horleys Bread Mix2		Icatibant		cannula, straight insertion with	
Horleys Flour2		Idarubicin hydrochloride		insertion device)	2:
Hormone Replacement Therapy -	•	Idursulfase		Insulin pump infusion set (teflon	
Systemic	81	Ifosfamide		cannula, straight insertion)	2:
HPV2		Igroton		Insulin pump reservoir	
Humalog		Ikorel		Insulin syringes, disposable with	
Humalog Mix 25		lloprost		attached needle	1/
Humalog Mix 50		Imatinib mesilate		Intelence	
Human papillomavirus (6, 11, 16, 18,		Imatinib-Rex		Interferon beta-1-alpha	
31, 33, 45, 52 and 58) vaccine		Imigran		Interferon beta-1-beta	12
[HPV]2	69	Imipramine hydrochloride		Intra-uterine device	
Humatin		Imiquimod		Invega Sustenna	
Humira1		Immune Modulators		IPCA-Frusemide	10i
HumiraPen1		Immunosuppressants		IPCA-Metoprolol	51
Humulin 30/70		Imuran		IPCA-Propranolol	
Humulin NPH		Incruse Ellipta		IPOL	
FIGHTURE IN TELESCOPE	īŪ	11 101 u3€ L111µta	201	II ∨∟	413

Ipratropium bromide	.230, 234	Klacid		Levomepromazine	
Iressa	158	Alimentary	9	hydrochloride	13
Irinotecan Actavis 100	148	Infection	90	Levonorgestrel	
Irinotecan hydrochloride	148	Kliogest	82	Genito-Urinary	74–7
Irinotecan-Rex	148	Kliovance	82	Hormone	
Iron (as ferric carboxymaltose)	35	Kogenate FS	41	Levothyroxine	8
Iron polymaltose	35	Konakion MM	41	Lidocaine [Lignocaine]	
Isentress		Konsyl-D		Lidocaine [Lignocaine]	
Isentress HD	106	Kuvan	30	hydrochloride	12
Ismo 20	57	-L-		Lidocaine [Lignocaine] with	
Ismo 40 Retard	57	Labetalol	50	chlorhexidine	12
Isoniazid	99	Lacosamide		Lidocaine [Lignocaine] with	
Isoniazid with rifampicin	99	Lactulose	<mark>26</mark>	prilocaine	12
Isoptin		Laevolac		Lidocaine-Baxter	
Isoptin Retard	52	Lagevrio	103	Lidocaine-Claris	12
Isoptin SR		Lamictal		Life Extension	
Isopto Carpine		Lamivudine		Lignocaine	
Isosorbide mononitrate		Lamivudine Alphapharm	105	Lioresal Intrathecal	
Isosource Standard		Lamotrigine		Lipid-Modifying Agents	
Isotretinoin		Lamprene		Liquigen	
Ispaghula (psyllium) husk		Lanoxin		Lisinopril	
Itch-Soothe		Lanoxin Paediatric Elixir		Litak	
Itraconazole		Lanoxin PG		Lithium carbonate	
Itrazole		Lanoxin S29		Livostin	
lvacaftor		Lansoprazole		LMX4	
lvermectin		Lantus		Locacorten-Viaform ED's	
- J -		Lantus SoloStar		Local preparations for Anal ar	
Jadelle	74	Lanvis		Rectal Disorders	
Jakavi	161	Lanzol Relief		Locasol	
Jardiamet	13	Lapatinib ditosylate	159	Locoid	64, 6
Jardiance	13	Largactil		Locoid Crelo	6
Jaydess	82	Laronidase	<mark>29</mark>	Locoid Lipocream	6
Jevity HiCal RTH		Lasix	53	Locorten-Vioform	
Jevity Plus		Latanoprost	239	Lodoxamide	23
Jevity RTH		Latanoprost with timolol		Logem	
Juno Pemetrexed		Lax-Suppositories		Lomide	
- K -		Laxatives		Lomustine	14
Kadcyla	220	Laxsol	25	Loniten	
Kaletra		Ledipasvir with sofosbuvir.		Loperamide hydrochloride	
Kalydeco		Leflunomide		Lopinavir with ritonavir	
Kemadrin		Lenalidomide		Lopinavir/Ritonavir Mylan	
Kenacomb		Letrole		Loprofin	
Kenacort-A 10		Letrozole		Loprofin Mix	
Kenacort-A 40	80	Leukeran FC	145	Lorafix	
Kenalog in Orabase		Leukotriene Receptor		Loratadine	22
Ketocal 3:1		Antagonists	233	Lorazepam	
KetoCal 4:1		Leuprorelin		Lorstat	
Ketoconazole		Leustatin		Losartan Actavis	
Dermatological	69	Levetiracetam		Losartan potassium	
Infection		Levetiracetam-AFT		Losartan potassium with	
Ketogenic Diet		Levlen ED		hydrochlorothiazide	4
Ketoprofen		Levocabastine		Lovir	10
KetoSens		Levocarnitine		Loxamine	
Keytruda		Levodopa with benserazide		Lucrin Depot 1-month	
Kindergen		Levodopa with carbidopa		Lucrin Depot 3-month	
Kivexa	105	Levomepromazine		Lyderm	
ι τιν ολα	100	Lovomopromazine		-yuo::::::::::::::::::::::::::::::::::::	

Lynparza152	Metabolic Disorder Agents26	Midazolam	13
Lyrica127	Metformin hydrochloride11	Midazolam-Baxter	136
- M -	Metformin Mylan11	Midodrine	4
m-Eslon	Methadone hydrochloride	Mifegyne	7
Mabthera200	Extemporaneous244	Mifepristone	7
Macro Organic Psyllium Husk25	Nervous122	Minerals	3
Macrobid109	Methatabs122	Mini-Wright AFS Low Range	23
Macrogol 3350 with potassium	Methenamine (hexamine)	Mini-Wright Standard	23
chloride, sodium bicarbonate and	hippurate108	Minidiab	1
sodium chloride26	Methopt239	MiniMed 1.8 Reservoir	
Macrogol 400 and propylene	Methotrexate148	MMT-326A	2
glycol 240	Methotrexate DBL Onco-Vial148	MiniMed 3.0 Reservoir	
Madopar 125118	Methotrexate Ebewe148	MMT-332A	2
Madopar 250118	Methotrexate Sandoz148	MiniMed 770G	
Madopar 62.5118	Methyl hydroxybenzoate244	MiniMed Mio MMT-921A	2
Madopar HBS118	Methylcellulose244	MiniMed Mio MMT-923A	2
Madopar Rapid118	Methylcellulose with glycerin and	MiniMed Mio MMT-925A	2
Magnesium hydroxide36	sodium saccharin244	MiniMed Mio MMT-941A	2
Magnesium sulphate36	Methylcellulose with glycerin and	MiniMed Mio MMT-943A	2
Mantoux276	sucrose244	MiniMed Mio MMT-945A	2
Marevan44	Methyldopa52	MiniMed Mio MMT-965A	2
Marine Blue Lotion SPF 50+70	Methyldopa Mylan52	MiniMed Mio MMT-975A	
Martindale Pharma241	Methyldopa Mylan S2952	MiniMed Quick-Set MMT-386A	2
Martindale Pharma S29241	Methylnaltrexone bromide25	MiniMed Quick-Set MMT-387A	
Mask for spacer device234	Methylphenidate ER - Teva138	MiniMed Quick-Set MMT-396A	2
Maviret102	Methylphenidate hydrochloride 138	MiniMed Quick-Set MMT-397A	2
Maxidex237	Methylphenidate hydrochloride	MiniMed Quick-Set MMT-398A	2
Maxitrol237	extended-release139	MiniMed Quick-Set MMT-399A	
MCT oil (Nutricia)248	Methylprednisolone79	MiniMed Silhouette MMT-368A	
Measles, mumps and rubella	Methylprednisolone (as sodium	MiniMed Silhouette MMT-377A	2
vaccine 272	succinate) 79	MiniMed Silhouette MMT-378A	2
Mebendazole89	Methylprednisolone aceponate64	MiniMed Silhouette MMT-381A	2
Mebeverine hydrochloride8	Methylprednisolone acetate80	MiniMed Silhouette MMT-382A	2
Medrol79	Methylxanthines233	MiniMed Silhouette MMT-383A	
Medroxyprogesterone acetate	Metoclopramide Actavis 10129	MiniMed Silhouette MMT-384A	
Genito-Urinary74	Metoclopramide hydrochloride 129	MiniMed Sure-T MMT-864A	
Hormone81–82	Metolazone53	MiniMed Sure-T MMT-866A	
Mefenamic acid110	Metopirone88	MiniMed Sure-T MMT-874A	
Megace164	Metoprolol IV Mylan50	MiniMed Sure-T MMT-876A	
Megestrol acetate164	Metoprolol succinate50	MiniMed Sure-T MMT-884A	
Melatonin	Metoprolol tartrate50	MiniMed Sure-T MMT-886A	
Melphalan146	Metrogyl98	Minims Pilocarpine	
Menactra272	Metronidazole98	Minims Prednisolone	
Meningococcal (groups A, C, Y and	Metyrapone88	Minirin	
W-135) conjugate vaccine 272	Mexiletine hydrochloride49	Minirin Melt	
Meningococcal B multicomponent	Miacalcic78	Mino-tabs	
vaccine	Micolette26	Minocycline hydrochloride	
Meningococcal C conjugate	Miconazole32	Minomycin	
vaccine	Miconazole nitrate	Minor Skin Infections	6
Menthol	Dermatological	Minoxidil	5
Mepolizumab	Genito-Urinary	Mirena	8 ^t
Mercaptopurine	Micreme	Mirtazapine	
Mercilon 28	Micreme H	Misoprostol	12
Mesalazine	Microgynon 20 ED74	Mitomycin C	
Mesna 152	Microgynon 30	Mitozantrone	15
Mestinon	Microlut74	Mitozantrone Ebewe	

Mixtard 30	10	Naphazoline hydrochloride	240	Factor IX]	4
MMR II		Naphcon Forte		Norethisterone	
Moclobemide	124	Naprosyn SR 1000		Genito-Urinary	<mark>7</mark>
Modafinil	140	Naprosyn SR 750		Hormone	8
Modavigil		Naproxen	110	Norflex	
Moduretic		Narcaricin mite		Norfloxacin	
Molaxole		Nasal Preparations		Noriday 28	
Molnupiravir		Natalizumab		Norimin	
Moments		Natulan		Normacol Plus	
Mometasone furoate		Nausafix		Normison	
Monogen	249	Nausicalm	129	Norpress	
Montelukast		Navelbine	156	Nortriptyline hydrochloride	
Montelukast Mylan		Nefopam hydrochloride		Norvir	
Moroctocog alfa [Recombinant f		Neisvac-C		Noumed	
VIII]		Neo-B12		Noumed Paracetamol	
Morphine hydrochloride		Neo-Mercazole		NovaSource Renal	
Morphine sulphate		Neocate Gold		Novatretin	
Motetis		Neocate Junior Unflavoured		NovoMix 30 FlexPen	
Mouth and Throat		Neocate Junior Vanilla		NovoRapid	
Movapo		Neocate SYNEO		NovoRapid FlexPen	
Moxifloxacin		Neoral		NovoRapid Penfill	
		Neostigmine metilsulfate		NovoSeven RT	
MSUD Maxamum	200	3		Noxafil	
Mucilaginous laxatives with	0.5	Nepro HP (strawberry)			
stimulants		Nepro HP (vanilla)		Nozinan	
Mucolytics		Nepro HP RTH		Nozinan (Swiss)	
Mucosoothe		Neulactil		Nucala	
Multiple Sclerosis Treatments		Neulastim		Nuelin	
Multivitamin renal		Neuraxpharm		Nuelin-SR	
Multivitamins		NeuroTabs		Nupentin	
Mupirocin		Nevirapine		Nutilis	
Muscle Relaxants		Nevirapine Alphapharm		Nutren Diabetes	
Mvite		Nicorandil		Nutrient Modules	
Myambutol		Nicotine		Nutrini Energy Multi Fibre	
Mycobutin		Nifedipine		Nutrini Energy RTH	
MycoNail	6 <mark>2</mark>	Nifuran	109	Nutrini Low Energy Multi Fibre	25
Mycophenolate mofetil	166	Nilotinib	159	Nutrini Peptisorb	26
Mydriacyl		Nilstat		Nutrini Peptisorb Energy	26
Mylan (12 hr release)		Alimentary	32	Nutrini RTH	25
Mylan (24 hr release)	51	Genito-Urinary	75	Nutrison 800 Complete Multi	
Mylan Atenolol	50	Infection	97	Fibre	25
Mylan Clomiphen	88	Nintedanib	231	Nutrison Advanced Diason	
Mylan Indapamide	54	Nipent	153	Nutrison Advanced Peptisorb	25
Mylan Italy (24 hr release)		Nirmatrelvir with ritonavir	103	Nutrison Concentrated	
Myleran	145	Nitrates	56	Nutrison Energy	25
Mylotarg	189	Nitroderm TTS	56	Nutrison Energy Multi Fibre	
Myometrial and Vaginal Hormor		Nitrofurantoin	109	Nutrison Multi Fibre	
Preparations		Nitrolingual Pump Spray	<u>56</u>	Nutrison Standard RTH	25
Myozyme		Nivestim		Nyefax Retard	5
- N -		Nivolumab	221	Nystatin	
Nadolol	50	Nizoral		Alimentary	3
Nadolol BNM		Nodia		Genito-Urinary	7
Naglazyme		Noflam 250		Infection	9
Nalcrom		Noflam 500		NZB Low Gluten Bread Mix	
Naloxone hydrochloride		Non-Steroidal Anti-Inflammator		- O -	
Naltraccord		Drugs		O/W Fatty Emulsion Cream	6
Naltrexone hydrochloride		Nonacog gamma, [Recombinar		Obinutuzumab	
		J J	-		

Obstetric Preparations	77	Other Oestrogen Preparations.	82	Paxlovid	103
Ocrelizumab	135	Other Progestogen		Pazopanib	
Ocrevus	135	Preparations	82	Peak flow meter	234
Octocog alfa [Recombinant fa	actor	Other Skin Preparations	70	Pedialyte - Bubblegum	
VIII] (Advate)		Ovestin		Pediasure	
Octocog alfa [Recombinant fa		Genito-Urinary	75	Pediasure RTH	
VIII] (Kogenate FS)		Hormone		Pegaspargase	
Octreotide		Oxaliplatin		Pegasys	
Octreotide Depot Teva		Oxaliplatin Accord		Pegfilgrastim	
Octreotide long-acting		Oxaliplatin Actavis 100		Pegylated interferon alfa-2a	
Oestradiol		Oxaliplatin Ebewe		Pembrolizumab	
Oestradiol valerate		Oxis Turbuhaler	229	Pemetrexed	
Oestradiol with norethisteron	e82	Oxpentifylline		Penicillamine	111
Oestriol		Oxybutynin		Penicillin G	
Genito-Urinary	75	Oxycodone hydrochloride		PenMix 30	
Hormone		Oxycodone Sandoz		PenMix 40	
Oestrogens		OxyNorm		PenMix 50	
Ofev		Oxytocin		Pentasa	
Oil in water emulsion		Oxytocin BNM		Pentostatin [Deoxycoformycin]	
Olanzapine		Oxytocin with ergometrine		Pentoxifylline [Oxpentifylline]	
Olaparib		maleate	75	Peptamen Junior	
Olbetam		Ozurdex		Peptisorb	252
Olbetam S29		- P -	207	Perhexiline maleate	252
Olopatadine		Pacifen	117	Pericyazine	
Olopatadine Teva		Pacimol		Perindopril	
Olsalazine		Paclitaxel		Perjeta	
				•	
Omalizumab		Paclitaxel Actavis Paclitaxel Ebewe		Permethrin	
Omeprazole actavia 10		Paediatric Seravit		Perrigo Pertuzumab	
Omeprazole actavis 10				Peteha	
Omeprazole actavis 20		Palbociclib			
Omeprazole actavis 40		Paliperidone		Pethidine hydrochloride	
Omnitrope		Palivizumab		Pevaryl	
Onbrez Breezhaler		Pamidronate disodium		Pexsig	
Oncaspar LYO		Pamisol		Pfizer Exemestane	
OncoTICE		Pancreatic enzyme		Pharmacy Health Sorbolene with	
Ondansetron		Pantoprazole		Glycerin	
Ondansetron ODT-DRLA		Panzop Relief		Pheburane	
One-Alpha		Panzytrat		Phenasen	
Onrex		Papaverine hydrochloride		Phenobarbitone	126
Opdivo		Para-amino salicylic acid		Phenobarbitone sodium	
Ora-Blend		Paracare		Extemporaneous	
Ora-Blend SF		Paracare Double Strength		Nervous	136
Ora-Plus		Paracetamol	121	Phenoxybenzamine	
Ora-Sweet		Paracetamol + Codeine		hydrochloride	
Ora-Sweet SF		(Relieve)		Phenoxymethylpenicillin (Penicilli	in
Orabase		Paracetamol with codeine		V)	
Oral and Enteral Feeds	249	Paraffin		Phenytoin sodium1	25, 127
Oratane	61	Paraffin liquid with wool fat	240	Phillips Milk of Magnesia	36
Ordine		Parasiticidal Preparations	66	Phlexy 10	
Orgran		Parlodel	118	Phosphate Phebra	
Ornidazole		Parnate	124	Phosphorus	
Orphenadrine citrate		Parnate S29	124	Phytomenadione	41
Ortho-tolidine		Paromomycin	95	Pilocarpine hydrochloride	239
Oruvail SR		Paroxetine		Pimafucort	
Osmolite RTH		Paser	99	Pimecrolimus	
Other Endocrine Agents	88	Paxam	133	Pine tar with trolamine laurilsulfat	te

and fluorescein		Pregnancy Tests - hCG Urine	76	hydrochlorothiazide	48
Pinetarsol		Premarin		Qvar	228
Pioglitazone		Prevenar 13		- R -	
Pirfenidone		Priadel		RA-Morph	
Pizotifen		Primaquine		Raloxifene hydrochloride	
PKU Anamix Infant		Primidone		Raltegravir potassium	
PKU Anamix Junior		Primidone Clinect		Ramipex	
PKU Anamix Junior Chocolate		Primolut N		Ranbaxy-Cefaclor	
PKU Anamix Junior LQ		Priorix		Ranbaxy-Cefaclor S29	
PKU Anamix Junior Orange		Probenecid		Rapamune	
PKU Anamix Junior Vanilla		Probenecid-AFT		Rasagiline	
PKU Lophlex LQ 10		Procaine penicillin		Reandron 1000	
PKU Lophlex LQ 20	261	Procarbazine hydrochloride		Recombinant factor IX	
PKU Lophlex Powder		Prochlorperazine		Recombinant factor VIIa	
PKU Lophlex Sensation 20		Proctofoam		Recombinant factor VIII	40–41
Plaquenil		Proctosedyl		Rectogesic	
Plendil ER		Procyclidine hydrochloride	119	Redipred	80
Pneumococcal (PCV10) conjug	gate	Progesterone		Relieve	110
vaccine		Proglicem		Relistor	25
Pneumococcal (PCV13) conjug	gate	Proglycem	10	Remicade	
vaccine	274	Progynova	81	Renilon 7.5	
Pneumococcal (PPV23)		Prolia		Resonium-A	
polysaccharide vaccine		Promethazine hydrochloride		Resource Beneprotein	
Pneumovax 23		Propafenone hydrochloride		Respigen	
Podophyllotoxin	70	Propamidine isethionate		Respiratory Devices	234
Polaramine		Propranolol		Respiratory Stimulants	
Poliomyelitis vaccine	275	Propylene glycol		Retinol palmitate	
Poloxamer		Propylthiouracil		ReTrieve	61
Poly-Gel		Prostacur	164	Retrovir	
Poly-Tears	239	Protaphane	10	Revlimid	151
Poly-Visc	240	Protaphane Penfill		Revolade	
Polycal	246	Protifar	248	Rexacrom	238
Ponstan	110	Protionamide	99	Riboflavin	29
Posaconazole	97	Provera	81	Ribomustin	144
Postinor-1		Provera HD	82	Ricit	76
Potassium chloride	45–46	PSM Citalopram	124	Rifabutin	100
Potassium citrate	76	Psoriasis and Eczema		Rifadin	100
Potassium iodate	34	Preparations	68	Rifampicin	100
Povidone iodine	66	PTU	83	Rifaximin	9
Pradaxa		Pulmicort Turbuhaler	228	Rifinah	99
Pramipexole hydrochloride	118	Pulmozyme	233	Rilutek	119
Pravastatin	54	Puri-nethol	148	Riluzole	119
Pravastatin Mylan	54	Puria		RINVOQ	226
Praziquantel	89	Pyrazinamide	100	Riodine	66
Prazosin	47	Pyridostigmine bromide	110	Risedronate Sandoz	113
Pred Forte	238	Pyridoxine hydrochloride	33	Risedronate sodium	113
Prednisolone	80	Pyridoxine multichem	33	Risperdal Consta	133
Prednisolone acetate	238	Pyrimethamine	95	Risperidone	131, 133
Prednisolone sodium	8	Pytazen SR		Risperidone (Teva)	131
Prednisolone sodium		- Q -		Risperon	131
phosphate	238	Quetapel	131	Ritalin	
Prednisolone-AFT	238	Quetiapine	131	Ritalin LA	
Prednisone		Quick-Set MMT-392	23	Ritonavir	
Prednisone Clinect	80	Quick-Set MMT-393	23	Rituximab (Mabthera)	200
Pregabalin	127	Quinapril	48	Rituximab (Riximyo)	202
Pregabalin Pfizer		Quinapril with		Rivaroxaban	

Rivastigmine	140	Seretide	230	Solgar2	27–29, 3°
Rivastigmine Patch BNM 10	140	Seretide Accuhaler		Solifenacin Mylan	<mark>7</mark>
Rivastigmine Patch BNM 5	140	Serevent	229	Solifenacin succinate	7
Rivotril		Serevent Accuhaler	229	Solu-Cortef	<mark>7</mark> 9
Riximyo	202	Sertraline	125	Solu-Medrol	<mark>7</mark> 9
RIXUBIS	40	Setrona	125	Solu-Medrol-Act-O-Vial	<mark>7</mark> 9
Rizamelt	128	Setrona AU	125	Somatropin (Omnitrope)	8
Rizatriptan	128	Sevredol	123	Sotalol	5
RoActemra S29	215	Sex Hormones Non		Spacer device	23
Ronapreve	188	Contraceptive	80	Span-K	
Ropin	118	Shield XL	72	Spiolto Respimat	23
Ropinirole hydrochloride	118	shingles vaccine	276	Spiractin	5
Rosuvastatin	55	SII-Onco-BCG	173	Spiriva	23
Rosuvastatin Viatris		Sildenafil		Spiriva Respimat	23
Rotarix	275	Silhouette MMT-373	23	Spironolactone	
Rotavirus oral vaccine	275	Siltuximab	214	Sporanox	90
Roxane	6	Simvastatin	55	Sprycel	15
Roxane-Propranolol	51	Simvastatin Mylan	55	Stemetil	130
Roxithromycin	91	Sinemet	118	SteroClear	23
Rubifen		Sinemet CR	118	Stesolid	12
Rubifen SR	138	Sirolimus	224	Stimulants/ADHD Treatments	130
Rugby Capsaicin Topical		Siterone	80	Stiripentol	12
Cream	120	Slow-Lopresor		Stocrin	104
Rulide D	91	Smith BioMed Rapid Pregna		Stomahesive	
Rurioctocog alfa pegol [Recon	nbinant	Test		Strattera	13
factor VIII]	41	Sodibic	46	Strides Shasun	9
Ruxolitinib	161	Sodium acid phosphate	26	Stromectol	6
Rythmodan	49	Sodium alginate		Sucralfate	
Rytmonorm		Sodium benzoate		Sulfadiazine Silver	6
- S -		Sodium bicarbonate		Sulfadiazine sodium	
Sabril	128	Blood	45–46	Sulfasalazine	
Sacubitril with valsartan	48	Extemporaneous	244	Sulphur	6
Sagent	156	Sodium calcium edetate	242	Sulprix	
SalAir		Sodium chloride		Sumagran	
Salazopyrin	8	Blood	45	Sumatriptan	
Salazopyrin EN		Respiratory	234	Sunitinib	16
Salbutamol		Sodium citrate with sodium I		Sunitinib Pfizer	16
Salbutamol with ipratropium		sulphoacetate	•	Sunscreens	70
bromide	230	Sodium citro-tartrate		Sunscreens, proprietary	
Salicylic acid	69	Sodium cromoglicate		Sure-T MMT-863	
Salmeterol	229	Alimentary	8	Sure-T MMT-873	
Sandomigran		Sensory		Sustagen Hospital Formula	
Sanofi Primaguine	98	Sodium fluoride	34	Sustagen Hospital Formula	
Sapropterin dihydrochloride	30	Sodium Fusidate [fusidic aci	d]	Active	250
Scalp Preparations		Dermatological	62	Sustanon Ampoules	
Scopoderm TTS		Infection		Sylvant	
Sebizole		Sensory		Symbicort Turbuhaler 100/6	
Secukinumab	212	Sodium hyaluronate [Hyaluro		Symbicort Turbuhaler 200/6	
Sedatives and Hypnotics		acid]	240	Symbicort Turbuhaler 400/12	
Seebri Breezhaler		Sodium phenylbutyrate		Symmetrel	
Selegiline hydrochloride	118	Sodium picosulfate		Sympathomimetics	
Senna		Sodium polystyrene sulphon		Synacthen	
Senokot		Sodium tetradecyl sulphate.		Synacthen Depot	
SensoCard		Sodium valproate		Synacthene Retard	
Serc		Sofradex		Synagis	
Serenace		Soframycin		Synflorix	

INDEX: Generic Chemicals and Brands

Synthroid	83	Thiotepa	146	Trimethoprim	9
Syntometrine		Thyroid and Antithyroid Agents	83	Trimethoprim with	
Syrup (pharmaceutical grade).		Ticagrelor		sulphamethoxazole	
Systane Unit Dose		Tilcotil		[Co-trimoxazole]	9
· т-		Timolol		Trisequens	
Tacrolimus		Timoptol XE		Trisul	
Dermatological	69	Tiotropium bromide		Trophic Hormones	
Oncology		Tiotropium bromide with		Tropicamide	
Tacrolimus Sandoz		olodaterol	231	Trulicity	1
Taliglucerase alfa		Tivicay		Trusopt	23
Tambocor		Tixagevimab with cilgavimab		TruSteel	
Tamoxifen citrate		TMP		Tuberculin PPD [Mantoux] test	
Tamoxifen Sandoz		Tobramycin		Tubersol	
Tamsulosin hydrochloride		,	05		
•		Infection		Two Cal HN	
Tamsulosin-Rex		Sensory		Tykerb	
Tandem Cartridge		Tobramycin BNM		Tysabri	13
Tandem t:slim X2 with Basal-IC		Tobramycin Mylan		- U -	
Tap water		Tobrex		Ultibro Breezhaler	
Tarceva		Tocilizumab		Ultraproct	
Tasigna	159	Tofranil	124	Umeclidinium	
Tasmar		Tolcapone	118	Umeclidinium with vilanterol	
Taurine	31	Topamax	127	Univent	.230, 23
Tecfidera	134	Topical Products for Joint and		Upadacitinib	22
Tegretol	125	Muscular Pain	111	Ural	7
Tegretol CR		Topiramate	127	Urea	6
Telfast	228	Topiramate Actavis	127	Urex Forte	5
Teligent		Total parenteral nutrition (TPN).		Urinary Agents	7
Temaccord		TPN		Urinary Tract Infections	10
Temazepam		Tramadol hydrochloride		Urinorm	
Temozolomide		Tramal SR 100		Uromitexan	
Tenofovir disoproxil		Tramal SR 150		Ursodeoxycholic acid	
Tenofovir Disoproxil Emtricitab		Tramal SR 200		Ursosan	
Mylan		Trandate		Utrogestan	
Tenofovir Disoproxil Mylan		Tranexamic acid		- V -	
Tenofovir Disoproxil Teva		Tranylcypromine sulphate		Vaccinations	26
Tenoxicam		Trastuzumab		Vaclovir	
		Trastuzumab emtansine		Valaciclovir	
Tensipine MR10					
Tepadina		Travatan		Valganciclovir	
Terbinafine		Travoprost		Valganciclovir Mylan	10
Terbutaline sulphate		Treatments for Dementia	140	Vancomycin	
Teriflunomide		Treatments for Substance		Vannair	
Teriparatide		Dependence		Varenicline Pfizer	
Testosterone		Trental 400	5/	Varenicline tartrate	14
Testosterone cipionate		Tretinoin		Varicella vaccine [Chickenpox	
Testosterone esters		Dermatological		vaccine]	
Testosterone undecanoate		Oncology		Varicella zoster virus (Oka strai	,
Tetrabenazine		Trexate	148	attenuated vaccine [shingles	3
Tetrabromophenol	77	Triamcinolone acetonide		vaccine]	
Tetracosactrin	80	Alimentary	32	Varivax	27
Tetracycline		Dermatological	64	Vasodilators	
Thalidomide	155	Hormone		Vasopressin Agonists	8
Thalomid		Triamcinolone acetonide with		Vasorex	
Theophylline		gramicidin, neomycin and ny	statin	Vedafil	
Thiamine hydrochloride		Dermatological		Veletri	
THIO-TEPA		Sensory		Venclexta	
Thioguanine		Triazolam		Venetoclax	
9~~					

Venlafaxine125	Xylocaine
Venomil227	Xylocaine 2% Jelly
VENOX	Xyntha
Ventavis 60	-Z-
	Zapril
Ventolin	
Vepesid	Zarontin
Verapamil hydrochloride52	Zaroxolyn
Vermox89	Zavedos
Versacloz130	Zeffix
Vesanoid155	Zematop
Vexazone11	Zetlam
Vfend97	Ziagen
Viaderm KC64	Zidovudine [AZT]
Vigabatrin128	Zidovudine [AZT] with
Vigisom135	lamivudine
Vildagliptin11	Zimybe
Vildagliptin with metformin	Zinc and castor oil
hydrochloride12	Zinc sulphate
Vimpat126	Zincaps
Vinblastine sulphate156	Zinnat
Vincristine sulphate	Ziprasidone
Vinorelbine	Zista
Vinorelbine Ebewe	Zithromax
Viramune Suspension	Zoledronic acid
ViruPOS236	Hormone
Vit.D3	Musculoskeletal
Vita-B1233	
	Zoledronic acid Mylan
VitA-POS	Zopiclone
Vitabdeck34	Zopiclone Actavis
Vital	Zostavax
Vitamin B complex33	Zostrix
Vitamin B6 2533	Zostrix HP
Vitamins33–34	Zuclopenthixol decanoate
Vivonex TEN252	Zuclopenthixol hydrochloride.
Voltaren110	Zusdone
Voltaren D110	Zyban
Voltaren Ophtha237	Zypine
Voltaren SR110	Zypine ODT
Volumatic234	Zyprexa Relprevv
Voriconazole97	Zytiga
Votrient160	, 0
Vttack97	
- W -	
Warfarin sodium44	
Wart Preparations70	
Wasp venom allergy treatment227	
Water	
Blood45	
Extemporaneous245	
Wool fat with mineral oil	
- X -	
Xarelto	
Xifaxan9 XMET Maxamum	
Xolair	
XP Mayamum 261	

Xylocaine	120
Xylocaine 2% Jelly	119
Xyntha	
- Z -	
Zapril	4
Zarontin	.120
Zaroxolyn	5
Zavedos	15
Zeffix	100
Zematop	69
Zetlam	. 100
Ziagen	. 10
Zidovudine [AZT]	10
Zidovudine [AZT] with	
lamivudine	
Zimybe	5
Zinc and castor oil	6
Zinc sulphate	
Zincaps	30
Zinnat	
Ziprasidone	13
Zista	
Zithromax	90
Zoledronic acid	
Hormone	
Musculoskeletal	
Zoledronic acid Mylan	78
Zopiclone	. 13
Zopiclone Actavis	. 13
Zostavax	27
Zostrix	
Zostrix HP	
Zuclopenthixol decanoate	. 13
Zuclopenthixol hydrochloride	
Zusdone	13
Zyban	14
Zypine	13
Zypine ODT	13
Zyprexa Relprevv	13
Zytiga	16