October 2022 Volume 29 Number 2

Editors:

Kaye Wilson, & Doris Chong email: enquiry@pharmac.govt.nz Telephone +64 4 460 4990 Level 9, 40 Mercer Street PO Box 10 254 Wellington

Freephone Information Line 0800 66 00 50 (9am – 5pm weekdays)

Circulation

You can register to have an electronic version of the Pharmaceutical Schedule (link to PDF copy) emailed to your nominated email address each month by subscribing at pharmac.govt.nz/subscribe.

Production

Typeset automatically from XML and T_EX. XML version of the Schedule available from schedule.pharmac.govt.nz/pub/schedule

Programmers

Anrik Drenth & John Geering email: texschedule@pharmac.govt.nz

©Pharmaceutical Management Agency

ISSN 1179-3686

This work is licensed under the Creative Commons Attribution 4.0 International licence. In essence, you are free to copy, distribute and adapt it, as long as you attribute the work to Pharmac and abide by the other licence terms. To view a copy of this licence, visit: creativecommons.org/licenses/by/4.0/. Attribution to Pharmac should be in written form and not by reproduction of the Pharmac logo. While care has been taken in compiling this Schedule, Pharmac takes no responsibility for any errors or omissions, and shall not be liable for any consequences arising there from.

Section A	General Rules	5
Section B		
Section b	Alimentary Tract & Metabolism	6
	Blood & Blood Forming Organs	38
	Cardiovascular System	48
	Dermatologicals	62
	Genito Urinary System	72
	Hormone Preparations – Systemic	79
	Infections – Agents For Systemic Use	90
	Musculoskeletal System	110
	Nervous System	118
	Oncology Agents & Immunosuppressants	144
	Respiratory System & Allergies	232
	Sensory Organs	241
	Various	246
Section C	Extemporaneous Compounds (ECPs)	248
Section D	Special Foods	251
Section I	National Immunisation Schedule	273
	lanka	222

Introducing Pharmac

Index

Introducing Pharmac

The Pharmaceutical Management Agency (Pharmac) makes decisions that help control Government spending on pharmaceuticals. This includes community pharmaceuticals, hospital pharmaceuticals, vaccines and increasingly, hospital medical devices. Pharmac negotiates prices, sets subsidy levels and conditions, and makes decisions on changes to the subsidised list.

Pharmac's role:

"to secure for eligible people in need of pharmaceuticals, the best health outcomes that are reasonably achievable from pharmaceutical treatment and from within the amount of funding provided."

Pae Ora (Healthy Futures) Act 2022

To ensure our decisions are as fair and robust as possible we use a decision-making process that incorporates clinical, economic and commercial issues. We also seek the views of users and the wider community through consultation. The processes we generally use are outlined in our Operating Policies and Procedures.

Further information about Pharmac and the way we make funding decisions can be found on the Pharmac website at https://www.pharmac.govt.nz/about.

Purpose of the Pharmaceutical Schedule

The purpose of the Schedule is to list:

- the Community Pharmaceuticals that are subsidised by the Government and to show the amount of the subsidy paid to contractors, as well as the manufacturer's price and any access conditions that may apply;
- the Hospital Pharmaceuticals that may be used in Health NZ Hospitals, as well as any access conditions that may apply; and
- the Pharmaceuticals, including Medical Devices, used in Health NZ Hospitals for which national prices have been negotiated by Pharmac.

The Schedule does not show the final cost to Government of subsidising each Community Pharmaceutical or to Health NZ Hospitals in purchasing each Pharmaceutical, since that will depend on any rebate and other arrangements Pharmac has with the supplier and, for Pharmaceuticals used in Health NZ Hospitals, on any logistics arrangements put in place.

This book contains sections A to D and Section I of the Pharmaceutical Schedule and lists the Pharmaceuticals funded for use in the community, including vaccines, as well as haemophilia and cancer treatments given in Health NZ hospitals. Section H lists the Pharmaceuticals that that can be used in Health NZ hospitals and is a separate publication.


The Pharmaceuticals in this book are listed by therapeutic group, which is based on the WHO Anatomical Therapeutic Chemical (ATC) system. The listings are displayed alphabetically under each heading.

The index lists both chemical entities and product brand names.

Explaining pharmaceutical entries

The Pharmaceutical Schedule lists pharmaceuticals subsidised by the Government, the subsidy, the supplier's price and the access conditions that may apply.

Example



Glossary

Units of Measure

gramg kilogramkg international unitiu	mi mi mi
Abbreviations	
AmpouleAmp	Ge
CapsuleCap	Gr
Cream	Inf
DeviceDev	Ini
DispersibleDisp	Lic
EffervescentEff	Lo
EmulsionEmul	Oi
Enteric Coated EC	Sa

microgrammilligrammillilitre	mg
Gelatinous	
Granules	
Infusion	Inf
Injection	Inj
Liquid	Liq
Long Acting	LA
Ointment	Oint
Sachet	Sach

millimoleunit	
Solution	Supp Tab
Trans Dermal Delivery System	TDDS

Read the <u>General Rules</u>: <u>https://www.pharmac.govt.nz/section-a</u>.

SECTION B: ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price \$	e) Sul Per	Fully osidised	Brand or Generic Manufacturer
Antacids and Antiflatulents				
Antacids and Reflux Barrier Agents				
LGINIC ACID Sodium alginate 225 mg and magnesium alginate 87.5 mg sachet	0 1	30	√ Ga	aviscon Infant
SODIUM ALGINATE * Tab 500 mg with sodium bicarbonate 267 mg and calcium carbonate 160 mg - peppermint flavour		60		aviscon Double Strength
Foral liq 500 mg with sodium bicarbonate 267 mg and calc carbonate 160 mg per 10 ml		500 ml		idex
Phosphate Binding Agents				
ALUMINIUM HYDROXIDE Tab 600 mg CALCIUM CARBONATE Oral liq 1,250 mg per 5 ml (500 mg elemental per 5 ml)		100	✓ Al	u-Tab
Subsidy by endorsementOnly when prescribed for patients unable to swallow of inappropriate and the prescription is endorsed according to the control of the control o	39.00 calcium carbonate table	500 ml ets or whe		oxane n carbonate tablets are
Antidiarrhoeals				
Agents Which Reduce Motility				
OPERAMIDE HYDROCHLORIDE – Up to 30 cap available of Tab 2 mg	10.75	400 400	✓ No	odia amide Relief
Rectal and Colonic Anti-inflammatories				
BUDESONIDE Cap 3 mg - Special Authority see SA1886 below - Retail pharmacy		90	√ Er	ntocort CIR
■ SA1886 Special Authority for Subsidy nitial application — (Crohn's disease) from any relevant pute following criteria: toth:		valid for 6	months fo	r applications meeting

- 1 Mild to moderate ileal, ileocaecal or proximal Crohn's disease; and
- 2 Any of the following:
 - 2.1 Diabetes; or
 - 2.2 Cushingoid habitus; or
 - 2.3 Osteoporosis where there is significant risk of fracture; or

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 2.4 Severe acne following treatment with conventional corticosteroid therapy; or
- 2.5 History of severe psychiatric problems associated with corticosteroid treatment; or
- 2.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
- 2.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated).

Initial application — (collagenous and lymphocytic colitis (microscopic colitis)) from any relevant practitioner. Approvals valid for 6 months where patient has a diagnosis of microscopic colitis (collagenous or lymphocytic colitis) by colonoscopy with biopsies.

Initial application — (gut Graft versus Host disease) from any relevant practitioner. Approvals valid for 6 months where patient has a gut Graft versus Host disease following allogenic bone marrow transplantation*.

Note: Indication marked with * is an unapproved indication.

Initial application — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has autoimmune hepatitis*; and
- 2 Patient does not have cirrhosis; and
- 3 Any of the following:
 - 3.1 Diabetes; or
 - 3.2 Cushingoid habitus; or
 - 3.3 Osteoporosis where there is significant risk of fracture; or
 - 3.4 Severe acne following treatment with conventional corticosteroid therapy; or
 - 3.5 History of severe psychiatric problems associated with corticosteroid treatment; or
 - 3.6 History of major mental illness (such as bipolar affective disorder) where the risk of conventional corticosteroid treatment causing relapse is considered to be high; or
 - 3.7 Relapse during pregnancy (where conventional corticosteroids are considered to be contraindicated); or
 - 3.8 Adolescents with poor linear growth (where conventional corticosteroid use may limit further growth) .

Note: Indication marked with * is an unapproved indication.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (non-cirrhotic autoimmune hepatitis) from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: Clinical trials for Entocort CIR use beyond three months demonstrated no improvement in relapse rate.

HYDROCORTISONE ACETATE

Rectal foam 10%, CFC-Free (14 applications)26.55	15 g OP 21.1 g OP	✓ Cortifoam S29✓ Colifoam
HYDROCORTISONE ACETATE WITH PRAMOXINE HYDROCHLORIDE		
Topical aerosol foam, 1% with pramoxine hydrochloride 1%26.55	10 g OP	✓ Proctofoam §29
MESALAZINE		
Tab 400 mg49.50	100	✓ Asacol
Tab long-acting 500 mg56.10	100	✓ Pentasa
Tab 800 mg85.50	90	✓ Asacol
Modified release granules, 1 g118.10	100 OP	✓ Pentasa
Enema 1 g per 100 ml41.30	7	✓ Pentasa
Suppos 500 mg22.80	20	✓ Asacol
Suppos 1 g50.96	28	✓ Pentasa

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
OLSALAZINE Tab 500 mg	56.02	60	✓	Atnahs Olsalazine \$29
Cap 250 mg	93.37 53.00	100 100		Dipentum Dipentum
PREDNISOLONE SODIUM Rectal foam 20 mg per dose (14 applications)	74.10	1 OP	•	Essential Prednisolone S29
SODIUM CROMOGLICATE Cap 100 mg(Nalcrom Cap 100 mg to be delisted 1 April 2023)	92.91	100		Nalcrom Ralicrom
SULFASALAZINE * Tab 500 mg Tab EC 500 mg		100 100		Salazopyrin Salazopyrin EN

Local preparations for Anal and Rectal Disorders

Antihaemorrhoidal Preparations

FLUOCORTOLONE CAPROATE WITH FLUOCORTOLONE PIVALATE AND C	NCHOCAINE	
Oint 950 mcg, with fluocortolone pivalate 920 mcg, and cinchocaine hydrochloride 5 mg per g11.06	30 g OP	✓ Ultraproct
Suppos 630 mcg, with fluocortolone pivalate 610 mcg, and cinchocaine hydrochloride 1 mg7.30	12	✓ Ultraproct
HYDROCORTISONE WITH CINCHOCAINE		
Oint 5 mg with cinchocaine hydrochloride 5 mg per g15.00	30 g OP	✓ Proctosedyl
Suppos 5 mg with cinchocaine hydrochloride 5 mg per g9.90	12	✓ Proctosedyl

Management of Anal Fissures

⇒SA1329 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has a chronic anal fissure that has persisted for longer than three weeks.

Antispasmodics and Other Agents Altering Gut Motility

GLYCOPYRRONIUM BROMIDE			
Inj 200 mcg per ml, 1 ml ampoule - Up to 10 inj available on a			
PSO	65.45	10	Max Health
HYOSCINE BUTYLBROMIDE			
* Tab 10 mg	6.35	100	✓ Buscopan
* Inj 20 mg, 1 ml - Up to 5 inj available on a PSO	6.35	5	✓ Buscopan
MEBEVERINE HYDROCHLORIDE			
* Tab 135 mg	9.20	90	✓ Colofac

		ALIMENTARY	TRACT	AND	METABOLISM
		Subsidy (Manufacturer's Price) \$	Subsid Per	Fully lised	Brand or Generic Manufacturer
Antiulcerants					
Antisecretory and	Cytoprotective				
MISOPROSTOL * Tab 200 mcg - Up to	o 120 tab available on a PSO	41.50	120	✓ C	ytotec
Helicobacter Pylor	ri Eradication				
a) Maximum of 2b) Subsidised of Note: the pre	dy by endorsement	i eradication and prescr		dorsec	
H2 Antagonists					
FAMOTIDINE – Only on * Tab 20 mg	a prescription	4.91	100	√ F	amotidine Hovid §29
* Tab 40 mg		8.48	100	√ F	amotidine Hovid \$29
	nl – Subsidy by endorsementrsement – Subsidised for patients rec		10 t of palliative		lylan S29
Proton Pump Inhil	bitors				
* Cap 30 mg OMEPRAZOLE	pension refer Standard Formulae, pag	5.26	100 100		anzol Relief anzol Relief
			90	√ 0	meprazole actavis 10
* Cap 20 mg		1.86	90	√ <u>0</u>	meprazole actavis 20
* Cap 40 mg		3.11	90	√ <u>0</u>	meprazole actavis 40
	mbinationraneously compounded omeprazole s		5 g	✓ N	lidwest
	ith diluent		5	✓ D	r Reddy's Omeprazole
			100 100		anzop Relief anzop Relief
Site Protective Ag	ents				
COLLOIDAL BISMUTH S Tab 120 mg	SUBCITRATE	14.51	50	√ G	astrodenol S29

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer
SUCRALFATE Tab 1 g	35.50 (48.28)	120		Carafate
Bile and Liver Therapy				
RIFAXIMIN – Special Authority see SA1461 below – Retail Tab 550 mg SA1461 Special Authority for Subsidy Initial application only from a gastroenterologist, hepatological application only from a gastroenterologist, hepatologist, hepa	625.00		mendation	
hepatologist. Approvals valid for 6 months where the patiel tolerated doses of lactulose. Renewal only from a gastroenterologist, hepatologist or Prahepatologist. Approvals valid without further renewal unles benefiting from treatment.	actitioner on the recomme	endatio	n of a gastr	oenterologist or
Diabetes				
Hyperglycaemic Agents				
DIAZOXIDE - Special Authority see SA1320 below - Reta Cap 25 mg Cap 100 mg Oral liq 50 mg per ml	110.00	100 100 30 ml 0	✓ OP ✓	Proglicem \$29 Proglicem \$29 Proglycem \$29 e5 Pharma \$29
➤ SA1320 Special Authority for Subsidy Initial application from any relevant practitioner. Approval hypoglycaemia caused by hyperinsulinism. Renewal from any relevant practitioner. Approvals valid with appropriate and the patient is benefiting from treatment.			d for the tre	eatment of confirmed
GLUCAGON HYDROCHLORIDE Inj 1 mg syringe kit – Up to 5 kit available on a PSO	32.00	1	✓	Glucagen Hypokit
Insulin - Short-acting Preparations				
INSULIN NEUTRAL ▲ Inj human 100 u per ml	25.26	10 ml (Actrapid Humulin B
▲ Inj human 100 u per ml, 3 ml	42.66	5	✓.	Actrapid Penfill Humulin R
Insulin - Intermediate-acting Preparations				
INSULIN ASPART WITH INSULIN ASPART PROTAMINE Inj 100 iu per ml, 3 ml prefilled pen	52.15	5	•	NovoMix 30 FlexPen
INSULIN ISOPHANE ▲ Inj human 100 u per ml	17.68	10 ml (Humulin NPH Protaphane

▲ Inj human 100 u per ml, 3 ml......29.86

5

✓ Humulin NPH

✓ Protaphane Penfill

	Subsidy		Fully Brand or
	(Manufacturer's Pric		sidised Generic
	\$	Per	✓ Manufacturer
NSULIN ISOPHANE WITH INSULIN NEUTRAL			
Inj human with neutral insulin 100 u per ml	25.26	10 ml OP	✓ Humulin 30/70✓ Mixtard 30
Inj human with neutral insulin 100 u per ml, 3 ml	42.66	5	✓ Humulin 30/70 ✓ PenMix 30 ✓ PenMix 40 ✓ PenMix 50
PenMix 40 Inj human with neutral insulin 100 u per ml, 3 ml to b	e delisted 1 Decem	ber 2022)	
NSULIN LISPRO WITH INSULIN LISPRO PROTAMINE			
Inj lispro 25% with insulin lispro protamine 75% 100 u per ml	l,		
3 ml	•	5	✓ Humalog Mix 25
▲ Inj lispro 50% with insulin lispro protamine 50% 100 u per m	l,		·
3 ml	42.66	5	Humalog Mix 50
Insulin - Long-acting Preparations			
NSULIN GLARGINE			
▲ Inj 100 u per ml, 10 ml	63.00	1	✓ Lantus
▲ Inj 100 u per ml, 3 ml	94.50	5	✓ Lantus
Inj 100 u per ml, 3 ml disposable pen	94.50	5	✓ Lantus SoloStar
Insulin - Rapid Acting Preparations			
NSULIN ASPART			
▲ Inj 100 u per ml, 10 ml	30.03	1	✓ NovoRapid
Inj 100 u per ml, 3 ml		5	✓ NovoRapid Penfill
▲ Inj 100 u per ml, 3 ml syringe	51.19	5	✓ NovoRapid FlexPen
NSULIN GLULISINE			
▲ Inj 100 u per ml, 10 ml	27.03	1	✓ Apidra
▲ Inj 100 u per ml, 3 ml		5	✓ Apidra
Inj 100 u per ml, 3 ml disposable pen	46.07	5	✓ Apidra SoloStar
NSULIN LISPRO			
▲ Inj 100 u per ml, 10 ml	34.92	10 ml OP	✓ Humalog
Inj 100 u per ml, 3 ml	59.52	5	✓ Humalog
Alpha Glucosidase Inhibitors			
ACARBOSE			
* Tab 50 mg	8 95	90	✓ Accarb
* Tab 100 mg		90	✓ Accarb
Oral Hypoglycaemic Agents			
GLIBENCLAMIDE	7.50	400	∠ December
* Tab 5 mg	/.50	100	✓ <u>Daonil</u>
GLICLAZIDE			
★ Tab 80 mg	15.18	500	✓ Glizide
9			
GLIPIZIDE * Tab 5 mg		100	✓ Minidiab

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
14.74	1,000		Metformin Mylan Metformin Viatris
11.28	500	1	Metformin Mylan
6.80	90	✓	Vexazone
	90	✓	Vexazone
12.25	90	✓	Vexazone
35.00	60	✓	Galvus
35.00 35.00	60 60	_	Galvumet Galvumet
	(Manufacturer's Price) \$14.7411.286.807.3012.2535.00	(Manufacturer's Price) Per 14.74 1,000	(Manufacturer's Price) Subsidised Per \$ Per ✓ ✓ 14.74 1,000 ✓ ✓ 11.28 500 ✓ ✓ 6.80 90 7.30 90 ✓ ✓ 35.00 60 ✓

GLP-1 Agonists

⇒SA2065 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Patient has previously received an initial approval for an SGLT-2 inhibitor; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Māori or any Pacific ethnicity*: or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
- 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

טט	LAGLUTIDE – Special Authority see SA2065 above – Retail pharm	nacy		
	Note: Not to be given in combination with a funded SGLT-2 inhibit	tor.		
*	Inj 1.5mg per 0.5 ml prefilled pen	115.23	4	✓ Trulicity

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

SGLT2 Inhibitors

⇒SA2068 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has previously received an initial approval for a GLP-1 agonist; or
- 2 All of the following:
 - 2.1 Patient has type 2 diabetes; and
 - 2.2 Any of the following:
 - 2.2.1 Patient is Maori or any Pacific ethnicity*; or
 - 2.2.2 Patient has pre-existing cardiovascular disease or risk equivalent (see note a)*; or
 - 2.2.3 Patient has an absolute 5-year cardiovascular disease risk of 15% or greater according to a validated cardiovascular risk assessment calculator*; or
 - 2.2.4 Patient has a high lifetime cardiovascular risk due to being diagnosed with type 2 diabetes during childhood or as a young adult*; or
 - 2.2.5 Patient has diabetic kidney disease (see note b)*; and
 - 2.3 Target HbA1c (of 53 mmol/mol or less) has not been achieved despite the regular use of at least one blood-glucose lowering agent (e.g. metformin, vildagliptin, or insulin) for at least 3 months.

Notes: * Criteria intended to describe patients at high risk of cardiovascular or renal complications of diabetes.

- a) Pre-existing cardiovascular disease or risk equivalent defined as: prior cardiovascular disease event (i.e. angina, myocardial infarction, percutaneous coronary intervention, coronary artery bypass grafting, transient ischaemic attack, ischaemic stroke, peripheral vascular disease), congestive heart failure or familial hypercholesterolaemia.
- b) Diabetic kidney disease defined as: persistent albuminuria (albumin:creatinine ratio greater than or equal to 3 mg/mmol, in at least two out of three samples over a 3-6 month period) and/or eGFR less than 60 mL/min/1.73m2 in the presence of diabetes, without alternative cause.

EMPAGLIFLOZIN – Special Authority see SA2068 above – Retail pharmacy

*	Tab 10 mg58.56	30	Jardiance
*	Tah 25 mg 58 56	30	✓ .lardiance

EMPAGLIFLOZIN WITH METFORMIN HYDROCHLORIDE – Special Authority see SA2068 above – Retail pharmacy Note: Not to be given in combination with a funded GLP-1 agonist.

*	Tab 5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 1,000 mg metformin hydrochloride	58.56	60	Jardiamet
*	Tab 12.5 mg with 500 mg metformin hydrochloride	58.56	60	Jardiamet

Subsidy Fully Brand or
(Manufacturer's Price) Subsidised Generic

\$ Per ✓ Manufacturer

Diabetes Management

Ketone Testing

BLOOD KETONE DIAGNOSTIC TEST STRIP - Subsidy by endorsement

- a) Not on a BSO
- b) Maximum of 20 strip per prescription
- c) Up to 10 strip available on a PSO
- d) Patient has any of the following:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes: or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly.

Dual Blood Glucose and Blood Ketone Testing

DUAL BLOOD GLUCOSE AND BLOOD KETONE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A dual blood glucose and blood ketone diagnostic test meter is subsidised for a patient who has:
 - 1) type 1 diabetes; or
 - 2) permanent neonatal diabetes; or
 - 3) undergone a pancreatectomy; or
 - 4) cystic fibrosis-related diabetes; or
 - 5) metabolic disease or epilepsy under the care of a paediatrician, neurologist or metabolic specialist.

The prescription must be endorsed accordingly. Only 1 meter per patient will be subsidised (no repeat prescriptions). For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Meter with 50 lancets, a lancing device and 10 blood glucose

diagnostic test strips ______20.00 1 OP ✓ CareSens Dual

				_
Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	ubsidised	Generic	
\$	Per	1	Manufacturer	

Blood Glucose Testing

BLOOD GLUCOSE DIAGNOSTIC TEST METER - Subsidy by endorsement

- a) Maximum of 1 pack per prescription
- b) Up to 1 pack available on a PSO
- c) A diagnostic blood glucose test meter is subsidised for a patient who:
 - 1) is receiving insulin or sulphonylurea therapy; or
 - 2) is pregnant with diabetes: or
 - 3) is on home TPN at risk of hypoglycaemia or hyperglycaemia; or
 - 4) has a genetic or an acquired disorder of glucose homeostasis, excluding type 1 or type 2 diabetes and metabolic syndrome.

The prescription must be endorsed accordingly. Only one CareSens meter per patient will be subsidised (no repeat prescriptions). Patients already using the CareSens N POP meter and CareSens N meter are not eligible for a new meter, unless they have:

- 1) type 1 diabetes; or
- 2) permanent neonatal diabetes: or
- 3) undergone a pancreatectomy; or
- 4) cystic fibrosis-related diabetes.

For the avoidance of doubt patients who have previously received a funded meter, other than CareSens, are eligible for a funded CareSens meter.

Note: Only 1 meter available per PSO

BLOOD GLUCOSE DIAGNOSTIC TEST STRIP - Up to 50 test available on a PSO

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Test strips	50 test OP	✓ CareSens N
		✓ CareSens PRO

BLOOD GLUCOSE TEST STRIPS (VISUALLY IMPAIRED)

The number of test strips available on a prescription is restricted to 50 unless:

- Prescribed for a patient on insulin or a sulphonylurea and endorsed accordingly. Pharmacists may annotate the
 prescription as endorsed where there exists a record of prior dispensing of insulin or sulphonylurea; or
- Prescribed on the same prescription as insulin or a sulphonylurea in which case the prescription is deemed to be endorsed; or
- 3) Prescribed for a pregnant woman with diabetes and endorsed accordingly; or
- 4) Prescribed for a patient on home TPN at risk of hypoglycaemia or hyperglycaemia and endorsed accordingly; or
- 5) Prescribed for a patient with a genetic or an acquired disorder of glucose homeostasis excluding type 1 or type 2 diabetes and metabolic syndrome and endorsed accordingly.

Blood glucose test strips26.	6.20 50 test	OP 🗸 🤋	SensoCard
------------------------------	--------------	--------	-----------

Subsidy	Fu	ly B	rand or
(Manufacturer's Price)	Subsidise	ed G	Generic
\$	Per	/ N	Manufacturer

Insulin Syringes and Needles

Subsidy is available for disposable insulin syringes, needles, and pen needles if prescribed on the same form as the one used for the supply of insulin or when prescribed for an insulin patient and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of insulin.

INSULIN PEN NEEDLES – Maximum of 200 (
--

IIV	ocini i chi necocco – maximum di 200 dev pei prescripti	1011		
*	29 g × 12.7 mm	10.95	100	✓ B-D Micro-Fine
*	31 g × 5 mm		100	✓ B-D Micro-Fine
*	31 g × 6 mm	9.50	100	✓ Berpu
*	31 g × 8 mm		100	✓ B-D Micro-Fine
*	32 g × 4 mm		100	✓ B-D Micro-Fine
INS	SULIN SYRINGES, DISPOSABLE WITH ATTACHED NEED	LE - Maximum of 2	200 dev per p	orescription
*	Syringe 0.3 ml with 29 g x 12.7 mm needle	13.56	100	B-D Ultra Fine
		1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.3 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.30	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 0.5 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 0.5 ml with 31 g x 8 mm needle	13.56	100	✓ B-D Ultra Fine II
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine II
*	Syringe 1 ml with 29 g x 12.7 mm needle	13.56	100	✓ B-D Ultra Fine
	, ,	1.36	10	
		(1.99)		B-D Ultra Fine
*	Syringe 1 ml with 31 g × 8 mm needle	13.56	100	✓ B-D Ultra Fine II
		1.36	10	
		(1.99)		B-D Ultra Fine II
		` '		

Insulin Pumps

INSULIN PUMP - Special Authority see SA1603 below - Retail pharmacy

- a) Maximum of 1 dev per prescription
- b) Only on a prescription

c) Maximum of 1 insulin pump per patient each four year	ar period.		
Min basal rate 0.025 U/h	8,800.00	1	✓ MiniMed 770G
Min basal rate 0.1 U/h	4,500.00	1	✓ Tandem t:slim
			X2 with Basal-IQ

⇒SA1603 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subsi	dised	Generic
\$	Per	✓	Manufacturer

continued...

education from an appropriate health professional); and

- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 It has been at least 4 years since the last insulin pump received by the patient or, in the case of patients qualifying under previous pump therapy for the initial application; the pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol: and
- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Fither:
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Fither:
 - 4.1 Applicant is a relevant specialist: or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and

			
	Subsidy	Fully	Brand or
	(Manufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol: and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Fither:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol: and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Fither
 - 3.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 3.2 The pump is due for replacement; and
- 4 Either:
 - 4.1 Applicant is a relevant specialist; or
 - 4.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating pump therapy; and
- 4 The patient is continuing to derive benefit from pump therapy; and
- 5 The patient had achieved and is maintaining a HbA1c of equal to or less than 80 mmol/mol on pump therapy; and
- 6 The patient has had no increase in severe unexplained hypoglycaemic episodes from baseline; and
- 7 The patient's HbA1c has not deteriorated more than 5 mmol/mol from baseline; and
- 8 Either:
 - 8.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 8.2 The pump is due for replacement; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

continued...

- 1 The patient is continuing to derive benefit according to the treatment plan and has maintained a HbA1c of equal to or less than 80 mmol/mol; and
- 2 The patient's HbA1c has not deteriorated more than 5 mmol/mol from the time of commencing pump treatment; and
- 3 The patient has not had an increase in severe unexplained hypoglycaemic episodes from baseline; and
- 4 Either:
 - 4.1 It has been at least 4 years since the last insulin pump was received by the patient; or
 - 4.2 The pump is due for replacement; and
- 5 Either:
 - 5.1 Applicant is a relevant specialist; or
 - 5.2 Applicant is a nurse practitioner working within their vocational scope.

Insulin Pump Consumables

⇒SA1985 Special Authority for Subsidy

Initial application — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has permanent neonatal diabetes; and
- 2 A MDI regimen trial is inappropriate; and
- 3 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 4 Patient/Parent/Guardian has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 5 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 6 Either:
 - 6.1 Applicant is a relevant specialist; or
 - 6.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (permanent neonatal diabetes) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction; and
- 2 Patient remains fully compliant and transition to MDI is considered inappropriate by the treating physician; and
- 3 Fither
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has had four severe unexplained hypoglycaemic episodes over a six month period (severe as defined as requiring the assistance of another person); and
- 6 Has an average HbA1c between the following range: equal to or greater than 53 mmol/mol and equal to or less than 90 mmol/mol; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	I Generic
\$	Per 🗸	Manufacturer

continued...

- 7 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 8 Either:
 - 8.1 Applicant is a relevant specialist; or
 - 8.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (severe unexplained hypoglycaemia) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of at least a 50% reduction from baseline in hypoglycaemic events; and
- 2 HbA1c has not increased by more than 5 mmol/mol from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Has undertaken carbohydrate counting education (either a carbohydrate counting course or direct education from an appropriate health professional); and
- 3 Applicant is part of a multidisciplinary team experienced in the management of type 1 diabetes care; and
- 4 Has adhered to an intensive MDI regimen using analogue insulins for at least six months; and
- 5 Has unpredictable and significant variability in blood glucose including significant hypoglycaemia affecting the ability to reduce HbA1: and
- 6 In the opinion of the treating clinician, HbA1c could be reduced by at least 10 mmol/mol using insulin pump treatment; and
- 7 Has typical HbA1c results between the following range: equal to or greater than 65 mmol/mol and equal to or less than 90 mmol/mol; and
- 8 Has been evaluated by the multidisciplinary team for their suitability for insulin pump therapy; and
- 9 Either:
 - 9.1 Applicant is a relevant specialist; or
 - 9.2 Applicant is a nurse practitioner working within their vocational scope.

Renewal — (HbA1c) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to derive benefit according to the treatment plan agreed at induction of achieving and maintaining a reduction in HbA1c from baseline of 10 mmol/mol; and
- 2 The number of severe unexplained recurrent hypoglycaemic episodes has not increased from baseline; and
- 3 Either:
 - 3.1 Applicant is a relevant specialist; or
 - 3.2 Applicant is a nurse practitioner working within their vocational scope.

Initial application — (Previous use before 1 September 2012) only from a relevant specialist or nurse practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has type 1 diabetes or has undergone a pancreatectomy or has cystic fibrosis-related diabetes; and
- 2 Was already on pump treatment prior to 1 September 2012 and had been evaluated by the multidisciplinary team for their suitability for insulin pump therapy at the time of initiating that pump treatment and continues to benefit from pump treatment; and
- 3 The patient has adhered to an intensive MDI regimen using analogue insulins for at least six months prior to initiating

1 OP

1 OP

✓ Sure-T MMT-863

✓ Sure-T MMT-873

ALIMENTARY TRACT AND METABOLISM				
	Subsidy (Manufacturer's Price \$	e) S Per	Fully ubsidised	Brand or Generic Manufacturer
continued				
pump therapy; and 4 The patient is continuing to derive benefit from pump th 5 The patient had achieved and is maintaining a HbA1c o 6 The patient has had no increase in severe unexplained 7 The patient's HbA1c has not deteriorated more than 5 r 8 Either: 8.1 Applicant is a relevant specialist; or	f equal to or less than hypoglycaemic episoo nmol/mol from baselin	des from		
8.2 Applicant is a nurse practitioner working within the Renewal — (Previous use before 1 September 2012) only f		iot or num		anar Annrovala valid for C
years for applications meeting the following criteria:	ioin a reievant specia	ist or riur	se praciiii	orier. Approvais valid for 2
All of the following:				
1 The patient is continuing to derive benefit according to t	he treatment plan and	l has mai	ntained a	HbA1c of equal to or less
than 80 mmol/mol; and 2 The patient's HbA1c has not deteriorated more than 5 r	nmal/mal from initial a	nnlication	o: and	
3 The patient has not had an increase in severe unexplain				ne: and
4 Either:				,
4.1 Applicant is a relevant specialist; or				
4.2 Applicant is a nurse practitioner working within the	neir vocational scope.			
INSULIN PUMP CARTRIDGE - Special Authority see SA1985	on page 19 - Retail	pharmacy	/	
a) Maximum of 3 sets per prescription				
b) Only on a prescriptionc) Maximum of 13 packs of cartridge sets will be funded p	or voar			
Cartridge 300 U, t:lock × 10	50.00	1 OP	✓ T	andem Cartridge
INSULIN PUMP INFUSION SET (STEEL CANNULA) - Specia		35 on pag	ge 19 – Re	etail pharmacy
a) Maximum of 3 sets per prescription	•			, ,
b) Only on a prescription				
 c) Maximum of 13 infusion sets will be funded per year. 10 mm steel needle; 60 cm tubing x 10 	130.00	1 OP	. .	/liniMed Sure-T
To thin steel needle, oo chi tubing x to	130.00	1 01	• 10	MMT-884A
10 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	// IniMed Sure-T
				MMT-886A
6 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	MiniMed Sure-T
Commented and allow 20 are taking at 40	100.00	4 OD		MMT-864A
6 mm steel needle; 80 cm tubing × 10	130.00	1 OP	• 1	/liniMed Sure-T MMT-866A
8 mm steel needle; 60 cm tubing × 10	130.00	1 OP	✓ N	MiniMed Sure-T
		-		MMT-874A
8 mm steel needle; 80 cm tubing × 10	130.00	1 OP	✓ N	IniMed Sure-T
				MMT-876A

6 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

8 mm steel needle; 29 G; manual insertion; 60 cm tubing ×

Subsidy	H	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
 \$	Per	•	Manufacturer

INSULIN PUMP INFUSION SET (STEEL CANNULA, STRAIGHT INSERTION) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 sets per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

6 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 80 cm line x 10 with 10 needles		1 OP	✓ TruSteel
6 mm steel cannula; straight insertion; 60 cm line × 10 with 10 needles	130.00	1 OP	✓ TruSteel
8 mm steel cannula; straight insertion; 60 cm line x 10 with	130 00	1 OP	✓ TruSteel

1 OP

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

INSULIN PUMP INFUSION SET (TEFLON CANNULA) - Special Authority see SA1985 on page 19 - Retail pharmacy

- a) Maximum of 3 set per prescription
- b) Only on a prescription
- c) Maximum of 13 infusion sets will be funded per year.

13 mm teflon needle, 45 cm tubing x 10130.00	1 OP
13 mm teflon needle, 60 cm tubing × 10130.00	1 OP

6 mm teflon needle, 80 cm clear tubing × 10130.00

- ✓ MiniMed Silhouette MMT-382A
- ✓ MiniMed Silhouette MMT-368A
 ✓ MiniMed Silhouette
- MMT-381A

 MiniMed Silhouette

✓ MiniMed Silhouette

- MMT-383A ✓ MiniMed Silhouette MMT-377A
- MMT-378A ✓ MiniMed Silhouette MMT-384A
- 6 mm teflon needle, 110 cm tubing × 10
 10 P

 6 mm teflon needle, 45 cm blue tubing × 10
 130.00
 1 OP
- ✓ MiniMed Quick-Set MMT-398A
 ✓ MiniMed Mio MMT-941A
- ✓ MiniMed Mio MMT-921A
 - ✓ MiniMed Mio MMT-943A
 - ✓ MiniMed Mio MMT-923A
 - ✓ MiniMed Quick-Set MMT-399A
 - ✓ MiniMed Mio MMT-945A
 - ✓ MiniMed Mio MMT-965A
 - ✓ MiniMed Mio MMT-925A
 - ✓ MiniMed Quick-Set MMT-387A
 - ✓ MiniMed Quick-Set MMT-396A
 - ✓ MiniMed Quick-Set MMT-397A
 - ✓ MiniMed Mio MMT-975A
 - ✓ MiniMed Quick-Set MMT-386A

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. *Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price	e) Sub	Fully	Brand or Generic
	\$	Per	1	Manufacturer
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II SA1985 on page 19 – Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year.	NSERTION WITH II	NSERTION	I DEVICE)	- Special Authority see
13 mm teflon cannula; angle insertion; insertion device; 110 line × 10 with 10 needles	cm 140.00	1 OP	√ Διι	toSoft 30
13 mm teflon cannula; angle insertion; insertion device; 60 cl	n	1 OP		toSoft 30
INSULIN PUMP INFUSION SET (TEFLON CANNULA, ANGLE II Retail pharmacy a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 17 mm teflon cannula; angle insertion; 60 cm line × 10 with	NSERTION) - Spe			
10 needles; luer lock	IT INSERTION WIT	1 OP H INSERT		houette MMT-373 ICE) – Special Authority
110 cm line × 10 with 10 needles	m	1 OP		toSoft 90
line x 10 with 10 needles		1 OP		toSoft 90 toSoft 90
9 mm teflon cannula; straight insertion; insertion device; 60 c	m	1 OP		toSoft 90
INSULIN PUMP INFUSION SET (TEFLON CANNULA, STRAIGH Retail pharmacy a) Maximum of 3 sets per prescription		-		
b) Only on a prescription c) Maximum of 13 infusion sets will be funded per year. 6 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock	130.00	1 OP	√ Qu	ick-Set MMT-393
9 mm teflon cannula; straight insertion; 60 cm tubing × 10 wi 10 needles; luer lock		1 OP	√ Qu	ick-Set MMT-392
INSULIN PUMP RESERVOIR – Special Authority see SA1985 c a) Maximum of 3 sets per prescription b) Only on a prescription c) Maximum of 13 packs of reservoir sets will be funded per	n page 19 – Retail _I	oharmacy	-	
10 × luer lock conversion cartridges 1.8 ml for Paradigm pun Cartridge for 5 and 7 series pump; 1.8 ml × 10	ps50.00	1 OP 1 OP	✔ Mir 1	R Cartridge 1.8 niMed .8 Reservoir //MT-326A
Cartridge for 7 series pump; 3.0 ml × 10	50.00	1 OP		niMed 8.0 Reservoir //MT-332A

	Subsidy (Manufacturer's Price \$	e) Per	Fully Subsidised	Brand or Generic Manufacturer	
Digestives Including Enzymes					
PANCREATIC ENZYME					
Cap pancreatin 150 mg (amylase 8,000 Ph Eur U, lipase 10,000 Ph Eur U, total protease 600 Ph Eur U)	34.93	100	v (Creon 10000	
Cap pancreatin (175 mg (25,000 U lipase, 22,500 U amylase 1,250 U protease))		100	✓ !	Panzytrat	
Cap pancreatin 300 mg (amylase 18,000 Ph Eur U, lipase 25,000 Ph Eur U, total protease 1,000 Ph Eur U)	94.38	100	1	Creon 25000	
Modified release granules pancreatin 60.12 mg (amylase 3,600 Ph Eur U, lipase 5,000 Ph Eur U, protease 200 Ph					
Eur U)		20 g O	-	Creon Micro	
(Panzytrat Cap pancreatin (175 mg (25,000 U lipase, 22,500 U al	mylase, 1,250 U pro	otease),) to be deli	isted 1 June 2023)	1
URSODEOXYCHOLIC ACID - Special Authority see SA1739 be	low - Retail pharma	асу			
Cap 250 mg	32.95	100	✓	Ursosan	

⇒SA1739 Special Authority for Subsidy

Initial application — (Alagille syndrome or progressive familial intrahepatic cholestasis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 Patient has been diagnosed with Alagille syndrome; or
- 2 Patient has progressive familial intrahepatic cholestasis.

Initial application — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic severe drug induced cholestatic liver injury; and
- 2 Cholestatic liver injury not due to Total Parenteral Nutrition (TPN) use in adults: and
- 3 Treatment with ursodeoxycholic acid may prevent hospital admission or reduce duration of stay.

Initial application — (**Primary biliary cholangitis**) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Primary biliary cholangitis confirmed by antimitochondrial antibody titre (AMA) > 1:80, and raised cholestatic liver enzymes with or without raised serum IgM or, if AMA is negative, by liver biopsy; and
- 2 Patient not requiring a liver transplant (bilirubin > 100 umol/l; decompensated cirrhosis).

Initial application — (**Pregnancy**) from any relevant practitioner. Approvals valid for 6 months where the patient diagnosed with cholestasis of pregnancy.

Initial application — (Haematological Transplant) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient at risk of veno-occlusive disease or has hepatic impairment and is undergoing conditioning treatment prior to allogenic stem cell or bone marrow transplantation; and
- 2 Treatment for up to 13 weeks.

Initial application — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Paediatric patient has developed abnormal liver function as indicated on testing which is likely to be induced by Total Parenteral Nutrition (TPN); and
- 2 Liver function has not improved with modifying the TPN composition.

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (Chronic severe drug induced cholestatic liver injury) from any relevant practitioner. Approvals valid for 6 months where the patient continues to benefit from treatment.

Renewal — (Pregnancy/Primary biliary cholangitis) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Total parenteral nutrition induced cholestasis) from any relevant practitioner. Approvals valid for 6 months where the paediatric patient continues to require TPN and who is benefiting from treatment, defined as a sustained improvement in bilirubin levels.

Laxatives

Bulk-forming Agents

* Powder for oral soln	6.00	250 g OP	✓ Macro Organic Psyllium Husk
	12.20	500 g OP	✓ Konsyl-D
MUCILAGINOUS LAXATIVES WITH STIMULANTS			
* Dry	6.02	500 g OP	
	(17.32)		Normacol Plus
	2.41	200 g OP	
	(8.72)	-	Normacol Plus

Faecal Softeners

* Tab 120 mg		100 100	✓ <u>Coloxyl</u> ✓ <u>Coloxyl</u>
DOCUSATE SODIUM WITH SENNOSIDES Tab 50 mg with sennosides 8 mg	3.50	200	✓ Laxsol
POLOXAMER – Only on a prescription Not funded for use in the ear.			
* Oral drops 10%	3.98	30 ml OP	✓ Coloxyl

Opioid Receptor Antagonists - Peripheral

ISPAGHULA (PSYLLIUM) HUSK - Only on a prescription

METHYLNALTREXONE BROMIDE - Special Authority	y see SA1691 below – Retail p	harmacy	
Inj 12 mg per 0.6 ml vial	36.00	1	✓ Relistor
	246 00	7	✓ Relistor

⇒SA1691 Special Authority for Subsidy

DOCUSATE SODIUM - Only on a prescription

Initial application — (Opioid induced constipation) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient is receiving palliative care; and
- 2 Either:
 - 2.1 Oral and rectal treatments for opioid induced constipation are ineffective; or
 - 2.2 Oral and rectal treatments for opioid induced constipation are unable to be tolerated.

Subsidised

Fully

Brand or

Generic

Subsidy

(Manufacturer's Price)

	\$	Per	✓ Manufacturer
Osmotic Laxatives			
GLYCEROL * Suppos 3.6 g - Only on a prescription * Suppos 4 g - Only on a prescription		20 20	✓ PSM ✓ Lax-suppositories Glycerol
(PSM Suppos 3.6 g to be delisted 1 February 2023) LACTULOSE – Only on a prescription			diyeeloi
* Oral liq 10 g per 15 ml		500 ml ND SODIUM C	✓ Laevolac HLORIDE
Powder for oral soln 13.125 g with potassium chloride 46.6 sodium bicarbonate 178.5 mg and sodium chloride 350.	mg,	30	✓ <u>Molaxole</u>
SODIUM ACID PHOSPHATE – Only on a prescription Enema 16% with sodium phosphate 8%	2.50	1	✓ Fleet Phosphate Enema
SODIUM CITRATE WITH SODIUM LAURYL SULPHOACETATI Enema 90 mg with sodium lauryl sulphoacetate 9 mg per mi	, ,	escription	Literia
5 ml	•	50	✓ Micolette✓ Micolette-S29 S29
Stimulant Laxatives			
BISACODYL – Only on a prescription * Tab 5 mg	5.80	200	✓ Bisacodyl Viatris✓ Pharmacy Health
* Suppos 10 mg(Pharmacy Health Tab 5 mg to be delisted 1 January 2023)	3.69	10	✓ <u>Lax-Suppositories</u>
SENNA – Only on a prescription * Tab, standardised	2.17 (8.21)	100	Senokot
	0.43 (2.06)	20	Senokot
SODIUM PICOSULFATE – Special Authority see SA2053 below Oral soln 7.5 mg per ml		cy 30 ml OP	✓ Dulcolax SP Drop
■ SA2053 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals val Both:	id for 6 months fo	r applications r	neeting the following criteria:
1 The patient is a child with problematic constipation despit macrogol where practicable; and			· · · · · · · · · · · · · · · · · · ·
2 The patient would otherwise require a high-volume bowe Renewal from any relevant practitioner. Approvals valid for 12 r	0		

Metabolic Disorder Agents

is benefiting from treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

⇒SA1986 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient is aged up to 24 months at the time of initial application and has been diagnosed with infantile Pompe disease; and
- 2 Any of the following:
 - 2.1 Diagnosis confirmed by documented deficiency of acid alpha-glucosidase by prenatal diagnosis using chorionic villus biopsies and/or cultured amniotic cells; or
 - 2.2 Documented deficiency of acid alpha-glucosidase, and urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides; or
 - 2.3 Documented deficiency of acid alpha-glucosidase, and documented molecular genetic testing indicating a disease-causing mutation in the acid alpha-glucosidase gene (GAA gene); or
 - 2.4 Documented urinary tetrasaccharide testing indicating a diagnostic elevation of glucose tetrasaccharides, and molecular genetic testing indicating a disease-causing mutation in the GAA gene; and
- 3 Patient has not required long-term invasive ventilation for respiratory failure prior to starting enzyme replacement therapy (ERT); and
- 4 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by ERT or might be reasonably expected to compromise a response to ERT; and
- 5 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks.

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Alglucosidase alfa to be administered at doses no greater than 20 mg/kg every 2 weeks; and
- 3 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 4 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by ERT; and
- 5 Patient has not developed another medical condition that might reasonably be expected to compromise a response to FRT: and
- 6 There is no evidence of life threatening progression of respiratory disease as evidenced by the needed for > 14 days of invasive ventilation; and
- 7 There is no evidence of new or progressive cardiomyopathy.

ARGININE – Special Authorit	y see SA2042 below – Retail p	oharmacy
-----------------------------	-------------------------------	----------

Tab 1,000 mg	CBS	90	Clinicians
Cap 500 mg		50	✓ Solgar
Powder		400 g	✓ Biomed

⇒SA2042 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to arginine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to arginine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

BETAINE – Special Authority see SA1987 on the next page – Retail pl	harmacy		
Powder for oral soln	575.00	180 g OP	✓ Cystadane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1987 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a confirmed diagnosis of homocystinuria; and
- 2 Any of the following:
 - 2.1 A cystathionine beta-synthase (CBS) deficiency; or
 - 2.2 A 5.10-methylene-tetrahydrofolate reductase (MTHFR) deficiency; or
 - 2.3 A disorder of intracellular cobalamin metabolism; and
- 3 An appropriate homocysteine level has not been achieved despite a sufficient trial of appropriate vitamin supplementation.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

COENZYME Q10 - Special Authority see SA2039 below -	- Retail pharmacy		
Cap 120 mg	CBS	30	✓ Solgar
Cap 160 mg	CBS	60	✓ Go Healthy

⇒SA2039 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to coenzyme Q10 supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to coenzyme Q10 supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

```
GALSULFASE – Special Authority see SA1988 below – Retail pharmacy
Inj 1 mg per ml, 5 ml vial.......2,234.00 1 ✓ Naglazyme
```

⇒SA1988 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has been diagnosed with mucopolysaccharidosis VI; and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of N-acetyl-galactosamine-4-sulfatase (arylsulfatase B) deficiency by either enzyme activity assay in leukocytes or skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations and patient has a sibling who is known to have mucopolysaccharidosis VI

Renewal only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The treatment remains appropriate for the patient and the patient is benefiting from treatment; and
- 2 Patient has not had severe infusion-related adverse reactions which were not preventable by appropriate pre-medication and/or adjustment of infusion rates; and
- 3 Patient has not developed another life threatening or severe disease where the long term prognosis is unlikely to be influenced by Enzyme Replacement Therapy (ERT); and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT.

DURSULFASE - Special Authority see SA1623 on the	e next page – Retail pharmacy		
Inj 2 mg per ml, 3 ml vial	4,608.30	1	Elaprase

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic
\$	Per	✓	Manufacturer

⇒SA1623 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hunter Syndrome (mucopolysaccharidosis II); and
- 2 Either:
 - 2.1 Diagnosis confirmed by demonstration of iduronate 2-sulfatase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of a disease causing mutation in the iduronate 2-sulfatase gene; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with idursulfase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Idursulfase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 weeks post-HSCT) at doses no greater than 0.5 mg/kg every week.

⇒SA1695 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 24 weeks for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with Hurler Syndrome (mucopolysacchardosis I-H); and
- 2 Fither:
 - 2.1 Diagnosis confirmed by demonstration of alpha-L-iduronidase deficiency in white blood cells by either enzyme assay in cultured skin fibroblasts; or
 - 2.2 Detection of two disease causing mutations in the alpha-L-iduronidase gene and patient has a sibling who is known to have Hurler syndrome; and
- 3 Patient is going to proceed with a haematopoietic stem cell transplant (HSCT) within the next 3 months and treatment with laronidase would be bridging treatment to transplant; and
- 4 Patient has not required long-term invasive ventilation for respiratory failure prior to starting Enzyme Replacement Therapy (ERT); and
- 5 Laronidase to be administered for a total of 24 weeks (equivalent to 12 weeks pre- and 12 post-HSCT) at doses no greater than 100 units/kg every week.

LEVOCARNITINE - Special Authority see SA2040 below - Retail pharmacy Tab 500 mgCBS 30 ✓ Solgar 30 ✓ Solgar 60 ✓ Balance Oral lig 1 g per 10 mlCBS ✓ Carnitor S29 118 ml Oral lig 500 mg per 10 mlCBS 300 ml ✓ Balance

⇒SA2040 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to carnitine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to carnitine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

RIBOFLAVIN - Special Authority see SA2041 on the next page - Retail pharmacy
Tab 100 mgCBS 100
Cap 100 mgCBS 100
✓ Country Life
✓ Solgar

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per Manufacturer

⇒SA2041 Special Authority for Subsidy

Initial application only from a metabolic physician or neurologist. Approvals valid for 6 months where patient has a suspected inborn error of metabolism that may respond to riboflavin supplementation.

Renewal only from a metabolic physician or neurologist. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The patient has a confirmed diagnosis of an inborn error of metabolism that responds to riboflavin supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

⇒SA1989 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 1 month for applications meeting the following criteria: All of the following:

- 1 Patient has phenylketonuria (PKU) and is pregnant or actively planning to become pregnant; and
- 2 Treatment with sapropterin is required to support management of PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Following the initial one-month approval, the patient has demonstrated an adequate response to a 2 to 4 week trial of sapropterin with a clinically appropriate reduction in phenylalanine levels to support management of PKU during pregnancy; or
 - 1.2 On subsequent renewal applications, the patient has previously demonstrated response to treatment with sapropterin and maintained adequate phenylalanine levels to support management of PKU during pregnancy; and
- 2 Any of the following:
 - 2.1 Patient continues to be pregnant and treatment with sapropterin will not continue after delivery; or
 - 2.2 Patient is actively planning a pregnancy and this is the first renewal for treatment with sapropterin; or
 - 2.3 Treatment with sapropterin is required for a second or subsequent pregnancy to support management of their PKU during pregnancy; and
- 3 Sapropterin to be administered at doses no greater than a total daily dose of 20 mg/kg; and
- 4 Sapropterin to be used alone or in combination with PKU dietary management; and
- 5 Total treatment duration with sapropterin will not exceed 22 months for each pregnancy (includes time for planning and becoming pregnant) and treatment will be stopped after delivery.

⇒SA1599 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM PHENYLBUTYRATE – Special Authority see SA1990 on the next page – Retail pharmacy Grans 483 mg per g.......2,016.00 174 g OP ✓ Pheburane

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1990 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months where the patient has a diagnosis of a urea cycle disorder involving a deficiency of carbamylphosphate synthetase, ornithine transcarbamylase or argininosuccinate synthetase.

Renewal only from a metabolic physician. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

TAURINE - Special Authority see SA2043 below - Retail pharmacy

Cap 500 mg	CBS	50	✓ Solgar
Cap 1,000 mg	CBS	90	✓ Life Extension
Powder	CBS	300 g	 Life Extension

⇒SA2043 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 6 months where patient has a suspected specific mitochondrial disorder that may respond taurine supplementation.

Renewal only from a metabolic physician. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 The patient has confirmed diagnosis of a specific mitochondrial disorder which responds to taurine supplementation; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Gaucher's Disease

TALIGLUCERASE ALFA - Special Authority see SA2137 below - Retail pharmacy ✓ Elelvso

⇒SA2137 Special Authority for Subsidy

Initial application only from a metabolic physician. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has a diagnosis of symptomatic type 1 or type 3* Gaucher disease confirmed by the demonstration of specific deficiency of glucocerebrosidase in leukocytes or cultured skin fibroblasts, and genotypic analysis; and
- 2 Patient does not have another life-threatening or severe disease where the prognosis is unlikely to be influenced by enzyme replacement therapy (ERT) or the disease might be reasonably expected to compromise a response to ERT; and
- 3 Any of the following:
 - 3.1 Patient has haematological complications of Gaucher disease; or
 - 3.2 Patient has skeletal complications of Gaucher disease; or
 - 3.3 Patient has significant liver dysfunction or hepatomegaly attributable to Gaucher disease; or
 - 3.4 Patient has reduced vital capacity from clinically significant or progressive pulmonary disease due to Gaucher disease: or
 - 3.5 Patient is a child and has experienced growth failure with significant decrease in percentile linear growth over a 6-12 month period; and
- 4 Taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Note: Indication marked with * is an unapproved indication

Renewal only from a metabolic physician or any relevant practitioner on the recommendation of a metabolic physician. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Patient has demonstrated a symptomatic improvement and has maintained improvements in the main symptom or symptoms for which therapy was started; and
- 2 Patient has demonstrated a clinically objective improvement or no deterioration in haemoglobin levels, platelet counts and liver and spleen size: and

Subsidy	ce)	Fully	Brand or	
(Manufacturer's Pri		Subsidised	Generic	
\$	Per	· /	Manufacturer	

continued...

- 3 Radiological (MRI) signs of bone activity performed at two years since initiation of treatment, and five yearly thereafter, demonstrate no deterioration shown by the MRI, compared with MRI taken immediately prior to commencement of therapy or adjusted dose; and
- 4 Patient has not developed another medical condition that might reasonably be expected to compromise a response to ERT; and
- 5 Patient is adherent with regular treatment and taliglucerase alfa is to be administered at a dose no greater than 30 unit/kg every other week rounded to the nearest whole vial (200 units).

Mouth and Throat

Agents Used in Mouth Ulceration

Agents used in Mouth diceration			
BENZYDAMINE HYDROCHLORIDE Soln 0.15% – Higher subsidy of \$20.31 per 500 ml with Endorsement	(20.31)	500 ml	Difflam eatment for cancer, and the
prescription is endorsed accordingly.			
CARMELLOSE SODIUM WITH GELATIN AND PECTIN			
Paste	17.20	56 g OP	✓ Stomahesive
	4.55	15 g OP	
	(7.90)		Orabase
	1.52	5 g OP	0.1
Douder	(3.60)	00 ~ OD	Orabase
Powder	(10.95)	28 g OP	Stomahesive
CHOLINE CALICYLATE WITH CETALIZONII IN CHILODIDE	(10.93)		Otomanesive
CHOLINE SALICYLATE WITH CETALKONIUM CHLORIDE * Adhesive gel 8.7% with cetalkonium chloride 0.01%	2.06	15 g OP	
Adilesive gel 6.7 % with cetalkoritum chiloride 0.01 %	(6.00)	15 g OF	Bonjela
TRIAMCINOLONE ACETONIDE	(0.00)		Donjoid
Paste 0.1%	5.33	5 g OP	✓ Kenalog in Orabase
1 4010 0.170		0 y 0.	- Itolialog III Olabado
Oropharyngeal Anti-infectives			
AMPHOTERICIN B			
Lozenges 10 mg	5.86	20	✓ Fungilin
MICONAZOLE			
Oral gel 20 mg per g	4.74	40 g OP	✓ <u>Decozol</u>
NYSTATIN			
Oral liq 100,000 u per ml	1.76	24 ml OP	✓ <u>Nilstat</u>

	(Manufacturer's Price	Per	✓ Manufacturer
Vitamins			
Vitamin B			
HYDROXOCOBALAMIN * Inj 1 mg per ml, 1 ml ampoule – Up to 6 inj available on a Pt	SO1.89 2.46	3	✓ Vita-B12✓ Hydroxocobalamin Panpharma
	2.84 3.15	5	✓ Neo-B12✓ Hydroxocobalamin Mercury Pharma
(Vita-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 Novemb (Neo-B12 Inj 1 mg per ml, 1 ml ampoule to be delisted 1 Novemb (Hydroxocobalamin Mercury Pharma Inj 1 mg per ml, 1 ml ampou PYRIDOXINE HYDROCHLORIDE a) No more than 100 mg per dose b) Only on a prescription	oer 2022)	November 20	022)
* Tab 25 mg - No patient co-payment payable Tab 50 mg		90 500	✓ <u>Vitamin B6 25</u> ✓ Pyridoxine multichem
THIAMINE HYDROCHLORIDE – Only on a prescription * Tab 50 mg VITAMIN B COMPLEX		100	✓ Max Health
* Tab, strong, BPC	7.15	500	✓ Bplex
Vitamin C			
ASCORBIC ACID a) No more than 100 mg per dose b) Only on a prescription			
* Tab 100 mg	12.50	500	✓ Cvite
Vitamin D			
# Cap 1 mcg		100 100	✓ One-Alpha ✓ One-Alpha ✓ One-Alpha S29 S29
* Oral drops 2 mcg per ml CALCITRIOL		20 ml OP	✓ One-Alpha
* Cap 0.25 mcg Calcitriol-AFT to be Principal Supply on 1 December 202		100	✓ Calcitriol-AFT
* Cap 0.5 mcg Calcitriol-AFT to be Principal Supply on 1 December 202 COLECALCIFEROL		100	✓ Calcitriol-AFT
** Cap 1.25 mg (50,000 iu) – Maximum of 12 cap per prescript ** Oral liq 188 mcg per ml (7,500 iu per ml)		12 4.8 ml OP	✓ <u>Vit.D3</u> ✓ Puria

Subsidy

(Manufacturer's Price)

Fully

Subsidised

Brand or

Generic

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer **Multivitamin Preparations** MULTIVITAMIN RENAL - Special Authority see SA1546 below - Retail pharmacy 30 ✓ Clinicians Renal Vit. ⇒SA1546 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Fither: 1 The patient has chronic kidney disease and is receiving either peritoneal dialysis or haemodialysis; or 2 The patient has chronic kidney disease grade 5, defined as patient with an estimated glomerular filtration rate of < 15 ml/min/1.73 m² body surface area (BSA). MULTIVITAMINS - Special Authority see SA1036 below - Retail pharmacy 200 g OP ✓ Paediatric Seravit ⇒SA1036 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where patient has had a previous approval for multivitamins. **VITAMINS** 1.000 ✓ Myite * Cap (fat soluble vitamins A, D, E, K) - Special Authority see Vitabdeck ⇒SA1720 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 Patient has cystic fibrosis with pancreatic insufficiency; or 2 Patient is an infant or child with liver disease or short gut syndrome; or 3 Patient has severe malabsorption syndrome. Minerals Calcium **CALCIUM CARBONATE** 250 ✓ Calci-Tab 500 * Tab eff 1.25 g (500 mg elemental) - Subsidy by endorsement......260.00 100 ✓ Calcium 500 mg Hexal S29 Subsidy by endorsement - Only when prescribed for paediatric patients (< 5 years) where calcium carbonate oral liquid is considered unsuitable. **CALCIUM GLUCONATE** 10 ✓ Max Health -Hameln S29 64.00 20 Max Health \$29 Fluoride

100

✓ PSM

(PSM Tab 1.1 mg (0.5 mg elemental) to be delisted 1 March 2023)

*Three months or six months, as applicable, dispensed all-at-once

SODIUM FLUORIDE

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
lodine				
POTASSIUM IODATE * Tab 253 mcg (150 mcg elemental iodine)	4.58	90	✓ <u>N</u>	euroTabs
Iron				
FERROUS FUMARATE * Tab 200 mg (65 mg elemental)	3.04	100	√ F	erro-tab
FERROUS FUMARATE WITH FOLIC ACID * Tab 310 mg (100 mg elemental) with folic acid 350 mcg		100	✓ F	erro-F-Tabs
FERROUS SULFATE			_	
* Tab long-acting 325 mg (105 mg elemental)* Oral liq 30 mg (6 mg elemental) per 1 ml		30 500 ml	_	errograd erodan
IRON (AS FERRIC CARBOXYMALTOSE) – Special Authority so Inj 50 mg per ml, 10 ml vial		etail p		erinject
⇒SA1840 Special Authority for Subsidy				•

Initial application — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Rapid correction of anaemia is required.

Renewal — (serum ferritin less than or equal to 20 mcg/L) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia with a serum ferritin level of less than or equal to 20 mcg/L; and
- 2 A re-trial with oral iron is clinically inappropriate.

Initial application — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with iron-deficiency anaemia; and
- 2 Any of the following:
 - 2.1 Patient has been compliant with oral iron treatment and treatment has proven ineffective; or
 - 2.2 Treatment with oral iron has resulted in dose-limiting intolerance; or
 - 2.3 Patient has symptomatic heart failure, chronic kidney disease stage 3 or more or active inflammatory bowel disease and a trial of oral iron is unlikely to be effective: or
 - 2.4 Rapid correction of anaemia is required.

Renewal — (iron deficiency anaemia) only from an internal medicine physician, obstetrician, gynaecologist, anaesthetist or medical practitioner on the recommendation of a internal medicine physician, obstetrician, gynaecologist or anaesthetist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient continues to have iron-deficiency anaemia; and
- 2 A re-trial with oral iron is clinically inappropriate.

IRON POLYMALTOSE

Principal Supply

ALIMENTARY TRACT AND METABOLISM

	Subsidy (Manufacturer's Price \$	e) Sub Per	Fully sidised	Brand or Generic Manufacturer
Magnesium				
MAGNESIUM HYDROXIDE Suspension 8%	33.60	355 ml	√ P	hillips Milk of Magnesia S29
MAGNESIUM SULPHATE * Inj 2 mmol per ml, 5 ml ampoule	25.53	10	✓ M	lartindale
Zinc				
ZINC SULPHATE * Cap 137.4 mg (50 mg elemental)	11.00	100	✓ Z	incaps

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

Antianaemics

Hypoplastic and Haemolytic

⇒SA1775 Special Authority for Subsidy

Initial application — (chronic renal failure) from any specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient in chronic renal failure: and
- 2 Haemoglobin is less than or equal to 100g/L; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient does not have diabetes mellitus; and
 - 3.1.2 Glomerular filtration rate is less than or equal to 30ml/min; or
 - 3.2 Both:
 - 3.2.1 Patient has diabetes mellitus; and
 - 3.2.2 Glomerular filtration rate is less than or equal to 45ml/min; or
 - 3.3 Patient is on haemodialysis or peritoneal dialysis.

Initial application — (myelodysplasia) from any specialist. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of myelodysplasia (MDS)*; and
- 2 Has had symptomatic anaemia with haemoglobin < 100g/L and is red cell transfusion-dependent; and
- 3 Patient has very low, low or intermediate risk MDS based on the WHO classification based prognostic scoring system for myelodysplastic syndrome (WPSS); and
- 4 Other causes of anaemia such as B12 and folate deficiency have been excluded; and
- 5 Patient has a serum epoetin level of < 500 IU/L; and
- 6 The minimum necessary dose of epoetin would be used and will not exceed 80,000 iu per week.

Note: Indication marked with * is an unapproved indication

Renewal — (chronic renal failure) from any specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (myelodysplasia) from any specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient's transfusion requirement continues to be reduced with erythropoietin treatment; and
- 2 Transformation to acute myeloid leukaemia has not occurred; and
- 3 The minimum necessary dose of epoetin would be used and will not exceed 80,000 ju per week.

Note: Indication marked with * is an unapproved indication

EPOETIN ALFA - Special Authority see SA1775 above - Retail pharmacy

Wastage claimable			
Inj 1,000 iu in 0.5 ml, syringe	250.00	6	Binocrit
Inj 2,000 iu in 1 ml, syringe		6	Binocrit
Inj 3,000 iu in 0.3 ml, syringe		6	Binocrit
Inj 4,000 iu in 0.4 ml, syringe		6	Binocrit
Inj 5,000 iu in 0.5 ml, syringe		6	Binocrit
Inj 6,000 iu in 0.6 ml, syringe		6	Binocrit
Inj 8,000 iu in 0.8 ml, syringe		6	Binocrit
Inj 10,000 iu in 1 ml, syringe		6	Binocrit
Inj 40,000 iu in 1 ml, syringe		1	Binocrit

	Subsidy (Manufacturer's Price) \$		ully sed	Brand or Generic Manufacturer
Megaloblastic				
FOLIC ACID Tab 0.8 mg	26.60	1,000		olic Acid multichem
* Tab 5 mg Oral liq 50 mcg per ml		100 5 ml OP		olic Acid Mylan iomed

Antifibrinolytics, Haemostatics and Local Sclerosants

EFTRENONACOG ALFA [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia B receiving prophylaxis treatment. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management group.

Treaters Group in conjunction with the National F	Haemophilia Management grou	ıp.	
Inj 250 iu vial	612.50	1	Alprolix
Inj 500 iu vial	1,225.00	1	✓ Alprolix
Inj 1,000 iu vial	2,450.00	1	✓ Alprolix
Inj 2,000 iu vial	4,900.00	1	✓ Alprolix
Inj 3,000 iu vial	7,350.00	1	✓ Alprolix
Inj 4,000 iu vial	9,800.00	1	✓ Alprolix
ELTROMBOPAG – Special Authority see SA1743 be Wastage claimable	elow – Retail pharmacy		
Tab 25 mg	1,550.00	28	Revolade
Tab 50 mg	3,100.00	28	Revolade

⇒SA1743 Special Authority for Subsidy

Initial application — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has had a splenectomy; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Any of the following:
 - 3.1 Patient has a platelet count of 20,000 to 30,000 platelets per microlitre and has evidence of significant mucocutaneous bleeding; or
 - 3.2 Patient has a platelet count of less than or equal to 20,000 platelets per microlitre and has evidence of active bleeding; or
 - 3.3 Patient has a platelet count of less than or equal to 10,000 platelets per microlitre.

Initial application — (idiopathic thrombocytopenic purpura - preparation for splenectomy) only from a haematologist. Approvals valid for 6 weeks where the patient requires eltrombopag treatment as preparation for splenectomy. Initial application — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a significant and well-documented contraindication to splenectomy for clinical reasons; and
- 2 Two immunosuppressive therapies have been trialled and failed after therapy of 3 months each (or 1 month for rituximab); and
- 3 Either:
 - 3.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per microliter: or
 - 3.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre

continued...

Subsic		Fully	Brand or
(Manufacture	r's Price) Su	ubsidised	Generic
\$	Per	✓	Manufacturer

continued...

and significant mucocutaneous bleeding.

Initial application — (severe aplastic anaemia) only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Two immunosuppressive therapies have been trialled and failed after therapy of at least 3 months duration; and
- 2 Either:
 - 2.1 Patient has severe aplastic anaemia with a platelet count of less than or equal to 20,000 platelets per microliter; or
 - 2.2 Patient has severe aplastic anaemia with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding.

Renewal — (idiopathic thrombocytopenic purpura - post-splenectomy) only from a haematologist. Approvals valid for 12 months where the patient has obtained a response (see Note) from treatment during the initial approval or subsequent renewal periods and further treatment is required.

Note: Response to treatment is defined as a platelet count of > 30,000 platelets per microlitre.

Renewal — (idiopathic thrombocytopenic purpura contraindicated to splenectomy) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient's significant contraindication to splenectomy remains; and
- 2 The patient has obtained a response from treatment during the initial approval period; and
- 3 Patient has maintained a platelet count of at least 50,000 platelets per microlitre on treatment; and
- 4 Further treatment with eltrombopag is required to maintain response.

Renewal — (severe aplastic anaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has obtained a response from treatment of at least 20,000 platelets per microlitre above baseline during the initial approval period; and
- 2 Platelet transfusion independence for a minimum of 8 weeks during the initial approval period.

EMICIZUMAB - [Xpharm] - Special Authority see SA1969 below

Inj 30 mg in 1 ml vial	3,570.00	1	✓ Hemlibra
Inj 60 mg in 0.4 ml vial	7,138.00	1	✓ Hemlibra
Inj 105 mg in 0.7 ml vial		1	✓ Hemlibra
Inj 150 mg in 1 ml vial	17,846.00	1	✓ Hemlibra

⇒SA1969 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe congenital haemophilia A and history of bleeding and bypassing agent usage within the last six months; and
- 2 Either:
 - 2.1 Patient has had greater than or equal to 6 documented and treated spontaneous bleeds within the last 6 months if on an on-demand bypassing agent regimen; or
 - 2.2 Patient has had greater than or equal to 2 documented and treated spontaneous bleeds within the last 6 months if on a bypassing agent prophylaxis regimen; and
- 3 Patient has a high-titre inhibitor to Factor VIII (greater than or equal to 5 Bethesda units per ml) which has persisted for six months or more; and
- 4 There is no immediate plan for major surgery within the next 12 months; and
- 5 Either:
 - 5.1 Patient has failed immune tolerance induction (ITI) after an initial period of 12 months; or

continued...

Subsidy (Manufacturer's Price)	Fı Subsidis	ully	Brand or Generic
\$	Per	✓	Manufacturer

continued...

- 5.2 The Haemophilia Treaters Group considers the patient is not a suitable candidate for ITI; and
 - 6 Treatment is to be administered at a maximum dose of 3 mg/kg weekly for 4 weeks followed by the equivalent of 1.5 mg/kg weekly.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient has had no more than two spontaneous and clinically significant treated bleeds after the end of the loading dose period (i.e. after the first four weeks of treatment until the end of the 24-week treatment period); and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

EPTACOG ALFA [RECOMBINANT FACTOR VIIA] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group. Rare Clinical Circumstances Brand of bypassing agent for > 14 days predicted use. Access to funded treatment for > 14 days predicted use is by named patient application to the Haemophilia Treaters Group, subject to access criteria.

Inj 1 mg syringe	1,178.30	1	✓ NovoSeven RT
Inj 2 mg syringe	·	1	✓ NovoSeven RT
Inj 5 mg syringe	5,891.50	1	✓ NovoSeven RT
Inj 8 mg syringe	9,426.40	1	✓ NovoSeven RT

FACTOR EIGHT INHIBITOR BYPASSING FRACTION - [Xpharm]

For patients with haemophilia. Preferred Brand of bypassing agent for > 14 days predicted use. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 U	1,315.00	1	✓ FĚIBA NF
Inj 1,000 U	2,630.00	1	✓ FEIBA NF
Inj 2,500 U	6,575.00	1	✓ FEIBA NF

MOROCTOCOG ALFA [RECOMBINANT FACTOR VIII] - [Xpharm]

For patients with haemophilia. Rare Clinical Circumstances Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group, subject to criteria

Inj 250 iu prefilled syringe	287.50	1	Xyntha
Inj 500 iu prefilled syringe		1	Xyntha
Inj 1,000 iu prefilled syringe		1	Xyntha
Inj 2,000 iu prefilled syringe		1	Xyntha
Inj 3,000 iu prefilled syringe		1	Xyntha

NONACOG GAMMA, [RECOMBINANT FACTOR IX] - [Xpharm]

For patients with haemophilia. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

Inj 500 iu vial	1	✓ RIXUBIS
Inj 1,000 iu vial	1	✓ RIXUBIS
Inj 2,000 iu vial	1	✓ RIXUBIS
Inj 3,000 iu vial2,610.00	1	✓ RIXUBIS

OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (ADVATE) - [Xpharm]

For patients with haemophilia. Preferred Brand of short half-life recombinant factor VIII. Access to funded treatment is managed by the Haemophilia Treaters Group in conjunction with the National Haemophilia Management Group.

managed by the naemophilia freaters Group in	i conjunction with the National H	aemopniia	Management G
Inj 250 iu vial	210.00	1	✓ Advate
Inj 500 iu vial	420.00	1	Advate
Inj 1,000 iu vial	840.00	1	Advate
Inj 1,500 iu vial	1,260.00	1	Advate
Inj 2,000 iu vial		1	Advate
Inj 3,000 iu vial	2,520.00	1	Advate
•			

	Subsidy (Manufacturer's Price)	Su	Fully bsidised	Brand or Generic
	\$	Per	√	Manufacturer
OCTOCOG ALFA [RECOMBINANT FACTOR VIII] (KOGENATE	FS) - [Xpharm]			
For patients with haemophilia. Rare Clinical Circumstances	Brand of short half-lif	e recom	binant fac	ctor VIII. Access to funder
treatment is managed by the Haemophilia Treaters Group in	conjunction with the	National	Haemop	hilia Management Group
subject to criteria.				
Inj 250 iu vial		1		Kogenate FS
Inj 500 iu vial		1		Cogenate FS
Inj 1,000 iu vial		1		Kogenate FS
Inj 2,000 iu vial		1		Cogenate FS
Inj 3,000 iu vial	•	1	✓ K	Cogenate FS
RURIOCTOCOG ALFA PEGOL [RECOMBINANT FACTOR VIII]				
For patients with haemophilia A receiving prophylaxis treatments		d treatm	ent is ma	anaged by the Haemophil
Treaters Group in conjunction with the National Haemophilia	0 0 1			
Inj 250 iu vial		1		Adynovate
Inj 500 iu vial		1		Adynovate
Inj 1,000 iu vial	*	1		Adynovate
Inj 2,000 iu vial	2,400.00	1	✓	Adynovate
SODIUM TETRADECYL SULPHATE				
* Inj 3% 2 ml		5		
	(73.00)		F	Fibro-vein
FRANEXAMIC ACID				
Tab 500 mg	9.45	60	✓ N	Mercury Pharma
Vitamin K				
PHYTOMENADIONE	0.00	_		
Inj 2 mg per 0.2 ml — Up to 5 inj available on a PSO		5		Conakion MM
Inj 10 mg per ml, 1 ml - Up to 5 inj available on a PSO	9.21	5	✓ K	Conakion MM
Antithrombotic Agents				
Antiplatelet Agents				
ASPIRIN				
% Таb 100 mg	10.00	990	./ 5	Ethioo Aonirin EC
•	10.80	990	• -	Ethics Aspirin EC
CLOPIDOGREL				
* Tab 75 mg	4.60	84	✓ (Clopidogrel
				Multichem
DIPYRIDAMOLE				
* Tab long-acting 150 mg	10.90	60	✓ F	Pytazen SR
FICAGRELOR – Special Authority see SA1955 below – Retail p	harmacy			
* Tab 90 mg	•	56	✓ T	Ticagrelor Sandoz
-	90.00		✓ E	Brilinta
Brilinta Tab 90 mg to be delisted 1 March 2023)				
⇒SA1955 Special Authority for Subsidy				
nitial application — (acute coronary syndrome) from any rel	levant practitioner Δι	nrovals	valid for	12 months for application

1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary

continued...

meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)			Brand or Generic	
(Waitulatule 31 Noe)	Per	oubsidised ✓	Manufacturer	

continued...

syndrome: and

2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Initial application — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Patient has had a neurological stenting procedure* in the last 60 days; or
 - 1.2 Patient is about to have a neurological stenting procedure performed*; and
- 2 Fither:
 - 2.1 Patient has demonstrated clopidogrel resistance using the P2Y12 (VerifyNow) assay or another appropriate platelet function assay and requires antiplatelet treatment with ticagrelor; or
 - 2.2 Fither:
 - 2.2.1 Clopidogrel resistance has been demonstrated by the occurrence of a new cerebral ischemic event; or
 - 2.2.2 Clopidogrel resistance has been demonstrated by the occurrence of transient ischemic attack symptoms referable to the stent.

Initial application — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Initial application — (Stent thrombosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has experienced cardiac stent thrombosis whilst on clopidogrel.

Renewal — (subsequent acute coronary syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient has recently (within the last 60 days) been diagnosed with an ST-elevation or a non-ST-elevation acute coronary syndrome; and
- 2 Fibrinolytic therapy has not been given in the last 24 hours and is not planned.

Renewal — (thrombosis prevention neurological stenting) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient is continuing to benefit from treatment; and
- 2 Treatment continues to be clinically appropriate.

Renewal — (Percutaneous coronary intervention with stent deployment) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone percutaneous coronary intervention; and
- 2 Patient has had a stent deployed in the previous 4 weeks; and
- 3 Patient is clopidogrel-allergic**.

Notes: indications marked with * are unapproved indications.

Note: ** Clopidogrel allergy is defined as a history of anaphylaxis, urticaria, generalised rash or asthma (in non-asthmatic patients) developing soon after clopidogrel is started and is considered unlikely to be caused by any other treatment.

	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

Heparin and Antagonist Preparations

ENOXAPARIN SODIUM - Special Authority see SA2	152 below – Retail pharmacy		
Inj 20 mg in 0.2 ml syringe	31.28	10	Clexane
Inj 40 mg in 0.4 ml syringe	42.49	10	Clexane
Inj 60 mg in 0.6 ml syringe	60.67	10	Clexane
Inj 80 mg in 0.8 ml syringe	80.89	10	Clexane
Inj 100 mg in 1 ml syringe		10	Clexane
Inj 120 mg in 0.8 ml syringe		10	 Clexane Forte
Inj 150 mg in 1 ml syringe	143.86	10	Clexane Forte

⇒SA2152 Special Authority for Subsidy

Initial application — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patients pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Initial application — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 For the short-term treatment of venous thromboembolism prior to establishing a therapeutic level of oral anti-coagulant treatment; or
- 2 For the prophylaxis and treatment of venous thromboembolism in high risk surgery; or
- 3 To enable cessation/re-establishment of existing oral anticoagulant treatment pre/post surgery; or
- 4 For the prophylaxis and treatment of venous thromboembolism in Acute Coronary Syndrome surgical intervention; or
- 5 To be used in association with cardioversion of atrial fibrillation.

Initial application — (Short-term use during treatment of COVID-19 with nirmatrelvir with ritonavir) from any relevant practitioner. Approvals valid for 2 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is receiving an anticoagulation treatment that has drug/drug interactions with ritonavir that increases risk of bleeding; and
- 2 Patient meets the Access Criteria for COVID-19 antivirals published on the Pharmac website*; and
- 3 Other antiviral treatments for COVID-19 have been considered and are not clinically suitable options.

Renewal — (Pregnancy, Malignancy or Haemodialysis) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Low molecular weight heparin treatment is required during a patient's pregnancy; or
- 2 For the treatment of venous thromboembolism where the patient has a malignancy; or
- 3 For the prevention of thrombus formation in the extra-corporeal circulation during haemodialysis.

Renewal — (Venous thromboembolism other than in pregnancy or malignancy) from any relevant practitioner. Approvals valid for 1 month where low molecular weight heparin treatment or prophylaxis is required for a second or subsequent event (surgery, ACS, cardioversion, or prior to oral anti-coagulation).

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	\$	Per	✓ Manufacturer
HEPARIN SODIUM			
Inj 1,000 iu per ml, 5 ml ampoule	72.84	50	✓ Pfizer
Inj 5,000 iu per ml, 1 ml	32.66	5	DBL Heparin
			Sodium S29
	70.33		✓ Hospira
Inj 5,000 iu per ml, 5 ml ampoule	289.05	50	✓ Pfizer
Inj 25,000 iu per ml, 0.2 ml		5	✓ Hospira
•	42.40		✓ Heparin DBL S29
	482.20	50	✓ Heparin DBL S29
HEPARINISED SALINE	.02.20	-	
Inj 10 iu per ml, 5 ml	65.49	50	✓ Pfizer
inj to iu per mi, 5 mi	03.40	50	▼ FIIZEI
Oral Anticoagulants			
DABIGATRAN			
Cap 75 mg - No more than 2 cap per day	76.36	60	✓ Pradaxa
Cap 110 mg	76.36	60	✓ Pradaxa
Cap 150 mg	76.36	60	✓ Pradaxa
RIVAROXABAN			
Tab 10 mg - No more than 1 tab per day	83.10	30	✓ Xarelto
Tab 15 mg - Up to 14 tab available on a PSO		28	✓ Xarelto
Tab 20 mg		28	✓ Xarelto
WARFARIN SODIUM			
Note: Marevan and Coumadin are not interchangeable.			
* Tab 1 mg	3.46	50	✓ Coumadin
· · · · · · · · · · · · · · · · · · ·	6.46	100	✓ Marevan
* Tab 2 mg	4.31	50	✓ Coumadin
* Tab 3 mg		100	✓ Marevan
* Tab 5 mg	5.93	50	✓ Coumadin
-	11.48	100	✓ Marevan

FILGRASTIM - Special Authority see SA1259 below - Retail pharmacy Inj 300 mcg per 0.5 ml prefilled syringe.......96.22 ✓ Nivestim 10 ✓ Nivestim 10

⇒SA1259 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 20%*); or
- 2 Peripheral blood stem cell mobilisation in patients undergoing haematological transplantation; or
- 3 Peripheral blood stem cell mobilisation or bone marrow donation from healthy donors for transplantation; or
- 4 Treatment of severe chronic neutropenia (ANC < 0.5 ×10⁹/L); or
- 5 Treatment of drug-induced prolonged neutropenia (ANC < 0.5 ×10⁹/L).

Note: *Febrile neutropenia risk greater than or equal to 20% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

PEGFILGRASTIM - Special Authority see SA1912 on the next page - Retail pharmacy

1 ✓ Neulastim

Subsidy		Fully	Brand or	
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1912 Special Authority for Subsidy

Initial application only from a relevant specialist, vocationally registered general practitioner or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where used for prevention of neutropenia in patients undergoing high risk chemotherapy for cancer (febrile neutropenia risk greater than or equal to 5%*). Note: *Febrile neutropenia risk greater than or equal to 5% after taking into account other risk factors as defined by the European Organisation for Research and Treatment of Cancer (EORTC) guidelines.

Fluids and Electrolytes

Intravenous Administration

GLUCOSE [DEXTROSE] * Inj 50%, 10 ml ampoule – Up to 5 inj available on a PSO * Inj 50%, 90 ml bottle – Up to 5 inj available on a PSO		5 1	✓ Biomed ✓ Biomed
POTASSIUM CHLORIDE * Inj 75 mg per ml, 10 ml	65.00	50	✓ Juno
SODIUM BICARBONATE Inj 8.4%, 50 ml	21.40	1	✓ Biomed
a) Up to 5 inj available on a PSO b) Not in combination Inj 8.4%, 100 ml	01 OF	1	✓ Biomed
a) Up to 5 inj available on a PSO b) Not in combination	21.95	1	• bioilled

SODIUM CHLORIDE

Not funded for use as a nasal drop. Not funded for nebuliser use except when used in conjunction with an antibiotic intended for nebuliser use.

Baxter	500 ml	PSO1.33	Inj 0.9%, bag - Up to 2000 ml available on a PSO
✓ Baxter	1.000 ml	1.36	

Only if prescribed on a prescription for renal dialysis, maternity or post-natal care in the home of the patient, or on a PSO for emergency use. (500 ml and 1,000 ml packs)

For Sodium chloride oral liquid formulation refer Standard Formulae, page 248 Inj 0.9%, 5 ml ampoule - Up to 5 inj available on a PSO4.00

✓ Biomed

 ✓ Fresenius Kabi ✓ Fresenius Kabi

✓ Fresenius Kabi

TOTAL PARENTERAL NUTRITION (TPN) Infusion......CBS

1 OP ✓ TPN

20

WATER

- 1) On a prescription or Practitioner's Supply Order only when on the same form as an injection listed in the Pharmaceutical Schedule requiring a solvent or diluent; or
- 2) On a bulk supply order: or
- 3) When used in the extemporaneous compounding of eye drops; or
- 4) When used for the dilution of sodium chloride soln 7% for cystic fibrosis patients only.

Inj 10 ml ampoule – Up to 5 inj available on a PSO7.19	50
Ini 20 ml ampoule – Up to 5 ini available on a PSO	20

✓ Pfizer

✓ Fresenius Kabi

✓ Multichem

(Multichem Inj 20 ml ampoule to be delisted 1 January 2023)

	Subsidy (Manufacturer's Pri	ice) Subsi	Fully idised	Brand or Generic Manufacturer
Oral Administration	<u> </u>			
CALCIUM POLYSTYRENE SULPHONATE Powder	169.85	300 g OP	√ 0	alcium Resonium
COMPOUND ELECTROLYTES Powder for oral soln — Up to 5 sach available on a PSO Electral to be Principal Supply on 1 December 2022	9.53	50	√ E	lectral
COMPOUND ELECTROLYTES WITH GLUCOSE [DEXTROSE] Soln with electrolytes (2 × 500 ml)		1,000 ml OP	√ P	edialyte - Bubblegum
PHOSPHORUS Tab eff 500 mg (16 mmol)	82.50	100	✓ P	hosphate Phebra
POTASSIUM CHLORIDE * Tab eff 548 mg (14 m eq) with chloride 285 mg (8 m eq)	5.26 (17.10)	60	C	hlorvescent
* Tab long-acting 600 mg (8 mmol)		200		pan-K
Cap 840 mg	8.52	100	_	odibic odibic
SODIUM POLYSTYRENE SULPHONATE				
Powder	84.65	454 a OP	✓ R	esonium-A

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Alpha-A	drenoceptor Blockers				
Alpha Ad	drenoceptor Blockers				
	g		500		Doxazosin Clinect
	9ENZAMINE HYDROCHLORIDE	20.94	500	•	Doxazosin Clinect
* Cap 10 r	ng	65.00 216.67	30 100		BNM S29 Dibenzyline S29
PRAZOSIN * Tab 1 m	g	5.53	100	•	Arrotex-Prazosin
* Tab 2 m	g	7.00	100	•	Arrotex-Prazosin
* Tab 5 m	g	11.70	100	•	Arrotex-Prazosin S29 S29
Agents A	Affecting the Renin-Angiotensin System	1			
ACE Inhi	ibitors				
	5 mg per mllliquid restricted to children under 12 years of age.	94.99 95	5 ml C)P ✓	Capoten
Subsidy endorse	. – Subsidy by endorsement by endorsement – Subsidised for patients who were d accordingly. Pharmacists may annotate the presci ng of cilazapril.				
	mg	2.09	90	1	Zapril
* Tab 2.5	mg	4.80	90	✓	Zapril
Tab 5 m	g	8.35	90	✓	Zapril
	g	1.82	100	1	Acetec
	ng		100		Acetec
	ng		100		Acetec
LISINOPRIL	· ·		. 50	,	
	g	11.07	90		Ethics Lisinopril Teva Lisinopril
* Tab 10 r	ng	11.67	90	/	Ethics Lisinopril Teva Lisinopril
* Tab 20 r	ng	14.69	90	•	Ethics Lisinopril Teva Lisinopril
PERINDOPE	RIL	4.50			_

30

30

30

✓ Coversyl

✓ Coversyl

✓ Coversyl

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
QUINAPRIL Tab 5 mg Tab 10 mg Tab 20 mg	5.18	90 90 90	✓	Arrow-Quinapril 5 Arrow-Quinapril 10 Arrow-Quinapril 20
ACE Inhibitors with Diuretics				
QUINAPRIL WITH HYDROCHLOROTHIAZIDE – Subsidy by e Subsidy by endorsement – Subsidised for patients who wel 2022 and the prescription is endorsed accordingly. Pharma exists a record of prior dispensing of quinapril with hydrochl Tab 10 mg with hydrochlorothiazide 12.5 mg	re taking quinapril with acists may annotate the lorothiazide4.10		scription a	
Angiotensin II Antagonists				
CANDESARTAN CILEXETIL * Tab 4 mg	2.28 3.31 5.26 1.56 1.84 2.25	90 90 90 90 84 84 84 84	\ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \ \	Candestar Candestar Candestar Candestar Candestar Losartan Actavis Losartan Actavis Losartan Actavis Losartan Actavis
Angiotensin II Antagonists with Diuretics				
LOSARTAN POTASSIUM WITH HYDROCHLOROTHIAZIDE * Tab 50 mg with hydrochlorothiazide 12.5 mg	4.00	30	•	Arrow-Losartan & Hydrochlorothiazide
Angiotensin II Antagonists with Neprilysin Inhi	ibitors			
SACUBITRIL WITH VALSARTAN – Special Authority see SA11 Tab 24.3 mg with valsartan 25.7 mg Tab 48.6 mg with valsartan 51.4 mg Tab 97.2 mg with valsartan 102.8 mg	190.00 190.00	rmad 56 56 56	· •	Entresto 24/26 Entresto 49/51 Entresto 97/103

⇒SA1905 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Patient has heart failure; and
- 2 Any of the following:
 - 2.1 Patient is in NYHA/WHO functional class II; or
 - 2.2 Patient is in NYHA/WHO functional class III; or
 - 2.3 Patient is in NYHA/WHO functional class IV; and
- 3 Fither
 - 3.1 Patient has a documented left ventricular ejection fraction (LVEF) of less than or equal to 35%; or

continued...

Subsidy (Manufacturer's Pric \$	ce) Subsi Per	Fully dised	Brand or Generic Manufacturer	
---------------------------------------	------------------	----------------	-------------------------------------	--

continued...

- 3.2 An ECHO is not reasonably practical, and in the opinion of the treating practitioner the patient would benefit from treatment; and
- 4 Patient is receiving concomitant optimal standard chronic heart failure treatments.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Antiarrhythmics

For lignocaine hydrochloride refer to NERVOUS SYSTEM, Anaesthetics, Lo	ocal, page 118	
AMIODARONE HYDROCHLORIDE		_
▲ Tab 100 mg	49 30	✓ Aratac
Aratac to be Principal Supply on 1 December 2022		
▲ Tab 200 mg	49 30	✓ Aratac
Aratac to be Principal Supply on 1 December 2022		
Inj 50 mg per ml, 3 ml ampoule — Up to 10 inj available on a PSO15.	22 10	✓ Max Health
Max Health to be Principal Supply on 1 December 2022	22 10	• Wax riealtii
ATROPINE SULPHATE		
* Inj 600 mcg per ml, 1 ml ampoule – Up to 5 inj available on a PSO	09 10	✓ Martindale
	09 10	<u> Martinuale</u>
DIGOXIN * Tab 62.5 mcg - Up to 30 tab available on a PSO	80 240	✓ Lanoxin PG
* Tab 250 mcg - Up to 30 tab available on a PSO		✓ Lanoxin
* Oral lig 50 mcg per ml		✓ Lanoxin
Total liq 50 flog por fill	00 001111	✓ Lanoxin Paediatric
		Elixir S29
		✓ Lanoxin S29 S29
DISOPYRAMIDE PHOSPHATE		- Lunoxiii O20
▲ Cap 100 mg23.	87 100	✓ Rythmodan
FLECAINIDE ACETATE	07 100	• Hydimodan
↑ Tab 50 mg19.	95 60	✓ Flecainide BNM
▲ Cap long-acting 100 mg		✓ Flecainide Bittin
Sup long doding foo mg	01 00	Controlled
		Release Teva
▲ Cap long-acting 200 mg61.	06 90	✓ Flecainide
		Controlled
		Release Teva
Inj 10 mg per ml, 15 ml ampoule100.	00 5	✓ Tambocor
MEXILETINE HYDROCHLORIDE		
▲ Cap 150 mg162.	00 100	✓ Teva S29
▲ Cap 250 mg	00 100	✓ Teva S29
PROPAFENONE HYDROCHLORIDE		
▲ Tab 150 mg40.	90 50	✓ Rytmonorm
		•
Antihypotensives		
MIDODRINE - Special Authority see SA1474 on the next page - Retail ph	armacy	
Tab 2.5 mg	,	✓ Gutron
Tab 5 mg79.		✓ Gutron

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

⇒SA1474 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where patient has disabling orthostatic hypotension not due to drugs.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Beta-Adrenoceptor Blockers

Beta Adrenoceptor Blockers

ATENOLOL	0 500	/ Modern Adam allal
* Tab 100 mg		 ✓ Mylan Atenolol ✓ Mylan Atenolol
* Tab 100 mg		✓ Atenolol AFT
The order lig 20 mg por 0 million	3 000 1111 01	S29 S29
38.2	0	✓ Essential
33.		Generics S29
49.8	5	✓ Atenolol AFT
Restricted to children under 12 years of age.	-	
BISOPROLOL FUMARATE		
* Tab 2.5 mg	4 90	✓ Bisoprolol Mylan
* Tab 5 mg	5 90	✓ Bisoprolol Mylan
* Tab 10 mg3.6	2 90	✓ Bisoprolol Mylan
CARVEDILOL		
* Tab 6.25 mg	4 60	✓ Carvedilol Sandoz
* Tab 12.5 mg2.3	0 60	✓ Carvedilol Sandoz
* Tab 25 mg2.9	5 60	✓ Carvedilol Sandoz
LABETALOL		
* Tab 100 mg14.5	0 100	✓ Trandate
* Tab 200 mg27.0	0 100	✓ <u>Trandate</u>
* Inj 5 mg per ml, 20 ml ampoule59.0	6 5	
(88.6	,	Trandate
* inj 5 mg per ml, 20 ml vial42.2		
(48.2	0)	Alvogen S29
METOPROLOL SUCCINATE		
* Tab long-acting 23.75 mg1.4		✓ Betaloc CR
* Tab long-acting 47.5 mg		✓ Betaloc CR
* Tab long-acting 95 mg		✓ Betaloc CR
* Tab long-acting 190 mg4.2	7 30	✓ Betaloc CR
METOPROLOL TARTRATE		4.
Tab 50 mg5.6		✓ IPCA-Metoprolol
Tab 100 mg		✓ IPCA-Metoprolol
* Tab long-acting 200 mg		✓ Slow-Lopresor✓ Metoprolol IV Mylan
* Inj 1 mg per ml, 5 ml vial	U 5	wietoprotot iv wytan
NADOLOL		
Tab 40 mg19.1		✓ Nadolol BNM S29
Tab 80 mg30.3	9 100	✓ Nadolol BNM S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
PROPRANOLOL				
Tab 10 mg	7.04	100	✓ [<u> Drofate</u>
Tab 40 mg	8.75	100	✓	PCA-Propranolol
* Cap long-acting 160 mg	18.17	100	✓ (Cardinol LA
* Oral lig 4 mg per ml - Special Authority see SA1327 be	low –			
Retail pharmacy	CBS	500 ml	✓ F	Roxane- Propranolol S29

⇒SA1327 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the treatment of a child under 12 years with an haemangioma causing functional impairment (not for cosmetic reasons only); or
- 2 For the treatment of a child under 12 years with cardiac arrthymias or congenital cardiac abnormalities.

SOTALOL

*	Tab 80 mg	.37.50	500	Mylan
*	Tab 160 mg	.14.00	100	Mylan

Calcium Channel Blockers

Dihydropyridine Calcium Channel Blockers

ΑN	ILODIPINE		
*	Tab 2.5 mg1.08	90	✓ Vasorex
*	Tab 5 mg	90	✓ Vasorex
*	Tab 10 mg1.19	90	✓ Vasorex
FE	LODIPINE		
*	Tab long-acting 2.5 mg1.45	30	✓ Plendil ER
*	Tab long-acting 5 mg4.07	90	✓ Felo 5 ER
*	Tab long-acting 10 mg4.32	90	✓ Felo 10 ER
NIF	EDIPINE		
*	Tab long-acting 10 mg	56	✓ Tensipine MR10 S29
*	Tab long-acting 20 mg9.12	50	✓ Mylan (12 hr release) S29
	17.72	100	✓ Nyefax Retard
*	Tab long-acting 30 mg4.78	14	✓ Mylan Italy (24 hr
			release) S29
	34.10	100	✓ Mylan (24 hr
			release) S29
*	Tab long-acting 60 mg52.81	100	✓ Mylan (24 hr
			release) S29

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
Other Calcium Channel Blockers				
DILTIAZEM HYDROCHLORIDE				
Cap extended-release 120 mg	44.40	100	1	Accord S29
* Cap long-acting 120 mg		500	✓	Apo-Diltiazem CD
* Cap long-acting 180 mg	7.00	30		Cardizem CD
* Cap long-acting 240 mg	9.30	30	✓	Cardizem CD
PERHEXILINE MALEATE				
* Tab 100 mg	62.90	100	1	Pexsig
/ERAPAMIL HYDROCHLORIDE				
* Tab 40 mg	7.01	100	1	Isoptin
* Tab 80 mg		100		Isoptin
* Tab long-acting 120 mg		100		Isoptin Retard \$29
* Tab long-acting 120 mg		100		Isoptin SR
* Tab long-acting 240 mg	15 12	30		Isoptin SR
* Inj 2.5 mg per ml, 2 ml ampoule – Up to 5 inj available on a		00	-	ioopiiii oii
PSOPSO		5	1	Isoptin
				Tooptiii
Centrally-Acting Agents				
, , ,				
CLONIDINE ** Poten 2.5 mg, 100 mgg, now down. Only on a proportion.	10.04	4	./	Mules
* Patch 2.5 mg, 100 mcg per day – Only on a prescription		4		Mylan Mylan
 Patch 5 mg, 200 mcg per day - Only on a prescription Patch 7.5 mg, 300 mcg per day - Only on a prescription 		4 4	_	Mylan Mylan
	10.93	4	•	<u>Mylan</u>
CLONIDINE HYDROCHLORIDE	0.75		,	O B
* Tab 25 mcg		112		Clonidine BNM
N. Tab 150	29.32	100		Clonidine Teva
* Tab 150 mcg		100 10		<u>Catapres</u>
* Inj 150 mcg per ml, 1 ml ampoule(Clonidine BNM Tab 25 mcg to be delisted 1 November 2022)	29.00	10	•	<u>Medsurge</u>
,				
METHYLDOPA	45.40	400	,	
* Tab 250 mg		100		Methyldopa Mylan
	52.85	500	•	Methyldopa Mylan
				S29 S29
Diuretics				
Didientes				
Loop Diuretics				
BUMETANIDE				
* Tab 1 mg	4.91	30	1	Burinex S29 S29
	16.36	100		Burinex
* Inj 500 mcg per ml, 4 ml vial		5		Burinex

	Subsidy		Fully Brand or
	(Manufacturer's F		sidised Generic
	\$	Per	✓ Manufacturer
FUROSEMIDE [FRUSEMIDE]	0.00	4 000	(IDOA E
Tab 40 mg - Up to 30 tab available on a PSO		1,000	✓ <u>IPCA-Frusemide</u> ✓ Urex Forte
* Tab 500 mg	25.00 89.48	50	✓ Urex Forte ✓ Furosemid-
	09.40		
			Ratiopharm S29
	169.96	100	✓ Furosemid-
			Ratiopharm \$29
			•
* Oral liq 10 mg per ml		30 ml OP	✓ Lasix
* Inj 10 mg per ml, 25 ml ampoule		6	✓ Lasix
Inj 10 mg per ml, 2 ml ampoule – Up to 5 inj availab	ole on a PSO2.40	5	✓ Furosemide-Baxter
Potassium Sparing Diuretics			
AMILORIDE HYDROCHLORIDE			
Oral liq 1 mg per ml	32.10	25 ml OP	✓ Biomed
EPLERENONE - Special Authority see SA1728 below -			
Tab 25 mg		30	✓ Inspra
Tab 50 mg	25.00	30	✓ Inspra
⇒SA1728 Special Authority for Subsidy			
Initial application from any relevant practitioner. Appro	vals valid without further	renewal unless	s notified for applications meetin
the following criteria:			
Both:			
 Patient has heart failure with ejection fraction less Either: 	s than 40%; and		
2.1 Patient is intolerant to optimal dosing of sp	oironolactone: or		
2.2 Patient has experienced a clinically signifi		on optimal dos	sing of spironolactone.
METOLAZONE			gp
Tab 5 mg	CBS	1	✓ Metolazone S29
Tab 5 Hig		=	
		50	✓ Zaroxolyn S29
SPIRONOLACTONE	0.00	400	40 ! !!
* Tab 25 mg		100	✓ <u>Spiractin</u>
* Tab 100 mg		100 25 ml OP	✓ <u>Spiractin</u> ✓ Biomed
Oral liq 5 mg per ml	30.60	25 MI OP	♥ Blomed
Potassium Sparing Combination Diuretic	s		
AMILORIDE HYDROCHLORIDE WITH FUROSEMIDE			
W. Tala Financials formanancials 40 man	0.00	00	/ F

 28

50

✓ Frumil

✓ Moduretic

	Subsidy		Fully	
	(Manufacturer's Price \$	e) Per	Subsidised <	Generic Manufacturer
Thiazide and Related Diuretics				
BENDROFLUMETHIAZIDE [BENDROFLUAZIDE]				
₭ Tab 2.5 mg — Up to 150 tab available on a PSO	20.00	500	✓	Arrow- Bendrofluazide
May be supplied on a PSO for reasons other than emerg	gency.			
≰ Tab 5 mg	34.55	500	✓	Arrow- Bendrofluazide
CHLOROTHIAZIDE				
Oral liq 50 mg per ml	27.82	25 ml C	P 🗸	Biomed
CHLORTALIDONE [CHLORTHALIDONE]				
Tab 25 mg		30		Igroton \$29
ND A DANIED	6.50	50	•	Hygroton
NDAPAMIDE 烙 Tab 2.5 mg	10.45	90	J	Dapa-Tabs
r 1au 2.0 mg	11.61	100		Mylan
				Indapamide \$29
Lipid-Modifying Agents Fibrates				
BEZAFIBRATE				
⊁ Tab 200 mg	19.46	90	1	Bezalip
₭ Tab long-acting 400 mg	21.21	30	✓	Bezalip Retard
Other Lipid-Modifying Agents				
CIPIMOX				
★ Cap 250 mg	21.56	30		Olbetam
			•	Olbetam S29 S29
Resins				
COLESTIPOL HYDROCHLORIDE				
Grans for oral liq 5 g	32.89	30	✓	Colestid
HMG CoA Reductase Inhibitors (Statins)				
TORVASTATIN				
k Tab 10 mg		500		Lorstat
lk Tab 20 mgk Tab 40 mg		500 500		Lorstat Lorstat
F Tab 40 mg		500		Lorstat
		500	-	
PRAVASTATIN				
PRAVASTATIN ★ Tab 20 mg	2.11	28	/	Pravastatin Mylan

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price)			Brand or Generic	
	\$			Manufacturer	
ROSUVASTATIN - Special Authority see SA2093 below - Ret	ail pharmacy				_
* Tab 5 mg	1.70	30	✓ R	osuvastatin Viatris	
* Tab 10 mg	2.42	30	✓ R	osuvastatin Viatris	
* Tab 20 mg	3.92	30	✓ R	osuvastatin Viatris	
* Tab 40 mg		30	✓ R	osuvastatin Viatris	

⇒SA2093 Special Authority for Subsidy

Initial application — (cardiovascular disease risk) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient is considered to be at risk of cardiovascular disease; and
 - 1.2 Patient is Māori or any Pacific ethnicity: or
- 2 Both:
 - 2.1 Patient has a calculated risk of cardiovascular disease of at least 15% over 5 years; and
 - 2.2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (familial hypercholesterolemia) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has familial hypercholesterolemia (defined as a Dutch Lipid Criteria score greater than or equal to 6); and
- 2 LDL cholesterol has not reduced to less than 1.8 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (established cardiovascular disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has proven coronary artery disease (CAD); or
 - 1.2 Patient has proven peripheral artery disease (PAD); or
 - 1.3 Patient has experienced an ischaemic stroke; and
- 2 LDL cholesterol has not reduced to less than 1.4 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

Initial application — (recurrent major cardiovascular events) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has experienced a recurrent major cardiovascular event (defined as myocardial infarction, ischaemic stroke, coronary revascularisation, hospitalisation for unstable angina) in the last 2 years; and
- 2 LDL cholesterol has not reduced to less than 1.0 mmol/litre with treatment with the maximum tolerated dose of atorvastatin and/or simvastatin.

SIN	//VASTATIN			
*	Tab 10 mg	1.23	90	✓ Simvastatin Mylan
*	Tab 20 mg	2.03	90	✓ Simvastatin Mylan
*	Tab 40 mg	3.58	90	✓ Simvastatin Mylan
*	Tab 80 mg	7.12	90	✓ Simvastatin Mylan

Selective Cholesterol Absorption Inhibitors

EZE	TIMIBE – Special Authority see SA1045 on the next page – Retail pharmacy		
*	Tab 10 mg1.95	30	✓ Ezetimibe Sandoz

Subsidy		Fully	Brand or
(Manufacturer's	Price) Sub	sidised	Generic
\$	Per	✓	Manufacturer

⇒SA1045 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 years; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 Any of the following:
 - 3.1 The patient has rhabdomyolysis (defined as muscle aches and creatine kinase more than $10 \times \text{normal}$) when treated with one statin; or
 - 3.2 The patient is intolerant to both simvastatin and atorvastatin; or
 - 3.3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atoryastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

5.15	30	✓ Zimybe
6.15	30	✓ Zimybe
7.15	30	✓ Zimybe
8.15	30	✓ Zimybe
	6.15 7.15	6.15 30 7.15 30

⇒SA1046 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 Patient has a calculated absolute risk of cardiovascular disease of at least 15% over 5 year; and
- 2 Patient's LDL cholesterol is 2.0 mmol/litre or greater; and
- 3 The patient has not reduced their LDL cholesterol to less than 2.0 mmol/litre with the use of the maximal tolerated dose of atorvastatin.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Nitrates

GLYCERYL TRINITRATE

*	Oral pump spray, 400 mcg per dose — Up to 250 dose			
	available on a PSO	6.09	250 dose OP	✓ Nitrolingual Pump Spray
*	Patch 25 mg, 5 mg per day	15.73	30	✓ Nitroderm TTS
	Patch 50 mg, 10 mg per day		30	✓ Nitroderm TTS
IS	OSORBIDE MONONITRATE			
*	Tab 20 mg	19.55	100	✓ Ismo 20
*	Tab long-acting 40 mg	8.20	30	✓ Ismo 40 Retard
	Tab long-acting 60 mg		90	✓ Duride

Sympathomimetics

ADRENALINE

Inj 1 in 1,000, 1 ml ampoule - Up to 5 inj available on a PSO4.98	5	✓ Aspen Adrenaline
10.76		✓ DBL Adrenaline
Inj 1 in 10,000, 10 ml ampoule - Up to 5 inj available on a PSO27.00	5	✓ Hospira
49.00	10	✓ Acnon Adronalina

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per		Manufacturer
Vasodilators				
HYDRALAZINE HYDROCHLORIDE				
* Tab 25 mg - Special Authority see SA1321 below - Retail				
pharmacy	CBS	1	1	Hydralazine
		56		Onelink \$29
		84		AMDIPHARM \$29
		100	_	Onelink S29
* Inj 20 mg ampoule	25.90	5	•	Apresoline
⇒SA1321 Special Authority for Subsidy				
nitial application from any relevant practitioner. Approvals valine following criteria: Either:	d without further rene	wal ui	nless notif	ied for applications meeting
 For the treatment of refractory hypertension; or For the treatment of heart failure in combination with a nit inhibitors and/or angiotensin receptor blockers. 	rate, in patients who a	re int	olerant or	have not responded to AC
MINOXIDIL				
▲ Tab 10 mg	70.00	100	✓	Loniten
NICORANDIL				
▲ Tab 10 mg	25.57	60	1	Ikorel
▲ Tab 20 mg	32.28	60	✓	Ikorel
PAPAVERINE HYDROCHLORIDE				
₭ Inj 12 mg per ml, 10 ml ampoule	257.12	5	✓	Hospira
PENTOXIFYLLINE [OXPENTIFYLLINE]				
Tab 400 mg	42.26	50	✓	Trental 400
Endothelin Receptor Antagonists				
MBRISENTAN - Special Authority see SA1702 below - Retail	pharmacy			
Tab 5 mg		30		Ambrisentan Mylan
Tab 10 mg	1,550.00	30		Ambrisentan Mylan
Ambricanton Mulan Tab 10 mg to be delicted 1 March 2022			•	Mylan
Ambrisentan Mylan Tab 10 mg to be delisted 1 March 2023)				
SA1702 Special Authority for Subsidy	on Donal			
Special Authority approved by the Pulmonary Arterial Hypertensi- lotes: Application details may be obtained from Pharmac's web		c aov	t nz/SAFo	rme or
The Coordinator, PAH Panel	ono <u>oonoudio.prianna</u>	o.gov	LIIZ/OAL O	iiio oi.
Pharmac, PO Box 10-254, WELLINGTON				
el: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac	c.govt.nz			
BOSENTAN - Special Authority see SA1991 on the next page -	Retail pharmacy			
Tab 62.5 mg	119.85	60	1	Bosentan Dr
				Reddy's
T-1-405	440.05	~~		December Do

Tab 125 mg119.85

60

✓ Bosentan Dr Reddy's

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA1991 Special Authority for Subsidy

Initial application only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 PAH is in Group 1, 4 or 5 of the WHO (Venice) clinical classifications; and
- 3 PAH is at NYHA/WHO functional class II, III, or IV; and
- 4 Any of the following:
 - 4.1 Both:
 - 4.1.1 Bosentan is to be used as PAH monotherapy; and
 - 4.1.2 Either:
 - 4.1.2.1 Patient is intolerant or contraindicated to sildenafil; or
 - 4.1.2.2 Patient is a child with idiopathic PAH or PAH secondary to congenital heart disease; or
 - 4.2 Both:
 - 4.2.1 Bosentan is to be used as PAH dual therapy; and
 - 4.2.2 Either:
 - 4.2.2.1 Patient has tried a PAH monotherapy for at least three months and failed to respond; or
 - 4.2.2.2 Patient deteriorated while on a PAH monotherapy, or
 - 4.3 Both:
 - 4.3.1 Bosentan is to be used as PAH triple therapy; and
 - 4.3.2 Any of the following:
 - 4.3.2.1 Patient is on the lung transplant list; or
 - 4.3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 4.3.2.3 Patient is deteriorating rapidly to NYHAWHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised; or
 - 4.3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Renewal only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory physician or cardiologist. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Bosentan is to be used as PAH monotherapy; and
 - 1.2 Patient is stable or has improved while on bosentan: or
- 2 Both:
 - 2.1 Bosentan is to be used as PAH dual therapy; and
 - 2.2 Patient has tried a PAH monotherapy for at least three months and either failed to respond or later deteriorated; or
- 3 Both:
 - 3.1 Bosentan is to be used as PAH triple therapy; and
 - 3.2 Any of the following:
 - 3.2.1 Patient is on the lung transplant list; or
 - 3.2.2 Patient is presenting acutely with idiopathic pulmonary arterial hypertension (IPAH) in New York Heart Association/World Health Organization (NYHA/WHO) Functional Class IV; or
 - 3.2.3 Patient is deteriorating rapidly to NYHA/WHO Functional Class IV who may be lung transplant recipients in the future, if their disease is stabilised: or
 - 3.2.4 Patient has PAH associated with the scleroderma spectrum of diseases (APAHSSD) who have no major morbidities and are deteriorating despite combination therapy.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

Phosphodiesterase Type 5 Inhibitors

SILDENAFIL – Special Authority see SA1992 below – Retail pharmacy		
Tab 25 mg	5 4	✓ Vedafil
Tab 50 mg) 4	✓ Vedafil
Tab 100 mg10.20) 12	✓ Vedafil

⇒SA1992 Special Authority for Subsidy

Initial application — (Raynaud's Phenomenon*) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has Raynaud's Phenomenon*; and
- 2 Patient has severe digital ischaemia (defined as severe pain requiring hospital admission or with a high likelihood of digital ulceration; digital ulcers; or gangrene); and
- 3 Patient is following lifestyle management (avoidance of cold exposure, sufficient protection, smoking cessation support, avoidance of sympathomimetic drugs); and
- 4 Patient is being treated with calcium channel blockers and nitrates (or these are contraindicated/not tolerated).

Initial application — (Pulmonary arterial hypertension*) only from a respiratory specialist, cardiologist or medical practitioner on the recommendation of a respiratory specialist or cardiologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has pulmonary arterial hypertension (PAH)*; and
- 2 Any of the following:
 - 2.1 PAH is in Group 1 of the WHO (Venice) clinical classifications; or
 - 2.2 PAH is in Group 4 of the WHO (Venice) clinical classifications; or
 - 2.3 PAH is in Group 5 of the WHO (Venice) clinical classifications; and
- 3 Any of the following:
 - 3.1 PAH is in NYHA/WHO functional class II; or
 - 3.2 PAH is in NYHA/WHO functional class III; or
 - 3.3 PAH is in NYHA/WHO functional class IV; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient has a pulmonary capillary wedge pressure (PCWP) less than or equal to 15 mmHg; and
 - 4.1.2 Fither:
 - 4.1.2.1 Patient has a mean pulmonary artery pressure (PAPm) > 25 mmHg; or
 - 4.1.2.2 Patient is peri Fontan repair; and
 - 4.1.3 Patient has a pulmonary vascular resistance (PVR) of at least 3 Wood Units or at least 240 International Units (dyn s cm-5); or
 - 4.2 Testing for PCWP, PAPm, or PVR cannot be performed due to the patient's young age.

Note: Indications marked with * are unapproved indications.

Initial application — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has a documented history of traumatic or non-traumatic spinal cord injury; and
- 2 Patient has erectile dysfunction secondary to spinal cord injury requiring pharmacological treatment.

Renewal — (erectile dysfunction due to spinal cord injury) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Subsidy		Fully	Brand or	Τ
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Prostacyclin Analogues

EPOPROSTENOL - Special Authority see SA1696 below - Retail pharm	nacy		
Inj 500 mcg vial3	6.61	1	✓ Veletri
Inj 1.5 mg vial7	3.21	1	✓ Veletri

⇒SA1696 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

✓ Vebulis✓ Ventavis

30

(Ventavis Nebuliser soln 10 mcg per ml, 2 ml to be delisted 1 March 2023)

⇒SA1705 Special Authority for Subsidy

Special Authority approved by the Pulmonary Arterial Hypertension Panel

Notes: Application details may be obtained from Pharmac's website schedule.pharmac.govt.nz/SAForms or:

The Coordinator, PAH Panel

Pharmac, PO Box 10-254, WELLINGTON

Tel: (04) 916 7561, Fax: (04) 974 4858, Email: PAH@pharmac.govt.nz



Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Antiacne Preparations

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

ADAPAI FNF

- a) Maximum of 30 g per prescription

b) Only on a prescription			
Crm 0.1%	22.89	30 g OP	Differin
Gel 0.1%	22.89	30 g OP	Differin
ISOTRETINOIN - Special Authority see SA2023 below - Reta	il pharmacy		
Cap 5 mg	11.26	60	Oratane
Cap 10 mg	18.75	120	✓ Oratane
Cap 20 mg	26.73	120	✓ Oratane

⇒SA2023 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around isotretinoin and is competent to prescribe isotretinoin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if isotretinoin is used during pregnancy and that they must not become pregnant during treatment and for a period of one month after the completion of treatment: or

(11.50)

2 Patient is not of child bearing potential.

TRFTINOIN

✓ ReTrieve Crm 0.5 mg per g - Maximum of 50 g per prescription15.57 50 q OP

Antibacterials Topical

For systemic antibacterials, refer to INFECTIONS, Antibacterials, page 90

HYDROGEN PEROXIDE

* Crm 1%	8.56		✓ Crystaderm✓ Crystaderm
MUPIROCIN Oint 2%	6.60	15 g OP	

- a) Only on a prescription
- b) Not in combination

Bactroban

	Subsidy (Manufacturer's Price \$	e) Sul Per	Fully osidised	Brand or Generic Manufacturer	
SODIUM FUSIDATE [FUSIDIC ACID]	<u> </u>				
Crm 2%	1.59	5 g OP	√ F	oban	
a) Maximum of 5 g per prescription		Ü	_		
b) Only on a prescription					
c) Not in combination		- 0-			
Oint 2%	1.59	5 g OP	✓ <u>F</u>	<u>oban</u>	
a) Maximum of 5 g per prescription b) Only on a prescription					
b) Only on a prescriptionc) Not in combination					
SULFADIAZINE SILVER					
Crm 1%	10.80	50 g OP	√ F	lamazine	
a) Up to 250 g available on a PSO		3 -			
b) Not in combination					
Antifuncale Tenical					
Antifungals Topical					
For systemic antifungals, refer to INFECTIONS, Antifungals, page 1	age 97				
AMOROLFINE					
a) Only on a prescription					
b) Not in combination Nail soln 5%	14.00	5 ml OP	./ 1	Avec Neil	
	14.93	5 IIII OP	<u> </u>	<u>/lycoNail</u>	
CLOTRIMAZOLE * Crm 1%	0.77	20 g OP	10	Clomazol	
a) Only on a prescription		20 g Oi	• •	,ioiiiazoi	
b) Not in combination					
* Soln 1%	4.36	20 ml OP			
	(7.55)		C	Canesten	
a) Only on a prescription					
b) Not in combination					
ECONAZOLE NITRATE	1.00	00 = OD			
Crm 1%	(7.48)	20 g OP	Р	Pevaryl	
a) Only on a prescription	(7.40)		į	Cvaryi	
b) Not in combination					
Foaming soln 1%, 10 ml sachets	9.89	3			
	(17.23)		P	Pevaryl	
a) Only on a prescription					
b) Not in combination					
MICONAZOLE NITRATE * Crm 2%	0.01	15 a OD	./ N	/lultichem	
a) Only on a prescription	0.01	15 g OP	<u> </u>	luitichem	
b) Not in combination					
* Lotn 2%	4.36	30 ml OP			
	(10.03)			Daktarin	
a) Only on a prescription					
b) Not in combination	4.00	30 ml OD			
* Tinct 2%	4.36 (12.10)	30 ml OP	г	Daktarin	
a) Only on a prescription	(12.10)		L	-unturn	
b) Not in combination					

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	(Manufacturer's Pric \$	e) Subs Per	idised •	Generic Manufacturer	
Antipruritic Preparations					
CALAMINE a) Only on a prescription b) Not in combination Crm, aqueous, BP	1.08	100 g	√ <u>c</u>	alamine-AFT	
CROTAMITON a) Only on a prescription b) Not in combination					
Crm 10% MENTHOL – Only in combination	3.29	20 g OP	✓ <u>lt</u>	ch-Soothe	
 Only in combination with a dermatological base or prop With or without other dermatological galenicals. 	rietary Topical Cort	icosteriod –	Plain		
Crystals	6.92	25 g	✓ M	idWest	

Subsidy

29.60

100 g

Fully

Brand or

MidWest

Corticosteroids Topical

Carticostaraide - Plain

For systemic corticosteroids, refer to CORTICOSTEROIDS AND RELATED AGENTS, page 80

Corticosteroids - Plain				
BETAMETHASONE DIPROPIC	DNATE			
Crm 0.05%		2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ Diprosone
Oint 0.05%		2.96	15 g OP	✓ Diprosone
		36.00	50 g OP	✓ <u>Diprosone</u>
Oint 0.05% in propylene gly	ycol base	4.33	30 g OP	Diprosone OV
BETAMETHASONE VALERAT	E			
* Crm 0.1%		4.53	50 g OP	✓ Beta Cream
* Oint 0.1%		5.84	50 g OP	✓ Beta Ointment
* Lotn 0.1%		25.00	50 ml OP	✓ Betnovate
CLOBETASOL PROPIONATE				
* Crm 0.05%		2.40	30 g OP	✓ Dermol
* Oint 0.05%		2.33	30 g OP	✓ Dermol
CLOBETASONE BUTYRATE				
Crm 0.05%		5.38	30 g OP	
		(10.00)	-	Eumovate
HYDROCORTISONE				
* Crm 1% – Only on a preso	pription	3.70	100 g OP	✓ Hydrocortisone (PSM)
		17.15	500 g	✓ Hydrocortisone (PSM)
* Powder - Only in combina	tion	49.95	25 g	✓ ABM
Up to 5% in a dermato galenicals	logical base (not proprietary Topical C	Corticosterio	d – Plain) with o	or without other dermatological
HYDROCORTISONE AND PAR	RAFFIN LIQUID AND LANOLIN			
Lotn 1% with paraffin liquid	15.9% and lanolin 0.6% - Only on			
		10.57	250 ml	✓ <u>DP Lotn HC</u>

	Subsidy		Fully	
	(Manufacturer's Pri \$	ice) Subs Per	idised •	I Generic Manufacturer
HYDROCORTISONE BUTYRATE	Ψ	101		manuacuro
Lipocream 0.1%	4 85	100 g OP	1	Locoid Lipocream
Oint 0.1%		100 g OP		Locoid
Milky emul 0.1%		100 ml OP		Locoid Crelo
METHYLPREDNISOLONE ACEPONATE				
Crm 0.1%	4.46	15 g OP	1	Advantan
Oint 0.1%	4.46	15 g OP	✓	Advantan
MOMETASONE FUROATE				
Crm 0.1%	1.95	15 g OP	1	Elocon Alcohol Free
	3.10	50 g OP		Elocon Alcohol Free
Oint 0.1%		15 g OP		Elocon
Lotn 0.1%	2.90	50 g OP 30 ml OP		Elocon
	4.50	JU IIII UP	٧	Elocon
TRIAMCINOLONE ACETONIDE	6.00	100 ~ OD	.1	Ariotocort
Crm 0.02% Oint 0.02%		100 g OP 100 g OP		Aristocort Aristocort
OHR 0.02 /0	0.00	100 g OF	•	Andiocont
Corticosteroids - Combination				
BETAMETHASONE VALERATE WITH SODIUM FUSIDATE [FUS	SIDIC ACID]			
Crm 0.1% with sodium fusidate (fusidic acid) 2%	3.49	15 g OP		
	(10.45)			Fucicort
a) Maximum of 15 g per prescriptionb) Only on a prescription				
HYDROCORTISONE WITH MICONAZOLE - Only on a prescript				
* Crm 1% with miconazole nitrate 2%	1.89	15 g OP	1	Micreme H
HYDROCORTISONE WITH NATAMYCIN AND NEOMYCIN - O	, , ,			
Crm 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		Pimafucort
Oint 1% with natamycin 1% and neomycin sulphate 0.5%		15 g OP		Pimafucort
(Pimafucort Crm 1% with natamycin 1% and neomycin sulphate C		'	<i>')</i>	
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCII		IN		
Crm 1 mg with nystatin 100,000 u, neomycin sulphate 2.5 mg and gramicidin 250 mcg per g - Only on a prescription	•	15 g OP		
and gramiolain 200 mag per g Only on a prescription	(9.28)	10 9 01		Viaderm KC
Barrier Creams and Emollients	(/			
Darrier Greatils and Emoliterits				
Barrier Creams				
DIMETHICONE				
* Crm 5% pump bottle	4.30	500 ml OP	✓	healthE
				Dimethicone 5%
healthE Dimethicone 5% to be Principal Supply on 1 Dec			_	
* Crm 10% pump bottle	4.52	500 ml OP	/	healthE
TIME AND GASTED ON				Dimethicone 10%
ZINC AND CASTOR OIL	4.65	E00 ~	./	Daushau
* Oint	4.05	500 g	•	Boucher

	Subsidy		Fully Brand or
	(Manufacturer's \$	Price) Subsi Per	idised Generic Manufacturer
	3	rei	- Ivianulaciulei
Emollients			
AQUEOUS CREAM			
* Crm	1.73	500 g	✓ <u>GEM Aqueous</u> <u>Cream</u>
CETOMACROGOL	4.00	500	
* Crm BP	1.99	500 g	✓ Cetomacrogol-AFT
CETOMACROGOL WITH GLYCEROL			4-
Crm 90% with glycerol 10%	2.35	500 ml OP	✓ Boucher ✓ Evara ✓ Pharmacy Health Sorbolene with Glycerin
	3.10	1,000 ml OP	✓ Boucher ✓ Evara
(Boucher Crm 90% with glycerol 10% to be delisted 1 March 20 (Boucher Crm 90% with glycerol 10% to be delisted 1 March 20 EMULSIFYING OINTMENT			
* Oint BP	3.40	500 g	✓ Emulsifying Ointment ADE
OIL IN WATER EMULSION			
* Crm	2.04	500 g	✓ Fatty Cream AFT
PARAFFIN Oint liquid paraffin 50% with white soft paraffin 50% UREA	5.35	500 ml OP	✓ healthE
* Crm 10%	1.37	100 g OP	✓ healthE Urea Cream
WOOL FAT WITH MINERAL OIL - Only on a prescription		3 -	
* Lotn hydrous 3% with mineral oil	5.60	1,000 ml	
•	(11.95)		DP Lotion
	1.40	250 ml OP	
	(4.53)	4 000	DP Lotion
	5.60	1,000 ml	Alaba Kari Latian
	(20.53) (23.91)		Alpha-Keri Lotion BK Lotion
	1.40	250 ml OP	DIX LOUDIT
	(7.73)	200 1111 01	BK Lotion
Other Dermatological Bases			
PARAFFIN			
White soft - Only in combination	4.99 19.99	450 g 2,500 g	✓ healthE✓ healthE
Only in combination with a dermatological galenical or	as a diluent for a		cal Corticosteroid - Plain.

✓ Stromectol

	Subsidy		Fully	Brand or
	(Manufacturer's Price	e) Subs Per	idised	Generic Manufacturer
	Ψ	rei		Ivianulacturei
Minor Skin Infections				
POVIDONE IODINE				
Oint 10%	7.40	65 g OP	✓ <u>E</u>	Betadine
 a) Maximum of 130 g per prescription 				
b) Only on a prescription				
Antiseptic Solution 10%	4.15	100 ml	✓ <u>F</u>	<u>Riodine</u>
Antiseptic soln 10%	3.83	15 ml	•	Riodine
	5.40	500 ml	✓ F	Riodine
Skin preparation, povidone iodine 10% with 30% alcohol		100 ml		
<u> </u>	(3.48)		E	Betadine Skin Prep
Skin preparation, povidone iodine 10% with 70% alcohol		100 ml	_	
	(7.78)		ŀ	Pfizer
Parasiticidal Preparations				
DIMETHICONE				
* Lotn 4%	4.25 2	200 ml OP	√ ł	healthE Dimethicone 4% Lotion
healthE Dimethicone 4% Lotion to be Principal Supply of	n 1 December 2022	2		
IVERMECTIN - Special Authority see SA1225 below - Retail ph	armacy			

- 1) PSO for institutional use only. Must be endorsed with the name of the institution for which the PSO is required and a valid Special Authority for patient of that institution.
- 2) Ivermectin available on BSO provided the BSO includes a valid Special Authority for a patient of the institution.
- For the purposes of subsidy of ivermectin, institution means age related residential care facilities, disability care facilities or prisons.

⇒SA1225 Special Authority for Subsidy

Initial application — (Scables) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
 Either:
- -.....
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and

Tab 3 mg - Up to 100 tab available on a PSO.......17.20

- 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
- 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:

continued...

DERMATOLOGICALS

	Subsidy		Fully	Brand or
1)	Manufacturer's Price)	Su	bsidised	Generic
	\$	Per	1	Manufacturer

continued...

- 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
- 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy; or
- 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Initial application — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

Renewal — (Scabies) from any relevant practitioner. Approvals valid for 1 month for applications meeting the following criteria: Both:

- 1 Applying clinician has discussed the diagnosis of scabies with a dermatologist, infectious disease physician or clinical microbiologist; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient is in the community; and
 - 2.1.2 Any of the following:
 - 2.1.2.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.1.2.2 The community patient is physically or mentally unable to comply with the application instructions of topical therapy; or
 - 2.1.2.3 The patient has previously tried and failed to clear infestation using topical therapy; or
 - 2.2 All of the following:
 - 2.2.1 The Patient is a resident in an institution; and
 - 2.2.2 All residents of the institution with scabies or at risk of carriage are to be treated for scabies concurrently; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 Patient has a severe scabies hyperinfestation (Crusted/ Norwegian scabies); or
 - 2.2.3.2 The patient is physically or mentally unable to comply with the application instructions of topical therapy: or
 - 2.2.3.3 Previous topical therapy has been tried and failed to clear the infestation.

Renewal — (Other parasitic infections) only from an infectious disease specialist, clinical microbiologist or dermatologist. Approvals valid for 1 month for applications meeting the following criteria:

Any of the following:

- 1 Filaricides: or
- 2 Cutaneous larva migrans (creeping eruption); or
- 3 Strongyloidiasis.

PERMETHRIN

Crm 5%	5.75	30 g OP	Lyderm
Lotn 5%	3.99	30 ml OP	✓ <u>A-Scabies</u>

Psoriasis and Eczema Preparations

ACITRETIN - Special Authority see SA2024 on the n	ext page – Retail pharmacy		
Cap 10 mg	17.86	60	✓ Novatretin
Cap 25 mg	41.36	60	✓ Novatretin

DERMATOLOGICAL

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA2024 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: All of the following:

- 1 Applicant is a vocationally registered dermatologist, vocationally registered general practitioner, or nurse practitioner working in a relevant scope of practice; and
- 2 Applicant has an up to date knowledge of the safety issues around acitretin and is competent to prescribe acitretin; and
- 3 Either:
 - 3.1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment: or
 - 3.2 Patient is not of child bearing potential.

Renewal from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Fither:

- 1 Patient is of child bearing potential and the possibility of pregnancy has been excluded prior to commencement of treatment and patient has been counselled and understands the risk of teratogenicity if acitretin is used during pregnancy and that they must not become pregnant during treatment and for a period of three years after the completion of treatment:
- 2 Patient is not of child bearing potential.

BETAMETHASONE DIPROPIONATE WITH CALCIPOTRIOL			
Foam spray 500 mcg with calcipotriol 50 mcg per g	59.95	60 g OP	Enstilar
Gel 500 mcg with calcipotriol 50 mcg per g	39.35	60 g OP	Daivobet
Oint 500 mcg with calcipotriol 50 mcg per g	15.90	30 g OP	✓ Daivobet
CALCIPOTRIOL			
Oint 50 mcg per g	40.00	120 g OP	Daivonex
COAL TAR			
Soln BP - Only in combination	36.25	200 ml	✓ Midwest
1) Up to 10% only in combination with a dermatological	base or propr	ietary Topical C	Corticosteriod – Pl

- Plain
- 2) With or without other dermatological galenicals.

COAL TAR WITH ALLANTOIN, MENTHOL, PHENOL AND SULPHUR Soln 5% with sulphur 0.5%, menthol 0.75%, phenol 0.5% and

allantoin crm 2.5%	6.59	75 g OP	
	(8.00)	Ü	Egopsoryl TA
	3.43	30 g OP	01 7
	(4.35)	3 3	Egopsoryl TA
COAL TAR WITH SALICYLIC ACID AND SULPHUR			
Soln 12% with salicylic acid 2% and sulphur 4% oint	4.97	25 g OP	✓ Coco-Scalp
	7.95	40 g OP	✓ Coco-Scalp

PIMECROLIMUS - Special Authority see SA1970 on the next page - Retail pharmacy

- a) Maximum of 15 g per prescription
- b) Note: a maximum of 15 g per prescription and no more than one prescription per 12 weeks

b) reservation of region presemption and resimilar	p. 000p	po=oc	,,,,,,
Cream 1%	28.50	15 g OP	Elidel



Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

⇒SA1970 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician, ophthalmologist or any relevant practitioner on the recommendation of a dermatologist, paediatrician or ophthalmologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has atopic dermatitis on the eyelid; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy, documented allergy to topical corticosteroids, cataracts, glaucoma, or raised intraocular pressure.

PINE TAR WITH TROLAMINE LAURILSULFATE AND FLUORESCEIN - Only on a prescription

* Soln 2.3% with trolamine laurilsulfate and fluorescein sodium...........4.44

500 ml

Pinetarsol

SALICYLIC ACID

Powder - Only in combination......18.88

250 g

100 a

✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain or collodion flexible
- 2) With or without other dermatological galenicals.

SULPHUR

✓ Midwest

- 1) Only in combination with a dermatological base or proprietary Topical Corticosteroid Plain
- 2) With or without other dermatological galenicals.

TACROLIMUS

Oint 0.1% - Special Authority see SA2074 below - Retail pharmacy.......33.00

30 q OP

✓ Zematop

- a) Maximum of 30 g per prescription
- b) Note: a maximum of 30 g per prescription and no more than one prescription per 12 weeks.

⇒SA2074 Special Authority for Subsidy

Initial application only from a dermatologist, paediatrician or any relevant practitioner on the recommendation of a dermatologist, paediatrician. . Approvals valid without further renewal unless notified for applications meeting the following criteria: Both:

- 1 Patient has atopic dermatitis on the face; and
- 2 Patient has at least one of the following contraindications to topical corticosteroids: periorificial dermatitis, rosacea, documented epidermal atrophy or documented allergy to topical corticosteroids.

Scalp Preparations

ETAMETHASONE VALERATE • Scalp app 0.1%	Beta Scalp
LOBETASOL PROPIONATE	Deta Scalp
	Dermol
YDROCORTISONE BUTYRATE Scalp lotn 0.1%	Locoid
ETOCONAZOLE	2000.0
Shampoo 2%	Sebizole Sebizole

- a) Maximum of 100 ml per prescription
- b) Only on a prescription

DERMATOLOGICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Sunscreens

SUNSCREENS, PROPRIETARY - Subsidy by endorsement

Only if prescribed for a patient with severe photosensitivity secondary to a defined clinical condition and the prescription is endorsed accordingly.

Wart Preparations

For salicylic acid preparations refer to PSORIASIS AND ECZEMA PREPARATIONS, page 68

IMIQUIMOD

PODOPHYLLOTOXIN

a) Maximum of 3.5 ml per prescription

b) Only on a prescription

Other Skin Preparations

Antineoplastics

FLUOROURACIL SODIUM

GENITO-URINARY SYSTEM

Subsidy		Fully	Brand or
(Manufacturer's Price)	,	Subsidised	Generic
\$	Per	1	Manufacturer

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✓ Manufacturer

Contraceptives - Non-hormonal

Condoms

CONDOMS				_
	vailable on a PSO		144	✓ Moments
€ 53 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa				_
53 mm, 0.05 mm thickness	S		10	✓ Moments
		11.42	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, chocolate, brown.			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
53 mm, strawberry, red			10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm			10	✓ Moments
		11.64	144	✓ Moments
 a) Maximum of 60 de 				
b) Up to 60 dev availa	able on a PSO			
56 mm, 0.05 mm thickness	S		12	✓ Gold Knight
		15.57	144	Gold Knight
 a) Up to 60 dev availa 				
b) Maximum of 60 de				
	s (bulk pack)	14.61	144	Gold Knight
 a) Maximum of 60 de 				
b) Up to 60 dev available				
56 mm, 0.08 mm thickness	S	0.97	10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de				
56 mm, 0.08 mm thickness	s, red	0.97	10	✓ Moments
		11.64	144	✓ Moments
 a) Up to 60 dev available 				
b) Maximum of 60 de	v per prescription			
56 mm, chocolate		1.30	12	Gold Knight
		15.57	144	Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de	v per prescription			
56 mm, strawberry		1.30	12	Gold Knight
•		15.57	144	✓ Gold Knight
a) Up to 60 dev availa	able on a PSO			
b) Maximum of 60 de				
		1.42	12	Gold Knight XL
		14.87	144	✓ Shield XL
		17.02		Gold Knight XL

GENITO-URINARY SYSTEM

		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
*	60 mm (bulk pack)	14.87	144	1	Gold Knight XL
	a) Maximum of 60 dev per prescription				
	b) Up to 60 dev available on a PSO				

Contraceptive Devices

INTRA-UTERINE DEVICE

- a) Up to 40 dev available on a PSO
- b) Only on a PSO

*	IÚD 29.1 mm length × 23.2 mm width	1	✓ 7 MED NSHA Silver/ Copper Short
			✓ Choice 380 7med Nsha Silver/ copper Short
			✓ Choice TT380 Short
*	IUD 33.6 mm length × 29.9 mm width	1	✓ Choice TT380 Standard
*	IUD 35.5 mm length × 19.6 mm width	1	✓ Choice Load 375

Contraceptives - Hormonal

Combined Oral Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Patient is on a Social Welfare benefit; or
 - 1.2 Patient has an income no greater than the benefit: and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit; or
 - 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

ETHINYLOESTRADIOL WITH DESOGESTREL

* Tab 20 mcg with desogestrel 150 mcg and 7 inert tab - Up to 84 tab available on a PSO 10.00 84 ✓ Mercilon 28

	Subsidy		Fully	Brand or
	(Manufacturer's Price)		Subsidised	I Generic
	\$	Per	•	Manufacturer
ETHINYLOESTRADIOL WITH LEVONORGESTREL				
* Tab 20 mcg with levonorgestrel 100 mcg and 7 inert tablets -	_			
Up to 112 tab available on a PSO	2.18	84	✓	Microgynon 20 ED
·	6.45	112		Femme-Tab ED
* Tab 30 mcg with levonorgestrel 150 mcg	6.62	63		
	(16.50)			Microgynon 30
a) Higher subsidy of \$15.00 per 63 tab with Special Auth	nority see SA0500 on	the	orevious p	age
b) Up to 63 tab available on a PSO	•			
* Tab 30 mcg with levonorgestrel 150 mcg and 7 inert tablets	_			
Up to 112 tab available on a PSO	1.77	84	1	Levlen ED
	6.45	112	✓	Femme-Tab ED
ETHINYLOESTRADIOL WITH NORETHISTERONE				
Tab 35 mcg with norethisterone 1 mg and 7 inert tab - Up to	1			
84 tab available on a PSO		84	/	Brevinor 1/28
Tab 35 mcg with norethisterone 500 mcg and 7 inert tab - U		•		
to 84 tab available on a PSO		84	/	Norimin
to on tab available on a 1 oo	29.32	112		Norimin
	23.32	112	•	Norman

Progestogen-only Contraceptives

⇒SA0500 Special Authority for Alternate Subsidy

Initial application from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Fither:
 - 1.1 Patient is on a Social Welfare benefit: or
 - 1.2 Patient has an income no greater than the benefit; and
- 2 Has tried at least one of the fully funded options and has been unable to tolerate it.

Renewal from any medical practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient is on a Social Welfare benefit: or
- 2 Patient has an income no greater than the benefit.

Notes: The approval numbers of Special Authorities approved after 1 November 1999 are interchangeable between Mercilon and Marvelon.

The additional subsidy will fund Mercilon and Marvelon up to the manufacturer's price for each of these products as identified on the Schedule at 1 November 1999.

Special Authorities approved before 1 November 1999 remain valid until the expiry date and can be renewed providing that women are still either:

- on a Social Welfare benefit; or
- have an income no greater than the benefit.

The approval numbers of Special Authorities approved before 1 November 1999 are interchangeable for products within the combined oral contraceptives and progestogen-only contraceptives groups, except Loette and Microgynon 20 ED

	Tab 30 mcg – Up to 84 tab available on a PSO		84 112	✓ Microlut✓ Microlut
*	Subdermal implant (2 × 75 mg rods) — Up to 3 pack available on a PSO10	06.92	1	✓ <u>Jadelle</u>
ME	DROXYPROGESTERONE ACETATE Inj 150 mg per ml, 1 ml syringe – Up to 5 inj available on a PSO	.7.98	1	✓ Depo-Provera

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
NORETHISTERONE Tab 350 mcg - Up to 84 tab available on a PSO	12.25	84	1	Noriday 28
Emergency Contraceptives				
LEVONORGESTREL * Tab 1.5 mg	4.95	1	•	Postinor-1
b) Up to 5 tab available on a PSO c) Note: Direct Provision by a pharmacist permitted ur	nder the provisions in F	Part I	of Section	ı A.

Antiandrogen Oral Contraceptives

Prescribers may code prescriptions "contraceptive" (code "O") when used as indicated for contraception. The period of supply and prescription charge will be as per other contraceptives, as follows:

- \$5.00 prescription charge (patient co-payment) will apply.
- prescription may be written for up to six months supply.

ACETIC ACID WITH HADDOAAOUINOLINE VAID DICINOLEIC VOID

Prescriptions coded in any other way are subject to the non contraceptive prescription charges, and the non-contraceptive period of supply. ie. Prescriptions may be written for up to three months supply.

CYPROTERONE ACETATE WITH ETHINYLOESTRADIOL

★ Tab 2 mg with ethinyloestradiol 35 mcg and 7 inert tabs – Up to 168 tab available on a PSO.......4.98 168 ✓ Ginet

Gynaecological Anti-infectives

Jelly with glacial acetic acid 0.94%, hydroxyguinoline sulphate		
0.025%, glycerol 5% and ricinoleic acid 0.75% with applicator8.43	100 g OP	
(24.15)		Aci-Jel
CLOTRIMAZOLE		
* Vaginal crm 1% with applicators2.50	35 g OP	✓ Clomazol
* Vaginal crm 2% with applicators3.00	20 g OP	✓ Clomazol
MICONAZOLE NITRATE		
* Vaginal crm 2% with applicator	40 g OP	✓ Micreme
NYSTATIN	J	
Vaginal crm 100,000 u per 5 q with applicator(s)4.00	75 g OP	✓ Nilstat
vaginar om 100,000 a por 0 g with applicator(o)	70 g Oi	· Milotat

Myometrial and Vaginal Hormone Preparations

ERGOMETRINE MALEATE			
Inj 500 mcg per ml, 1 ml ampoule – Up to 5 inj available on a			
PSOPSO	160.00	5	✓ DBL Ergometrine
DESTRIOL			
★ Crm 1 mg per g with applicator		15 g OP	✓ Ovestin
★ Pessaries 500 mcg	6.86	15	✓ Ovestin
DXYTOCIN - Up to 5 inj available on a PSO			
Inj 5 iu per ml, 1 ml ampoule	3.98	5	 Oxytocin BNM
Inj 10 iu per ml, 1 ml ampoule	4.98	5	 Oxytocin BNM
DXYTOCIN WITH ERGOMETRINE MALEATE - Up to 5 inj available	on a PSO		
Inj 5 iu with ergometrine maleate 500 mcg per ml, 1 ml ampoule Syntometrine to be Principal Supply on 1 December 2022		5	✓ Syntometrine

GENITO-URINARY SYSTEM

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Pregnancy Tests - hCG Urine

PREGNANCY TESTS - HCG URINE

- a) Up to 200 test available on a PSO
- b) Only on a PSO

16.00

✓ Smith BioMed Rapid Pregnancy Test

✓ David One Step Cassette Pregnancy Test

Urinary Agents

For urinary tract Infections refer to INFECTIONS, Antibacterials, page 107

5-Alpha Reductase Inhibitors

FINASTERIDE – Special Authority see SA0928 below – Retail pharmacy

★ Tab 5 mg4.81 100 ✓ Ricit

⇒SA0928 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 Either:
 - 2.1 The patient is intolerant of non-selective alpha blockers or these are contraindicated; or
 - 2.2 Symptoms are not adequately controlled with non-selective alpha blockers.

Alpha-1A Adrenoreceptor Blockers

⇒SA1032 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Patient has symptomatic benign prostatic hyperplasia; and
- 2 The patient is intolerant of non-selective alpha blockers or these are contraindicated.

Other Urinary Agents

OXYBUTYNIN

POTASSIUM CITRATE

GENITO-URINARY SYSTEM

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

⇒SA1083 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

1 The patient has recurrent calcium oxalate urolithiasis; and

2 The patient has had more than two renal calculi in the two years prior to the application.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

SODIUM CITRO-TARTRATE * Grans eff 4 g sachets	2 28	✓ <u>Ural</u>
SOLIFENACIN SUCCINATE		
Tab 5 mg2.0	5 30	✓ Solifenacin Mylan
Tab 10 mg3.7	2 30	✓ Solifenacin Mylan

Detection of Substances in Urine

\sim	D -	-1	_	-	 4.1		18	
u	R	ιн		- I	"	11)	ШΛ	11

٠.	THO TOLIDITE			
*	Compound diagnostic sticks	7.50	50 test OP	
	· •	(8.25)		Hemastix
TF	TRABROMOPHENOL			

Obstetric Preparations

Antiprogesterones

		MIFEPRISTONE	MIF
✓ Mifegyne	1	Tab 200 mg60.00	
✓ Mifegyne	3	180.00	

- a) Up to 15 tab available on a PSO
- b) Only on a PSO

100 test OP

✓ Albustix

Subsidy		Fully	Brand or	Τ
(Manufacturer's Price)	5	Subsidised	Generic	
\$	Per	✓	Manufacturer	

Calcium Homeostasis

0 4 1	_			
CA	()	1()	N	IΝ

CINACALCET – Special Authority see SA1618 below – Retail pharmacy
Tab 30 mg – Wastage claimable......42.06

 Tab 30 mg − Wastage claimable
 42.06
 28
 ✓ Cinacalet Devatis

 Tab 60 mg − Wastage claimable
 84.12
 28
 ✓ Cinacalet Devatis

⇒SA1618 Special Authority for Subsidy

Initial application only from a nephrologist or endocrinologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has been diagnosed with a parathyroid carcinoma (see Note); and
 - 1.2 The patient has persistent hypercalcaemia (serum calcium greater than or equal to 3 mmol/L) despite previous first-line treatments including sodium thiosulfate (where appropriate) and bisphosphonates; and
 - 1.3 The patient is symptomatic; or
- 2 All of the following:
 - 2.1 The patient has been diagnosed with calciphylaxis (calcific uraemic arteriolopathy); and
 - 2.2 The patient has symptomatic (e.g. painful skin ulcers) hypercalcaemia (serum calcium greater than or equal to 3 mmol/L); and
 - 2.3 The patient's condition has not responded to previous first-line treatments including bisphosphonates and sodium thiosulfate.

Renewal only from a nephrologist or endocrinologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient's serum calcium level has fallen to < 3mmol/L; and
- 2 The patient has experienced clinically significant symptom improvement.

Note: This does not include parathyroid adenomas unless these have become malignant.

ZOLEDRONIC ACID

⇒SA2109 Special Authority for Subsidy

Initial application — (bone metastases) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Patient has hypercalcaemia of malignancy; or
- 2 Both:
 - 2.1 Patient has bone metastases or involvement; and
 - 2.2 Patient has severe bone pain resistant to standard first-line treatments; or
- 3 Both:
 - 3.1 Patient has bone metastases or involvement; and
 - 3.2 Patient is at risk of skeletal-related events pathological fracture, spinal cord compression, radiation to bone or surgery to bone.

Initial application — (early breast cancer*) from any relevant practitioner. Approvals valid for 3 years for applications meeting

continued...

Viatris

	ubsidy cturer's Price) Subs	Fully	Brand or Generic
·	\$ Per	•	Manufacturer

continued...

the following criteria:

All of the following:

- 1 Treatment to be used as adjuvant therapy for early breast cancer; and
- 2 Patient has been amenorrhoeic for 12 months or greater, either naturally or induced, with endocrine levels consistent with a postmenopausal state; and
- 3 Treatment to be administered at a minimum interval of 6-monthly for a maximum of 3 years.

Note: Indications marked with * are unapproved indications.

Initial application — (symptomatic hypercalcaemia*) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has symptomatic hypercalcaemia .

Note: Indications marked with * are unapproved indications.

Corticosteroids and Related Agents for Systemic Use

Inj 3.9 mg with betamethasone acetate 3 mg per ml, 1 ml	TE 5	
(36.96)	J	Celestone
()		Chronodose
EXAMETHASONE		
Tab 0.5 mg - Up to 60 tab available on a PSO	30	Dexmethsone
Tab 4 mg - Up to 30 tab available on a PSO2.65	30	✓ Dexmethsone
Oral liq 1 mg per ml48.15	25 ml OP	✓ Biomed
EXAMETHASONE PHOSPHATE		
Dexamethasone phosphate injection will not be funded for oral use.		
Inj 4 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSO7.86	10	✓ Hameln
9.25		 Dexamethasone
		Phosphate
		Panpharma
Inj 4 mg per ml, 2 ml ampoule – Up to 5 inj available on a PSO 13.10	10	✓ Hameln
16.37		✓ Dexamethason
		Phosphate
		Panpharma
Novamathacana Phaanhata Pannharma Ini 1 mg nar ml. 1 ml amnaula ta ha da		m, 2022)
, , , , , , , , , , , , , , , , , , , ,		• '
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de		• '
examethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de UDROCORTISONE ACETATE	listed 1 Februa	ry 2023)
lexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de .UDROCORTISONE ACETATE Tab 100 mcg11.46		• '
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg11.46 Florinef to be Principal Supply on 1 December 2022	listed 1 Februa	ry 2023)
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg11.46 Florinef to be Principal Supply on 1 December 2022 YDROCORTISONE	listed 1 Februa. 100	ry 2023) ✓ Florinef
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg	listed 1 Februa 100 100	ry 2023) ✓ Florinef ✓ Douglas
Lexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg	100 100 100 100 100	ry 2023) ✓ Florinef ✓ Douglas ✓ Douglas
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg	listed 1 Februa 100 100	ry 2023) ✓ Florinef ✓ Douglas
Nexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de	100 100 100 100 100	ry 2023) ✓ Florinef ✓ Douglas ✓ Douglas
Dexamethasone Phosphate Panpharma Inj 4 mg per ml, 2 ml ampoule to be de LUDROCORTISONE ACETATE Tab 100 mcg	100 100 100 100 100	ry 2023) ✓ Florinef ✓ Douglas ✓ Douglas
Florinef to be Principal Supply on 1 December 2022 YDROCORTISONE Tab 5 mg	100 100 100 100 100	ry 2023) ✓ Florinef ✓ Douglas ✓ Douglas

Redipred Restricted to children under 12 years of age. REDNISONE RETAB 1 mg		Subsidy		Fully	Brand or
Inj 40 mg vial					
Inj 40 mg vial		\$	Per		Manutacturer
Inj 125 mg vial					
Inj 1500 mg vial	Inj 40 mg vial	22.30	1	✓ ;	
Inj 1 g vial	Inj 125 mg vial	34.10	1	✓ 9	
METHYLPREDNISOLONE ACETATE	Inj 500 mg vial	26.88	1	✓ (
Inj 40 mg per ml, 1 ml vial	, ,	32.84	1	√ 9	Solu-Medrol
PREDNISOLONE Prednisone Prednisone			_		
Redipred Restricted to children under 12 years of age. RREDNISONE R Tab 1 mg	Inj 40 mg per ml, 1 ml vial	47.06	5	/ I	Depo-Medrol
Restricted to children under 12 years of age. PREDNISONE Tab 1 mg	PREDNISOLONE			_	
Tab 1 mg	Restricted to children under 12 years of age.	6.00	30 ml OP	√ <u>I</u>	<u>Redipred</u>
★ Tab 2.5 mg 21.04 500 ✓ Apo-Prednisone Clinect ★ Tab 5 mg Up to 30 tab available on a PSO 19.30 500 ✓ Apo-Prednisone Clinect ★ Tab 20 mg Up to 30 tab available on a PSO 50.51 500 ✓ Apo-Prednisone Clinect ✔ Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) ✓ Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) ✓ Apo-Prednisone Clinect ✔ Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) ✓ Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) ✔ Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) ✓ Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) ✔ Inj 250 mcg per ml, 1 ml ampoule 75.00 1 ✓ AU Synacthen ★ Inj 1 mg per ml, 1 ml ampoule 690.00 1 ✓ Synacthen Depot ✔ Synacthen Depot ✓ Synacthen Retard \$200 ✓ Synacthen Depot ✔ Synacthen Inj 250 mcg per ml, 1 ml ampoule to be delisted 1 March 2023) Telegrate New Year Contracted to the Year Contracted To the New Year Contracted To the New Year Contracted To the New Year Contracted To the Year Contracted To the Y		10.50			
★ Tab 2.5 mg 21.04 500 ✓ Apo-Prednisone Prednisone Clinect ★ Tab 5 mg Up to 30 tab available on a PSO 19.30 500 ✓ Apo-Prednisone Clinect ★ Tab 20 mg Up to 30 tab available on a PSO 50.51 500 ✓ Apo-Prednisone Clinect ★ Tab 20 mg Up to 30 tab available on a PSO 50.51 500 ✓ Apo-Prednisone Clinect ★ Prednisone Tab 1 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.0 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Clinect Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) **ETRACOSACTRIN * Ju Synacthen ** Syna	* Tab 1 mg	18.58	500		•
★ Tab 5 mg — Up to 30 tab available on a PSO	¥- Tah 2.5 mg	21.04	500	_	
** Tab 5 mg — Up to 30 tab available on a PSO	1 ab 2.3 mg	21.07	300		•
Frednisone Clinect Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) TETRACOSACTRIN Inj 250 mcg per ml, 1 ml ampoule	* Tab 5 mg - Up to 30 tab available on a PSO	19.30	500		
Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) TETRACOSACTRIN k Inj 250 mcg per ml, 1 ml ampoule					
Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) Apo-Prednisone Tab 20 mg to be delisted 1 November 2022) TETRACOSACTRIN Inj 250 mcg per ml, 1 ml ampoule	* Tab 20 mg - Up to 30 tab available on a PSO	50.51	500		•
k Inj 250 mcg per ml, 1 ml ampoule	(Apo-Prednisone Tab 1 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 2.5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 5 mg to be delisted 1 November 2022) (Apo-Prednisone Tab 20 mg to be delisted 1 November 2022)				
✓ Synacthen ✓ UK Synacthen ✓ Synacthen ✓ UK Synacthen Depot ✓ Synacthen Depot ✓ Synacthen Depot ✓ Synacthen Depot ✓ Synacthene Retard S29 RIAMCINOLONE ACETONIDE Inj 10 mg per ml, 1 ml ampoule Inj 40 mg per ml, 1 ml ampoule Inj 40 mg per ml, 1 ml ampoule Sex Hormones Non Contraceptive Androgen Agonists and Antagonists CYPROTERONE ACETATE Tab 50 mg Tab 100 mg 28.03 50 ✓ Siterone Siterone Siterone TESTOSTERONE					
Inj 1 mg per ml, 1 ml ampoule	** Inj 250 mcg per mi, 1 mi ampoule	75.00	1	✓ 9	Synacthen
AU Synacthen Inj 250 mcg per ml, 1 ml ampoule to be delisted 1 March 2023) TRIAMCINOLONE ACETONIDE Inj 10 mg per ml, 1 ml ampoule 20.80 5 Kenacort-A 10 lnj 40 mg per ml, 1 ml ampoule 51.10 5 Kenacort-A 40 Sex Hormones Non Contraceptive Androgen Agonists and Antagonists EYPROTERONE ACETATE Tab 50 mg 14.37 50 Siterone Tab 100 mg 28.03 50 Siterone TESTOSTERONE	* Inj 1 mg per ml, 1 ml ampoule	690.00	1	✓ 9	Synacthen Depot Synacthene
RIAMCINOLONE ACETONIDE Inj 10 mg per ml, 1 ml ampoule	(AU Synacthen Ini 250 mcg per ml. 1 ml ampoule to be delisted 1	March 2023)			Tiotaru -
Inj 10 mg per ml, 1 ml ampoule	, , , ,				
Inj 40 mg per ml, 1 ml ampoule		20.80	5	✓ I	Cenacort-Δ 10
Sex Hormones Non Contraceptive Androgen Agonists and Antagonists CYPROTERONE ACETATE Tab 50 mg					
Androgen Agonists and Antagonists CYPROTERONE ACETATE Tab 50 mg	, . 91. , 1			-	
CYPROTERONE ACETATE 14.37 50 Siterone Tab 50 mg 28.03 50 Siterone TESTOSTERONE Siterone 14.37	Sex Hormones Non Contraceptive				
Tab 50 mg 14.37 50 ✓ Siterone Tab 100 mg 28.03 50 ✓ Siterone	Androgen Agonists and Antagonists				
Tab 100 mg	CYPROTERONE ACETATE				
ESTOSTERONE	Tab 50 mg	14.37	50		
	Tab 100 mg	28.03	50	√ 9	<u>Siterone</u>
Patch 5 mg per day90.00 30	TESTOSTERONE				
	Patch 5 mg per day	90.00	30	1	Androderm

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price)		Fully Subsidised	I Generic
	\$	Per		Manufacturer
TESTOSTERONE CIPIONATE				
Inj 100 mg per ml, 10 ml vial	85.00	1	✓	Depo-Testosterone
	393.00		✓	Taro-
				Testosterone S29
TESTOSTERONE ESTERS				
Inj 250 mg per ml, 1 ml	12.98	1	✓	Sustanon Ampoules
TESTOSTERONE UNDECANOATE				
Cap 40 mg - Subsidy by endorsement	21.00	60	✓	Andriol Testocaps
	35.00	100	1	Steril-Gene S29
Subsidy by endorsement – subsidised for patients who was 1 November 2021 and the prescription is endorsed accomplete where there exists a record of prior dispensing of testost Inj 250 mg per ml, 4 ml vial	rdingly. Pharmacists erone undecanoate c	may	annotate mg in the	the prescription as endorsed

Hormone Replacement Therapy - Systemic

Oestrogens

ESTRADIOL		
₹ Tab 1 mg4.12	28 OP	
(11.10)		Estrofem
₹ Tab 2 mg4.12	28 OP	
(11.10)		Estrofem
Patch 50 mcg per 24 hours	4	✓ Climara
a) No more than 1 patch per week		
b) Only on a prescription		
Patch 25 mcg per day6.12	8	✓ Estradot
13.50		✓ Estraderm MX S29
a) No more than 2 patch per week		- Louddilli MX
b) Only on a prescription		
, , ,	8	✓ Estradot 50 mcg
Patch 50 mcg per day7.04 9.22	0	✓ Estradiol TDP
9.22		
		Mylan S29
14.50		✓ Estraderm MX S29
a) No more than 2 patch per week		
b) Only on a prescription		
Patch 75 mcg per day7.91	8	Estradot
10.60		Estradiol TDP
		Mylan S29
a) No more than 2 patch per week		-
b) Only on a prescription		
Patch 100 mcg per day7.91	8	✓ Estradot
15.50		✓ Estraderm MX S29
a) No more than 2 patch per week		
b) Only on a prescription		
, , , , ,		
ESTRADIOL VALERATE	0.4	4.5
F Tab 1 mg	84	✓ Progynova
F Tab 2 mg	84	Progynova

	Subsidy (Manufacturer's Pr \$	ice) Sub Per	Fully Brand or sidised Generic Manufacturer
ESTROGENS			
Conjugated, equine tab 300 mcg	3.01	28	
	(17.50)		Premarin
Conjugated, equine tab 625 mcg	4.12	28	
	(17.50)		Premarin
Progestogens			
EDROXYPROGESTERONE ACETATE			
F Tab 2.5 mg	4.69	30	✓ Provera
· ·	8.75	56	✓ Provera
F Tab 5 mg	9.80	56	✓ Provera
	17.50	100	✓ Provera
Tab 10 mg	8.94	30	✓ Provera
Progestogen and Oestrogen Combined Pre	parations		
ESTRADIOL WITH NORETHISTERONE			
Tab 1 mg with 0.5 mg norethisterone acetate	5.40	28 OP	
ů ů	(18.10)		Kliovance
Tab 2 mg with 1 mg norethisterone acetate	5.40	28 OP	
	(18.10)		Kliogest
Tab 2 mg with 1 mg norethisterone acetate (10), and 2	mg		-
oestradiol tab (12) and 1 mg oestradiol tab (6)	5.40	28 OP	
	(18.10)		Trisequens
Other Oestrogen Preparations			
THINYLOESTRADIOL – Subsidy by endorsement			
Subsidy by endorsement – Subsidised for patients who	ware taking ethinyloes	tradial prior t	o 1 March 2022 and the
prescription is endorsed accordingly. Pharmacists may			
prior dispensing of ethinyloestradiol.	annotate the prescript	ion as chaoi	SCU WHOLE HIGHE CAISIS & 1000
Tab 10 mcg	17.60	100	✓ NZ Medical and
			Scientific
NZ Medical and Scientific Tab 10 mcg to be delisted 1 Feb	ruary 2023)		
ESTRIOL	, ,		
: Tab 2 mg	7.00	30	✓ Ovestin
·			<u> </u>
Other Progestogen Preparations			
EVONORGESTREL			
Intra-uterine device 52 mg	269.50	1	✓ Mirena
Intra-uterine device 13.5 mg		1	✓ Jaydess
			
EDBOXYPROGESTERONE ACETATE			4 B UD
EDROXYPROGESTERONE ACETATE Tab 100 mg	116 15	100	✓ Provera HD
Tab 100 mg	116.15	100	✓ Provera HD
Tab 100 mg ORETHISTERONE			
Tab 100 mg ORETHISTERONE : Tab 5 mg – Up to 30 tab available on a PSO		100 30	✓ Primolut N
Tab 100 mg ORETHISTERONE Tab 5 mg – Up to 30 tab available on a PSO ROGESTERONE	5.49		
	5.49 ext page		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy	Fully	Brand or
(Man	ufacturer's Price)	Subsidised	Generic
	\$ P	Per 🗸	Manufacturer

⇒SA1609 Special Authority for Subsidy

Initial application only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 For the prevention of pre-term labour*; and
- 2 Fither:
 - 2.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 2.2 The patient has a history of pre-term birth at less than 28 weeks.

Renewal only from an obstetrician or gynaecologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 For the prevention of pre-term labour*; and
- 2 Treatment is required for second or subsequent pregnancy; and
- 3 Either:
 - 3.1 The patient has a short cervix on ultrasound (defined as < 25 mm at 16 to 28 weeks); or
 - 3.2 The patient has a history of pre-term birth at less than 28 weeks.

Note: Indications marked with * are unapproved indications.

Thyroid and Antithyroid Agents

	RBIMAZOLE Tab 5 mg	7.56	100	✓ Neo-Mercazole
	VOTHYROXINE			
*	Tab 25 mcg	5.55	90	✓ Synthroid
*	Tab 50 mcg	1.71	28	✓ Mercury Pharma
	•	5.79	90	✓ Synthroid
		64.28	1,000	✓ Eltroxin
*	Tab 100 mcg	1.78	28	✓ Mercury Pharma
	•	6.01	90	✓ Synthroid
		66.78	1,000	✓ Eltroxin
PF	OPYLTHIOURACIL - Special Authority see S	A1199 below – Retail pharmacy		
	Tab 50 mg	35.00	100	✓ PTU S29

⇒SA1199 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The patient has hyperthyroidism; and
- 2 The patient is intolerant of carbimazole or carbimazole is contraindicated.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from the treatment.

Trophic Hormones

Growth Hormones

SOMATROPIN (OMNITROPE) - Special Authority see SA2032 on the next page - Retail pharmacy

-		poolar ridiriority coo or Look page	- Hotali pila	maoy
*	Inj 5 mg cartridge	69.75	1	Omnitrope
*	Inj 10 mg cartridge	69.75	1	✓ Omnitrope
		139.50	1	✓ Omnitrope

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA2032 Special Authority for Subsidy

Initial application — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Growth hormone deficiency causing symptomatic hypoglycaemia, or with other significant growth hormone deficient sequelae (e.g. cardiomyopathy, hepatic dysfunction) and diagnosed with GH < 5 mcg/l on at least two random blood samples in the first 2 weeks of life, or from samples during established hypoglycaemia (whole blood glucose < 2 mmol/l using a laboratory device); or</p>
- 2 All of the following:
 - 2.1 Height velocity < 25th percentile for age adjusted for bone age/pubertal status if appropriate over 6 or 12 months using the standards of Tanner and Davies (1985); and
 - 2.2 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
 - 2.3 Peak growth hormone value of < 5.0 mcg per litre in response to two different growth hormone stimulation tests. In children who are 5 years or older, GH testing with sex steroid priming is required; and
 - 2.4 If the patient has been treated for a malignancy, they should be disease free for at least one year based upon follow-up laboratory and radiological imaging appropriate for the malignancy, unless there are strong medical reasons why this is either not necessary or appropriate; and
 - 2.5 Appropriate imaging of the pituitary gland has been obtained.

Renewal — (growth hormone deficiency in children) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 2 Height velocity is greater than or equal to 25th percentile for age (adjusted for bone age/pubertal status if appropriate) while on growth hormone treatment, as calculated over six months using the standards of Tanner and Davis (1985); and
- 3 Height velocity is greater than or equal to 2.0 cm per year, as calculated over 6 months; and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (**Turner syndrome**) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a post-natal genotype confirming Turner Syndrome; and
- 2 Height velocity is < 25th percentile over 6-12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is < 14 years.

Renewal — (Turner syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile for age (while on growth hormone calculated over 6 to 12 months using the Ranke's Turner Syndrome growth velocity charts); and
- 2 Height velocity is greater than or equal to 2 cm per year, calculated over six months; and
- 3 A current bone age is 14 years or under; and
- 4 No serious adverse effect that the specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed since starting growth hormone.

Initial application — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient's height is more than 3 standard deviations below the mean for age or for bone age if there is marked growth acceleration or delay; and
- 2 Height velocity is < 25th percentile for age (adjusted for bone age/pubertal status if appropriate), as calculated over 6 to 12 months using the standards of Tanner and Davies(1985); and
- 3 A current bone age is < 14 years or under (female patients) or < 16 years (male patients); and
- 4 The patient does not have severe chronic disease (including malignancy or recognized severe skeletal dysplasia) and is not receiving medications known to impair height velocity.

Renewal — (short stature without growth hormone deficiency) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred.

Initial application — **(short stature due to chronic renal insufficiency)** only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient's height is more than 2 standard deviations below the mean; and
- 2 Height velocity is < 25th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 3 A current bone age is to 14 years or under (female patients) or to 16 years or under (male patients); and
- 4 The patient is metabolically stable, has no evidence of metabolic bone disease and absence of any other severe chronic disease; and
- 5 The patient is under the supervision of a specialist with expertise in renal medicine; and
- 6 Either:
 - 6.1 The patient has a GFR less than or equal to 30 ml/min/1.73m² as measured by the Schwartz method (Height(cm)/plasma creatinine (umol/l) × 40 = corrected GFR (ml/min/1.73m² in a child who may or may not be receiving dialysis; or
 - 6.2 The patient has received a renal transplant and has received < 5mg/ m²/day of prednisone or equivalent for at least 6 months..

Renewal — (short stature due to chronic renal insufficiency) only from a paediatric endocrinologist, endocrinologist or renal physician on the recommendation of a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patients specialist considers is likely to be attributable to growth hormone has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not experienced significant biochemical or metabolic deterioration confirmed by diagnostic results; and
- 7 The patient has not received renal transplantation since starting growth hormone treatment; and
- 8 If the patient requires transplantation, growth hormone prescription should cease before transplantation and a new application should be made after transplantation based on the above criteria.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of Prader-Willi syndrome that has been confirmed by genetic testing or clinical scoring criteria; and
- 2 The patient is aged six months or older; and
- 3 A current bone age is < 14 years (female patients) or < 16 years (male patients); and
- 4 Sleep studies or overnight oximetry have been performed and there is no obstructive sleep disorder requiring treatment, or if an obstructive sleep disorder is found, it has been adequately treated under the care of a paediatric respiratory physician and/or ENT surgeon; and
- 5 Fither:
 - 5.1 Both:
 - 5.1.1 The patient is aged two years or older; and
 - 5.1.2 There is no evidence of type II diabetes or uncontrolled obesity defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months; or
 - 5.2 The patient is aged between six months and two years and a thorough upper airway assessment is planned to be undertaken prior to treatment commencement and at six to 12 weeks following treatment initiation.

Renewal — (Prader-Willi syndrome) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Height velocity is greater than or equal to 50th percentile (adjusted for bone age/pubertal status if appropriate) as calculated over 6 to 12 months using the standards of Tanner and Davies (1985); and
- 2 Height velocity is greater than or equal to 2 cm per year as calculated over six months; and
- 3 A current bone age is 14 years or under (female patients) or 16 years or under (male patients); and
- 4 No serious adverse effect that the patient's specialist considers is likely to be attributable to growth hormone treatment has occurred; and
- 5 No malignancy has developed after growth hormone therapy was commenced; and
- 6 The patient has not developed type II diabetes or uncontrolled obesity as defined by BMI that has increased by greater than or equal to 0.5 standard deviations in the preceding 12 months.

Initial application — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a medical condition that is known to cause growth hormone deficiency (e.g. surgical removal of the pituitary for treatment of a pituitary tumour); and
- 2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
- 3 The patient has severe growth hormone deficiency (see notes); and
- 4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
- 5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test.

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Subsid	ly Fu	y Brand or
(Manufacture)	,	
\$	Per	Manufacturer

continued...

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

Renewal — (adults and adolescents) only from a paediatric endocrinologist or endocrinologist. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 All of the following:
 - 1.1 The patient has been treated with somatropin for < 12 months; and
 - 1.2 There has been an improvement in Quality of Life defined as a reduction of at least 8 points on the Quality of Life Assessment of Growth Hormone Deficiency in Adults (QoL-AGHDA®) score from baseline; and
 - 1.3 Serum IGF-I levels have been increased within ±1SD of the mean of the normal range for age and sex; and
 - 1.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients, or 1 mg per day for female patients; or
- 2 All of the following:
 - 2.1 The patient has been treated with somatropin for more than 12 months; and
 - 2.2 The patient has not had a deterioration in Quality of Life defined as a 6 point or greater increase from their lowest QoL-AGHDA® score on treatment (other than due to obvious external factors such as external stressors); and
 - 2.3 Serum IGF-I levels have continued to be maintained within ±1SD of the mean of the normal range for age and sex (other than for obvious external factors); and
 - 2.4 The dose of somatropin has not exceeded 0.7 mg per day for male patients or 1 mg per day for female patients; or
- 3 All of the following:
 - 3.1 The patient has had a Special Authority approval for somatropin for childhood deficiency in children and no longer meets the renewal criteria under this indication; and
 - 3.2 The patient has undergone appropriate treatment of other hormonal deficiencies and psychological illnesses; and
 - 3.3 The patient has severe growth hormone deficiency (see notes); and
 - 3.4 The patient's serum IGF-I is more than 1 standard deviation below the mean for age and sex; and
 - 3.5 The patient has poor quality of life, as defined by a score of 16 or more using the disease-specific quality of life questionnaire for adult growth hormone deficiency (QoL-AGHDA®).

Notes: For the purposes of adults and adolescents, severe growth hormone deficiency is defined as a peak serum growth hormone level of less than or equal to 3 mcg per litre during an adequately performed insulin tolerance test (ITT) or glucagon stimulation test

Patients with one or more additional anterior pituitary hormone deficiencies and a known structural pituitary lesion only require one test. Patients with isolated growth hormone deficiency require two growth hormone stimulation tests, of which, one should be ITT unless otherwise contraindicated. Where an additional test is required, an arginine provocation test can be used with a peak serum growth hormone level of less than or equal to 0.4 mcg per litre.

The dose of somatropin should be started at 0.2 mg daily and be titrated by 0.1 mg monthly until the serum IGF-I is within 1 standard deviation of the mean normal value for age and sex; and

Dose of somatropin not to exceed 0.7 mg per day for male patients, or 1 mg per day for female patients.

At the commencement of treatment for hypopituitarism, patients must be monitored for any required adjustment in replacement doses of corticosteroid and levothyroxine.

GnRH Analogues

	DEI	

Implant 3.6 mg, syringe	65.68	1	✓ Teva
Implant 10.8 mg, syringe		1	✓ Teva

Subsidy

Fully

Brand or

	(Manager at the part of a Date of		Out-idia	Oi-
	(Manufacturer's Price) \$	Per	Subsidised <	Generic Manufacturer
LEUPRORELIN				
Additional subsidy by endorsement where the patient is a chil goserelin and the prescription is endorsed accordingly.	ld or adolescent and	is una	able to tolera	ate administration of
Inj 3.75 mg prefilled dual chamber syringe - Higher subsidy	of			
\$221.60 per 1 inj with Endorsement	66.48	1		
	(221.60)		L	ucrin Depot 1-month
Inj 11.25 mg prefilled dual chamber syringe - Higher subsidy	1			
of \$591.68 per 1 inj with Endorsement	177.50	1		
	(591.68)		L	ucrin Depot 3-month.
Vasopressin Agonists				
DESMOPRESSIN				
Wafer 120 mcg	47.00	30	✓ N	Minirin Melt
DESMOPRESSIN ACETATE				
Tab 100 mcg	25.00	30	✓ N	Minirin
Tab 200 mcg		30	✓ N	Minirin
A Nasal spray 10 mcg per dose	27.95	3 ml O	₽ ✓ <u>[</u>	<u>PH&T</u>
Inj 4 mcg per ml, 1 ml	67.18	10	✓ N	<i>l</i> inirin

CABERGOLINE

Γ

		Tab 0.5 mg - Maximum of 2 tab per prescription; can be
Dostinex	2	waived by Special Authority see SA2070 below
✓ Dostinex	8	15.20

⇒SA2070 Special Authority for Waiver of Rule

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 Hyperprolactinemia; or

Other Endocrine Agents

- 2 Acromegaly*; or
- 3 Inhibition of lactation.

Renewal — (for patients who have previously been funded under Special Authority form SA1031) from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has previously held a valid Special Authority which has expired and the treatment remains appropriate and the patient is benefiting from treatment.

Note: Indication marked with * is an unapproved indication.

α	N 41			Δ		$\overline{}$
CLC	ж	FFI	٧F	GH	IKA	11

Tab 50 mg	29.84	10	✓ Mylan Clomiphen S29
METYRAPONE Cap 250 mg	558.00	50	✓ Metopirone

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	1	Manufacturer	

Anthelmintics

ALBENDAZOLE - Special Authority see SA1318 below	- Retail pharmacy		
Tab 400 mg	469.20	60	✓ Eskazole S29

⇒SA1318 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the patient has hydatids.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

MEBENDAZOLE - Only on a prescription

Tab 100 mg	7.97	6	✓ Vermox
Oral liq 100 mg per 5 ml	2.18	15 ml	
	(7.53)		Vermox
PRAZIQUANTEL			
Tab 600 mg	68.00	8	✓ Biltricide

Antibacterials

CEFACLOR MONOHYDRATE

- a) For topical antibacterials, refer to DERMATOLOGICALS, page 62
- b) For anti-infective eye preparations, refer to SENSORY ORGANS, page 241

Cephalosporins and Cephamycins

Cap 250 mg24.70	100	 ✓ Ranbaxy-Cefaclor ✓ Ranbaxy-Cefaclor S29 S29
Grans for oral liq 125 mg per 5 ml - Wastage claimable3.53	100 ml	 ✓ Ranbaxy-Cefaclor ✓ Ranbaxy-Cefaclor S29 S29
CEFALEXIN		
Cap 250 mg3.33	20	 Cephalexin ABM
Cap 500 mg5.95	20	✓ Cephalexin ABM
Grans for oral liq 25 mg per ml - Wastage claimable7.88	100 ml	✓ Flynn
8.75		✓ Cefalexin Sandoz
Grans for oral liq 50 mg per ml - Wastage claimable10.38	100 ml	✓ Flynn
11.75		✓ Cefalexin Sandoz
(Cefalexin Sandoz Grans for oral liq 25 mg per ml to be delisted 1 January 2023)		

(Cetalexin Sandoz Grans for oral liq 25 mg per ml to be delisted 1 January 2023) (Cefalexin Sandoz Grans for oral liq 50 mg per ml to be delisted 1 January 2023)

CEFAZOLIN - Subsidy by endorsement

Only if prescribed for dialysis or cellulitis in accordance with a Health NZ Hospital approved protocol and the prescription is endorsed accordingly.

Inj 500 mg vial	3.39	5	✓ AFT
lnj 1 g vial	3.49	5	✓ <u>AFT</u>

	Subsidy (Manufacturer's Price) \$	Sub Per	Fully osidised	Brand or Generic Manufacturer
CEFTRIAXONE – Subsidy by endorsement				
a) Up to 10 inj available on a PSO				
b) Subsidised only if prescribed for a dialysis or cystic fibros	sis patient, or the treat	ment of	gonorrho	ea, or the treatment o
pelvic inflammatory disease, or the treatment of suspecte	ed meningococcal dise	ease, and	the pres	scription or PSO is
endorsed accordingly.	-			
Inj 500 mg vial	0.89	1	√ C	eftriaxone-AFT
Inj 1 g vial	3.99	5	✓ C	eftriaxone-AFT
CEFUROXIME AXETIL - Subsidy by endorsement				
Only if prescribed for prophylaxis of endocarditis and the pre	scription is endorsed	accordin	alv.	
Tab 250 mg		50	0,	innat

Macrolides

AZITHROMYCIN – Maximum of 5 days treatment per prescription; can be waived by Special Authority see SA1683 below A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised on Special Authority.

Tab 250 mg8.	19 30	✓ Apo-Azithromycin
Tab 500 mg - Up to 8 tab available on a PSO2.5		✓ Zithromax
Grans for oral liq 200 mg per 5 ml (40 mg per ml) - Wastage		
claimable16.9	97 15 ml	✓ Zithromax

⇒SA1683 Special Authority for Waiver of Rule

(Zinnat Tab 250 mg to be delisted 1 March 2024)

Initial application — (bronchiolitis obliterans syndrome, cystic fibrosis and atypical Mycobacterium infections) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following:

- 1 Patient has received a lung transplant, stem cell transplant, or bone marrow transplant and requires treatment for bronchiolitis obliterans syndrome*; or
- 2 Patient has received a lung transplant and requires prophylaxis for bronchiolitis obliterans syndrome*; or
- 3 Patient has cystic fibrosis and has chronic infection with Pseudomonas aeruginosa or Pseudomonas-related gram negative organisms*; or
- 4 Patient has an atypical Mycobacterium infection.

Note: Indications marked with * are unapproved indications.

Initial application — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 For prophylaxis of exacerbations of non-cystic fibrosis bronchiectasis*; and
- 2 Patient is aged 18 and under; and
- 3 Fither:
 - 3.1 Patient has had 3 or more exacerbations of their bronchiectasis, within a 12 month period; or
 - 3.2 Patient has had 3 acute admissions to hospital for treatment of infective respiratory exacerbations within a 12 month period.

Note: Indications marked with * are unapproved indications.

Renewal — (non-cystic fibrosis bronchiectasis*) only from a respiratory specialist or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has completed 12 months of azithromycin treatment for non-cystic fibrosis bronchiectasis; and
- 2 Following initial 12 months of treatment, the patient has not received any further azithromycin treatment for non-cystic fibrosis bronchiectasis for a further 12 months, unless considered clinically inappropriate to stop treatment; and

 Subsidy Manufacturer's Price)	S	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

3 The patient will not receive more than a total of 24 months' azithromycin cumulative treatment (see note).

The patient must not have had more than 1 prior approval.

Note: No further renewals will be subsidised. A maximum of 24 months of azithromycin treatment for non-cystic fibrosis bronchiectasis will be subsidised. Indications marked with * are unapproved indications

CLARITHROMYCIN - Maximum of 500 mg per prescription; can be waived by Special Authority see SA1857 below

 Tab 250 mg
 8.53
 14
 ✓ Klacid

 Grans for oral liq 250 mg per 5 ml
 − Wastage claimable
 192.00
 50 ml
 ✓ Klacid

⇒SA1857 Special Authority for Waiver of Rule

Initial application — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Atypical mycobacterial infection; or
- 2 Mycobacterium tuberculosis infection where there is drug-resistance or intolerance to standard pharmaceutical agents.

Initial application — (Helicobacter pylori eradication) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 For the eradication of helicobacter pylori in a patient unable to swallow tablets; and
- 2 For use only in combination with omeprazole and amoxicillin as part of a triple therapy regimen.

Initial application — (Prophylaxis of infective endocarditis) from any relevant practitioner. Approvals valid for 3 months where prophylaxis of infective endocarditis associated with surgical or dental procedures if amoxicillin is contra-indicated. Renewal — (Mycobacterial infections) only from a respiratory specialist, infectious disease specialist or paediatrician. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Inj 1 g vial10.00	1	Erythrocin IV
Erythrocin IV to be Principal Supply on 1 December 2022		
ERYTHROMYCIN ETHYL SUCCINATE		
Tab 400 mg16.95	100	✓ E-Mycin
a) Up to 20 tab available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
Grans for oral liq 200 mg per 5 ml5.00	100 ml	E-Mycin
a) Up to 300 ml available on a PSO		
b) Up to 2 x the maximum PSO quantity for RFPP		
c) Wastage claimable		
Grans for oral liq 400 mg per 5 ml6.77	100 ml	E-Mycin
a) Up to 200 ml available on a PSO		
b) Wastage claimable		
ROXITHROMYCIN		
Tab disp 50 mg8.29	10	Rulide D
Restricted to children under 12 years of age.		
Tab 150 mg8.28	50	✓ Arrow-
		Roxithromycin
Tab 300 mg16.33	50	✓ Arrow-
	30	Roxithromycin

(Rulide D Tab disp 50 mg to be delisted 1 March 2023)

Penicillins AMOXICILLIN Cap 250 mg		Subsidy (Manufacturer's Pr \$	ice) Subs Per	Fully idised	
Cap 250 mg	Penicillins				
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg. a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml. a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 125 mg vift clavulanic acid 12.5 mg per ml. 4 Jup to 300 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable lnj 250 mg vial lnj 1 g vial — Up to 5 inj available on a PSO 21.64 10	AMOXICILLIN				
b) Up to 10 x the maximum PSO quantity for RFPP Cap 500 mg,		22.50	500	✓	Alphamox
Cap 500 mg					
a) Up to 30 cap available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml. a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 250 mg per 5 ml. 1.73 100 ml Alphamox 250 a) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable liq 250 mg vial 1.743 10 Alphamox 250 b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable liq 250 mg vial 1.743 10 Alphamox 250 b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable liq 250 mg vial 1.743 10 Alphamox 250 b) Wastage claimable liq 250 mg with clavulanic acid 125 mg liq yial - Up to 5 in javailable on a PSO AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg - Up to 30 tab available on a PSO. Bornas for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml. Bornas for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml - Up to 200 ml available on a PSO. BENZATHINE BENZYLPENICILLIN liq 900 mg (1, million units) via Up to 2 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600 mg (1 million units) via Up to 5 inj available on a PSO. BENZYLPENICILLIN SODIUM [PENICILLIN G] liq 600	b) Up to 10 x the maximum PSO quantity for RFPP				
Dict 10 x the maximum PSO quantity for RFPP		36.98	500	1	Alphamox
Grans for oral liq 125 mg per 5 ml					
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 250 mg per 5 ml				_	
b) Wastage claimable Grans for oral liq 250 mg per 5 ml		1.40	100 ml		Alphamox 125
Grans for oral liq 250 mg per 5 ml					
a) Up to 300 ml available on a PSO b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable lnj 250 mg vial					
b) Up to 10 x the maximum PSO quantity for RFPP c) Wastage claimable Inj 250 mg vial		1.73	100 ml	•	Alphamox 250
c) Wastage claimable Inj 250 mg vial					
Inj 250 mg vial					
Inj 500 mg vial		15.07	10	./	Ihiamay
Inj 1 g vial − Up to 5 inj available on a PSO					
AMOXICILLIN WITH CLAVULANIC ACID Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO					
Tab 500 mg with clavulanic acid 125 mg − Up to 30 tab available on a PSO		21.07	10	•	IDIGITION
available on a PSO					
Grans for oral liq amoxicillin 25 mg with clavulanic acid 6.25 mg per ml		0.00	10		O D 500/405
per ml			10	•	Curam Duo 500/125
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO			100 ml	./	Accommenting
b) Wastage claimable Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO	•	6.50	100 1111	•	Augmenun
Grans for oral liq amoxicillin 50 mg with clavulanic acid 12.5 mg per ml – Up to 200 ml available on a PSO	, ·				
per ml − Up to 200 ml available on a PSO	, •	~~			
BENZATHINE BENZYLPENICILLIN Inj 900 mg (1.2 million units) in 2.3 ml syringe − Up to 5 inj			100 ml OD	1	Curam
Inj 900 mg (1.2 million units) in 2.3 ml syringe − Up to 5 inj available on a PSO	·	2.20	100 IIII OF	•	Curain
available on a PSO					
BENZYLPENICILLIN SODIUM [PENICILLIN G] Inj 600 mg (1 million units) vial − Up to 5 inj available on a PSO11.09 FLUCLOXACILLIN Cap 250 mg − Up to 30 cap available on a PSO	, , , , , , , , , , , , , , , , , , , ,	075 07	40	,	District A
Inj 600 mg (1 million units) vial − Up to 5 inj available on a PSO11.09 10		3/5.9/	10	•	BICIIIIN LA
FLUCLOXACILLIN Cap 250 mg − Up to 30 cap available on a PSO					
Cap 250 mg − Up to 30 cap available on a PSO	Inj 600 mg (1 million units) vial – Up to 5 inj available on a P	SO 11.09	10	/	Sandoz
Cap 500 mg − Up to 30 cap available on a PSO 52.99 500 ✓ Flucloxacillin-AFT Grans for oral liq 25 mg per ml 3.29 100 ml ✓ AFT a) Up to 200 ml available on a PSO 3.68 100 ml ✓ AFT a) Up to 200 ml available on a PSO 3.68 100 ml ✓ AFT a) Up to 200 ml available on a PSO 50 mg vial 17.56 10 ✓ Flucloxin Inj 250 mg vial 18.87 10 ✓ Flucloxin	FLUCLOXACILLIN				
Grans for oral liq 25 mg per ml					
a) Up to 200 ml available on a PSO b) Wastage claimable Grans for oral liq 50 mg per ml					
b) Wastage claimable Grans for oral liq 50 mg per ml		3.29	100 ml	/	<u>AFT</u>
Grans for oral liq 50 mg per ml	, ,				
a) Up to 200 ml available on a PSO b) Wastage claimable Inj 250 mg vial	b) Wastage claimable	0.00	4001	,	A = T
b) Wastage claimable Inj 250 mg vial		3.ხგ	IUU MI	•	<u>AFI</u>
Inj 250 mg vial 17.56 10 ✓ Flucloxin Inj 500 mg vial 18.87 10 ✓ Flucloxin					
Inj 500 mg vial		17.56	10	_/	Eluciovin
			5		

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Generic
PHENOXYMETHYLPENICILLIN (PENICILLIN V)				
Cap 250 mg - Up to 30 cap available on a PSO	3.84	50	1	Cilicaine VK
Cap 500 mg	6.86	50	✓	Cilicaine VK
a) Up to 20 cap available on a PSO b) Up to 2 x the maximum PSO quantity for RFPP Grans for oral liq 125 mg per 5 ml		100 m	nl 🗸	AFT
Grans for oral liq 250 mg per 5 ml	4.24	100 m	nl 🗸	AFT
PROCAINE PENICILLIN Inj 1.5 g in 3.4 ml syringe — Up to 5 inj available on a PSO (Cilicaine Inj 1.5 g in 3.4 ml syringe to be delisted 1 February 202		5	•	Cilicaine

Tetracyclines

DOXYCYCLINE	
-------------	--

*	Tab 100 mg - Up to 30 tab available on a PSO	64.43	500	Doxine
MIN	NOCYCLINE HYDROCHLORIDE			
*	Tab 50 mg - Additional subsidy by Special Authority see			
	SA1355 below – Retail pharmacy	5.79	60	
		(12.05)		Mino-tabs
*	Cap 100 mg	19.32	100	
		(52.04)		Minomycin

⇒SA1355 Special Authority for Manufacturers Price

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has rosacea.

TETRACYCLINE - Special Authority see SA1332 below - Retail pharmacy

⇒SA1332 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 For the eradication of helicobacter pylori following unsuccessful treatment with appropriate first-line therapy; and
- 2 For use only in combination with bismuth as part of a quadruple therapy regimen.

Other Antibiotics

For topical antibiotics, refer to DERMATOLOGICALS, page 62

CIPROFLOXACIN

Recommended for patients with any of the following:

- i) microbiologically confirmed and clinically significant pseudomonas infection; or
- ii) prostatitis; or
- iii) pyelonephritis; or
- iv) gonorrhoea.

Tab 250 mg - Up to 5 tab available on a PSO	2.42	28	✓ Cipflox
Tab 500 mg - Up to 5 tab available on a PSO	3.40	28	✓ Cipflox
Tab 750 mg	5.95	28	✓ Cipflox

	Subsidy (Manufacturer's Price)		Fully Brand or Subsidised Generic
	(Manufacturer's Frice)	Per	
CLINDAMYCIN			
Cap hydrochloride 150 mg	4.61	24	Dalacin C
Inj phosphate 150 mg per ml, 4 ml ampoule	39.00	10	Dalacin C
COLISTIN SULPHOMETHATE – Retail pharmacy-Specialist – S Only if prescribed for dialysis or cystic fibrosis patient and the	e prescription is endo		
Inj 150 mg	65.00	1	✓ Colistin-Link
GENTAMICIN SULPHATE Inj 10 mg per ml, 1 ml ampoule — Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.		5 trac	✓ DBL Gentamicin of infection and the prescription is
Inj 10 mg per ml, 2 ml ampoule - Subsidy by endorsement	91.00	5	✓ Wockhardt S29
	182.00	10	✓ Teligent S29
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	ct infection and the prescription is
Inj 40 mg per ml, 2 ml ampoule - Subsidy by endorsement	17.50	10	✓ Pfizer
	87.50	50	✓ Pfizer
Only if prescribed for a dialysis or cystic fibrosis patient of endorsed accordingly.	or complicated urinary	trac	ct infection and the prescription is
MOXIFLOXACIN – Special Authority see SA1740 below – Retail No patient co-payment payable	pharmacy		
Tab 400 mg	42.00	5	✓ <u>Avelox</u>

⇒SA1740 Special Authority for Subsidy

Initial application — (Tuberculosis) only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Active tuberculosis*: and
 - 1.2 Any of the following:
 - 1.2.1 Documented resistance to one or more first-line medications; or
 - 1.2.2 Suspected resistance to one or more first-line medications (tuberculosis assumed to be contracted in an area with known resistance), as part of regimen containing other second-line agents; or
 - 1.2.3 Impaired visual acuity (considered to preclude ethambutol use); or
 - 1.2.4 Significant pre-existing liver disease or hepatotoxicity from tuberculosis medications; or
 - 1.2.5 Significant documented intolerance and/or side effects following a reasonable trial of first-line medications;
- 2 Mycobacterium avium-intracellulare complex not responding to other therapy or where such therapy is contraindicated.*; or
- 3 Patient is under five years of age and has had close contact with a confirmed multi-drug resistant tuberculosis case.

Note: Indications marked with * are unapproved indications.

Renewal only from a respiratory specialist or infectious disease specialist. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Mycoplasma genitalium) only from a sexual health specialist or Practitioner on the recommendation of a sexual health specialist. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Has nucleic acid amplification test (NAAT) confirmed Mycoplasma genitalium* and is symptomatic; and
- 2 Fither:
 - 2.1 Has tried and failed to clear infection using azithromycin; or
 - 2.2 Has laboratory confirmed azithromycin resistance; and

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer continued... 3 Treatment is only for 7 days. Initial application — (Penetrating eye injury) only from an ophthalmologist. Approvals valid for 1 month where the patient requires prophylaxis following a penetrating eve injury and treatment is for 5 days only. Note: Indications marked with * are unapproved indications. PAROMOMYCIN - Special Authority see SA1689 below - Retail pharmacy ✓ Humatin S29 16 ⇒SA1689 Special Authority for Subsidy Initial application only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Either: 1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage. Renewal only from an infectious disease specialist, clinical microbiologist or gastroenterologist. Approvals valid for 1 month for applications meeting the following criteria: Fither: 1 Patient has confirmed cryptosporidium infection; or 2 For the eradication of Entamoeba histolyica carriage. PYRIMETHAMINE - Special Authority see SA1328 below - Retail pharmacy 30 ✓ Daraprim S29 ⇒SA1328 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. SODIUM FUSIDATE [FUSIDIC ACID] 36 ✓ Fucidin Tab 250 mg67.85 SULFADIAZINE SODIUM - Special Authority see SA1331 below - Retail pharmacy 56 Tab 500 mg543.20 ✓ Wockhardt \$29 ⇒SA1331 Special Authority for Subsidy Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Any of the following: 1 For the treatment of toxoplasmosis in patients with HIV for a period of 3 months; or 2 For pregnant patients for the term of the pregnancy; or 3 For infants with congenital toxoplasmosis until 12 months of age. **TOBRAMYCIN** Inj 40 mg per ml, 2 ml vial - Subsidy by endorsement......18.50 ✓ Tobramvcin Mylan Only if prescribed for dialysis or cystic fibrosis patient and the prescription is endorsed accordingly. Solution for inhalation 60 mg per ml, 5 ml - Subsidy by ✓ Tobramvcin BNM endorsement......395.00 56 dose a) Wastage claimable b) Only if prescribed for a cystic fibrosis patient and the prescription is endorsed accordingly. TRIMETHOPRIM * Tab 300 mg - Up to 30 tab available on a PSO......18.55 TMP

	Subsidy (Manufacturer's Price \$	e) Subs	Fully sidised	Brand or Generic Manufacturer
TRIMETHOPRIM WITH SULPHAMETHOXAZOLE [CO-TRIMOX	AZOLE]			
* Tab trimethoprim 80 mg and sulphamethoxazole 400 mg – L to 30 tab available on a PSO	64.80	500	√ <u>T</u>	<u>risul</u>
** Oral liq 8 mg sulphamethoxazole 40 mg per ml - Up to 200 r available on a PSO		100 ml	✓ D	eprim
VANCOMYCIN – Subsidy by endorsement Only if prescribed for a dialysis or cystic fibrosis patient or for difficile following metronidazole failure and the prescription is Inj 500 mg vial	endorsed accordin			tment of Clostridium

Antifungals

- a) For topical antifungals refer to DERMATOLOGICALS, page 63
- b) For topical antifungals refer to GENITO URINARY, page 76

FLUCONAZOLE

Cap 50 mg	28	✓ Dizole✓ Mylan
Cap 150 mg	1	✓ Mylan
Cap 200 mg12.89	28	✓ Mylan
Powder for oral suspension 10 mg per ml - Special Authority		
see SA1359 below – Retail pharmacy109.34	35 ml	Diflucan
Wastage claimable		

⇒SA1359 Special Authority for Subsidy

Initial application — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Initial application — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Patient is at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

Renewal — (Systemic candidiasis) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires prophylaxis for, or treatment of systemic candidiasis; and
- 2 Patient is unable to swallow capsules.

Renewal — (Immunocompromised) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient remains immunocompromised; and
- 2 Patient remains at moderate to high risk of invasive fungal infection; and
- 3 Patient is unable to swallow capsules.

ITRACONAZOLE

Cap 100 mg4.2	7 15	✓ Itrazole
Oral lig 10 mg per ml - Special Authority see SA1322 on the		
next page – Retail pharmacy141.8	0 150 ml OP	✓ Sporanox

Subsidy Fully		Fully	Brand or
(Manufacturer's Price)	S	ubsidised	Generic
\$	Per	✓	Manufacturer

⇒SA1322 Special Authority for Subsidy

Initial application only from an infectious disease specialist, clinical microbiologist, clinical immunologist or any relevant practitioner on the recommendation of a infectious disease physician, clinical microbiologist or clinical immunologist. Approvals valid for 6 months where the patient has a congenital immune deficiency.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefitting from the treatment.

KETOCONAZOI E

Tab 200 mg - PCT	CBS	30	✓ Link Healthcare S29
			✓ Nizoral S29
		100	✓ Strides Shasun S29
NYSTATIN			
Tab 500,000 u	14.16	50	
	(17.09)		Nilstat
Cap 500,000 u	12.81	50	
•	(15.47)		Nilstat
POSACONAZOLE - Special Authority see SA1285 be	elow – Retail pharmacy		
Tab modified-release 100 mg	869.86	24	✓ Noxafil
Oral liq 40 mg per ml	761.13	105 ml OP	✓ Noxafil

⇒SA1285 Special Authority for Subsidy

Initial application only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation chemotherapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppressive therapy*.

Renewal only from a haematologist or infectious disease specialist. Approvals valid for 6 weeks for applications meeting the following criteria:

Either:

- 1 Patient has acute myeloid leukaemia and is to be treated with high dose remission induction, re-induction or consolidation therapy; or
- 2 Patient has received a stem cell transplant and has graft versus host disease and is on significant immunosuppression* and requires on going posaconazole treatment.

Note: * Graft versus host disease (GVHD) on significant immunosuppression is defined as acute GVHD, grade II to IV, or extensive chronic GVHD, or if they were being treated with intensive immunosuppressive therapy consisting of either high-dose corticosteroids (1 mg or greater per kilogram of body weight per day for patients with acute GVHD or 0.8 mg or greater per kilogram every other day for patients with chronic GVHD), antithymocyte globulin, or a combination of two or more immunosuppressive agents or types of treatment.

TERRINAFINE

* Tab 250 mg	8.15	84	✓ Deolate
VORICONAZOLE - Special Authority see SA1273 on the next pa		macy	
Tab 50 mg	•	[*] 56	✓ Vttack
Tab 200 mg	350.00	56	✓ Vttack
Powder for oral suspension 40 mg per ml - Wastage			
claimable	1,523.22	70 ml	✓ Vfend

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA1273 Special Authority for Subsidy

Initial application — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient has proven or probable invasive aspergillus infection; or
 - 3.2 Patient has possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis; or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Renewal — (invasive fungal infection) only from a haematologist, infectious disease specialist or clinical microbiologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient is immunocompromised; and
- 2 Applicant is part of a multidisciplinary team including an infectious disease specialist; and
- 3 Any of the following:
 - 3.1 Patient continues to require treatment for proven or probable invasive aspergillus infection; or
 - 3.2 Patient continues to require treatment for possible invasive aspergillus infection; or
 - 3.3 Patient has fluconazole resistant candidiasis: or
 - 3.4 Patient has mould strain such as Fusarium spp. and Scedosporium spp.

Antimalarials

PRIMAQUINE - Special Authority see SA1684 below - R	etail pharmacy		
Tab 15 mg	400.00	100	✓ Sanofi
			Primaguine \$29

⇒SA1684 Special Authority for Subsidy

Initial application only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has vivax or ovale malaria; and
- 2 Primaquine is to be given for a maximum of 21 days.

Renewal only from an infectious disease specialist or clinical microbiologist. Approvals valid for 1 month for applications meeting the following criteria:

Both:

- 1 The patient has relapsed vivax or ovale malaria; and
- 2 Primaguine is to be given for a maximum of 21 days.

Antitrichomonal Agents

METRONIDAZOLE			
Tab 200 mg - Up to 30 tab available on a PSO	33.15	250	✓ Metrogyl
Tab 400 mg - Up to 15 tab available on a PSO	5.23	21	✓ Metrogyl
Oral liq benzoate 200 mg per 5 ml	25.00	100 ml	✓ Flagyl-S
Suppos 500 mg		10	✓ Flagyl
ORNIDAZOLE			
Tab 500 mg	36.16	10	✓ Arrow-Ornidazole

	Subsidy (Manufacturer's Price) \$	S Per	Fully Subsidised	Brand or Generic Manufacturer
Antituberculotics and Antileprotics				
Note: There is no co-payment charge for all pharmaceuticals lis immigration status.	ted in the Antitubercul	lotics a	nd Antilepi	rotics group regardless of
CLOFAZIMINE - Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda dermatologist. 	tion of, an infectious d	lisease	physician,	clinical microbiologist or
* Cap 50 mg	442.00	100	✓ L	amprene S29
CYCLOSERINE - Retail pharmacy-Specialist				•
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician. 				-
Cap 250 mg	344.00	60	✓ (Cyclorin S29
DAPSONE - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda dermatologist				
Tab 25 mg		100		Dapsone
Tab 100 mg		100	V L	Dapsone
ETHAMBUTOL HYDROCHLORIDE - Retail pharmacy-Special	st			
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician 	tion of, an infectious d	lisease	physician,	clinical microbiologist or
Tab 100 mg	85.73	100	✓ E	MB Fatol S29
Tab 400 mg	49.34	56	✓ N	Myambutol S29
ISONIAZID - Retail pharmacy-Specialist				
 a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician 	tion of, an internal me	dicine ¡	ohysician,	paediatrician, clinical
* Tab 100 mg	23.00	100	√ F	PSM
ISONIAZID WITH RIFAMPICIN - Retail pharmacy-Specialist			_	
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda microbiologist, dermatologist or public health physician	tion of, an internal me	dicine ¡	ohysician,	paediatrician, clinical
* Tab 100 mg with rifampicin 150 mg	89.82	100	√ F	Rifinah
* Tab 150 mg with rifampicin 300 mg		100	✓ <u>F</u>	Rifinah
PARA-AMINO SALICYLIC ACID - Retail pharmacy-Specialist				
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician	tion of, an infectious d	lisease	specialist,	clinical microbiologist or
Grans for oral liq 4 g sachet	280.00	30	√ F	Paser \$29
PROTIONAMIDE – Retail pharmacy-Specialist		00		4001
a) No patient co-payment payable b) Prescriptions must be written by, or on the recommenda respiratory physician	tion of, an infectious d	lisease	specialist,	clinical microbiologist or
Tab 250 mg	305.00	100	✓ F	Peteha S29
✓ fully subsidised	\$29 Unapprove	d modici	ing gunnlind	under Coation 20

	IIA	IEECTIONS A	CENT	C EOD (EVETEMIC LICE
	IIV	IFECTIONS - A	GENI	3 FUR 8	SYSTEMIC USE
		Subsidy (Manufacturer's Price) \$	S Per	Fully ubsidised	Brand or Generic Manufacturer
PY	RAZINAMIDE - Retail pharmacy-Specialist				
	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation	on of, an infectious of	disease	physician,	clinical microbiologist or
*	respiratory physician Tab 500 mg	64.95	100	√ ∆	FT-Pyrazinamide
	FABUTIN – Retail pharmacy-Specialist	04.00	100	• •	i i-i yrazmamiac
1 1111	a) No patient co-payment payable				
	b) Prescriptions must be written by, or on the recommendation gastroenterologist	on of, an infectious o	disease	physician,	respiratory physician or
*	Cap 150 mg	299.75	30	✓ M	lycobutin
RIF	AMPICIN – Subsidy by endorsement				
	 a) No patient co-payment payable b) For confirmed recurrent Staphylococcus aureus infection in antimicrobial based on susceptibilities and the prescription Retail pharmacy - Specialist. Specialist must be an interna paediatrician, or public health physician. 	is endorsed accord	dingly; c	an be waiv	red by endorsement -
*	Cap 150 mg	58.54	100	_	<u>ifadin</u>
	Cap 300 mg		100		ifadin
*	Oral liq 100 mg per 5 ml	12.60	60 ml	✓ <u>R</u>	<u>ifadin</u>
A	ntivirals				
For	eye preparations refer to Eye Preparations, Anti-Infective Prep	parations, page 241			
Н	epatitis B Treatment				
	TECAVIR				
	Tab 0.5 mg		30	✓ E	ntecavir Sandoz
LA	MIVUDINE - Special Authority see SA1685 below - Retail pha				
	Tab 100 mg Oral liq 5 mg per ml		28 10 ml Ol		<u>etlam</u> effix
	SA1685 Special Authority for Subsidy	270.00 25	+0 IIII OI	• 2	CIIIA
Init App Re	ial application only from a relevant specialist or medical practi provals valid for 1 year where used for the treatment or prevent newal from any relevant practitioner. Approvals valid for 2 yea	ion of hepatitis B.			
TE	NOFOVIR DISOPROXIL	-t	⊷! امضامنا	46	of our to A collectation of
	Tenofovir disoproxil prescribed under endorsement for the treantiretrovirals for the purposes of Special Authority SA2139.,		iuaea in	tne count	of up to 4 subsidised
*	Tab 245 mg (300 mg as a maleate)		30	✓ T	enofovir Disoproxil Mylan
	Tenofovir Disoproxil Mylan to be Principal Supply on 1 De	ecember 2022			•
*	Tab 245 mg (300.6 mg as a succinate)	38.10	30	✓ T	enofovir Disoproxil

Herpesvirus Treatments

AC	ICLOVIR		
*	Tab dispersible 200 mg1.78	25	✓ Lovir
	Tab dispersible 400 mg5.38	56	✓ Lovir
	Tab dispersible 800 mg5.98	35	✓ Lovir

(Tenofovir Disoproxil Teva Tab 245 mg (300.6 mg as a succinate) to be delisted 1 December 2022)

Teva

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
VALACICLOVIR Tab 500 mg Tab 1,000 mg		30 30		aclovir aclovir
VALGANCICLOVIR - Special Authority see SA1993 below - Ret Tab 450 mg		60	✓ <u>v</u>	alganciclovir Mylan

⇒SA1993 Special Authority for Subsidy

Initial application — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months where the patient has undergone a solid organ transplant and requires valganciclovir for CMV prophylaxis.

Renewal — (transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has undergone a solid organ transplant and received anti-thymocyte globulin and requires valganciclovir therapy for CMV prophylaxis; and
 - 1.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following anti-thymocyte globulin; or
- 2 Both:
 - 2.1 Patient has received pulse methylprednisolone for acute rejection and requires further valganciclovir therapy for CMV prophylaxis; and
 - 2.2 Patient is to receive a maximum of 90 days of valganciclovir prophylaxis following pulse methylprednisolone.

Initial application — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 Patient has undergone a solid organ transplant and received valganciclovir under Special Authority more than 2 years ago (27 months): and
- 2 Patient has received anti-thymocyte globulin and requires valganciclovir for CMV prophylaxis.

Renewal — (cytomegalovirus prophylaxis following anti-thymocyte globulin) only from a relevant specialist. Approvals valid for 3 months where the patient has received a further course of anti-thymocyte globulin and requires valganciclovir for CMV

Initial application — (Lung transplant cytomegalovirus prophylaxis) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undergone a lung transplant; and
- 2 Either:
 - 2.1 The donor was cytomegalovirus positive and the patient is cytomegalovirus negative; or
 - 2.2 The recipient is cytomegalovirus positive: and
- 3 Patient has a high risk of CMV disease.

Initial application — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 Patient is immunocompromised; and
- 2 Any of the following:
 - 2.1 Patient has cytomegalovirus syndrome or tissue invasive disease; or
 - 2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or
 - 2.3 Patient has cytomegalovirus retinitis.

Renewal — (Cytomegalovirus in immunocompromised patients) only from a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria: Both:

(Manufacturer's Price) Subsidised Generic	,		201010		
	s Per ✓ Manufacturer ontinued 1 Patient is immunocompromised; and		Subsidy	Fully	Brand or
\$ Per ✓ Manufacturer	ontinued 1 Patient is immunocompromised; and		(Manutacturer's Price)	Subsidised	
ψ 1 Cl • Widification	1 Patient is immunocompromised; and		\$	Per 🗸	Manufacturer
	2 Any of the following:	 Patient is immunocompromised; and 			
1 Patient is immunocompromised; and		2 Any of the following:			
· · · · · · · · · · · · · · · · · · ·		0.0 0.0 1.1 1.1 1.1 1.1 0.00.0	ALA to alcasona at discussion		

2.2 Patient has rapidly rising plasma CMV DNA in absence of disease; or

2.3 Patient has cytomegalovirus retinitis.

Note: for the purpose of this Special Authority "immunocompromised" includes transplant recipients, patients with immunosuppressive diseases (e.g. HIV) or those receiving immunosuppressive treatment for other conditions.

Hepatitis C Treatment

GLECAPREVIR WITH PIBRENTASVIR - [Xpharm]

Note the supply of treatment is via Pharmac's approved direct distribution supply. Further details can be found on Pharmac's website https://pharmac.govt.nz/maviret

Tab 100 mg with pibrentasvir 40 mg24,750.00 ✓ Maviret 84 OP

LEDIPASVIR WITH SOFOSBUVIR - [Xpharm] - Special Authority see SA1605 below

No patient co-payment payable

Tab 90 mg with sofosbuvir 400 mg......24,363.46 ✓ Harvoni

⇒SA1605 Special Authority for Subsidy

Special Authority approved by the Hepatitis C Treatment Panel (HepCTP)

Notes: By application to the Hepatitis C Treatment Panel (HepCTP).

Applications will be considered by HepCTP and approved subject to confirmation of eligibility.

Application details may be obtained from Pharmac's website http://www.pharmac.govt.nz/maviret or:

The Coordinator, Hepatitis C Treatment Panel

Pharmac, PO Box 10-254, WELLINGTON Tel: (04) 460 4990.

Email: hepcpanel@pharmac.govt.nz

HIV Prophylaxis and Treatment

EMTRICITABINE WITH TENOFOVIR DISOPROXIL - Subsidy by endorsement; can be waived by Special Authority see SA2138 on the next page

Endorsement for treatment of HIV: Prescription is deemed to be endorsed if emtricitabine with tenofovir disoproxil is co-prescribed with another antiretroviral subsidised under Special Authority SA2139 and the prescription is annotated accordingly by the Pharmacist or endorsed by the prescriber.

Note: Emtricitabine with tenofovir disoproxil prescribed under endorsement for the treatment of HIV is included in the count of up to 4 subsidised antiretrovirals, and counts as two antiretroviral medications, for the purposes of Special Authority SA2139, page 104 There is an approval process to become a named specialist to prescribe antiretroviral therapy in New Zealand. Further information is available on the Pharmac website.

*	Tab 200 mg with tenofovir disoproxil 245 mg (300 mg as a maleate)	15.45	30	✓ Tenofovir Disoproxil Emtricitabine Mylan
	Tenofovir Disoproxil Emtricitabine Mylan to be Principal Su	ipply on 1 Dece	ember 2022	
*	Tab 200 mg with tenofovir disoproxil 245 mg (300.6 mg as a			
	succinate)	61.15	30	✓ Teva
/Te	eva Tah 200 mg with tenofovir disoprovil 245 mg (300 6 mg as a	succinate) to h	e delisted 1	December 2022)

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2138 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

Renewal from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria: Both:

- 1 Patient has tested HIV negative, does not have signs or symptoms of acute HIV infection and has been assessed for HIV seroconversion; and
- 2 The Practitioner considers the patient is at elevated risk of HIV exposure and use of PrEP is clinically appropriate.

Notes: Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines:

https://ashm.org.au/HIV/PrEP/

COVID-19 Treatments

MOLNUPIRAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

NIRMATRELVIR WITH RITONAVIR - [Xpharm] - Subsidy by endorsement

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for oral antiviral COVID-19 treatments (as on <u>Pharmac's website</u>) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Antiretrovirals

⇒SA2139 Special Authority for Subsidy

Initial application — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the patient has confirmed HIV infection.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Confirmed HIV) only from a named specialist. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Prevention of maternal transmission) only from a named specialist. Approvals valid for 1 year for applications meeting the following criteria:

Either:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Prevention of maternal foetal transmission; or
- 2 Treatment of the newborn for up to eight weeks.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Some antiretrovirals are unapproved or contraindicated for this indication. Practitioners prescribing these medications should exercise their own skill, judgement, expertise and discretion, and make their own prescribing decisions with respect to the use of a Pharmaceutical for an indication for which it is not approved or contraindicated.

Initial application — (post-exposure prophylaxis following exposure to HIV) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Refer to local health pathways or the Australasian Society for HIV, Viral Hepatitis and Sexual Health Medicine clinical guidelines for PEP (https://www.ashm.org.au/hiv/hiv-management/pep/).

Renewal — (second or subsequent post-exposure prophylaxis) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

Both:

- 1 Treatment course to be initiated within 72 hours post exposure; and
- 2 Any of the following:
 - 2.1 Patient has had condomless anal intercourse or receptive vaginal intercourse with a known HIV positive person with an unknown or detectable viral load greater than 200 copies per ml; or
 - 2.2 Patient has shared intravenous injecting equipment with a known HIV positive person; or
 - 2.3 Patient has had non-consensual intercourse and the clinician considers that the risk assessment indicates prophylaxis is required; or
 - 2.4 Patient has had condomless anal intercourse with a person from a high HIV prevalence country or risk group whose HIV status is unknown.

Initial application — (Percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

Notes: Tenofovir disoproxil prescribed under endorsement for HIV is included in the count of up to 4 subsidised antiretrovirals. Subsidies apply for a combination of up to four antiretroviral medications. The combination of a protease inhibitor and low-dose ritonavir given as a booster (either as part of a combination product or separately) will be counted as one protease inhibitor for the purpose of accessing funding to antiretrovirals.

Renewal — (Second or subsequent percutaneous exposure) only from a named specialist. Approvals valid for 6 weeks where the patient has percutaneous exposure to blood known to be HIV positive.

	Subsidy (Manufacturer's P \$	rice) Subs Per	Fully idised	Brand or Generic Manufacturer
Non-nucleosides Reverse Transcriptase Inhib	itors			
EFAVIRENZ – Special Authority see SA2139 on page 104 – F Tab 200 mg Tab 600 mg	190.15	90 30	_	tocrin tocrin
ETRAVIRINE – Special Authority see SA2139 on page 104 – Tab 200 mg		60	✓ In	telence
NEVIRAPINE - Special Authority see SA2139 on page 104 - Tab 200 mg		60		evirapine Alphapharm
Oral suspension 10 mg per ml	203.55	240 ml OP	✓ Vi	ramune Suspension
Nucleosides Reverse Transcriptase Inhibitors	•			
ABACAVIR SULPHATE – Special Authority see SA2139 on p Tab 300 mg Oral liq 20 mg per ml ABACAVIR SULPHATE WITH LAMIVUDINE – Special Author Note: abacavir with lamivudine (combination tablets) cour		60 240 ml OP page 104 – Re	✓ Zi tail pha	
anti-retroviral Special Authority. Tab 600 mg with lamivudine 300 mg		30		ivexa
pharmacy Note: Efavirenz with emtricitabine and tenofovir disoproxi anti-retroviral Special Authority Tab 600 mg with emtricitabine 200 mg and tenofovir disoproxi 245 mg (300 mg as a maleate)	counts as three an			s for the purposes of the
EMTRICITABINE - Special Authority see SA2139 on page 10 Cap 200 mg		y 30	✓ Eı	mtriva
LAMIVUDINE - Special Authority see SA2139 on page 104 - Tab 150 mg		60		a <u>mivudine</u> Alphapharm
Oral liq 10 mg per ml ZIDOVUDINE [AZT] – Special Authority see SA2139 on page	104 – Retail pharm	240 ml OP acy	√ 31	rc
Cap 100 mg Oral liq 10 mg per ml	30.45	100 200 ml OP	✓ R	etrovir etrovir
ZIDOVUDINE [AZT] WITH LAMIVUDINE – Special Authority: Note: zidovudine [AZT] with lamivudine (combination table the anti-retroviral Special Authority.				
Tab 300 mg with lamivudine 150 mg	33.00	60	✓ A	lphapharm
Protease Inhibitors				
ATAZANAVIR SULPHATE – Special Authority see SA2139 or Cap 150 mg	141.68 188.91	pharmacy 60 60	✓ Te	
Tab 400 mg	132.00	60 60		arunavir Mylan arunavir Mylan
✓ fully subsidised	S29 Unapp	roved medicine s	upplied i	under Section 29

	Subsidy		Fully	Brand or
(Manu	ufacturer ['] s Price	e) Subs	sidised	Generic
	\$	Per	✓	Manufacturer
LOPINAVIR WITH RITONAVIR - Special Authority see SA2139 on pa	ge 104 – Reta	ail pharmacy	/	
Tab 100 mg with ritonavir 25 mg	150.00	60	✓ <u>L</u>	opinavir/Ritonavir Mylan
Tab 200 mg with ritonavir 50 mg	295.00	120	✓ <u>L</u>	opinavir/Ritonavir Mylan
Oral liq 80 mg with ritonavir 20 mg per ml	735.00	300 ml OP	✓ K	Caletra
RITONAVIR – Special Authority see SA2139 on page 104 – Retail pha Tab 100 mg	•	30	✓ N	lorvir
Strand Transfer Inhibitors				
DOLUTEGRAVIR - Special Authority see SA2139 on page 104 - Reta Tab 50 mg1,1		30	✓ T	ïvicay
RALTEGRAVIR POTASSIUM - Special Authority see SA2139 on page	e 104 – Retai	l pharmacv		-
Tab 400 mg		60	√ s	sentress
Tab 600 mg		60		sentress HD

Immune Modulators

PEGYLATED INTERFERON ALFA-2A - Special Authority see SA2034 below - Retail pharmacy

⇒SA2034 Special Authority for Subsidy

Initial application — (chronic hepatitis C - genotype 1, 4, 5 or 6 infection or co-infection with HIV or genotype 2 or 3 post liver transplant) from any specialist. Approvals valid for 18 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 Patient has chronic hepatitis C, genotype 1, 4, 5 or 6 infection; or
 - 1.2 Patient has chronic hepatitis C and is co-infected with HIV; or
 - 1.3 Patient has chronic hepatitis C genotype 2 or 3 and has received a liver transplant; and
- 2 Maximum of 48 weeks therapy.

Renewal — (Chronic hepatitis C - genotype 1 infection) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and
- 3 Either:
 - 3.1 Patient has responder relapsed: or
 - 3.2 Patient was a partial responder; and
- 4 Patient is to be treated in combination with boceprevir; and
- 5 Maximum of 48 weeks therapy.

Initial application — (Chronic Hepatitis C - genotype 1 infection treatment more than 4 years prior) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic hepatitis C, genotype 1; and
- 2 Patient has had previous treatment with pegylated interferon and ribavirin; and

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	-
(Wallulaciulei S Filce)	Per	Jubsidised	Manufacturer	
φ	rei	•	Manuaciurei	

continued...

- 3 Any of the following:
 - 3.1 Patient has responder relapsed; or
 - 3.2 Patient was a partial responder; or
 - 3.3 Patient received interferon treatment prior to 2004; and
 - 4 Patient is to be treated in combination with boceprevir; and
 - 5 Maximum of 48 weeks therapy.

Initial application — (chronic hepatitis C - genotype 2 or 3 infection without co-infection with HIV) from any specialist. Approvals valid for 12 months for applications meeting the following criteria:

- Both:
 - 1 Patient has chronic hepatitis C, genotype 2 or 3 infection; and
 - 2 Maximum of 6 months therapy.

Initial application — (Hepatitis B) only from a gastroenterologist, infectious disease specialist or general physician. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed Hepatitis B infection (HBsAg positive for more than 6 months); and
- 2 Patient is Hepatitis B treatment-naive; and
- 3 ALT > 2 times Upper Limit of Normal: and
- 4 HBV DNA < 10 log10 IU/ml; and
- 5 Fither:
 - 5.1 HBeAg positive; or
 - 5.2 serum HBV DNA greater than or equal to 2.000 units/ml and significant fibrosis (Metavir Stage F2 or greater or moderate fibrosis); and
- 6 Compensated liver disease; and
- 7 No continuing alcohol abuse or intravenous drug use; and
- 8 Not co-infected with HCV, HIV or HDV; and
- 9 Neither ALT nor AST > 10 times upper limit of normal; and
- 10 No history of hypersensitivity or contraindications to pegylated interferon; and
- 11 Maximum of 48 weeks therapy.

Initial application — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 Patient has a cutaneous T cell lymphoma*; or
- 2 All of the following:
 - 2.1 Patient has a myeloproliferative disorder*; and
 - 2.2 Patient is intolerant of hydroxyurea; and
 - 2.3 Treatment with an agrelide and busulfan is not clinically appropriate; or
- 3 Both:
 - 3.1 Patient has a myeloproliferative disorder; and
 - 3.2 Patient is pregnant, planning pregnancy or lactating.

Renewal — (myeloproliferative disorder or cutaneous T cell lymphoma) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment; and
- 3 Either:
 - 3.1 Patient has a cutaneous T cell lymphoma*; or
 - 3.2 Both:

INFECTIONS - AGENTS FOR SYSTEMIC USE

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 3.2.1 Patient has a myeloproliferative disorder*; and
- 3.2.2 Either:
 - 3.2.2.1 Remains intolerant of hydroxyurea and treatment with anagrelide and busulfan remains clinically inappropriate; or
 - 3.2.2.2 Patient is pregnant, planning pregnancy or lactating.

Note: Indications marked with * are unapproved indications.

Initial application — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient has received an allogeneic bone marrow transplant* and has evidence of disease relapse.

Renewal — (post-allogenic bone marrow transplant) from any relevant practitioner. Approvals valid for 3 months where patient is responding and ongoing treatment remains appropriate.

Note: Indications marked with * are unapproved indications.

Urinary Tract Infections

METHENAMINE (HEXAMINE) HIPPURATE			
* Tab 1 g	19.95	100	✓ Hiprex
NITROFURANTOIN			
* Tab 50 mg - Up to 30 tab available on a PSO	22.20	100	✓ Nifuran
Nifuran to be Principal Supply on 1 December 2022			
* Tab 100 mg	37.50	100	✓ Nifuran
Nifuran to be Principal Supply on 1 December 2022			
* Cap modified-release 100 mg - Up to 15 cap available on a			
PSO	86.40	100	✓ <u>Macrobid</u>
NORFLOXACIN			
Tab 400 mg - Subsidy by endorsement	245.00	100	Arrow-Norfloxacin

Only if prescribed for a patient with an uncomplicated urinary tract infection that is unresponsive to a first line agent or with proven resistance to first line agents and the prescription is endorsed accordingly.

	0.4.44.		E. II.	December 1
	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	Jubsidised	Manufacturer
Anticholinesterases				
EOSTIGMINE METILSULFATE				
Inj 2.5 mg per ml, 1 ml ampoule	33.81	10	/	Max Health
/RIDOSTIGMINE BROMIDE				
Tab 60 mg	45.79	100	✓	Mestinon
Non-Steroidal Anti-Inflammatory Drugs				
CLOFENAC SODIUM	4.00		,	D
Tab EC 25 mg		50		Diclofenac Sandoz
Tab 50 mg dispersible		20		Voltaren D
Tab EC 50 mg		50		Diclofenac Sandoz
Tab long-acting 75 mg		100		Voltaren SR
Inj 25 mg per ml, 3 ml ampoule – Up to 5 inj available on a	PSO 13.20	5		Voltaren
Suppos 12.5 mg	2.04	10	•	Voltaren
Suppos 25 mg	2.44	10	✓	Voltaren
Suppos 50 mg - Up to 10 supp available on a PSO	4.22	10	✓	Voltaren
Suppos 100 mg	7.00	10	✓	Voltaren
UPROFEN				
Tab 200 mg	21.40	1.000	•	Relieve
Tab long-acting 800 mg		30		Brufen SR
Oral lig 20 mg per ml		200 m		Ethics
Oral liq 20 mg per mi	11.29	200 11		Fenpaed 100 mg per
	11.29		•	5 ml
ETOPROFEN				
Cap long-acting 200 mg	12.07	28	✓	Oruvail SR
EFENAMIC ACID				
Cap 250 mg	1 25	50		
Oup 200 mg	(9.16)	00		Ponstan
	0.50	20		i Olistali
	(7.50)	20		Ponstan
	(7.50)			FUISIAII
APROXEN			_	
Tab 250 mg		500		Noflam 250
Tab 500 mg		250		Noflam 500
Tab long-acting 750 mg	6.47	28	/	Naprosyn SR 750
Tab long-acting 1 g	8.62	28	✓	Naprosyn SR 1000
ENOXICAM				
Tab 20 mg	18.50	100	/	Tilcotil
Inj 20 mg vial		1		AFT
11 20 11g Vid		'	•	Al I
NSAIDs Other				
ELECOXIB				
Cap 100 mg	3.45	60	✓	Celebrex
			1	Celecoxib Pfizer
Cap 200 mg	3.20	30	1	Celebrex

Subsidy
(Manufacturer's Price) Subsidy

\$ Per

Fully Subsidised Brand or Generic Manufacturer

Topical Products for Joint and Muscular Pain

CAPSAICIN

Crm 0.025% - Special Authority see SA1289 below - Retail

⇒SA1289 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified where the patient has osteoarthritis that is not responsive to paracetamol and oral non-steroidal anti-inflammatories are contraindicated.

Antirheumatoid Agents

HYDROXYCHLOROQUINE - Subsidy by endorsement

Subsidised only if prescribed for rheumatoid arthritis, systemic or discoid lupus erythematosus, malaria treatment or suppression, relevant dermatological conditions (cutaneous forms of lupus and lichen planus, cutaneous vasculitides and mucosal ulceration)*, sarcoidosis (pulmonary and non-pulmonary)*, and the prescription is endorsed accordingly.

Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of hydroxychloroquine. Note: Indication marked with a * is an unapproved indication.

* Tab 200 mg	8.78	100	Plaquenil
LEFLUNOMIDE			
Tab 10 mg	6.00	30	✓ Arava
Tab 20 mg		30	✓ Arava
PENICILLAMINE			
Tab 125 mg	67.23	100	✓ D-Penamine
Tab 250 mg	110.12	100	✓ D-Penamine

Drugs Affecting Bone Metabolism

Alendronate for Osteoporosis

ALENDRONATE SODIUM			
* Tab 70 mg	2.44	4	✓ Fosamax
ALENDRONATE SODIUM WITH COLECAL CIFEROL			

Other Treatments

AL ENDOONATE CODULA

⇒SA1777 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 Fither
 - 2.1 The patient is female and postmenopausal; or
 - 2.2 The patient is male or non-binary; and
- 3 Any of the following:
 - 3.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density

Subsidy		Fully	Brand or
(Manufacturer's Price)	Subs	sidised	Generic
\$	Per	✓	Manufacturer

continued...

- (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
- 3.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons; or
- 3.3 History of two significant osteoporotic fractures demonstrated radiologically; or
- 3.4 Documented T-Score less than or equal to -3.0 (see Note); or
- 3.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
- 3.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 4 Zoledronic acid is contraindicated because the patient's creatinine clearance is less than 35 mL/min; and
- 5 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes); and
- 6 The patient must not receive concomitant treatment with any other funded antiresorptive agent for this condition or teriparatide.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with denosumab
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body
- e) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: risedronate sodium tab 35 mg once weekly; alendronate sodium tab 70 mg or tab 70 mg with cholecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy

PAMIDRONATE DISODIUM

inj 3 mg per mi, 10 mi viai	27.53	1	Pamisoi
Inj 6 mg per ml, 10 ml vial	74.67	1	✓ Pamisol
Inj 9 mg per ml, 10 ml vial	79.95	1	✓ Pamisol
RALOXIFENE HYDROCHLORIDE - Special Authority see SA177	79 below – Retail	pharmacy	
* Tab 60 mg	53.76	28	Evista

⇒SA1779 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Notes); or

Subsic (Manufacture	. ,	,	nd or neric
\$	Per	✓ Mar	nufacturer

continued...

- 2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
- 3 History of two significant osteoporotic fractures demonstrated radiologically: or
- 4 Documented T-Score less than or equal to -3.0 (see Notes); or
- 5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Notes); or
- 6 Patient has had a Special Authority approval for zoledronic acid (Underlying cause Osteoporosis) or has had a Special Authority approval for alendronate (Underlying cause - Osteoporosis) prior to 1 February 2019.

Notes:

- a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).
 Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.
- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for raloxifene funding.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis, and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

⇒SA1139 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 18 months for applications meeting the following criteria: All of the following:

- 1 The patient has severe, established osteoporosis; and
- 2 The patient has a documented T-score less than or equal to -3.0 (see Notes); and
- 3 The patient has had two or more fractures due to minimal trauma; and
- 4 The patient has experienced at least one symptomatic new fracture after at least 12 months' continuous therapy with a funded antiresorptive agent at adequate doses (see Notes).

Notes:

- a) The bone mineral density (BMD) measurement used to derive the T-score must be made using dual-energy x-ray absorptiometry (DXA). Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable
- b) Antiresorptive agents and their adequate doses for the purposes of this Special Authority are defined as: alendronate sodium tab 70 mg or tab 70 mg with colecalciferol 5,600 iu once weekly; raloxifene hydrochloride tab 60 mg once daily; zoledronic acid 5 mg per year. If an intolerance of a severity necessitating permanent treatment withdrawal develops during the use of one antiresorptive agent, an alternate antiresorptive agent must be trialled so that the patient achieves the minimum requirement of 12 months' continuous therapy.
- c) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.
- d) A maximum of 18 months of treatment (18 cartridges) will be subsidised.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

70I FDRONIC ACID

Inj 0.05 mg per ml, 100 ml, vial - Special Authority see

SA2110 below – Retail pharmacy60.00 100 ml OP 🗸 Aclasta

⇒SA2110 Special Authority for Subsidy

Initial application — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Paget's disease; and
- 2 Any of the following:
 - 2.1 Bone or articular pain; or
 - 2.2 Bone deformity; or
 - 2.3 Bone, articular or neurological complications; or
 - 2.4 Asymptomatic disease, but risk of complications; or
 - 2.5 Preparation for orthopaedic surgery; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Initial application — (Underlying cause - Osteoporosis) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note); or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 Patient has had a Special Authority approval for alendronate (Underlying cause Osteoporosis) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (Underlying cause - glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is receiving systemic glucocorticosteroid therapy (greater than or equal to 5 mg per day prednisone equivalents) and has already received or is expected to receive therapy for at least three months; and
- 2 Any of the following:
 - 2.1 The patient has documented BMD greater than or equal to 1.5 standard deviations below the mean normal value in voung adults (i.e. T-Score less than or equal to -1.5) (see Note): or
 - 2.2 The patient has a history of one significant osteoporotic fracture demonstrated radiologically; or
 - 2.3 The patient has had a Special Authority approval for alendronate (Underlying cause glucocorticosteroid therapy) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

Renewal — (Paget's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

1 Any of the following:

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 1.1 The patient has relapsed (based on increases in serum alkaline phosphatase); or
- 1.2 The patient's serum alkaline phosphatase has not normalised following previous treatment with zoledronic acid; or
- 1.3 Symptomatic disease (prescriber determined); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was, and remains, glucocorticosteroid therapy) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The patient is continuing systemic glucocorticosteriod therapy (greater than or equal to 5 mg per day prednisone equivalents); and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in the 12-month approval period.

The patient must not have had more than 1 prior approval in the last 12 months.

Renewal — (Underlying cause was glucocorticosteroid therapy but patient now meets the 'Underlying cause - osteoporosis' criteria) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 History of one significant osteoporotic fracture demonstrated radiologically and documented bone mineral density (BMD) greater than or equal to 2.5 standard deviations below the mean normal value in young adults (i.e. T-Score less than or equal to -2.5) (see Note): or
 - 1.2 History of one significant osteoporotic fracture demonstrated radiologically, and either the patient is elderly, or densitometry scanning cannot be performed because of major logistical, technical or pathophysiological reasons. It is unlikely that this provision would apply to many patients under 75 years of age; or
 - 1.3 History of two significant osteoporotic fractures demonstrated radiologically; or
 - 1.4 Documented T-Score less than or equal to -3.0 (see Note); or
 - 1.5 A 10-year risk of hip fracture greater than or equal to 3%, calculated using a published risk assessment algorithm (e.g. FRAX or Garvan) which incorporates BMD measurements (see Note); or
 - 1.6 The patient has had a Special Authority approval for alendronate (Underlying was glucocorticosteroid therapy but patient now meets the 'Underlying cause Osteoporosis' criteria) prior to 1 February 2019 or has had a Special Authority approval for raloxifene; and
- 2 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Initial application — (spinal cord injury*) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Patient has experienced an acute traumatic spinal cord injury in the last six months; and
- 2 Patient is being managed by a specialist spinal acute care and rehabilitation unit; and
- 3 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period.

Note: Indications marked with * are unapproved indications.

Renewal — (spinal cord injury*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient will not be prescribed more than 5 mg of zoledronic acid in a 12-month period; and
- 2 The patient has not received more than two doses of zoledronic acid for this indication.

The patient must not have had more than 1 prior approval.

Notes: No further renewals will be subsidised. A maximum of 2 vials of zoledronic acid treatment for spinal cord injury will be subsidised. Indications marked with * are unapproved indications.

a) BMD (including BMD used to derive T-Score) must be measured using dual-energy x-ray absorptiometry (DXA).

Subsidy (Manufacturer's Price)	,	Fully Subsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Quantitative ultrasound and quantitative computed tomography (QCT) are not acceptable.

- b) Evidence suggests that patients aged 75 years and over who have a history of significant osteoporotic fracture demonstrated radiologically are very likely to have a T-Score less than or equal to -2.5 and, therefore, do not require BMD measurement for treatment with bisphosphonates.
- c) Osteoporotic fractures are the incident events for severe (established) osteoporosis and can be defined using the WHO definitions of osteoporosis and fragility fracture. The WHO defines severe (established) osteoporosis as a T-score below -2.5 with one or more associated fragility fractures. Fragility fractures are fractures that occur as a result of mechanical forces that would not ordinarily cause fracture (minimal trauma). The WHO has quantified this as forces equivalent to a fall from a standing height or less.
- d) A vertebral fracture is defined as a 20% or greater reduction in height of the anterior or mid portion of a vertebral body relative to the posterior height of that body, or a 20% or greater reduction in any of these heights compared to the vertebral body above or below the affected vertebral body.

Hyperuricaemia and Antigout

ALLOPURINOL			
* Tab 100 mg	11.47	500	✓ DP-Allopurinol
* Tab 300 mg		500	✓ DP-Allopurinol
BENZBROMARONE - Special Authority see SA196	3 below - Retail pharmacy		
Tab 50 mg	22.50	100	✓ Narcaricin mite \$29
Tab 100 mg	13.50	30	✓ Desuric S29
			✓ Urinorm S29
	45.00	100	✓ Benzbromaron AL
			100 S29

⇒SA1963 Special Authority for Subsidy

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate and the patient is benefitting from the treatment; and
- 2 There is no evidence of liver toxicity and patient is continuing to receive regular (at least every three months) liver function tests.

COLCHICINE

* Tab 500 mcg	6.00	100	✓ Colgout
FEBUXOSTAT - Special Authority see SA2054 below - Retail	pharmacy		
Tab 80 mg	20.00	28	✓ Febuxostat
			multichem
Tab 120 mg	20.00	28	✓ Febuxostat
			multichem

⇒SA2054 Special Authority for Subsidy

Initial application — (Gout) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has been diagnosed with gout; and
- 2 Any of the following:
 - 2.1 The patient has a serum urate level greater than 0.36 mmol/l despite treatment with allopurinol at doses of at least 600 mg/day and addition of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
 - 2.2 The patient has experienced intolerable side effects from allopurinol such that treatment discontinuation is required

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
\$	Per	✓	Manufacturer

continued...

- and serum urate remains greater than 0.36 mmol/l despite use of probenecid at doses of up to 2 g per day or maximum tolerated dose; or
- 2.3 The patient has renal impairment such that probenecid is contraindicated or likely to be ineffective and serum urate remains greater than 0.36 mmol/l despite optimal treatment with allopurinol (see Note); or
- 2.4 The patient has previously had an initial Special Authority approval for benzbromarone for treatment of gout...

Initial application — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient is scheduled to receive cancer therapy carrying an intermediate or high risk of tumour lysis syndrome; and
- 2 Patient has a documented history of allopurinol intolerance.

Renewal — (Gout) from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (Tumour lysis syndrome) only from a haematologist or oncologist. Approvals valid for 6 weeks where the treatment remains appropriate and the patient is benefitting from treatment.

PROBENECID

BACLOFEN

*	Tab 500 mg	66.95	100	✓ [Probenecid-AFT
---	------------	-------	-----	------------	----------------

Muscle Relaxants

	· =· ·			
∗ Ta	b 10 mg	4.20	100	✓ Pacifen
Inj	0.05 mg per ml, 1 ml ampoule - Subsidy by endorsement	11.55	1	✓ Lioresal Intrathecal
•	Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo	s where oral an		ents have been ineffective or have
Inj	2 mg per ml, 5 ml ampoule - Subsidy by endorsement	306.82	5	✓ Medsurge
•	Subsidised only for use in a programmable pump in patient caused intolerable side effects and the prescription is endo			ents have been ineffective or have
DANTE	ROLENE			
Ca	p 25 mg	97.50	100	✓ Dantrium
				✓ Dantrium S29 S29
Ca	p 50 mg	77.00	100	✓ Dantrium
ORPHI	ENADRINE CITRATE			
Ta	b 100 mg	20.76	100	✓ <u>Norflex</u>

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Agents for Parkinsonism and Related Disorders

Dopamine Agonists a	nd Related Agents
----------------------------	-------------------

AMANTADINE HYDROCHLORIDE			
▲ Cap 100 mg		60	✓ Symmetrel
	63.73	100	✓ Symmetrel S29 S29
APOMORPHINE HYDROCHLORIDE			
▲ Inj 10 mg per ml, 2 ml ampoule	59.50	5	✓ Movapo
▲ Inj 10 mg per ml, 5 ml ampoule	121.84	5	✓ Movapo
ENTACAPONE			
▲ Tab 200 mg	18.04	100	✓ Comtan
LEVODOPA WITH BENSERAZIDE			
* Tab dispersible 50 mg with benserazide 12.5 mg	13.25	100	✓ Madopar Rapid
* Cap 50 mg with benserazide 12.5 mg		100	✓ Madopar 62.5
* Cap 100 mg with benserazide 25 mg	15.80	100	✓ Madopar 125
* Cap long-acting 100 mg with benserazide 25 mg	22.85	100	✓ Madopar HBS
* Cap 200 mg with benserazide 50 mg	26.25	100	Madopar 250
LEVODOPA WITH CARBIDOPA			
* Tab 100 mg with carbidopa 25 mg	21.11	100	✓ Sinemet
* Tab long-acting 200 mg with carbidopa 50 mg	43.65	100	✓ Sinemet CR
* Tab 250 mg with carbidopa 25 mg	38.39	100	✓ Sinemet
PRAMIPEXOLE HYDROCHLORIDE			
▲ Tab 0.25 mg	5.51	100	✓ Ramipex
Ramipex to be Principal Supply on 1 December 2022			
▲ Tab 1 mg	18.66	100	✓ Ramipex
Ramipex to be Principal Supply on 1 December 2022			
RASAGILINE			
* Tab 1 mg	53.50	30	✓ Azilect S29
ROPINIROLE HYDROCHLORIDE			
▲ Tab 0.25 mg	3.39	100	✓ Mylan S29
· · · · · · · · · ·	4.05	84	✓ Ropin
▲ Tab 1 mg	4.70	100	✓ Mylan S29
•	4.95	84	✓ Ropin
▲ Tab 2 mg	6.48	84	✓ Ropin
▲ Tab 5 mg	14.50	84	✓ Ropin
(Mylan \$29 Tab 0.25 mg to be delisted 1 January 2023)			

(Mylan S29 Tab 0.25 mg to be delisted 1 January 2023)

(Mylan S29) Tab 1 mg to be delisted 1 December 2022)

SELEGILINE HYDROCHLORIDE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking selegiline hydrochloride prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of selegiline hydrochloride.

*	Tab 5 mg48.00	100	✓ Eldepryl S29
TC	DLCAPONE		
\blacktriangle	Tab 100 mg	100	✓ Tasmar

Anticholinergics BENZATROPINE MESYLATE Tab 2 mg		Per 60 5	Fully bsidised	Generic Manufacturer
BENZATROPINE MESYLATE Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 10 inj available on a PSO				enztron
Tab 2 mg Inj 1 mg per ml, 2 ml a) Up to 10 inj available on a PSO				enztron
, .			₹ <u>P</u>	<u>hebra</u>
PROCYCLIDINE HYDROCHLORIDE Tab 5 mg	7.40	100	✓ K	emadrin
Agents for Essential Tremor, Chorea and Related	d Disorders			
RILUZOLE – Special Authority see SA1403 below – Retail pharm Wastage claimable Tab 50 mg	•	56	√ <u>R</u>	ilutek
■ SA1403 Special Authority for Subsidy Initial application only from a neurologist or respiratory specialist following criteria: All of the following: 1 The patient has amyotrophic lateral sclerosis with disease 2. The patient has at least 60 percent of predicted forced vital 3. The patient has not undergone a tracheostomy; and 4. The patient has not experienced respiratory failure; and 5. Any of the following: 5.1 The patient is ambulatory; or 5.2 The patient is able to use upper limbs; or 5.3 The patient is able to swallow. Renewal from any relevant practitioner. Approvals valid for 18 mc All of the following: 1 The patient has not undergone a tracheostomy; and 2. The patient has not experienced respiratory failure; and 3. Any of the following: 3.1 The patient is ambulatory; or 3.2 The patient is able to use upper limbs; or 3.3 The patient is able to swallow. TETRABENAZINE	duration of 5 years o	or less; a onths pr	nd ior to the	initial application; and
Tab 25 mg	91.10	112	✓ M	otetis
Anaesthetics				
Local				
LIDOCAINE [LIGNOCAINE] Gel 2%, tube — Subsidy by endorsement	dministration and the	10	otion is er	stillagel Lido

accordingly.

	Subsidy (Manufacturer's Price		Fully	
	\$	Per	JISEU ✓	Manufacturer
LIDOCAINE [LIGNOCAINE] HYDROCHLORIDE				
Oral (gel) soln 2%	38.00	200 ml	1	Mucosoothe
Inj 1%, 5 ml ampoule - Up to 25 inj available on a PSO	9.50	25	✓	Lidocaine-Baxter
	17.50	50		
	(35.00)			Xylocaine
Inj 2%, 5 ml ampoule - Up to 5 inj available on a PSO	8.25	25	✓	Lidocaine-Baxter
Inj 1%, 20 ml ampoule - Up to 5 inj available on a PSO	12.00	5		
	(20.00)			Xylocaine
Inj 1%, 20 ml vial - Up to 5 inj available on a PSO	6.20	5	✓	Lidocaine-Baxter
			✓	Lidocaine-Claris
Inj 2%, 20 ml vial - Up to 5 inj available on a PSO	6.45	5	✓	Lidocaine-Baxter
LIDOCAINE [LIGNOCAINE] WITH CHLORHEXIDINE				
Gel 2% with chlorhexidine 0.05%, 10 ml urethral syringes -				
Subsidy by endorsement	103.32	10	1	Pfizer
a) Up to 5 each available on a PSO				
b) Subsidised only if prescribed for urethral or cervical a	administration and th	e prescriptio	n is	endorsed accordingly.

Topical Local Anaesthetics

⇒SA0906 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years where the patient is a child with a chronic medical condition requiring frequent injections or venepuncture.

Renewal from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

LIDOCAINE [LIGNOCAINE] – Special Authority see	SA0906 above – Retail pharm	acy	
Crm 4%	5.40	5 g OP	✓ LMX4
	27.00	30 g OP	✓ LMX4
LIDOCAINE [LIGNOCAINE] WITH PRILOCAINE - S	Special Authority see SA0906	above – Reta	il pharmacy
Crm 2.5% with prilocaine 2.5%	45.00	30 g OP	✓ EMLA
Crm 2.5% with prilocaine 2.5% (5 g tubes)	45.00	5	EMLA

Analgesics

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

Non-opioid Analgesics

ASPIRIN			
* Tab dispersible 300 mg - Up to 30 tab available on a PSO	4.50	100	Ethics Aspirin
CAPSAICIN - Subsidy by endorsement			
Subsidised only if prescribed for post-herpetic neuralgia or diab accordingly.	etic periphera	al neuropathy a	nd the prescription is endorsed
Crm 0.075%	11.95	45 g OP	✓ Zostrix HP
	15.83	57 g OP	✓ Rugby Capsaicin Topical Cream \$29
NEFOPAM HYDROCHLORIDE Tab 30 mg	23.40	90	✓ Acupan

	Subsidy (Manufacturer's Price		Fully Brand	
	\$	Per		facturer
PARACETAMOL				
Tab 500 mg - blister pack	19.75	1,000	✓ Pacimo	<u>I</u>
 a) Maximum of 300 tab per prescription; can be waived b) Up to 30 tab available on a PSO c) 	by endorsement			
1) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater, at annotate the prescription as endorsed where die 2) Maximum of 100 tab per dispensing for non-endorsed patients), then dispense in re Tab 500 mg - bottle pack — Maximum of 300 tab per	nd the prescription spensing history surprised patients. If peat dispensings n	is annotated a apports a long- quantities pre	accordingly. -term condition scribed for m	Pharmacists may on. nore than 100 tabs
prescription; can be waived by endorsement	17.92	1,000	✓ Noumed Parac	<u>d</u> etamol
 Subsidy by endorsement for higher quantities is aw daily dosing for one month or greater, and the pres prescription as endorsed where dispensing history Maximum of 100 tab per dispensing for non-endors non-endorsed patients), then dispense in repeat dispense in repeat dispense. 	cription is annotate supports a long-ter ed patients. If qua	d accordingly m condition. Intities prescri	. Pharmacis bed for more	ts may annotate the than 100 tabs (for
Oral liq 120 mg per 5 ml		1,000 ml 200 ml OP	✓ <u>Paracar</u> ✓ Avallon	_
 a) Maximum of 600 ml per prescription; can be waived b b) Up to 200 ml available on a PSO c) Not in combination d) 	y endorsement			
 Maximum of 200 ml per dispensing for non-end non-endorsed patients), then dispense in repea Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater an Pharmacists may annotate the prescription as e condition. 	t dispensing not ex available for patien d the prescription i	ceeding 200 r nts with long t s endorsed or	nl per disper erm conditio annotated a	nsing. ns who require accordingly.
Oral liq 240 mg per 5 ml		200 ml OP	✓ Avallon	\$29
1) Maximum of 200 ml per dispensing for non-end non-endorsed patients), then dispense in repea 2) Subsidy by endorsement for higher quantities is regular daily dosing for one month or greater ar Pharmacists may annotate the prescription as e condition.	t dispensing not ex available for patiend the prescription i	ceeding 200 r nts with long t s endorsed or	nl per disper erm conditio annotated a	nsing. ns who require accordingly.
Oral lig 250 mg per 5 ml	6.25	1 000 ml	✓ Paracar	e Double

Strength

Subsidy		rully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

- a) Maximum of 600 ml per prescription; can be waived by endorsement
- b) Up to 100 ml available on a PSO
- c) Not in combination
- 1) Maximum of 200 ml per dispensing for non-endorsed patients. If quantities prescribed exceed 200 ml (for non-endorsed patients), then dispense in repeat dispensing not exceeding 200 ml per dispensing.
- 2) Subsidy by endorsement for higher quantities is available for patients with long term conditions who require regular daily dosing for one month or greater and the prescription is endorsed or annotated accordingly. Pharmacists may annotate the prescription as endorsed where dispensing history supports a long-term condition.

*	Suppos 125 mg	10	✓ Gacet
	Suppos 250 mg		✓ Gacet
	Suppos 500 mg		✓ Gacet

CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency

Tab 15 mg6.25

Tab 30 mg7.45

Opioid Analgesics

32.80 Tab 60 mg14.25 DIHYDROCODEINE TARTRATE	100	✓ Aspen S29 ✓ PSM
Tab long-acting 60 mg8.60 DHC Continus to be Principal Supply on 1 December 2022	60	✓ DHC Continus
FENTANYL		
a) Only on a controlled drug form		
b) No patient co-payment payable		
c) Safety medicine; prescriber may determine dispensing frequency		
Inj 50 mcg per ml, 2 ml ampoule3.75	10	✓ Boucher and Muir
Inj 50 mcg per ml, 10 ml ampoule9.41	10	✓ Boucher and Muir
Patch 12.5 mcg per hour6.99	5	✓ Fentanyl Sandoz
Patch 25 mcg per hour7.99	5	✓ Fentanyl Sandoz
Patch 50 mcg per hour9.49	5	✓ Fentanyl Sandoz
Patch 75 mcg per hour17.99	5	✓ Fentanyl Sandoz
Patch 100 mcg per hour18.59	5	✓ Fentanyl Sandoz

METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency
- d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

е) For methadone h	ydrochloride oral lic	uid refer Standard	Formulae, page 2	248
---	-------------------	-----------------------	--------------------	------------------	-----

e) For methadone hydrochloride oral liquid refer S	Standard Formulae, page 24	₽Ŗ	
Tab 5 mg	1.40	10	✓ Methatabs
•	1.45		Methadone BNM
Oral liq 2 mg per ml	6.40	200 ml	✓ Biodone
Oral liq 5 mg per ml	6.40	200 ml	✓ Biodone Forte
Oral liq 10 mg per ml		200 ml	✓ Biodone Extra Forte
Inj 10 mg per ml, 1 ml	68.90	10	✓ AFT

(Methatabs Tab 5 mg to be delisted 1 February 2023)

✓ PSM

✓ PSM

100 100

	Subsidy		Fully	Brand or
	(Manufacturer's Price) Su	bsidised	Generic
	\$	Per	1	Manufacturer
ORPHINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing	frequency			
Oral lig 1 mg per ml		200 ml	√ F	RA-Morph
Oral liq 2 mg per ml		200 ml		RA-Morph
Oral lig 5 mg per ml		200 ml		Ordine \$29
Ording orning por minimum.		200 1111	-	RA-Morph
Oral liq 10 mg per ml	27.74	200 ml		Ordine \$29
Oral liq 10 mg per mi	27.74	200 1111		RA-Morph
			• ,	A-MOIPH
ORPHINE SULPHATE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
 Safety medicine; prescriber may determine dispensing 			_	
Tab immediate-release 10 mg		10	_	Sevredol
Tab immediate-release 20 mg		10	_	Sevredol
Cap long-acting 10 mg		10		n-Eslon
Cap long-acting 30 mg		10		n-Eslon
Cap long-acting 60 mg		10		n-Eslon
Cap long-acting 100 mg		10	-	n-Eslon
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a		5		/ledsurge
	6.99		✓ [DBL Morphine
				Sulphate
Inj 10 mg per ml, 1 ml ampoule - Up to 5 inj available on	a PSO4.68	5	✓ N	/ledsurge
	5.61		✓ [BL Morphine
				Sulphate
Inj 15 mg per ml, 1 ml ampoule - Up to 5 inj available on	a PSO5.53	5	✓ N	/ledsurge
	7.08		✓ [OBL Morphine
				Sulphate
Inj 30 mg per ml, 1 ml ampoule - Up to 5 inj available on	a PSO6.28	5	✓ N	Nedsurge
, , , , , , , , , , , , , , , , , , , ,	7.28			BL Morphine
				Sulphate

(DBL Morphine Sulphate Inj 5 mg per ml, 1 ml ampoule to be delisted 1 March 2023) (DBL Morphine Sulphate Inj 10 mg per ml, 1 ml ampoule to be delisted 1 March 2023) (DBL Morphine Sulphate Inj 15 mg per ml, 1 ml ampoule to be delisted 1 March 2023) (DBL Morphine Sulphate Inj 30 mg per ml, 1 ml ampoule to be delisted 1 March 2023)

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

-	Subsidy	.\	Fully	
(M	anufacturer's Price \$	e) Per	Subsidised	Generic Manufacturer
XYCODONE HYDROCHLORIDE	· · · · · · · · · · · · · · · · · · ·			
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequ	encv			
Tab controlled-release 5 mg	•	20	1	Oxycodone Sando
Tab controlled-release 10 mg		20		Oxycodone Sando
Tab controlled-release 20 mg		20		Oxycodone Sando
Tab controlled-release 40 mg		20		Oxycodone Sando
Tab controlled-release 80 mg.		20		Oxycodone Sando
Cap immediate-release 5 mg		20		OxyNorm
Cap immediate-release 10 mg		20		OxyNorm
Cap immediate-release 20 mg		20		OxyNorm
1 5				
Oral liq 5 mg per 5 ml		250 m		OxyNorm
Inj 10 mg per ml, 1 ml ampoule		5		Hameln
Inj 10 mg per ml, 2 ml ampoule		5		Hameln
Inj 50 mg per ml, 1 ml ampoule	22.92	5	•	<u>Hameln</u>
RACETAMOL WITH CODEINE - Safety medicine; prescriber ma	ay determine disp	pensing	frequenc	γ
Tab paracetamol 500 mg with codeine phosphate 8 mg		1,000		Paracetamol +
				Codeine (Relieve
THIDINE HYDROCHLORIDE				
a) Only on a controlled drug form				
b) No patient co-payment payable				
c) Safety medicine; prescriber may determine dispensing frequ	•		_	
Tab 50 mg		10		<u>PSM</u>
Inj 50 mg per ml, 1 ml ampoule - Up to 5 inj available on a PSC)29.88	5	✓	DBL Pethidine
				Hydrochloride
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a PSC)30.72	5	1	DBL Pethidine
				Hydrochloride
RAMADOL HYDROCHLORIDE				•
Tab sustained-release 100 mg	1.50	20	./	Tramal SR 100
		20		
Tab sustained-release 150 mg				Tramal SR 150
Tab sustained-release 200 mg		20		Tramal SR 200
Cap 50 mg	2.80	100	•	Arrow-Tramadol
Antidepressants				
Cyclic and Related Agents				
MITRIPTYLINE – Safety medicine; prescriber may determine disp	0 1 3	,		
Tab 10 mg	2.49	100		Arrow-Amitriptylin
Tab 25 mg	1.51	100	•	Arrow-Amitriptylin
Tab 50 mg	2.51	100		Arrow-Amitriptylin
OMIPRAMINE HYDROCHLORIDE - Safety medicine; prescriber	may determine		sing frequ	ency
Tab 10 mg		30		Clomipramine Teva
Tab 25 mg	44.00	30		Clomipramine Teva

					IIVOOS STSTEM
		Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
DC	SULEPIN [DOTHIEPIN] HYDROCHLORIDE - Subsidy by er	ndorsement			
	Safety medicine; prescriber may determine dispensing fr Subsidy by endorsement – Subsidised for patients who v 2019 and the prescription is endorsed accordingly. Phar exists a record of prior dispensing of dosulepin [dothiepin]	requency were taking dosulepin rmacists may annotate			
	Tab 75 mg		30		Dosulepin Mylan Dosulepin Viatris
	Cap 25 mg	7.83	50	•	Dosulepin Mylan \$29
IMI	PRAMINE HYDROCHLORIDE - Safety medicine; prescriber	r mav determine dispe	nsino	ı frequency	•
	Tab 10 mg		50		Tofranil
		10.96	100		Tofranil
	Tab 25 mg	8.80	50	1	Tofranil
NC	RTRIPTYLINE HYDROCHLORIDE - Safety medicine; presc	criber may determine d	lisper	าsing frequ	uency
	Tab 10 mg		100	_	Norpress
	Tab 25 mg	5.98	180	/	Norpress
N	Ionoamine-Oxidase Inhibitors (MAOIs) - Non S	Selective			
TR	ANYLCYPROMINE SULPHATE				
	Tab 10 mg	12.85	28	/	Parnate S29 S29
	ů	22.94	50	✓	Parnate
		45.88	100	✓	Parnate S29 S29
		96.00		1	Parnate S29 S29
N	Ionoamine-Oxidase Type A Inhibitors				
MC	OCLOBEMIDE				
	Tab 150 mg	11.80	60	1	Aurorix
	Tab 300 mg		60		Aurorix
			-		
S	elective Serotonin Reuptake Inhibitors				
	TALOPRAM HYDROBROMIDE			_	
*	Tab 20 mg		84		PSM Citalopram
		2.86		•	Celapram
_	CITALOPRAM				
	Tab 10 mg		28	•	Escitalopram (Ethics)
*	Tab 20 mg	1.92	28	/	Escitalopram (Ethics)
FLI	JOXETINE HYDROCHLORIDE				
	Tab dispersible 20 mg, scored – Subsidy by endorsement Subsidised by endorsement	2.50	28	•	Fluox
	When prescribed for a patient who cannot swallow	whole tablets or caps	sules	and the pr	rescription is endorsed
	accordingly; orWhen prescribed in a daily dose that is not a multi endorsed. Note: Tablets should be combined wit				
	Cap 20 mg	2.91	84	1	Fluox

	Subsidy (Manufacturer's Price \$) Per	Fully Subsidised	
AROXETINE	Ψ	1 01		Waltulactarer
Anoxemine ← Tab 20 mg	4 11	90	/	Loxamine
ERTRALINE		00		L OXUIIIIO
€ Tab 50 mg	0.02	30	_	Setrona
• Tab 50 mg		00		Setrona AU
€ Tab 100 mg	1.61	30		Setrona
			✓	Setrona AU
Other Antidepressants				
IIRTAZAPINE				
Tab 30 mg	2.60	28	/	Noumed
Tab 45 mg		28	•	Noumed
ENLAFAXINE				
€ Cap 37.5 mg	6.38	84	1	Enlafax XR
€ Cap 75 mg		84	•	Enlafax XR
Cap 150 mg	11.16	84	✓	Enlafax XR
Antiepilepsy Drugs				
Agents for Control of Status Epilepticus				
IAZEPAM – Safety medicine; prescriber may determine dispe	nsing frequency			
Inj 5 mg per ml, 2 ml ampoule — Subsidy by endorsement		5	/	Hospira
a) Up to 5 inj available on a PSO		·		
b) Only on a PSO				
c) PSO must be endorsed "not for anaesthetic proced	ıres".			
Rectal tubes 5 mg - Up to 5 tube available on a PSO	54.58	5	1	Stesolid
HENYTOIN SODIUM				
Inj 50 mg per ml, 2 ml ampoule - Up to 5 inj available on a				
PSO	104.58	5	•	Hospira
Ini FO ma nor ml. E ml. omnoulo Un to F ini quailable an o				
Inj 50 mg per ml, 5 ml ampoule – Up to 5 inj available on a				
PSOPSO	154.01	5	✓	Hospira
PSO	154.01	5	•	Hospira
PSO	154.01	5	•	Hospira
PSO Control of Epilepsy ARBAMAZEPINE		100		Hospira Tegretol
PSO Control of Epilepsy ARBAMAZEPINE Tab 200 mg	14.53		•	
PSO Control of Epilepsy ARBAMAZEPINE Tab 200 mg Tab long-acting 200 mg	14.53 16.98	100	<i>y</i>	Tegretol
PSO Control of Epilepsy ARBAMAZEPINE Tab 200 mg Tab long-acting 200 mg Tab 400 mg	14.53 16.98 34.58	100 100	<i>y y y y</i>	Tegretol Tegretol CR Tegretol Tegretol CR
PSO Control of Epilepsy ARBAMAZEPINE Tab 200 mg Tab long-acting 200 mg Tab 400 mg Tab long-acting 400 mg	14.53 16.98 34.58 39.17	100 100 100	<i>y y y y</i>	Tegretol Tegretol CR Tegretol
PSO		100 100 100 100	<i>y y y y</i>	Tegretol Tegretol CR Tegretol Tegretol CR
PSO		100 100 100 100		Tegretol Tegretol CR Tegretol Tegretol CR
PSO		100 100 100 100 250 ml		Tegretol Tegretol CR Tegretol Tegretol CR Tegretol CR Tegretol
PSO		100 100 100 100 250 ml		Tegretol Tegretol CR Tegretol Tegretol CR Tegretol CR Tegretol
PSO		100 100 100 100 250 ml		Tegretol Tegretol CR Tegretol Tegretol CR Tegretol CR Tegretol
PSO		100 100 100 100 250 ml	/ / / / / / / / / / / / / / / / / / /	Tegretol Tegretol CR Tegretol Tegretol CR Tegretol CR Tegretol
PSO		100 100 100 100 250 ml 50	/ / / / / / / / / / / / / / / / / / /	Tegretol Tegretol CR Tegretol Tegretol CR Tegretol CR Tegretol Frisium
PSO		100 100 100 100 250 ml 50		Tegretol Tegretol CR Tegretol CR Tegretol CR Tegretol Frisium Rivotril

	Subsidy		Fully	Brand or
	(Manufacturer's Pric	ce) Su Per	ıbsidised •	Generic Manufacturer
	J	rei		Manuacturer
ABAPENTIN				
Note: Not subsidised in combination with	subsidised pregabalin			
★ Cap 100 mg	6.45	100	✓ N	<u>lupentin</u>
★ Cap 300 mg	8.45	100	✓ N	lupentin
★ Cap 400 mg	10.26	100	✓ N	lupentin
ACOSAMIDE - Special Authority see SA112	25 below – Retail pharmacy			
·		4.4		
▲ Tab 50 mg	25.04	14	V V	/impat
▲ Tab 50 mg		14 14		/impat /impat
· ·			✓ V	•
Tab 100 mg	50.06 200.24	14	✓ V ✓ V	/impat
· ·	50.06 200.24	14 56	✓ V ✓ V	/impat /impat

SA1125 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Patient has partial-onset epilepsy; and
- 2 Seizures are not adequately controlled by, or patient has experienced unacceptable side effects from, optimal treatment with all of the following: sodium valproate, topiramate, levetiracetam and any two of carbamazepine, lamotrigine and phenytoin sodium (see note).

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient has demonstrated a significant and sustained improvement in seizure rate or severity and/or quality of life compared with that prior to starting lacosamide treatment.

LAMOTRIGINE

	Tab dispersible 2 mg55.00	30	✓ Lamictal
	Tab dispersible 5 mg50.00	30	✓ Lamictal
*	Tab dispersible 25 mg2.76	56	✓ Logem
*	Tab dispersible 50 mg3.31	56	✓ Logem
*	Tab dispersible 100 mg4.40	56	✓ Logem
ΙF	VETIRACETAM		•
	Tab 250 mg4.99	60	✓ Everet
	Tab 500 mg8.79	60	✓ Everet
	Tab 750 mg14.39	60	✓ Everet
	Tab 1,000 mg18.59	60	✓ Everet
	Oral liq 100 mg per ml44.78	300 ml OP	✓ Levetiracetam-AFT
PH	IENOBARBITONE		
	For phenobarbitone oral liquid refer Standard Formulae, page 248		
*	Tab 15 mg40.00	500	✓ PSM
*	Tab 30 mg40.00	500	✓ PSM
PH	IENYTOIN SODIUM		
*	Tab 50 mg	200	✓ Dilantin Infatab
	Cap 30 mg	200	✓ Dilantin
	Cap 100 mg37.00	200	✓ Dilantin
*	Oral liq 30 mg per 5 ml	500 ml	✓ Dilantin
PF	EGABALIN		
• •	Note: Not subsidised in combination with subsidised gabapentin		
	Cap 25 mg	56	✓ Pregabalin Pfizer
*		56	✓ Pregabalin Pfizer
•	Cap 150 mg	56	✓ Lyrica
		30	✓ Pregabalin Pfizer
	Cap 300 mg7.38	56	✓ Pregabalin Pfizer
			-

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	I Generic
PRIMIDONE				
* Tab 250 mg	37.35	100		Apo-Primidone Primidone Clinect
(Apo-Primidone Tab 250 mg to be delisted 1 January 2023)				
SODIUM VALPROATE				
Tab 100 mg	13.65	100	✓	Epilim Crushable
Tab 200 mg EC	27.44	100	✓	Epilim
Tab 500 mg EC	52.24	100	1	Epilim
* Oral lig 200 mg per 5 ml	20.48	300 m		Epilim S/F Liquid
			1	Epilim Syrup
* Inj 100 mg per ml, 4 ml	41.50	1	1	Epilim IV
STIRIPENTOL - Special Authority see SA1330 below - Retail pl	harmacy			-
Cap 250 mg	509.29	60	1	Diacomit S29
Powder for oral liq 250 mg sachet	509.29	60	✓	Diacomit \$29

⇒SA1330 Special Authority for Subsidy

TODIDAMATE

Initial application only from a paediatric neurologist or Practitioner on the recommendation of a paediatric neurologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed diagnosis of Dravet syndrome; and
- 2 Seizures have been inadequately controlled by appropriate courses of sodium valproate, clobazam and at least two of the following: topiramate, levetiracetam, ketogenic diet.

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified where the patient continues to benefit from treatment as measured by reduced seizure frequency from baseline.

Tab 25 mg	11.07	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	26.04		✓ Topamax
Tab 50 mg	18.81	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	44.26		✓ Topamax
Tab 100 mg	31.99	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	75.25		✓ Topamax
Tab 200 mg	55.19	60	✓ Arrow-Topiramate
•			✓ Topiramate Actavis
	129.85		✓ Topamax
Sprinkle cap 15 mg	20.84	60	✓ Topamax
Sprinkle cap 25 mg	26.04	60	✓ Topamax
GABATRIN – Special Authority see SA2088 below			•
Tab 500 mg		100	✓ Sabril

⇒SA2088 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 15 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 Patient has infantile spasms; or
 - 1.2 Both:
 - 1.2.1 Patient has epilepsy; and

Subsic (Manufacture	. ,	,	nd or neric
\$	Per	✓ Mar	nufacturer

continued...

- 1.2.2 Either:
 - 1.2.2.1 Seizures are not adequately controlled with optimal treatment with other antiepilepsy agents; or
 - 1.2.2.2 Seizures are controlled adequately but the patient has experienced unacceptable side effects from optimal treatment with other antiepilepsy agents; or
- 1.3 Patient has tuberous sclerosis complex; and
- 2 Either:
 - 2.1 Patient is, or will be, receiving regular automated visual field testing (ideally before starting therapy and on a 6-monthly basis thereafter): or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Renewal from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 The patient has demonstrated a significant and sustained improvement in seizure rate or severity and or quality of life; and
- 2 Fither:
 - 2.1 Patient is receiving regular automated visual field testing (ideally every 6 months) on an ongoing basis for duration of treatment with vigabatrin; or
 - 2.2 It is impractical or impossible (due to comorbid conditions) to monitor the patient's visual fields...

Antimigraine Preparations

For Anti-inflammatory NSAIDS refer to MUSCULOSKELETAL, page 110

3.65	30	✓ Rizamelt
14.41	90	✓ Sumagran
22.68	90	✓ Sumagran
		_
34.00	2 OP	Imigran
	14.41 22.68	14.41 90 22.68 90

Prophylaxis of Migraine

For Beta Adrenoceptor Blockers refer to CARDIOVASCULAR SYSTEM, page 51

PIZOTIFEN

* Tab 500 mcg......23.21 100 Sandomigran

Antinausea and Vertigo Agents

For Antispasmodics refer to ALIMENTARY TRACT, page 8

APREPITANT - Special Authority see SA0987 below - Retail pharmacy

3 OP ✓ Emend Tri-Pack

⇒SA0987 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

Renewal from any relevant practitioner. Approvals valid for 12 months where the patient is undergoing highly emetogenic chemotherapy and/or anthracycline-based chemotherapy for the treatment of malignancy.

BETAHISTINE DIHYDROCHLORIDE

100 ✓ Serc

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
CYCLIZINE HYDROCHLORIDE				
Tab 50 mg	0.49	10	✓ V	<u>Nausicalm</u>
CYCLIZINE LACTATE Inj 50 mg per ml, 1 ml ampoule Hameln to be Principal Supply on 1 December 2022	16.36	10	✓ H	Hameln
DOMPERIDONE * Tab 10 mg	2.85	100	✓ <u>F</u>	Pharmacy Health
HYOSCINE HYDROBROMIDE				
* Inj 400 mcg per ml, 1 ml ampoule	93.00	10	✓ N	Martindale \$29
Patch 1.5 mg – Special Authority see SA1998 below – Retail pharmacy		2	√ 9	Scopoderm TTS

⇒SA1998 Special Authority for Subsidy

METOCLOPRAMIDE HYDROCHLORIDE

Initial application from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria: Either:

- 1 Control of intractable nausea, vomiting, or inability to swallow saliva in the treatment of malignancy or chronic disease where the patient cannot tolerate or does not adequately respond to oral anti-nausea agents; or
- 2 Control of clozapine-induced hypersalivation where trials of at least two other alternative treatments have proven ineffective.

Renewal from any relevant practitioner. Approvals valid for 1 year where the treatment remains appropriate and the patient is benefiting from treatment.

*	Tab 10 mg - Up to 30 tab available on a PSO1.30	100	✓ Metoclopramide Actavis 10
*	Inj 5 mg per ml, 2 ml ampoule $-$ Up to 5 inj available on a PSO7.00 9.50	10	✓ Baxter ✓ Pfizer
,	Baxter to be Principal Supply on 1 December 2022 zer Inj 5 mg per ml, 2 ml ampoule to be delisted 1 December 2022)		
-	DANSETRON		
*	Tab 4 mg	50	✓ Onrex
	Tab disp 4 mg - Up to 10 tab available on a PSO	10	✓ Ondansetron ODT-DRLA
*	Tab 8 mg4.57	50	✓ Onrex
	Tab disp 8 mg - Up to 10 tab available on a PSO1.13	10	✓ <u>Ondansetron</u> <u>ODT-DRLA</u>
PRO	OCHLORPERAZINE		
*	Tab 3 mg buccal	50	
	(30.00)		Buccastem
*	Tab 5 mg – Up to 30 tab available on a PSO8.00	250	✓ Nausafix
	Inj 12.5 mg per ml, 1 ml - Up to 5 inj available on a PSO25.81	10	✓ Stemetil

Antipsychotics

General

AMISULPRIDE - Safety medicine; prescriber may determine dispensing frequency		
Tab 100 mg5.15	30	✓ Sulprix
Tab 200 mg14.96	60	✓ Sulprix
Tab 400 mg29.78	60	✓ Sulprix

	Subsidy		Fully Brand or
	(Manufacturer's Price)	Per	Subsidised Generic Manufacturer
	Ψ	rei	ivianulacturei
ARIPIPRAZOLE – Safety medicine; prescriber may determine d			
Tab 5 mg		30	✓ Aripiprazole Sandoz
Tab 10 mg		30	Aripiprazole Sandoz
Tab 15 mg		30	Aripiprazole Sandoz
Tab 20 mg		30	Aripiprazole Sandoz
Tab 30 mg	10.50	30	Aripiprazole Sandoz
CHLORPROMAZINE HYDROCHLORIDE - Safety medicine; pr	escriber may determi	ne dis	spensing frequency
Tab 10 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Tab 25 mg - Up to 30 tab available on a PSO	15.62	100	✓ Largactil
Tab 100 mg - Up to 30 tab available on a PSO		100	✓ Largactil
Inj 25 mg per ml, 2 ml - Up to 5 inj available on a PSO		10	✓ Largactil
CLOZAPINE – Hospital pharmacy [HP4]			3
Safety medicine; prescriber may determine dispensing frequ	lonov.		
	•	50	✓ Clopino
Tab 25 mg	0.09	50	✓ Clopine✓ Clozaril
	10.07	100	
	13.37	100	✓ Clopine
T 50	0.07		✓ Clozaril
Tab 50 mg		50	Clopine
	17.33	100	Clopine
Tab 100 mg	17.33	50	Clopine
			✓ Clozaril
	34.65	100	Clopine
			✓ Clozaril
Tab 200 mg	34.65	50	✓ Clopine
	69.30	100	✓ Clopine
Suspension 50 mg per ml	67.62	100 m	nl ✓ Versacloz
HALOPERIDOL - Safety medicine; prescriber may determine di	ispensing frequency		
Tab 500 mcg – Up to 30 tab available on a PSO		100	✓ Serenace
Tab 1.5 mg – Up to 30 tab available on a PSO		100	_
Tab 5 mg - Up to 30 tab available on a PSO		50	✓ Serenace
Tub o mg op to oo tub available on a roo	29.72	100	✓ Serenace
Oral lig 2 mg per ml - Up to 200 ml available on a PSO		100 m	
Inj 5 mg per ml, 1 ml ampoule – Up to 5 inj available on a P		100 11	✓ Serenace
LEVOMEPROMAZINE – Safety medicine; prescriber may deter			
Tab 25 mg (33.8 mg as a maleate)		100	,
Tab 25 mg as a maleate		100	
Tab 100 mg (135 mg as a maleate)		100	Nozinan (Swiss)
Tab 100 mg as a maleate	41.75	100	Nozinan
LEVOMEPROMAZINE HYDROCHLORIDE - Safety medicine;	orescriber may deterr	nine d	dispensing frequency
Inj 25 mg per ml, 1 ml ampoule	-	5	✓ Neuraxpharm \$29
ing to mg por mi, i mi amposio		Ū	✓ Nozinan S29 S29
	33.50	10	✓ Nozinan
LITHIUM CARBONATE - Safety medicine; prescriber may dete		quency	
Tab long-acting 400 mg	72.00	100	
Cap 250 mg	9.42	100	✓ Douglas
OLANZAPINE - Safety medicine; prescriber may determine dis	pensina frequency		
Tab 2.5 mg		28	✓ Zypine
Tab 5 mg		28	✓ Zypine
Tab orodispersible 5 mg		28	✓ Zypine ODT
Tab 10 mg		28	✓ Zypine OD1 ✓ Zypine
Tab orodispersible 10 mg		28	✓ Zypine ODT
rab orodisporsible to mg	2.00	20	- Lypine OD1

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist. ★Three months or six months, as applicable, dispensed all-at-once

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
PERICYAZINE - Safety medicine; prescriber may determine dis	pensing frequency			
Tab 2.5 mg	10.49	84	1	Neulactil
•	12.49	100	1	Neulactil
Tab 10 mg	37.34	84	✓	Neulactil
•	44.45	100	1	Neulactil
QUETIAPINE - Safety medicine; prescriber may determine disp	ensina frequency			
Tab 25 mg		90	1	Quetapel
Tab 100 mg		90		Quetapel
Tab 200 mg		90		Quetapel
Tab 300 mg		90	1	Quetapel
RISPERIDONE – Safety medicine; prescriber may determine dis	spensing frequency			
Tab 0.5 mg		60	1	Risperidone (Teva)
Tab 1 mg		60		Risperidone (Teva)
Tab 2 mg		60		Risperidone (Teva)
Tab 3 mg		60		Risperidone (Teva)
Tab 4 mg		60		Risperidone (Teva)
Oral lig 1 mg per ml		30 m	· •	Risperon
ZIPRASIDONE - Safety medicine; prescriber may determine dis	spensing frequency			
Cap 20 mg		60	1	Zusdone
Cap 40 mg		60		Zusdone
Cap 60 mg		60		Zusdone
Cap 80 mg		60	_	Zusdone
ZUCLOPENTHIXOL HYDROCHLORIDE – Safety medicine; pre	•	100		equency Clopixol
Tab 10 mg	31.45	100	•	Ciopixoi

Depot Injections

FLUPENTHIXOL DECANOATE – Safety medicine; prescriber may determine disp Inj 20 mg per ml, 1 ml – Up to 5 inj available on a PSO	ensing freq 5 5	uency ✓ Fluanxol ✓ Fluanxol
Inj 100 mg per ml, 1 ml — Up to 5 inj available on a PSO40.87	5	✓ Fluanxol
HALOPERIDOL DECANOATE - Safety medicine; prescriber may determine dispe	nsing frequ	ency
Inj 50 mg per ml, 1 ml - Up to 5 inj available on a PSO28.39	5	✓ Haldol
Inj 100 mg per ml, 1 ml – Up to 5 inj available on a PSO55.90	5	✓ Haldol Concentrate✓ Haldol
		Decanoas S29
OLANZAPINE - Special Authority see SA1428 below - Retail pharmacy		
Safety medicine; prescriber may determine dispensing frequency		
Inj 210 mg vial252.00	1	✓ Zyprexa Relprevv
Inj 300 mg vial414.00	1	✓ Zyprexa Relprevv
Inj 405 mg vial504.00	1	✓ Zyprexa Relprevv

⇒SA1428 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or risperidone depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia; and

(Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	Subsidy	Fully	Brand or
\$ Per ✓ Manufacturer	(Manufacturer's Price)		
	 \$	Per 🗸	Manufacturer

continued...

- 2.2 The patient has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
- 2.3 The patient has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in the last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of olanzapine depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

PALIPERIDONE - Special Authority see SA1429 below - Retail pharmacy

Safety medicine; prescriber may determine dispensing	g frequency		
Inj 25 mg syringe	194.25	1	✓ Invega Sustenna
Inj 50 mg syringe	271.95	1	✓ Invega Sustenna
Inj 75 mg syringe		1	✓ Invega Sustenna
Inj 100 mg syringe	435.12	1	✓ Invega Sustenna
Inj 150 mg syringe	435.12	1	✓ Invega Sustenna

⇒SA1429 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for risperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of paliperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

RISPERIDONE - Special Authority see SA1427 below - Retail pharmacy

Inj 25 mg vial	Risperdal Consta
Inj 37.5 mg vial	 Risperdal Consta
Inj 50 mg vial217.56	✓ Risperdal Consta

⇒SA1427 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Either:

- 1 The patient has had an initial Special Authority approval for paliperidone depot injection or olanzapine depot injection; or
- 2 All of the following:
 - 2.1 The patient has schizophrenia or other psychotic disorder; and
 - 2.2 Has tried but failed to comply with treatment using oral atypical antipsychotic agents; and
 - 2.3 Has been admitted to hospital or treated in respite care, or intensive outpatient or home-based treatment for 30 days or more in last 12 months.

Renewal from any relevant practitioner. Approvals valid for 12 months where the initiation of risperidone depot injection has been associated with fewer days of intensive intervention than was the case during a corresponding period of time prior to the initiation of an atypical antipsychotic depot injection.

ZUCLOPENTHIXOL DECANOATE - Safety medicine; prescriber may determine dispensing frequency



	Subsidy	F	ully	Brand or
(Ma	nufacturer's Price)	Subsidi	sed	Generic
	\$	Per	<u> </u>	Manufacturer

Anxiolytics

BUSPIRONE HYDROCHLORIDE			
* Tab 5 mg	18.50	100	 Buspirone Viatris
* Tab 10 mg	12.50	100	✓ Buspirone Viatris
CLONAZEPAM - Safety medicine; prescriber may determine dis	spensing frequency		
Tab 500 mcg	5.64	100	✓ Paxam
Tab 2 mg	10.78	100	✓ Paxam
DIAZEPAM - Safety medicine; prescriber may determine disper	nsing frequency		
Tab 2 mg	61.07	500	✓ Arrow-Diazepam
Tab 5 mg	73.60	500	✓ Arrow-Diazepam
LORAZEPAM - Safety medicine; prescriber may determine disp	ensing frequency		
Tab 1 mg	9.72	250	✓ Ativan
Tab 2.5 mg	12.50	100	✓ <u>Ativan</u>

Multiple Sclerosis Treatments

⇒SA2140 Special Authority for Subsidy

Initial application — (Multiple sclerosis) only from a neurologist or general physician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Diagnosis of multiple sclerosis (MS) meets the McDonald 2017 diagnostic criteria for MS and has been confirmed by a neurologist: and
- 2 Patients has an EDSS score between 0 6.0; and
- 3 Patient has had at least one significant attack of MS in the previous 12 months or two significant attacks in the past 24 months; and
- 4 All of the following:
 - 4.1 Each significant attack must be confirmed by the applying neurologist or general physician (the patient may not necessarily have been seen by them during the attack, but the neurologist/physician must be satisfied that the clinical features were characteristic); and
 - 4.2 Each significant attack is associated with characteristic new symptom(s)/sign(s) or substantially worsening of previously experienced symptoms(s)/sign(s); and
 - 4.3 Each significant attack has lasted at least one week and has started at least one month after the onset of a previous attack (where relevant); and
 - 4.4 Each significant attack can be distinguished from the effects of general fatigue; and is not associated with a fever (T> 37.5°C); and
 - 4.5 Fither:
 - 4.5.1 Each significant attack is severe enough to change either the EDSS or at least one of the Kurtze Functional System scores by at least 1 point; or
 - 4.5.2 Each significant attack is a recurrent paroxysmal symptom of multiple sclerosis (tonic seizures/spasms, trigeminal neuralgia. Lhermitte's symptom): and
- 5 Evidence of new inflammatory activity on an MRI scan within the past 24 months; and
- 6 Any of the following:
 - 6.1 A sign of that new inflammatory activity on MRI scanning (in criterion 5 immediately above) is a gadolinium enhancing lesion: or
 - 6.2 A sign of that new inflammatory activity is a lesion showing diffusion restriction; or
 - 6.3 A sign of that new inflammatory is a T2 lesion with associated local swelling; or

		-		
	Subsidy (Manufacturer's Price)	Subsid Per	ully ised	Brand or Generic Manufacturer
continued 6.4 A sign of that new inflammatory activity is a promin a recent attack that occurred within the last 2 years 6.5 A sign of that new inflammatory activity is new T2 le Note: Natalizumab can only be dispensed from a pharmacy regis operated by the supplier. Treatment on two or more funded multi	; or esions compared with tered in the Tysabri <i>F</i>	n a previous Australasian	MRI s	can. cribing Programme
Renewal — (Multiple sclerosis) only from a neurologist or gene nad an EDSS score of 0 to 6.0 (inclusive) with or without the use i.e. the patient has walked 100 metres or more with or without at Note: Natalizumab can only be dispensed from a pharmacy regis operated by the supplier. Treatment on two or more funded multi DIMETHYL FUMARATE — Special Authority see SA2140 on the	ral physician. Appro of unilateral or bilater ds in the last six mon tered in the Tysabri A ple sclerosis treatme	vals valid fo al aids at ar aths). Australasian nts simultan	r 12 m ny time Presc eously	onths where patient has in the last six months cribing Programme
a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Cap 120 mg Cap 240 mg	s treatments simultan		ot perm	nitted. ecfidera ecfidera
FINGOLIMOD – Special Authority see SA2140 on the previous p a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Cap 0.5 mg	s treatments simultan		•	nitted. lenya
GLATIRAMER ACETATE – Special Authority see SA2140 on the Note: Treatment on two or more funded multiple sclerosis tre Inj 40 mg prefilled syringe	atments simultaneou		ermitte	d. opaxone
NTERFERON BETA-1-ALPHA – Special Authority see SA2140 Note: Treatment on two or more funded multiple sclerosis tre Inj 6 million iu prefilled syringe	eatments simultaneou 1,170.00		ermitte	
NTERFERON BETA-1-BETA — Special Authority see SA2140 or Note: Treatment on two or more funded multiple sclerosis tre Inj 8 million iu per 1 ml	atments simultaneou		ermitte	d. etaferon
NATALIZUMAB – Special Authority see SA2140 on the previous Note: Treatment on two or more funded multiple sclerosis tre Inj 20 mg per ml, 15 ml vial	atments simultaneou			d. rsabri
OCRELIZUMAB - Special Authority see SA2140 on the previous Note: Treatment on two or more funded multiple sclerosis tre Inj 30 mg per ml, 10 ml vial	atments simultaneou			d. crevus
TERIFLUNOMIDE – Special Authority see SA2140 on the previo a) Wastage claimable b) Note: Treatment on two or more funded multiple sclerosis Tab 14 mg	s treatments simultan	•	٠	nitted. ubagio
Sedatives and Hypnotics				
MELATONIN – Special Authority see SA1666 on the next page – Tab modified-release 2 mg – No more than 5 tab per day		30	✓ Vi	gisom



Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

⇒SA1666 Special Authority for Subsidy

Initial application only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with persistent and distressing insomnia secondary to a neurodevelopmental disorder (including, but not limited to, autism spectrum disorder or attention deficit hyperactivity disorder)*; and
- 2 Behavioural and environmental approaches have been tried and were unsuccessful, or are inappropriate; and
- 3 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day; and
- 4 Patient is aged 18 years or under*.

Renewal only from a psychiatrist, paediatrician, neurologist, respiratory specialist or medical practitioner on the recommendation of a psychiatrist, paediatrician, neurologist or respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is aged 18 years or under*; and
- 2 Patient has demonstrated clinically meaningful benefit from funded modified-release melatonin (clinician determined); and
- 3 Patient has had a trial of funded modified-release melatonin discontinuation within the past 12 months and has had a recurrence of persistent and distressing insomnia; and
- 4 Funded modified-release melatonin is to be given at doses no greater than 10 mg per day.

MIDAZOLAM Cofety modicine: properitor may determine dispensing frequency

Note: Indications marked with * are unapproved indications.

MIDAZOLAM – Safety medicine; prescriber may determine dispens	sing frequency		
Inj 1 mg per ml, 5 ml ampoule	6.10	10	✓ Midazolam-Baxter
Inj 1 mg per ml, 5 ml plastic ampoule - Up to 10 inj available			
on a PSO	17.28	10	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be en	dorsed for statu	s epilepticu	us use only.
Inj 5 mg per ml, 3 ml ampoule	5.00	5	✓ Midazolam-Baxter
Inj 5 mg per ml, 3 ml plastic ampoule - Up to 5 inj available on	1		
a PSO	13.09	5	✓ Pfizer
On a PSO for status epilepticus use only. PSO must be en	dorsed for statu	ıs epilepticı	us use only.
PHENOBARBITONE SODIUM - Special Authority see SA1386 bel	ow – Retail pha	rmacy	
Inj 200 mg per ml, 1 ml ampoule	103.30	10	✓ Max Health S29
SA1386 Special Authority for Subsidy			

SA1386 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

- 1 For the treatment of terminal agitation that is unresponsive to other agents; and
- 2 The applicant is part of a multidisciplinary team working in palliative care.

TEMAZEPAM – Safety medicine; prescriber may determine Tab 10 mg	1 0 1 ,	25	✓ Normison
TRIAZOLAM – Safety medicine; prescriber may determine d			
Tab 125 mcg	5.10	100	
·	(9.85)		Hypam
Tab 250 mcg	4.10 [°]	100	
Ç	(11.20)		Hypam
ZOPICLONE - Safety medicine; prescriber may determine of	lispensing frequency		
Tab 7.5 mg		500	✓ Zopiclone Actavis

	Subsidy (Manufacturer's Price) \$	Per	Fully Brand or Subsidised Generic Manufacturer
Stimulants/ADHD Treatments			
ATOMOXETINE			
Cap 10 mg	18.41	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			S29 S29
			✓ Generic Partners
	107.03		✓ Strattera
Cap 18 mg	27.06	28	✓ APO-Atomoxetine
			Generic Partners
	107.03		✓ Strattera
Cap 25 mg	29.22	28	✓ APO-Atomoxetine
			✓ Generic Partners
Cap 40 mg	29.22	28	✓ APO-Atomoxetine
	407.00		✓ Generic Partners
0.00 00 00	107.03	00	✓ Strattera
Cap 60 mg	46.51	28	✓ APO-Atomoxetine ✓ APO-Atomoxetine
			•
			S29 S29
0.00 00 00	50.45	00	✓ Generic Partners
Cap 80 mg	56.45	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			S29 S29
0	50.40	00	✓ Generic Partners
Cap 100 mg	58.48	28	✓ APO-Atomoxetine
			✓ APO-Atomoxetine
			\$29 S29
			Generic Partners
DEXAMFETAMINE SULFATE - Special Authority see SA1149	•	су	
a) Brand switch fee payable (Pharmacode 2641356) - see	page 246 for details		
b) Only on a controlled drug form			
c) Safety medicine; prescriber may determine dispensing fr			
Tab 5 mg	21.00	100	✓ <u>PSM</u>

⇒SA1149 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

28.50


All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

continued...

✓ Aspen S29



Subsidy		Fully	Brand or	
(Manufacturer's Price)	S	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Both:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist or medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Fither:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

METHYLPHENIDATE HYDROCHLORIDE - Special Authority see SA1964 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency ✓ Rubifen Tab immediate-release 5 mg.......3.20 30 ✓ Ritalin 30 ✓ Rubifen Tab extended-release 18 mg......7.75 ✓ Methylphenidate ER 30 - Teva ✓ Rubifen 30 ✓ Rubifen SR 30 ✓ Methylphenidate ER Tab extended-release 27 mg.......11.45 30 - Teva Tab extended-release 36 mg......15.50 30 ✓ Methylphenidate ER - Teva ✓ Methylphenidate ER 30 - Teva

⇒SA1964 Special Authority for Subsidy

Initial application — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients aged 5 years or over; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Initial application — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

	Subsidy	Fully	
(N	flanufacturer's Price)	Subsidised	Generic
	\$	Per 🗸	Manufacturer

continued...

- 1 ADHD (Attention Deficit and Hyperactivity Disorder) patients under 5 years of age; and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria.

Initial application — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the patient suffers from narcolepsy.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

Renewal — (ADHD in patients 5 or over) only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

Renewal — (ADHD in patients under 5) only from a paediatrician or psychiatrist. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (Narcolepsy*) only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Note: *narcolepsy is not a registered indication for Methylphenidate ER – Teva.

METHYLPHENIDATE HYDROCHLORIDE EXTENDED-RELEASE - Special Authority see SA1965 below - Retail pharmacy

- a) Only on a controlled drug form
- b) Safety medicine; prescriber may determine dispensing frequency

Tab extended-release 18 mg	58.96	30	Concerta
Tab extended-release 27 mg	65.44	30	Concerta
Tab extended-release 36 mg	71.93	30	Concerta
Tab extended-release 54 mg		30	Concerta
Cap modified-release 10 mg		30	Ritalin LA
Cap modified-release 20 mg		30	Ritalin LA
Cap modified-release 30 mg		30	Ritalin LA
Cap modified-release 40 mg		30	Ritalin LA

⇒SA1965 Special Authority for Subsidy

Initial application only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 ADHD (Attention Deficit and Hyperactivity Disorder); and
- 2 Diagnosed according to DSM-IV or ICD 10 criteria; and
- 3 Either:
 - 3.1 Applicant is a paediatrician or psychiatrist; or
 - 3.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing; and
- 4 Either:
 - 4.1 Patient is taking a currently subsidised formulation of methylphenidate hydrochloride (immediate-release or sustained-release) which has not been effective due to significant administration and/or compliance difficulties; or
 - 4.2 There is significant concern regarding the risk of diversion or abuse of immediate-release methylphenidate hydrochloride.

Renewal only from a paediatrician, psychiatrist, medical practitioner on the recommendation of a paediatrician or psychiatrist (in

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

writing) or nurse practitioner on the recommendation of a paediatrician or psychiatrist (in writing). Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 Either:
 - 2.1 Applicant is a paediatrician or psychiatrist; or
 - 2.2 Applicant is a medical practitioner or nurse practitioner and confirms that a paediatrician or psychiatrist has been consulted within the last 2 years and has recommended treatment for the patient in writing.

MODAFINIL - Special Authority see SA1999 below - Retail pharmacy

⇒SA1999 Special Authority for Subsidy

Initial application only from a neurologist or respiratory specialist. Approvals valid for 24 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a diagnosis of narcolepsy and has excessive daytime sleepiness associated with narcolepsy occurring almost daily for three months or more; and
- 2 Either:
 - 2.1 The patient has a multiple sleep latency test with a mean sleep latency of less than or equal to 10 minutes and 2 or more sleep onset rapid eye movement periods; or
 - 2.2 The patient has at least one of: cataplexy, sleep paralysis or hypnagogic hallucinations; and
- 3 Either:
 - 3.1 An effective dose of a subsidised formulation of methylphenidate or dexamfetamine has been trialled and discontinued because of intolerable side effects; or
 - 3.2 Methylphenidate and dexamfetamine are contraindicated.

Renewal only from a neurologist or respiratory specialist. Approvals valid for 24 months where the treatment remains appropriate and the patient is benefiting from treatment.

Treatments for Dementia

DONEPEZIL HYDROCHLORIDE			
* Tab 5 mg	4.34	90	✓ Donepezil-Rex
* Tab 10 mg	6.64	90	✓ Donepezil-Rex
RIVASTIGMINE - Special Authority see SA1488 below - Reta	il pharmacy		
Patch 4.6 mg per 24 hour	38.00	30	✓ <u>Rivastigmine Patch</u> <u>BNM 5</u>
Patch 9.5 mg per 24 hour	38.00	30	✓ Rivastigmine Patch BNM 10

⇒SA1488 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient has been diagnosed with dementia; and
- 2 The patient has experienced intolerable nausea and/or vomiting from donepezil tablets.

Renewal from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The treatment remains appropriate; and
- 2 The patient has demonstrated a significant and sustained benefit from treatment.

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

Treatments for Substance Dependence

BUPRENORPHINE WITH NALOXONE - Special Authority see SA1203 below - Retail pharmacy

- a) No patient co-payment payable
- b) Safety medicine; prescriber may determine dispensing frequency

Tab sublingual 2 mg with naloxone 0.5 mg.......11.76 28

✓ Buprenorphine

Naloxone BNM

Buprenorphine Naloxone BNM to be Principal Supply on 1 December 2022 Tab sublingual 8 mg with naloxone 2 mg......34.00

......34.00 28 Suprenorphine
Naloxone BNM

Buprenorphine Naloxone BNM to be Principal Supply on 1 December 2022

⇒SA1203 Special Authority for Subsidy

Initial application — (Detoxification) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health...

Initial application — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient will not be receiving methadone; and
- 3 Patient is currently enrolled in an opioid substitution treatment program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (**Detoxification**) from any medical practitioner. Approvals valid for 1 month for applications meeting the following criteria:

All of the following:

- 1 Patient is opioid dependent; and
- 2 Patient has previously trialled but failed detoxification with buprenorphine with naloxone with relapse back to opioid use and another attempt is planned; and
- 3 Patient is currently engaged with an opioid treatment service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

Renewal — (Maintenance treatment) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is or has been receiving maintenance therapy with buprenorphine with naloxone (and is not receiving methadone); and
- 2 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 3 Applicant works in an opioid treatment service approved by the Ministry of Health or is a medical practitioner authorised by the service to manage treatment in this patient.

Renewal — (Maintenance treatment where the patient has previously had an initial application for detoxification) from any medical practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient received but failed detoxification with buprenorphine with naloxone; and
- 2 Maintenance therapy with buprenorphine with naloxone is planned (and patient will not be receiving methadone); and
- 3 Patient is currently enrolled in an opioid substitution program in a service approved by the Ministry of Health; and
- 4 Applicant works in an opioid treatment service approved by the Ministry of Health.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer			
BUPROPION HYDROCHLORIDE Tab modified-release 150 mg	11.00	30	✓ <u>Z</u>	yban			
DISULFIRAM							
Tab 200 mg	236.40	100	✓ A	ntabuse S29			
NALTREXONE HYDROCHLORIDE - Special Authority see SA1408 below - Retail pharmacy							
Tab 50 mg	133.33	30	✓ N	altraccord			
- CA1400 Chariel Authority for Cyloidy							

⇒SA1408 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Patient is currently enrolled in a recognised comprehensive treatment programme for alcohol dependence; and
- 2 Applicant works in or with a community Alcohol and Drug Service contracted to Health NZ or accredited against the New Zealand Alcohol and Other Drug Sector Standard or the National Mental Health Sector Standard.

Renewal from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Compliance with the medication (prescriber determined); and
- 2 Any of the following:
 - 2.1 Patient is still unstable and requires further treatment; or
 - 2.2 Patient achieved significant improvement but requires further treatment; or
 - 2.3 Patient is well controlled but requires maintenance therapy.

NICOTINE

a) Nicotine will not be funded in amounts less than 4 weeks of treatment.

b) Note: Direct Provision by a pharmacist permitted under the pro	visions in Part	I of Section A	٨.
Patch 7 mg - Up to 28 patch available on a PSO	.18.14	28	✓ Habitrol
Patch 7 mg for direct distribution only - [Xpharm]	3.94	7	✓ Habitrol
Patch 14 mg - Up to 28 patch available on a PSO	.19.95	28	Habitrol
Patch 14 mg for direct distribution only - [Xpharm]	4.52	7	Habitrol
Patch 21 mg - Up to 28 patch available on a PSO	.22.86	28	Habitrol
Patch 21 mg for direct distribution only - [Xpharm]	5.18	7	Habitrol
Lozenge 1 mg - Up to 216 loz available on a PSO	.19.18	216	Habitrol
Lozenge 1 mg for direct distribution only - [Xpharm]	3.20	36	Habitrol
Lozenge 2 mg - Up to 216 loz available on a PSO	.21.02	216	Habitrol
Lozenge 2 mg for direct distribution only - [Xpharm]	3.24	36	Habitrol
Gum 2 mg (Fruit) - Up to 384 piece available on a PSO		384	Habitrol
Gum 2 mg (Fruit) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 2 mg (Mint) - Up to 384 piece available on a PSO	.38.21	384	Habitrol
Gum 2 mg (Mint) for direct distribution only - [Xpharm]	8.64	96	Habitrol
Gum 4 mg (Fruit) - Up to 384 piece available on a PSO		384	Habitrol
Gum 4 mg (Fruit) for direct distribution only - [Xpharm]	.10.01	96	Habitrol
Gum 4 mg (Mint) – Up to 384 piece available on a PSO	.44.17	384	Habitrol
Gum 4 mg (Mint) for direct distribution only - [Xpharm]	.10.01	96	Habitrol

VARENICLINE TARTRATE - Special Authority see SA1845 on the next page - Retail pharmacy

- a) A maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval, including the starter pack
- b) Varenicline will not be funded in amounts less than 4 weeks of treatment.
- c) The 6-month time period in which a patient can receive a funded 12-week course of varenicline tartrate starts from the date the Special Authority is approved.

Tab 0.5 mg × 11 and 1 mg × 42	16.67	53 OP	✓ Varenicline Pfizer
Tab 1 mg	17.62	56	✓ Varenicline Pfizer

Subsidy		Fully	Brand or	_
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

⇒SA1845 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 Either:
 - 3.1 The patient has tried but failed to quit smoking after at least two separate trials of nicotine replacement therapy, at least one of which included the patient receiving comprehensive advice on the optimal use of nicotine replacement therapy; or
 - 3.2 The patient has tried but failed to quit smoking using bupropion or nortriptyline; and
- 4 The patient has not had a Special Authority for varenicline approved in the last 6 months; and
- 5 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 6 The patient is not pregnant; and
- 7 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

Renewal from any relevant practitioner. Approvals valid for 5 months for applications meeting the following criteria: All of the following:

- 1 Short-term therapy as an aid to achieving abstinence in a patient who has indicated that they are ready to cease smoking; and
- 2 The patient is part of, or is about to enrol in, a comprehensive support and counselling smoking cessation programme, which includes prescriber or nurse monitoring; and
- 3 It has been 6 months since the patient's previous Special Authority was approved; and
- 4 Varenicline is not to be used in combination with other pharmacological smoking cessation treatments and the patient has agreed to this; and
- 5 The patient is not pregnant; and
- 6 The patient will not be prescribed more than 12 weeks' funded varenicline (see note).

The patient must not have had an approval in the past 6 months.

Notes: a maximum of 12 weeks' varenicline will be subsidised on each Special Authority approval.

This includes the 4-week 'starter' pack.

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer

Chemotherapeutic Agents

Alkylating Agents

BENDAMUSTINE HYDROCHI ORIDE - PCT only - Specialist - Special Authority see SA2153 below

Inj 25 mg vial	 	77.00	1	✓ Ribomustin
Inj 100 mg vial	 	308.00	1	✓ Ribomustin
Inj 1 mg for ECP	 	3.23	1 mg	✓ Baxter

⇒SA2153 Special Authority for Subsidy

Initial application — (treatment naive CLL) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has Binet stage B or C, or progressive stage A chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is chemotherapy treatment naive; and
- 3 The patient is unable to tolerate toxicity of full-dose FCR; and
- 4 Patient has ECOG performance status 0-2; and
- 5 Patient has a Cumulative Illness Rating Scale (CIRS) score of < 6: and
- 6 Bendamustine is to be administered at a maximum dose of 100 mg/m² on days 1 and 2 every 4 weeks for a maximum of 6 cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL). Chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 The patient has indolent low grade NHL requiring treatment; and
- 2 Patient has a WHO performance status of 0-2; and
- 3 Any of the following:
 - 3.1 Both:
 - 3.1.1 Patient is treatment naive; and
 - 3.1.2 Bendamustine is to be administered for a maximum of 6 cycles (in combination with rituximab when CD20+); or
 - 3.2 Both:
 - 3.2.1 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen; and
 - 3.2.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or
 - 3.3 All of the following:
 - 3.3.1 The patient has not received prior bendamustine therapy; and
 - 3.3.2 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 3.3.3 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 3.4 Bendamustine is to be administered as monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Renewal — (Indolent, Low-grade lymphomas) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: Fither:

- 1 Both:
 - 1.1 Patient is refractory to or has relapsed within 12 months of rituximab in combination with bendamustine; and
 - 1.2 Bendamustine is to be administered in combination with obinutuzumab for a maximum of 6 cycles; or

	Subsidy acturer's Price) Subs	Fully	Brand or Generic	
(wanua	\$ Per	uiseu •	Manufacturer	

continued...

- 2 Both:
 - 2.1 Patients have not received a bendamustine regimen within the last 12 months; and
 - 2.2 Either:
 - 2.2.1 Both:
 - 2.2.1.1 Bendamustine is to be administered for a maximum of 6 cycles in relapsed patients (in combination with rituximab when CD20+); and
 - 2.2.1.2 Patient has had a rituximab treatment-free interval of 12 months or more; or
 - 2.2.2 Bendamustine is to be administered as a monotherapy for a maximum of 6 cycles in rituximab refractory patients.

Note: 'indolent, low-grade lymphomas' includes follicular, mantle cell, marginal zone and lymphoplasmacytic/ Waldenstrom's macroglobulinaemia.

Initial application — (Hodgkin's lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

- All of the following:
 - 1 Patient has Hodgkin's lymphoma requiring treatment; and
 - 2 Patient has a ECOG performance status of 0-2; and
 - 3 Patient has received one prior line of chemotherapy; and
 - 4 Patient's disease relapsed or was refractory following prior chemotherapy; and
 - 5 Bendamustine is to be administered in combination with gemcitabine and vinorelbine (BeGeV) at a maximum dose of no greater than 90 mg/m2 twice per cycle, for a maximum of four cycles.

Note: Indications marked with * are unapproved indications.

DUCULEAN DCT Patail pharmany Chanielist

BUSULFAN - PCT - Retail pharmacy-Specialist	100	✓ Myleran
Tab 2 mg	100	♥ Wylcian
CARBOPLATIN – PCT only – Specialist		C DDL Oanhandatin
Inj 10 mg per ml, 45 ml vial	1	✓ DBL Carboplatin
45.20		✓ Carboplatin Ebewe
48.50	4	✓ Carbaccord
Inj 1 mg for ECP0.10	1 mg	✓ Baxter
CARMUSTINE - PCT only - Specialist		
Inj 100 mg vial710.00	1	✓ <u>BiCNU</u>
Inj 100 mg for ECP710.00	100 mg OP	✓ Baxter
CHLORAMBUCIL - PCT - Retail pharmacy-Specialist		
Tab 2 mg29.06	25	✓ Leukeran FC
CISPLATIN - PCT only - Specialist		
Inj 1 mg per ml, 50 ml vial15.00	1	✓ Cisplatin Ebewe
Inj 1 mg per ml, 100 ml vial21.00	1	✓ Cisplatin Ebewe
29.66	'	✓ DBL Cisplatin
Inj 1 mg for ECP	1 mg	✓ Baxter
CYCLOPHOSPHAMIDE	9	
	50	√ Cycloney
Tab 50 mg - PCT - Retail pharmacy-Specialist	1	✓ <u>Cyclonex</u> ✓ Endoxan
127.80	6	✓ Cytoxan
1-11-1	1	✓ Endoxan
Inj 2 g vial – PCT only – Specialist	1 mg	✓ Baxter
, ,	ring	Daxiei
IFOSFAMIDE – PCT only – Specialist		
lnj 1 g96.00	1	✓ Holoxan
lnj 2 g	. 1	✓ Holoxan
Inj 1 mg for ECP0.10	1 mg	✓ Baxter

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	
LOMUSTINE - PCT - Retail pharmacy-Specialist	Ψ	1 01		Managadio
Cap 10 mg	132.50	20	1	CeeNU
Cap 40 mg		20		CeeNU
			•	Occino
MELPHALAN	40.70	0.5		Alkeran
Tab 2 mg - PCT - Retail pharmacy-Specialist		25		
Inj 50 mg - PCT only - Specialist	67.80	1		Alkeran
			•	Alkeran S29 S29
OXALIPLATIN - PCT only - Specialist				
Inj 100 mg vial	25.01	1	•	Oxaliplatin Actavis 100
	110.00		1	Oxaliplatin Ebewe
Inj 5 mg per ml, 20 ml vial	46.32	1	/	Oxaliplatin Accord
Inj 1 mg for ECP		1 mg	_	Baxter
THIOTEPA - PCT only - Specialist		·	•	
Inj 15 mg vial	CBS	1	/	Bedford S29
, · ··-g			/	Max Health S29
				THIO-TEPA S29
	000			Tepadina S29
Inj 100 mg vial	CBS	1	•	Max Health S29
			✓	Tepadina S29

Antimetabolites

AZACITIDINE - PCT only - Specialist - Special Authority see SA214	41 below		
Inj 100 mg vial	75.06	1	✓ Azacitidine Dr
			Reddy's
Inj 1 mg for ECP	0.83	1 mg	✓ Baxter

⇒SA2141 Special Authority for Subsidy

Initial application only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has International Prognostic Scoring System (IPSS) intermediate-2 or high risk myelodysplastic syndrome; or
 - 1.2 The patient has chronic myelomonocytic leukaemia (10%-29% marrow blasts without myeloproliferative disorder); or
 - 1.3 The patient has acute myeloid leukaemia with 20-30% blasts and multi-lineage dysplasia, according to World Health Organisation Classification (WHO); and
- 2 The patient has performance status (WHO/ECOG) grade 0-2; and
- 3 The patient has an estimated life expectancy of at least 3 months.

Renewal only from a haematologist or medical practitioner on the recommendation of a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

	Subsidy		Fully Brand or
	(Manufacturer's Pri	ice) Subs Per	idised Generic Manufacturer
CALCIUM FOLINATE			manadator
Tab 15 mg - PCT - Retail pharmacy-Specialist	114 60	10	✓ DBL Leucovorin
Tab 13 mg - 1 01 - Hetali phamacy-opecialist	114.03	10	Calcium
Inj 3 mg per ml, 1 ml - PCT - Retail pharmacy-Specialist	17.10	5	✓ Hospira
Inj 10 mg per ml, 5 ml vial - PCT - Retail pharmacy-Speciali		1	✓ Calcium Folinate
			Sandoz
			Calcium Folinate
			Sandoz S29 S29
Inj 50 mg - PCT - Retail pharmacy-Specialist	72.80	10	✓ Leucovorin
			Pharmacia \$29
Inj 10 mg per ml, 10 ml vial - PCT only - Specialist	9.49	1	✓ Calcium Folinate
lai 100 ann DOT anh Chaoinlint	7.00		Sandoz
Inj 100 mg - PCT only - Specialist	1.33	1	✓ Calcium Folinate Ebewe
	94.90	10	✓ Leucovorin
	01.00	10	Pharmacia \$29
Inj 300 mg - PCT only - Specialist	22.51	1	✓ Calcium Folinate
ng ood ing it or only openation in the control of t		·	Ebewe
	25.14		✓ Leucovorin DBL S29
Litto Los Lil BOT L O LIN	05.44		
Inj 10 mg per ml, 35 ml vial – PCT only – Specialist	25.14	1	✓ Calcium Folinate Sandoz
			✓ Calcium Folinate
			Sandoz S29 S29
Inj 1 g - PCT only - Specialist	67 51	1	✓ Calcium Folinate
ing i g i or only openialist			Ebewe
Inj 10 mg per ml, 100 ml vial - PCT only - Specialist	72.00	1	✓ Calcium Folinate
, .			Sandoz
Inj 1 mg for ECP - PCT only - Specialist	0.06	1 mg	✓ Baxter
CAPECITABINE - Retail pharmacy-Specialist			
Tab 150 mg		60	✓ Capercit
Tab 500 mg	49.00	120	✓ Capercit
CLADRIBINE – PCT only – Specialist			
Inj 2 mg per ml, 5 ml		1	✓ Litak S29
Inj 1 mg per ml, 10 ml		1	✓ Leustatin
Inj 10 mg for ECP	/49.96	10 mg OP	✓ Baxter
CYTARABINE	int 400.00	5	✓ Pfizer
Inj 20 mg per ml, 5 ml vial – PCT – Retail pharmacy-Speciali Inj 100 mg per ml, 20 ml vial – PCT – Retail	ist 400.00	5	▼ Pilzer
pharmacy-Specialist	41.36	1	✓ Pfizer
Inj 1 mg for ECP - PCT only - Specialist		10 mg	✓ Baxter
Inj 100 mg intrathecal syringe for ECP - PCT only - Speciali		100 mg OP	✓ Baxter
FLUDARABINE PHOSPHATE			
Tab 10 mg - PCT - Retail pharmacy-Specialist		20	✓ Fludara Oral
Inj 50 mg vial – PCT only – Specialist		5	✓ Fludarabine Ebewe
Inj 50 mg for ECP - PCT only - Specialist	126.80	50 mg OP	✓ Baxter

	Subsidy (Manufacturer's Pri	ica) Subs	Fully	
	\$	Per	olui3eu √	Manufacturer
FLUOROURACIL				
Inj 50 mg per ml, 20 ml vial - PCT only - Specialist	10.51	1	1	Fluorouracil Accord
Inj 50 mg per ml, 100 ml vial - PCT only - Specialist	29.44	1	1	Fluorouracil Accord
Inj 1 mg for ECP - PCT only - Specialist	0.62	100 mg	1	Baxter
GEMCITABINE HYDROCHLORIDE - PCT only - Specialist				
Inj 1 g, 26.3 ml vial	62.50	1	1	DBL Gemcitabine
Inj 1 g		1	1	Gemcitabine Ebewe
Inj 1 mg for ECP	0.02	1 mg	1	Baxter
IRINOTECAN HYDROCHLORIDE - PCT only - Specialist				
Inj 20 mg per ml, 5 ml vial	52.57	1	1	Accord
	71.44		1	Irinotecan Actavis 100
	100.00		1	Irinotecan-Rex
Inj 1 mg for ECP	0.54	1 mg	1	Baxter
MERCAPTOPURINE				
Tab 50 mg - PCT - Retail pharmacy-Specialist Puri-nethol to be Principal Supply on 1 December 2022	25.90	25	•	Puri-nethol
Oral suspension 20 mg per ml - Retail pharmacy-Specialist	_			
Special Authority see SA1725 below	428.00	100 ml OP	1	Allmercap

⇒SA1725 Special Authority for Subsidy

Initial application only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where the patient requires a total dose of less than one full 50 mg tablet per day.

Renewal only from a paediatric haematologist or paediatric oncologist. Approvals valid for 12 months where patient still requires a total dose of less than one full 50 mg tablet per day.

METHOTREXATE

ME	THOTREXATE		
*	Tab 2.5 mg - PCT - Retail pharmacy-Specialist9.98	90	✓ Trexate
*	Tab 10 mg - PCT - Retail pharmacy-Specialist33.71	90	✓ Trexate
*	Inj 2.5 mg per ml, 2 ml - PCT - Retail pharmacy-Specialist47.50	5	✓ Methotrexate DBL
*		1	Methotrexate Sandoz
*	Inj 10 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 15 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 20 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 30 mg prefilled syringe	1	Methotrexate Sandoz
*	Inj 25 mg per ml, 2 ml vial - PCT - Retail pharmacy-Specialist30.00	5	Methotrexate DBL Onco-Vial
*	Inj 25 mg per ml, 20 ml vial - PCT - Retail pharmacy-Specialist45.00	1	✓ DBL Methotrexate Onco-Vial
*	Inj 100 mg per ml, 10 ml - PCT - Retail pharmacy-Specialist25.00	1	✓ Methotrexate Ebewe
*	Inj 100 mg per ml, 50 ml vial – PCT – Retail		
	pharmacy-Specialist79.99	1	✓ Methotrexate Ebewe
*	Inj 1 mg for ECP - PCT only - Specialist	1 mg	✓ Baxter
*	Inj 5 mg intrathecal syringe for ECP - PCT only - Specialist4.73	5 mg ÖP	✓ Baxter

	Subsidy (Manufacturer's Price)		Fully Subsidised	
	\$	Per	•	Manufacturer
PEMETREXED - PCT only - Specialist - Special Authority see	SA1679 below			
Inj 100 mg vial	60.89	1	✓	Juno Pemetrexed
Inj 500 mg vial	217.77	1	✓	Juno Pemetrexed
Inj 1 mg for ECP	0.55	1 mg	✓	Baxter

⇒SA1679 Special Authority for Subsidy

Initial application — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

Roth:

- 1 Patient has been diagnosed with mesothelioma; and
- 2 Pemetrexed to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles.

Renewal — (mesothelioma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

All of the following:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed to be administered at a dose of 500mg/m² every 21 days for a maximum of 6 cycles.

Initial application — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria: Both:

- 1 Patient has locally advanced or metastatic non-squamous non-small cell lung carcinoma; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has chemotherapy-naïve disease; and
 - 2.1.2 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days in combination with cisplatin or carboplatin for a maximum of 6 cycles; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had first-line treatment with platinum based chemotherapy; and
 - 2.2.2 Patient has not received prior funded treatment with pemetrexed; and
 - 2.2.3 Pemetrexed is to be administered at a dose of 500 mg/m² every 21 days for a maximum of 6 cycles.

Renewal — (non-small cell lung carcinoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 8 months for applications meeting the following criteria:

- All of the following:
 - 1 No evidence of disease progression; and

THIOGUANINE - PCT - Retail pharmacy-Specialist

- 2 The treatment remains appropriate and the patient is benefitting from treatment; and
- 3 Pemetrexed is to be administered at a dose of 500mg/m² every 21 days.

Tab 40 mg	126.31	25	Lanvis
Other Cytotoxic Agents			
AMSACRINE - PCT only - Specialist			
Inj 50 mg per ml, 1.5 ml ampoule	1,500.00	6	✓ Amsidine S29
	4,736.00		✓ Amsidine S29
Inj 75 mg	1,250.00	5	✓ AmsaLyo S29
ANAGRELIDE HYDROCHLORIDE - PCT - Retail pharm.	acy-Specialist		
Cap 0.5 mg		100	✓ Agrylin
ARSENIC TRIOXIDE - PCT only - Specialist			
Inj 1 mg per ml, 10 ml vial	4,817.00	10	✓ Phenasen
Inj 10 mg for ECP	481.70	10 mg OP	✓ Baxter

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Price	ce) Sub	Fully	Brand or Generic	
	\$	Per	1	Manufacturer	
BLEOMYCIN SULPHATE - PCT only - Specialist					
Inj 15,000 iu, vial	185.16	1	✓ [DBL Bleomycin	
				Sulfate	
Inj 1,000 iu for ECP	14.32	1,000 iu	✓ [Baxter	
BORTEZOMIB - PCT only - Specialist - Special Authority see S	SA1889 below				
Inj 3.5 mg vial		1	✓ [Bortezomib	
•				Dr-Reddy's	
Inj 1 mg for ECP	31.20	1 mg	✓ [Baxter	

⇒SA1889 Special Authority for Subsidy

Initial application — (multiple myeloma/amyloidosis) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has symptomatic multiple myeloma; or
- 2 The patient has symptomatic systemic AL amyloidosis *.

Note: Indications marked with * are unapproved indications.

DACARBAZINE - PCT only - Specialist			
Inj 200 mg vial	62.70	1	 DBL Dacarbazine
	580.60	10	✓ Dacarbazine APP \$29
Inj 200 mg for ECP	62.70	200 mg OP	✓ Baxter
DACTINOMYCIN [ACTINOMYCIN D] - PCT only - Specialist			
Inj 0.5 mg vial	255.00	1	✓ Cosmegen
Inj 0.5 mg for ECP		0.5 mg OP	✓ Baxter
DAUNORUBICIN - PCT only - Specialist		-	
Inj 2 mg per ml, 10 ml	149.50	1	✓ Pfizer
Inj 20 mg vial		10	✓ Daunorubicin
, - 3	,		Zentiva S29
Inj 20 mg for ECP	149.50	20 mg OP	✓ Baxter
DOCETAXEL - PCT only - Specialist		3 -	
Inj 20 mg	48 75	1	✓ Docetaxel Sandoz
Inj 10 mg per ml, 8 ml vial		1	✓ DBL Docetaxel
Inj 20 mg per ml, 4 ml vial		1	✓ Docetaxel
ing 20 mg per mi, 4 mi viai	20.33	'	Accord \$29
Inj 80 mg	195.00	1	✓ Docetaxel Sandoz
Inj 1 mg for ECP	0.65	1 mg	✓ Baxter
DOXORUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	10.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Doxorubicin Ebewe
, 3, 4	17.00		✓ Arrow-Doxorubicin
Inj 2 mg per ml, 50 ml vial	23.00	1	✓ Doxorubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Arrow-Doxorubicin
, 01	69.99		Doxorubicin Ebewe
Inj 1 mg for ECP	0.35	1 mg	✓ Baxter
EPIRUBICIN HYDROCHLORIDE - PCT only - Specialist			
Inj 2 mg per ml, 5 ml vial	25.00	1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 25 ml vial		1	✓ Epirubicin Ebewe
Inj 2 mg per ml, 100 ml vial		1	✓ Epirubicin Ebewe
Inj 1 mg for ECP		1 mg	✓ Baxter

	Subsidy		Fully	
	(Manufacturer's Price)	Subsid		
	<u> </u>	Per		Manufacturer
ETOPOSIDE				
Cap 50 mg - PCT - Retail pharmacy-Specialist	340.73	20	1	Vepesid
Cap 100 mg - PCT - Retail pharmacy-Specialist		10	1	Vepesid
Inj 20 mg per ml, 5 ml vial - PCT - Retail pharmacy-Specialis	t7.90	1	1	Rex Medical
Inj 1 mg for ECP - PCT only - Specialist	0.09	1 mg	1	Baxter
ETOPOSIDE PHOSPHATE - PCT only - Specialist		•		
Inj 100 mg (of etoposide base)	40.00	1	1	Etopophos
Inj 1 mg (of etoposide base) for ECP		1 mg		Baxter
, , ,		9		
HYDROXYUREA [HYDROXYCARBAMIDE] – PCT – Retail pharm		100	./	Devetie
Cap 500 mg	23.82	100	٧	Devatis
IDARUBICIN HYDROCHLORIDE				
Inj 5 mg vial – PCT only – Specialist	109.74	1	1	Zavedos
Inj 10 mg vial - PCT only - Specialist	233.64	1	✓	Zavedos
Inj 1 mg for ECP - PCT only - Specialist	25.77	1 mg	1	Baxter
LENALIDOMIDE - Retail pharmacy-Specialist - Special Authority	see SA2047 below			
Wastage claimable				
Cap 5 mg	5.122.76	28	1	Revlimid
Cap 10 mg		21	1	Revlimid
	6.207.00	28		Revlimid
Cap 15 mg	-,	21		Revlimid
	7.239.18	28		Revlimid
Cap 25 mg	,	21		Revlimid
	,527.00		-	

⇒SA2047 Special Authority for Subsidy

Initial application — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has relapsed or refractory multiple myeloma with progressive disease; and
- 2 Patient has not previously been treated with lenalidomide; and
- 3 Either:
 - 3.1 Lenalidomide to be used as third line* treatment for multiple myeloma; or
 - 3.2 Both:
 - 3.2.1 Lenalidomide to be used as second line treatment for multiple myeloma; and
 - 3.2.2 The patient has experienced severe (grade 3 or higher), dose limiting, peripheral neuropathy with either bortezomib or thalidomide that precludes further treatment with either of these treatments; and
- 4 Lenalidomide to be administered at a maximum dose of 25 mg/day in combination with dexamethasone.

Initial application — (Maintenance following first-line autologous stem cell transplant (SCT)) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed symptomatic multiple myeloma and has undergone first-line treatment that included an autologous stem cell transplantation; and
- 2 Patient has at least a stable disease response in the first 100 days after transplantation; and
- 3 Lenalidomide maintenance is to be commenced within 6 months of transplantation; and
- 4 Lenalidomide to be administered at a maximum dose of 15 mg/day.

Renewal — (Relapsed/refractory disease) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

- 1 No evidence of disease progression; and
 - 2 The treatment remains appropriate and patient is benefitting from treatment.

Renewal — (Maintenance following first line autologous SCT) only from a haematologist or any relevant practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and patient is benefitting from treatment.

Note: Indication marked with * is an unapproved indication. A line of treatment is considered to comprise either: a) a known therapeutic chemotherapy regimen and supportive treatments or b) a transplant induction chemotherapy regimen, stem cell transplantation and supportive treatments. Prescriptions must be written by a registered prescriber in the lenalidomide risk management programme operated by the supplier.

MESNA

Tab 400 mg - PCT - Retail pharmacy-Specialist	314.00	50	✓ Uromitexan
Tab 600 mg - PCT - Retail pharmacy-Specialist	448.50	50	✓ Uromitexan
Inj 100 mg per ml, 4 ml ampoule - PCT only - Specialist	177.45	15	✓ Uromitexan
Inj 100 mg per ml, 10 ml ampoule - PCT only - Specialist	407.40	15	✓ Uromitexan
Inj 1 mg for ECP - PCT only - Specialist	2.96	100 mg	✓ Baxter
MITOMYCIN C - PCT only - Specialist			
Inj 5 mg vial	641.70	1	✓ Accord S29
lnj 20 mg vial		1	✓ Teva
, -	3,275.00		✓ Omegapharm S29
Inj 1 mg for ECP	269.85	1 mg	✓ Baxter
(Omegapharm 129 Inj 20 mg vial to be delisted 1 November 2022,)		
MITOZANTRONE - PCT only - Specialist			
Inj 2 mg per ml, 10 ml vial	97.50	1	✓ Mitozantrone Ebewe
Inj 1 mg for ECP	5.51	1 mg	✓ Baxter
OLAPARIB - Retail pharmacy-Specialist - Special Authority see S.	A2148 below		
Tab 100 mg	3,701.00	56	✓ Lynparza
Tab 150 mg		56	✓ Lynparza
- CAO440 One stal Authority for Cubaldy			

⇒SA2148 Special Authority for Subsidy

Initial application — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 Patient is currently on treatment with olaparib and met all remaining criteria (criterion 2) below prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has a high-grade serous* epithelial ovarian, fallopian tube, or primary peritoneal cancer; and
 - 2.2 There is documentation confirming pathogenic germline BRCA1 or BRCA2 gene mutation; and
 - 2.3 Either:
 - 2.3.1 All of the following:
 - 2.3.1.1 Patient has newly diagnosed, advanced disease; and
 - 2.3.1.2 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 2.3.1.3 Patient's disease must have experienced a partial or complete response to the first-line platinum-based regimen; or
 - 2.3.2 All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.3.2.1 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy; and
- 2.3.2.2 Patient has platinum sensitive disease defined as disease progression occurring at least 6 months after the last dose of the penultimate line** of platinum-based chemotherapy; and
- 2.3.2.3 Patient's disease must have experienced a partial or complete response to treatment with the immediately preceding platinum-based regimen; and
- 2.3.2.4 Patient has not previously received funded olaparib treatment; and
- 2.4 Treatment will be commenced within 12 weeks of the patient's last dose of the immediately preceding platinum-based regimen; and
- 2.5 Treatment to be administered as maintenance treatment; and
- 2.6 Treatment not to be administered in combination with other chemotherapy.

Renewal — (Ovarian cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from treatment; and
- 2 Either:
 - 2.1 No evidence of progressive disease; or
 - 2.2 Evidence of residual (not progressive) disease and the patient would continue to benefit from treatment in the clinician's opinion; and
- 3 Treatment to be administered as maintenance treatment; and
- 4 Treatment not to be administered in combination with other chemotherapy; and
- 5 Either:
 - 5.1 Both:
 - 5.1.1 Patient has received one line** of previous treatment with platinum-based chemotherapy; and
 - 5.1.2 Documentation confirming that the patient has been informed and acknowledges that the funded treatment period of olaparib will not be continued beyond 2 years if the patient experiences a complete response to treatment and there is no radiological evidence of disease at 2 years; or
 - 5.2 Patient has received at least two lines** of previous treatment with platinum-based chemotherapy.

Notes: *Note "high-grade serous" includes tumours with high-grade serous features or a high-grade serous component.
**A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

PACLITAXEL - PCT only - Specialist			
Inj 30 mg	47.30	5	✓ Paclitaxel Ebewe
Inj 100 mg	24.00	1	✓ Paclitaxel Ebewe
•	91.67		✓ Paclitaxel Actavis
Inj 150 mg	26.69	1	✓ Paclitaxel Ebewe
	137.50		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 300 mg	44.00	1	✓ Paclitaxel Ebewe
	275.00		✓ Anzatax
			✓ Paclitaxel Actavis
Inj 1 mg for ECP	0.20	1 mg	✓ Baxter
PEGASPARGASE - PCT only - Special Authority se	ee SA1979 on the next page		
Inj 750 iu per ml, 5 ml vial	3,455.00	1	✓ Oncaspar LYO S29

Sul	bsidy Fu	ully Brand or
(Manufacti	urer's Price) Subsidis	sed Generic
	\$ Per	✓ Manufacturer

⇒SA1979 Special Authority for Subsidy

Initial application — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has newly diagnosed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Initial application — (Lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months where the patient has lymphoma requiring L-asparaginase containing protocols (e.g. SMILE).

Renewal — (Acute lymphoblastic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has relapsed acute lymphoblastic leukaemia; and
- 2 Pegaspargase to be used with a contemporary intensive multi-agent chemotherapy treatment protocol.

Inj 10 mg	CBS	1	✓ Nipent S29
PROCARBAZINE HYDROCHLORIDE - PCT - Retail	pharmacy-Specialist		
Cap 50 mg	980.00	50	✓ Natulan S29
TEMOZOLOMIDE - Special Authority see SA1741 be	elow - Retail pharmacy		
Cap 5 mg	9.13	5	✓ Temaccord
Cap 20 mg	16.38	5	✓ Temaccord
	18.30		✓ Apo-Temozolomide
	136.00	14	✓ Accord S29
Cap 100 mg	35.98	5	✓ Temaccord
, ,	40.20		✓ Apo-Temozolomide
	532.00	14	✓ Accord S29
Cap 140 mg	50.12	5	✓ Temaccord
	400.00		✓ Amneal S29
Cap 180 mg	620.00	14	✓ Accord S29
Cap 250 mg	86.34	5	✓ Temaccord
	688.00		✓ Amneal S29

(Accord \$29 Cap 20 mg to be delisted 1 December 2022)

(Accord S29 Cap 100 mg to be delisted 1 December 2022)

(Amneal S29 Cap 140 mg to be delisted 1 December 2022)

(Amneal S29 Cap 250 mg to be delisted 1 December 2022)

⇒SA1741 Special Authority for Subsidy

Initial application — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has newly diagnosed glioblastoma multiforme; or
 - 1.2 Patient has newly diagnosed anaplastic astrocytoma*; and
- 2 Temozolomide is to be (or has been) given concomitantly with radiotherapy; and
- 3 Following concomitant treatment temozolomide is to be used for a maximum of 5 days treatment per cycle, at a maximum dose of 200 mg/m² per day.

Subsidy	Ful	y Brand or	
(Manufacturer's Price)	Subsidise	d Generic	
\$	Per •	 Manufacturer 	

continued...

Initial application — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with metastatic or unresectable well-differentiated neuroendocrine tumour*; and
- 2 Temozolomide is to be given in combination with capecitabine; and
- 3 Temozolomide is to be used in 28 day treatment cycles for a maximum of 5 days treatment per cycle at a maximum dose of 200 mg/m² per day; and
- 4 Temozolomide to be discontinued at disease progression.

Initial application — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 9 months where the patient has relapsed/refractory Ewing's sarcoma.

Renewal — (high grade gliomas) only from a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has glioblastoma multiforme; and
 - 1.2 The treatment remains appropriate and the patient is benefitting from treatment; or
- 2 All of the following:
 - 2.1 Patient has anaplastic astrocytoma*; and
 - 2.2 The treatment remains appropriate and the patient is benefitting from treatment; and
 - 2.3 Adjuvant temozolomide is to be used for a maximum of 24 months.

Renewal — (neuroendocrine tumours) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Renewal — (ewing's sarcoma) only from a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indication marked with a * is an unapproved indication. Temozolomide is not subsidised for the treatment of relapsed high grade glioma.

THALIDOMIDE – Retail pharmacy-Specialist – Special A	uthority see SA1124 below		
Cap 50 mg	378.00	28	Thalomid
Cap 100 mg	756.00	28	Thalomid

⇒SA1124 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 The patient has multiple myeloma; or
- 2 The patient has systemic AL amyloidosis*.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid without further renewal unless notified where the patient has obtained a response from treatment during the initial approval period.

Notes: Prescription must be written by a registered prescriber in the thalidomide risk management programme operated by the supplier.

Maximum dose of 400 mg daily as monotherapy or in a combination therapy regimen.

Indication marked with * is an unapproved indication.

	Subsidy (Manufacturer's Price) \$	S Per	Fully subsidised	Brand or Generic Manufacturer	
TRETINOIN	· · · · · · · · · · · · · · · · · · ·				_
Cap 10 mg - PCT - Retail pharmacy-Specialist	479.50	100	✓ V	/esanoid	
VENETOCLAX - Retail pharmacy-Specialist - Special Authority s	see SA1868 below				
Tab 14×10 mg, 7×50 mg, 21×100 mg	1,771.86	42 OP	✓ V	/enclexta	
Tab 10 mg	95.78	14 OP	✓ V	/enclexta	
Tab 50 mg	239.44	7 OP	✓ V	/enclexta	
Tab 100 mg - Wastage claimable	8,209.41	120	✓ V	/enclexta	

⇒SA1868 Special Authority for Subsidy

Initial application — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 Patient has chronic lymphocytic leukaemia requiring treatment; and
- 2 Patient has received at least one prior therapy for chronic lymphocytic leukaemia; and
- 3 Patient has not previously received funded venetoclax; and
- 4 The patient's disease has relapsed within 36 months of previous treatment; and
- 5 Venetoclax to be used in combination with six 28-day cycles of rituximab commencing after the 5-week dose titration schedule with venetoclax; and
- 6 Patient has an ECOG performance status of 0-2.

Renewal — (relapsed/refractory chronic lymphocytic leukaemia) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment; and
- 2 Venetoclax is to be discontinued after a maximum of 24 months of treatment following the titration schedule unless earlier discontinuation is required due to disease progression or unacceptable toxicity.

Initial application — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has previously untreated chronic lymphocytic leukaemia; and
- 2 There is documentation confirming that patient has 17p deletion by FISH testing or TP53 mutation by sequencing; and
- 3 Patient has an ECOG performance status of 0-2.

Renewal — (previously untreated chronic lymphocytic leukaemia with 17p deletion or TP53 mutation*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where the treatment remains clinically appropriate and the patient is benefitting from and tolerating treatment.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma (SLL)* and B-cell prolymphocytic leukaemia (B-PLL)*. Indications marked with * are Unapproved indications.

VINBLASTINE SULPHATE

Inj 1 mg per ml, 10 ml vial – PCT – Retail pharmacy-Specialist270.37	5	✓ Hospira
Inj 1 mg for ECP - PCT only - Specialist6.00	1 mg	✓ Baxter
VINCRISTINE SULPHATE		
Inj 1 mg per ml, 1 ml vial - PCT - Retail pharmacy-Specialist74.52	5	DBL Vincristine Sulfate
Inj 1 mg per ml, 2 ml vial – PCT – Retail pharmacy-Specialist 102.73	5	DBL Vincristine Sulfate
Ini 1 mg for FCP - PCT only - Specialist 12.60	1 ma	✓ Baxter

	Subsidy (Manufacturer's Price \$) Sı Per	Fully ubsidised	
VINORELBINE - PCT only - Specialist				
Inj 10 mg per ml, 1 ml vial	12.00	1	1	Navelbine
	42.00		/	Vinorelbine Ebewe
Inj 10 mg per ml, 5 ml vial	56.00	1	/	Navelbine
	210.00		/	Vinorelbine Ebewe
	328.65		/	Sagent S29
Inj 1 mg for ECP	1.25	1 mg		Baxter
Inj 50 mg for ECP	328.65	0 mg OP	•	Baxter (Sagent)

Protein-tyrosine Kinase Inhibitors

⇒SA1870 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced, or metastatic, unresectable, non-small cell lung cancer; and
- 2 There is documentation confirming that the patient has an ALK tyrosine kinase gene rearrangement using an appropriate ALK test; and
- 3 Patient has an ECOG performance score of 0-2.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 No evidence of progressive disease according to RECIST criteria; and
- 2 The patient is benefitting from and tolerating treatment.

DASATINIB - Special Authority see SA1805 below - Retail pharmacy

Wastage claimable			
Tab 20 mg	3,774.06	60	✓ Sprycel
Tab 50 mg	6,214.20	60	✓ Sprycel
Tab 70 mg	7,692.58	60	✓ Sprycel

⇒SA1805 Special Authority for Subsidy

Initial application only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 The patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis or accelerated phase; and
 - 1.2 Maximum dose of 140 mg/day; or
- 2 Both:
 - 2.1 The patient has a diagnosis of Philadelphia chromosome-positive acute lymphoid leukaemia (Ph+ ALL); and
 - 2.2 Maximum dose of 140 mg/day; or
- 3 All of the following:
 - 3.1 The patient has a diagnosis of CML in chronic phase; and
 - 3.2 Maximum dose of 100 mg/day; and
 - 3.3 Any of the following:
 - 3.3.1 Patient has documented treatment failure* with imatinib; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 3.3.2 Patient has experienced treatment-limiting toxicity with imatinib precluding further treatment with imatinib; or
- 3.3.3 Patient has high-risk chronic-phase CML defined by the Sokal or EURO scoring system; or
- 3.3.4 Patients is enrolled in the KISS study** and requires dasatinib treatment according to the study protocol.

Renewal only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on dasatinib*; and
- 2 Dasatinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum dasatinib dose of 140 mg/day for accelerated or blast phase CML and Ph+ ALL, and 100 mg/day for chronic phase CML.

Note: *treatment failure for CML as defined by Leukaemia Net Guidelines. **Kinase-Inhibition Study with Sprycel Start-up https://www.cancertrialsnz.ac.nz/kiss/

ERLOTINIB - Retail pharmacy-Specialist - Special A	uthority see SA2115 below		
Tab 100 mg	329.70	30	✓ Alchemy
-	764.00		✓ Tarceva
Tab 150 mg	569.70	30	✓ Alchemy
	1.146.00		✓ Tarceva

(Tarceva Tab 100 mg to be delisted 1 February 2023) (Tarceva Tab 150 mg to be delisted 1 February 2023)

⇒SA2115 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has locally advanced or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 There is documentation confirming that the disease expresses activating mutations of EGFR tyrosine kinase; and
- 3 Fitho
 - 3.1 Patient is treatment naive: or
 - 3.2 Both:
 - 3.2.1 The patient has discontinued gefitinib due to intolerance; and
 - 3.2.2 The cancer did not progress while on gefitinib; and
- 4 Erlotinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Erlotinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

GEFITINIB − Retail pharmacy-Specialist − Special Authority see SA2116 below
Tab 250 mg918.00 30 ✓ Iressa

⇒SA2116 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 Patient has locally advanced, or metastatic, unresectable, non-squamous Non Small Cell Lung Cancer (NSCLC); and
- 2 Either
 - 2.1 Patient is treatment naive: or
 - 2.2 Both:
 - 2.2.1 The patient has discontinued erlotinib due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on erlotinib; and
- 3 There is documentation confirming that disease expresses activating mutations of EGFR tyrosine kinase; and
- 4 Gefitinib is to be given for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months where radiological assessment (preferably including CT scan) indicates NSCLC has not progressed.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Gefitinib to be discontinued at progression; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

IMATINIB MESILATE

Note: The Glivec brand of imatinib mesilate (supplied by Novartis) remains fully subsidised under Special Authority for patients with unresectable and/or metastatic malignant GIST only, see SA1460 in Section B of the Pharmaceutical Schedule.

	Tab 100 mg - [Xpharm] - Special Authority see SA1460		
	below2,400.00	60	✓ Glivec
*	Cap 100 mg58.23	60	✓ Imatinib-Rex
*	Cap 400 mg84.79	30	✓ Imatinib-Rex

⇒SA1460 Special Authority for Subsidy

Special Authority approved by the CML/GIST Co-ordinator

Notes: Application details may be obtained from Pharmac's website <u>schedule.pharmac.govt.nz/SAForms</u>, and prescriptions should be sent to:

The CML/GIST Co-ordinator Phone: (04) 460 4990 Pharmac Facsimile: (04) 916 7571

PO Box 10 254 Email: cmlgistcoordinator@pharmac.govt.nz

Wellington

Special Authority criteria for GIST – access by application

Funded for patients:

- a) With a diagnosis (confirmed by an oncologist) of unresectable and/or metastatic malignant gastrointestinal stromal tumour (GIST).
- b) Maximum dose of 400 mg/day.
- c) Applications to be made and subsequent prescriptions can be written by an oncologist.
- d) Initial and subsequent applications are valid for one year. The re-application criterion is an adequate clinical response to the treatment with imatinib (prescriber determined).

LAPATINIB DITOSYLATE - Special Authority see SA2035 on the next page - Retail pharmacy

Note – no new patients to be initiated on lapatinib ditosylate.

	,	ully Brand or	
(Manufact	turer's Price) Subsidi	sed Generic	
	\$ Per	 Manufac 	turer

⇒SA2035 Special Authority for Subsidy

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on lapatinib; and
- 3 Lapatinib not to be given in combination with trastuzumab; and
- 4 Lapatinib to be discontinued at disease progression.

NILOTINIB - Special Authority see SA1489 below - Retail pharmacy

Wastage claimable

Cap 150 mg	4,680.00	120	✓ Tasigna
Cap 200 mg	6,532.00	120	Tasigna

⇒SA1489 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has a diagnosis of chronic myeloid leukaemia (CML) in blast crisis, accelerated phase, or in chronic phase; and
- 2 Fither
 - 2.1 Patient has documented CML treatment failure* with imatinib: or
 - 2.2 Patient has experienced treatment limiting toxicity with imatinib precluding further treatment with imatinib; and
- 3 Maximum nilotinib dose of 800 mg/day: and
- 4 Subsidised for use as monotherapy only.

Note: *treatment failure as defined by Leukaemia Net Guidelines.

Renewal only from a haematologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Lack of treatment failure while on nilotinib as defined by Leukaemia Net Guidelines; and
- 2 Nilotinib treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Maximum nilotinib dose of 800 mg/day; and
- 4 Subsidised for use as monotherapy only.

PALBOCICLIB - Retail pharmacy-Specialist - Special Authority see SA1894 below

Wastage claimable

Tab 75 mg	4,000.00	21	Ibrance
Tab 100 mg	4,000.00	21	✓ Ibrance
Tab 125 mg	4.000.00	21	✓ Ibrance

⇒SA1894 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has unresectable locally advanced or metastatic breast cancer; and
- 2 There is documentation confirming disease is hormone-receptor positive and HER2-negative; and
- 3 Patient has an ECOG performance score of 0-2; and
- 4 Either:

second or subsequent line setting

- 4.1 Disease has relapsed or progressed during prior endocrine therapy; or
- 4.2 Both:

first line setting

Subsidy		Fully	Brand or
(Manufacturer's Price)	Sı	ubsidised	Generic
 \$	Per	✓	Manufacturer

continued...

- 4.2.1 Patient is amenorrhoeic, either naturally or induced, with endocrine levels consistent with a postmenopausal state: and
- 4.2.2 Either:
 - 4.2.2.1 Patient has not received prior systemic treatment for metastatic disease; or
 - 4.2.2.2 All of the following:
 - 4.2.2.2.1 Patient commenced treatment with palbociclib in combination with an endocrine agent prior to 1 April 2020; and
 - 4.2.2.2.2 Patient has not received prior systemic endocrine treatment for metastatic disease; and
 - 4.2.2.2.3 There is no evidence of progressive disease; and
- 5 Treatment must be used in combination with an endocrine partner.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a Medical oncologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment must be used in combination with an endocrine partner; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains appropriate and the patient is benefitting from treatment.

PAZOPANIB - Special Authority see SA1190 below - Retail pharmacy

Tab 200 mg	1,334.70	30	✓ Votrient
Tab 400 mg	2,669.40	30	✓ Votrient

⇒SA1190 Special Authority for Subsidy

Initial application only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist.

Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic renal cell carcinoma; and
- 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 Both:
- 2.3.1 The patient has discontinued sunitinib within 3 months of starting treatment due to intolerance; and
 - 2.3.2 The cancer did not progress whilst on sunitinib; and
- 3 The patient has good performance status (WHO/ECOG grade 0-2); and
- 4 The disease is of predominant clear cell histology; and

The patient has intermediate or poor prognosis defined as:

- 5 Any of the following:
 - 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
 - 5.2 Haemoglobin level < lower limit of normal; or
 - 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
 - 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
 - 5.5 Karnofsky performance score of less than or equal to 70; or
 - 5.6 2 or more sites of organ metastasis; and
- 6 Pazopanib to be used for a maximum of 3 months.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Pazopanib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6.

	Subsidy (Manufacturer's Price) \$	Per	Fully Subsidised	Brand or Generic Manufacturer
RUXOLITINIB - Special Authority see SA1890 below - Reta Wastage claimable	il pharmacy			
Tab 5 mg	2,500.00	56	√ Ja	kavi
Tab 10mg	5,000.00	56	✓ Ja	ıkavi
Tab 15 mg	5,000.00	56	✓ Ja	ıkavi
Tab 20 mg	5.000.00	56	✓ Ja	kavi

⇒SA1890 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 The patient has primary myelofibrosis or post-polycythemia vera myelofibrosis or post-essential thrombocythemia myelofibrosis; and
- 2 Either:
 - 2.1 A classification of risk of intermediate-2 or high-risk myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS; or
 - 2.2 Both:
 - 2.2.1 A classification of risk of intermediate-1 myelofibrosis according to either the International Prognostic Scoring System (IPSS), Dynamic International Prognostic Scoring System (DIPSS), or the Age-Adjusted DIPSS: and
 - 2.2.2 Patient has severe disease-related symptoms that are resistant, refractory or intolerant to available therapy; and
- 3 A maximum dose of 20 mg twice daily is to be given.

Renewal only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 A maximum dose of 20 mg twice daily is to be given.

SUNITINIB - Special Authority see SA2117 below - Retail pharmacy

Cap 12.5 mg208.38	28	Sunitinib Pfizer
Cap 25 mg416.77	28	✓ Sunitinib Pfizer
Cap 50 mg	28	✓ Sunitinib Pfizer

⇒SA2117 Special Authority for Subsidy

Initial application — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

- All of the following:
 - 1 The patient has metastatic renal cell carcinoma; and
 - 2 Any of the following:
 - 2.1 The patient is treatment naive; or
 - 2.2 The patient has only received prior cytokine treatment; or
 - 2.3 The patient has only received prior treatment with an investigational agent within the confines of a bona fide clinical trial which has Ethics Committee approval; or
 - 2.4 Both:
 - 2.4.1 The patient has discontinued pazopanib within 3 months of starting treatment due to intolerance; and
 - 2.4.2 The cancer did not progress whilst on pazopanib; and
 - 3 The patient has good performance status (WHO/ECOG grade 0-2); and
 - 4 The disease is of predominant clear cell histology; and The patient has intermediate or poor prognosis defined as:
 - 5 Any of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 5.1 Lactate dehydrogenase level > 1.5 times upper limit of normal; or
- 5.2 Haemoglobin level < lower limit of normal; or
- 5.3 Corrected serum calcium level > 10 mg/dL (2.5 mmol/L); or
- 5.4 Interval of < 1 year from original diagnosis to the start of systemic therapy; or
- 5.5 Karnofsky performance score of less than or equal to 70; or
- 5.6 2 or more sites of organ metastasis; and
- 6 Sunitinib to be used for a maximum of 2 cycles.

Initial application — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal tumour (GIST); and
- 2 Either:
 - 2.1 The patient's disease has progressed following treatment with imatinib; or
 - 2.2 The patient has documented treatment-limiting intolerance, or toxicity to, imatinib.

Renewal — (RCC) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

- 1 No evidence of disease progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Notes: Sunitinib treatment should be stopped if disease progresses.

Poor prognosis patients are defined as having at least 3 of criteria 5.1-5.6. Intermediate prognosis patients are defined as having 1 or 2 of criteria 5.1-5.6

Renewal — (GIST) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

The patient has responded to treatment or has stable disease as determined by Choi's modified CT response evaluation criteria as follows:

- 1 Any of the following:
 - 1.1 The patient has had a complete response (disappearance of all lesions and no new lesions); or
 - 1.2 The patient has had a partial response (a decrease in size of 10% or more or decrease in tumour density in Hounsfield Units (HU) of 15% or more on CT and no new lesions and no obvious progression of non measurable disease); or
 - 1.3 The patient has stable disease (does not meet criteria the two above) and does not have progressive disease and no symptomatic deterioration attributed to tumour progression; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: It is recommended that response to treatment be assessed using Choi's modified CT response evaluation criteria (J Clin Oncol, 2007, 25:1753-1759). Progressive disease is defined as either: an increase in tumour size of 10% or more and not meeting criteria of partial response (PR) by tumour density (HU) on CT; or: new lesions, or new intratumoral nodules, or increase in the size of the existing intratumoral nodules.

Renewal — (GIST pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has unresectable or metastatic malignant gastrointestinal stromal (GIST); and
- 2 The patient is clinically benifiting from treatment and continued treatment remains appropriate; and
- 3 Sunitinib is to be discontinued at progression; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Day /	Manufacturer

Endocrine Therapy

For GnRH ANALOGUES - refer to HORMONE PREPARATIONS, Trophic Hormones, page 84

ABIRATERONE ACETATE – Retail pharmacy-Specialist – Special Authority see SA2118 below Wastage claimable

⇒SA2118 Special Authority for Subsidy

Initial application only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has prostate cancer; and
- 2 Patient has metastases; and
- 3 Patient's disease is castration resistant; and
- 4 Either:
 - 4.1 All of the following:
 - 4.1.1 Patient is symptomatic; and
 - 4.1.2 Patient has disease progression (rising serum PSA) after second line anti-androgen therapy; and
 - 4.1.3 Patient has ECOG performance score of 0-1; and
 - 4.1.4 Patient has not had prior treatment with taxane chemotherapy; or
 - 4.2 All of the following:
 - 4.2.1 Patient's disease has progressed following prior chemotherapy containing a taxane; and
 - 4.2.2 Patient has ECOG performance score of 0-2; and
 - 4.2.3 Patient has not had prior treatment with abiraterone.

Renewal — (abiraterone acetate) only from a medical oncologist, radiation oncologist, urologist or medical practitioner on the recommendation of a medical oncologist, radiation oncologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Significant decrease in serum PSA from baseline; and
- 2 No evidence of clinical disease progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The treatment remains appropriate and the patient is benefiting from treatment.

Renewal — (pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 2 Abiraterone acetate to be discontinued at progression; and
- 3 No initiation of taxane chemotherapy with abiraterone; and
- 4 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Tab 50 mg	4.21	28	✓ Binarex
FLUTAMIDE			
Tab 250 mg	107.55	90	✓ Prostacur S29
Ÿ	119.50	100	✓ Flutamin
FULVESTRANT - Retail pharmacy-Specialist - Special Aut	hority see SA1895 on	the next pag	e
Ini 50 mg per ml. 5 ml prefilled syringe	,	, ,	

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
•	Por 🗸	Manufacturer

⇒SA1895 Special Authority for Subsidy

Initial application only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has oestrogen-receptor positive locally advanced or metastatic breast cancer; and
- 2 Patient has disease progression following prior treatment with an aromatase inhibitor or tamoxifen for their locally advanced or metastatic disease; and
- 3 Treatment to be given at a dose of 500 mg monthly following loading doses; and
- 4 Treatment to be discontinued at disease progression.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains appropriate and patient is benefitting from treatment; and
- 2 Treatment to be given at a dose of 500 mg monthly; and
- 3 There is no evidence of disease progression.

MEGESTROL ACETATE - Subsidy by endorsement

Subsidy by endorsement – Subsidised for patients who were taking megestrol acetate prior to 1 August 2021 and the prescription is endorsed accordingly. Pharmacists may annotate the prescription as endorsed where there exists a record of prior dispensing of megestrol acetate.

Tab 160 mg	48.80	30	✓ Megace S29
(Megace S29 Tab 160 mg to be delisted 1 February 202	23)		
OCTREOTIDE			
Inj 50 mcg per ml, 1 ml ampoule	27.58	5	✓ Max Health
			✓ Octreotide GH S29
Inj 100 mcg per ml, 1 ml ampoule	32.71	5	✓ Max Health
Inj 500 mcg per ml, 1 ml ampoule	113.10	5	✓ Max Health
OCTREOTIDE LONG-ACTING - Special Authority see \$	SA2119 below – Retail pha	armacy	
Inj depot 10 mg prefilled syringe	439.97	ĺ	✓ Octreotide Depot
			Teva
Inj depot 20 mg prefilled syringe	647.03	1	✓ Octreotide Depot
			<u>Teva</u>
Inj depot 30 mg prefilled syringe	718.55	1	✓ Octreotide Depot
			<u>Teva</u>

⇒SA2119 Special Authority for Subsidy

Initial application — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 2 months for applications meeting the following criteria:

All of the following:

- 1 The patient has nausea* and vomiting* due to malignant bowel obstruction*; and
- 2 Treatment with antiemetics, rehydration, antimuscarinic agents, corticosteroids and analgesics for at least 48 hours has failed; and
- 3 Octreotide to be given at a maximum dose 1500 mcg daily for up to 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Malignant Bowel Obstruction) from any relevant practitioner. Approvals valid for 3 months where the treatment remains appropriate and the patient is benefiting from treatment.

Initial application — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 3 months for applications meeting the following criteria:

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidis	sed	Generic
	Per	1	Manufacturer

continued...

Both:

- 1 The patient has acromegaly; and
- 2 Any of the following:
 - 2.1 Treatment with surgery, radiotherapy and a dopamine agonist has failed; or
 - 2.2 Treatment with octreotide is for an interim period while awaiting the effects of radiotherapy and a dopamine agonist has failed: or
 - 2.3 The patient is unwilling, or unable, to undergo surgery and/or radiotherapy.

Renewal — (Acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 IGF1 levels have decreased since starting octreotide; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment.

Note: In patients with Acromegaly octreotide treatment should be discontinued if IGF1 levels have not decreased after 3 months treatment. In patients treated with radiotherapy octreotide treatment should be withdrawn every 2 years, for 1 month, for assessment of remission. Octreotide treatment should be stopped where there is biochemical evidence of remission (normal IGF1 levels) following octreotide treatment withdrawal for at least 4 weeks

Renewal — (Acromegaly - pandemic circumstances) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has acromegaly; and
- 2 The patient is clinically benefiting from treatment and continued treatment remains appropriate; and
- 3 The regular Special Authority renewal requirements cannot be met due to COVID-19 constraints on the health sector.

Initial application — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:
Any of the following:

- 1 VIPomas and Glucagonomas for patients who are seriously ill in order to improve their clinical state prior to definitive surgery; or
- 2 Both:
 - 2.1 Gastrinoma; and
 - 2.2 Fither:
 - 2.2.1 Patient has failed surgery; or
 - 2.2.2 Patient in metastatic disease after H2 antagonists (or proton pump inhibitors) have failed; or
- 3 Both:
 - 3.1 Insulinomas; and
 - 3.2 Surgery is contraindicated or has failed: or
- 4 For pre-operative control of hypoglycaemia and for maintenance therapy; or
- 5 Both:
 - 5.1 Carcinoid syndrome (diagnosed by tissue pathology and/or urinary 5HIAA analysis); and
 - 5.2 Disabling symptoms not controlled by maximal medical therapy.

Note: The use of octreotide in patients with fistulae, oesophageal varices, miscellaneous diarrhoea and hypotension will not be funded as a Special Authority item

Renewal — (Other Indications) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment. Initial application — (pre-operative acromegaly) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

1 Patient has acromegaly; and

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS					
	Subsidy (Manufacturer's Price)	Sub:	Fully sidised	Brand or Generic Manufacturer	
continued					
2 Patient has a large pituitary tumour, greater than 10 mm a3 Patient is scheduled to undergo pituitary surgery in the ne					
TAMOXIFEN CITRATE					
* Tab 10 mg		60	_	amoxifen Sandoz	
* Tab 20 mg	6.65	60	✓ <u>I</u>	amoxifen Sandoz	
Aromatase Inhibitors					
NASTROZOLE					
₭ Tab 1 mg	4.55	30	✓ <u>F</u>	<u>Inatrole</u>	
EXEMESTANE	44.50	00		NG	
≰ Tab 25 mg	14.50	30	•	fizer Exemestane	
.ETROZOLE ★ Tab 2.5 mg	E 0.4	30	./ 1	.etrole	
r 1ab 2.5 mg	5.04	30	<u> </u>	<u>.etrole</u>	
Immunosuppressants					
Cytotoxic Immunosuppressants					
ZATHIOPRINE					
₭ Tab 25 mg		60	-	zamun	
★ Tab 50 mg		100		zamun	
k Inj 50 mg vial	199.00	1	✓ II	muran	
Imuran Inj 50 mg vial to be delisted 1 January 2023)					
MYCOPHENOLATE MOFETIL	25.00	50	10	`allaant	
Tab 500 mg Cap 250 mg		100		Cellcept Cellcept	
Powder for oral liq 1 g per 5 ml – Subsidy by endorsement		35 ml OP		Cellcept	
Mycophenolate powder for oral liquid is subsidised only				•	

Fusion Proteins

ETANERCEPT - Special Authority see SA2103 below -	- Retail pharmacy		
Inj 25 mg	690.00	4	Enbrel
Inj 25 mg autoinjector	690.00	4	✓ Enbrel
Inj 50 mg autoinjector	1,050.00	4	✓ Enbrel
Inj 50 mg prefilled syringe		4	✓ Enbrel

⇒SA2103 Special Authority for Subsidy

the prescription is endorsed accordingly.

Initial application — (adult-onset Still's disease) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Either:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and

Subsidy	,	Fully	Brand or	
(Manufacturer's Price	·) :	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or tocilizumab; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or tocilizumab such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg. non-steroidal anti-inflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 The patient has a sustained improvement in inflammatory markers and functional status.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for ankylosing spondylitis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for ankylosing spondylitis: or
- 2 All of the following:
 - 2.1 Patient has a confirmed diagnosis of ankylosing spondylitis present for more than six months; and
 - 2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.3 Patient has bilateral sacroillitis demonstrated by plain radiographs, CT or MRI scan; and
 - 2.4 Patient's ankylosing spondylitis has not responded adequately to treatment with two or more non-steroidal anti-inflammatory drugs (NSAIDs), in combination with anti-ulcer therapy if indicated, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
 - 2.5 Either:
 - 2.5.1 Patient has limitation of motion of the lumbar spine in the sacittal and the frontal planes as determined by the following Bath Ankylosing Spondylitis Metrology Index (BASMI) measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender (see Notes); and
 - 2.6 A Bath Ankylosing Spondylitis Disease Activity Index (BASDAI) of at least 6 on a 0-10 scale.

Notes: The BASDAI must have been determined at the completion of the 3 month exercise trial, but prior to ceasing NSAID treatment. The BASDAI measure must be no more than 1 month old at the time of initial application.

Average normal chest expansion corrected for age and gender:

18-24 years - Male: 7.0 cm; Female: 5.5 cm 25-34 years - Male: 7.5 cm; Female: 5.5 cm 35-44 years - Male: 6.5 cm; Female: 4.5 cm

Subsidy (Manufacturer's Price) Sub \$ Per

Fully Subsidised Brand or Generic Manufacturer

continued...

45-54 years - Male: 6.0 cm; Female: 5.0 cm 55-64 years - Male: 5.5 cm; Female: 4.0 cm 65-74 years - Male: 4.0 cm; Female: 4.0 cm 75+ years - Male: 3.0 cm; Female: 2.5 cm

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Following 12 weeks' initial treatment and for subsequent renewals, treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 3 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 4 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (polyarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for polyarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose): or
 - 2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (polyarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	1	

continued...

Initial application — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for oligoarticular course juvenile idiopathic arthritis (JIA): and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for oligoarticular course JIA; or
- 2 All of the following:
 - 2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.3 Any of the following:
 - 2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.3.3 High disease activity (cJADAS10 score greater than 4) after a 6-month trial of methotrexate.

Renewal — (oligoarticular course juvenile idiopathic arthritis) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Subsidised as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both
 - 1.1 The patient has had an initial Special Authority approval for adalimumab or secukinumab for psoriatic arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab or secukinumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab or secukinumab to meet the renewal criteria for adalimumab or secukinumab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Fither:

continued...

- 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints; or
- 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2.5 Any of the following:
 - 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — **(psoriatic arthritis)** only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*: and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for rheumatoid arthritis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects; or
 - 1.2.2 The patient has received insufficient benefit to meet the renewal criteria for rheumatoid arthritis; or
- 2 All of the following:

Subsidy (Manufacturer's Price)	Ful Subsidise	,	
\$	Per •	Manufacturer	

continued...

- 2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
- 2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses unless contraindicated); and
- 2.5 Either:
 - 2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
- 2.6 Either:
 - 2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Either:
 - 2.1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Initial application — (severe chronic plaque psoriasis) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe chronic plaque psoriasis; and
 - 1.2 Fither:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Either:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	osidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 2.4 The most recent PASI or DLQI assessment is no more than 1 month old at the time of application.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Renewal — (severe chronic plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 Applicant is a dermatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a dermatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and
 - 2.1.2 Fither:
 - 2.1.2.1 Following each prior etanercept treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-treatment baseline value; or
 - 2.1.2.2 Following each prior etanercept treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 2.2 Both:
 - 2.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2.2 Fither:
 - 2.2.2.1 Following each prior etanercept treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2.2 Following each prior etanercept treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-treatment baseline value; and
- 3 Etanercept to be administered at doses no greater than 50 mg every 7 days.

Note: A treatment course is defined as a minimum of 12 weeks of etanercept treatment

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day (or maximum tolerated dose); and
- 4 Patient has tried and not responded to at least three months of leflunomide at a dose of up to 20 mg daily (or maximum tolerated dose); and
- 5 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour measured no more than one month prior to the date of this application; or
- 5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications.

Renewal — (undifferentiated spondyloarthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Applicant is a rheumatologist; or
 - 1.2 Applicant is a Practitioner and confirms that a rheumatologist has provided a letter, email or fax recommending that the patient continues with etanercept treatment; and
- 2 Either:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior etanercept treatment in the opinion of the treating physician; and
- 3 Etanercept to be administered at doses no greater than 50 mg dose every 7 days.

Immune Modulators

ANTITHYMOCYTE GLOBUL	_IN (EQUINE) – PCT only – S	Specialist			
Inj 50 mg per ml, 5 ml		2,774.48	5	✓ ATGAM	
BACILLUS CALMETTE-GUE	ERIN (BCG) VACCINE - PCT	only - Specialist			
Subsidised only for blad	der cancer.				
Inj 2-8 × 100 million CFU	J	149.37	1	✓ OncoTICE	
Inj 40 mg per ml, vial		176.90	3	✓ SII-Onco-BCG S29	

Monoclonal Antibodies

ADALIMUMAB (AMGEVITA) - Special Authority see SA2142 below - Retail pharmacy

		645165) - see page 246 for details	Brand switch fee payable (Pharmacode 2645165
Amgevita	1	190.00	Inj 20 mg per 0.4 ml prefilled syringe
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled pen
✓ Amgevita	2	375.00	Inj 40 mg per 0.8 ml prefilled syringe

⇒SA2142 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 Both:
 - 2.1 The patient has severe Behcet's disease* that is significantly impacting the patient's quality of life; and
 - 2.2 Either:
 - 2.2.1 The patient has severe ocular, neurological, and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s); or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.2.2 The patient has severe gastrointestinal, rheumatological, and/or mucocutaneous symptoms and has not responded adequately to two or more treatments appropriate for the particular symptom(s).

Note: Indications marked with * are unapproved indications.

Initial application — (Hidradenitis suppurativa) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has hidradenitis suppurativa Hurley Stage II or Hurley Stage III lesions in distinct anatomic areas; and
 - 2.2 Patient has tried, but had an inadequate response to at least a 90 day trial of systemic antibiotics or has demonstrated intolerance to or has contraindications for systemic antibiotics; and
 - 2.3 Patient has 3 or more active lesions: and
 - 2.4 The patient has a DLQI of 10 or more and the assessment is no more than 1 month old at time of application.

Renewal — (Hidradenitis suppurativa) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline; and
- 2 The patient has a DLQI improvement of 4 or more from baseline.

Initial application — (Plaque psoriasis - severe chronic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept for severe chronic plaque psoriasis; and 2.1.2 Either:
 - 2.1.2.1 Patient has experienced intolerable side effects; or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for etanercept for severe chronic plaque psoriasis; or
 - 2.2 All of the following:
 - 2.2.1 Either:
 - 2.2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a PASI score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 2.2.1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2.2 Patient has tried, but had an inadequate response to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate. ciclosporin. or acitretin: and
 - 2.2.3 A PASI assessment or DLQI assessment has been completed for at least the most recent prior treatment course but no longer than 1 month following cessation of each prior treatment course and is no more than 1 month old at the time of application.

Renewal — (Plaque psoriasis - severe chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient had "whole body" severe chronic plaque psoriasis at the start of treatment; and

Subsidy (Manufacturer's Price)	Fully Subsidised	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

- 1.2 Either:
 - 1.2.1 The patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre treatment baseline value; or
 - 1.2.2 The patient has a DLQI improvement of 5 or more, when compared with the pre-treatment baseline value; or
- 2 Both:
 - 2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 2.2 Fither:
 - 2.2.1 The patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 2.2.2 The patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre treatment baseline value.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Both:
 - 2.1 Patient has pyoderma gangrenosum*; and
 - 2.2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporin, azathioprine, or methotrexate) and has not received an adequate response.

Note: Indications marked with * are unapproved indications.

Initial application — (Crohn's disease - adults) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira: or
- 2 All of the following:
 - 2.1 Patient has severe active Crohn's disease; and
 - 2.2 Any of the following:
 - 2.2.1 Patient has a CDAI score of greater than or equal to 300, or HBI score of greater than or equal to 10; or
 - 2.2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection: or
 - 2.2.4 Patient has an ileostomy or colostomy and has intestinal inflammation; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - adults) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 CDAI score has reduced by 100 points from the CDAI score, or HBI score has reduced by 3 points, from when the patient was initiated on adalimumab; or
- 2 CDAI score is 150 or less, or HBI is 4 or less; or
- 3 The patient has demonstrated an adequate response to treatment, but CDAI score and/or HBI score cannot be assessed.

Initial application — (Crohn's disease - children) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Fither:

1 The patient has previously had an approval for Humira; or

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsidi	sed	Generic
\$	Per	•	Manufacturer

continued...

- 2 All of the following:
 - 2.1 Paediatric patient has active Crohn's disease; and
 - 2.2 Fither:
 - 2.2.1 Patient has a PCDAI score of greater than or equal to 30; or
 - 2.2.2 Patient has extensive small intestine disease; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior therapy with immunomodulators and corticosteroids; and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (Crohn's disease - children) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
- 2 PCDAI score is 15 or less; or
- 3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed.

Initial application — **(Crohn's disease - fistulising)** only from a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has confirmed Crohn's disease: and
 - 2.2 Any of the following:
 - 2.2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2.2 Patient has one or more rectovaginal fistula(e); or
 - 2.2.3 Patient has complex peri-anal fistula; and
 - 2.3 A Baseline Fistula Assessment has been completed and is no more than 1 month old at the time of application.

Renewal — (Crohn's disease - fistulising) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain.

Initial application — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 The patient has had an initial Special Authority approval for infliximab for chronic ocular inflammation; or
 - 2.2 Both:
 - 2.2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective: or
 - 2.2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	

continued...

achieving a therapeutic dose of methotrexate.

Renewal — (Ocular inflammation - chronic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 12 weeks' initial treatment; or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Patient has had an initial Special Authority approval for infliximab for severe ocular inflammation; or
 - 2.2 Both:
 - 2.2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2.2 Any of the following:
 - 2.2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (Ocular inflammation - severe) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses: or
- 2 Following each 2 year treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 2 year treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Initial application — (ankylosing spondylitis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept for ankylosing spondylitis; and
 - 2.1.2 Either:
 - 2.1.2.1 The patient has experienced intolerable side effects; or
 - 2.1.2.2 The patient has received insufficient benefit to meet the renewal criteria for ankylosing spondylitis; or
 - 2.2 All of the following:
 - 2.2.1 Patient has a confirmed diagnosis of ankylosing spondylitis for more than six months; and
 - 2.2.2 Patient has low back pain and stiffness that is relieved by exercise but not by rest; and
 - 2.2.3 Patient has bilateral sacroiliitis demonstrated by radiology imaging; and

Subsidy	Fı	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacture 	r

continued...

- 2.2.4 Patient has not responded adequately to treatment with two or more NSAIDs, while patient was undergoing at least 3 months of a regular exercise regimen for ankylosing spondylitis; and
- 2.2.5 Either:
 - 2.2.5.1 Patient has limitation of motion of the lumbar spine in the sagittal and the frontal planes as determined by the following BASMI measures: a modified Schober's test of less than or equal to 4 cm and lumbar side flexion measurement of less than or equal to 10 cm (mean of left and right); or
 - 2.2.5.2 Patient has limitation of chest expansion by at least 2.5 cm below the average normal values corrected for age and gender; and
- 2.2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment and is no more than 1 month old at the time of application.

Renewal — (ankylosing spondylitis) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (Arthritis - oligoarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Fither:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept for oligoarticular course juvenile idiopathic arthritis (JIA); and
 - 2.1.2 Fither:
 - 2.1.2.1 Patient has experienced intolerable side effects; or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for oligoarticular course JIA; or
 - 2.2 All of the following:
 - 2.2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.2 Patient has had oligoarticular course JIA for 6 months duration or longer; and
 - 2.2.3 Either:
 - 2.2.3.1 At least 2 active joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.2 Moderate or high disease activity (cJADAS10 score greater than 1.5) with poor prognostic features after a 3-month trial of methotrexate (at the maximum tolerated dose).

Renewal — (Arthritis - oligoarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline: or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Roth:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 2.1.1 Patient has had an initial Special Authority approval for etanercept for polyarticular course juvenile idiopathic arthritis (JIA): and
- 2.1.2 Either:
 - 2.1.2.1 Patient has experienced intolerable side effects; or
 - 2.1.2.2 Patient has received insufficient benefit to meet the renewal criteria for polyarticular course JIA: or
- 2.2 All of the following:
 - 2.2.1 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.2.3 Any of the following:
 - 2.2.3.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.2.3.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Patient has had an initial Special Authority approval for etanercept or secukinumab for psoriatic arthritis; and
 - 2.1.2 Fither:
 - 2.1.2.1 The patient has experienced intolerable side effects; or
 - 2.1.2.2 The patient has received insufficient benefit from to meet the renewal criteria for psoriatic arthritis; or
 - 2.2 All of the following:
 - 2.2.1 Patient has had active psoriatic arthritis for six months duration or longer; and
 - 2.2.2 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.2.3 Patient has tried and not responded to at least three months of sulfasalazine or leflunomide at maximum tolerated doses (unless contraindicated); and
 - 2.2.4 Either:
 - 2.2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints; or
 - 2.2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.2.5 Any of the following:
 - 2.2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.2.5.2 Patient has an ESR greater than 25 mm per hour; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

2.2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (Arthritis - psoriatic) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

- 1 Following initial treatment, the patient has at least a 50% decrease in swollen joint count from baseline and a clinically significant response in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in swollen joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (Arthritis - rheumatoid) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept for rheumatoid arthritis; and
 - 2.1.2 Fither:
 - 2.1.2.1 The patient has experienced intolerable side effects: or
 - 2.1.2.2 The patient has received insufficient benefit from etanercept to meet the renewal criteria for rheumatoid arthritis: or
 - 2.2 All of the following:
 - 2.2.1 Patient has had rheumatoid arthritis (either confirmed by radiology imaging, or the patient is CCP antibody positive) for six months duration or longer; and
 - 2.2.2 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.2.3 Patient has tried and not responded to at least three months of methotrexate at a maximum tolerated dose (unless contraindicated); and
 - 2.2.4 Patient has tried and not responded to at least three months of methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate at maximum tolerated doses (unless contraindicated); and
 - 2.2.5 Either:
 - 2.2.5.1 Patient has tried and not responded to at least three months of methotrexate in combination with the maximum tolerated dose of ciclosporin; or
 - 2.2.5.2 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with methotrexate; and
 - 2.2.6 Either:
 - 2.2.6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen joints;
 - 2.2.6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip.

Renewal — (Arthritis - rheumatoid) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy (Manufacturer's Price)	Subs	Fully	Brand or Generic	
\$	Per	1	Manufacturer	

continued...

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 Either:
 - 2.1 Both:
 - 2.1.1 The patient has had an initial Special Authority approval for etanercept and/or tocilizumab for AOSD; and
 - 2.1.2 Fither:
 - 2.1.2.1 Patient has experienced intolerable side effects from etanercept and/or tocilizumab; or
 - 2.1.2.2 Patient has received insufficient benefit from at least a three-month trial of etanercept and/or tocilizumab; or
 - 2.2 All of the following:
 - 2.2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria; and
 - 2.2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, NSAIDs and methotrexate; and
 - 2.2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (ulcerative colitis) only from a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira; or
- 2 All of the following:
 - 2.1 Patient has histologically confirmed ulcerative colitis; and
 - 2.2 Either
 - 2.2.1 Patient's SCCAI score is greater than or equal to 4: or
 - 2.2.2 Patient's PUCAI score is greater than or equal to 65; and
 - 2.3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from prior therapy with immunomodulators and systemic corticosteroids: and
 - 2.4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 The SCCAI score has reduced by 2 points or more from the SCCAI score since initiation on adalimumab; or
- 2 The PUCAI score has reduced by 30 points or more from the PUCAI score since initiation on adalimumab.

Initial application — (undifferentiated spondyloarthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira: or
- 2 All of the following:
 - 2.1 Patient has undifferentiated peripheral spondyloarthritis* with active peripheral joint arthritis in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.2 Patient has tried and not responded to at least three months of each of methotrexate, sulfasalazine and leflunomide, at maximum tolerated doses (unless contraindicated); and
 - 2.3 Any of the following:
 - 2.3.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.3.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

2.3.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Note: Indications marked with * are unapproved indications

Renewal — (undifferentiated spondyloarthritis) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Either:

- 1 Following initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response in the opinion of the treating physician.

Initial application — (inflammatory bowel arthritis – axial) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira: or
- 2 All of the following:
 - 2.1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
 - 2.2 Patient has axial inflammatory pain for six months or more; and
 - 2.3 Patient is unable to take NSAIDs; and
 - 2.4 Patient has bilateral sacroiliitis demonstrated by radiological imaging; and
 - 2.5 Patient has not responded adequately to prior treatment consisting of at least 3 months of an exercise regime supervised by a physiotherapist; and
 - 2.6 A BASDAI of at least 6 on a 0-10 scale completed after the 3 month exercise trial, but prior to ceasing any previous pharmacological treatment.

Renewal — (inflammatory bowel arthritis – axial) from any relevant practitioner. Approvals valid for 2 years where treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less.

Initial application — (inflammatory bowel arthritis – peripheral) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 The patient has previously had an approval for Humira: or
- 2 All of the following:
 - 2.1 Patient has a diagnosis of active ulcerative colitis or active Crohn's disease; and
 - 2.2 Patient has active arthritis in at least four joints from the following: hip, knee, ankle, subtalar, tarsus, forefoot, wrist, elbow, shoulder, sternoclavicular; and
 - 2.3 Patient has tried and not responded to at least three months of methotrexate, or azathioprine at a maximum tolerated dose; and
 - 2.4 Patient has tried and not responded to at least three months of sulfasalazine at a maximum tolerated dose; and
 - 2.5 Any of the following:
 - 2.5.1 Patient has a CRP level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 2.5.2 Patient has an ESR greater than 25 mm per hour measured no more than one month prior to the date of this application; or
 - 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (inflammatory bowel arthritis – peripheral) from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria:

Fither:

(Man	Subsidy ufacturer's Price)	Fu Subsidis	ılly	Brand or Generic
(wich	. ,	Per	✓	Manufacturer

continued...

- 1 Following initial treatment, patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 Patient demonstrates at least a continuing 30% improvement in active joint count from baseline in the opinion of the treating physician.

ADALIMUMAB (HUMIRA - ALTERNATIVE BRAND) - Special Authority see SA2157 below - Retail pharmacy

Inj 20 mg per 0.2 ml prefilled syringe	1,599.96	2	✓ Humira
Inj 20 mg per 0.4 ml prefilled syringe		2	✓ Humira
Inj 40 mg per 0.8 ml prefilled pen		2	✓ HumiraPen
Inj 40 mg per 0.8 ml prefilled syringe	1,599.96	2	✓ Humira

(Humira Inj 20 mg per 0.4 ml prefilled syringe to be delisted 1 December 2022)

⇒SA2157 Special Authority for Subsidy

Initial application — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Behcet's disease - severe) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient has had a good clinical response to treatment with measurably improved quality of life; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 7 days. Fortnightly dosing has been considered.

Renewal — (Hidradenitis suppurativa) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has a reduction in active lesions (e.g. inflammatory nodules, abscesses, draining fistulae) of 25% or more from baseline: and
- 2 The patient has a Dermatology Quality of Life Index improvement of 4 or more from baseline; and
- 3 Adalimumab is to be administered at doses no greater than 40mg every 7 days. Fortnightly dosing has been considered.

Initial application — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Psoriasis - severe chronic plaque) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.1.2 Fither:
 - 1.1.2.1 Following each prior adalimumab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-adalimumab treatment baseline value; or
 - 1.1.2.2 Following each prior adalimumab treatment course the patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, when compared with the pre-treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior adalimumab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior adalimumab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-adalimumab treatment baseline value; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 A maximum of 8 doses.

Renewal — (Pyoderma gangrenosum) only from a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

Subsidy (Manufacturer's Price)		Fully Subsidised	Brand or Generic	
<u> </u>	Per	✓	Manufacturer	

continued...

- 1 The patient has demonstrated clinical improvement and continues to require treatment; and
- 2 A maximum of 8 doses.

Initial application — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevitat; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - adult) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on adalimumab; or
 - 1.2 CDAI score is 150 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment, but CDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - children) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on adalimumab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment, but PCDAI score cannot be assessed; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price	e)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has Crohn's and is considered to be at risk of disease destabilisation if there were to be a change to current treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Crohn's disease - fistulising) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
 - 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline as demonstrated by a reduction in the Fistula Assessment score, together with less induration and patient-reported pain; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
 - 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – chronic) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 12 weeks' initial treatment; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment, and a maximum of 6 months treatment with Amgevita; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with Amgevita, and a maximum of 6 months treatment with Amgevita and clinician attributes this loss of disease response to a change in treatment regimen; or
- 1.3 Patient has uveitis and is considered to be at risk of vision loss if they were to change treatment; and
- 2 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 3 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Ocular inflammation – severe) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 The patient has had a good clinical response following 3 initial doses; or
 - 1.2 Following each 12-month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or
 - 1.3 Following each 12-month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old; and</p>
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita); and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Treatment has resulted in an improvement in BASDAI of 4 or more points from pre-treatment baseline on a 10 point scale, or an improvement in BASDAI of 50%, whichever is less; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment; or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis – oligoarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Arthritis - polyarticular course juvenile idiopathic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months where the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Initial application — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Renewal — (Arthritis - psoriatic) only from a named specialist, rheumatologist or Practitioner on the recommendation of a named specialist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Adalimumab to be administered at doses no greater than 40 mg every 14 days.

Initial application — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication; and
- 4 Either:
 - 4.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Por 🗸	Manufacturer

continued...

4.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Renewal — (Arthritis – rheumatoid) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior adalimumab treatment in the opinion of the treating physician; and
- 2 Fither:
 - 2.1 Adalimumab to be administered at doses no greater than 40 mg every 14 days; or
 - 2.2 Patient cannot take concomitant methotrexate and requires doses of adalimumab higher than 40 mg every 14 days to maintain an adequate response.

Initial application — (Still's disease - adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 The patient has experienced intolerable side effects from adalimumab (Amgevita) following a minimum of 4 weeks treatment: or
 - 1.2 Patient has developed symptoms of loss of disease control following a minimum of 4 weeks treatment with adalimumab (Amgevita) and clinician attributes this loss of disease response to a change in treatment regimen; and
- 2 Patient has received a maximum of 6 months treatment with Amgevita; and
- 3 Patient has previously had a Special Authority approval for the Humira brand of adalimumab for this indication.

Renewal — (Still's disease – adult-onset (AOSD)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has demonstrated a sustained improvement in inflammatory markers and functional status.

AFLIBERCEPT - Special Authority see SA1772 below - Retail pharmacy

1 ✓ Eylea

⇒SA1772 Special Authority for Subsidy

Initial application — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 3 months for applications meeting the following criteria:

- Fither:
 - 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Wet age-related macular degeneration (wet AMD); or
 - 1.1.2 Polypoidal choroidal vasculopathy: or
 - 1.1.3 Choroidal neovascular membrane from causes other than wet AMD; and
 - 1.2 Fither:
 - 1.2.1 The patient has developed severe endophthalmitis or severe posterior uveitis following treatment with bevacizumab: or
 - 1.2.2 There is worsening of vision or failure of retina to dry despite three intraocular injections of bevacizumab four weeks apart; and
 - 1.3 There is no structural damage to the central fovea of the treated eye; and
 - 1.4 Patient has not previously been treated with ranibizumab for longer than 3 months; or
 - 2 Fither:
 - 2.1 Patient has current approval to use ranibizumab for treatment of wAMD and was found to be intolerant to ranibizumab within 3 months: or
 - 2.2 Patient has previously* (*before June 2018) received treatment with ranibizumab for wAMD and disease was stable

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

while on treatment.

Initial application — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has centre involving diabetic macular oedema (DMO); and
- 2 Patient's disease is non responsive to 4 doses of intravitreal bevacizumab when administered 4-6 weekly; and
- 3 Patient has reduced visual acuity between 6/9 6/36 with functional awareness of reduction in vision; and
- 4 Patient has DMO within central OCT (ocular coherence tomography) subfield > 350 micrometers; and
- 5 There is no centre-involving sub-retinal fibrosis or foveal atrophy.

Renewal — (wet age related macular degeneration) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented benefit must be demonstrated to continue; and
- 2 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 3 There is no structural damage to the central fovea of the treated eye.

Renewal — (diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 There is stability or two lines of Snellen visual acuity gain; and
- 2 There is structural improvement on OCT scan (with reduction in intra-retinal cysts, central retinal thickness, and sub-retinal fluid); and
- 3 Patient's vision is 6/36 or better on the Snellen visual acuity score; and
- 4 There is no centre-involving sub-retinal fibrosis or foveal atrophy; and
- 5 After each consecutive 12 months treatment with (2nd line anti-VEGF agent), patient has retrialled with at least one injection of bevacizumab and had no response.

BENRALIZUMAB - Special Authority see SA2151 below - Retail pharmacy

⇒SA2151 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and
- 4 Patient has a blood eosinophil count of greater than 0.5 × 10⁹ cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long-acting beta-2 agonist, or budesonide/formoterol as part of the anti-inflammatory reliever therapy plus maintenance regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months: and
- 7 Treatment is not to be used in combination with subsidised mepolizumab; and

ONCOLOGY AGENTS AND IMMUNOSUPPRESSANTS				
	Subsidy (Manufacturer's Price) \$	Fully Subsidised Per	Brand or Generic Manufacturer	
continued				
 8 Patient has an Asthma Control Test (ACT) score of using the ACT and oral corticosteroid dose must be the first dose to assess response to treatment; and 9 Either: 				
9.1 Patient has not previously received an anti-IL9.2 Both:	.5 biological therapy for the	ir severe eosinop	philic asthma; or	
9.2.1 Patient was refractory or intolerant to 9.2.2 Patient was not eligible to continue tre within 12 months of commencing trea	eatment with previous anti-litment.	L5 biological the		
Renewal — (Severe eosinophilic asthma) only from a re years for applications meeting the following criteria: Both:	spiratory physician or clinic	al immunologist.	Approvals valid for 2	
1 An increase in the Asthma Control Test (ACT) score2 Either:	of at least 5 from baseline;	and		
2.1 Exacerbations have been reduced from base2.2 Reduction in continuous oral corticosteroid u control.	•		·	
CASIRIVIMAB AND IMDEVIMAB - [Xpharm] - Special Au	thority see SA2096 below			
Inj 120 mg per ml casirivimab, 11.1 ml vial (1) and inj 1 per ml imdevimab, 11.1 ml vial (1)	•	1 OP ✓ F	Ronapreve	
SA2096 Special Authority for Subsidy Initial application — (Treatment of profoundly immunod valid for 2 weeks for applications meeting the following crite All of the following:		om any relevant	practitioner. Approvals	
 Patient has confirmed (or probable) COVID-19; and The patient is in the community with mild to moderat Patient is profoundly immunocompromised** and is against COVID-19 or is unvaccinated; and 		ed an adequate r	esponse to vaccination	
 4 Patient's symptoms started within the last 10 days; a 5 Patient is not receiving high flow oxygen or assisted 6 Casirivimab and imdevimab is to be administered at 	/mechanical ventilation; and a maximum dose of no gre	ater than 2,400 i	mg.	
Notes: * Mild to moderate disease severity as described or ** Examples include B-cell depletive illnesses or patients re				
CETUXIMAB – PCT only – Specialist – Special Authority s Inj 5 mg per ml, 20 ml vial Inj 5 mg per ml, 100 ml vial Inj 1 mg for ECP	364.00 1,820.00	1 🗸 E	Erbitux Erbitux Baxter	
⇒SA1697 Special Authority for Subsidy				
Initial application only from a medical oncologist or medical Approvals valid for 6 months for applications meeting the for All of the following:		mendation of a n	nedical oncologist.	
Patient has locally advanced, non-metastatic, squan Patient is contraindicated to, or is intolerant of, cisple Patient has good performance status; and To be administered in combination with radiation the	atin; and	d and neck; and		

GEMTUZUMAB OZOGAMICIN - PCT only - Specialist - Special Authority see SA2158 on the next page

✓ Mylotarg

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

⇒SA2158 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 Patient has not received prior chemotherapy for this condition; and
- 2 Patient has de novo CD33-positive acute myeloid leukaemia; and
- 3 Patient does not have acute promyelocytic leukaemia; and
- 4 Gemtuzumab ozogamicin will be used in combination with standard anthracycline and cytarabine (AraC); and
- 5 Patient is being treated with curative intent; and
- 6 Patient's disease risk has been assessed by cytogenetic testing to be good or intermediate; and
- 7 Patient must be considered eligible for standard intensive remission induction chemotherapy with standard anthracycline and cytarabine (AraC); and
- 8 Gemtuzumab ozogamicin to be funded for one course only (one dose at 3 mg per m² body surface area or up to 2 vials of 5 mg as separate doses).

Note: Acute myeloid leukaemia excludes acute promyelocytic leukaemia and acute myeloid leukaemia that is secondary to another haematological disorder (eg myelodysplasia or myeloproliferative disorder).

INFLIXIMAB - PCT only - Special Authority see SA2082 below

 Inj 100 mg
 806.00
 1
 ✓ Remicade

 Inj 1 mg for ECP
 8.29
 1 mg
 ✓ Baxter

⇒SA2082 Special Authority for Subsidy

Initial application — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe active Crohn's disease; and
- 2 Any of the following:
 - 2.1 Patient has a Crohn's Disease Activity Index (CDAI) score of greater than or equal to 300; or
 - 2.2 Patient has extensive small intestine disease affecting more than 50 cm of the small intestine; or
 - 2.3 Patient has evidence of short gut syndrome or would be at risk of short gut syndrome with further bowel resection; or
 - 2.4 Patient has an ileostomy or colostomy, and has intestinal inflammation; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (adults)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Any of the following:
 - 1.1 CDAI score has reduced by 100 points from the CDAI score when the patient was initiated on infliximab; or
 - 1.2 CDAI score is 150 or less; or
 - 1.3 The patient has demonstrated an adequate response to treatment but CDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or
(Manufacturer's Pric	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

- 1 Paediatric patient has severe active Crohn's disease; and
- - 2.1 Patient has a Paediatric Crohn's Disease Activity Index (PCDAI) score of greater than or equal to 30; or
 - 2.2 Patient has extensive small intestine disease; and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate; and
- 5 Patient must be reassessed for continuation after 3 months of therapy.

Renewal — (Crohn's disease (children)) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Any of the following:
 - 1.1 PCDAI score has reduced by 10 points from the PCDAI score when the patient was initiated on infliximab; or
 - 1.2 PCDAI score is 15 or less: or
 - 1.3 The patient has demonstrated an adequate response to treatment but PCDAI score cannot be assessed; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (Graft vs host disease) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has steroid-refractory acute graft vs. host disease of the gut.

Initial application — (Pulmonary sarcoidosis) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has life-threatening pulmonary sarcoidosis diagnosed by a multidisciplinary team that is refractory to other treatments.

Initial application — (acute severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 weeks for applications meeting the following criteria: Both:

- 1 Patient has acute, severe fulminant ulcerative colitis; and
- 2 Treatment with intravenous or high dose oral corticosteroids has not been successful.

Initial application — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Following 12 weeks of infliximab treatment, BASDAI has improved by 4 or more points from pre-infliximab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefited from treatment and that continued treatment is appropriate; and
- 3 Infliximab to be administered at doses no greater than 5 mg/kg every 6-8 weeks.

Initial application — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria: Fither:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

continued...

1 Both:

- 1.1 The patient has had an initial Special Authority approval for adalimumab for chronic ocular inflammation; and
- 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for chronic ocular inflammation; or

2 Both:

- 2.1 Patient has severe uveitis uncontrolled with treatment of steroids and other immunosuppressants with a severe risk of vision loss; and
- 2.2 Any of the following:
 - 2.2.1 Patient is 18 years or older and treatment with at least two other immunomodulatory agents has proven ineffective; or
 - 2.2.2 Patient is under 18 years and treatment with methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or
 - 2.2.3 Patient is under 8 years and treatment with steroids or methotrexate has proven ineffective or is not tolerated at a therapeutic dose; or disease requires control to prevent irreversible vision loss prior to achieving a therapeutic dose of methotrexate.

Renewal — (chronic ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 Patient has confirmed Crohn's disease: and
- 2 Either:
 - 2.1 Patient has one or more complex externally draining enterocutaneous fistula(e); or
 - 2.2 Patient has one or more rectovaginal fistula(e).

Renewal — (fistulising Crohn's disease) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

1 Either:

- 1.1 The number of open draining fistulae have decreased from baseline by at least 50%; or
- 1.2 There has been a marked reduction in drainage of all fistula(e) from baseline (in the case of adult patients, as demonstrated by a reduction in the Fistula Assessment score), together with less induration and patient reported pain; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Subsidy		Fully	Brand or
(Manufacturer's Price	e)	Subsidised	Generic
\$	Per	✓	Manufacturer

continued...

Initial application — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with neurosarcoiosis by a multidisciplinary team; and
- 2 Patient has CNS involvement: and
- 3 Patient has steroid-refractory disease; and
- 4 Fither:
 - 4.1 IV cyclophosphamide has been tried; or
 - 4.2 Treatment with IV cyclophosphamide is clinically inappropriate.

Renewal — (neurosarcoidosis) only from a neurologist or Practitioner on the recommendation of a neurologist. Approvals valid for 18 months for applications meeting the following criteria:

Either:

- 1 A withdrawal period has been tried and the patient has relapsed; or
- 2 All of the following:
 - 2.1 A withdrawal period has been considered but would not be clinically appropriate; and
 - 2.2 There has been a marked reduction in prednisone dose; and
 - 2.3 Either:
 - 2.3.1 There has been an improvement in MRI appearances; or
 - 2.3.2 Marked improvement in other symptomology.

Initial application — (plaque psoriasis) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 Both:
 - 1.1 The patient has had an initial Special Authority approval for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; and
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or secukinumab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or secukinumab to meet the renewal criteria for adalimumab, etanercept or secukinumab for severe chronic plaque psoriasis; or
- 2 All of the following:
 - 2.1 Fither:
 - 2.1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis: or
 - 2.1.2 Patient has severe chronic plague psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
 - 2.2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from. at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin; and
 - 2.3 A PASI assessment has been completed for at least the most recent prior treatment course (but preferably all prior treatment courses), preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
 - 2.4 The most recent PASI assessment is no more than 1 month old at the time of initiation.

Note: "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom subscores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot,

Subsidy	Fu	ılly	Brand or	
(Manufacturer's Price)	Subsidis	ed	Generic	
\$	Per	✓	Manufacturer	

continued...

as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment. **Renewal — (plaque psoriasis)** only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

- 1 Fither:
 - 1.1 Both:
 - 1.1.1 Patient had "whole body" severe chronic plague psoriasis at the start of treatment; and
 - 1.1.2 Following each prior infliximab treatment course the patient has a PASI score which is reduced by 75% or more, or is sustained at this level, when compared with the pre-infliximab treatment baseline value; or
 - 1.2 Both:
 - 1.2.1 Patient had severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot at the start of treatment; and
 - 1.2.2 Either:
 - 1.2.2.1 Following each prior infliximab treatment course the patient has a reduction in the PASI symptom subscores for all 3 of erythema, thickness and scaling, to slight or better, or sustained at this level, as compared to the treatment course baseline values; or
 - 1.2.2.2 Following each prior infliximab treatment course the patient has a reduction of 75% or more in the skin area affected, or sustained at this level, as compared to the pre-infliximab treatment baseline value: and
 - 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with infliximab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 Rheumatoid arthritis; or
 - 2.2 Ankylosing spondylitis; or
 - 2.3 Psoriatic arthritis; or
 - 2.4 Severe ocular inflammation: or
 - 2.5 Chronic ocular inflammation: or
 - 2.6 Crohn's disease (adults): or
 - 2.7 Crohn's disease (children); or
 - 2.8 Fistulising Crohn's disease: or
 - 2.9 Severe fulminant ulcerative colitis; or
 - 2.10 Severe ulcerative colitis; or
 - 2.11 Plaque psoriasis; or
 - 2.12 Neurosarcoidosis; or
 - 2.13 Severe Behcet's disease.

Initial application — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept and/or secukinumab; or
 - 2.2 Following 3-4 months' initial treatment with adalimumab and/or etanercept and/or secukinumab, the patient did not meet the renewal criteria for adalimumab and/or etanercept and/or secukinumab for psoriatic arthritis.

S	Subsidy	Fully	Brand or
(Manufa	acturer's Price) Si	ubsidised	Generic
	\$ Per	✓	Manufacturer

continued...

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior infliximab treatment in the opinion of the treating physician; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Initial application — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept; and
- 3 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance.

Renewal — (rheumatoid arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 2.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 3 Infliximab to be administered at doses no greater than 3 mg/kg every 8 weeks.

Initial application — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe Behcet's disease which is significantly impacting the patient's quality of life (see Notes); and
- 2 Fither:
 - 2.1 The patient has severe ocular, neurological and/or vasculitic symptoms and has not responded adequately to one or more treatment(s) appropriate for the particular symptom(s) (see Notes); or
 - 2.2 The patient has severe gastrointestinal, rheumatologic and/or mucocutaneous symptoms and has not responded adequately to two or more treatment appropriate for the particular symptom(s) (see Notes); and
- 3 The patient is experiencing significant loss of quality of life.

Notes: Behcet's disease diagnosed according to the International Study Group for Behcet's Disease. Lancet 1990;335(8697):1078-80. Quality of life measured using an appropriate quality of life scale such as that published in Gilworth et al J Rheumatol. 2004;31:931-7.

Treatments appropriate for the particular symptoms are those that are considered standard conventional treatments for these symptoms, for example intravenous/oral steroids and other immunosuppressants for ocular symptoms; azathioprine, steroids, thalidomide, interferon alpha and ciclosporin for mucocutaneous symptoms; and colchicine, steroids and methotrexate for rheumatological symptoms.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	1	Manufacturer	

continued...

Renewal — (severe Behcet's disease) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has had a good clinical response to initial treatment with measurably improved quality of life; and
- 2 Infliximab to be administered at doses no greater than 5 mg/kg every 8 weeks.

Renewal — (severe fulminant ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Where maintenance treatment is considered appropriate, infliximab should be used in combination with immunomodulators and reassessed every 6 months; and
- 2 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 4 months for applications meeting the following criteria:

Either: 1 Both:

- 4.4 The second of the second
 - 1.1 The patient has had an initial Special Authority approval for adalimumab for severe ocular inflammation; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab; or
 - 1.2.2 The patient has received insufficient benefit from adalimumab to meet the renewal criteria for adalimumab for severe ocular inflammation: or
- 2 Both:
 - 2.1 Patient has severe, vision-threatening ocular inflammation requiring rapid control; and
 - 2.2 Any of the following:
 - 2.2.1 Treatment with high-dose steroids (intravenous methylprednisolone) followed by high dose oral steroids has proven ineffective at controlling symptoms; or
 - 2.2.2 Patient developed new inflammatory symptoms while receiving high dose steroids; or
 - 2.2.3 Patient is aged under 8 years and treatment with high dose oral steroids and other immunosuppressants has proven ineffective at controlling symptoms.

Renewal — (severe ocular inflammation) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Any of the following:

- 1 The patient has had a good clinical response following 3 initial doses; or
- 2 Following each 12 month treatment period, the patient has had a sustained reduction in inflammation (Standardisation of Uveitis Nomenclature (SUN) criteria < ½+ anterior chamber or vitreous cells, absence of active vitreous or retinal lesions, or resolution of uveitic cystoid macular oedema); or</p>
- 3 Following each 12 month treatment period, the patient has a sustained steroid sparing effect, allowing reduction in prednisone to < 10mg daily, or steroid drops less than twice daily if under 18 years old.

Note: A trial withdrawal should be considered after every 24 months of stability, unless the patient is deemed to have extremely high risk of irreversible vision loss if infliximab is withdrawn.

Initial application — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has histologically confirmed ulcerative colitis; and
- 2 Fither:

	Subsidy		Fully	Brand or
(M	anufacturer's Price)	Subs	idised	Generic
	\$	Per	✓	Manufacturer

continued...

- 2.1 Patient is 18 years or older and the Simple Clinical Colitis Activity Index (SCCAI) is greater than or equal to 4; or
- 2.2 Patient is under 18 years and the Paediatric Ulcerative Colitis Activity Index (PUCAI) score is greater than or equal to 65: and
- 3 Patient has tried but had an inadequate response to, or has experienced intolerable side effects from, prior systemic therapy with immunomodulators at maximum tolerated doses for an adequate duration (unless contraindicated) and corticosteroids; and
- 4 Surgery (or further surgery) is considered to be clinically inappropriate.

Renewal — (ulcerative colitis) only from a gastroenterologist or Practitioner on the recommendation of a gastroenterologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient is continuing to maintain remission and the benefit of continuing infliximab outweighs the risks; and
- 2 Either:
 - 2.1 Patient is 18 years or older and the SCCAI score has reduced by 2 points or more from the SCCAI score when the patient was initiated on infliximab; or
 - 2.2 Patient is under 18 years and the PUCAI score has reduced by 30 points or more from the PUCAI score when the patient was initiated on infliximab; and
- 3 Infliximab to be administered at doses up to 5 mg/kg every 8 weeks. Up to 10 mg/kg every 8 weeks (or equivalent) can be used for up to 3 doses if required for secondary non-response to treatment for re-induction. Another re-induction may be considered sixteen weeks after completing the last re-induction cycle. Up to 10 mg/kg every 8 weeks (or equivalent) may be used for patients treated with this dose prior to 1 February 2019.

Initial application — (pyoderma gangrenosum) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has pyoderma gangrenosum*; and
- 2 Patient has received three months of conventional therapy including a minimum of three pharmaceuticals (e.g. prednisone, ciclosporine, azathioprine, or methotrexate) and not received an adequate response; and
- 3 A maximum of 8 doses.

Note: Note: Indications marked with * are unapproved indications.

Renewal — (pyoderma gangrenosum) only from a dermatologist or Practitioner on the recommendation of a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient has shown clinical improvement; and
- 2 Patient continues to require treatment; and
- 3 A maximum of 8 doses.

MEPOLIZUMAB - Special Authority see SA2154 below - Retail pharmacy

✓ Nucala	1	1,638.00 mg prefilled pen	
✓ Nucala	1	1,638.00	Ini 100 ma

⇒SA2154 Special Authority for Subsidy

Initial application — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 12 years or older; and
- 2 Patient must have a diagnosis of severe eosinophilic asthma documented by a respiratory physician or clinical immunologist; and
- 3 Conditions that mimic asthma eg. vocal cord dysfunction, central airway obstruction, bronchiolitis etc. have been excluded: and

Subsidy (Manufacturer's Price)	Fully Subsidised	
 \$	Per 🗸	Manufacturer

continued...

- 4 Patient has a blood eosinophil count of greater than 0.5 x 10^9 cells/L in the last 12 months; and
- 5 Patient must be adherent to optimised asthma therapy including inhaled corticosteroids (equivalent to at least 1000 mcg per day of fluticasone propionate) plus long acting beta-2 agonist, or budesonide/formoterol as part of the single maintenance and reliever therapy regimen, unless contraindicated or not tolerated; and
- 6 Either:
 - 6.1 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral corticosteroids; or
 - 6.2 Patient has received continuous oral corticosteroids of at least the equivalent of 10 mg per day over the previous 3 months; and
- 7 Treatment is not to be used in combination with subsidised benralizumab: and
- 8 Patient has an Asthma Control Test (ACT) score of 10 or less. Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 52 weeks after the first dose to assess response to treatment; and
- 9 Either:
 - 9.1 Patient has not previously received an anti-IL5 biological therapy for their severe eosinophilic asthma; or
 - 9.2 Both:
 - 9.2.1 Patient was refractory or intolerant to previous anti-IL5 biological therapy; and
 - 9.2.2 Patient was not eligible to continue treatment with previous anti-IL5 biological therapy and discontinued within 12 months of commencing treatment.

Renewal — (Severe eosinophilic asthma) only from a respiratory physician or clinical immunologist. Approvals valid for 2 years for applications meeting the following criteria:

Roth:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 Either:
 - 2.1 Exacerbations have been reduced from baseline by 50% as a result of treatment with mepolizumab; or
 - 2.2 Reduction in continuous oral corticosteroid use by 50% or by 10 mg/day while maintaining or improving asthma control.

OBINUTUZUMAB - PCT only - Specialist - Special Author	ority see SA2155 below		
Inj 25 mg per ml, 40 ml vial	5,910.00	1	Gazyva
Inj 1 mg for ECP	6.21	1 mg	✓ Baxter

⇒SA2155 Special Authority for Subsidy

Initial application — (chronic lymphocytic leukaemia) only from a haematologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C CD20+ chronic lymphocytic leukaemia requiring treatment; and
- 2 The patient is obinutuzumab treatment naive; and
- 3 The patient is not eligible for full dose FCR due to comorbidities with a score > 6 on the Cumulative Illness Rating Scale (CIRS) or reduced renal function (creatinine clearance < 70mL/min); and
- 4 Patient has adequate neutrophil and platelet counts* unless the cytopenias are a consequence of marrow infiltration by CLL: and
- 5 Patient has good performance status; and
- 6 Obinutuzumab to be administered at a maximum cumulative dose of 8,000 mg and in combination with chlorambucil for a maximum of 6 cycles.

Notes: Chronic lymphocytic leukaemia includes small lymphocytic lymphoma. Comorbidity refers only to illness/impairment other than CLL induced illness/impairment in the patient. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with obinutuzumab

Subsidy		Fully	Brand or
(Manufacturer's F	,	Subsidised	Generic
\$	Per		Manutacturer

continued...

is expected to improve symptoms and improve ECOG score to < 2.

* Neutrophil greater than or equal to 1.5×10^9 /L and platelets greater than or equal to 75×10^9 /L.

Initial application — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 9 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 Patient has follicular lymphoma; or
 - 1.2 Patient has marginal zone lymphoma; and
- 2 Patient is refractory to or has relapsed within 12 months of a rituximab containing combined chemo-immunotherapy regimen*: and
- 3 Patient has an ECOG performance status of 0-2; and
- 4 Patient has been previously treated with no more than four chemotherapy regimens; and
- 5 Obinutuzumab to be administered at a maximum dose of 1000 mg for a maximum of 6 cycles in combination with chemotherapy*.

Note: * includes unapproved indications

Renewal — (follicular / marginal zone lymphoma) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 24 months for applications meeting the following criteria: All of the following:

- 1 Patient has no evidence of disease progression following obinutuzumab induction therapy; and
- 2 Obinutuzumab to be administered at a maximum of 1000 mg every 2 months for a maximum of 2 years; and
- 3 Obinutuzumab to be discontinued at disease progression.

OMALIZUMAB - Special Authority see SA1744 below - Retail pharmacy

Inj 150 mg prefilled syringe	450.00	1	Xolair
Inj 150 mg vial	450.00	1	✓ Xolair

⇒SA1744 Special Authority for Subsidy

Initial application — (severe asthma) only from a respiratory specialist or clinical immunologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient must be aged 6 years or older; and
- 2 Patient has a diagnosis of severe asthma; and
- 3 Past or current evidence of atopy, documented by skin prick testing or RAST; and
- 4 Total serum human immunoglobulin E (IgE) between 76 IU/mL and 1300 IU/ml at baseline; and
- 5 Proven adherence with optimal inhaled therapy including high dose inhaled corticosteroid (budesonide 1,600 mcg per day or fluticasone propionate 1,000 mcg per day or equivalent), plus long-acting beta-2 agonist therapy (at least salmeterol 50 mcg bd or eformoterol 12 mcg bd) for at least 12 months, unless contraindicated or not tolerated; and
- 6 Fither:
 - 6.1 Patient has received courses of systemic corticosteroids equivalent to at least 28 days treatment in the past 12 months, unless contraindicated or not tolerated; or
 - 6.2 Patient has had at least 4 exacerbations needing systemic corticosteroids in the previous 12 months, where an exacerbation is defined as either documented use of oral corticosteroids for at least 3 days or parenteral steroids;
- 7 Patient has an Asthma Control Test (ACT) score of 10 or less; and
- 8 Baseline measurements of the patient's asthma control using the ACT and oral corticosteroid dose must be made at the time of application, and again at around 26 weeks after the first dose to assess response to treatment.

Initial application — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer

continued...

- 1 Patient must be aged 12 years or older; and
- 2 Fither:
 - 2.1 Both:
 - 2.1.1 Patient is symptomatic with Urticaria Activity Score 7 (UAS7) of 20 or above; and
 - 2.1.2 Patient has a Dermatology life quality index (DLQI) of 10 or greater; or
 - 2.2 Patient has a Urticaria Control Test (UCT) of 8 or less; and
- 3 Any of the following:
 - 3.1 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and ciclosporin (> 3 mg/kg day) for at least 6 weeks; or
 - 3.2 Patient has been taking high dose antihistamines (e.g. 4 times standard dose) and at least 3 courses of systemic corticosteroids (> 20 mg prednisone per day for at least 5 days) in the previous 6 months; or
 - 3.3 Patient has developed significant adverse effects whilst on corticosteroids or ciclosporin; and
- 4 Fither:
 - 4.1 Treatment to be stopped if inadequate response* following 4 doses: or
 - 4.2 Complete response* to 6 doses of omalizumab.

Renewal — (severe asthma) only from a clinical immunologist or respiratory specialist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 An increase in the Asthma Control Test (ACT) score of at least 5 from baseline; and
- 2 A reduction in the maintenance oral corticosteroid dose or number of exacerbations of at least 50% from baseline.

Renewal — (severe chronic spontaneous urticaria) only from a clinical immunologist or dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Patient has previously adequately responded* to 6 doses of omalizumab; or
- 2 Both:
 - 2.1 Patient has previously had a complete response* to 6 doses of omalizumab; and
 - 2.2 Patient has relapsed after cessation of omalizumab therapy.

Note: *Inadequate response defined as less than 50% reduction in baseline UAS7 and DLQI score, or an increase in Urticaria Control Test (UCT) score of less than 4 from baseline. Patient is to be reassessed for response after 4 doses of omalizumab. Complete response is defined as UAS7 less than or equal to 6 and DLQI less than or equal to 5; or UCT of 16. Relapse of chronic urticaria on stopping prednisone/ciclosporin does not justify the funding of omalizumab.

PALIVIZUMAB - PCT only - Specialist - Special Authority see SA2143 below

(Synagis Inj 100 mg per ml, 1 ml vial to be delisted 1 January 2024)

⇒SA2143 Special Authority for Subsidy

Initial application — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Infant was born in the last 2 years and has severe lung, airway, neurological or neuromuscular disease that requires ongoing, life-sustaining community ventilation; or
- 2 Both:
 - 2.1 Infant was born in the last 12 months; and
 - 2.2 Any of the following:
 - 2.2.1 Patient was born at less than 28 weeks gestation; or
 - 2.2.2 Both:
 - 2.2.2.1 Patient was born at less than 32 weeks gestation; and

Subsidy (Manufacturer's Price)	Sul	Fully bsidised	Brand or Generic
\$	Per	✓	Manufacturer

continued...

2.2.2.2 Either:

2.2.2.2.1 Patient has chronic lung disease; or

2.2.2.2. Patient is Māori or any Pacific ethnicity: or

2.2.3 Both:

2.2.3.1 Patient has haemodynamically significant heart disease; and

2.2.3.2 Any of the following:

2.2.3.2.1 Patient has unoperated simple congenital heart disease with significant left to right shunt (see

2.2.3.2.2 Patient has unoperated or surgically palliated complex congenital heart disease; or

2.2.3.2.3 Patient has severe pulmonary hypertension (see note b); or

2.2.3.2.4 Patient has moderate or severe LV failure (see note c).

Notes:

- a) Patient requires/will require heart failure medication, and/or patient has significant pulmonary hypertension, and/or patient will require surgical palliation/definitive repair within the next 3 months.
- b) Mean pulmonary artery pressure more than 25 mmHg.
- c) LV Eiection Fraction less than 40%.

Renewal — (RSV prophylaxis for the 2022/2023 RSV seasons, in the context of COVID-19) only from a paediatrician. Approvals valid for 6 months where patient still meets initial criteria.

PERTUZUMAB - PCT only - Specialist - Special Authority see SA1606 below

Inj 30 mg per ml, 14 ml vial	3,927.00	1	Perjeta
Inj 420 mg for ECP	3,927.00	420 mg OP	Baxter

⇒SA1606 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Fither:
 - 2.1 Patient is chemotherapy treatment naïve; or
 - 2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
- 3 The patient has good performance status (ECOG grade 0-1); and
- 4 Pertuzumab to be administered in combination with trastuzumab; and
- 5 Pertuzumab maximum first dose of 840 mg, followed by maximum of 420 mg every 3 weeks; and
- 6 Pertuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on pertuzumab and trastuzumab.

1 075 50

RITUXIMAB (MABTHERA) - PCT only - Specialist - Special Authority see SA1976 on the next page . Mahthara

iiij 100 iiig pei 10 iiii viai		~	• IVIADUICIA
Inj 500 mg per 50 ml vial	2,688.30	1	✓ Mabthera
Inj 1 mg for ECP	5.64	1 mg	✓ Baxter (Mabthera)

Ini 100 ma nor 10 ml vial

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic

\$ Per ✔ Manufacturer

⇒SA1976 Special Authority for Subsidy

Initial application — (rheumatoid arthritis - TNF inhibitors contraindicated) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Treatment with a Tumour Necrosis Factor alpha inhibitor is contraindicated: and
- 2 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and
- 3 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
- 4 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with sulfasalazine and hydroxychloroguine sulphate (at maximum tolerated doses); and
- 5 Any of the following:
 - 5.1 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with the maximum tolerated dose of ciclosporin: or
 - 5.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate in combination with intramuscular gold; or
 - 5.3 Patient has tried and not responded to at least three months of therapy at the maximum tolerated dose of leflunomide alone or in combination with oral or parenteral methotrexate; and
- 6 Either:
 - 6.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 swollen, tender joints; or
 - 6.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 7 Either:
 - 7.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 7.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months; and
- 8 Fither:
 - 8.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 8.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 9 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Initial application — (rheumatoid arthritis - prior TNF inhibitor use) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Both:
 - 1.1 The patient has had an initial community Special Authority approval for at least one of etanercept and/or adalimumab for rheumatoid arthritis: and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 1.2.2 Following at least a four month trial of adalimumab and/or etanercept, the patient did not meet the renewal criteria for adalimumab and/or etanercept for rheumatoid arthritis: and
- 2 Fither:
 - 2.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 2.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 3 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

5	Subsidy	Fully	Brand or
(Manufa	acturer's Price)	Subsidised	Generic
	\$ Per	•	Manufacturer

continued...

Renewal — (rheumatoid arthritis - re-treatment in 'partial responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had between a 30% and 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.3 At 4 months following the third and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
- 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

Renewal — (rheumatoid arthritis - re-treatment in 'responders' to rituximab) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Either:
 - 1.1 At 4 months following the initial course of rituximab infusions the patient had at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 At 4 months following the second and subsequent courses of rituximab infusions, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; and
- 2 Rituximab re-treatment not to be given within 6 months of the previous course of treatment; and
- 3 Either:
 - 3.1 Rituximab to be used as an adjunct to methotrexate or leflunomide therapy; or
 - 3.2 Patient is contraindicated to both methotrexate and leflunomide, requiring rituximab monotherapy to be used; and
- 4 Maximum of two 1,000 mg infusions of rituximab given two weeks apart.

RITUXIMAB (RIXIMYO) - PCT only - Specialist - Special A	uthority see SA2114 b	elow	
Inj 100 mg per 10 ml vial	275.33	2	✓ Riximyo
Inj 500 mg per 50 ml vial	688.20	1	✓ Riximyo
Inj 1 mg for ECP	1.38	1 mg	✓ Baxter (Riximyo)

⇒SA2114 Special Authority for Subsidy

Initial application — (ABO-incompatible organ transplant) from any relevant practitioner. Approvals valid without further renewal unless notified where patient is to undergo an ABO-incompatible solid organ transplant*.

Note: Indications marked with * are unapproved indications.

Initial application — **(ANCA associated vasculitis)** from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks; and
- 3 Any of the following:

Subsidy	Fu	,	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

- 3.1 Induction therapy with daily oral or pulse intravenous cyclophosphamide has failed to achieve significant improvement of disease after at least 3 months; or
- 3.2 Patient has previously had a cumulative dose of cyclophosphamide > 15 g or a further repeat 3 month induction course of cyclophosphamide would result in a cumulative dose > 15 g; or
- 3.3 Cyclophosphamide and methotrexate are contraindicated; or
- 3.4 Patient is a female of child-bearing potential; or
- 3.5 Patient has a previous history of haemorrhagic cystitis, urological malignancy or haematological malignancy.

Note: Indications marked with * are unapproved indications.

Renewal — (ANCA associated vasculitis) from any relevant practitioner. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with ANCA associated vasculitis*; and
- 2 Patient has previously responded to treatment with rituximab but is now experiencing an acute flare of vasculitis; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² of body-surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Antibody-mediated organ transplant rejection) from any relevant practitioner. Approvals valid without further renewal unless notified where patient has been diagnosed with antibody-mediated organ transplant rejection*.

Note: Indications marked with * are unapproved indications.

Initial application — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has progressive Binet stage A, B or C chronic lymphocytic leukaemia (CLL) requiring treatment; and
- 2 Any of the following:
 - 2.1 The patient is rituximab treatment naive; or
 - 2.2 Either:
 - 2.2.1 The patient is chemotherapy treatment naive; or
 - 2.2.2 Both:
 - 2.2.2.1 The patient's disease has relapsed following no more than three prior lines of chemotherapy treatment: and
 - 2.2.2.2 The patient has had a treatment-free interval of 12 months or more if previously treated with fludarabine and cyclophosphamide chemotherapy; or
 - 2.3 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; and
- 3 The patient has good performance status; and
- 4 Either:
 - 4.1 The patient does not have chromosome 17p deletion CLL; or
 - 4.2 Rituximab treatment is to be used in combination with funded venetoclax for relapsed/refractory chronic lymphocytic leukaemia: and
- 5 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles; and
- 6 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration), bendamustine or venetoclax.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments. 'Good performance status' means ECOG score of 0-1, however, in patients temporarily debilitated by their CLL disease symptoms a higher ECOG (2 or 3) is acceptable where treatment with rituximab is expected to improve symptoms and improve ECOG score to < 2.

Subsidy	Fu	lly Brand or	
(Manufacturer's Price)	Subsidis	ed Generic	
\$	Per	 Manufacturer 	

continued...

Renewal — (Chronic lymphocytic leukaemia) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 The patient's disease has relapsed within 36 months of previous treatment and rituximab treatment is to be used in combination with funded venetoclax; or
 - 1.2 All of the following:
 - 1.2.1 The patient's disease has relapsed following no more than one prior line of treatment with rituximab for CLL;
 - 1.2.2 The patient has had an interval of 36 months or more since commencement of initial rituximab treatment; and
 - 1.2.3 The patient does not have chromosome 17p deletion CLL; and
 - 1.2.4 It is planned that the patient receives full dose fludarabine and cyclophosphamide (orally or dose equivalent intravenous administration) or bendamustine; and
- 2 Rituximab to be administered in combination with fludarabine and cyclophosphamide, bendamustine or venetoclax for a maximum of 6 treatment cycles.

Note: 'Chronic lymphocytic leukaemia (CLL)' includes small lymphocytic lymphoma. A line of chemotherapy treatment is considered to comprise a known standard therapeutic chemotherapy regimen and supportive treatments.

Initial application — (Neuromyelitis Optica Spectrum Disorder(NMOSD)) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 Either:
 - 2.1 The patient has experienced a severe episode or attack of NMOSD (rapidly progressing symptoms and clinical investigations supportive of a severe attack of NMOSD); or
 - 2.2 All of the following:
 - 2.2.1 The patient has experienced a breakthrough attack of NMOSD; and
 - 2.2.2 The patient is receiving treatment with mycophenolate; and
 - 2.2.3 The patients is receiving treatment with corticosteroids.

Renewal — (Neuromyelitis Optica Spectrum Disorder) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 One of the following dose regimens is to be used: 2 doses of 1,000 mg rituximab administered fortnightly, or 4 doses of 375 mg/m2 administered weekly for four weeks; and
- 2 The patients has responded to the most recent course of rituximab; and
- 3 The patient has not received rituximab in the previous 6 months.

Initial application — (Post-transplant) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 2 To be used for a maximum of 8 treatment cycles.

Note: Indications marked with * are unapproved indications.

Renewal — (Post-transplant) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

All of the following:

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has B-cell post-transplant lymphoproliferative disorder*; and
- 3 To be used for no more than 6 treatment cycles.

Note: Indications marked with * are unapproved indications.

Initial application — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

Both:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 Fither:
 - 2.1 Treatment with corticosteroids and at least one other immunosuppressant for at least a period of 12 months has been ineffective; or
 - 2.2 Both:
 - 2.2.1 Treatment with at least one other immunosuppressant for a period of at least 12 months; and
 - 2.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Renewal — (Severe Refractory Myasthenia Gravis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Either:
 - 3.1 The patient has relapsed despite treatment with corticosteroids and at least one other immunosuppressant for a period of at least 12 months; or
 - 3.2 Both:
 - 3.2.1 The patient's myasthenia gravis has relapsed despite treatment with at least one immunosuppressant for a period of at least 12 months; and
 - 3.2.2 Corticosteroids have been trialed for at least 12 months and have been discontinued due to unacceptable side effects.

Initial application — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient is a child with SDNS* or FRNS*; and
- 2 Treatment with steroids for at least a period of 3 months has been ineffective or associated with evidence of steroid toxicity; and
- 3 Treatment with ciclosporin for at least a period of 3 months has been ineffective and/or discontinued due to unacceptable side effects; and
- 4 Treatment with mycophenolate for at least a period of 3 months with no reduction in disease relapses; and
- 5 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid dependent nephrotic syndrome (SDNS) or frequently relapsing nephrotic syndrome (FRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

Subsidy		Fully	Brand or	_
(Manufacturer's Price))	Subsidised	Generic	
\$	Per	1	Manufacturer	

continued...

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient is a child with SRNS* where treatment with steroids and ciclosporin for at least 3 months have been ineffective; and
- 2 Treatment with tacrolimus for at least 3 months has been ineffective; and
- 3 Genetic causes of nephrotic syndrome have been excluded; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (Steroid resistant nephrotic syndrome (SRNS)) only from a nephrologist or Practitioner on the recommendation of a nephrologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient who was previously treated with rituximab for nephrotic syndrome*; and
- 2 Treatment with rituximab was previously successful and has demonstrated sustained response for greater than 6 months, but the condition has relapsed and the patient now requires repeat treatment; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

Fither:

- 1 All of the following:
 - 1.1 The patient has treatment naive aggressive CD20 positive NHL; and
 - 1.2 To be used with a multi-agent chemotherapy regimen given with curative intent; and
 - 1.3 To be used for a maximum of 8 treatment cycles; or
- 2 Both:
 - 2.1 The patient has aggressive CD20 positive NHL with relapsed disease following prior chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Renewal — (aggressive CD20 positive NHL) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has relapsed refractory/aggressive CD20 positive NHL; and
- 3 To be used with a multi-agent chemotherapy regimen given with curative intent; and
- 4 To be used for a maximum of 4 treatment cycles.

Note: 'Aggressive CD20 positive NHL' includes large B-cell lymphoma and Burkitt's lymphoma/leukaemia

Initial application — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria: Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

- 1 Patient has mild congenital haemophilia complicated by inhibitors; or
- 2 Patient has severe congenital haemophilia complicated by inhibitors and has failed immune tolerance therapy; or
- 3 Patient has acquired haemophilia.

Renewal — (haemophilia with inhibitors) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for haemophilia with inhibitors; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment.

Initial application — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Fither:
 - 1.1 Patient has immune thrombocytopenic purpura* with a platelet count of less than or equal to 20,000 platelets per
 - 1.2 Patient has immune thrombocytopenic purpura* with a platelet count of 20,000 to 30,000 platelets per microlitre and significant mucocutaneous bleeding; and
- 2 Any of the following:
 - 2.1 Treatment with steroids and splenectomy have been ineffective; or
 - 2.2 Treatment with steroids has been ineffective and splenectomy is an absolute contraindication; or
 - 2.3 Other treatments including steroids have been ineffective and patient is being prepared for elective surgery (e.g. splenectomy); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (immune thrombocytopenic purpura (ITP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
 - 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for immune thrombocytopenic purpura*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 The patient has indolent low grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has indolent, low grade lymphoma or hairy cell leukaemia* requiring first-line systemic chemotherapy;
 - 2.2 To be used for a maximum of 6 treatment cycles.

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	✓	

continued...

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Renewal — (indolent, low-grade lymphomas or hairy cell leukaemia*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had a rituximab treatment-free interval of 12 months or more; and
- 2 The patient has indolent, low-grade NHL or hairy cell leukaemia* with relapsed disease following prior chemotherapy; and
- 3 To be used for no more than 6 treatment cycles.

Note: 'Indolent, low-grade lymphomas' includes follicular, mantle, marginal zone and lymphoplasmacytic/Waldenstrom macroglobulinaemia. *Unapproved indication. 'Hairy cell leukaemia' also includes hairy cell leukaemia variant.

Initial application — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient has autoimmune pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder.

Note: Indications marked with * are unapproved indications.

Renewal — (pure red cell aplasia (PRCA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 6 weeks where patient was previously treated with rituximab for pure red cell aplasia* associated with a demonstrable B-cell lymphoproliferative disorder and demonstrated an initial response lasting at least 12 months.

Note: Indications marked with * are unapproved indications.

Initial application — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: All of the following:

- 1 Patient has cold haemagglutinin disease*; and
- 2 Patient has severe disease which is characterized by symptomatic anaemia, transfusion dependence or disabling circulatory symptoms; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of

Note: Indications marked with * are unapproved indications.

Renewal — (severe cold haemagglutinin disease (CHAD)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Either:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for severe cold haemagglutinin disease*: and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Initial application — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Both:

- 1 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks: and
- 2 Fither:
 - 2.1 Patient has thrombotic thrombocytopenic purpura* and has experienced progression of clinical symptoms or persistent thrombocytopenia despite plasma exchange; or
 - 2.2 Patient has acute idiopathic thrombotic thrombocytopenic purpura* with neurological or cardiovascular pathology.

Note: Indications marked with * are unapproved indications.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (thrombotic thrombocytopenic purpura (TTP)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for thrombotic thrombocytopenic purpura*; and
- 2 An initial response lasting at least 12 months was demonstrated; and
- 3 Patient now requires repeat treatment; and
- 4 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Initial application — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 7 months for applications meeting the following criteria:

All of the following:

- 1 The patient has severe, immediately life- or organ-threatening SLE*; and
- 2 The disease has proved refractory to treatment with steroids at a dose of at least 1 mg/kg; and
- 3 The disease has relapsed following prior treatment for at least 6 months with maximal tolerated doses of azathioprine, mycophenolate mofetil and high dose cyclophosphamide, or cyclophosphamide is contraindicated; and
- 4 Maximum of four 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Renewal — (treatment refractory systemic lupus erythematosus (SLE)) only from a rheumatologist, nephrologist or Practitioner on the recommendation of a rheumatologist or nephrologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's SLE* achieved at least a partial response to the previous round of prior rituximab treatment; and
- 2 The disease has subsequently relapsed; and
- 3 Maximum of two 1000 mg infusions of rituximab.

Note: Indications marked with * are unapproved indications.

Initial application — (warm autoimmune haemolytic anaemia (warm AlHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has warm autoimmune haemolytic anaemia*: and
- 2 One of the following treatments has been ineffective: steroids (including if patient requires ongoing steroids at doses equivalent to > 5 mg prednisone daily), cytotoxic agents (e.g. cyclophosphamide monotherapy or in combination), intravenous immunoglobulin; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Note: Indications marked with * are unapproved indications.

Renewal — (warm autoimmune haemolytic anaemia (warm AIHA)) only from a haematologist or Practitioner on the recommendation of a haematologist. Approvals valid for 8 weeks for applications meeting the following criteria: Fither:

- 1 Previous treatment with lower doses of rituximab (100 mg weekly for 4 weeks) have proven ineffective and treatment with higher doses (375 mg/m² weekly for 4 weeks) is now planned; or
- 2 All of the following:
 - 2.1 Patient was previously treated with rituximab for warm autoimmune haemolytic anaemia*; and
 - 2.2 An initial response lasting at least 12 months was demonstrated; and
 - 2.3 Patient now requires repeat treatment.

Note: Indications marked with * are unapproved indications.

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

continued...

Initial application — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed antisynthetase syndrome; and
- 2 Patient has severe, immediately life or organ threatening disease, including interstitial lung disease; and
- 3 Fither
 - 3.1 Treatment with at least 3 immunosuppressants (oral steroids, cyclophosphamide, methotrexate, mycophenolate, ciclosporin, azathioprine) has not be effective at controlling active disease; or
 - 3.2 Rapid treatment is required due to life threatening complications; and
- 4 Maximum of four 1,000mg infusions of rituximab.

Renewal — (severe antisynthetase syndrome) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in inflammatory markers, muscle strength and pulmonary function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 Maximum of two cycles of 2 x 1,000mg infusions of rituximab given two weeks apart.

Initial application — (graft versus host disease) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has refractory graft versus host disease following transplant; and
- 2 Treatment with at least 3 immunosuppressants (oral steroids, ciclosporin, tacrolimus, mycophenolate, sirolimus) has not be effective at controlling active disease; and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m² of body surface area per week for a total of 4 weeks

Initial application — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Patient has severe chronic inflammatory demyelinating polyneuropathy (CIPD); and
- 2 Either:
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (severe chronic inflammatory demyelinating polyneuropathy) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function compared to baseline; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Initial application — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe anti-NMDA receptor autoimmune encephalitis: and
- 2 Fither
 - 2.1 Both:
 - 2.1.1 Treatment with steroids and intravenous immunoglobulin and/or plasma exchange has not been effective at controlling active disease; and
 - 2.1.2 At least one other immunosuppressant (cyclophosphamide, ciclosporin, tacrolimus, mycophenolate) has not been effective at controlling active disease; or
 - 2.2 Rapid treatment is required due to life threatening complications; and
- 3 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Renewal — (anti-NMDA receptor autoimmune encephalitis) only from a neurologist or medical practitioner on the recommendation of a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient's disease has responded to the previous rituximab treatment with demonstrated improvement in neurological function; and
- 2 The patient has not received rituximab in the previous 6 months; and
- 3 The patient has experienced a relapse and now requires further treatment; and
- 4 One of the following dose regimens is to be used: 375 mg/m2 of body surface area per week for a total of four weeks, or 500 mg once weekly for four weeks, or two 1,000 mg doses given two weeks apart.

Initial application — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 9 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 The patient has CD20+ low grade or follicular B-cell NHL with relapsed disease following prior chemotherapy; and
 - 1.2 To be used for a maximum of 6 treatment cycles; or
- 2 Both:
 - 2.1 The patient has CD20+ low grade or follicular B-cell NHL requiring first-line systemic chemotherapy; and
 - 2.2 To be used for a maximum of 6 treatment cycles.

Renewal — (CD20+ low grade or follicular B-cell NHL) from any relevant practitioner. Approvals valid for 24 months for applications meeting the following criteria:

Both:

- 1 Rituximab is to be used for maintenance in CD20+ low grade or follicular B-cell NHL following induction with first-line systemic chemotherapy; and
- 2 Patient is intended to receive rituximab maintenance therapy for 2 years at a dose of 375 mg/m2 every 8 weeks (maximum of 12 cycles).

Initial application — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has biopsy-proven primary/idiopathic membranous nephropathy*; or
 - 1.2 Patient has PLA2 antibodies with no evidence of secondary cause, and an eGFR of > 60ml/min/1.73m2; and
- 2 Patient remains at high risk of progression to end-stage kidney disease despite more than 3 months of treatment with conservative measures (see Note): and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
	Por 🗸	Manufacturor

continued...

3 The total rituximab dose would not exceed the equivalent of 375mg/m2 of body surface area per week for a total of 4 weeks

Renewal — (Membranous nephropathy) only from a nephrologist or any relevant practitioner on the recommendation of a nephrologist. Approvals valid for 6 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient was previously treated with rituximab for membranous nephropathy*; and
- 2 Either:
 - 2.1 Treatment with rituximab was previously successful, but the condition has relapsed, and the patient now requires repeat treatment; or
 - 2.2 Patient achieved partial response to treatment and requires repeat treatment (see Note); and
- 3 The total rituximab dose used would not exceed the equivalent of 375 mg/m2 of body surface area per week for a total of 4 weeks.

Notes:

- a) Indications marked with * are unapproved indications.
- b) High risk of progression to end-stage kidney disease defined as > 5g/day proteinuria.
- c) Conservative measures include renin-angiotensin system blockade, blood-pressure management, dietary sodium and protein restriction, treatment of dyslipidaemia, and anticoagulation agents unless contraindicated or the patient has experienced intolerable side effects.
- d) Partial response defined as a reduction of proteinuria of at least 50% from baseline, and between 0.3 grams and 3.5 grams per 24 hours.

Initial application — (B-cell acute lymphoblastic leukaemia/lymphoma*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria:

All of the following:

- 1 Patient has newly diagnosed B-cell acute lymphoblastic leukaemia/lymphoma*; and
- 2 Treatment must be in combination with an intensive chemotherapy protocol with curative intent; and
- 3 The total rituximab dose would not exceed the equivalent of 375 mg/m² per dose for a maximum of 18 doses.

Note: Indications marked with * are unapproved indications.

Initial application — (desensisation prior to transplant) from any relevant practitioner. Approvals valid for 6 weeks for applications meeting the following criteria:

Both:

- 1 Patient requires desensitisation prior to mismatched allogenic stem cell transplant*; and
- 2 Patient would receive no more than two doses at 375 mg/m2 of body-surface area.

Note: Indications marked with * are unapproved indications.

Initial application — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Patient has severe rapidly progressive pemphigus; and
 - 1.2 Is used in combination with systemic corticosteroids (20 mg/day); and
 - 1.3 Any of the following:
 - 1.3.1 Skin involvement is at least 5% body surface area; or
 - 1.3.2 Significant mucosal involvement (10 or more mucosal erosions) or diffuse gingivitis or confluent large erosions; or
 - 1.3.3 Involvement of two or more mucosal sites; or
- 2 Both:
 - 2.1 Patient has pemphigus; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

2.2 Patient has not experienced adequate clinical benefit from systemic corticosteroids (20 mg/day) in combination with a steroid sparing agent, unless contraindicated.

Note: Indications marked with * are unapproved indications.

Renewal — (pemiphigus*) only from a dermatologist or relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has experienced adequate clinical benefit from rituximab treatment, with improvement in symptoms and healing of skin ulceration and reduction in corticosteroid requirement; and
- 2 Patient has not received rituximab in the previous 6 months.

Note: Indications marked with * are unapproved indications.

SECUKINUMAB - Special Authority see SA2084 below - Retail pharmacy

⇒SA2084 Special Authority for Subsidy

Initial application — (severe chronic plaque psoriasis – second-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab or etanercept, or has trialled infliximab in a Health NZ Hospital, for severe chronic plaque psoriasis; and
- 2 Fither:
 - 2.1 The patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 2.2 The patient has received insufficient benefit from adalimumab, etanercept or infliximab; and
- 3 A Psoriasis Area and Severity Index (PASI) assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Initial application — (severe chronic plaque psoriasis – first-line biologic) only from a dermatologist. Approvals valid for 4 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient has "whole body" severe chronic plaque psoriasis with a Psoriasis Area and Severity Index (PASI) score of greater than 10, where lesions have been present for at least 6 months from the time of initial diagnosis; or
 - 1.2 Patient has severe chronic plaque psoriasis of the face, or palm of a hand or sole of a foot, where the plaque or plaques have been present for at least 6 months from the time of initial diagnosis; and
- 2 Patient has tried, but had an inadequate response (see Note) to, or has experienced intolerable side effects from, at least three of the following (at maximum tolerated doses unless contraindicated): phototherapy, methotrexate, ciclosporin, or acitretin: and
- 3 A PASI assessment or Dermatology Quality of Life Index (DLQI) assessment has been completed for at least the most recent prior treatment course, preferably while still on treatment but no longer than 1 month following cessation of each prior treatment course; and
- 4 The most recent PASI or DQLI assessment is no more than 1 month old at the time of application.

Note: A treatment course is defined as a minimum of 12 weeks of treatment. "Inadequate response" is defined as: for whole body severe chronic plaque psoriasis, a PASI score of greater than 10, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment; for severe chronic plaque psoriasis of the face, hand or foot, at least 2 of the 3 PASI symptom sub scores for erythema, thickness and scaling are rated as severe or very severe, and the skin area affected is 30% or more of the face, palm of a hand or sole of a foot, as assessed preferably while still on treatment but no longer than 1 month following cessation of the most recent prior treatment.

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per	✓	Manufacturer	

continued...

Renewal — (severe chronic plaque psoriasis – first and second-line biologic) only from a dermatologist or medical practitioner on the recommendation of a dermatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Fither:
 - 1.1 Patient's PASI score has reduced by 75% or more (PASI 75) as compared to baseline PASI prior to commencing secukinumab: or
 - 1.2 Patient has a Dermatology Quality of Life Index (DLQI) improvement of 5 or more, as compared to baseline DLQI prior to commencing secukinumab; and
- 2 Secukinumab to be administered at a maximum dose of 300 mg monthly.

Initial application — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 3 months for applications meeting the following criteria: Both:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for ankylosing spondylitis; and
- - 2.1 The patient has experienced intolerable side effects from a reasonable trial of adalimumab and/or etanercept; or
 - 2.2 Following 12 weeks of adalimumab and/or etanercept treatment, the patient did not meet the renewal criteria for adalimumab and/or etanercept for ankylosing spondylitis.

Renewal — (ankylosing spondylitis – second-line biologic) only from a rheumatologist or medical practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria: All of the following:

- 1 Following 12 weeks initial treatment of secukinumab treatment, BASDAI has improved by 4 or more points from pre-secukinumab baseline on a 10 point scale, or by 50%, whichever is less; and
- 2 Physician considers that the patient has benefitted from treatment and that continued treatment is appropriate; and
- 3 Secukinumab to be administered at doses no greater than 150 mg monthly.

Initial application — (psoriatic arthritis) only from a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Both:
 - 1.1 Patient has had an initial Special Authority approval for adalimumab, etanercept or infliximab for psoriatic arthritis;
 - 1.2 Either:
 - 1.2.1 Patient has experienced intolerable side effects from adalimumab, etanercept or infliximab; or
 - 1.2.2 Patient has received insufficient benefit from adalimumab, etanercept or infliximab to meet the renewal criteria for adalimumab, etanercept or infliximab for psoriatic arthritis; or
- 2 All of the following:
 - 2.1 Patient has had severe active psoriatic arthritis for six months duration or longer; and
 - 2.2 Patient has tried and not responded to at least three months of oral or parenteral methotrexate at a dose of at least 20 mg weekly or a maximum tolerated dose; and
 - 2.3 Patient has tried and not responded to at least three months of sulfasalazine at a dose of at least 2 g per day or leflunomide at a dose of up to 20 mg daily (or maximum tolerated doses); and
 - 2.4 Either:
 - 2.4.1 Patient has persistent symptoms of poorly controlled and active disease in at least 15 swollen, tender joints;
 - 2.4.2 Patient has persistent symptoms of poorly controlled and active disease in at least four joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
 - 2.5 Any of the following:

Subsidy		Fully	Brand or	
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 2.5.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
- 2.5.2 Patient has an elevated erythrocyte sedimentation rate (ESR) greater than 25 mm per hour; or
- 2.5.3 ESR and CRP not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Renewal — (psoriatic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
 - 1.2 The patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to prior secukinumab treatment in the opinion of the treating physician; and
- 2 Secukinumab to be administered at doses no greater than 300 mg monthly.

SILTUXIMAB - Special Authority see SA1596 below - Retail pharmacy

Note: Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Inj 100 mg vial	770.57	1	Sylvant
Inj 400 mg vial	3,082.33	1	✓ Sylvant

⇒SA1596 Special Authority for Subsidy

Initial application only from a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's Disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Siltuximab is to be administered at doses no greater than 11 mg/kg every 3 weeks.

Renewal only from a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has sustained improvement in inflammatory markers and functional status.

TIXAGEVIMAB WITH CILGAVIMAB - [Xpharm] - Subsidy by endorsement

4. Francisco de La della collega describa 400 como conse

- a) No patient co-payment payable
- b) Treatment is funded only if patient meets access criteria for tixagevimab with cilgavimab (as per https://pharmac.govt.nz/Evusheld) and has been endorsed accordingly by the prescriber. The supply of treatment is via Pharmac's approved distribution process. Refer to the Pharmac website for more information about this and stock availability.

Inj 100 mg per mi, 1.5 ml viai with cilgavimab 100 mg per ml,1.5 ml viai	.0.00	1	✓ Evusheld
TOCILIZUMAB - PCT only - Special Authority see SA2159 on the next	page		
Inj 20 mg per ml, 4 ml vial2	20.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
8	30.00	4	✓ RoActemra S29 S29
Inj 20 mg per ml, 10 ml vial5	50.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
Inj 20 mg per ml, 20 ml vial1,1	00.00	1	✓ Actemra
			✓ Actemra S29 S29
			✓ RoActemra S29 S29
4,4'	00.00	4	✓ RoActemra S29 S29
Inj 1 mg for ECP	.2.85	1 mg	✓ Baxter

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

⇒SA2159 Special Authority for Subsidy

Initial application — (cytokine release syndrome) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 The patient is enrolled in the Children's Oncology Group AALL1731 trial; and
 - 1.2 The patient has developed grade 3 or 4 cytokine release syndrome associated with the administration of blinatumomab for the treatment of acute lymphoblastic leukaemia; and
 - 1.3 Tocilizumab is to be administered at doses no greater than 8 mg/kg IV for a maximum of 3 doses (if less than 30kg, maximum of 12 mg/kg); or
- 2 All of the following:
 - 2.1 The patient is enrolled in the Malaghan Institute of Medical Research Phase I ENABLE trial; and
 - 2.2 The patient has developed CRS or CAR T-Cell Related Encephalopathy Syndrome (CRES) associated with the administration of CAR T-cell therapy for the treatment of relapsed or refractory B-cell non-Hodgkin lymphoma; and
 - 2.3 Tocilizumab is to be administered according to the consensus guidelines for CRS and CRES for CAR T-cell therapy (Neelapu et al. Nat Rev Clin Oncol 2018;15:47-62) at doses no greater than 8 mg/kg IV for a maximum of 3 doses.

Initial application — (previous use) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient was being treated with tocilizumab prior to 1 February 2019; and
- 2 Any of the following:
 - 2.1 rheumatoid arthritis: or
 - 2.2 systemic juvenile idiopathic arthritis; or
 - 2.3 adult-onset Still's disease; or
 - 2.4 polyarticular juvenile idiopathic arthritis; or
 - 2.5 idiopathic multicentric Castleman's disease.

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Initial application — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

1 Patient has had severe and active erosive rheumatoid arthritis (either confirmed by radiology imaging, or the patient is cyclic citrullinated peptide (CCP) antibody positive) for six months duration or longer; and

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subsidised		Generic	
\$	Per	✓	Manufacturer	

continued...

- 2 Tocilizumab is to be used as monotherapy; and
- 3 Fither:
 - 3.1 Treatment with methotrexate is contraindicated; or
 - 3.2 Patient has tried and did not tolerate oral and/or parenteral methotrexate; and
- 4 Fither:
 - 4.1 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of ciclosporin alone or in combination with another agent; or
 - 4.2 Patient has tried and not responded to at least three months therapy at the maximum tolerated dose of leflunomide alone or in combination with another agent; and
- 5 Either:
 - 5.1 Patient has persistent symptoms of poorly controlled and active disease in at least 20 active, swollen, tender joints; or
 - 5.2 Patient has persistent symptoms of poorly controlled and active disease in at least four active joints from the following: wrist, elbow, knee, ankle, and either shoulder or hip; and
- 6 Either:
 - 6.1 Patient has a C-reactive protein level greater than 15 mg/L measured no more than one month prior to the date of this application; or
 - 6.2 C-reactive protein levels not measured as patient is currently receiving prednisone therapy at a dose of greater than 5 mg per day and has done so for more than three months.

Initial application — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient diagnosed with systemic juvenile idiopathic arthritis; and
- 2 Patient has tried and not responded to a reasonable trial of all of the following, either alone or in combination: oral or parenteral methotrexate; non-steroidal anti-inflammatory drugs (NSAIDs); and systemic corticosteroids.

Initial application — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Fither:
 - 1.1.1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for adult-onset Still's disease (AOSD); or
 - 1.1.2 The patient has been started on tocilizumab for AOSD in a Health NZ Hospital; and
 - 1.2 Either:
 - 1.2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 1.2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for AOSD; or
- 2 All of the following:
 - 2.1 Patient diagnosed with AOSD according to the Yamaguchi criteria (J Rheumatol 1992;19:424-430); and
 - 2.2 Patient has tried and not responded to at least 6 months of glucocorticosteroids at a dose of at least 0.5 mg/kg, non-steroidal antiinflammatory drugs (NSAIDs) and methotrexate; and
 - 2.3 Patient has persistent symptoms of disabling poorly controlled and active disease.

Initial application — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 4 months for applications meeting the following criteria: Either:

1 Both:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
 \$	Per	√	

continued...

- 1.1 The patient has had an initial Special Authority approval for both etanercept and adalimumab for polyarticular course juvenile idiopathic arthritis (JIA); and
- 1.2 The patient has experienced intolerable side effects, or has received insufficient benefit from, both etanercept and adalimumab; or
- 2 All of the following:
 - 2.1 Treatment with a tumour necrosis factor alpha inhibitor is contraindicated; and
 - 2.2 Patient has had polyarticular course JIA for 6 months duration or longer; and
 - 2.3 To be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
 - 2.4 Any of the following:
 - 2.4.1 At least 5 active joints and at least 3 joints with limited range of motion, pain or tenderness after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.2 Moderate or high disease activity (cJADAS10 score of at least 2.5) after a 3-month trial of methotrexate (at the maximum tolerated dose); or
 - 2.4.3 Low disease activity (cJADAS10 score between 1.1 and 2.5) after a 6-month trial of methotrexate.

Initial application — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe HHV-8 negative idiopathic multicentric Castleman's disease; and
- 2 Treatment with an adequate trial of corticosteroids has proven ineffective; and
- 3 Tocilizumab to be administered at doses no greater than 8 mg/kg IV every 3-4 weeks.

Initial application — (moderate to severe COVID-19) from any relevant practitioner. Approvals valid for 4 weeks for applications meeting the following criteria:

All of the following:

- 1 Patient has confirmed (or probable) COVID-19; and
- 2 Oxygen saturation of < 92% on room air, or requiring supplemental oxygen; and
- 3 Patient is receiving adjunct systemic corticosteroids, or systemic corticosteroids are contraindicated; and
- 4 Tocilizumab is to be administered at doses no greater than 8mg/kg IV for a maximum of one dose; and
- 5 Tocilizumab is not to be administered in combination with barcitinib.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Fither:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Renewal — (systemic juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following up to 6 months' initial treatment, the patient has achieved at least an American College of Rheumatology paediatric 30% improvement criteria (ACR Pedi 30) response from baseline; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing ACR Pedi 30 response from baseline.

Renewal — (adult-onset Still's disease) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months where the patient has a sustained improvement in inflammatory markers and functional status. Renewal — (polyarticular juvenile idiopathic arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer
--

continued...

Both:

- 1 Treatment is to be used as an adjunct to methotrexate therapy or monotherapy where use of methotrexate is limited by toxicity or intolerance; and
- 2 Fither:
 - 2.1 Following 3 to 4 months' initial treatment, the patient has at least a 50% decrease in active joint count and an improvement in physician's global assessment from baseline; or
 - 2.2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count and continued improvement in physician's global assessment from baseline.

Renewal — (idiopathic multicentric Castleman's disease) only from a haematologist, rheumatologist or Practitioner on the recommendation of a haematologist or rheumatologist. Approvals valid for 12 months where the treatment remains appropriate and the patient has a sustained improvement in inflammatory markers and functional status.

TRASTUZUMAB - PCT only - Specialist - Special Authority see SA1632 below

Inj 150 mg vial	1,350.00	1	✓ Herceptin
Inj 440 mg vial	3,875.00	1	✓ Herceptin
Inj 1 mg for ECP	· ·	1 mg	✓ Baxter

⇒SA1632 Special Authority for Subsidy

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Either:
 - 2.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 2.2 Both:
 - 2.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 2.2.2 The cancer did not progress whilst on lapatinib; and
- 3 Either:
 - 3.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 3.2 All of the following:
 - 3.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 3.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer; and
 - 3.2.3 The patient has good performance status (ECOG grade 0-1); and
- 4 Trastuzumab not to be given in combination with lapatinib; and
- 5 Trastuzumab to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 3 Trastuzumab not to be given in combination with lapatinib; and
- 4 Trastuzumab to be discontinued at disease progression.

Initial application — **(early breast cancer)** only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 15 months for applications meeting the following criteria:

Subsidy	Fully		Brand or
(Manufacturer's Price)	Subsidised		Generic
<u> </u>	Per	1	Manufacturer

continued...

All of the following:

- 1 The patient has early breast cancer expressing HER 2 IHC 3+ or ISH + (including FISH or other current technology); and
- 2 Maximum cumulative dose of 106 mg/kg (12 months' treatment); and
- 3 Any of the following:
 - 3.1 9 weeks' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.2 12 months' concurrent treatment with adjuvant chemotherapy is planned; or
 - 3.3 12 months' sequential treatment following adjuvant chemotherapy is planned; or
 - 3.4 12 months' treatment with neoadjuvant and adjuvant chemotherapy is planned; or
 - 3.5 Other treatment regimen, in association with adjuvant chemotherapy, is planned.

Renewal — (early breast cancer*) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 The patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 The patient received prior adjuvant trastuzumab treatment for early breast cancer; and
- 3 Any of the following:
 - 3.1 The patient has not previously received lapatinib treatment for HER-2 positive metastatic breast cancer; or
 - 3.2 Both:
 - 3.2.1 The patient started lapatinib treatment for metastatic breast cancer but discontinued lapatinib within 3 months of starting treatment due to intolerance; and
 - 3.2.2 The cancer did not progress whilst on lapatinib; or
 - 3.3 The cancer has not progressed at any time point during the previous 12 months whilst on trastuzumab; and
- 4 Fither:
 - 4.1 Trastuzumab will not be given in combination with pertuzumab; or
 - 4.2 All of the following:
 - 4.2.1 Trastuzumab to be administered in combination with pertuzumab; and
 - 4.2.2 Patient has not received prior treatment for their metastatic disease and has had a treatment-free interval of at least 12 months between prior (neo)adjuvant chemotherapy treatment and diagnosis of metastatic breast cancer: and
 - 4.2.3 The patient has good performance status (ECOG grade 0-1); and
- 5 Trastuzumab not to be given in combination with lapatinib; and
- 6 Trastuzumab to be discontinued at disease progression.

Note: * For patients with relapsed HER-2 positive disease who have previously received adjuvant trastuzumab for early breast cancer.

TRASTUZUMAB EMTANSINE - PCT only - Specialist - Special Authority see SA2144 below

Inj 100 mg vial	 2,320.00	1	✓ Kadcyla
Inj 160 mg vial	 3,712.00	1	✓ Kadcyla
Inj 1 mg for ECP	 24.52	1 mg	✓ Baxter

⇒SA2144 Special Authority for Subsidy

Initial application — (early breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has early breast cancer expressing HER2 IHC3+ or ISH+; and
- 2 Documentation of pathological invasive residual disease in the breast and/or auxiliary lymph nodes following completion of surgery; and
- 3 Patient has completed systemic neoadjuvant therapy with trastuzumab and chemotherapy prior to surgery; and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

- 4 Disease has not progressed during neoadjuvant therapy; and
- 5 Patient has left ventricular ejection fraction of 45% or greater; and
- 6 Adjuvant treatment with trastuzumab emtansine to be commenced within 12 weeks of surgery; and
- 7 Trastuzumab emtansine to be discontinued at disease progression; and
- 8 Total adjuvant treatment duration must not exceed 42 weeks (14 cycles).

Initial application — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has metastatic breast cancer expressing HER-2 IHC 3+ or ISH+ (including FISH or other current technology); and
- 2 Patient has previously received trastuzumab and chemotherapy, separately or in combination; and
- 3 Either:
 - 3.1 The patient has received prior therapy for metastatic disease*; or
 - 3.2 The patient developed disease recurrence during, or within six months of completing adjuvant therapy*; and
- 4 Patient has a good performance status (ECOG 0-1); and
- 5 Either:
 - 5.1 Patient does not have symptomatic brain metastases; or
 - 5.2 Patient has brain metastases and has received prior local CNS therapy; and
- 6 Patient has not received prior funded trastuzumab emtansine treatment; and
- 7 Treatment to be discontinued at disease progression.

Renewal — (metastatic breast cancer) only from a relevant specialist or medical practitioner on the recommendation of a relevant specialist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 The cancer has not progressed at any time point during the previous approval period whilst on trastuzumab emtansine; and
- 2 Treatment to be discontinued at disease progression.

Note: *Note: Prior or adjuvant therapy includes anthracycline, other chemotherapy, biological drugs, or endocrine therapy.

Programmed Cell Death-1 (PD-1) Inhibitors

DURVALUMAB - PCT only - Specialist - Special Authority se	ee SA2147 below		
Inj 50 mg per ml, 10 ml vial	4,700.00	1	Imfinzi
Inj 50 mg per ml, 2.4 ml vial	1,128.00	1	Imfinzi
Inj 1 mg for ECP	9.59	1 mg	✓ Baxter

⇒SA2147 Special Authority for Subsidy

Initial application — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria: Either:

- 1 Patient is currently on treatment with durvalumab and met all remaining criteria (criterion 2) below prior to commencing treatment; or
- 2 All of the following:
 - 2.1 Patient has histologically or cytologically documented stage III, locally advanced, unresectable non-small cell lung cancer (NSCLC); and
 - 2.2 Patient has received two or more cycles of platinum-based chemotherapy concurrently with definitive radiation therapy; and
 - 2.3 Patient has no disease progression following the second or subsequent cycle of platinum-based chemotherapy with definitive radiation therapy treatment; and
 - 2.4 Patient has a ECOG performance status of 0 or 1; and

Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidi	sed	Generic	
\$	Per	✓	Manufacturer	

continued...

- 2.5 Patient has completed last radiation dose within 8 weeks of starting treatment with durvalumab; and
- 2.6 Patient must not have received prior PD-1 or PD-L1 inhibitor therapy for this condition; and
- 2.7 Either:
 - 2.7.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.7.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 2.8 Treatment with durvalumab to cease upon signs of disease progression.

Renewal — (Non-small cell lung cancer) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 The treatment remains clinically appropriate and the patient is benefitting from treatment; and
- 2 Either:
 - 2.1 Durvalumab is to be used at a maximum dose of no greater than 10 mg/kg every 2 weeks; or
 - 2.2 Durvalumab is to be used at a flat dose of 1500 mg every 4 weeks; and
- 3 Treatment with durvalumab to cease upon signs of disease progression; and
- 4 Total continuous treatment duration must not exceed 12 months.

NIVOLUMAB - PCT	only – Specialist – Special Authority see SA	2120 below
Ini 10 ma nor ml	1 ml vial	1 051 08

Inj 10 mg per ml, 4 ml vial	1,051.98	1	Opdivo
Inj 10 mg per ml, 10 ml vial	2,629.96	1	Opdivo
Inj 1 mg for ECP	27.62	1 mg	Baxter

⇒SA2120 Special Authority for Subsidy

Initial application only from a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded pembrolizumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for pembrolizumab and has discontinued pembrolizumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on pembrolizumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with nivolumab will not be continued if their disease progresses.

Renewal only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria:

Either:

- 1 All of the following:
 - 1.1 Any of the following:
 - 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
 - 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
 - 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
 - 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
 - 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
 - 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or

Subsidy (Manufacturer's Price)	Subsi	Fully	Brand or Generic
 \$	Per	1	Manufacturer

continued...

- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with nivolumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with nivolumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

PEMBROLIZUMAB - PCT only - Specialist - Special A	Authority see SA2121 below		
Inj 25 mg per ml, 4 ml vial	4,680.00	1	✓ Keytruda
Inj 1 mg for ECP	49.14	1 mg	✓ Baxter

⇒SA2121 Special Authority for Subsidy

Initial application — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: All of the following:

- 1 Patient has metastatic or unresectable melanoma (excluding uveal) stage III or IV; and
- 2 Patient has measurable disease as defined by RECIST version 1.1; and
- 3 The patient has ECOG performance score of 0-2; and
- 4 Either:
 - 4.1 Patient has not received funded nivolumab; or
 - 4.2 Both:
 - 4.2.1 Patient has received an initial Special Authority approval for nivolumab and has discontinued nivolumab within 12 weeks of starting treatment due to intolerance; and
 - 4.2.2 The cancer did not progress while the patient was on nivolumab; and
- 5 Baseline measurement of overall tumour burden is documented (see Note); and
- 6 Documentation confirming that the patient has been informed and acknowledges that funded treatment with pembrolizumab will not be continued if their disease progresses.

Renewal — (unresectable or metastatic melanoma) only from a medical oncologist or medical practitioner on the recommendation of a medical oncologist. Approvals valid for 4 months for applications meeting the following criteria: Fither:

- 1 All of the following:
 - 1.1 Any of the following:

Subsidy (Manufacturer's Pri	ice)	Fu Subsidis	,	Brand or Generic
\$	P	er	/	Manufacturer

continued...

- 1.1.1 Patient's disease has had a complete response to treatment according to RECIST criteria (see Note); or
- 1.1.2 Patient's disease has had a partial response to treatment according to RECIST criteria (see Note); or
- 1.1.3 Patient has stable disease according to RECIST criteria (see Note); and
- 1.2 Patient's disease has not progressed clinically and disease response to treatment has been clearly documented in patient notes; and
- 1.3 No evidence of progressive disease according to RECIST criteria (see Note); and
- 1.4 The treatment remains clinically appropriate and the patient is benefitting from the treatment; or
- 2 All of the following:
 - 2.1 Patient has previously discontinued treatment with pembrolizumab for reasons other than severe toxicity or disease progression; and
 - 2.2 Patient has signs of disease progression; and
 - 2.3 Disease has not progressed during previous treatment with pembrolizumab.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer EA, et al. Eur J Cancer 2009;45:228-47). Assessments of overall tumour burden and measurable disease to be undertaken on a minimum of one lesion and maximum of 5 target lesions (maximum two lesions per organ). Target lesions should be selected on the basis of their size (lesions with the longest diameter), be representative of all involved organs, and suitable for reproducible repeated measurements. Measurable disease includes by CT or MRI imaging or caliper measurement by clinical exam. Target lesion measurements should be assessed using the same method of assessment and the same technique used to characterise each identified and reported lesion at baseline and every 12 weeks. Response definitions as follows:

- Complete Response: Disappearance of all target lesions. Any pathological lymph nodes (whether target or non-target)
 must have reduction in short axis to < 10 mm.
- Partial Response: At least a 30% decrease in the sum of diameters of target lesions, taking as reference the baseline sum diameters.
- Progressive Disease: At least a 20% increase in the sum of diameters of target lesions, taking as reference the smallest sum on study (this includes the baseline sum if that is the smallest on study). In addition to the relative increase of 20%, the sum must also demonstrate an absolute increase of at least 5 mm. (Note: the appearance of one or more new lesions is also considered progression).
- Stable Disease: Neither sufficient shrinkage to qualify for partial response nor sufficient increase to qualify for progressive disease.

Other Immunosuppressants

CICLOSPORIN			
Cap 25 mg	44.63	50	✓ Neoral
Cap 50 mg	88.91	50	✓ Neoral
Cap 100 mg	177.81	50	✓ Neoral
Oral liq 100 mg per ml	198.13	50 ml OP	✓ Neoral
EVEROLIMUS - Special Authority see SA2008 below - Retail phar	macy		
Wastage claimable			
Tab 10 mg	6,512.29	30	✓ Afinitor
Tab 5 mg	4,555.76	30	✓ Afinitor

⇒SA2008 Special Authority for Subsidy

Initial application only from a neurologist or oncologist. Approvals valid for 3 months for applications meeting the following criteria:

Both:

1 Patient has tuberous sclerosis: and

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
 \$	Per 🗸	Manufacturer

continued...

2 Patient has progressively enlarging sub-ependymal giant cell astrocytomas (SEGAs) that require treatment.

Renewal only from a neurologist or oncologist. Approvals valid for 12 months for applications meeting the following criteria: All of the following:

- 1 Documented evidence of SEGA reduction or stabilisation by MRI within the last 3 months; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 Everolimus to be discontinued at progression of SEGAs.

SIROLIMUS - Special Authority see SA2005 below - Retail pharmacy

Tab 1 mg	749.99	100	Rapamune
Tab 2 mg	1,499.99	100	✓ Rapamune
Oral liq 1 mg per ml	449.99	60 ml OP	✓ Rapamune

⇒SA2005 Special Authority for Subsidy

Initial application from any medical practitioner. Approvals valid without further renewal unless notified where the drug is to be used for rescue therapy for an organ transplant recipient.

Notes: Rescue therapy defined as unresponsive to calcineurin inhibitor treatment as defined by refractory rejection; or intolerant to calcineurin inhibitor treatment due to any of the following:

- GFR< 30 ml/min: or
- Rapidly progressive transplant vasculopathy; or
- Rapidly progressive obstructive bronchiolitis; or
- . HUS or TTP; or
- · Leukoencepthalopathy; or
- · Significant malignant disease

Initial application — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has severe non-malignant lymphovascular malformation*; and
- 2 Any of the following:
 - 2.1 Malformations are not adequately controlled by sclerotherapy and surgery; or
 - 2.2 Malformations are widespread/extensive and sclerotherapy and surgery are not considered clinically appropriate; or
 - 2.3 Sirolimus is to be used to reduce malformation prior to consideration of surgery; and
- 3 Patient is being treated by a specialist lymphovascular malformation multi-disciplinary team; and
- 4 Patient has measurable disease as defined by RECIST version 1.1 (see Note).

Renewal — (severe non-malignant lymphovascular malformations*) from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Either:
 - 1.1 Patient's disease has had either a complete response or a partial response to treatment, or patient has stable disease according to RECIST version 1.1 (see Note); or
 - 1.2 Patient's disease has stabilised or responded clinically and disease response to treatment has been clearly documents in patient notes; and
- 2 No evidence of progressive disease; and
- 3 The treatment remains clinically appropriate and the patient is benefitting from the treatment.

Notes: Baseline assessment and disease responses to be assessed according to the Response Evaluation Criteria in Solid Tumours (RECIST) version 1.1 (Eisenhauer et al. Eur J Cancer 2009;45:228-47)

Indications marked with * are unapproved indications

Initial application — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) only from a nephrologist or urologist. Approvals valid for 6 months for applications meeting the following criteria:

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	✓	Manufacturer

continued...

Both:

- 1 Patient has tuberous sclerosis complex*; and
- 2 Evidence of renal angiomyolipoma(s) measuring 3 cm or greater and that have shown interval growth.

Renewal — (renal angiomyolipoma(s) associated with tuberous sclerosis complex*) from any relevant practitioner.

Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Documented evidence of renal angiomyolipoma reduction or stability by magnetic resonance imaging (MRI) or ultrasound; and
- 2 Demonstrated stabilisation or improvement in renal function; and
- 3 The patient has not experienced angiomyolipoma haemorrhage or significant adverse effects to sirolimus treatment; and
- 4 The treatment remains appropriate and the patient is benefitting from treatment.

Note: Indications marked with * are unapproved indications

Initial application — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has epilepsy with a background of documented tuberous sclerosis complex; and
- Patient nas 2 Either:
 - 2.1 Both:
 - 2.1.1 Vigabatrin has been trialled and has not adequately controlled seizures; and
 - 2.1.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least two of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); or
 - 2.2 Both:
 - 2.2.1 Vigabatrin is contraindicated; and
 - 2.2.2 Seizures are not adequately controlled by, or the patient has experienced unacceptable side effects from, optimal treatment with at least three of the following: sodium valproate, topiramate, levetiracetam, carbamazepine, lamotrigine, phenytoin sodium, and lacosamide (see Note); and
- 3 Seizures have a significant impact on quality of life; and
- 4 Patient has been assessed and surgery is considered inappropriate for this patient, or the patient has been assessed and would benefit from mTOR inhibitor treatment prior to surgery.

Note: Patients of childbearing potential are not required to have a trial of sodium valporate

Renewal — (refractory seizures associated with tuberous sclerosis complex*) only from a neurologist. Approvals valid for 12 months where demonstrated significant and sustained improvement in seizure rate (e.g. 50% reduction in seizure frequency) or severity and/or patient quality of life compared with baseline prior to starting sirolimus treatment.

Note: Indications marked with * are unapproved indications

TACROLIMUS - Special Authority see SA1745 below - Retail pharmacy

		iow i iciali priarriacy	Of IOLINIOO Opecial Authority 300 OA 1743 belov
✓ Tacrolimus Sandoz	100	49.60	Cap 0.5 mg
✓ Tacrolimus Sandoz	100		Cap 0.75 mg
✓ Tacrolimus Sandoz	100		Cap 1 mg
✓ Tacrolimus Sandoz	50		Cap 5 mg
			- 1 - 3

⇒SA1745 Special Authority for Subsidy

Initial application — (organ transplant) only from a relevant specialist. Approvals valid without further renewal unless notified where the patient is an organ transplant recipient.

Note: Subsidy applies for either primary or rescue therapy.

Initial application — (non-transplant indications*) only from a relevant specialist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Both:

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Su	bsidised	Generic	
\$	Per	•	Manufacturer	

continued...

- 1 Patient requires long-term systemic immunosuppression; and
- 2 Ciclosporin has been trialled and discontinued treatment because of unacceptable side effects or inadequate clinical response.

Note: Indications marked with * are unapproved indications

JAK inhibitors

UPADACITINIB – Special Authority see SA2079 below – Retail pharmacy
Tab 15 mg1,271.00 28 ✓ RINVOQ

⇒SA2079 Special Authority for Subsidy

Initial application — (Rheumatoid Arthritis (patients previously treated with adalimumab or etanercept)) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 The patient has had an initial Special Authority approval for adalimumab and/or etanercept for rheumatoid arthritis; and
- 2 Either:
 - 2.1 The patient has experienced intolerable side effects from adalimumab and/or etanercept; or
 - 2.2 The patient has received insufficient benefit from at least a three-month trial of adalimumab and/or etanercept such that they do not meet the renewal criteria for rheumatoid arthritis; and
- 3 Either:
 - 3.1 The patient is seronegative for both anti-cyclic citrullinated peptide (CCP) antibodies and rheumatoid factor; or
 - 3.2 Both:
 - 3.2.1 The patient has been started on rituximab for rheumatoid arthritis in a Health NZ Hospital; and
 - 3.2.2 Either:
 - 3.2.2.1 The patient has experienced intolerable side effects from rituximab; or
 - 3.2.2.2 At four months following the initial course of rituximab the patient has received insufficient benefit such that they do not meet the renewal criteria for rheumatoid arthritis.

Renewal — (Rheumatoid Arthritis) only from a rheumatologist or Practitioner on the recommendation of a rheumatologist. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Following 6 months' initial treatment, the patient has at least a 50% decrease in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician; or
- 2 On subsequent reapplications, the patient demonstrates at least a continuing 30% improvement in active joint count from baseline and a clinically significant response to treatment in the opinion of the physician.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per ✓ Manufacturer

Antiallergy Preparations

Allergic Emergencies

ICATIBANT - Special Authority see SA1558 below - Retail pharmacy

Inj 10 mg per ml, 3 ml prefilled syringe..................2,668.00 1 ✓ Firazyr

⇒SA1558 Special Authority for Subsidy

Initial application only from a clinical immunologist or relevant specialist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Supply for anticipated emergency treatment of laryngeal/oro-pharyngeal or severe abdominal attacks of acute hereditary angioedema (HAE) for patients with confirmed diagnosis of C1-esterase inhibitor deficiency; and
- 2 The patient has undergone product training and has agreed upon an action plan for self-administration.

Renewal from any relevant practitioner. Approvals valid for 12 months where the treatment remains appropriate and the patient is benefiting from treatment.

Allergy Desensitisation

⇒SA1367 Special Authority for Subsidy

Initial application only from a relevant specialist. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 RAST or skin test positive; and
- 2 Patient has had severe generalised reaction to the sensitising agent.

Renewal only from a relevant specialist. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

REF_VENOM_ALLERGY_TREATMENT — Special Authority see SA1367 above — Retail pharmacy.

BEE VENOM ALLERGY TREATMENT - Special Authority see SAT	367 above – Heta	aii pnarmacy	
Initiation kit - 5 vials freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 1 vial freeze dried venom with diluent	305.00	1 OP	✓ VENOX S29
Maintenance kit - 6 vials 120 mcg freeze dried venom, with			
diluent	285.00	1 OP	✓ Venomil S29
Treatment kit - 1 vial 550 mcg freeze dried venom, 1 diluent			
9 ml, 3 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit - 1 vial 550 mcg freeze dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
WASP VENOM ALLERGY TREATMENT - Special Authority see SA	1367 above – R	etail pharma	су
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze		•	•
dried polister venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Paper wasp venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Paper wasp venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29
Treatment kit (Yellow Jacket venom) - 1 vial 550 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Hymenoptera S29
Treatment kit (Yellow jacket venom) - 1 vial 550 mcg freeze			
dried vespula venom, 1 diluent 9 ml, 1 diluent 1.8 ml	305.00	1 OP	✓ Albey
Treatment kit (Yellow jacket venom) - 6 vials 120 mcg freeze			
dried venom, with diluent	305.00	1 OP	✓ Venomil S29

	Subsidy		Fully Brand or
	(Manufacturer's Price		
	\$	Per	✓ Manufacturer
Autibistaniuss			
Antihistamines			
CETIRIZINE HYDROCHLORIDE			
* Tab 10 mg	1.12	100	✓ Zista
* Oral liq 1 mg per ml	2.84	200 ml	✓ Histaclear
CHLORPHENIRAMINE MALEATE			
* Oral liq 2 mg per 5 ml	0.37	500 ml	✓ Histafen
		300 1111	Tilstaleli
DEXTROCHLORPHENIRAMINE MALEATE	0.00	40	
* Tab 2 mg	4	40	Delevenine
	(8.40)	00	Polaramine
	1.01	20	Polaramine
Y Over lie 0 mg new E ml	(5.99)	100 ml	Polaramine
* Oral liq 2 mg per 5 ml		100 1111	Polaramine
	(10.29)		Polaramine
FEXOFENADINE HYDROCHLORIDE			
* Tab 60 mg		20	
	(8.23)		Telfast
* Tab 120 mg		10	-
	(8.23)		Telfast
	14.22	30	T-161
	(26.44)		Telfast
LORATADINE			
* Tab 10 mg		100	✓ Lorafix
* Oral liq 1 mg per ml	1.43	100 ml	Haylor syrup
PROMETHAZINE HYDROCHLORIDE			
* Tab 10 mg	1.39	50	✓ Allersoothe
* Tab 25 mg	1.58	50	✓ Allersoothe
* Oral liq 1 mg per 1 ml		100 ml	 Allersoothe
* Inj 25 mg per ml, 2 ml ampoule - Up to 5 inj available on a	PSO 17.87	5	✓ Hospira
Inhaled Corticosteroids			
BECLOMETHASONE DIPROPIONATE			
Aerosol inhaler, 50 mcg per dose	14.01 2	00 dose OP	✓ Qvar
Aerosol inhaler, 50 mcg per dose CFC-free		200 dose OF	✓ Beclazone 50
Aerosol inhaler, 100 mcg per dose		200 dose OP	✓ Qvar
Aerosol inhaler, 100 mcg per dose CFC-free		200 dose OP	✓ Beclazone 100
Aerosol inhaler, 250 mcg per dose CFC-free		200 dose OP	✓ Beclazone 250
•		.00 0000 01	5 Decideone 200
BUDESONIDE Powder for inhelation, 100 mag pay does	17.00	100 daga OD	./ Dulminant
Powder for inhalation, 100 mcg per dose	17.00 2	00 dose OP	✓ Pulmicort
Decides for interlating 000 seconds and	40.00	00 -1 05	Turbuhaler
Powder for inhalation, 200 mcg per dose	19.00 2	00 dose OP	✓ Pulmicort
D 1 (' 1 1 1' 400 ' 1	00.00		Turbuhaler
Powder for inhalation, 400 mcg per dose	32.00 2	00 dose OP	✓ Pulmicort
			Turbuhaler

	Subsidy	Duine) O. I	Fully Brand or
	(Manufacturer's \$	Price) Sub Per	sidised Generic Manufacturer
FLUTICASONE			
Aerosol inhaler, 50 mcg per dose	7.19	120 dose OP	✓ Flixotide
Powder for inhalation, 50 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Powder for inhalation, 100 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Aerosol inhaler, 125 mcg per dose		120 dose OP	✓ Flixotide
Aerosol inhaler, 250 mcg per dose		120 dose OP	✓ Flixotide
Powder for inhalation, 250 mcg per dose		60 dose OP	✓ Flixotide Accuhaler
Inhaled Long-acting Beta-adrenoceptor Agoni	sts		
FORMOTEROL FUMARATE			
Powder for inhalation, 12 mcg per dose, and monodose de	vice20.64	60 dose	
T	(35.80)	"	Foradil
Foradil Powder for inhalation, 12 mcg per dose, and monodose	e aevice to be del	istea 1 July 20.	23)
FORMOTEROL FUMARATE DIHYDRATE			
Powder for inhalation 4.5 mcg per dose, breath activated			
(equivalent to eformoterol fumarate 6 mcg metered dos	se)10.32	60 dose OP	
-	(16.90)		Oxis Turbuhaler
NDACATEROL			
Powder for inhalation 150 mcg	61.00	30 dose OP	✓ Onbrez Breezhaler
Powder for inhalation 300 mcg		30 dose OP	✓ Onbrez Breezhaler
SALMETEROL			
Aerosol inhaler CFC-free, 25 mcg per dose	26.25	120 dose OP	✓ Serevent
Powder for inhalation, 50 mgg per dose, breath activated		60 dose OP	✓ Serevent Accuhale
rowder for illifatation, 50 mcg per dose, breath activated	20.20	00 dose or	Serevent Accumates
Inhaled Corticosteroids with Long-Acting Beta	ı-Adrenocept	or Agonists	5
BUDESONIDE WITH EFORMOTEROL			
Powder for inhalation 160 mcg with 4.5 mcg eformoterol			
fumarate per dose (equivalent to 200 mcg budesonide	with		
6 mcg eformoterol fumarate metered dose)		120 dose OP	 DuoResp Spiromax
Powder for inhalation 320 mcg with 9 mcg eformoterol fuma			•
per dose (equivalent to 400 mcg budesonide with 12 m			
eformoterol fumarate metered dose) - No more than 2			
dose per day		120 dose OP	✓ DuoResp Spiromax
		120 dose OP	
Aerosol inhaler 100 mcg with eformoterol fumarate 6 mcg		120 dose OP	✓ Symbicort
Aerosol innaler 100 mcg with eformoterol tumarate 6 mcg Powder for inhalation 100 mcg with eformoterol fumarate 6	mcg33.74		
· · · · · · · · · · · · · · · · · · ·	mcg33.74		Turbuhaler 100/6
Powder for inhalation 100 mcg with eformoterol fumarate 6	· ·	120 dose OP	
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg	21.40		✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6	21.40	120 dose OP	✓ Vannair
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg. Powder for inhalation 200 mcg with eformoterol fumarate 6	21.40	120 dose OP	✓ Vannair ✓ Symbicort
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg. Powder for inhalation 200 mcg with eformoterol fumarate 6 Powder for inhalation 400 mcg with eformoterol fumarate	21.40 mcg33.74	120 dose OP 120 dose OP	✓ Vannair ✓ Symbicort Turbuhaler 200/6
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg. Powder for inhalation 200 mcg with eformoterol fumarate 6	21.40 mcg33.74	120 dose OP	✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg. Powder for inhalation 200 mcg with eformoterol fumarate 6 Powder for inhalation 400 mcg with eformoterol fumarate 12 mcg – No more than 2 dose per day	21.40 mcg33.74	120 dose OP 120 dose OP	✓ Vannair ✓ Symbicort Turbuhaler 200/6
Powder for inhalation 100 mcg with eformoterol fumarate 6 Aerosol inhaler 200 mcg with eformoterol fumarate 6 mcg. Powder for inhalation 200 mcg with eformoterol fumarate 6 Powder for inhalation 400 mcg with eformoterol fumarate	21.40 mcg33.74	120 dose OP 120 dose OP	✓ Vannair ✓ Symbicort Turbuhaler 200/6 ✓ Symbicort

	Subsidy		Fully Brand or
	(Manufacturer's	Price) Subsi	idised Generic Manufacturer
UTICASONE WITH SALMETEROL	Ψ		
Aerosol inhaler 50 mcg with salmeterol 25 mcg	25.79	120 dose OP	✓ Seretide
Aerosol inhaler 125 mcg with salmeterol 25 mcg		120 dose OP	✓ Seretide
Powder for inhalation 100 mcg with salmeterol 50 mcg - No			
more than 2 dose per day		60 dose OP	Seretide Accuhaler
Powder for inhalation 250 mcg with salmeterol 50 mcg – No more than 2 dose per day		60 dose OP	✓ Seretide Accuhaler
more than 2 dose per day	44.00	00 003e O1	• Serelide Acculialer
Beta-Adrenoceptor Agonists			
ALBUTAMOL			
Oral liq 400 mcg per ml		150 ml	✓ <u>Ventolin</u>
Infusion 1 mg per ml, 5 ml		10	✓ Ventolin
Inj 500 mcg per ml, 1 ml - Up to 5 inj available on a PSO	53.00	5	✓ Ventolin
Inhaled Beta-Adrenoceptor Agonists			
ALBUTAMOL			
Aerosol inhaler, 100 mcg per dose CFC free - Up to 1000			
dose available on a PSO	3.80	200 dose OP	✓ Respigen
	(0.00)		✓ SalAir
Nebuliser soln, 1 mg per ml, 2.5 ml ampoule – Up to 30 neb	(6.20)		Ventolin
available on a PSO		20	✓ Asthalin
Nebuliser soln, 2 mg per ml, 2.5 ml ampoule – Up to 30 neb		_0	<u>- 10 11 11 11 11 11 11 11 11 11 11 11 11 </u>
available on a PSO		20	✓ <u>Asthalin</u>
ERBUTALINE SULPHATE			
Powder for inhalation, 200 mcg per dose (equivalent to			
250 mcg metered dose), breath activated	22.20	120 dose OP	Bricanyl Turbuhaler
Anticholinergic Agents			
RATROPIUM BROMIDE			
Aerosol inhaler, 20 mcg per dose CFC-free – Up to 400 dose	e		
available on a PSO		200 dose OP	✓ Atrovent
Nebuliser soln, 250 mcg per ml, 2 ml ampoule - Up to 40 ne	eb		
available on a PSO	11.73	20	✓ Univent
	28.20		✓ Accord S29
nhaled Beta-Adrenoceptor Agonists with Antic	holinergic A	Agents	
ALBUTAMOL WITH IPRATROPIUM BROMIDE			
Aerosol inhaler, 100 mcg with ipratropium bromide, 20 mcg p	er		
dose CFC-free		200 dose OP	✓ Duolin HFA
Nebuliser soln, 2.5 mg with ipratropium bromide 0.5 mg per			
tial O.F. and amenda	44.04		/ Deceller

20

✓ Duolin

vial, 2.5 ml ampoule - Up to 20 neb available on a PSO 11.04

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

Long-Acting Muscarinic Antagonists

GLYCOPYRRONIUM - Subsidy by endorsement

- a) Inhaled glycopyrronium treatment will not be subsidised if patient is also receiving treatment with subsidised tiotropium or umeclidinium
- b) Glycopyrronium powder for inhalation 50 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

TIOTROPIUM BROMIDE - Subsidy by endorsement

- a) Tiotropium treatment will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or umeclidinium.
- b) Tiotropium bromide is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly. Patients who had tiotropium dispensed before 1 October 2018 with a valid Special Authority are deemed endorsed.

UMECLIDINIUM - Subsidy by endorsement

- a) Umeclidinium will not be subsidised if patient is also receiving treatment with subsidised inhaled glycopyrronium or tiotropium bromide.
- b) Umeclidinium powder for inhalation 62.5 mcg per dose is subsidised only for patients who have been diagnosed as having COPD using spirometry if spirometry is possible, and the prescription is endorsed accordingly.

Long-Acting Muscarinic Antagonists with Long-Acting Beta-Adrenoceptor Agonists

Combination long acting muscarinic antagonist and long acting beta-2 agonist will not be subsidised if patient is also receiving treatment with a combination inhaled corticosteroid and long acting beta-2 agonist.

⇒SA1584 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient has been stabilised on a long acting muscarinic antagonist; and
- 2 The prescriber considers that the patient would receive additional benefit from switching to a combination product.

Renewal from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Both:

- 1 Patient is compliant with the medication; and
- 2 Patient has experienced improved COPD symptom control (prescriber determined).

GLYCOPYRRONIUM WITH INDACATEROL - Special Authority see SA1584 above - Retail pharmacy

Powder for Inhalation 50 mcg with indacaterol 110 mcg81.00 30 dose OP

Ultibro Breezhaler

TIOTROPIUM BROMIDE WITH OLODATEROL - Special Authority see SA1584 above - Retail pharmacy

UMECLIDINIUM WITH VILANTEROL - Special Authority see SA1584 above - Retail pharmacy

Antifibrotics

NINTEDANIB - Special Authority see SA2012 on the next page - Retail pharmacy

Note: Nintedanib not subsidised in combination with subsidised pirfenidone.

 Cap 100 mg
 2,554.00
 60 OP
 ✓ Ofev

 Cap 150 mg
 3,870.00
 60 OP
 ✓ Ofev

✓ fully subsidised S29 Unapproved n
Principal Supply Sole Subsidised St

Subsidy		Fully	Brand or	
(Manufacturer's Price)	:	Subsidised	Generic	
\$	Per	✓	Manufacturer	

⇒SA2012 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Nintedanib is to be discontinued at disease progression (See Note); and
- 4 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with pirfenidone; or
 - 5.2 Patient has previously received pirfenidone, but discontinued pirfenidone within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received pirfenidone, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with pirfenidone).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Nintedanib is not to be used in combination with subsidised pirfenidone; and
- 3 Nintedanib is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

PIRFENIDONE – Retail pharmacy-Specialist – Special Authority see SA2013 below

Note: Pirfenidone is not subsidised in combination with subsidised nintedanib

Troto: I monidono lo not cabolaleca in combination with cabol	alood milloddino.		
Tab 801 mg	3,645.00	90	Esbriet
Tab 267 mg	1,215.00	90	Esbriet

⇒SA2013 Special Authority for Subsidy

Initial application — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with idiopathic pulmonary fibrosis by a multidisciplinary team including a radiologist; and
- 2 Forced vital capacity is between 50% and 90% predicted; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note); and
- 4 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 5 Any of the following:
 - 5.1 The patient has not previously received treatment with nintedanib; or
 - 5.2 Patient has previously received nintedanib, but discontinued nintedanib within 12 weeks due to intolerance; or
 - 5.3 Patient has previously received nintedanib, but the patient's disease has not progressed (disease progression defined as 10% or more decline in predicted FVC within any 12 month period since starting treatment with nintedanib).

Renewal — (idiopathic pulmonary fibrosis) only from a respiratory specialist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Treatment remains clinically appropriate and patient is benefitting from and tolerating treatment; and
- 2 Pirfenidone is not to be used in combination with subsidised nintedanib; and
- 3 Pirfenidone is to be discontinued at disease progression (See Note).

Note: disease progression is defined as a decline in percent predicted FVC of 10% or more within any 12 month period.

	Subsidy (Manufacturer's Price) \$	Sul Per	Fully bsidised	Brand or Generic Manufacturer	
Leukotriene Receptor Antagonists					
MONTELUKAST					

Louitotiiciic	ricocptor	Ailtag	OHIOU

*	Tab 4 mg	28	✓ Montelukast Mylan
	Montelukast Mylan to be Principal Supply on 1 December 2022		·
*	Tab 5 mg	28	✓ Montelukast Mylan
	Montelukast Mylan to be Principal Supply on 1 December 2022		·
*	Tab 10 mg	28	✓ Montelukast Mylan
	Montelukast Mylan to be Principal Supply on 1 December 2022		•

Methylxanthines

AMINOPHYLLINE

* Inj 25 mg per ml, 10 ml ampoule - Up to 5 inj available	on a		
PSO	180.00	5	DBL Aminophylline
THEOPHYLLINE			
* Tab long-acting 250 mg	23.02	100	✓ Nuelin-SR
* Oral lig 80 mg per 15 ml	16.60	500 ml	✓ Nuelin

Mucolytics

DORNASE ALFA - Special Authority see S	A1978 below – Retail pharmacy		
Nebuliser soln, 2.5 mg per 2.5 ml ampo	ule250.00	6	✓ Pulmozyme

⇒SA1978 Special Authority for Subsidy

Initial application — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a confirmed diagnosis of cystic fibrosis; and
- 2 Patient has previously undergone a trial with, or is currently being treated with, hypertonic saline; and
- 3 Any of the following:
 - 3.1 Patient has required one or more hospital inpatient respiratory admissions in the previous 12 month period; or
 - 3.2 Patient has had 3 exacerbations due to CF, requiring oral or intravenous (IV) antibiotics in the previous 12 month
 - 3.3 Patient has had 1 exacerbation due to CF, requiring oral or IV antibiotics in the previous 12 month period and a Brasfield score of < 22/25: or
 - 3.4 Patient has a diagnosis of allergic bronchopulmonary aspergillosis (ABPA).

Renewal — (cystic fibrosis) only from a respiratory physician or paediatrician. Approvals valid without further renewal unless notified where the treatment remains appropriate and the patient continues to benefit from treatment.

IVACAFTOR - PCT only - Specialist - Special Authority see SA2017 below

		olalist opeolal halfforty see on 2017 below	V/10/11 TOTE TOTOTHY
✓ Kalydeco	56	29,386.00	Tab 150 mg
✓ Kalydeco	56	t29,386.00	Oral granules 50 mg, s
✓ Kalydeco	56	t29,386.00	Oral granules 75 mg, s

⇒SA2017 Special Authority for Subsidy

Initial application only from a respiratory specialist or paediatrician. Approvals valid without further renewal unless notified for applications meeting the following criteria:

All of the following:

- 1 Patient has been diagnosed with cystic fibrosis; and
- 2 Fither:

	RESPIRA	TORY SYSTE	EM AI	ND ALLERGIES
	Subsidy (Manufacturer's \$	Price) Subs Per	Fully idised	Brand or Generic Manufacturer
continued				
Patient must have G551D mutation in the cystic least 1 allele; or Patient must have other gating (class III) mutation.	on (G1244E, G134			
and S549R) in the CFTR gene on at least 1 allel 3 Patients must have a sweat chloride value of at least 60	•	itative pilocarpin	e ionto	phoresis or by Macrodu
sweat collection system; and 4 Treatment with ivacaftor must be given concomitantly w 5 Patient must not have an acute upper or lower respirato (including antibiotics) for pulmonary disease in the last 6 The dose of ivacaftor will not exceed one tablet or one of 7 Applicant has experience and expertise in the manager	ory infection, pulm 4 weeks prior to c sachet twice daily:	onary exacerbat ommencing treat and	ion, or	changes in therapy
SODIUM CHLORIDE Not funded for use as a nasal drop. Soln 7%	24.50	90 ml OP	√ B	iomed
Nasal Preparations				
Allergy Prophylactics				
BUDESONIDE Metered aqueous nasal spray, 50 mcg per dose Metered aqueous nasal spray, 100 mcg per dose		200 dose OP 200 dose OP	_	teroClear teroClear
FLUTICASONE PROPIONATE Metered aqueous nasal spray, 50 mcg per dose	1.98	120 dose OP	√ <u>F</u>	lixonase Hayfever & Allergy
PRATROPIUM BROMIDE				
Aqueous nasal spray, 0.03%	5.23	15 ml OP	√ <u>U</u>	<u>Inivent</u>
Respiratory Devices				
MASK FOR SPACER DEVICE a) Up to 50 dev available on a PSO b) Only on a PSO				
c) Only for children aged six years and under Small	2.20	1	√ e	-chamber Mask
PEAK FLOW METER a) Up to 25 dev available on a PSO b) Only on a PSO				
Low range	9.54	1	✓ N	lini-Wright AFS Low Range
Normal range	9.54	1	✓ N	lini-Wright Standard
SPACER DEVICE				
a) Up to 50 dev available on a PSO				
b) Only on a PSO 220 ml (single patient)	2.95	1	√ e	-chamber Turbo
510 ml (single patient)		1		-chamber I a

✓ e-chamber La

Grande

✓ Volumatic

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic
\$ Per Manufacturer

Respiratory Stimulants

CAFFEINE CITRATE

Oral liq 20 mg per ml (10 mg base per ml)......15.10 25 ml OP **✔ Biomed**

	Subsidy		Fully Brand or
	(Manufacturer's Pri	ce) Subs	idised Generic
	\$	Per	 Manufacturer
Ear Preparations			
ELLINATIA CONTE DIVALATE			
FLUMETASONE PIVALATE			
Ear drops 0.02% with clioquinol 1%	4.46	7.5 ml OP	✓ Locacorten-Viaform
			ED's
			✓ Locorten-Vioform
TRIAMCINOLONE ACETONIDE WITH GRAMICIDIN, NEOMYCI	N AND NYSTATII	N	
Ear drops 1 mg with nystatin 100,000 u, neomycin sulphate			
2.5 mg and gramicidin 250 mcg per g	5 16	7.5 ml OP	✓ Kenacomb
2.0 mg and gramolam 200 mog por g		7.0 1111 01	- Itoliacomb
Ear/Eye Preparations			
Edit Lyc i Toparations			
DEXAMETHASONE WITH FRAMYCETIN AND GRAMICIDIN			
Ear/Eye drops 500 mcg with framycetin sulphate 5 mg and			
gramicidin 50 mcg per ml	4 50	8 ml OP	
gramoum oo mog por mi	(9.27)	0 1111 01	Sofradex
EDAM/OETIN OUR BLIATE	(0.27)		Gondaex
FRAMYCETIN SULPHATE	4.40	0l OD	
Ear/Eye drops 0.5%		8 ml OP	0 - (
	(8.65)		Soframycin
Eve Drenerations			
Eye Preparations			
Eye preparations are only funded for use in the eye, unless explic	ritly stated otherwi	isa	
	nily stated offici w	130.	
Anti-Infective Preparations			
ACICLOVID			
ACICLOVIR * Eye oint 3%	44.00	4.5 = OD	/ Vim.poc
•	14.00	4.5 g OP	✓ <u>ViruPOS</u>
CHLORAMPHENICOL			
Eye oint 1%	1.09	5 g OP	Devatis
Devatis to be Principal Supply on 1 December 2022			
Eye drops 0.5%		10 ml OP	✓ Chlorafast
Funded for use in the ear*. Indications marked with * are	e unapproved indi	cations.	
CIPROFLOXACIN			
Eye drops 0.3% - Subsidy by endorsement	9.73	5 ml OP	✓ Ciprofloxacin Teva
When prescribed for the treatment of bacterial keratitis o		conjunctivitis	resistant to chloramphenicol; or
for the second line treatment of chronic suppurative otitis			
Note: Indication marked with a * is an unapproved indication		•	,
GENTAMICIN SULPHATE			
Eye drops 0.3%	11.40	5 ml OP	✓ Genoptic
(Genoptic Eye drops 0.3% to be delisted 1 August 2023)	11.40	3 1111 01	• delloptic
PROPAMIDINE ISETHIONATE			
* Eye drops 0.1%		10 ml OP	5 .
	(14.55)		Brolene
SODIUM FUSIDATE [FUSIDIC ACID]			
Eye drops 1%	5.29	5 g OP	✓ Fucithalmic
TOBRAMYCIN		-	
Eye oint 0.3%	10.45	3.5 g OP	✓ Tobrex
Eye drops 0.3%		5.5 g Oi 5 ml OP	✓ Tobrex
_, o alopo 0.0 /0	11.70	3 1111 01	- 100104



Subsidy	F	ully	Brand or	
(Manufacturer's Price)	Subsidis	sed	Generic	
\$	Per	•	Manufacturer	
				ě

Corticosteroids and Other Anti-Inflammatory Preparations

DEXAMETHASONE			
* Eye oint 0.1%	5.86	3.5 g OP	✓ Maxidex
* Eye drops 0.1%	4.50	5 ml OP	Maxidex
Ocular implant 700 mcg - Special Authority see SA1680 below			
- Retail pharmacy1	.444.50	1	Ozurdex

⇒SA1680 Special Authority for Subsidy

Initial application — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema with pseudophakic lens: and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Fither
 - 3.1 Patient's disease has progressed despite 3 injections with bevacizumab; or
 - 3.2 Patient is unsuitable or contraindicated to treatment with anti-VEGF agents; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

Both:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Initial application — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient has diabetic macular oedema; and
- 2 Patient has reduced visual acuity of between 6/9 6/48 with functional awareness of reduction in vision; and
- 3 Patient is of child bearing potential and has not yet completed a family; and
- 4 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

Renewal — (Women of child bearing age with diabetic macular oedema) only from an ophthalmologist. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient's vision is stable or has improved (prescriber determined); and
- 2 Patient is of child bearing potential and has not yet completed a family; and
- 3 Dexamethasone implants are to be administered not more frequently than once every 4 months into each eye, and up to a maximum of 3 implants per eye per year.

DEXAMETHASONE WITH NEOMYCIN SULPHATE AND POLYMYXIN B SULPHATE

* Eye oint 0.1% with neomycin sulphate 0.35% and polymyxin b		
sulphate 6,000 u per g5.39	3.5 g OP	✓ Maxitrol
* Eye drops 0.1% with neomycin sulphate 0.35% and polymyxin		
b sulphate 6,000 u per ml4.50	5 ml OP	✓ Maxitrol
DICLOFENAC SODIUM		
Eye drops 0.1%	5 ml OP	✓ Voltaren Ophtha
FLUOROMETHOLONE		
	5l OD	/ F141
* Eye drops 0.1%3.09	5 ml OP	✓ FML
5.20		✓ Flucon

Betoptic S

5 ml OP

				
	Subsidy		Fully	Brand or
	(Manufacturer's F	,	sidised	Generic
	\$	Per		Manufacturer
LEVOCABASTINE				
Eye drops 0.5 mg per ml	8.71	4 ml OP		
, , , , , , , , , , , , , , , , , , , ,	(10.34)		L	ivostin
LODOXAMIDE				
Eye drops 0.1%	8.71	10 ml OP	✓ L	.omide
PREDNISOLONE ACETATE				
Eye drops 1%	6.92	10 ml OP	√ F	Prednisolone-AFT
•	7.00	5 ml OP	√ F	red Forte
PREDNISOLONE SODIUM PHOSPHATE - Special Author	ity see SA1715 below	v – Retail phar	macy	
Eye drops 0.5%, single dose (preservative free)	38.50	20 dose	٧	Minims
				Prednisolone
⇒SA1715 Special Authority for Subsidy				
- Control opposition of outside				

Initial application only from an ophthalmologist or optometrist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Patient has severe inflammation; and
- 2 Patient has a confirmed allergic reaction to preservative in eye drops.

Renewal from any relevant practitioner. Approvals valid for 6 months where the treatment remains appropriate and the patient is benefiting from treatment.

SODIUM CROMOGLICATE	
---------------------	--

Eye drops 2%	1.79	5 ml OP	Rexacrom
	2.62	10 ml OP	Allerfix
(Daysaram Five drang 00/ to be delicted 1 March 0000)			

(Rexacrom Eye drops 2% to be delisted 1 March 2023)

ΔΩΕΤΔΖΟΙ ΔΜΙΩΕ

木	Eye drops 0.5%	5 MI OP	•	веторис
TIN	NOLOL			
*	Eye drops 0.25%	5 ml OP	1	Arrow-Timolol
*	Eye drops 0.5%2.04	5 ml OP	1	Arrow-Timolol
*	Eye drops 0.5%, gel forming	2.5 ml OP	1	Timoptol XE

Glaucoma Preparations - Carbonic Anhydrase Inhibitors

* Tab 250 mg	'.03	100	✓ Diamox
BRINZOLAMIDE	7.00	E ml OD	. Amont
* Eye drops 1%	.30	5 IIII OP	<u>Azopi</u>
* Eye drops 2%).77	5 ml OP	

DORZOLAMIDE WITH TIMOLOL (17.44) Trusopt

Glaucoma Preparations - Prostaglandin Analogues

DΙ	M٨	т,	\cap	D		\cap	C1	ľ
DI	IVI <i>F</i>	١I١	U	ГΙ	יח	U	O	

[▲]Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

	Subsidy (Manufacturer's Pr		
	\$	Per	✓ Manufacturer
LATANOPROST			
* Eye drops 0.005%	1.82	2.5 ml OP	✓ Teva
TRAVOPROST			
* Eye drops 0.004%	9.75	2.5 ml OP	✓ Travatan
7			
Glaucoma Preparations - Other			
·			
BRIMONIDINE TARTRATE			
* Eye drops 0.2%	4.29	5 ml OP	✓ <u>Arrow-Brimonidine</u>
BRIMONIDINE TARTRATE WITH TIMOLOL MALEATE			
* Eye drops 0.2% with timolol maleate 0.5%	18.50	5 ml OP	✓ Combigan
LATANOPROST WITH TIMOLOL			_
Eye drops 0.005% with timolol 0.5%	2.49	2.5 ml OP	✓ Arrow - Lattim
PILOCARPINE HYDROCHLORIDE			
	4.06	15 ml OP	✓ Isopto Carpine
7		15 ml OP	
7		15 ml OP	✓ Isopto Carpine
* Eye drops 4%		15 1111 OF	✓ Isopto Carpine
•	a c .		
* Eye drops 2% single dose – Special Authority see SA0895	04.05	00 dasa	/ Minima Dilanamina
below – Retail pharmacy	31.95	20 dose	Minims Pilocarpine

⇒SA0895 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 2 years for applications meeting the following criteria: Either:

- 1 Patient has to use an unpreserved solution due to an allergy to the preservative; or
- 2 Patient wears soft contact lenses.

Note: Minims for a general practice are considered to be "tools of trade" and are not approved as special authority items. **Renewal** from any relevant practitioner. Approvals valid for 2 years where the treatment remains appropriate and the patient is benefiting from treatment.

Mydriatics and Cycloplegics		
* Eye drops 1%	15 ml OP	✓ Atropt
* Eye drops 1%	15 ml OP	✓ Cyclogyl
* Eye drops 0.5%	15 ml OP 15 ml OP	MydriacylMydriacyl
Preparations for Tear Deficiency		
For acetylcysteine eye drops refer Standard Formulae, page 248 HYPROMELLOSE		
* Eye drops 0.5%	15 ml OP	✓ Methopt
* Eye drops 0.3% with dextran 0.1%2.30	15 ml OP	✓ Poly-Tears

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	1	Manufacturer

Preservative Free Ocular Lubricants

⇒SA2134 Special Authority for Subsidy

Initial application from any relevant practitioner. Approvals valid for 12 months for applications meeting the following criteria: Both:

- 1 Confirmed diagnosis by slit lamp or Schirmer test of severe secretory dry eye; and
- 2 Either:
 - 2.1 Patient is using eye drops more than four times daily on a regular basis; or
 - 2.2 Patient has had a confirmed allergic reaction to preservative in eye drop.

Renewal from any relevant practitioner. Approvals valid for 24 months where the patient continues to require lubricating eye drops and has benefited from treatment.

CARBOMER – Special Authority see SA2134 above – Retail pha	rmacy		
Ophthalmic gel 0.3%, 0.5 g	8.25	30	✓ Poly-Gel
MACROGOL 400 AND PROPYLENE GLYCOL - Special Authori	ty see SA2134 al	bove – Reta	il pharmacy
Eye drops 0.4% and propylene glycol 0.3%, 0.4 ml	4.30	24	 Systane Unit Dose
Eye drops 0.4% and propylene glycol 0.3%, 0.8 ml	10.78	30	✓ Systane Unit Dose
(Systane Unit Dose Eye drops 0.4% and propylene glycol 0.3%, 0	0.4 ml to be delist	ed 1 June 20	023)
SODIUM HYALURONATE [HYALURONIC ACID] - Special Author	ority see SA2134	above - Re	tail pharmacy
Eye drops 1 mg per ml	13.85	10 ml OP	✓ Hylo-Fresh
Hylo-Fresh has a 6 month expiry after opening. The Pha	rmacy Procedure	es Manual re	striction allowing one bottle per
month is not relevant and therefore only the prescribed de	osage to the near	rest OP may	be claimed.

Other Eye Preparations

NAPHAZOLINE HYDROCHLORIDE * Eye drops 0.1%	15 ml OP	✓ Naphcon Forte
OLOPATADINE Eye drops 0.1%	5 ml OP	✓ Olopatadine Teva
PARAFFIN LIQUID WITH WOOL FAT * Eye oint 3% with wool fat 3%	3.5 g OP	✓ Poly-Visc
RETINOL PALMITATE Eye oint 138 mcg per g3.80	5 g OP	✓ VitA-POS



Subsidy		Fully	Brand or	_
(Manufacturer's Price)		Subsidised	Generic	
\$	Per	/	Manufacturer	

Various

PHARMACY SERVICES

May only be claimed once per patient.

1 fee ✓ BSF Amgevita ✓ BSF Aspen

a) The Pharmacode for BSF Aspen is 2641356 - see also page 137

b) The Pharmacode for BSF Amgevita is 2645165 - see also page 174

(BSF Amgevita Brand switch fee to be delisted 1 January 2023)

(BSF Aspen Brand switch fee to be delisted 1 November 2022)

Agents Used in the Treatment of Poisonings

Antidotes

ACETYLCYSTEINE

Inj 200 mg per ml, 10 ml ampoule52.88

58.76

✓ Martindale Pharma

✓ DBL Acetvlcvsteine

✓ Martindale Pharma S29 S29

Martindale Pharma to be Principal Supply on 1 December 2022

(DBL Acetylcysteine Inj 200 mg per ml, 10 ml ampoule to be delisted 1 December 2022)

(Martindale Pharma S29 S29 Inj 200 mg per ml, 10 ml ampoule to be delisted 1 December 2022)

NALOXONE HYDROCHLORIDE

- a) Up to 5 inj available on a PSO
- b) Only on a PSO

* Inj 400 mcg per ml, 1 ml ampoule22.60

✓ DBL Naloxone Hydrochloride

10

✓ Hameln

(DBL Naloxone Hydrochloride Inj 400 mcg per ml, 1 ml ampoule to be delisted 1 February 2023)

Removal and Elimination

CHARCOAL

* Oral liq 50 g per 250 ml43.50 250 ml OP ✓ Carbosorb-X

a) Up to 250 ml available on a PSO

b) Only on a PSO

DEFERASIROX - Special Authority see SA1492 below - Retail pharmacy

Wastage claimable

28 Exiade 28 ✓ Exiade 28 Exiade

⇒SA1492 Special Authority for Subsidy

Initial application only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: All of the following:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; and
- 2 Deferasirox is to be given at a daily dose not exceeding 40 mg/kg/day; and

Subsidy)	Fully	Brand or
(Manufacturer's Price		Subsidised	Generic
\$	Per	•	Manufacturer

continued...

- 3 Any of the following:
 - 3.1 Treatment with maximum tolerated doses of deferiprone monotherapy or deferiprone and desferrioxamine combination therapy have proven ineffective as measured by serum ferritin levels. liver or cardiac MRI T2*; or
 - 3.2 Treatment with deferiprone has resulted in severe persistent vomiting or diarrhoea; or
 - 3.3 Treatment with deferiprone has resulted in arthritis: or
 - 3.4 Treatment with deferiprone is contraindicated due to a history of agranulocytosis (defined as an absolute neutrophil count (ANC) of < 0.5 cells per μL) or recurrent episodes (greater than 2 episodes) of moderate neutropenia (ANC 0.5 1.0 cells per μL).</p>

Renewal only from a haematologist. Approvals valid for 2 years for applications meeting the following criteria: Fither:

- 1 For the first renewal following 2 years of therapy, the treatment has been tolerated and has resulted in clinical improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels; or
- 2 For subsequent renewals, the treatment has been tolerated and has resulted in clinical stability or continued improvement in all three parameters namely serum ferritin, cardiac MRI T2* and liver MRI T2* levels.

DEFERIPRONE - Special Authority see SA1480 below - Re	tail pharmacy		
Tab 500 mg	533.17	100	✓ Ferriprox
Oral liq 100 mg per 1 ml	266.59	250 ml OP	✓ Ferriprox

⇒SA1480 Special Authority for Subsidy

DECEEDBIOVAMINE MECH ATE

Initial application only from a haematologist. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Fither:

- 1 The patient has been diagnosed with chronic iron overload due to congenital inherited anaemia; or
- 2 The patient has been diagnosed with chronic iron overload due to acquired red cell aplasia.

* Inj 500 mg vial	151.31	10	✓ DBL Desferrioxamine Mesylate for Inj BP ✓ Deferoxamine Pfizer S29 \$29
SODIUM CALCIUM EDETATE * Inj 200 mg per ml, 5 ml	53.31	6	
,	(156.71)		Calcium Disodium Versenate

Standard Formulae

Standard Formulae			
ACETYLCYSTEINE EYE DROPS Acetylcysteine inj 200 mg per ml, 10 ml Suitable eye drop base	qs qs	PHENOBARBITONE ORAL LIQUID Phenobarbitone Sodium Glycerol BP Water	1 g 70 ml to 100 ml
CODEINE LINCTUS (3 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	60 mg 40 ml qs to 100 ml	PHENOBARBITONE SODIUM PAEDIATRIC ORAL mg per ml) Phenobarbitone Sodium Glycerol BP Water	400 mg 4 ml to 40 ml
CODEINE LINCTUS (15 mg per 5 ml) Codeine phosphate Glycerol Preservative Water	300 mg 40 ml qs to 100 ml	PILOCARPINE ORAL LIQUID Pilocarpine 4% eye drops Preservative Water	qs qs to 500 ml
FOLINIC MOUTHWASH Calcium folinate 15 mg tab Preservative Water (Preservative should be used if quantity supplied is f than 5 days. Maximum 500 ml per prescription.)	1 tab qs to 500 ml or more	(Preservative should be used if quantity supplied is than 5 days.) SALIVA SUBSTITUTE FORMULA Methylcellulose Preservative Water	5 g qs to 500 ml
METHADONE MIXTURE Methadone powder Glycerol Water	qs qs to 100 ml	(Preservative should be used if quantity supplied is than 5 days. Maximum 500 ml per prescription.) SODIUM CHLORIDE ORAL LIQUID Sodium chloride inj 23.4%, 20 ml Water	qs qs
METHYL HYDROXYBENZOATE 10% SOLUTION Methyl hydroxybenzoate Propylene glycol (Use 1 ml of the 10% solution per 100 ml of oral liqui	10 g to 100 ml id mixture)	(Only funded if prescribed for treatment of hyponatra VANCOMYCIN ORAL SOLUTION (50 mg per ml) Vancomycin 500 mg injection	
OMEPRAZOLE SUSPENSION Omeprazole capsules or powder Sodium bicarbonate powder BP Water	qs 8.4 g to 100 ml	Glycerol BP Water (Only funded if prescribed for treatment of Clostridiu following metronidazole failure)	to 100 ml

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Manufacturer

Per **Extemporaneously Compounded Preparations and Galenicals**

CODEINE PHOSPHATE - Safety medicine; prescriber may determine dispensing frequency

(90.09)

Douglas

✓ AFT

1 a

Only in extemporaneously compounded codeine linctus.

COLLODION FLEXIBLE

Note: This product is no longer being manufactured by the supplier and will be delisted from the Schedule at a date to be determined.

100 ml ✓ PSM

COMPOUND HYDROXYBENZOATE - Only in combination

Only in extemporaneously compounded oral mixtures.

100 ml Midwest

GLYCERIN WITH SODIUM SACCHARIN - Only in combination

Only in combination with Ora-Plus.

473 ml ✓ Ora-Sweet SF

GLYCERIN WITH SUCROSE - Only in combination

Only in combination with Ora-Plus.

Suspension.......30.95 473 ml Ora-Sweet

GI YCFROI

500 ml ✓ healthE Glycerol BP Only in extemporaneously compounded oral liquid preparations.

METHADONE HYDROCHLORIDE

- a) Only on a controlled drug form
- b) No patient co-payment payable
- c) Safety medicine; prescriber may determine dispensing frequency

d) Extemporaneously compounded methadone will only be reimbursed at the rate of the cheapest form available (methadone powder, not methadone tablets).

METHYL HYDROXYBENZOATE Powder	8.98	25 g	✓ Midwest
METHYLCELLULOSE			
Powder	36.95	100 g	✓ MidWest
Suspension - Only in combination	30.95	473 ml	Ora-Plus
METHYLCELLULOSE WITH GLYCERIN AND SODIUM SACCHAR	IN - Only in o	combination	

473 ml Ora-Blend SF

METHYLCELLULOSE WITH GLYCERIN AND SUCROSE - Only in combination 473 ml

Ora-Blend

PHENOBARRITONE SODIUM

Powder - Only in combination......52.50 ✓ MidWest 10 a 325.00 ✓ MidWest 100 g

Only in children up to 12 years

PROPYLENE GLYCOL

Only in extemporaneously compounded methyl hydroxybenzoate 10% solution.

500 ml Midwest Liq......11.25

SODIUM BICARBONATE

Midwest 500 a Only in extemporaneously compounded omeprazole and lansoprazole suspension.

▲Three months supply may be dispensed at one time if endorsed "certified exemption" by the prescriber or pharmacist.

EXTEMPORANEOUSLY COMPOUNDED PREPARATIONS AND GALENICALS

	Subsidy (Manufacturer's Price \$	e) Su Per	Fully bsidised	Brand or Generic Manufacturer	
SYRUP (PHARMACEUTICAL GRADE) – Only in combination Only in extemporaneously compounded oral liquid preparatio Liq		500 ml	✓ M	lidwest	
WATER Tap - Only in combination	0.00	1 ml	✓ Ta	ap water	

Subsidy (Manufacturer's Price) \$

Subsidised Per 🗸

Fully

Brand or Generic Manufacturer

Nutrient Modules

Carbohydrate

⇒SA1930 Special Authority for Subsidy

Initial application — (Cystic fibrosis or kidney disease) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Fither:

- 1 cystic fibrosis; or
- 2 chronic kidney disease.

Initial application — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 cancer in children: or
- 2 cancers affecting alimentary tract where there are malabsorption problems in patients over the age of 20 years; or
- 3 faltering growth in an infant/child; or
- 4 bronchopulmonary dysplasia; or
- 5 premature and post premature infant; or
- 6 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified where the patient has inborn errors of metabolism. Renewal — (Cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis or renal failure) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

CARBOHYDRATE SUPPLEMENT - Special Authority see SA1930 above - Hospital pharmacy [HP3]

Carbohydrate And Fat

⇒SA1376 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

continued...

✓ fully subsidised 251



	Subsidy (Manufacturer's Price)	C.	Fully ubsidised	Brand or Generic
((Manufacturer's Frice)		ubsidised	
	\$	Per	✓	Manufacturer

continued...

- 1 Infant or child aged four years or under; and
- 2 cystic fibrosis.

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 infant or child aged four years or under; and
- 2 Any of the following:
 - 2.1 cancer in children; or
 - 2.2 faltering growth; or
 - 2.3 bronchopulmonary dysplasia; or
 - 2.4 premature and post premature infants.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Fat

⇒SA1523 Special Authority for Subsidy

Initial application — (Inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has inborn errors of metabolism.

Initial application — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria: Any of the following:

- 1 faltering growth in an infant/child; or
- 2 bronchopulmonary dysplasia; or
- 3 fat malabsorption; or
- 4 lymphangiectasia; or
- 5 short bowel syndrome: or
- 6 infants with necrotising enterocolitis; or
- 7 biliary atresia: or
- 8 for use in a ketogenic diet; or
- 9 chyle leak; or

Subsidy (Manufacturer's Price)	Sı	Fully ubsidised	Brand or Generic
 \$	Per	1	Manufacturer

- 10 ascites: or
- 11 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal — (Inborn errors of metabolism)** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than inborn errors of metabolism) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT SUPPLEMENT - Special Authority see SA1523 on the previous page - Hospital pharmacy [HP3]

Emulsion (neutral)	12.30	200 ml OP	✓ Calogen
	30.75	500 ml OP	✓ Calogen
Emulsion (strawberry)	12.30	200 ml OP	✓ Calogen
Oil	30.00	500 ml OP	✓ MCT oil (Nutricia)
Oil, 250 ml	114.92	4 OP	✓ Liquigen

Protein

⇒SA1524 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 protein losing enteropathy; or
- 2 high protein needs; or
- 3 for use as a component in a modular formula made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule or breast milk.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

	rmacy [HP3]	PROTEIN SUPPLEMENT — Special Authority see SA1524 above — Hospital pha
✓ Protifar	225 g OP	Powder7.90
✓ Resource	227 g OP	8.95
Beneprotein	•	

Subsidy (Manufacturer's Price) \$ Per

Fully Subsidised Brand or Generic Manufacturer

Oral and Enteral Feeds

Diabetic Products

⇒SA1095 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is a type I or and II diabetic who is suffering weight loss and malnutrition that requires nutritional support. Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

DIABETIC ENTERAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid3.75	500 ml OP	✓ Glucerna Select
7.50	1,000 ml OP	Diason RTH
		✓ Nutrison Advanced
		Diason

(Diason RTH Liquid to be delisted 1 December 2022)

DIABETIC ORAL FEED 1KCAL/ML - Special Authority see SA1095 above - Hospital pharmacy [HP3]

Liquid (strawberry)1.50	200 ml OP	Diasip
Liquid (vanilla)1.50	200 ml OP	✓ Diasip
2 10		✓ Nutren Diahe

Fat Modified Products

⇒SA1525 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Patient has metabolic disorders of fat metabolism; or
- 2 Patient has a chyle leak; or
- 3 Modified as a modular feed, made from at least one nutrient module and at least one further product listed in Section D of the Pharmaceutical Schedule, for adults.

Note: Patients are required to meet any Special Authority criteria associated with all of the products used in the modular formula. **Renewal** only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

FAT MODIFIED FEED - Special Authority see SA1525 above - Hospital pharmacy [HP3]

Subsidy (Manufacturer's Price) \$ Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children Awaiting Liver Transplant

⇒SA1098 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) who requires a liver transplant.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1098 above - Hospital pharmacy [HP3]

Paediatric Products For Children With Chronic Renal Failure

⇒SA1099 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient is a child (up to 18 years) with acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL FEED 1KCAL/ML - Special Authority see SA1099 above - Hospital pharmacy [HP3]

Paediatric Products

⇒SA1379 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child is aged one to ten years; and
- 2 Any of the following:
 - 2.1 the child is being fed via a tube or a tube is to be inserted for the purposes of feeding; or
 - 2.2 any condition causing malabsorption; or
 - 2.3 faltering growth in an infant/child; or
 - 2.4 increased nutritional requirements; or
 - 2.5 the child is being transitioned from TPN or tube feeding to oral feeding.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	1	Manufacturer	

applications meeting the following criteria:

Roth:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

practitioner and date contacted.			
PAEDIATRIC ENTERAL FEED 1.5KCAL/ML – Special Authority se Liquid		the previous pa 500 ml OP	
PAEDIATRIC ENTERAL FEED 1KCAL/ML - Special Authority see Liquid		e previous page 500 ml OP	 Hospital pharmacy [HP3] Nutrini RTH Pediasure RTH
PAEDIATRIC ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Specipharmacy [HP3]	al Authority se	e SA1379 on th	e previous page – Hospital
Liquid	6.00	500 ml OP	✓ Nutrini Energy Multi Fibre
PAEDIATRIC ORAL FEED 1.5KCAL/ML - Special Authority see Soliquid (strawberry)	1.60	previous page – 200 ml OP 200 ml OP 500 ml OP	Hospital pharmacy [HP3] Fortini Fortini Pediasure Plus
PAEDIATRIC ORAL FEED 1KCAL/ML - Special Authority see SA1 Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.07 1.07	evious page – H 200 ml OP 200 ml OP 200 ml OP 250 ml OP	lospital pharmacy [HP3] ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure ✓ Pediasure
PAEDIATRIC ORAL FEED WITH FIBRE 1.5KCAL/ML - Special Aupharmacy [HP3]	uthority see S/	A1379 on the pre	evious page – Hospital
Liquid (unflavoured) Liquid (chocolate) Liquid (strawberry) Liquid (vanilla)	1.60 1.60	200 ml OP 200 ml OP 200 ml OP 200 ml OP	 ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre ✓ Fortini Multi Fibre
PEPTIDE-BASED ORAL FEED – Special Authority see SA1379 or Powder		page – Hospital 400 g OP	pharmacy [HP3] ✓ Peptamen Junior

Renal Products

⇒SA1101 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years where the patient has acute or chronic kidney disease.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

RENAL ENTERAL FEED 1.8 KCAL/ML	 Special Authority see SA1101 above - 	 Hospital phari 	macy [HP3]
Liquid	6.08	500 ml OP	✓ Nepro HP RTH

	Subsidy (Manufacturer's Pric		Fully dised	Brand or Generic Manufacturer
RENAL ORAL FEED 1.8 KCAL/ML – Special Authority see SA1 Liquid		s page – Hosp 220 ml OP	✓ N	narmacy [HP3] lepro HP (strawberry) lepro HP (vanilla)
RENAL ORAL FEED 2 KCAL/ML - Special Authority see SA110 Liquid, 200 ml bottle	11.52 (13.24)	page – Hospit 4 OP	N	lovaSource Renal
Liquid (apricot) 125 ml Liquid (caramel) 125 ml		4 OP 4 OP	-	Renilon 7.5 Renilon 7.5

Specialised And Elemental Products

⇒SA1377 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 malabsorption: or
- 2 short bowel syndrome; or
- 3 enterocutaneous fistulas: or
- 4 eosinophilic oesophagitis; or
- 5 inflammatory bowel disease: or
- 6 patients with multiple food allergies requiring enteral feeding.

Notes: Each of these products is highly specialised and would be prescribed only by an expert for a specific disorder. The alternative is hospitalisation.

Elemental 028 Extra is more expensive than other products listed in this section and should only be used where the alternatives have been tried first and/or are unsuitable.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL/ORAL SEMI-ELEMENTAL FEED 1.5KCAL/ML – Special Authority s Liquid18.06	ee SA1377 above – Hospital pharmacy [HP3] 1,000 ml OP Vital
ORAL ELEMENTAL FEED 0.8KCAL/ML - Special Authority see SA1377 above	e – Hospital pharmacy [HP3]
Liquid (grapefruit), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
Liquid (pineapple & orange), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
Liquid (summer fruits), 250 ml carton171.00	18 OP ✓ Elemental 028 Extra
ORAL ELEMENTAL FEED 1KCAL/ML - Special Authority see SA1377 above - Powder (unflavoured)4.50	- Hospital pharmacy [HP3] 80 α OP ✓ Vivonex TEN
SEMI-ELEMENTAL ENTERAL FEED 1KCAL/ML – Special Authority see SA13 Liquid	
T ·	Peptisorb
	✓ Peptisorb

(Peptisorb Liquid to be delisted 1 June 2023)

Subsidy (Manufacturer's Price) Fully Subsidised Per

Brand or Generic Manufacturer

Paediatric Products For Children With Low Energy Requirements

⇒SA1196 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 Child aged one to eight years; and
- 2 The child has a low energy requirement but normal protein and micronutrient requirements.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

- Both:
 - 1 The treatment remains appropriate and the patient is benefiting from treatment; and
 - 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

PAEDIATRIC ENTERAL FEED WITH FIBRE 0.76 KCAL/ML –	 Special Authority 	see SA1196 ab	ove -	 Hospital pharmacy [HP3]
Liquid	4.00	500 ml OP	1	Nutrini Low Energy
				Multi Fibre

Standard Supplements

⇒SA1859 Special Authority for Subsidy

Initial application — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 Any of the following:
 - 2.1 The patient has a condition causing malabsorption; or
 - 2.2 The patient has failure to thrive; or
 - 2.3 The patient has increased nutritional requirements; and
- 3 Nutrition goal has been set (eg reach a specific weight or BMI).

Renewal — (Children - indications other than exclusive enteral nutrition for Crohn's disease) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 The patient is under 18 years of age; and
- 2 The treatment remains appropriate and the patient is benefiting from treatment; and
- 3 A nutrition goal has been set (eg reach a specific weight or BMI).

Initial application — (Children - exclusive enteral nutrition for Crohn's disease) only from a gastroenterologist or dietitian on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria: All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 Dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Renewal — (Children - exclusive enteral nutrition for Crohn's disease) from any relevant practitioner on the recommendation of a gastroenterologist. Approvals valid for 3 months for applications meeting the following criteria:

continued...

Subsidy (Manufacturer's Price) Subsid		Fully	Brand or Generic	
\$	Per	1	Manufacturer	

All of the following:

- 1 The patient is under 18 years of age; and
- 2 It is to be used as exclusive enteral nutrition for the treatment of Crohn's disease; and
- 3 General Practitioners and dietitians must include the name of the gastroenterologist recommending treatment and the date the gastroenterologist was contacted.

Initial application — (Adults) from any relevant practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

1 Any of the following:

Patient is Malnourished

- 1.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 1.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 1.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months; and
- 2 Any of the following:

Patient has not responded to first-line dietary measures over a 4 week period by:

- 2.1 Increasing their food intake frequency (eg snacks between meals); or
- 2.2 Using high-energy foods (e.g. milkshakes, full fat milk, butter, cream, cheese, sugar etc); or
- 2.3 Using over the counter supplements (e.g. Complan); and
- 3 A nutrition goal has been set (e.g. to reach a specific weight or BMI).

Renewal — (Adults) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 A nutrition goal has been set (eg reach a specific weight or BMI); and
- 2 Any of the following:

Patient is Malnourished

- 2.1 Patient has a body mass index (BMI) of less than 18.5 kg/m²; or
- 2.2 Patient has unintentional weight loss greater than 10% within the last 3-6 months; or
- 2.3 Patient has a BMI of less than 20 kg/m² and unintentional weight loss greater than 5% within the last 3-6 months.

Initial application — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube or a nasogastric tube is to be inserted for feeding; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Is undergoing a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

continued...



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Subs	idised	Generic	
\$	Per	1	Manufacturer	

Renewal — (Short-term medical condition) from any relevant practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a nasogastric tube; or
- 2 Malignancy and is considered likely to develop malnutrition as a result; or
- 3 Has undergone a bone marrow transplant; or
- 4 Tempomandibular surgery or glossectomy; or
- 5 Both:
 - 5.1 Pregnant; and
 - 5.2 Any of the following:
 - 5.2.1 Patient is in early pregnancy (< 13 weeks) and has severe clinical hyperemesis gravidarum requiring admission to hospital and is unlikely to meet her nutritional requirements due to continuing hyperemesis gravidarum; or
 - 5.2.2 Patient has clinical hyperemesis gravidarum continuing past 13 weeks and either there is concern that the patient is unlikely to meet the Institute of Medicine's (1990) recommended weight gain guidelines for pregnancy or the patient's weight has not increased past her booking/pre-pregnancy weight; or
 - 5.2.3 Patient is having multiple births and is under the care of an obstetric team who consider the nutritional needs of the patient are not being met.

Initial application — (Long-term medical condition) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease: or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome: or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions: or
- 10 Epidermolysis bullosa; or
- 11 AIDS (CD4 count < 200 cells/mm³); or
- 12 Chronic pancreatitis.

Renewal — (Chronic disease OR tube feeding for patients who have previously been funded under Special Authority forms SA0702 or SA0583) from any relevant practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Is being fed via a tube or a tube is to be inserted for the purpose of feeding (not nasogastric tube refer to specific medical condition criteria); or
- 2 Cystic Fibrosis; or
- 3 Liver disease; or
- 4 Chronic Renal failure; or
- 5 Inflammatory bowel disease; or
- 6 Chronic obstructive pulmonary disease with hypercapnia; or
- 7 Short bowel syndrome; or
- 8 Bowel fistula; or
- 9 Severe chronic neurological conditions.

	Subsidy (Manufacturer's \$		Fully Brand or lised Generic Manufacturer
ENTERAL FEED 1.5KCAL/ML – Special Authority see SA1859 (Liquid		lospital pharmacy 250 ml OP 1,000 ml OP	/ [HP3] ✓ Ensure Plus HN ✓ Ensure Plus RTH ✓ Nutrison Energy
ENTERAL FEED 1KCAL/ML - Special Authority see SA1859 or Liquid		spital pharmacy [250 ml OP 1,000 ml OP	HP3] Isosource Standard Nutrison Standard RTH Osmolite RTH
ENTERAL FEED WITH FIBRE 0.83 KCAL/ML - Special Authori Liquid	•	on page 258 – Ho 1,000 ml OP	ospital pharmacy [HP3] Nutrison 800 Complete Multi Fibre
ENTERAL FEED WITH FIBRE 1 KCAL/ML - Special Authority s Liquid		page 258 – Hospi 1,000 ml OP	tal pharmacy [HP3] ✓ Jevity RTH ✓ Nutrison Multi Fibre
ENTERAL FEED WITH FIBRE 1.2KCAL/ML - Special Authority Liquid		page 258 - Hosp 1,000 ml OP	oital pharmacy [HP3] ✓ Jevity Plus
ENTERAL FEED WITH FIBRE 1.5KCAL/ML - Special Authority Liquid		page 258 – Hosp 1,000 ml OP	oital pharmacy [HP3] ✓ Jevity HiCal RTH ✓ Nutrison Energy Multi Fibre
ORAL FEED (POWDER) - Special Authority see SA1859 on pa Powder (chocolate)	•	al pharmacy [HP3 840 g OP	3] ✓ Sustagen Hospital Formula
Powder (vanilla)	26.00 14.00	850 g OP 840 g OP	✓ Ensure ✓ Sustagen Hospital Formula Active
	26.00	850 g OP	✓ Ensure

Subsidy		Fully	Brand or	
(Manufacturer's Pric	ce)	Subsidised	Generic	
\$	Per	/	Manufacturer	

ORAL FEED 1.5KCAL/ML - Special Authority see SA1859 on page 258 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, who have severe epidermolysis bullosa, or as exclusive enteral nutrition in children under the age of 18 years for the treatment of Crohn's disease, or for patients with COPD and hypercapnia, defined as CO2 value exceeding 55mmHg. The prescription must be endorsed accordingly.

Liquid (banana) – Higher subsidy of \$1.26 per 200 ml with Endorsement	0.72	200 ml OP	
Lituoisement	(1.26) (1.26)	200 IIII OF	Ensure Plus Fortisip
Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26) (1.26)		Ensure Plus Fortisip
Liquid (fruit of the forest) - Higher subsidy of \$1.26 per 200 ml			
with Endorsement	0.72	200 ml OP	
	(1.26)		Ensure Plus
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement		200 ml OP	
	(1.26)		Fortisip
Liquid (vanilla) – Higher subsidy of up to \$1.33 per 237 ml with			
Endorsement		237 ml OP	
	(1.33)		Ensure Plus
	0.72	200 ml OP	
	(1.26)		Ensure Plus
	(1.26)		Fortisip

ORAL FEED WITH FIBRE 1.5 KCAL/ML - Special Authority see SA1859 on page 258 - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (chocolate) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (strawberry) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisip Multi Fibre
Liquid (vanilla) - Higher subsidy of \$1.26 per 200 ml with			
Endorsement	0.72	200 ml OP	
	(1.26)		Fortisin Multi Fibre

High Calorie Products

⇒SA1195 Special Authority for Subsidy

Initial application — (Cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

All of the following:

- 1 Cystic fibrosis; and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements.

continued...

0.1.1		F. "	D 1	
Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sı	ubsidised	Generic	
\$	Per	✓	Manufacturer	

Initial application — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

All of the following:

- 1 Any of the following:
 - 1.1 any condition causing malabsorption; or
 - 1.2 faltering growth in an infant/child; or
 - 1.3 increased nutritional requirements: or
 - 1.4 fluid restricted: and
- 2 other lower calorie products have been tried; and
- 3 patient has substantially increased metabolic requirements or is fluid restricted.

Renewal — (Cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 3 years for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

Renewal — (Indications other than cystic fibrosis) only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

ENTERAL FEED 2 KCAL/ML - Special Authority see SA1195	on the previous p	age - Hospital p	harmacy [HP3]
Liquid	5.50	500 ml OP	✓ Nutrison
			Concentrated
	11.00	1,000 ml OP	✓ Ensure Two Cal HN
			RTH

ORAL FEED 2 KCAL/ML - Special Authority see SA1195 on the previous page - Hospital pharmacy [HP3]

Additional subsidy by endorsement is available for patients being bolus fed through a feeding tube, or who have severe epidermolysis bullosa. The prescription must be endorsed accordingly.

Liquid (vanilla) - Higher subsidy of \$1.90 per 200 ml with

Food Thickeners

⇒SA1106 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient has motor neurone disease with swallowing disorder.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

SPECIAL FOODS

	Subsidy		Fully	Brand or	
	(Manufacturer's Pric	e) Sub	sidised	Generic	
	\$	Per	•	Manufacturer	
FOOD THICKENER	- Special Authority see SA1106 on the previous page - Hospi	tal pharmac	y [HP3]		
Powder	6.53	300 g OP	✓ N	lutilis	
	7.25	380 g OP	√ F	eed Thickener	
				Karicare Antamil	

Gluten Free Foods

The funding of gluten free foods is no longer being actively managed by Pharmac from 1 April 2011. This means that we are no longer considering the listing of new products, or making subsidy, or other changes to the existing listings. As a result we anticipate that the range of funded items will reduce over time. Management of Coeliac disease with a gluten free diet is necessary for good outcomes. A range of gluten free options are available through retail outlets.

⇒SA1729 Special Authority for Subsidy

Initial application — (all patients) only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria: Either:

- 1 Gluten enteropathy has been diagnosed by biopsy; or
- 2 Patient suffers from dermatitis herpetiformis.

Initial application — (paediatric patients diagnosed by ESPGHAN criteria) only from a paediatric gastroenterologist. Approvals valid without further renewal unless notified where the paediatric patient fulfils ESPGHAN criteria for biopsy free diagnosis of coeliac disease.

GLUTEN FREE BAKING MIX – Special Authority see SA1729 abo		I pharmacy [HP3] 1,000 g OP	
	(5.15)	, 3	Healtheries Simple Baking Mix
GLUTEN FREE BREAD MIX - Special Authority see SA1729 abo	ve – Hospital	pharmacy [HP3]	
Powder	3.93	1,000 g OP	
	(7.32)	-	NZB Low Gluten Bread Mix
	3.51		
	(10.87)		Horleys Bread Mix
GLUTEN FREE FLOUR - Special Authority see SA1729 above -	Hospital phan	macy [HP3]	
Powder	5.62	2,000 g OP	
	(18.10)		Horleys Flour

	Subsidy		Fully	Brand or
	(Manufacturer's Price		idised	Generic
	\$	Per	•	Manufacturer
GLUTEN FREE PASTA - Special Authority see SA1729 on the	previous page – H	ospital pharm	acy [Hi	P3]
Buckwheat Spirals		250 g OP		
	(3.11)	•	О)rgran
Corn and Vegetable Shells	2.00 [°]	250 g OP		
·	(2.92)	· ·	О)rgran
Corn and Vegetable Spirals	, ,	250 g OP		ŭ
ů i	(2.92)	Ü	О)rgran
Rice and Corn Lasagne Sheets	, ,	200 g OP		ŭ
ů	(3.82)	Ü	О)rgran
Rice and Corn Macaroni		250 g OP	_	··· g ····
	(2.92)		C)rgran
Rice and Corn Penne	, ,	250 g OP	_	··· g ····
	(2.92)		C)rgran
Rice and Maize Pasta Spirals	, ,	250 g OP	_	··· g ····
	(2.92)		0)rgran
Rice and Millet Spirals	, ,	250 g OP		· · · · · ·
	(3.11)	_00 g 0.	0)rgran
Rice and corn spaghetti noodles	` ,	375 g OP	Ŭ	rigian
Thoo and com spagnout necessor	(2.92)	0,0 g 0.	0)rgran
Vegetable and Rice Spirals	, ,	250 g OP	Ŭ	rigian
rogotable and rive opilate	(2.92)	200 g 0.	0)rgran
Italian long style spaghetti	, ,	220 g OP	Č	ngian
naman long style spagnetti	(3.11)	220 g Oi	0)rgran
	(0.11)		·	/igiaii

Foods And Supplements For Inborn Errors Of Metabolism

⇒SA1108 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid without further renewal unless notified for applications meeting the following criteria:

Any of the following:

- 1 Dietary management of homocystinuria; or
- 2 Dietary management of maple syrup urine disease; or
- 3 Dietary management of phenylketonuria (PKU); or
- 4 For use as a supplement to the Ketogenic diet in patients diagnosed with epilepsy.

Supplements For Homocystinuria

Supplements For MSUD

AMINOACID FORMULA WITHOUT VALINE, LEUCINE AND ISOLEUCINE - Special Authority see SA1108 above - Hospital pharmacy [HP3]

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
` \$	Per 🗸	Manufacturer	

Supplements For PKU

AMINOACID FORMULA WITHOUT PHENYLALANINE - Special Authority see SA1108 on the previous page - Hospital pharmacy [HP3]

Tabs	99.00	75 OP	✓ Phlexy 10
Powder (berry) 28 g sachets		30	✓ PKU Lophlex Powder
Powder (chocolate) 36 g sachet	393.00	30	✓ PKU Anamix Junior Chocolate
Powder (neutral) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (neutral) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (orange) 28 g sachets		30	✓ PKU Lophlex Powder
Powder (orange) 36 g sachet	393.00	30	✓ PKU Anamix Junior Orange
Powder (unflavoured) 28 g sachets	936.00	30	✓ PKU Lophlex Powder
Powder (unflavoured) 36 g sachets	393.00	30	✓ PKU Anamix Junior
Powder (vanilla) 36 g sachet	393.00	30	✓ PKU Anamix Junior Vanilla
Infant formula	174.72	400 g OP	✓ PKU Anamix Infant
Powder (orange)	320.00	500 g OP	✓ XP Maxamum
Powder (unflavoured)		500 g OP	✓ XP Maxamum
Liquid (berry)		125 ml OP	✓ PKU Anamix Junior LQ
Liquid (orange)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (unflavoured)	13.10	125 ml OP	✓ PKU Anamix Junior LQ
Liquid (forest berries), 250 ml carton	540.00	18 OP	 Easiphen Liquid
Liquid (juicy tropical) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20
Oral semi-solid (berries) 109 g		36 OP	✓ PKU Lophlex Sensation 20
Liquid (juicy berries) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy citrus) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy orange) 62.5 ml	939.00	60 OP	✓ PKU Lophlex LQ 10
Liquid (juicy berries) 125 ml		30 OP	✓ PKU Lophlex LQ 20
Liquid (juicy orange) 125 ml	936.00	30 OP	✓ PKU Lophlex LQ 20

(PKU Lophlex Powder Powder (unflavoured) 28 g sachets to be delisted 1 December 2022) (PKU Anamix Junior Powder (unflavoured) 36 g sachets to be delisted 1 December 2022)

Foods

	Subsidy (Manufacturer's P	rice) Subsi	Fully idised	Brand or Generic Manufacturer
LOW PROTEIN PASTA - Special Authority see SA1108 on page	e 265 – Hospital	pharmacy [HP3	3]	
Animal shapes		500 g OP		.oprofin
Lasagne	5.95	250 g OP	√ L	.oprofin
Low protein rice pasta	11.91	500 g OP	√ L	.oprofin
Macaroni	5.95	250 g OP	√ L	.oprofin
Penne	11.91	500 g OP	√ L	.oprofin
Spaghetti	11.91	500 g OP	√ L	oprofin
Spirals	11.91	500 g OP	√ L	.oprofin

Infant Formulae

For Williams Syndrome

⇒SA1110 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year where the patient is an infant suffering from Williams Syndrome and associated hypercalcaemia.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 1 year for applications meeting the following criteria:

Both:

- 1 The treatment remains appropriate and the patient is benefiting from treatment; and
- 2 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and date contacted.

LOW CALCIUM INFANT FORMULA - Special Authority see SA1110 above - Hospital pharmacy [HP3]
Powder44.40 400 g OP ✓ Locasol

Gastrointestinal and Other Malabsorptive Problems

MINO ACID FORMULA – Special Authority see SA2092 below – Hospital phal Powder43.60	400 g OP	✓ Alfamino
Powder (unflavoured)53.00	400 g OP	✓ Alfamino Junior ✓ Elecare ✓ Elecare LCP
		 ✓ Neocate Gold ✓ Neocate Junior Unflavoured
Powder (vanilla)53.00	400 g OP	✓ Neocate SYNEO✓ Elecare✓ Neocate Junior✓ Vanilla

⇒SA2092 Special Authority for Subsidy

Initial application — (Infants under 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 History of anaphylaxis to cow's milk protein formula or dairy products; or
- 2 Eosinophilic oesophagitis; or
- 3 Ultra-short gut; or
- 4 Severe Immune deficiency; or

continued...

Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sul	bsidised	Generic	
\$	Per	✓	Manufacturer	

- 5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
- 6 Both:
 - 6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 6.2 Fither:
 - 6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or 6.2.2 Patient has IgE mediated allergy.

Initial application — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria:

Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products: or
 - 2.2 Eosinophilic oesophagitis; or
 - 2.3 Ultra-short gut: or
 - 2.4 Severe Immune deficiency; or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Either:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number; or
 - 2.6.2.2 Patient has IgE mediated allergy.

Renewal — (Infants up to 12 months of age) from any relevant practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Either:

- 1 Both:
 - 1.1 Patient has IgE mediated allergy; and
 - 1.2 All of the following:
 - 1.2.1 Patient remains allergic to cow's milk; and
 - 1.2.2 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy or extensively hydrolysed infant formula has been undertaken; and
 - 1.2.3 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and 1.2.4 Amino acid formula is required for a nutritional deficit; and
 - 1.2.5 It has been more than three months from the previous approval; or
- 2 Both:
 - 2.1 Patient has non IgE mediated severe gastrointestinal intolerance (including eosinophilic oesophagitis, ultra-short gut and severe immune deficiency); and
 - 2.2 All of the following:
 - 2.2.1 An assessment as to whether the infant can be transitioned to a cow's milk protein, soy, or extensively hydrolysed infant formula has been undertaken; and
 - 2.2.2 The outcome of the assessment is that the infant continues to require an amino acid infant formula; and

continued...

Subsidy	Fully	Brand or
(Manufacturer's Price)	Subsidised	Generic
\$	Per 🗸	Manufacturer

- 2.2.3 Amino acid formula is required for a nutritional deficit; and
- 2.2.4 It has been more than three months from the previous approval.

Renewal — (Children 12 months of age and over) only from a paediatrician, paediatric gastroenterologist, paediatric immunologist or dietitian on the recommendation of a paediatrician, paediatric gastroenterologist or paediatric immunologist. Approvals valid for 6 months for applications meeting the following criteria: Both:

- 1 Either:
 - 1.1 Applicant is a paediatrician, paediatric gastroenterologist or paediatric immunologist; or
 - 1.2 Applicant is a dietitian and confirms that a paediatrician, paediatric gastroenterologist or paediatric immunologist has been consulted within the last 12 months and has recommended treatment for the patient; and
- 2 Any of the following:
 - 2.1 History of anaphylaxis to cow's milk protein formula or dairy products; or
 - 2.2 Eosinophilic oesophagitis: or
 - 2.3 Ultra-short gut; or
 - 2.4 Severe Immune deficiency: or
 - 2.5 Extensively hydrolysed formula has been trialled in an inpatient setting and is clinically inappropriate; or
 - 2.6 Both:
 - 2.6.1 Extensively hydrolysed formula has been reasonably trialled for 2-4 weeks and is inappropriate due to documented severe intolerance or allergy or malabsorption; and
 - 2.6.2 Fither:
 - 2.6.2.1 The patient has a valid Special Authority approval for extensively hydrolysed formula: approval number: or
 - 2.6.2.2 Patient has IgE mediated allergy.

Initial application — (for patients who have a current funding under Special Authority form SA1557) only from a dietitian. relevant specialist or vocationally registered general practitioner. Approvals valid for 3 months for applications meeting the following criteria:

All of the following:

- 1 Patient has a valid Special Authority approval for extensively hydrolysed formula (SA1557); and
- 2 Extensively hydrolysed formula (Aptamil Gold+ Pepti Junior, AllerPro SYNEO 1 and 2) is unable to be supplied at this time;
- 3 The approval only applies to funded dispensings of Neocate Gold and Neocate Syneo.

Note: This criteria is short term funding to cover an out-of-stock situation on some extensively hydrolysed formula powder funded under Special Authority form SA1557. There is no renewal criteria under this restriction.

ENTERAL LIQUID PEPTIDE FORMULA - Special Authority see SA1953 below - Hospital pharmacy [HP3]

Liquid 1 kcal/ml10.45	500 ml OP	✓ Nutrini Peptisorb
Liquid 1.5 kcal/ml15.68	500 ml OP	✓ Nutrini Peptisorb
		Energy

⇒SA1953 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient has impaired gastrointestinal function and either cannot tolerate polymeric feeds, or polymeric feeds are unsuitable: and
- 2 Any of the following:
 - 2.1 Severe malabsorption; or
 - 2.2 Short bowel syndrome: or

continued...



Subsidy		Fully	Brand or	
(Manufacturer's Price)	Sub	sidised	Generic	
\$	Per	/	Manufacturer	

- 2.3 Intractable diarrhoea; or
- 2.4 Biliary atresia; or
- 2.5 Cholestatic liver diseases causing malabsorption; or
- 2.6 Cystic fibrosis; or
- 2.7 Proven fat malabsorption; or
- 2.8 Severe intestinal motility disorders causing significant malabsorption; or
- 2.9 Intestinal failure: or
- 2.10 Both:
 - 2.10.1 The patient is currently receiving funded amino acid formula; and
 - 2.10.2 The patient is to be trialled on, or transitioned to, an enteral liquid peptide formula; and
- 3 Either:
 - 3.1 A semi-elemental or partially hydrolysed powdered feed has been reasonably trialled and considered unsuitable; or
 - 3.2 For step down from intravenous nutrition.

Note: A reasonable trial is defined as a 2-4 week trial.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the patient can be transitioned to a cows milk protein or soy infant formula or extensively hydrolysed formula has been undertaken; and
- 2 The outcome of the assessment is that the patient continues to require an enteral liquid peptide formula; and
- 3 General practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

EXTENSIVELY HYDROLYSED FORMULA - Special Authority see SA1557 below - Hospital pharmacy [HP3]

(Aptamil Gold+ Pepti Junior Powder to be delisted 1 November 2022) (Aptamil AllerPro SYNEO 1 Powder to be delisted 1 November 2022) (Aptamil AllerPro SYNEO 2 Powder to be delisted 1 November 2022)

⇒SA1557 Special Authority for Subsidy

Initial application only from a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

Any of the following:

- 1 Both:
 - 1.1 Cows milk formula is inappropriate due to severe intolerance or allergy to its protein content; and
 - 1.2 Either:
 - 1.2.1 Soy milk formula has been reasonably trialled without resolution of symptoms: or
 - 1.2.2 Soy milk formula is considered clinically inappropriate or contraindicated; or
- 2 Severe malabsorption; or

continued...

Subsidy		Fully	Brand or
(Manufacturer's Price)		Subsidised	Generic
\$	Per	/	Manufacturer

- 3 Short bowel syndrome; or
 - 4 Intractable diarrhoea; or
 - 5 Biliary atresia; or
- 6 Cholestatic liver diseases causing malsorption; or
- 7 Cystic fibrosis; or
- 8 Proven fat malabsorption; or
- 9 Severe intestinal motility disorders causing significant malabsorption; or
- 10 Intestinal failure; or
- 11 All of the following:
 - 11.1 For step down from Amino Acid Formula; and
 - 11.2 The infant is currently receiving funded amino acid formula; and
 - 11.3 The infant is to be trialled on, or transitioned to, an extensively hydrolysed formula; and
 - 11.4 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Note: A reasonable trial is defined as a 2-4 week trial, or signs of an immediate IqE mediated allergic reaction.

Renewal only from a dietitian, relevant specialist, vocationally registered general practitioner or general practitioner on the recommendation of a dietitian, relevant specialist or vocationally registered general practitioner. Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 An assessment as to whether the infant can be transitioned to a cows milk protein or soy infant formula has been undertaken; and
- 2 The outcome of the assessment is that the infant continues to require an extensively hydrolysed infant formula; and
- 3 General Practitioners must include the name of the dietitian, relevant specialist or vocationally registered general practitioner and the date contacted.

Fluid Restricted

⇒SA1698 Special Authority for Subsidy

Initial application only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian. Approvals valid for 12 months for applications meeting the following criteria:

All of the following:

- 1 Patient is fluid restricted or volume intolerant and has been diagnosed with faltering growth; and
- 2 Patient is under the care of a paediatrician or dietitian who has recommended treatment with a high energy infant formula; and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

Renewal only from a paediatrician, dietitian or general practitioner on the recommendation of a paediatrician or dietitian.

Approvals valid for 6 months for applications meeting the following criteria:

All of the following:

- 1 Patient continues to be fluid restricted or volume intolerant and has faltering growth; and
- 2 Patient is under the care of a hospital paediatrician or dietitian who has recommended treatment with a high energy infant formula: and
- 3 Patient is under 18 months of age or weighs less than 8 kg.

Note: 'Volume intolerant' patients are those who are unable to tolerate an adequate volume of infant formula to achieve expected growth rate. These patients should have first trialled appropriate clinical alternative treatments, such as concentrating, fortifying and adjusting the frequency of feeding.

SPECIAL FOODS

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per ✓ Manufacturer

Ketogenic Diet

⇒SA1197 Special Authority for Subsidy

Initial application only from a metabolic physician or paediatric neurologist. Approvals valid for 3 months where the patient has intractable epilepsy, pyruvate dehydrogenase deficiency or glucose transported type-1 deficiency and other conditions requiring a ketogenic diet.

Renewal only from a metabolic physician or paediatric neurologist. Approvals valid for 2 years where the patient is on a ketogenic diet and the patient is benefiting from the diet.

HIGH FAT LOW CARBOHYDRATE FORMULA - Special Authority see SA1197 above - Retail pharmacy

Powder (unflavoured)35.50	300 g OP	✓ KetoCal 4:1
		✓ Ketocal 3:1
Powder (vanilla)35.50	300 a OP	✓ KetoCal 4:1

SECTION I: NATIONAL IMMUNISATION SCHEDULE

Subsidy (Manufacturer's Price) \$

Subsidised Per

Fully

Brand or Generic Manufacturer

Vaccinations

BACILLUS CALMETTE-GUERIN VACCINE - [Xpharm]

For infants at increased risk of tuberculosis. Increased risk is defined as:

- 1) living in a house or family with a person with current or past history of TB; or
- 2) having one or more household members or carers who within the last 5 years lived in a country with a rate of TB > or equal to 40 per 100,000 for 6 months or longer; or
- 3) during their first 5 years will be living 3 months or longer in a country with a rate of TB > or equal to 40 per 100,000 Note a list of countries with high rates of TB are available at www.health.govt.nz/tuberculosis (search for downloads) or www.bcgatlas.org/index.php.
- Inj Mycobacterium bovis BCG (Bacillus Calmette-Guerin),

Danish strain 1331, live attenuated, vial with diluent.......................0.00

10 **✓ BCG Vaccine**

DIPHTHERIA. TETANUS AND PERTUSSIS VACCINE

- a) Only on a prescription
- b) No patient co-payment payable
- c)
- A) Funded for any of the following criteria:
 - 1) A single dose for pregnant women in the second or third trimester of each pregnancy; or
 - 2) A single dose for parents or primary caregivers of infants admitted to a Neonatal Intensive Care Unit or Specialist Care Baby Unit for more than 3 days, who had not been exposed to maternal vaccination at least 14 days prior to birth; or
 - A course of up to four doses is funded for children from age 7 up to the age of 18 years inclusive to complete full primary immunisation; or
 - 4) An additional four doses (as appropriate) are funded for (re-)immunisation for patients post haematopoietic stem cell transplantation or chemotherapy; pre or post splenectomy; pre- or post solid organ transplant, renal dialysis and other severely immunosuppressive regimens; or
 - 5) A single dose for vaccination of patients aged from 65 years old; or
 - 6) A single dose for vaccination of patients aged from 45 years old who have not had 4 previous tetanus doses; or
 - 7) For vaccination of previously unimmunised or partially immunised patients; or
 - 8) For revaccination following immunosuppression; or
 - 9) For boosting of patients with tetanus-prone wounds.

Notes: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes.

- B) Contractors will be entitled to claim payment from the Funder for the supply of diphtheria, tetanus and pertussis vaccine to patients eligible under the above criteria pursuant to their contract with Te Whatu Ora Health New Zealand for subsidised immunisation, and they may only do so in respect of the diphtheria, tetanus and pertussis vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraphs 1 – 9 above.
- Inj 2 IU diphtheria toxoid with 20 IU tetanus toxoid, 8 mcg

pertussis toxoid. 8 mcg pertussis filamentous

10

✓ <u>Boostrix</u>
✓ Boostrix

	Subsidy	Fully	
	(Manufacturer's Price) \$	Subsidised Per 🗸	d Generic Manufacturer
DIDUTUEDIA TETANILIS DEDTLISSIS AND DOLLO VACCINE			manadata o
DIPHTHERIA, TETANUS, PERTUSSIS AND POLIO VACCINE – Funded for any of the following:	[xpriarrii]		
A single dose for children up to the age of 7 who have c	ompleted primary imr	munication: or	
A course of four vaccines is funded for catch up program primary immunisation; or			years) to complete full
 An additional four doses (as appropriate) are funded for pre- or post splenectomy; pre- or post solid organ transpregimens; or 			
Five doses will be funded for children requiring solid org	an transplantation.		
Note: Please refer to the Immunisation Handbook for approp	•	ch up program	mes.
Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg		p p - 9	
pertussis toxoid, 25 mcg pertussis filamentous haemagglutinin, 8 mcg pertactin and 80 D-antigen units			
poliomyelitis virus in 0.5ml syringe	0.00	10	Infanrix IPV
DIPHTHERIA, TETANUS, PERTUSSIS, POLIO, HEPATITIS B AN			
Xpharm]	ND HAEIWOPHILUS I	NFLUENZAE	I THE B VACCINE -
Funded for patients meeting any of the following criteria:			
1) Up to four doses for children up to and under the age of	10 for primary immur	nisation; or	
2) An additional four doses (as appropriate) are funded for			
10 who are patients post haematopoietic stem cell trans			
post solid organ transplant, renal dialysis and other seve			
3) Up to five doses for children up to and under the age of			
Note: A course of up-to four vaccines is funded for catch up p			
to complete full primary immunisation. Please refer to the Improgrammes	nunisation handbook	tior the approp	priate scriedule for catch up
programmes. Inj 30 IU diphtheria toxoid with 40 IU tetanus toxoid, 25 mcg			
pertussis toxoid, 25 mcg pertussis filamentous			
haemagglutinin, 8 mcg pertactin, 80 D-Ag U polio virus,			
10 mcg hepatitis B surface antigen in 0.5 ml syringe	0.00	10	Infanrix-hexa
HAEMOPHILUS INFLUENZAE TYPE B VACCINE - [Xpharm]			
One dose for patients meeting any of the following:			
For primary vaccination in children; or			
2) An additional dose (as appropriate) is funded for (re-)imi	munisation for patient	ts post haemat	topoietic stem cell
transplantation, or chemotherapy; functional asplenic; p			
or post cochlear implants, renal dialysis and other sever			
For use in testing for primary immunodeficiency disease	s, on the recommend	lation of an into	ernal medicine physician or
paediatrician.			
Haamanhilua Influenzaa tuna P naluasaaharida 10 maa			
Haemophilus Influenzae type B polysaccharide 10 mcg conjugated to tetanus toxoid as carrier protein 20-40 mcg			
prefilled syringe plus vial 0.5 ml		1 🗸	Hiberix
HEPATITIS A VACCINE – [Xpharm]		•	
Funded for patients meeting any of the following criteria:			
Two vaccinations for use in transplant patients; or			
Two vaccinations for use in children with chronic liver dia	sease: or		
One dose of vaccine for close contacts of known hepatit			
Inj 1440 ELISA units in 1 ml syringe			<u>Havrix</u>
Inj 720 ELISA units in 0.5 ml syringe	0.00	1	Havrix Junior

	NATIONAL	. I IVI IVI	IUNISATI	ON SCHEDULE
	Subsidy (Manufacturer's Price) \$) Per	Fully Subsidised	Brand or Generic Manufacturer
HEPATITIS B RECOMBINANT VACCINE – [Xpharm] Inj 10 mcg per 0.5 ml prefilled syringe Funded for patients meeting any of the followin		1	✓ E	Engerix-B
 for household or sexual contacts of know for children born to mothers who are hepe for children up to and under the age of 18 serology and require additional vaccination for HIV positive patients; or for hepatitis C positive patients; or for patients following non-consensual sex for patients following immunosuppression for solid organ transplant patients; or for post-haematopoietic stem cell transplant following needle stick injury. 	n acute hepatitis B patients or atitis B surface antigen (HBsA B years inclusive who are cons on or require a primary course tual intercourse; or n; or	g) pos sidered	itive; or I not to have	e achieved a positive
Inj 20 mcg per 1 ml prefilled syringe	ng criteria: n acute hepatitis B patients or atitis B surface antigen (HBsA g years inclusive who are cons on or require a primary course atual intercourse; or n; or	g) pos idered	titis B carrier itive; or I not to have	e achieved a positive
HUMAN PAPILLOMAVIRUS (6, 11, 16, 18, 31, 33, 45, Any of the following: 1) Maximum of two doses for children aged 14 y 2) Maximum of three doses for patients meeting 1) People aged 15 to 26 years inclusive; or 2) Either: People aged 9 to 26 years inclusive 1) Confirmed HIV infection; or 2) Transplant (including stem cell) pa 3) Maximum of four doses for people aged 9 to 2	rears and under; or any of the following criteria: r tients: or 26 years inclusive post chemo		·	
Inj 270 mcg in 0.5 ml syringe	0.00	10	√ <u>c</u>	Gardasil 9

Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic \$ Per ✔ Manufacturer	
--	--

INFLUENZA VACCINE

Inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine)

A) INFLUENZA VACCINE - child aged 6 months to 35 months

is available each year for patients aged 6 months to 35 months who meet the following criteria, as set by Pharmac:

- i) have any of the following cardiovascular diseases
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
- ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
- iii) have diabetes; or
- iv) have chronic renal disease; or
- v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
- vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - j) pre and post splenectomy, or
 - k) down syndrome, or
- vii) have been hospitalised for respiratory illness or have a history of significant respiratory illness;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy.
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Doctors are the only Contractors entitled to claim payment for the supply of influenza vaccine inj 30 mcg in 0.25 ml syringe (paediatric quadrivalent vaccine) to patients eligible under the above criteria for subsidised immunisation and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.

Inj 60 mcg in 0.5 ml syringe (quadrivalent vaccine)	110.00	10	Afluria Quad
			(2022 formulation)

Subsidy	Fully	Brand or	
(Manufacturer's Price)	Subsidised	Generic	
\$	Per 🗸	Manufacturer	

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) INFLUENZA VACCINE - people 3 years and over

is available each year for patients aged 3 years and over who meet the following criteria, as set by Pharmac:

- a) all people 65 years of age and over; or
- b) People 55 to 64 years of age (inclusive) and is Māori or any Pacific ethnicity; or
- c) people under 65 years of age who:
 - i) have any of the following cardiovascular diseases:
 - a) ischaemic heart disease, or
 - b) congestive heart failure, or
 - c) rheumatic heart disease, or
 - d) congenital heart disease, or
 - e) cerebo-vascular disease; or
 - ii) have either of the following chronic respiratory diseases:
 - a) asthma, if on a regular preventative therapy, or
 - b) other chronic respiratory disease with impaired lung function; or
 - iii) have diabetes; or
 - iv) have chronic renal disease; or
 - v) have any cancer, excluding basal and squamous skin cancers if not invasive; or
 - vi) have any of the following other conditions:
 - a) autoimmune disease, or
 - b) immune suppression or immune deficiency, or
 - c) HIV, or
 - d) transplant recipients, or
 - e) neuromuscular and CNS diseases/disorders, or
 - f) haemoglobinopathies, or
 - g) are children on long term aspirin, or
 - h) have a cochlear implant, or
 - i) errors of metabolism at risk of major metabolic decompensation, or
 - i) pre and post splenectomy, or
 - k) down syndrome, or
 - vii) are pregnant; or
- d) children 3 and 4 years of age (inclusive) who have been hospitalised for respiratory illness or have a history of significant respiratory illness; or
- e) people under 65 years of age who:
 - i) have any of the following serious mental health conditions:
 - a) schizophrenia, or
 - b) major depressive disorder, or
 - c) bipolar disorder, or
 - d) schizoaffective disorder, or
 - ii) are currently accessing secondary or tertiary mental health and addiction services; or
- f) children 3 to 12 years of age (inclusive), from 1 July 2022 to 31 December 2022;

Unless meeting the criteria set out above, the following conditions are excluded from funding:

- a) asthma not requiring regular preventative therapy,
- b) hypertension and/or dyslipidaemia without evidence of end-organ disease.
- B) Contractors will be entitled to claim payment for the supply of influenza vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the influenza vaccine listed in the Pharmaceutical Schedule.
- C) Contractors may only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

	Subsidy	Fully	Brand or
(Manu	facturer's Price)	Subsidised	Generic
	\$ Pe	er 🗸	Manufacturer

MEASLES, MUMPS AND RUBELLA VACCINE

- a) Only on a prescription
- b) No patient co-payment payable

c)

A) Measles, mumps and rubella vaccine

A maximum of two doses for any patient meeting the following criteria:

- 1) For primary vaccination in children; or
- 2) For revaccination following immunosuppression; or
- 3) For any individual susceptible to measles, mumps or rubella; or
- 4) A maximum of three doses for children who have had their first dose prior to 12 months.

Note: Please refer to the Immunisation Handbook for appropriate schedule for catch up programmes. Although a price is listed for the vaccine, doctors can still order measles mumps and rubella vaccine free of charge, as with other Schedule vaccines.

- B) Contractors will be entitled to claim payment for the supply of measles, mumps and rubella vaccine to patients eligible under the above criteria pursuant to their contract with Health NZ for subsidised immunisation, and they may only do so in respect of the measles, mumps and rubella vaccine listed in the Pharmaceutical Schedule.
- C) Contractors can only claim for patient populations within the criteria that are covered by their contract, which may be a sub-set of the population described in paragraph A above.

Inj, measles virus 1,000 CCID50, mumps virus 5,012 CCID50,

Rubella virus 1,000 CCID50; prefilled syringe/ampoule of

MENINGOCOCCAL (GROUPS A, C, Y AND W-135) CONJUGATE VACCINE - [Xpharm]

Either:

- A) Any of the following:
 - Up to three doses and a booster every five years for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre or post solid organ transplant: or
 - 2) One dose for close contacts of meningococcal cases of any group; or
 - 3) One dose for person who has previously had meningococcal disease of any group; or
 - 4) A maximum of two doses for bone marrow transplant patients; or
 - 5) A maximum of two doses for person pre- and post-immunosuppression*; or
- B) Both
 - 1) Person is aged between 13 and 25 years, inclusive; and
 - 2) Either:
 - i) One dose for individuals who are entering within the next three months, or in their first year of living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons; or
 - ii) One dose for individuals who are currently living in boarding school hostels, tertiary education halls of residence, military barracks, or prisons, from 1 December 2019 to 30 November 2021.

Note: children under seven years of age require two doses 8 weeks apart, a booster dose three years after the primary series and then five yearly.

*Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days.

Inj 4 mcg of each meningococcal polysaccharide conjugated to

a total of approximately 48 mcg of diphtheria toxoid carrier

NATIONAL IMMUNISATION SCHEDULE Subsidy Fully Brand or (Manufacturer's Price) Subsidised Generic Per Manufacturer MENINGOCOCCAL B MULTICOMPONENT VACCINE - [Xpharm] Either: A) Both: 1) Child is under one year of age; and 2) Any of the following: i) up to three doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to three doses for close contacts of meningococcal cases of any group; or iii) up to three doses for child who has previously had meningococcal disease of any group; or iv) up to three doses for bone marrow transplant patients; or v) up to three doses for child pre- and post-immunosuppression*; or B) Both: 1) Person is one year of age or over; and 2) Any of the following: i) up to two doses and a booster every five years for patients pre- and post-splenectomy and for patients with functional or anatomic asplenia, HIV, complement deficiency (acquired or inherited), or pre- or post-solid organ transplant; or ii) up to two doses for close contacts of meningococcal cases of any group; or iii) up to two doses for person who has previously had meningococcal disease of any group; or iv) up to two doses for bone marrow transplant patients: or v) up to two doses for person pre- and post-immunosuppression*. *Immunosuppression due to corticosteroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Bexsero MENINGOCOCCAL C CONJUGATE VACCINE - [Xpharm] Both: 1) The child is under 9 months of age; and 2) Any of the following: 1) Up to three doses for patients pre- and post splenectomy and for patients with functional or anatomic asplenia, HIV. complement deficiency (acquired or inherited), or pre or post solid organ transplant; or 2) Two doses for close contacts of meningococcal cases of any group; or 3) Two doses for child who has previously had meningococcal disease of any group; or 4) A maximum of two doses for bone marrow transplant patients; or 5) A maximum of two doses for child pre- and post-immunosuppression*. Note: children under nine months of age require two doses 8 weeks apart. Refer to the Immunisation Handbook for booster schedules with meningococcal ACWY vaccine. *Immunosuppression due to steroid or other immunosuppressive therapy must be for a period of greater than 28 days. ✓ Neisvac-C PNEUMOCOCCAL (PCV10) CONJUGATE VACCINE - [Xpharm] 1) A primary course of three doses for previously unvaccinated individuals up to the age of 59 months inclusive Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes Ini 1 mcg of pneumococcal polysaccharide serotypes 1, 5, 6B. 7F, 9V, 14 and 23F; 3 mcg of pneumococcal polysaccharide serotypes 4, 18C and 19F in 0.5 ml

10

Synflorix

Subsidy	F	ully	Brand or
(Manufacturer's Price)	Subsid	ised	Generic
\$	Per	1	Manufacturer

PNEUMOCOCCAL (PCV13) CONJUGATE VACCINE - [Xpharm]

Any of the following:

- Two doses are funded for high risk children (over the age of 12 months and under 18 years) who have previously received two doses of the primary course of PCV10; or
- 2) Up to an additional four doses (as appropriate) are funded for high risk children aged under 5 years for (re-)immunisation of patients with any of the following:
 - a) on immunosuppressive therapy or radiation therapy, vaccinate when there is expected to be a sufficient immune response; or
 - b) with primary immune deficiencies; or
 - c) with HIV infection; or
 - d) with renal failure, or nephrotic syndrome; or
 - e) who are immune-suppressed following organ transplantation (including haematopoietic stem cell transplant); or
 - f) with cochlear implants or intracranial shunts; or
 - g) with cerebrospinal fluid leaks; or
 - h) receiving corticosteroid therapy for more than two weeks, and who are on an equivalent daily dosage of prednisone of 2 mg/kg per day or greater, or children who weigh more than 10 kg on a total daily dosage of 20 mg or greater; or
 - i) with chronic pulmonary disease (including asthma treated with high-dose corticosteroid therapy); or
 - j) pre term infants, born before 28 weeks gestation; or
 - k) with cardiac disease, with cyanosis or failure; or
 - I) with diabetes; or
 - m) with Down syndrome; or
 - n) who are pre-or post-splenectomy, or with functional asplenia; or
- 3) Up to an additional four doses (as appropriate) are funded for (re-)immunisation of patients 5 years and over with HIV, for patients pre or post haematopoietic stem cell transplantation, or chemotherapy; pre- or post splenectomy; functional asplenia, pre- or post- solid organ transplant, renal dialysis, complement deficiency (acquired or inherited), cochlear implants, intracranial shunts, cerebrospinal fluid leaks or primary immunodeficiency; or
- 4) For use in testing for primary immunodeficiency diseases, on the recommendation of an internal medicine physician or paediatrician.

Note: please refer to the Immunisation Handbook for the appropriate schedule for catch up programmes

Ini 30.8 mcg of pneumococcal polysaccharide serotypes 1, 3, 4,

	NATIONAL	IMMUNISA	TION SCHEDULE
	Subsidy (Manufacturer's Price) \$	Ful Subsidise Per •	,
PNEUMOCOCCAL (PPV23) POLYSACCHARIDE VACCINE	– [Xpharm]		
Either:			
 Up to three doses (as appropriate) for patients with I chemotherapy; pre- or post-splenectomy or with function complement deficiency (acquired or inherited), cochi All of the following: a) Patient is a child under 18 years for (re-)immur 	ctional asplenia, pre- or lear implants, or primary	post-solid orga	in transplant, renal dialysis,
b) Treatment is for a maximum of two doses; and			
c) Any of the following: i) on immunosuppressive therapy or radiati immune response; or ii) with primary immune deficiencies; or iii) with HIV infection; or iv) with renal failure, or nephrotic syndrome; v) who are immune-suppressed following or or vi) with cochlear implants or intracranial shu vii) with cerebrospinal fluid leaks; or viii) receiving corticosteroid therapy for more prednisone of 2 mg/kg per day or greater	or gan transplantation (inc nts; or than two weeks, and wh	cluding haemat no are on an ec	opoietic stem cell transplant); quivalent daily dosage of
20 mg or greater; or ix) with chronic pulmonary disease (including x) pre term infants, born before 28 weeks grail with cardiac disease, with cyanosis or fair xii) with diabetes; or xiii) with Down syndrome; or xiv) who are pre-or post-splenectomy, or with	g asthma treated with hi estation; or lure; or		, ,
Inj 575 mcg in 0.5 ml prefilled syringe (25 mcg of each 23 pneumococcal serotype)	0.00	1 🗸	Pneumovax 23
POLIOMYELITIS VACCINE - [Xpharm] Up to three doses for patients meeting either of the followi 1) For partially vaccinated or previously unvaccinated in 2) For revaccination following immunosuppression.	ng: ndividuals; or		
Note: Please refer to the Immunisation Handbook for app Inj 80D antigen units in 0.5 ml syringe			nmes. ′ IPOL
ROTAVIRUS ORAL VACCINE – [Xpharm] Maximum of two doses for patients meeting the following: 1) first dose to be administered in infants aged under 1 2) no vaccination being administered to children aged 2	•		
Oral susp live attenuated human rotavirus 1,000,000 CCID50 per dose, prefilled oral applicator	0.00	10	Rotarix

	Subsidy (Manufacturer's Price) \$	Subs Per	Fully idised	Brand or Generic Manufacturer
VARICELLA VACCINE [CHICKENPOX VACCINE] – [Xpharm] Either:				
1) Maximum of one dose for primary vaccination for eithe	r:			
a) Any infant born on or after 1 April 2016; or				
For previously unvaccinated children turning 11 y varicella infection (chickenpox), or	rears old on or after 1	July 2017,	who ha	ave not previously had a
2) Maximum of two doses for any of the following:				
a) Any of the following for non-immune patients:				
i) with chronic liver disease who may in future		nsplantatio	n; or	
ii) with deteriorating renal function before tran	splantation; or			
iii) prior to solid organ transplant; or				
iv) prior to any elective immunosuppression*, or				
v) for post exposure prophylaxis who are imm				
b) For patients at least 2 years after bone marrow tr				
c) For patients at least 6 months after completion of				
 d) For HIV positive non immune to varicella with mil e) For patients with inborn errors of metabolism at r 				
varicella, or	ion of major metabolio	accompci	ioution	, with the climbal filetory of
f) For household contacts of paediatric patients wh	o are immunocompron	nised, or u	ndergo	ing a procedure leading to
immune compromise where the household conta				
g) For household contacts of adult patients who have immunocompromised, or undergoing a procedure has no clinical history of varicella.				
* immunosuppression due to steroid or other immunosuppre	ssive therapy must be	for a treat	tment p	eriod of greater than
28 days				.
Inj 1350 PFU prefilled syringe	0.00	1	_	<u>arivax</u>
		10	✓ V	arivax
VARICELLA ZOSTER VACCINE [SHINGLES VACCINE] - [Xph	narm]			
Funded for patients meeting the following criteria:				
1) Two doses for all people aged 65 years				
Inj 50 mcg per 0.5 ml vial plus vial	0.00	1	√ S	hingrix
VARICELLA ZOSTER VIRUS (OKA STRAIN) LIVE ATTENUATI	ED VACCINE [SHINGI	LES VACO	CINE) -	- [Xpharm]
Funded for patients meeting the following criteria:	•		•	
1) One dose for all people aged 65 years				
Inj 19,400 PFU prefilled syringe plus vial	0.00	1		ostavax
		10	✓ Z	ostavax
Diagnostic Agents				
TUBERCULIN PPD [MANTOUX] TEST - [Xpharm]				
Inj 5 TU per 0.1 ml, 1 ml vial	0.00	1	√ T	ubersol
, 5 . 5 ps. 5.1 m, 1 m man		•		

- Symbols -	Poisonings	246	Amsacrine	149
3TC10	3,		AmsaLyo	
7 MED NSHA Silver/Copper	Albendazole	90	Amsidine	149
Short	74 Albey	232	Amzoate	31
- A -	Albustix	78	Anaesthetics	119
A-Scabies	68 Alchemy	158	Anagrelide hydrochloride	149
Abacavir sulphate1	O6 Alchemy Oxybutynin	<mark>77</mark>	Analgesics	120
Abacavir sulphate with	Aldurazyme	30	Anastrozole	167
lamivudine1	06 Alecensa	157	Anatrole	
Abiraterone acetate1	64 Alectinib	157	Andriol Testocaps	82
Acarbose	11 Alendronate sodium	111	Androderm	81
Accarb	11 Alendronate sodium with		Anoro Ellipta	236
Accuretic 10	49 colecalciferol	111	Antabuse	142
Accuretic 20	49 Alfacalcidol	34	Antacids and Antiflatulents	6
Acetazolamide2	43 Alfamino	267	Anthelmintics	
Acetec	48 Alfamino Junior	267	Antiacne Preparations	
Acetic acid with hydroxyquinoline and	Alginic acid	6	Antiallergy Preparations	
ricinoleic acid	76 Alglucosidase alfa	27	Antianaemics	
Acetylcysteine24			Antiandrogen Oral	
Aci-Jel		146	Contraceptives	76
Aciclovir	Allerfix	243	Antiarrhythmics	
Infection1			Antibacterials	
Sensory24			Antibacterials Topical	62
Acidex			Anticholinergic Agents	
Acipimox			Anticholinesterases	
Acitretin	•		Antidepressants	
Aclasta1	•		Antidiarrhoeals	
Actemra2	•		Antiepilepsy Drugs	
Actemra S292	•		Antifibrinolytics, Haemostatics and	
Actinomycin D1			Local Sclerosants	
Actrapid			Antifibrotics	
Actrapid Penfill			Antifungals	
Acupan1	•		Antifungals Topical	
Adalimumab (Amgevita)1			Antihistamines	
Adalimumab (Humira - Alternative	Alvogen		Antihypotensives	
brand) 1	ŭ		Antimalarials	
Adapalene	•		Antimigraine Preparations	
ADR Cartridge 1.8			Antinausea and Vertigo Agents	
Adrenaline			Antipruritic Preparations	
Advantan	•		Antipsychotics	
Advate	•		Antiretrovirals	
Adynovate		54	Antirheumatoid Agents	
Afinitor2			Antispasmodics and Other Agents	
Aflibercept1	•	54	Altering Gut Motility	
Afluria Quad	Aminophylline		Antithrombotic Agents	
(2022 formulation) 2			Antithymocyte globulin	
Afluria Quad Junior	Amisulpride		(equine)	174
(2022 formulation)			Antitrichomonal Agents	
AFT-Pyrazinamide1			Antituberculotics and	
Agents Affecting the	Amneal		Antileprotics	100
Renin-Angiotensin System			Antiulcerants	
Agents for Parkinsonism and Related	Amoxicillin		Antivirals	
Disorders 1			Anxiolytics	
Agents Used in the Treatment of	Amphotericin B		Anzatax	
•	1			

INDEX: Generic Chemicals and Brands

Apidra	11	Atorvastatin	55	Beta Ointment	64
Apidra SoloStar	11	Atropine sulphate		Beta Scalp	70
APO-Atomoxetine	137	Cardiovascular	50	Beta-Adrenoceptor Agonists	235
APO-Atomoxetine S29	137	Sensory	244	Beta-Adrenoceptor Blockers	51
Apo-Azithromycin	91	Atropt	244	Betadine	67
Apo-Diltiazem CD	53	Atrovent	235	Betadine Skin Prep	67
Apo-Prednisone	81	AU Synacthen	81	Betaferon	135
Apo-Primidone		Aubagio	135	Betahistine dihydrochloride	129
Apo-Temozolomide		Augmentin		Betaine	
Apomorphine hydrochloride		Aurorix		Betaloc CR	51
Aprepitant		AutoSoft 30	24	Betamethasone dipropionate	64
Apresoline		AutoSoft 90	24	Betamethasone dipropionate with	
Aptamil AllerPro SYNEO 1	270	Avallon	121	calcipotriol	69
Aptamil AllerPro SYNEO 2		Avelox	95	Betamethasone sodium phosphate	
Aptamil Gold+ Pepti Junior		Avonex	135	with betamethasone acetate	
Aqueous cream		Avonex Pen		Betamethasone valerate6	
Aratac		Azacitidine		Betamethasone valerate with sodiu	
Arava		Azacitidine Dr Reddy's		fusidate [fusidic acid]	
Arginine		Azamun		Betaxolol	
Aripiprazole		Azathioprine		Betnovate	
Aripiprazole Sandoz		Azilect		Betoptic	
Aristocort		Azithromycin		Betoptic S	
Arrotex-Prazosin S29		Azopt		Bexsero	
Arrow - Lattim		AZT		Bezafibrate	
Arrow-Amitriptyline		-B-	100	Bezalip	
Arrow-Bendrofluazide		B-D Micro-Fine	16	Bezalip Retard	
Arrow-Brimonidine		B-D Ultra Fine		Bicalutamide	
Arrow-Diazepam		B-D Ultra Fine II		Bicillin LA	
•			10		
Arrow-Doxorubicin Arrow-Losartan &	130	Bacillus Calmette-Guerin (BCG)	174	BiCNU	
	40	vaccine	174	Bile and Liver Therapy	
Hydrochlorothiazide		Bacillus Calmette-Guerin	070	Biltricide	
Arrow-Norfloxacin		vaccine		Bimatoprost Multiphon	
Arrow-Ornidazole		Baclofen		Bimatoprost Multichem	
Arrow-Quinapril 10		Bactroban		Binarex	
Arrow-Quinapril 20		Balance		Binocrit	
Arrow-Quinapril 5		Barrier Creams and Emollients		Biodone	
Arrow-Roxithromycin		BCG Vaccine		Biodone Extra Forte	
Arrow-Timolol		Beclazone 100		Biodone Forte	
Arrow-Topiramate		Beclazone 250		Bisacodyl	
Arrow-Tramadol		Beclazone 50		Bisacodyl Viatris	
Arsenic trioxide		Beclomethasone dipropionate		Bisoprolol fumarate	
Asacol		Bee venom allergy treatment		Bisoprolol Mylan	
Ascorbic acid		Bendamustine hydrochloride		BK Lotion	
Aspen Adrenaline	57	Bendrofluazide	55	Bleomycin sulphate	150
Aspirin		Bendroflumethiazide		Blood Colony-stimulating	
Blood		[Bendrofluazide]		Factors	45
Nervous		Benralizumab		Blood glucose diagnostic test	
Asthalin		Benzathine benzylpenicillin	93	meter	15
Atazanavir sulphate	106	Benzatropine mesylate		Blood glucose diagnostic test	
Atenolol	51	Benzbromaron AL 100		strip	15
Atenolol AFT		Benzbromarone		Blood glucose test strips (visually	
Atenolol AFT S29	51	Benztrop		impaired)	15
ATGAM	174	Benzydamine hydrochloride	33	Blood Ketone Diagnostic Test	
Ativan	134	Benzylpenicillin sodium [Penicillin		Strip	14
Atnahs Olsalazine	8	G]	93	Bonjela	
Atomoxetine	137	Beta Cream		Boostrix	

Bortezomib150	Candestar	49	Chlorafast	24
Bortezomib Dr-Reddy's150	Canesten	63	Chlorambucil	14
Bosentan58	Capecitabine	147	Chloramphenicol	24
Bosentan Dr Reddy's58	Capercit		Chlorothiazide	
Bplex34	Capoten		Chlorpheniramine maleate	
Breo Ellipta234	Capsaicin		Chlorpromazine hydrochloride	
Brevinor 1/2875	Musculoskeletal	111	Chlortalidone [Chlorthalidone]	
Bricanyl Turbuhaler235	Nervous	120	Chlorthalidone	
Brilinta42	Captopril	48	Chlorvescent	
Brimonidine tartrate244	Carafate		Choice 380 7med Nsha Silver/co	pper
Brimonidine tartrate with timolol	Carbaccord		Short	
maleate244	Carbamazepine		Choice Load 375	
Brinzolamide243	Carbimazole		Choice TT380 Short	74
Brolene241	Carbomer		Choice TT380 Standard	
Brufen SR110	Carboplatin		Choline salicylate with cetalkoniu	
BSF Amgevita246	Carboplatin Ebewe		chloride	
BSF Aspen246	Carbosorb-X		Ciclosporin	
Buccastem130	Cardinol LA		Cilazapril	
Budesonide	Cardizem CD		Cilicaine	
Alimentary6	CareSens Dual		Cilicaine VK	
Respiratory233, 239	CareSens N		Cinacalcet	
Budesonide with eformoterol234	CareSens N POP		Cipflox	
Bumetanide53	CareSens N Premier		Ciprofloxacin	
Buprenorphine Naloxone BNM141	CareSens PRO		Infection	9,
Buprenorphine with naloxone141	Carmellose sodium with gelati		Sensory	
Bupropion hydrochloride142	pectin		Ciprofloxacin Teva	
Burinex53	Carmustine		Cisplatin	
Burinex S29	Carnitor		Cisplatin Ebewe	
Buscopan8	Carvedilol		Citalopram hydrobromide	
Buspirone hydrochloride134	Carvedilol Sandoz		Cladribine	
Buspirone Viatris	Casirivimab and imdevimab		Clarithromycin	17
Busulfan145	Catapres		Alimentary	
- C -	CeeNU		Infection	
Cabergoline89	Cefaclor monohydrate		Clexane	
Caffeine citrate240	Cefalexin		Clexane Forte	
Calamine 64	Cefalexin Sandoz		Climara	
Calamine-AFT	Cefazolin		Clindamycin	
Calci-Tab 500	Ceftriaxone		Clinicians	
Calcipotriol 69	Ceftriaxone-AFT		Clinicians Renal Vit	
Calcitonin	Cefuroxime axetil		Clobazam	
Calcitriol 34	Celapram		Clobetasol propionate	
Calcitriol-AFT34	Celebrex		Clobetasone butyrate	
Calcium 500 mg Hexal	Celecoxib		Clofazimine	
Calcium carbonate	Celecoxib Pfizer		Clomazol	
Calcium Channel Blockers52	Celestone Chronodose		Dermatological	6
Calcium Disodium Versenate247	Cellcept		Genito-Urinary	
Calcium folinate147	Centrally-Acting Agents		Clomifene citrate	
Calcium Folinate Ebewe147			Clomipramine hydrochloride	
Calcium Folinate Ebewe147	Cephalexin ABM Cetirizine hydrochloride		Clomipramine Teva	
Calcium Folinate Sandoz S29 147	Cetomacrogol		Clonazepam1	
			Clonidine	20, 13
Calcium gluconate	Cetomacrogol with glycerol		Clonidine BNM	
	Cetomacrogol-AFT Cetuximab			
Calcium polystyrene sulphonate47 Calcium Resonium47	Charcoal		Clonidine hydrochloride	
Calogen253	Chemotherapeutic Agents		Clopidogrel	
Calogen253 Candesartan cilexetil49	Chickenpox vaccine		Clopidogrel Multichem	
Canuesanan Chexelli49	OHICKEHPOX VACCINE		Olopidogref Widilichefff	4

INDEX: Generic Chemicals and Brands

Clopine	131	Curam	93	DBL Leucovorin Calcium	14
Clopixol13	32-133	Curam Duo 500/125	93	DBL Methotrexate Onco-Vial	14
Clotrimazole		Cvite	34	DBL Morphine Sulphate	12
Dermatological	63	Cyclizine hydrochloride	130	DBL Naloxone Hydrochloride	
Genito-Urinary		Cyclizine lactate		DBL Pethidine Hydrochloride	
Clozapine		Cyclogyl		DBL Vincristine Sulfate	
Clozaril		Cyclonex		Decozol	
Co-trimoxazole		Cyclopentolate hydrochloride		Deferasirox	
Coal tar		Cyclophosphamide		Deferiprone	
Coal tar with allantoin, menthol,		Cyclorin		Deferoxamine Pfizer S29	
phenol and sulphur	69	Cycloserine		Denosumab	
Coal tar with salicylic acid and		Cyproterone acetate		Deolate	
sulphur	69	Cyproterone acetate with		Deoxycoformycin	
Coco-Scalp		ethinyloestradiol	76	Depo-Medrol	
Codeine phosphate	03	Cystadane		Depo-Provera	
	240	Cytarabine			
Extemporaneous				Depo-Testosterone	
Nervous		Cytotec		Deprim	
Coenzyme Q10		Cytoxan	145	Dermol	
Colonaloiforal		-	444	Desferrioxamine mesilate	
Colecalciferol		D-Penamine		Desmopressin	
Colestid		Dabigatran		Desmopressin acetate	
Colestipol hydrochloride		Dacarbazine		Desmopressin-PH&T	
Colgout		Dacarbazine APP		Desuric	11
Colifoam		Dactinomycin [Actinomycin D].		Detection of Substances in	_
Colistin sulphomethate		Daivobet		Urine	7
Colistin-Link		Daivonex		Dexamethasone	
Collodion flexible		Daktarin		Hormone	
Colloidal bismuth subcitrate		Dalacin C		Sensory	
Colofac		Dantrium		Dexamethasone phosphate	8
Coloxyl		Dantrium S29		Dexamethasone Phosphate	
Combigan		Dantrolene	117	Panpharma	
Compound electrolytes		Daonil	11	Dexamethasone with framycetin a	and
Compound electrolytes with gluco	se	Dapa-Tabs	55	gramicidin	24
[Dextrose]	47	Dapsone		Dexamethasone with neomycin	
Compound hydroxybenzoate	249	Daraprim	96	sulphate and polymyxin B	
Comtan	118	Darunavir	106	sulphate	24
Concerta	139	Darunavir Mylan	106	Dexamfetamine sulfate	13
Condoms	73	Dasatinib	157	Dexmethsone	8
Condyline	71	Daunorubicin	150	Dextrochlorpheniramine	
Contraceptives - Hormonal	74	Daunorubicin Zentiva	150	maleate	23
Contraceptives - Non-hormonal	72	David One Step Cassette Preg	nancy	Dextrose	. 46–4
Copaxone		Test	77	DHC Continus	
Corticosteroids and Related Agen		DBL Acetylcysteine	246	Diabetes	1
for Systemic Use		DBL Adrenaline		Diabetes Management	
Corticosteroids Topical		DBL Aminophylline		Diacomit	12
Cortifoam		DBL Bleomycin Sulfate		Diagnostic Agents	
Cosentyx		DBL Carboplatin		Diamide Relief	
Cosmegen		DBL Cisplatin		Diamox	
Coumadin		DBL Dacarbazine		Diasip	
Country Life		DBL Desferrioxamine Mesylate		Diason RTH	
Coversyl		BP	•	Diazepam1	
Creon 10000	25	DBL Docetaxel		Diazoxide	
Creon 25000		DBL Ergometrine		Dibenzyline	
Creon Micro		DBL Gemcitabine		Diclofenac Sandoz	11
Crotamiton		DBL Gentamicin		Diclofenac sodium	
Crystaderm		DBL Heparin Sodium		Musculoskeletal	11
or you adding		DDE Hopaili Oddiaili	T J	เขเนอบนเบอกบเซเนิเ	1 1

Sensory	242	Dr Reddy's Omeprazole	9	Emulsifying ointment	60
Differin	62	Drofate	52	Emulsifying Ointment ADE	
Difflam	33	Drugs Affecting Bone		Enalapril maleate	
Diflucan	97	Metabolism	111	Enbrel	
Digestives Including Enzymes	25	Dual blood glucose and blood ket	one	Endocrine Therapy	
Digoxin	50	diagnostic test meter	14	Endoxan	14
Dihydrocodeine tartrate		Dulaglutide		Engerix-B	27
Dilantin		Dulcolax SP Drop		Enlafax XR	12
Dilantin Infatab	127	Duocal Super Soluble Powder	252	Enoxaparin sodium	4
Diltiazem hydrochloride	53	Duolin		Enstilar	69
Dimethicone	65, 67	Duolin HFA	235	Ensure	
Dimethyl fumarate	135	DuoResp Spiromax	234	Ensure Plus	26
Dipentum	8	Duride	57	Ensure Plus HN	26
Diphtheria, tetanus and pertuss	sis	Durvalumab	225	Ensure Plus RTH	26
vaccine		-E-		Ensure Two Cal HN RTH	26
Diphtheria, tetanus, pertussis a	ind	e-chamber La Grande	239	Entacapone	118
polio vaccine	274	e-chamber Mask	239	Entecavir	10
Diphtheria, tetanus, pertussis, p		e-chamber Turbo	239	Entecavir Sandoz	
hepatitis B and haemophilus		E-Mycin		Entocort CIR	
influenzae type B vaccine		e5 Pharma		Entresto 24/26	
Diprosone		Ear Preparations		Entresto 49/51	49
Diprosone OV		Ear/Eye Preparations		Entresto 97/103	
Dipyridamole		Easiphen Liquid		Epilim	12
Disopyramide phosphate		Econazole nitrate		Epilim Crushable	12
Disulfiram		Efavirenz		Epilim IV	
Diuretics	53	Efavirenz with emtricitabine and		Epilim S/F Liquid	12
Dizole	97	tenofovir disoproxil	106	Epilim Syrup	
Docetaxel		Eformoterol fumarate		Epirubicin Ebewe	
Docetaxel Accord	150	Eformoterol fumarate dihydrate	234	Epirubicin hydrochloride	
Docetaxel Sandoz	150	Eftrenonacog alfa [Recombinant		Eplerenone	
Docusate sodium		factor IX]	39	Epoetin alfa	3
Docusate sodium with		Efudix		Epoprostenol	6
sennosides	26	Egopsoryl TA	69	Eptacog alfa [Recombinant factor	
Dolutegravir		Elaprase		VIIa]	4
Domperidone		Eldepryl		Erbitux	
Donepezil hydrochloride		Elecare		Ergometrine maleate	
Donepezil-Rex		Elecare LCP		Erlotinib	
Dornase alfa		Electral		Erythrocin IV	
Dortimopt		Elelyso		Erythromycin (as lactobionate)	
Dorzolamide hydrochloride		Elemental 028 Extra		Erythromycin ethyl succinate	
Dorzolamide with timolol		Elidel		Esbriet	
Dostinex		Elocon		Escitalopram	
Dosulepin [Dothiepin]		Elocon Alcohol Free		Escitalopram (Ethics)	
hydrochloride	125	Eltrombopag		Eskazole	
Dosulepin Mylan	125	Eltroxin		Essential Ethosuximide	
Dosulepin Viatris		EMB Fatol		Essential Generics	
Dothiepin		Emend Tri-Pack	129	Essential Prednisolone	
Doxazosin		Emicizumab	40	Estraderm MX	
Doxazosin Clinect		EMLA		Estradiol TDP Mylan	
Doxine		Empagliflozin		Estradot	
Doxorubicin Ebewe		Empagliflozin with metformin		Estradot 50 mcg	8
Doxorubicin hydrochloride		hydrochloride	13	Estrofem	
Doxycycline		Emtricitabine		Etanercept	16
DP Lotion		Emtricitabine with tenofovir		Ethambutol hydrochloride	
DP Lotn HC		disoproxil	103	Ethics Aspirin	
DP-Allopurinol		Emtriva		Ethics Aspirin EC	
•				•	

INDEX: Generic Chemicals and Brands

Ethics Lisinopril	48	Ferrous fumarate	36	Folic Acid Mylan	39
Ethinyloestradiol	83	Ferrous fumarate with folic acid	36	Food Thickeners	263
Ethinyloestradiol with		Ferrous sulfate	36	Foods And Supplements For Ir	nborn
desogestrel	74	Fexofenadine hydrochloride	233	Errors Of Metabolism	265
Ethinyloestradiol with		Fibro-vein	42	Foradil	234
levonorgestrel	75	Filgrastim	45	Forteo	113
Ethinyloestradiol with		Finasteride	77	Fortini	256
norethisterone	75	Fingolimod	135	Fortini Multi Fibre	256
Ethosuximide	126	Firazyr	232	Fortisip	262
Etopophos	151	Flagyl	99	Fortisip Multi Fibre	262
Etoposide	151	Flagyl-S	99	Fosamax	111
Etoposide phosphate	151	Flamazine	63	Fosamax Plus	111
Etravirine	106	Flecainide acetate	<mark>50</mark>	Framycetin sulphate	241
Eumovate	64	Flecainide BNM	50	Frisium	
Evara	66	Flecainide Controlled Release		Frumil	54
Everet	127	Teva	50	Frusemide	54
Everolimus	228	Fleet Phosphate Enema	<mark>27</mark>	Fucicort	65
Evista	112	Flixonase Hayfever & Allergy	239	Fucidin	96
Evusheld	219	Flixotide		Fucithalmic	241
Exemestane	167	Flixotide Accuhaler	234	Fulvestrant	164
Exjade	246	Florinef	80	Fungilin	33
Extemporaneously Compound	ed	Fluanxol	132	Furosemid-Ratiopharm	54
Preparations and		Flucil		Furosemide [Frusemide]	
Galenicals	249	Flucloxacillin	93	Furosemide-Baxter	
Eye Preparations	241	Flucloxacillin-AFT	93	fusidic acid	
Eylea	190	Flucloxin	93	Dermatological	63, 65
Ezetimibe		Flucon	242	Infection	
Ezetimibe Sandoz	56	Fluconazole	97	Sensory	241
LZCIIIIDC OdiidOZ		i idooriazoio		Oct 1301 y	
Ezetimibe with simvastatin		Fludara Oral		- G -	
			147	•	
Ezetimibe with simvastatin	57	Fludara Oral	147 147	- G -	127
Ezetimibe with simvastatin	57	Fludara OralFludarabine Ebewe	147 147 147	- G - Gabapentin	127 122
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing	57 g 41	Fludara Oral Fludarabine Ebewe Fludarabine phosphate	147 147 147 80	- G - Gabapentin	127 122 29
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction	57 g 41 9	Fludara Oral Fludarabine Ebewe Fludarabine phosphate Fludrocortisone acetate	147 147 147 80 46	- G - Gabapentin	127 122 29
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine	57 9 41 9	Fludara Oral Fludarabine Ebewe Fludarabine phosphate Fludrocortisone acetate Fluids and Electrolytes	147 147 147 80 46	Gabapentin Gacet Galsulfase Galvumet Galvumet	127 122 29 12
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid	57 9 41 9 9	Fludara Oral	147 147 147 80 46	Gabapentin	127 29 12 12
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex	57 g41999	Fludara Oral	147 147 80 46 241	Gabapentin	127 29 12 12 12 275
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra	57 9 41 9 9 191 164 66	Fludara Oral	147 147 147 80 46 241	Gabapentin	127 29 12 12 12 275 9
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex Fatty Cream AFT	57 99919116466116	Fludara Oral	147 147 80 46 241	Gabapentin	127 29 12 12 275 9 6
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex Fatty Cream AFT Febuxostat.	57 99919116466116	Fludara Oral	147 147 80 46 241 8 8 242	Gabapentin	127 122 12 12 275 9 6 6
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex Fatty Cream AFT Febuxostat Febuxostat multichem	57 9419919116466116	Fludara Oral	147 147 80 46 241 8 8 242 148	Gabapentin	127 122 12 12 12 275 9 6 6 201 158
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex Fatty Cream AFT Febuxostat Febuxostat multichem Feed Thickener Karicare	57 9419919116466116116264	Fludara Oral	14714714780462418242148	Gabapentin	127 29 12 12 12
Ezetimibe with simvastatin F - Factor eight inhibitor bypassing fraction Famotidine Famotidine Hovid Fasenra Faslodex Fatty Cream AFT Febuxostat Febuxostat multichem Feed Thickener Karicare Aptamil	57 941991916611611626441	Fludara Oral	1471471478046241824214814871125	Gabapentin	127 29 12 275 9 66 66 201 158 66 66
Ezetimibe with simvastatin	57 g419919116466116116264415252	Fludara Oral	147147804624188242148125125	Gabapentin	127 29 12 29 275 66 66 2011 158 666 148 148
Ezetimibe with simvastatin		Fludara Oral	147147804624188242148125125132	Gabapentin	127 29 25 275 66 66 201 158 666 148 148 148 192
Ezetimibe with simvastatin		Fludara Oral	14714780462418 8242148125125132164	Gabapentin	127 29 25 275 66 66 201 158 666 148 148 148 192
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127291229122756666201128666148148148192241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	12729122912275996660158666148148192241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127291229122756666666148148148192241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	12729122912
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127292912275966662011586614814819224124195241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127292912275966662011586614814819224124195241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127292912275966666615866148148148192241241
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	12729292759666620115866148148192241952411357676
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127292927596666201158661481481922419524113576
Ezetimibe with simvastatin		Fludara Oral		Gabapentin	127291229275

Glizide11	Holoxan145	Ibuprofen	110
Glucagen Hypokit10	Horleys Bread Mix264	lcatibant	
Glucagon hydrochloride10	Horleys Flour264	Idarubicin hydrochloride	
Glucerna Select254	Hormone Replacement Therapy -	Idursulfase	
Glucose [Dextrose]46	Systemic 82	Ifosfamide	
Gluten Free Foods264	HPV275	Igroton	5
Glycerin with sodium saccharin249	Humalog11	Ikorel	
Glycerin with sucrose249	Humalog Mix 2511	Iloprost	
Glycerol	Humalog Mix 5011	Imatinib mesilate	
Alimentary27	Human papillomavirus (6, 11, 16, 18,	Imatinib-Rex	15
Extemporaneous249	31, 33, 45, 52 and 58) vaccine	Imfinzi	22
Glyceryl trinitrate	[HPV]275	Imigran	129
Alimentary8	Humatin96	Imipramine hydrochloride	12
Cardiovascular57	Humira184	Imiquimod	
Glycopyrronium236	HumiraPen184	Immune Modulators	
Glycopyrronium bromide8	Humulin 30/7011	Immunosuppressants	16
Glycopyrronium with	Humulin NPH10	Imuran	16
indacaterol236	Humulin R10	Incruse Ellipta	
Go Healthy29	Hyaluronic acid245	Indacaterol	
Gold Knight73	Hydralazine58	Indapamide	
Gold Knight XL73–74	Hydralazine hydrochloride58	Infanrix IPV	
Goserelin	Hydrocortisone	Infanrix-hexa	
Gutron50	Dermatological64	Infant Formulae	26
Gynaecological Anti-infectives76	Hormone80	Infatrini	27
´ -H-	Hydrocortisone (PSM)64	Infliximab	19
Habitrol142	Hydrocortisone acetate7	Influenza vaccine	27
Haemophilus influenzae type B	Hydrocortisone acetate with	Inhaled Corticosteroids	23
vaccine274	pramoxine hydrochloride	Inhaled Long-acting	
Haldol132	Hydrocortisone and paraffin liquid	Beta-adrenoceptor Agonists	23
Haldol Concentrate132	and lanolin64	Inspra	
Haldol Decanoas132	Hydrocortisone butyrate65, 70	Instillagel Lido	
Haloperidol131	Hydrocortisone with cinchocaine8	Insulin aspart	
Haloperidol decanoate132	Hydrocortisone with miconazole65	Insulin aspart with insulin aspart	
Harvoni103	Hydrocortisone with natamycin and	protamine	10
Havrix274	neomycin 65	Insulin glargine	1
Havrix Junior274	Hydrogen peroxide62	Insulin glulisine	1
Haylor syrup233	Hydroxocobalamin34	Insulin isophane	
healthE Dimethicone 10%65	Hydroxocobalamin Mercury	Insulin isophane with insulin	
healthE Dimethicone 4% Lotion 67	Pharma34	neutral	1 [.]
healthE Dimethicone 5%65	Hydroxocobalamin Panpharma34	Insulin lispro	1
healthE Glycerol BP249	hydroxycarbamide151	Insulin lispro with insulin lispro	
healthE Urea Cream66	Hydroxychloroquine111	protamine	1°
Healtheries Simple Baking Mix264	Hydroxyurea	Insulin neutral	
Hemastix78	[hydroxycarbamide] 151	Insulin pen needles	10
Hemlibra40	Hygroton55	Insulin pump	10
Heparin sodium45	Hylo-Fresh245	Insulin pump cartridge	
Heparinised saline45	Hymenoptera232	Insulin pump infusion set (steel	
Heparon Junior255	Hyoscine butylbromide8	cannula)	2
Hepatitis A vaccine274	Hyoscine hydrobromide130	Insulin pump infusion set (steel	
Hepatitis B recombinant	Hypam 136	cannula, straight insertion)	2
vaccine275	Hyperuricaemia and Antigout 116	Insulin pump infusion set (teflon	
Herceptin223	Hypromellose244	cannula)	2
Hiberix274	Hypromellose with dextran244	Insulin pump infusion set (teflon	
Hiprex109	-1-	cannula, angle insertion with	
Histaclear233	Ibiamox93	insertion device)	2
Histafen233	Ibrance160	Insulin pump infusion set (teflon	

cannula, angle insertion)	24	Kaletra	107	Laxsol	2
Insulin pump infusion set (teflon		Kalydeco		Ledipasvir with sofosbuvir	10
cannula, straight insertion with		Kemadrin	119	Leflunomide	11
insertion device)	24	Kenacomb	241	Lenalidomide	15
Insulin pump infusion set (teflon		Kenacort-A 10	81	Letrole	16
cannula, straight insertion)	24	Kenacort-A 40	81	Letrozole	16
Insulin pump reservoir	24	Kenalog in Orabase	33	Leukeran FC	14
Insulin syringes, disposable with		Ketocal 3:1	272	Leukotriene Receptor	
attached needle	16	KetoCal 4:1	272	Antagonists	23
Intelence	106	Ketoconazole		Leuprorelin	8
Interferon beta-1-alpha	135	Dermatological	70	Leustatin	14
Interferon beta-1-beta	135	Infection	98	Levetiracetam	
Intra-uterine device	74	Ketogenic Diet		Levetiracetam-AFT	
Invega Sustenna	133	Ketoprofen	110	Levlen ED	
IPCA-Frusemide	54	KetoSens	14	Levocabastine	24
IPCA-Metoprolol	51	Keytruda	227	Levocarnitine	3
IPCA-Propranolol	52	Kindergen	255	Levodopa with benserazide	11
IPOL	281	Kivexa	106	Levodopa with carbidopa	11
Ipratropium bromide2	35, 239	Klacid		Levomepromazine	13
Iressa		Alimentary	9	Levomepromazine	
Irinotecan Actavis 100	148	Infection	92	hydrochloride	13
Irinotecan hydrochloride	148	Kliogest	83	Levonorgestrel	
Irinotecan-Rex	148	Kliovance	83	Genito-Urinary	75–7
Iron (as ferric carboxymaltose)	36	Kogenate FS	42	Hormone	8
Iron polymaltose		Konakion MM	42	Levothyroxine	8
Isentress	107	Konsyl-D	26	Lidocaine [Lignocaine]	119-12
Isentress HD	107	Kuvan	31	Lidocaine [Lignocaine]	
Ismo 20	57	-L-		hydrochloride	12
Ismo 40 Retard	57	Labetalol	51	Lidocaine [Lignocaine] with	
Isoniazid	100	Lacosamide	127	chlorhexidine	12
Isoniazid with rifampicin	100	Lactulose	27	Lidocaine [Lignocaine] with	
Isoptin	53	Laevolac	27	prilocaine	12
Isoptin Retard	53	Lagevrio	104	Lidocaine-Baxter	12
Isoptin SR	53	Lamictal		Lidocaine-Claris	12
Isopto Carpine		Lamivudine	101, 106	Life Extension	3
Isosorbide mononitrate		Lamivudine Alphapharm	106	Lignocaine	
Isosource Standard	261	Lamotrigine		Lioresal Intrathecal	
Isotretinoin	62	Lamprene		Lipid-Modifying Agents	5
Ispaghula (psyllium) husk	26	Lanoxin		Liquigen	
Itch-Soothe		Lanoxin Paediatric Elixir	50	Lisinopril	
Itraconazole	97	Lanoxin PG	50	Litak	
Itrazole	97	Lanoxin S29	50	Lithium carbonate	13
lvacaftor	238	Lansoprazole		Livostin	24
Ivermectin	67	Lantus		LMX4	12
- J -		Lantus SoloStar	11	Locacorten-Viaform ED's	24
Jadelle	75	Lanvis	149	Local preparations for Anal an	ıd
Jakavi	162	Lanzol Relief	9	Rectal Disorders	
Jardiamet		Lapatinib ditosylate	159	Locasol	
Jardiance		Largactil		Locoid	
Jaydess	83	Laronidase		Locoid Crelo	
Jevity HiCal RTH		Lasix		Locoid Lipocream	
Jevity Plus		Latanoprost		Locorten-Vioform	24
Jevity RTH		Latanoprost with timolol		Lodoxamide	
Juno Pemetrexed	149	Lax-Suppositories		Logem	
- K -	-	Lax-suppositories Glycerol		Lomide	
Kadcyla	224	Laxatives		Lomustine	
,					

Loniten	58	Mefenamic acid110	Metoclopramide hydrochloride	130
Loperamide hydrochloride	6	Megace165	Metolazone	54
Lopinavir with ritonavir	107	Megestrol acetate165	Metopirone	89
Lopinavir/Ritonavir Mylan	107	Melatonin135	Metoprolol IV Mylan	51
Loprofin		Melphalan146	Metoprolol succinate	51
Loprofin Mix	266	Menactra278	Metoprolol tartrate	
Lorafix	233	Meningococcal (groups A, C, Y and	Metrogyl	99
Loratadine	233	W-135) conjugate vaccine 278	Metronidazole	
Lorazepam	134	Meningococcal B multicomponent	Metyrapone	89
Lorstat	55	vaccine279	Mexiletine hydrochloride	50
Losartan Actavis	49	Meningococcal C conjugate	Miacalcic	79
Losartan potassium	49	vaccine279	Micolette	
Losartan potassium with		Menthol64	Micolette-S29	27
hydrochlorothiazide	49	Mepolizumab200	Miconazole	33
Lovir		Mercaptopurine148	Miconazole nitrate	
Loxamine	126	Mercilon 2874	Dermatological	63
Lucrin Depot 1-month	89	Mesalazine7	Genito-Urinary	
Lucrin Depot 3-month		Mesna152	Micreme	
Lyderm		Mestinon110	Micreme H	
Lynparza		Metabolic Disorder Agents27	Microgynon 20 ED	
Lyrica		Metformin hydrochloride12	Microgynon 30	
- M -		Metformin Mylan12	Microlut	
m-Eslon	123	Metformin Viatris12	Midazolam	
Mabthera	204	Methadone BNM122	Midazolam-Baxter	136
Macro Organic Psyllium Husk	26	Methadone hydrochloride	Midodrine	50
Macrobid		Extemporaneous249	Mifegyne	
Macrogol 3350 with potassium		Nervous122	Mifepristone	
chloride, sodium bicarbonate	and	Methatabs122	Minerals	
sodium chloride		Methenamine (hexamine)	Mini-Wright AFS Low Range	
Macrogol 400 and propylene		hippurate109	Mini-Wright Standard	
glycol	245	Methopt244	Minidiab	
Madopar 125	118	Methotrexate148	MiniMed 1.8 Reservoir	
Madopar 250		Methotrexate DBL Onco-Vial148	MMT-326A	24
Madopar 62.5		Methotrexate Ebewe148	MiniMed 3.0 Reservoir	
Madopar HBS		Methotrexate Sandoz148	MMT-332A	24
Madopar Rapid		Methyl hydroxybenzoate249	MiniMed 770G	
Magnesium hydroxide		Methylcellulose	MiniMed Mio MMT-921A	
Magnesium sulphate		Methylcellulose with glycerin and	MiniMed Mio MMT-923A	
Mantoux		sodium saccharin249	MiniMed Mio MMT-925A	
Marevan		Methylcellulose with glycerin and	MiniMed Mio MMT-941A	
Marine Blue Lotion SPF 50+		sucrose249	MiniMed Mio MMT-943A	
Martindale Pharma		Methyldopa53	MiniMed Mio MMT-945A	
Martindale Pharma S29		Methyldopa Mylan53	MiniMed Mio MMT-965A	
Mask for spacer device		Methyldopa Mylan S2953	MiniMed Mio MMT-975A	
Maviret		Methylnaltrexone bromide	MiniMed Quick-Set MMT-386A	
Maxidex		Methylphenidate ER - Teva138	MiniMed Quick-Set MMT-387A	
Maxitrol		Methylphenidate bydrochloride 138	MiniMed Quick-Set MMT-396A	
MCT oil (Nutricia)		Methylphenidate hydrochloride	MiniMed Quick-Set MMT-397A	
Measles, mumps and rubella	200	extended-release139	MiniMed Quick-Set MMT-398A	
vaccine	278	Methylprednisolone80	MiniMed Quick-Set MMT-399A	
Mebendazole		Methylprednisolone (as sodium	MiniMed Guick-Set MMT-399A	
Mebeverine hydrochloride		succinate) 81	MiniMed Silhouette MMT-377A	
Medrol		Methylprednisolone aceponate65	MiniMed Silhouette MMT-378A	
Medroxyprogesterone acetate	00	Methylprednisolone acetate81	MiniMed Silhouette MMT-381A	
	75	Methylxanthines238	MiniMed Silhouette MMT-382A	
Genito-Urinary Hormone		Metoclopramide Actavis 10	MiniMed Silhouette MMT-383A	
I IUIIIIUII		INICIOCIOPIAITING ACIAVIS TO 130	IVIII IIIVIEU OIII IOUELLE IVIIVI I -303A	40

MiniMed Silhouette MMT-384A	23	MycoNail	63	Nifuran	10
MiniMed Sure-T MMT-864A	21	Mycophenolate mofetil		Nilotinib	16
MiniMed Sure-T MMT-866A		Mydriacyl		Nilstat	
MiniMed Sure-T MMT-874A	21	Mylan (12 hr release)		Alimentary	3
MiniMed Sure-T MMT-876A		Mylan (24 hr release)		Genito-Urinary	
MiniMed Sure-T MMT-884A		Mylan Atenolol		Infection	
MiniMed Sure-T MMT-886A		Mylan Clomiphen		Nintedanib	
Minims Pilocarpine		Mylan Indapamide		Nipent	
Minims Prednisolone		Mylan Italy (24 hr release)		Nirmatrelvir with ritonavir	
Minirin		Myleran		Nitrates	
Minirin Melt		Mylotarg	192	Nitroderm TTS	
Mino-tabs		Myometrial and Vaginal Horm		Nitrofurantoin	
Minocycline hydrochloride		Preparations		Nitrolingual Pump Spray	
Minomycin		Myozyme		Nivestim	4
Minor Skin Infections		- N -		Nivolumab	
Minoxidil		Nadolol	51	Nizoral	
Mirena		Nadolol BNM		Nodia	
Mirtazapine		Naglazyme		Noflam 250	
Misoprostol		Nalcrom		Noflam 500	
Mitomycin C		Naloxone hydrochloride		Non-Steroidal Anti-Inflammatory	
Mitozantrone		Naltraccord		Drugs	11
Mitozantrone Ebewe		Naltrexone hydrochloride		Nonacog gamma, [Recombinant	
Mixtard 30		Naphazoline hydrochloride		Factor IX]	4
MMR II		Naphcon Forte		Norethisterone	
Moclobemide		Naprosyn SR 1000		Genito-Urinary	7
Modafinil		Naprosyn SR 750		Hormone	
Modavigil		Naproxen		Norflex	
Moduretic		Narcaricin mite		Norfloxacin	
Molaxole		Nasal Preparations		Noriday 28	
Molnupiravir		Natalizumab		Norimin	
Moments		Natulan		Normacol Plus	
Mometasone furoate		Nausafix		Normison	
Monogen		Nausicalm		Norpress	
Montelukast		Navelbine		Nortriptyline hydrochloride	
Montelukast Mylan		Nefopam hydrochloride		Norvir	
Moroctocog alfa [Recombinant		Neisvac-C		Noumed	
VIII]		Neo-B12		Noumed Paracetamol	
Morphine hydrochloride		Neo-Mercazole		NovaSource Renal	
Morphine sulphate		Neocate Gold		Novatretin	
Motetis		Neocate Junior Unflavoured		NovoMix 30 FlexPen	
Mouth and Throat		Neocate Junior Vanilla		NovoRapid	
Movapo		Neocate SYNEO		NovoRapid FlexPen	
Moxifloxacin		Neoral		NovoRapid Penfill	
MSUD Maxamum		Neostigmine metilsulfate		NovoSeven RT	
Mucilaginous laxatives with	200	Nepro HP (strawberry)		Noxafil	
stimulants	26	Nepro HP (vanilla)		Nozinan	
Mucolytics		Nepro HP RTH		Nozinan (Swiss)	
Mucosoothe		Neulactil		Nozinan S29	
Multiple Sclerosis Treatments		Neulastim		Nucala	
Multivitamin renal		Neuraxpharm		Nuelin	
Multivitamins		NeuroTabs		Nuelin-SR	
Mupirocin		Nevirapine		Nupentin	
Muscle Relaxants	117	Nevirapine Alphapharm	106	Nutilis	
Mvite		Nicorandil		Nutren Diabetes	
Myambutol		Nicotine		Nutrient Modules	20 2F
Mycobutin		Nifedipine		Nutrini Energy Multi Fibre	
1717 OODUUIT				TAGETTE ELECTORY WIGHT FIDE CO	<u>~</u> U

Nutrini Energy RTH25	6 Oncaspar LYO	153	Pamidronate disodium	112
Nutrini Low Energy Multi Fibre25			Pamisol	112
Nutrini Peptisorb26			Pancreatic enzyme	25
Nutrini Peptisorb Energy26	Ondansetron ODT-DRLA	130	Pantoprazole	9
Nutrini RTH25		34	Panzop Relief	
Nutrison 800 Complete Multi	One-Alpha S29	34	Panzytrat	
Fibre26			Papaverine hydrochloride	
Nutrison Advanced Diason25			Para-amino salicylic acid	100
Nutrison Advanced Peptisorb25	7 Ora-Blend	249	Paracare	
Nutrison Concentrated26	3 Ora-Blend SF	249	Paracare Double Strength	121
Nutrison Energy26	1 Ora-Plus	249	Paracetamol	
Nutrison Energy Multi Fibre26	1 Ora-Sweet	249	Paracetamol + Codeine	
Nutrison Multi Fibre26		249	(Relieve)	124
Nutrison Standard RTH26	1 Orabase	33	Paracetamol with codeine	
Nyefax Retard5			Paraffin	66
Nystatin	Oratane	62	Paraffin liquid with wool fat	
Alimentary3			Parasiticidal Preparations	
Genito-Urinary7			Parnate	125
Infection9			Parnate S29	
NZB Low Gluten Bread Mix26			Paromomycin	
-0-	Ortho-tolidine		Paroxetine	126
Obinutuzumab20			Paser	100
Obstetric Preparations7			Paxam	
Ocrelizumab13			Paxlovid	
Ocrevus13	- J		Pazopanib	
Octocog alfa [Recombinant factor	Other Progestogen		Peak flow meter	
VIII] (Advate)4		83	Pedialyte - Bubblegum	
Octocog alfa [Recombinant factor	Other Skin Preparations		Pediasure	
VIII] (Kogenate FS)4			Pediasure Plus	
Octreotide16		76	Pediasure RTH	
Octreotide Depot Teva16			Pegaspargase	
Octreotide GH16			Pegasys	
Octreotide long-acting16			Pegfilgrastim	
Oestradiol			Pegylated interferon alfa-2a	
Oestradiol valerate	•		Pembrolizumab	
Oestradiol with norethisterone			Pemetrexed	
Oestriol	Oxpentifylline		Penicillamine	
Genito-Urinary7			Penicillin G	
Hormone8			PenMix 30	
Oestrogens			PenMix 40	
Ofev23			PenMix 50	
Oil in water emulsion		76	Pentasa	
Olanzapine			Pentostatin [Deoxycoformycin]	
Olaparib			Pentoxifylline [Oxpentifylline]	
Olbetam5	,	76	Peptamen Junior	
Olbetam S295			Pepti-Junior	
Olopatadine24			Peptisorb	
Olopatadine Teva24	•	117	Perhexiline maleate	
Olsalazine			Pericyazine	
Omalizumab20			Perindopril	132
Omeprazole			Perjeta	1 0
Omeprazole actavis 10			Permethrin	
Omeprazole actavis 20	9 Paediatric Seravit		Perrigo	
Omeprazole actavis 40			Pertuzumab	
Omnitrope8			Peteha	
Onbrez Breezhaler23			Pethidine hydrochloride	
OIIDIOZ DIEGZIIAIGI	- I alivizuiliau	200	r outliante riyarocritoriae	124

Pevaryl	63	Poly-Visc	245	Protaphane Penfill	1
Pexsig	53	Polycal	251	Protifar	25
Pfizer Exemestane	167	Ponstan	110	Protionamide	10
Pharmacy Health Sorbolene with	1	Posaconazole	98	Provera	8
Glycerin		Postinor-1	76	Provera HD	
Pharmacy Services		Potassium chloride	46–47	PSM Citalopram	12
Pheburane	31	Potassium citrate	77	Psoriasis and Eczema	
Phenasen	149	Potassium iodate		Preparations	6
Phenobarbitone		Povidone iodine	67	PTU	
Phenobarbitone sodium		Pradaxa	45	Pulmicort Turbuhaler	
Extemporaneous	249	Pramipexole hydrochloride	118	Pulmozyme	23
Nervous		Pravastatin	55	Puri-nethol	
Phenoxybenzamine		Pravastatin Mylan		Puria	
hydrochloride	48	Praziquantel		Pyrazinamide	10
Phenoxymethylpenicillin (Penicill		Prazosin		Pyridostigmine bromide	
V)		Pred Forte		Pyridoxine hydrochloride	
Phenytoin sodium1		Prednisolone		Pyridoxine multichem	
Phillips Milk of Magnesia		Prednisolone acetate		Pyrimethamine	
Phlexy 10		Prednisolone sodium		Pytazen SR	4
Phosphate Phebra		Prednisolone sodium		- Q -	···········
Phosphorus		phosphate	243	Quetapel	13
Phytomenadione		Prednisolone-AFT		Quetiapine	
Pilocarpine hydrochloride		Prednisone		Quick-Set MMT-392	
Pimafucort		Prednisone Clinect		Quick-Set MMT-393	
Pimecrolimus				Quinapril	
		Pregabalin			4
Pine tar with trolamine laurilsulfa		Pregabalin Pfizer		Quinapril with	4
and fluorescein		Pregnancy Tests - hCG Urine		hydrochlorothiazide Qvar	
		Premarin			20
Pioglitazone	12	Prevenar 13	280	- R -	
Pioglitazone	12 237	Prevenar 13 Priadel	280 131	- R - RA-Morph	12
Pioglitazone	12 237 129	Prevenar 13 Priadel Primaquine	280 131 99	- R - RA-Morph Ralicrom	12
Pioglitazone	12 237 129 266	Prevenar 13 Priadel Primaquine Primidone	280 131 99 128	- R - RA-MorphRalicromRaloxifene hydrochloride	12
Pioglitazone	12 237 129 266	Prevenar 13 Priadel Primaquine Primidone Primidone Clinect	280 131 99 128	RA-MorphRalicromRaloxifene hydrochlorideRaltegravir potassium	12
Pioglitazone	12 237 129 266 266	Prevenar 13 Priadel Primaquine Primidone Primidone Clinect Primolut N	280 131 99 128 128	RA-Morph	12 11 10
Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ	12 237 266 266 266 266	Prevenar 13 Priadel Primaquine Primidone Primidone Clinect Primolut N Priorix	280 131 99 128 128 83 278	RA-Morph	12
Pioglitazone	12 237 266 266 266 266 266	Prevenar 13 Priadel Primaquine Primidone Primolut N Priorix Probenecid	280 131 99 128 128 83 278 117	RA-Morph	12
Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ PKU Anamix Junior Orange PKU Anamix Junior Vanilla	12 237 266 266 266 266 266 266	Prevenar 13 Priadel Primaquine Primidone Clinect Primolut N Priorix Probenecid Probenecid	2801319912812883278117	RA-Morph	12 11; 10 11; 9
Pioglitazone Pirfenidone Pizotifen PKU Anamix Infant PKU Anamix Junior PKU Anamix Junior Chocolate PKU Anamix Junior LQ PKU Anamix Junior Orange PKU Anamix Junior Vanilla PKU Lophlex LQ 10	12 237 266 266 266 266 266 266 266	Prevenar 13	2801319912812883278117117	RA-Morph	12 11 10 11 9 9
Pioglitazone	12 237 266 266 266 266 266 266 266	Prevenar 13		RA-Morph	12 11; 10 11; 9 9 22; 11; 8
Pioglitazone	12 237 266 266 266 266 266 266 266 266 266	Prevenar 13		RA-Morph	12 10 11 9 9 9 22 11 8
Pioglitazone	12237266266266266266266266266266266266266266266266266266266266	Prevenar 13		RA-Morph	12 11: 10 9 99 22 11: 8 39, 4
Pioglitazone	12237129266266266266266266266266266266266266266266266	Prevenar 13		RA-Morph	12 11 10 9 9 9
Pioglitazone	12237129266266266266266266266266266266266266266266266	Prevenar 13		RA-Morph	12 11 10 9 9 9
Pioglitazone	12237129266266266266266266266266266266266266266266266266	Prevenar 13		RA-Morph	12 11 10 9 9 9
Pioglitazone	12237129266266266266266266266266266266266266266266266266	Prevenar 13		RA-Morph	12: 11: 10: 99: 99: 8:
Pioglitazone	12237129266266266266266266266266266266266266266266266266266266279	Prevenar 13		RA-Morph	12: 11: 10: 99: 99: 8:
Pioglitazone Pirfenidone Pizotifen	12237129266266266266266266266266266266266266266266266266269279	Prevenar 13		RA-Morph	12:
Pioglitazone	12237129266266266266266266266266266266266266266266266266269279	Prevenar 13		RA-Morph	
Pioglitazone Pirfenidone Pizotifen	12237129266	Prevenar 13		RA-Morph	12
Pioglitazone Pirfenidone Pizotifen	12237129266268281	Prevenar 13		RA-Morph	12
Pioglitazone Pirfenidone Pizotifen	12237129266266266266266266266266266266266266269260260260260260260260281	Prevenar 13		RA-Morph	12
Pioglitazone	12237129266266266266266266266266266266266266269260260260260260260280279	Prevenar 13		RA-Morph	12
Pioglitazone	1212237129266266266266266266266	Prevenar 13		RA-Morph	12
Pioglitazone	12237129266266266266266266266266266266266269279 te279 te280281	Prevenar 13		RA-Morph	12
Pioglitazone	12237129266266266266266266266266266266266266	Prevenar 13		RA-Morph	12
Pioglitazone	1212237129266266266266266266266	Prevenar 13		RA-Morph	12

Paylimid	151	-S-		Codium honzooto	2.
Revlimid		-	100	Sodium benzoate	3
Revolade		Sabril		Sodium bicarbonate	46.4
Rexacrom		Sacubitril with valsartan		Blood	
Riboflavin		Sagent		Extemporaneous	
Ribomustin		SalAir		Sodium calcium edetate	24
Ricit		Salazopyrin		Sodium chloride	
Rifabutin		Salazopyrin EN		Blood	
Rifadin		Salbutamol	235	Respiratory	
Rifampicin		Salbutamol with ipratropium		Sodium citrate with sodium laury	
Rifaximin		bromide		sulphoacetate	
Rifinah		Salicylic acid		Sodium citro-tartrate	7
Rilutek		Salmeterol		Sodium cromoglicate	
Riluzole		Sandomigran		Alimentary	
RINVOQ		Sanofi Primaquine		Sensory	
Riodine	67	Sapropterin dihydrochloride		Sodium fluoride	3
Risedronate Sandoz	113	Scalp Preparations		Sodium Fusidate [fusidic acid]	
Risedronate sodium	113	Scopoderm TTS	130	Dermatological	6
Risperdal Consta	133	Sebizole	70	Infection	9
Risperidone		Secukinumab	217	Sensory	24
Risperidone (Teva)	132	Sedatives and Hypnotics	135	Sodium hyaluronate [Hyaluronic	
Risperon		Seebri Breezhaler	236	acid]	24
Ritalin	138	Selegiline hydrochloride	118	Sodium phenylbutyrate	3
Ritalin LA	139	Senna	27	Sodium picosulfate	
Ritonavir	107	Senokot	27	Sodium polystyrene sulphonate	4
Rituximab (Mabthera)	204	SensoCard	15	Sodium tetradecyl sulphate	4
Rituximab (Riximyo)		Serc	129	Sodium valproate	
Rivaroxaban	45	Serenace		Sofradex	
Rivastigmine		Seretide		Soframycin	
Rivastigmine Patch BNM 10		Seretide Accuhaler	235	Solgar28	
Rivastigmine Patch BNM 5	140	Serevent	234	Solifenacin Mylan	
Rivotril		Serevent Accuhaler	234	Solifenacin succinate	
Riximyo		Sertraline	126	Solu-Cortef	
RIXUBIS		Setrona	126	Solu-Medrol	
Rizamelt		Setrona AU		Solu-Medrol-Act-O-Vial	
Rizatriptan		Sevredol		Somatropin (Omnitrope)	
RoActemra S29		Sex Hormones Non		Sotalol	
Ronapreve		Contraceptive	81	Spacer device	
Ropin		Shield XL		Span-K	
Ropinirole hydrochloride		Shingles vaccine		Spiolto Respimat	23
Rosuvastatin		Shingrix		Spiractin	
Rosuvastatin Viatris		SII-Onco-BCG		Spiriva	23
Rotarix		Sildenafil		Spiriva Respimat	
Rotavirus oral vaccine		Silhouette MMT-373		Spironolactone	
Roxane		Siltuximab		Sporanox	
Roxane-Propranolol		Simvastatin		Sprycel	15
Roxithromycin	92	Simvastatin Mylan		Stemetil	130
Rubifen		Sinemet		Steril-Gene	
Rubifen SR		Sinemet CR		SteroClear	
Rugby Capsaicin Topical		Sirolimus		Stesolid	
Cream	120	Siterone		Stimulants/ADHD Treatments	
Rulide D		Slow-Lopresor		Stiripentol	
Rurioctocog alfa pegol [Reco		Smith BioMed Rapid Pregnancy		Stocrin	
factor VIII]		Test		Stomahesive	
Ruxolitinib		Sodibic		Strattera	
Rythmodan		Sodium acid phosphate		Strides Shasun	
,				Stromectol	
Rytmonorm		Sodium alginate	0	Stromettor	0

Sucralfate	10	Telfast	233	Muscular Pain	11
Sulfadiazine Silver	63	Teligent	95	Topiramate	12
Sulfadiazine sodium	96	Temaccord	154	Topiramate Actavis	12
Sulfasalazine	8	Temazepam	136	Total parenteral nutrition (TPN)	
Sulphur		Temozolomide		TPN	
Sulprix		Tenofovir disoproxil		Tramadol hydrochloride	
Sumagran		Tenofovir Disoproxil Emtricitabin		Tramal SR 100	
Sumatriptan		Mylan		Tramal SR 150	
Sunitinib		Tenofovir Disoproxil Mylan		Tramal SR 200	
Sunitinib Pfizer		Tenofovir Disoproxil Teva		Trandate	
Sunscreens		Tenoxicam		Tranexamic acid	
Sunscreens, proprietary		Tensipine MR10		Tranylcypromine sulphate	
		•			
Sure-T MMT-863		Tepadina		Trastuzumab	
Sure-T MMT-873		Terbinafine		Trastuzumab emtansine	
Sustagen Hospital Formula	261	Terbutaline sulphate		Travatan	
Sustagen Hospital Formula		Teriflunomide		Travoprost	
Active		Teriparatide		Treatments for Dementia	14
Sustanon Ampoules		Testosterone		Treatments for Substance	
Sylvant	219	Testosterone cipionate	82	Dependence	14
Symbicort Turbuhaler 100/6	234	Testosterone esters	82	Trental 400	5
Symbicort Turbuhaler 200/6	234	Testosterone undecanoate	82	Tretinoin	
Symbicort Turbuhaler 400/12	234	Tetrabenazine	119	Dermatological	6
Symmetrel		Tetrabromophenol	78	Oncology	15
Symmetrel S29		Tetracosactrin		Trexate	
Sympathomimetics		Tetracycline		Triamcinolone acetonide	
Synacthen		Teva Lisinopril		Alimentary	3
Synacthen Depot		Thalidomide		Dermatological	
Synacthene Retard		Thalomid		Hormone	
Synadic Netard	202	Theophylline		Triamcinolone acetonide with	0
Synagis	203				totio
Synflorix		Thiamine hydrochloride		gramicidin, neomycin and nys	
Synthroid		THIO-TEPA		Dermatological	
Syntometrine		Thioguanine		Sensory	
Syrup (pharmaceutical grade)		Thiotepa		Triazolam	
Systane Unit Dose	245	Thyroid and Antithyroid Agents		Trimethoprim	9
- T -		Ticagrelor		Trimethoprim with	
Tacrolimus		Ticagrelor Sandoz	42	sulphamethoxazole	
Dermatological	70	Tilcotil	110	[Co-trimoxazole]	9
Oncology	230	Timolol	243	Trisequens	8
Tacrolimus Sandoz	230	Timoptol XE	243	Trisul	9
Taliglucerase alfa	32	Tiotropium bromide	236	Trophic Hormones	8
Tambocor	50	Tiotropium bromide with		Tropicamide	24
Tamoxifen citrate		olodaterol	236	Trulicity	
Tamoxifen Sandoz		Tivicay		Trusopt	
Tamsulosin hydrochloride		Tixagevimab with cilgavimab		TruSteel	
Tamsulosin-Rex		TMP		Tuberculin PPD [Mantoux] test	
Tandem Cartridge		Tobramycin		Tubersol	
Tandem t:slim X2 with Basal-IC		Infection	06	Two Cal HN	
Tap water		Sensory	241	Tykerb	
Tarceva		Tobramycin BNM		Tysabri	13
Taro-Testosterone		Tobramycin Mylan		- U -	_
Tasigna		Tobrex		UK Synacthen	
Tasmar		Tocilizumab		Ultibro Breezhaler	
Taurine		Tofranil		Ultraproct	
Tecfidera	135	Tolcapone		Umeclidinium	
Tegretol		Topamax	128	Umeclidinium with vilanterol	
Tegretol CR	126	Topical Products for Joint and		Univent2	235, 23

Upadacitinib	231	Vimpat	127	Zincaps	37
Ural		Vinblastine sulphate		Zinnat	
Urea		Vincristine sulphate		Ziprasidone	
Urex Forte		Vinorelbine		Zista	
Urinary Agents		Vinorelbine Ebewe		Zithromax	
Urinary Tract Infections		Viramune Suspension		Zoledronic acid	
Urinorm		ViruPOS		Hormone	70
Uromitexan		Vit.D3		Musculoskeletal	
Ursodeoxycholic acid		Vita-B12		Zoledronic acid Mylan	
Ursosan		VitA-POS		Zoledronic acid Viatris	
Utrogestan		Vitabdeck		Zopiclone	
- V -	00	Vital		Zopiclone Actavis	
Vaccinations	272	Vitamin B complex		Zostavax	
Vaclovir		Vitamin B6 25		Zostrix	
		Vitamins		Zostrix HP	
Valaciclovir				Zuclopenthixol decanoate	
Valganciclovir		Vivonex TEN		•	
Valganciclovir Mylan	102	Voltaren		Zuclopenthixol hydrochloride	
Vancomycin		Voltaren D		Zusdone	
Vannair		Voltaren Ophtha		Zyban	
Varenicline Pfizer		Voltaren SR		Zypine	131
Varenicline tartrate	142	Volumatic		Zypine ODT	
Varicella vaccine [Chickenpox		Voriconazole		Zyprexa Relprevv	
vaccine]		Votrient		Zytiga	164
Varicella zoster vaccine [Shing		Vttack	98		
vaccine]		- W -			
Varicella zoster virus (Oka stra		Warfarin sodium			
attenuated vaccine [shingles		Wart Preparations			
vaccine]		Wasp venom allergy treatm	ent232		
Various		Water			
Varivax		Blood			
Vasodilators		Extemporaneous			
Vasopressin Agonists		Wool fat with mineral oil	66		
Vasorex		- X -			
Vebulis		Xarelto			
Vedafil		Xifaxan			
Veletri	61	XMET Maxamum	265		
Venclexta	156	Xolair	202		
Venetoclax	156	XP Maxamum	266		
Venlafaxine	126	Xylocaine	120		
Venomil	232	Xylocaine 2% Jelly	119		
VENOX	232	Xyntha	41		
Ventavis	61	- Z -			
Ventolin		Zapril			
Vepesid	151	Zarontin	126		
Verapamil hydrochloride	53	Zaroxolyn	54		
Vermox	90	Zavedos	151		
Versacloz	131	Zeffix	101		
Vesanoid	156	Zematop	70		
Vexazone	12	Zetlam			
Vfend	98	Ziagen	106		
Viaderm KC		Zidovudine [AZT]	106		
Vigabatrin	128	Zidovudine [AZT] with			
Vigisom		lamivudine	106		
Vildagliptin		Zimybe	57		
Vildagliptin with metformin		Zinc and castor oil	65		
hydrochloride	12	Zinc sulphate	37		